



Australian Government

Department of Health and Ageing

**SCHEDULE OF PHARMACEUTICAL
BENEFITS FOR APPROVED
PHARMACISTS AND MEDICAL
PRACTITIONERS**

**OPERATIVE FROM 1 DECEMBER 2005
(ALL PREVIOUS EDITIONS CANCELLED)**

© Commonwealth of Australia 2005
ISSN 1037-3667

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced by any process without written permission from the Commonwealth available from Information Services. Requests and inquiries concerning reproduction and rights should be addressed to the Manager, Copyright Services, Information Services, GPO Box 1920, Canberra ACT 2601 or by e-mail Cwealthcopyright@finance.gov.au.

This Schedule provides information on the arrangements for the prescribing of pharmaceutical benefits by medical practitioners and participating dental practitioners, and the supply of pharmaceutical benefits by approved pharmacists, approved medical practitioners and approved hospital authorities. These arrangements operate under the *National Health Act 1953*. However, at the time of printing, the relevant legislation giving authority for the changes included in this issue of the Schedule may still be subject to the usual Parliamentary scrutiny. This book is not a legal document, and, in cases of discrepancy, the legislation will be the source document for payment for the supply of pharmaceutical benefits.

CONTENTS

	Page		Page
SUMMARY OF CHANGES	5	<i>Repeat authorisations for injectables and solvents</i>	<i>40</i>
ADDRESSES — MEDICARE AUSTRALIA	17	<i>Repeat authorisations for deferred supply</i>	<i>40</i>
AUTHORITY PRESCRIPTION APPLICATIONS	18	Authority PBS prescriptions	41
REQUESTS FOR DRUGS VIA THE SPECIAL ACCESS SCHEME (SAS)	18	Urgent cases	41
POISONS INFORMATION CENTRES	19	Receipts	41
DRUG INFORMATION CENTRES	19	Emergency drug (doctor's bag) supplies	41
LIST OF CONTACT OFFICERS FOR RECALLS OF THERAPEUTIC GOODS	20		
INDEX OF MANUFACTURERS' CODES	21		
SECTION 1 — EXPLANATORY NOTES — YELLOW PAGES			
INTRODUCTION	31		
1. THE SCHEDULE — WHERE TO FIND WHAT	31	4. PATIENT CHARGES	42
Section 1 (yellow pages)	31	Type of patient	42
Section 2 (white, lavender, pink and orange pages)	31	Establishing entitlement	42
Section 3 (blue pages)	32	What to charge	43
Section 4 (green pages)	32	<i>Patient contribution</i>	<i>43</i>
Repatriation Schedule of Pharmaceutical Benefits	32	<i>Patient contributions for early supply of some PBS medicines</i>	<i>43</i>
		<i>Special patient contributions, brand premiums and therapeutic group premiums</i>	<i>44</i>
2. PRESCRIBING MEDICINES — INFORMATION FOR DOCTORS AND DENTISTS	32	<i>Solvents</i>	<i>44</i>
Eligible prescribers	32	<i>Increased quantities</i>	<i>44</i>
PBS prescription forms	32	<i>Regulation 24</i>	<i>44</i>
<i>Ordering forms</i>	<i>33</i>	<i>After hours</i>	<i>44</i>
Preparing general PBS prescriptions	33	<i>Delivery</i>	<i>44</i>
<i>Do's and don't's</i>	<i>33</i>		
<i>Writing the PBS prescription</i>	<i>34</i>	5. THE SAFETY NET SCHEME	45
Restrictions	35	Safety net thresholds	45
Authority PBS prescriptions	35	<i>Safety net cross-over arrangements</i>	<i>46</i>
<i>Writing authority PBS prescriptions</i>	<i>35</i>	Recording PBS prescriptions	46
Maximum quantities and repeats	36	Hospital PRF's	47
Regulation 24	37	Multi-item forms	47
Urgent cases	37	Qualifying PBS prescriptions	47
Drugs of addiction	37	Lost PRF's	47
Emergency drug (doctor's bag) supplies	37	Retrospective entitlement and patient refunds	48
		Applying for a Safety Net Entitlement/Concession Card	48
3. SUPPLYING MEDICINES — WHAT PHARMACISTS NEED TO KNOW	38	Issuing a Safety Net Entitlement/Concession Card	49
Eligible suppliers	38	Issuing supplementary cards	49
<i>Approval conditions for pharmacists</i>	<i>38</i>	Notification to Medicare Australia and claim for payment	49
Before supplying pharmaceutical benefits	38	Lost Safety Net Entitlement/Concession Cards	49
Supplying pharmaceutical benefits	39	Pharmacy record of issued cards	49
Do's and don't's	39		
What to do if the Schedule changes	39	6. MEDICARE AUSTRALIA ENTITLEMENT CHECKS	50
Suspected forgery	40	<i>General</i>	<i>50</i>
Regulation 24	40	<i>Concessional</i>	<i>50</i>
Repeat authorisations	40	Entitlement checking procedures	50
<i>Preparing repeat authorisation forms</i>	<i>40</i>	<i>General</i>	<i>50</i>
		<i>Concessional</i>	<i>50</i>
		<i>Step by step</i>	<i>51</i>
		7. HOW PHARMACISTS CLAIM REIMBURSEMENT: INFORMATION REQUIRED	51
		PBS prescription identification	51
		Serial numbers	52

	<i>Page</i>
<i>Repeat authorisations for authority</i>	
<i>PBS prescriptions</i>	52
<i>Repeat authorisations for deferred supply</i>	52
<i>Injectable item ordered with a solvent</i>	52
Dropper containers	52
Extemporaneously-prepared pharmaceutical	
benefits not included in the Standard Formulae	
List	52
<i>PBS prescriptions paid on an average price</i>	
<i>basis</i>	53
<i>Pricing non-pre-priced extemporaneous</i>	
<i>preparations</i>	53
RPBS prescriptions for items not included in either	
the PBS or RPBS Schedule	53
8. HOW PHARMACISTS CLAIM REIMBURSEMENT:	
DOCUMENTS TO BE SUBMITTED	53
Completing the claim form	54
Lodging claims	54
Reconciliation statements	54
9. PRICING PBS PRESCRIPTIONS	55
Pricing principles	55
Pricing dates	55
Pricing ready-prepared items	55
<i>For maximum quantities</i>	55
<i>For lesser quantities</i>	55
Wastage table percentage	56
Pricing extemporaneously-prepared items	56
<i>General</i>	56
<i>Pricing of ingredients</i>	57
<i>Pricing PBS prescriptions where extra</i>	
<i>ingredients are added to a formula</i>	57
<i>Containers</i>	57
<i>Special provisions for extemporaneous</i>	
<i>PBS prescriptions outside the Standard</i>	
<i>Formulae List</i>	57
10. MISCELLANEOUS	58
References	58
Standards	58
Legislation	58

THERAPEUTIC INDEX — MAUVE PAGES 59

	<i>Page</i>
SECTION 2 — READY-PREPARED	
PHARMACEUTICAL BENEFITS —	
WHITE, LAVENDER, PINK AND ORANGE PAGES	67
Symbols	68
Emergency Drug (Doctor's Bag) Supplies	71
Special Pharmaceutical Benefits	74
General Pharmaceutical Benefits	77
Pharmaceutical Benefits for Palliative Care	319
Pharmaceutical Benefits for Dental Use	333
Items Available under Special Arrangements	
(section 100)	357
SECTION 3 — STANDARD PACKS AND PRICES —	
BLUE PAGES	433
Container Prices and Fees for Ready Prepared	
Pharmaceutical Benefits	433
Standard Packs and Prices for Ready Prepared	
Pharmaceutical Benefits	434
SECTION 4 — EXTEMPOREANEOUSLY-PREPARED	
PHARMACEUTICAL BENEFITS —	
GREEN PAGES	447
Drug Tariff	448
Container Prices	452
Standard Formula Preparations	453
Table of Codes, Maximum Quantities and Number	
of Repeats for Extemporaneously Prepared	
Pharmaceutical Benefits	455
REPATRIATION PHARMACEUTICAL BENEFITS —	
WHITE, MAUVE AND BUFF PAGES	457
Beneficiaries' Entitlement Cards	458
Explanatory Notes	460
Summary of Changes	465
Therapeutic Index	467
SECTION 1 — Drugs, Medicines and Dressings	471
SECTION 2 — Standard Packs and Prices	513
GENERIC/PROPRIETARY INDEX —	
MAUVE PAGES	517

PHARMACEUTICAL BENEFITS

This Schedule will take effect on 1 December 2005 and all previous issues are cancelled. New Schedules will take effect on 1 April, 1 August and 1 December each year.

Change of Address

Notification of change of address should be directed to the Pharmaceutical Branch, Medicare Australia, GPO Box 9826, in your Capital City, or telephone 132 290.

Internet

The Schedule of Pharmaceutical Benefits is also available on the Department of Health and Ageing's internet site at www.health.gov.au.

The address of the Schedule is <http://www1.health.gov.au/pbs/index.htm>.

Fees, Patient Contributions and Safety Net Thresholds

The following fees, patient contributions and safety net thresholds apply as at 1 December 2005 and are included, where applicable, in prices published in the Schedule—

Dispensing Fees:	Ready-prepared	\$4.75
	Dangerous drug fee	\$2.61
	Extemporaneously-prepared	\$6.78
Additional Fees (for safety net prices):	Ready-prepared	\$0.97
	Extemporaneously-prepared	\$1.38
Patient Co-payments:	General	\$28.60
	Concessional	\$4.60
Safety Net Thresholds:	General	\$874.90
	Concessional	\$239.20
Safety Net Card Issue Fee:		\$7.21

SUMMARY OF CHANGES

ADDITIONS

Additions — Items

- 8865N **Buprenorphine**, transdermal patch 5 mg (releasing approximately 5 micrograms per hour) (*Norspan*)
- 8866P **Buprenorphine**, transdermal patch 10 mg (releasing approximately 10 micrograms per hour) (*Norspan*)
- 8867Q **Buprenorphine**, transdermal patch 20 mg (releasing approximately 20 micrograms per hour) (*Norspan*)
- 8864M **Coal Tar—Prepared**, gel 10 mg per g (1%), 100 mL (*Exorex*)
- 8858F **Epirubicin Hydrochloride**, solution for injection 200 mg in 100 mL (*Epirubicin Ebewe*)
- 8861J **Etanercept**, injection set containing 4 vials powder for injection 50 mg and 4 pre-filled syringes solvent 1 mL (*Enbrel*)
- 8862K **Etanercept**, injection set containing 4 vials powder for injection 50 mg and 4 pre-filled syringes solvent 1 mL (*Enbrel*) (**Diff. Max. Rpts**)
- 8835B **Follitropin Beta**, solution for injection 200 i.u. in 0.5 mL (*Puregon 200 IU/0.5 mL*) (effective 1 August 2005)
- 8871X **Follitropin Beta**, solution for injection 900 i.u. in 1.08 mL multi-dose cartridge (*Puregon 900 IU/1.08 mL*)
- 8873B **Granisetron Hydrochloride**, tablet 2 mg (base) (*Kytril*)
- 8859G **Leuporelin Acetate**, suspension for subcutaneous injection (modified release), 45 mg injection set (*Eligard 6 month*)
- 2875H **Lignocaine Hydrochloride**, injection 100 mg in 5 mL (*PF*) (effective 1 October 2005)
- 5142P **Lignocaine Hydrochloride**, injection 100 mg in 5 mL (*PF*) (**Dental**) (effective 1 October 2005)
- 3474W **Lignocaine Hydrochloride**, injection 100 mg in 5 mL (*PF*) (**Doctor's Bag**) (effective 1 October 2005)
- 8863L **Methotrexate**, solution concentrate for I.V. infusion 500 mg in 20 mL (*MX*)
- 8855C **Mirtazapine**, tablet 15 mg (orally disintegrating) (*Avanza SolTab*)
- 8856D **Mirtazapine**, tablet 30 mg (orally disintegrating) (*Avanza SolTab*)
- 8857E **Mirtazapine**, tablet 45 mg (orally disintegrating) (*Avanza SolTab*)
- 8860H **Quinagolide Hydrochloride**, pack containing 3 tablets 25 micrograms (base) and 3 tablets 50 micrograms (base) (*Norprolac*)
- 8869T **Risperidone**, tablet 0.5 mg (*Risperdal*)
- 8870W **Risperidone**, tablet 0.5 mg (orally disintegrating) (*Risperdal Quicklet*)
- 8836C **Sertraline Hydrochloride**, tablet 50 mg (base) (*Eleva 50, Xydep 50, Zoloft*)
- 8837D **Sertraline Hydrochloride**, tablet 100 mg (base) (*Eleva 100, Xydep 100, Zoloft*)
- 8868R **Venlafaxine Hydrochloride**, capsule 37.5 mg (base) (modified release) (*Efexor-XR*)

The following items, which may be prescribed for patients receiving palliative care, have been added to the Palliative Care section in the Schedule:

- 5357Y **Diazepam**, tablet 2 mg (*Antenex 2; Valpam 2; Ducene; Valium*)
- 5355W **Diazepam**, tablet 2 mg (*Antenex 2; Valpam 2; Ducene; Valium*) (**Diff. Max. Rpts**)
- 5358B **Diazepam**, tablet 5 mg (*Antenex 5; Diazepam-DP; Valpam 5; Ducene; Valium*)
- 5356X **Diazepam**, tablet 5 mg (*Antenex 5; Diazepam-DP; Valpam 5; Ducene; Valium*) (**Diff. Max. Rpts**)
- 5364H **Diclofenac Sodium**, tablet 25 mg (enteric coated) (*Chem mart Diclofenac; Clonac 25; Diclofenac-BC; Diclohexal; Dinac; GenRx Diclofenac; Terry White Chemists Diclofenac; Fenac 25; Voltaren 25*)

- 5361E **Diclofenac Sodium**, tablet 25 mg (enteric coated) (*Chem mart Diclofenac; Clonac 25; Diclofenac-BC; Diclohexal; Dinac; GenRx Diclofenac; Terry White Chemists Diclofenac; Fenac 25; Voltaren 25*) (**Diff. Max. Rpts**)
- 5365J **Diclofenac Sodium**, tablet 50 mg (enteric coated) (*Chem mart Diclofenac; Clonac 50; Diclohexal; Dinac; Fenac; GenRx Diclofenac; Terry White Chemists Diclofenac; Voltaren 50*)
- 5362F **Diclofenac Sodium**, tablet 50 mg (enteric coated) (*Chem mart Diclofenac; Clonac 50; Diclohexal; Dinac; Fenac; GenRx Diclofenac; Terry White Chemists Diclofenac; Voltaren 50*) (**Diff. Max. Rpts**)
- 5366K **Diclofenac Sodium**, suppository 100 mg (*Voltaren 100*)
- 5363G **Diclofenac Sodium**, suppository 100 mg (*Voltaren 100*) (**Diff. Max. Rpts**)
- 5369N **Ibuprofen**, tablet 200 mg (*Rafen 200*)
- 5367L **Ibuprofen**, tablet 200 mg (*Rafen 200*) (**Diff. Max. Rpts**)
- 5370P **Ibuprofen**, tablet 400 mg (*Brufen*)
- 5368M **Ibuprofen**, tablet 400 mg (*Brufen*) (**Diff. Max. Rpts**)
- 5379D **Indomethacin**, capsule 25 mg (*Arthrexin; Indocid*)
- 5377B **Indomethacin**, capsule 25 mg (*Arthrexin; Indocid*) (**Diff. Max. Rpts**)
- 5380E **Indomethacin**, suppository 100 mg (*Indocid*)
- 5378C **Indomethacin**, suppository 100 mg (*Indocid*) (**Diff. Max. Rpts**)
- 5349M **Naproxen**, tablet 250 mg (*Inza 250; Naprosyn*)
- 5345H **Naproxen**, tablet 250 mg (*Inza 250; Naprosyn*) (**Diff. Max. Rpts**)
- 5350N **Naproxen**, tablet 500 mg (*Inza 500; Naprosyn*)
- 5346J **Naproxen**, tablet 500 mg (*Inza 500; Naprosyn*) (**Diff. Max. Rpts**)
- 5351P **Naproxen**, tablet 750 mg (sustained release) (*Proxen SR 750; Naprosyn SR750*)
- 5347K **Naproxen**, tablet 750 mg (sustained release) (*Proxen SR 750; Naprosyn SR750*) (**Diff. Max. Rpts**)
- 5352Q **Naproxen**, tablet 1 g (sustained release) (*Proxen SR 1000; Naprosyn SR1000*)
- 5348L **Naproxen**, tablet 1 g (sustained release) (*Proxen SR 1000; Naprosyn SR1000*) (**Diff. Max. Rpts**)
- 5354T **Naproxen Sodium**, tablet 550 mg (*Anaprox 550; Crysanal*)
- 5353R **Naproxen Sodium**, tablet 550 mg (*Anaprox 550; Crysanal*) (**Diff. Max. Rpts**)
- 5360D **Nitrazepam**, tablet 5 mg (*Alodorm; Mogadon*)
- 5359C **Nitrazepam**, tablet 5 mg (*Alodorm; Mogadon*) (**Diff. Max. Rpts**)
- 5373T **Oxazepam**, tablet 15 mg (*Alepam 15; Serepax*)
- 5371Q **Oxazepam**, tablet 15 mg (*Alepam 15; Serepax*) (**Diff. Max. Rpts**)
- 5374W **Oxazepam**, tablet 30 mg (*Alepam 30; Murelax; Serepax*)
- 5372R **Oxazepam**, tablet 30 mg (*Alepam 30; Murelax; Serepax*) (**Diff. Max. Rpts**)
- 5383H **Sulindac**, tablet 100 mg (*Aclin*)
- 5381F **Sulindac**, tablet 100 mg (*Aclin*) (**Diff. Max. Rpts**)
- 5384J **Sulindac**, tablet 200 mg (*Aclin 200*)
- 5382G **Sulindac**, tablet 200 mg (*Aclin 200*) (**Diff. Max. Rpts**)
- 5376Y **Temazepam**, tablet 10 mg (*Temaze; Temtabs; Normison*)
- 5375X **Temazepam**, tablet 10 mg (*Temaze; Temtabs; Normison*) (**Diff. Max. Rpts**)

Additions — Brands

- 8511Y *Alendro Once Weekly, AW* — **Alendronate Sodium**, tablet equivalent to 70 mg alendronic acid
- 1259G *Lax-Tab, AE* — **Bisacodyl**, tablet 5 mg
- 5305F *Lax-Tab, AE* — **Bisacodyl**, tablet 5 mg (**Palliative Care**)
- 5301B *Lax-Tab, AE* — **Bisacodyl**, tablet 5 mg (**Palliative Care Diff. Max. Rpts**)
- 8220P *Ciazil, DP* — **Citalopram Hydrobromide**, tablet 20 mg (base)

- 8703C *GenRx Citalopram, GX* — **Citalopram Hydrobromide**, tablet 40 mg (base)
- 8318T *Clarihexal, HX* — **Clarithromycin**, tablet 250 mg
- 3196F *Chem mart Oral Rehydration Salts, CH; Terry White Chemists Oral Rehydration Salts, TW* — **Electrolyte Replacement (Oral)**, sachets containing powder for oral solution 4.9 g, 10
- 1375J *Epirubicin Ebewe, IT* — **Epirubicin Hydrochloride**, solution for injection 10 mg in 5 mL
- 1377L *Epirubicin Ebewe, IT* — **Epirubicin Hydrochloride**, solution for injection 50 mg in 25 mL
- 8817C *Epirubicin Ebewe, IT* — **Epirubicin Hydrochloride**, solution for injection 100 mg in 50 mL
- 1390E *Etoposide Ebewe, IT* — **Etoposide**, solution for I.V. infusion 100 mg in 5 mL
- 2413B *Frusehexal, HX* — **Frusemide**, injection 20 mg in 2 mL
- 3466K *Frusehexal, HX* — **Frusemide**, injection 20 mg in 2 mL (**Doctor's Bag**)
- 1459T *Nitrostat, LM* — **Glyceryl Trinitrate**, tablet 600 micrograms, 100 (effective 2 September 2005)
- 5108W *Nitrostat, LM* — **Glyceryl Trinitrate**, tablet 600 micrograms, 100 (**Dental**) (effective 2 September 2005)
- 8063J *Elmendos, ME* — **Lamotrigine**, tablet 5 mg
- 2430X *Formet 500, AW* — **Metformin Hydrochloride**, tablet 500 mg
- 1801T *Formet 850, AW* — **Metformin Hydrochloride**, tablet 850 mg
- 8607B *Formet 1000, AW* — **Metformin Hydrochloride**, tablet 1 g
- 8851W *MX* — **Methotrexate**, solution concentrate for I.V. infusion 1000 mg in 10 mL
- 8839F *Attenta, AF* — **Methylphenidate Hydrochloride**, tablet 10 mg
- 1746X *Terry White Chemists Paracetamol, TW* — **Paracetamol**, tablet 500 mg
- 5196L *Terry White Chemists Paracetamol, TW* — **Paracetamol**, tablet 500 mg (**Dental**)
- 8784H *Terry White Chemists Paracetamol, TW* — **Paracetamol**, tablet 500 mg (**Diff. Max. Qty and Rpts**)
- 5224Y *Terry White Chemists Paracetamol, TW* — **Paracetamol**, tablet 500 mg (**Diff. Max. Qty (Dental)**)
- 2833D *Lipostat 10, AW* — **Pravastatin Sodium**, tablet 10 mg
- 2834E *Lipostat 20, AW* — **Pravastatin Sodium**, tablet 20 mg
- 8197K *Lipostat 40, AW* — **Pravastatin Sodium**, tablet 40 mg
- 8829Q *Lipostat 80, AW* — **Pravastatin Sodium**, tablet 80 mg
- 2236Q *Chem mart Sertraline, CH; Concorz, HX; Eleva 50, AF; GenRx Sertraline, GX; Sertraline-DP, DP; Terry White Chemists Sertraline, TW* — **Sertraline Hydrochloride**, tablet 50 mg (base)
- 2237R *Chem mart Sertraline, CH; Concorz, HX; Eleva 100, AF; GenRx Sertraline, GX; Sertraline-DP, DP; Terry White Chemists Sertraline, TW* — **Sertraline Hydrochloride**, tablet 100 mg (base)
- 2013Y *Simvastatin-RL, RE* — **Simvastatin**, tablet 5 mg
- 2011W *Simvastatin-RL, RE* — **Simvastatin**, tablet 10 mg
- 2012X *Simvastatin-RL, RE* — **Simvastatin**, tablet 20 mg
- 8173E *Simvastatin-RL, RE* — **Simvastatin**, tablet 40 mg
- 8313M *Simvastatin-RL, RE* — **Simvastatin**, tablet 80 mg
- 2804N *GenRx Terbinafine, GX; Tamsil, AW; Terbihexal, HX; Terbinafine-DP, DP* — **Terbinafine Hydrochloride**, tablet 250 mg (base)

Additions of Extemporaneous Preparations —Drug Tariff

Boric Acid, Olive Oil and Zinc Oxide Ointment QHF

Gentian Alkaline Mixture APF

Kaolin Mixture BPC 1968

Kaolin and Opium Mixture APF 14

RETENTION OF ITEM

Contrary to previous advice, the following items will continue to be available as a pharmaceutical benefits from 1 December 2005:

8760C **Captopril**, oral solution 5 mg per mL, 95 mL (*Capoten*)
 3483H **Pethidine Hydrochloride**, injection 100 mg in 2 mL (MX)

BIOEQUIVALENCE INDICATORS

The bioequivalence indicator (a) has been added to the following brands:

8511Y *Fosamax Once Weekly, MK* — **Alendronate Sodium**, tablet equivalent to 70 mg alendronic acid
 1390E MX — **Etoposide**, solution for I.V. infusion 100 mg in 5 mL
 2413B *Lasix, AV* — **Furosemide**, injection 20 mg in 2 mL
 3466K *Lasix, AV* — **Furosemide**, injection 20 mg in 2 mL (**Doctor's Bag**)
 8851W *Methotrexate Ebewe, IT* — **Methotrexate**, solution concentrate for I.V. infusion 1000 mg in 10 mL
 8839F *Ritalin 10, NV* — **Methylphenidate Hydrochloride**, tablet 10 mg
 2833D *Pravachol, BQ* — **Pravastatin Sodium**, tablet 10 mg
 2834E *Pravachol, BQ* — **Pravastatin Sodium**, tablet 20 mg
 8197K *Pravachol, BQ* — **Pravastatin Sodium**, tablet 40 mg
 8829Q *Pravachol, BQ* — **Pravastatin Sodium**, tablet 80 mg

DELETIONS

Deletions — Items

1005X **Acetazolamide**, injection 500 mg (solvent required) (*Diamox*)
 2603B **Allopurinol**, capsule 300 mg (*Capurate-300*)
 1068F **Gentamicin Sulfate**, injection 40 mg (base) in 1 mL (MX)
 1168L **Gentamicin Sulfate**, injection 60 mg (base) in 1.5 mL (MX)
 2952J **Hypromellose**, eye drops 10 mg per mL (1%), 15 mL (*Methopt Forte*)
 2586D **Isosorbide Dinitrate**, tablet 20 mg (*Isordil*)
 2854F **Demeclocycline Hydrochloride**, capsule 150 mg (*Ledermycin*)
 1458R **Levonorgestrel with Ethinylloestradiol**, pack containing 11 tablets 50 micrograms-50 micrograms, 10 tablets 125 micrograms-50 micrograms and 7 inert tablets (*Sequilar ED*)
 1391F **Levonorgestrel with Ethinylloestradiol**, pack containing 6 tablets 50 micrograms-30 micrograms, 5 tablets 75 micrograms-40 micrograms and 10 tablets 125 micrograms-30 micrograms (*Triquilar*)
 2731R **Megestrol Acetate**, tablet 40 mg (*Megace*)

- 1687T **Nicotinic Acid**, tablet 250 mg (*AL*)
- 1765X **Oestradiol and Oestradiol with Norethisterone Acetate**, pack containing 12 tablets oestradiol 4 mg, 10 tablets oestradiol 4 mg with norethisterone acetate 1 mg and 6 tablets oestradiol 1 mg (*Trisequens Forte*)
- 8429P **Oestradiol Valerate and Oestradiol Valerate with Cyproterone Acetate**, pack containing 16 tablets oestradiol valerate 2 mg and 12 tablets oestradiol valerate 2 mg with cyproterone acetate 1 mg (*Climen 28*)
- 8210D **Oestrogens—Conjugated and Oestrogens—Conjugated with Medroxyprogesterone Acetate**, pack containing 14 tablets conjugated oestrogens 625 micrograms and 14 tablets conjugated oestrogens 625 micrograms with medroxyprogesterone acetate 5 mg (*Premia 5*)
- 8538J **Oestrogens—Conjugated and Oestrogens—Conjugated with Medroxyprogesterone Acetate**, pack containing 14 tablets conjugated oestrogens 625 micrograms and 14 tablets conjugated oestrogens 625 micrograms with medroxyprogesterone acetate 10 mg (*Premia 10*)
- 8680W **Oestrogens—Conjugated with Medroxyprogesterone Acetate**, tablets 450 micrograms-1.5 mg, 28 (*Premia Low*)
- 2496J **Pancrelipase**, capsule providing not less than 5,000 BP units of lipase activity (*Pancrease*)
- 2653P **Procainamide Hydrochloride**, capsule 250 mg (*Pronestyl*)
- 2024M **Sodium Chloride**, injection 9 mg per mL (0.9%), 2 mL (*AP*)
- 5209E **Sodium Chloride**, injection 9 mg per mL (0.9%), 2 mL (*AP*) (**Dental**)
- 2026P **Sodium Chloride**, injection 9 mg per mL (0.9%), 10 mL (*AP*)
- 5211G **Sodium Chloride**, injection 9 mg per mL (0.9%), 10 mL (*AP*) (**Dental**)
- 3090P **Spectinomycin**, injection 2 g with 3.2 mL diluent (*Trobicin*)
- 2145X **Tetracycline Hydrochloride (Buffered)**, capsule 250 mg (*Tetrex*)
- 2146Y **Tetracycline Hydrochloride (Buffered)**, capsule 250 mg (*Tetrex*) (**Diff. Max. Qty and Rpts**)
- 3386F **Tetracycline Hydrochloride (Buffered)**, capsule 250 mg (*Tetrex*) (**Dental**)
- 2218R **Water For Injections, Sterilised**, injection 10 mL (*AP*)
- 3396R **Water For Injections, Sterilised**, injection 10 mL (*AP*) (**Dental**)

Deletion of codes for injectable items with allowable solvents:

Previously the entry in the Schedule of Pharmaceutical Benefits for certain injectable items which require a solvent included codes for the items when prescribed with approved solvents. As all allowable solvents have been deleted from this edition of the Schedule due to their discontinuation by the manufacturer, the entries for these injectable items have been amended to delete reference to the codes which relate to the injectable with the approved solvents.

Deletions — Brands

- 2157M *Sigma Liquid Antacid, SI* — **Aluminium Hydroxide with Magnesium Hydroxide**, oral suspension 200 mg-200 mg per 5 mL, 500 mL
- 1169M *Cefaclor CD Hexal, HX* — **Cefaclor**, tablet 375 mg (sustained release)
- 5045M *Cefaclor CD Hexal, HX* — **Cefaclor**, tablet 375 mg (sustained release) (**Dental**)
- 1312C *Diltahexal CD, HX* — **Diltiazem Hydrochloride**, capsule 180 mg (controlled delivery)
- 1313D *Diltahexal CD, HX* — **Diltiazem Hydrochloride**, capsule 240 mg (controlled delivery)
- 1375J *MX* — **Epirubicin Hydrochloride**, solution for injection 10 mg in 5 mL
- 1376K *MX* — **Epirubicin Hydrochloride**, solution for injection 20 mg in 10 mL
- 1377L *MX* — **Epirubicin Hydrochloride**, solution for injection 50 mg in 25 mL

8817C	<i>MX</i> — Epirubicin Hydrochloride , solution for injection 100 mg in 50 mL
2487X	<i>Amfamox 20, AD</i> — Famotidine , tablet 20 mg
2488Y	<i>Amfamox 40, AD</i> — Famotidine , tablet 40 mg
1417N	<i>Fugerel, EX</i> — Flutamide , tablet 250 mg
2412Y	<i>Frusemide-BC, BG</i> — Frusemide , tablet 40 mg
1453L	<i>Gemfibrozil-BC, BG</i> — Gemfibrozil , tablet 600 mg

Deletions of Extemporaneous Preparations — Drug Tariff

Ammonia Aromatic Spirit BP
Cetostearyl Alcohol BP
Chlorinated Lime BP
Gentian Compound Infusion Concentrated 1 in 10 BP
Hydrochloric Acid Dilute BP
Ipecacuanha Tincture BP
Kaolin, Light or Kaolin, Light (Natural) BP
Opium Tincture BP
Orange Syrup BP
Orange Tincture BP
Paraffin Ointment (white) BP
Paraffin Ointment (yellow) BP 1968
Zinc Ointment BP

Deletions of Extemporaneous Preparations — Standard Formulae List

7599Y	Mixtures, Other — Gentian Alkaline (extemporaneous formula) BP
7539T	Solutions — Calcium Hypochlorite (Eusol) APF 15

ALTERATIONS

Restriction Changes

The following items have changed from “restricted benefit” to “authority required”:

8560M	Calcium , tablet 250 mg (as citrate) (<i>Citracal</i>)
3116B	Calcium , tablet (chewable) 500 mg (as carbonate) (<i>Cal-Sup</i>)
3117C	Calcium , tablet 600 mg (as carbonate) (<i>Caltrate</i>)

Other restriction changes:

1007B	Aciclovir , tablet 200 mg (<i>Acihexal; Acyclo-V 200; Chem mart Aciclovir; GenRx Aciclovir; Lovir; Terry White Chemists Aciclovir; Zyclir 200; Zovirax 200 mg</i>)
8560M	Calcium , tablet 250 mg (as citrate) (<i>Citracal</i>)
3116B	Calcium , tablet (chewable) 500 mg (as carbonate) (<i>Cal-Sup</i>)
3117C	Calcium , tablet 600 mg (as carbonate) (<i>Caltrate</i>)
8092X	Famciclovir , tablet 125 mg (<i>Famvir</i>)
8217L	Famciclovir , tablet 250 mg (<i>Famvir</i>)
8565T	Follitropin Beta , solution for injection 300 i.u. in 0.36 mL multi-dose cartridge (<i>Puregon 300 IU/0.36 mL</i>)
8566W	Follitropin Beta , solution for injection 600 i.u. in 0.72 mL multi-dose cartridge (<i>Puregon 600 IU/0.72 mL</i>)

8539K	Oxaliplatin , powder for I.V. infusion 50 mg (<i>Eloxatin</i>) (effective 1 October 2005)
8847P	Oxaliplatin , solution concentrate for I.V. infusion 50 mg in 10 mL (<i>Eloxatin</i>) (effective 1 October 2005)
8540L	Oxaliplatin , powder for I.V. infusion 100 mg (<i>Eloxatin</i>) (effective 1 October 2005)
8848Q	Oxaliplatin , solution concentrate for I.V. infusion 100 mg in 20 mL (<i>Eloxatin</i>) (effective 1 October 2005)
8694N	Pioglitazone Hydrochloride , tablet 15 mg (base) (<i>Actos</i>)
8695P	Pioglitazone Hydrochloride , tablet 30 mg (base) (<i>Actos</i>)
8696Q	Pioglitazone Hydrochloride , tablet 45 mg (base) (<i>Actos</i>)
8689H	Rosiglitazone Maleate , tablet 4 mg (base) (<i>Avandia</i>)
8690J	Rosiglitazone Maleate , tablet 8 mg (base) (<i>Avandia</i>)
2804N	Terbinafine Hydrochloride , tablet 250 mg (<i>GenRx Terbinafine, Lamisil, Tamsil, Terbihexal, Terbinafine-DP, Zabil</i>)
8134D	Valaciclovir Hydrochloride , tablet 500 mg (base)

Alterations — Notes

Notes have been added in respect of the following:

Oxaliplatin (effective 1 October 2005)

Notes have been deleted in respect of the following:

Aciclovir
Famciclovir
Valaciclovir Hydrochloride

Notes have been amended in respect of the following:

Hydromorphone Hydrochloride
Methadone Hydrochloride
Morphine Hydrochloride
Morphine Sulfate
Oxycodone
Oxycodone Hydrochloride

Alterations — Item Code and Item Description

From:	
1356J	Tobramycin Sulfate , injection 80 mg (base) (<i>PU</i>)
To:	
8872Y	Tobramycin Sulfate , injection 80 mg (base) in 2 mL (without preservative) (<i>PU</i>)

Alterations — Item Description

From:	
3486L	Benzympenicillin , injection 600 mg (with sterilised Water for Injections 10 mL) (<i>BenPen</i>)
To:	
3486L	Benzympenicillin , powder for injection 600 mg (<i>BenPen</i>)
From:	
3487M	Benzympenicillin , injection 3 g (with sterilised Water for Injections 10 mL) (<i>BenPen</i>)
To:	
3487M	Benzympenicillin , powder for injection 3 g (<i>BenPen</i>)

From:
8289G **Interferon Beta-1a**, injection set comprising 1 vial powder for injection 30 micrograms (6,000,000 i.u.) and 1 ampoule solvent 2 mL (*Avonex*)

To:
8289G **Interferon Beta-1a**, injection set comprising 1 vial powder for injection 30 micrograms (6,000,000 i.u.) with diluent (*Avonex*)

From:
1356J **Tobramycin Sulfate**, injection 80 mg (base) (*MX, Nebcin*)

To:
1356J **Tobramycin Sulfate**, injection 80 mg (base) in 2 mL (with preservative) (*MX, Nebcin*)

Alterations — Proprietary Name

2576N **Aluminium Hydroxide with Magnesium Hydroxide**, tablet 200 mg-200 mg

From: *Mylanta*
To: *Mylanta P*

Alterations — Manufacturer's Code

		From	To
2157M	Aluminium Hydroxide with Magnesium Hydroxide , oral suspension 200 mg-200 mg per 5 mL, 500 mL (<i>Mylanta P</i>)	WR	PC
2576N	Aluminium Hydroxide with Magnesium Hydroxide , tablet 200 mg-200 mg (<i>Mylanta P</i>)	WR	PC
1891M	Amoxycillin with Clavulanic Acid , tablet 500 mg-125 mg (<i>Clamoxyl Duo</i>)	AF	AL
5008N	Amoxycillin with Clavulanic Acid , tablet 500 mg-125 mg (<i>Clamoxyl Duo</i>) (Dental)	AF	AL
8254K	Amoxycillin with Clavulanic Acid , tablet 875 mg-125 mg (<i>Clamoxyl Duo forte</i>)	AF	AL
5006L	Amoxycillin with Clavulanic Acid , tablet 875 mg-125 mg (<i>Clamoxyl Duo forte</i>) (Dental)	AF	AL
1892N	Amoxycillin with Clavulanic Acid , powder for syrup 125 mg-31.25 mg per 5 mL, 75 mL (<i>Clamoxyl</i>)	AF	AL
5009P	Amoxycillin with Clavulanic Acid , powder for syrup 125 mg-31.25 mg per 5 mL, 75 mL (<i>Clamoxyl</i>) (Dental)	AF	AL
8319P	Amoxycillin with Clavulanic Acid , powder for syrup 400 mg-57 mg per 5 mL, 60 mL (<i>Clamoxyl Duo 400</i>)	AF	AL
5011R	Amoxycillin with Clavulanic Acid , powder for syrup 400 mg-57 mg per 5 mL, 60 mL (<i>Clamoxyl Duo 400</i>) (Dental)	AF	AL
8167W	Benzathine Penicillin , injection 900 mg in 2 mL cartridge-needle unit (for use with Tubex Injector) (<i>Bicillin L-A Tubex</i>)	WY	AS
8743E	Benzathine Penicillin , injection 900 mg in 2 mL cartridge-needle unit (for use with Tubex Injector) (<i>Bicillin L-A Tubex</i>) (Diff. Max Qty)	WY	AS
5025L	Benzathine Penicillin , injection 900 mg in 2 mL cartridge-needle unit (for use with Tubex Injector) (<i>Bicillin L-A Tubex</i>) (Dental)	WY	AS
2315W	Bleomycin Sulfate , powder for injection 15,000 i.u. (<i>Blenamax</i>)	ZH	SI
1215Y	Codeine Phosphate with Paracetamol , tablet 30 mg-500 mg (<i>Codapane Forte</i>)	AF	AL
8785J	Codeine Phosphate with Paracetamol , tablet 30 mg-500 mg (<i>Codapane Forte</i>) (Diff. Max. Qty)	AF	AL
3316M	Codeine Phosphate with Paracetamol , tablet 30 mg-500 mg (<i>Codapane Forte</i>) (Dental)	AF	AL

3196F	Electrolyte Replacement (Oral) , sachets containing powder for oral solution 4.9 g, 10 (O.R.S.)	GR	AS
1824B	Mefenamic Acid , capsule 250 mg (<i>Mefic</i>)	WW	PC
2430X	Metformin Hydrochloride , tablet 500 mg (<i>Glucophage</i>)	AW	MQ
1801T	Metformin Hydrochloride , tablet 850 mg (<i>Glucophage</i>)	AW	MQ
1207M	Metoclopramide Hydrochloride , tablet 10 mg (<i>Maxolon</i>)	ID	VT
5151D	Metoclopramide Hydrochloride , tablet 10 mg (<i>Maxolon</i>) (Dental)	ID	VT
1205K	Metoclopramide Hydrochloride , syrup 5 mg per 5 mL, 100 mL (<i>Maxolon</i>)	ID	VT
5152E	Metoclopramide Hydrochloride , syrup 5 mg per 5 mL, 100 mL (<i>Maxolon</i>) (Dental)	ID	VT
1206L	Metoclopramide Hydrochloride , injection 10 mg in 2 mL	ID	VT
3476Y	Metoclopramide Hydrochloride , injection 10 mg in 2 mL (<i>Maxolon</i>) (Doctor's Bag)	ID	VT
5153F	Metoclopramide Hydrochloride , injection 10 mg in 2 mL (<i>Maxolon</i>) (Dental)	ID	VT
8370M	Naltrexone Hydrochloride , tablet 50 mg (<i>ReVia</i>)	OA	BQ
2723H	Nitrazepam , tablet 5 mg (<i>Mogadon</i>)	ID	VT
2732T	Nitrazepam , tablet 5 mg (<i>Mogadon</i>) (Diff. Max. Qty)	ID	VT
5189D	Nitrazepam , tablet 5 mg (<i>Mogadon</i>) (Dental)	ID	VT
1728Y	Olsalazine Sodium , capsule 250 mg (<i>Dipentum</i>)	PH	UC
8086N	Olsalazine Sodium , tablet 500 mg (<i>Dipentum</i>)	PH	UC
1746X	Paracetamol , tablet 500 mg (<i>Dymadon P</i>)	WR	PC
5196L	Paracetamol , tablet 500 mg (<i>Dymadon P</i>) (Dental)	WR	PC
8784H	Paracetamol , tablet 500 mg (<i>Dymadon P</i>) (Diff. Max. Qty)	WR	PC
5224Y	Paracetamol , tablet 500 mg (<i>Dymadon P</i>) (Diff. Max. Qty) (Dental)	WR	PC
3054R	Permethrin , cream 50 mg per g (5%), 30 g (<i>Lyclear</i>)	WR	PC
1910M	Polymyxin B Sulfate with Bacitracin and Neomycin Sulfate , eye ointment 5,000 units-400 units-5 mg per g, 4 g (<i>Neosporin</i>)	WR	PC
2724J	Pyridostigmine Bromide , tablet 10 mg (<i>Mestinon</i>)	ID	VT
1959D	Pyridostigmine Bromide , tablet 60 mg (<i>Mestinon</i>)	ID	VT
2608G	Pyridostigmine Bromide , tablet 180 mg (modified release) (<i>Mestinon Timespan</i>)	ID	VT
2878L	Sodium Cromoglycate , capsule containing powder for oral inhalation 20 mg (for use in Intal Spinhaler or Intal Halermatic) (<i>Intal Spincaps</i>)	AV	DP

SECTION 100 — HIGHLY SPECIALISED DRUGS PROGRAM ADDITIONS

Additions — Items

6458X **Abacavir Sulfate with Lamivudine**, tablet 600 mg (base)-300 mg (*Kivexa*)

DELETIONS

Deletions — Items

6159E **Ganciclovir**, capsule 250 mg (*Cymevene*)

ALTERATIONS

Restriction Changes

6423C **Lanreotide Acetate**, injection 60 mg (base) in single dose pre-filled syringe
(*Somatuline Autogel*)

6424D **Lanreotide Acetate**, injection 90 mg (base) in single dose pre-filled syringe
(*Somatuline Autogel*)

6425E **Lanreotide Acetate**, injection 120 mg (base) in single dose pre-filled syringe
(*Somatuline Autogel*)

Alterations — Proprietary Name

6101D **Clozapine**, tablet 25 mg

From: Clozapine Synthon

To: CloSyn

6102E **Clozapine**, tablet 100 mg

From: Clozapine Synthon

To: CloSyn

SECTION 100 — HUMAN GROWTH HORMONE PROGRAM ADDITIONS

Additions — Items

6465G **Somatropin**, solution for injection 5 mg (15 i.u.) in 1.5 mL cartridge (with preservative)
(*Norditropin NordiFlex*)

6466H **Somatropin**, solution for injection 10 mg (30 i.u.) in 1.5 mL cartridge (with preservative)
(*Norditropin NordiFlex*)

6467J **Somatropin**, solution for injection 15 mg (45 i.u.) in 1.5 mL cartridge (with preservative)
(*Norditropin NordiFlex*)

SECTION 100 — IVF/GIFT PROGRAM

Additions — Items

6464F **Follitropin Beta**, solution for injection 900 i.u. in 1.08 mL multi-dose cartridge
(*Puregon 900 IU/ 1.08 mL*)

Alterations — Notes

Notes have been amended in respect of the following:

Progesterone (effective 1 August 2005)

SECTION 100 — SPECIAL AUTHORITY PROGRAM ALTERATIONS

Restriction Changes

- 6444E **Imatinib Mesylate**, tablet 100 mg (base) (*Glivec*)
 6445F **Imatinib Mesylate**, tablet 400 mg (base) (*Glivec*)

Alterations — Notes

Notes have been amended in respect of the following:

- 6444E **Imatinib Mesylate**, tablet 100 mg (base) (*Glivec*)
 6445F **Imatinib Mesylate**, tablet 400 mg (base) (*Glivec*)

ADVANCE NOTICES

Advance Notices — Deletion of Items

The following items will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006:

Items discontinued by the manufacturer —

- 1319K **Diflunisal**, tablet 250 mg (*Dolobid*)
 5080J **Diflunisal**, tablet 250 mg (*Dolobid*) (**Dental**)
 1320L **Diflunisal**, tablet 500 mg (*Dolobid*)
 5081K **Diflunisal**, tablet 500 mg (*Dolobid*) (**Dental**)
 8539K **Oxaliplatin**, powder for I.V. infusion 50 mg (*Eloxatin*)
 8540L **Oxaliplatin**, powder for I.V. infusion 100 mg (*Eloxatin*)
 8068P **Venlafaxine Hydrochloride**, tablet 37.5 mg (base) (*Efexor*)
 8069Q **Venlafaxine Hydrochloride**, tablet 75 mg (base) (*Efexor*)

Deletion recommended by the Pharmaceutical Benefits Advisory Committee —

- 3483H **Pethidine Hydrochloride**, injection 100 mg in 2 mL (*MX*) (**Doctor's Bag**)

Advance Notices — Deletion of Brands

The following brands will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006:

Brands discontinued by the manufacturer —

- 2418G *Tryptanol*, MK — **Amitriptyline Hydrochloride**, tablet 25 mg
 3010K *Insensye*, FR — **Norfloracin**, tablet 400 mg

Addresses—Medicare Australia

Medicare Australia has responsibility for the operational aspects of the Pharmaceutical Benefits Scheme (PBS). This responsibility covers the processing of pharmaceutical benefit and safety net claims and authority applications together with the distribution of the Schedule of Pharmaceutical Benefits and PBS stationery used by medical practitioners, participating dental practitioners and approved pharmacists.

Requests for copies of the Schedule and/or notification of change of address should be directed to the Pharmaceutical Branch, Medicare Australia, GPO Box 9826, in your capital city.

Procedures for ordering prescription forms are set out in Section 1 (yellow pages) of this Schedule.

NEW SOUTH WALES and AUSTRALIAN CAPITAL TERRITORY

Pharmaceutical Benefits Branch
Colonial State Tower
150 George Street
Parramatta NSW 2150

General enquiries—
IME enquiries—

Tel: 132 290
Tel: 1300 302 122

Orange Service Centre
204-206 Lords Place
Orange NSW 2800

General enquiries—
IME enquiries—

Tel: 132 290
Tel: 1300 302 122

VICTORIA

Pharmaceutical Branch
Medibank House
460 Bourke Street
Melbourne Vic 3000

General enquiries—
IME enquiries—

Tel: 132 290
Tel: 1300 302 122

QUEENSLAND

Pharmaceutical Services Branch
444 Queen Street
Brisbane Qld 4000

General enquiries—
IME enquiries

Tel: 132 290
Tel: 1300 302 122

SOUTH AUSTRALIA and NORTHERN TERRITORY

Pharmaceutical Services Branch
209 Greenhill Road
Eastwood SA 5063

General enquiries—
IME enquiries—

Tel: 132 290
Tel: 1300 302 122

WESTERN AUSTRALIA

Pharmaceutical Benefits Branch
11th Floor, Bankwest Tower
108 St George's Terrace
Perth WA 6000

General enquiries—
IME enquiries—

Tel: 132 290
Tel: 1300 302 122

TASMANIA

Pharmaceutical Branch
242 Liverpool Street
Hobart Tas 7000

General enquiries—
IME enquiries—

Tel: 132 290
Tel: 1300 302 122

NATIONAL PROGRAM MANAGEMENT

Pharmaceutical Benefits Branch
Medicare Australia
134 Reed Street
Tuggeranong ACT 2900
Telephone— (02) 6124 6333

Website— www.medicareaustralia.gov.au
Email— pbs.enq@medicareaustralia.gov.au

Prior approval is required for all “Authority required” items or requests for increased quantities and/or repeats. Authorisation is obtained by lodging an application using the REPLY PAID mail service or by using Medicare Australia’s FREECALL telephone number:

REPLY PAID No. 9857
PBS Authorities Section
Medicare Australia
GPO Box 9857
In your capital city

Free call 1800 888 333
Australia-wide—24 hour service

Patient:	Medicare number
	Surname
	First name
	Full residential address (including postcode)

Your Prescriber Number: Located below your address block on the personalised forms

Drug Information:	PBS item
	Quantity required and number of repeats
	Daily dose
	Disease or purpose information

Requests for individual patient approval to obtain drugs that are available only through the SAS may be directed to a delegate within the Drug Safety and Evaluation Branch, Therapeutic Goods Administration, telephone (02) 6232 8111, facsimile (02) 6232 8112, or by mail to PO Box 100 Woden ACT 2606.

Details of addresses and telephone numbers of the approving authorities within the State offices of the Department of Veterans' Affairs are listed at the front of the Repatriation Schedule of Pharmaceutical Benefits.

A 24-hour, seven days a week telephone service is available by contacting 131 450.

The translating service (TIS) can provide immediate assistance over the telephone or arrange for an interpreter to go to a location specified in either city or country areas. The TIS service has access to 2000 professional interpreters, covering over 100 languages and dialects.

Poisons Information Centres

Phone 131 126 from anywhere in Australia — 24 hours — for information and advice on the treatment of poisoning, bites and stings.

NSW

The New Children's Hospital
Hawkesbury Road
Westmead NSW 2148
Tel: (02) 9845 3111

VIC

Royal Children's Hospital
Flemington Road
Parkville Vic 3052
Tel: (03) 9345 5680

QLD

Pharmacy Department
Royal Children's Hospital
Herston Qld 4029
Tel: see 24 hour contact above

WA

Sir Charles Gairdner Hospital
Hospital Avenue
Nedlands WA 6009
Tel: see 24 hour contact above

TAS

Royal Hobart Hospital
Liverpool Street
Hobart Tas 7000
Tel: see 24 hour contact above

NT

Royal Darwin Hospital
PO Box 41326
Casuarina NT 0811
Non-emergency contact number
Tel: (08) 8922 8424

ACT

Canberra Hospital
Yamba Drive
Garran ACT 2605
Tel: (02) 6285 2852

Drug Information Centres

NSW

Drug Information Pharmacist
New South Wales Medicines
Information Centre
PO Box 766
Darlinghurst NSW 2010
Tel: (02) 8382 2136

Drug Information Pharmacists
Hunter Drug Information Service
Newcastle Mater Misericordiae
Hospital
Locked Bag 7
Hunter Regional Mail Centre NSW
2310
Tel: (02) 4921 1278
(02) 4921 1328

VIC

Drug Information Pharmacist
Austin & Repatriation Medical
Centre
Studley Road
Heidelberg Vic 3084
Tel: (03) 9496 5668

Drug Information Pharmacist
Drug Information Centre
Southern Health Care Network
Monash Medical Centre
246 Clayton Road
Clayton Vic 3168
Tel: (03) 9594 2361

QLD

Assistant Director of Pharmacy
Queensland Drug Information Ctr
Royal Brisbane Hospital
E Floor, Block 7
Herston Road
Herston Qld 4029
Tel: (07) 3636 7098
(07) 3636 7599

SA

Drug Information Pharmacist
Royal Adelaide Hospital
North Terrace
Adelaide SA 5000
Tel: (08) 8222 5546

Drug Information Pharmacist
Flinders Medical Centre
Bedford Park SA 5042
Tel: (08) 8204 5301

Drug Information Pharmacist
Queen Elizabeth Hospital
Woodville Road
Woodville SA 5011
Tel: (08) 8222 6777

WA

Drug Information Pharmacist
Sir Charles Gairdner Hospital
Hospital Avenue
Nedlands WA 6009
Tel: (08) 9346 2923

TAS

Drug Information Pharmacist
Royal Hobart Hospital
GPO Box 1061L
Hobart Tas 7001
Tel: (03) 6222 8737

NT

Drug Information Pharmacist
Royal Darwin Hospital
PO Box 41326
Casuarina NT 0811
Tel: (08) 8922 8424

ACT

Drug Information Pharmacist
Canberra Hospital
Yamba Drive
Garran ACT 2605
Tel: (02) 6244 3333

National Prescribing Service (NPS)

Therapeutic Advice and Information
Service (TAIS)
Level 7, 418A Elizabeth Street
Surry Hills NSW 2010
Tel: 1300 138 677
Fax: (03) 9459 4546
Email: tais@nps.org.au
Web: www.nps.org.au

List of Contact Officers for Recalls of Therapeutic Goods

For details of consumer level recalls only — telephone 1800 020 512

These officers may be contacted—

- to obtain information about current recalls
- to report suspected problems relating to the quality, safety or efficacy of a therapeutic good

Australian Recall Coordinator

Office of Devices, Blood and Tissues
Therapeutic Goods Administration
Department of Health and Ageing
PO Box 100
Woden ACT 2606

Mr P K Harrison (02) 6232 8636
Mr T Byrne (02) 6232 8637

Australian Capital Territory

ACT Health
GPO Box 825
Canberra ACT 2601

Ms J Strang (02) 6205 0961

New South Wales

Department of Health, NSW
PO Box 103
Gladesville NSW 1675

Mr J E Lumby (02) 9879 3214

Victoria

Department of Human Services
Drugs and Poisons Unit
GPO Box 1670N
Melbourne Vic 3001

Mr K Moyle 1300 364 545
Mr R Bell 1300 364 545

Queensland

Queensland Department of Health
GPO Box 48
Brisbane Qld 4001

Drugs—

Mr A Hawkins (07) 3234 0349
Mr C Healey (07) 3234 0960

Therapeutic Devices—

Mr C Healey (07) 3234 0960
Mr D Jones (07) 3406 8068

South Australia

Drug Policies and Programs
Metropolitan Health Division
South Australian Department of Health
PO Box 287
Rundle Mall SA 5000

Mr W Dollman (08) 8226 7110
Ms E Anear (08) 8226 7387

Western Australia

Health Department of WA
PO Box 8172, Perth Business Centre
Perth WA 6849

Mr M Patterson (08) 9388 4980

Tasmania

Department of Health and Human Services
GPO Box 125B
Hobart Tas 7001

Drugs—

Mr J Galloway (03) 6233 2064
Ms M Sharpe (03) 6233 3766

Therapeutic Devices—

Mr A L Wilkins (03) 6233 3913

Northern Territory

Department of Health and Community Services
PO Box 40596
Casuarina NT 0811

Ms H Stone (08) 8922 7035

Index of Manufacturers' Codes

<i>Code</i>	<i>Manufacturer</i>	<i>Code</i>	<i>Manufacturer</i>
AB	Abbott Australasia Pty Ltd Captain Cook Drive Kurnell NSW 2231 Tel: (02) 9668 9711 Fax: (02) 9668 8459	AP	AstraZeneca Pty Ltd Alma Road North Ryde NSW 2113 Tel: (02) 9978 3500 Fax: (02) 9978 3700
AC	Alberto Culver Company 14 Loyalty Road North Rocks NSW 2151 Tel: (02) 9630 5099 Fax: (02) 9683 5026	AQ	Alcon Laboratories (Australia) Pty Ltd Allambie Grove Park 25 Frenchs Forest Road East Frenchs Forest NSW 2086 Tel: 1800 025 004 Fax: (02) 9452 5209
AD	Amrad Pharmaceuticals Pty Ltd 54-68 Ferndell Street South Granville NSW 2142 Tel: (02) 9795 9500 Fax: (02) 9795 9595	AS	Aspen Pharmacare Australia Pty Ltd First Floor 34-36 Chandos Street St Leonards NSW 2065 Tel: (02) 8436 8300 Fax: (02) 9901 3540
AE	AFT Pharmaceuticals Pty Limited 9 Water Street Caringbah NSW 2229 Tel: 1800 097 639 Fax: 1800 097 810	AT	Actelion Pharmaceuticals Australia Pty Ltd Level 2 West, Suites 48-50 7 Narabang Way Belrose NSW 2085 Tel: (02) 9486 4600 Fax: (02) 9986 1344
AF	Alphapharm Pty Limited Chase Building 2 Wentworth Park Road Glebe NSW 2037 Tel: (02) 9298 3999 Fax: (02) 9566 4686	AV	Aventis Pharma Pty Limited 27 Sirius Road Lane Cove NSW 2066 Tel: 1800 555 995 Fax: 1800 555 335
AG	Allergan Australia Pty Ltd 77 Ridge Street Gordon NSW 2072 Tel: 1800 252 224 Fax: (02) 9498 0290	AW	Arrow Pharmaceuticals Limited 24 Rothschild Avenue Rosebery NSW 2018 Tel: (02) 8344 8344 Fax: (02) 8344 8355
AH	Altana Pharma Pty Ltd Ground Floor, 2 Lyonpark Road North Ryde NSW 2113 Tel: (02) 9859 6900 Fax: (02) 9859 6950	AX	Aventis Pasteur Pty Limited Unit 12, 12-14 Columbia Way Baulkham Hills NSW 2153 Tel: (02) 8852 4200 Fax: (02) 8852 4299
AL	Alphapharm Medical A Division of Alphapharm Pty Limited Chase Building 2 Wentworth Park Road Glebe NSW 2037 Tel: (02) 9298 3999 Fax: (02) 9566 4686	BB	Blackmores Ltd 23 Roseberry Street Balgowlah NSW 2093 Tel: (02) 9951 0111 Fax: (02) 9949 1954
AN	Amgen Australia Pty Ltd Level 7, 123 Epping Road North Ryde NSW 2113 Tel: (02) 9870 1333 Fax: (02) 9870 1344	BC	Bristol Laboratories A Division of Bristol-Myers Squibb Australia Pty Ltd 556 Princes Highway Noble Park Vic 3174 Tel: (03) 9213 4000 Fax: (03) 9701 1518

<i>Code</i>	<i>Manufacturer</i>
BD	Biogen Idec Australia Pty Ltd Suite 2601, Level 26 100 Miller Street North Sydney NSW 2060 Tel: (02) 8907 3300 Fax: (02) 8907 3332
BE	BDF Australia Ltd 112-118 Talavera Road North Ryde NSW 2113 Tel: 1800 269 933 Fax: (02) 9887 3487
BF	Bellwether Pharma Limited Suite 2, Level 2 71 Epping Road North Ryde NSW 2113 Tel: (02) 8875 5700 Fax: (02) 9889 2250
BG	Biochemie Australia A Division of Sandoz Pty Ltd 54 Waterloo Road North Ryde NSW 2113 Tel: (02) 9888 8550 Fax: (02) 9888 8557
BI	Biotech Pharmaceuticals Pty Ltd 100 Antimony Street Carole Park Qld 4300 Tel: (07) 3271 9600 Fax: (07) 3271 1315
BK	Becton Dickinson Pty Ltd 80 Rushdale Street Knoxfield Vic 3180 Tel: (03) 9764 2444 Fax: (03) 9764 2550
BN	Bayer Australia Limited 875-893 Pacific Highway Pymble NSW 2073 Tel: (02) 9391 6000 Fax: (02) 9988 3311
BP	British Pharmaceuticals Unit A, 31-33 Sirius Road Lane Cove NSW 2066 Tel: (02) 9428 9411 Fax: (02) 9428 1732
BQ	Bristol-Myers Squibb Pharmaceuticals A Division of Bristol-Myers Squibb Australia Pty Ltd 556 Princes Highway Noble Park Vic 3174 Tel: (03) 9213 4000 Fax: (03) 9701 1518

<i>Code</i>	<i>Manufacturer</i>
BR	B. Braun Australia Pty Ltd Norwest Business Park 17 Lexington Drive Bella Vista NSW 2153 Tel: (02) 9629 0200 Fax: (02) 9629 0299
BU	Bausch & Lomb Surgical A Division of Bausch & Lomb (Australia) Pty Ltd Level 4, 113 Wicks Road North Ryde NSW 2113 Tel: (02) 9887 1444 Fax: (02) 9888 9642
BV	B.S.N. 315 Ferntree Gully Road Mount Waverley Vic 3149 Tel: (03) 8540 6777 Fax: 1800 671 000
BX	Baxter Healthcare Pty Limited 1 Baxter Drive Old Toongabbie NSW 2146 Tel: (02) 9848 1111 Fax: (02) 9848 1123
BY	Boehringer Ingelheim Pty Limited 85 Waterloo Road North Ryde NSW 2113 Tel: (02) 8875 8800 Fax: (02) 8875 8801
CC	ConvaTec A Division of Bristol-Myers Squibb Australia Pty Ltd 606 Hawthorn Road East Brighton Vic 3187 Tel: 1800 335 276 Fax: (03) 9525 0920
CH	Chem mart Pty Limited 115 Sherrieff Street Underdale SA 5032 Tel: (08) 8408 3200 Fax: (08) 8408 3383
CO	Chemists' Own Pty Ltd A member of Sigma Group of Companies 96 Merrindale Drive Croydon Vic 3136 Tel: (03) 9839 2800 Fax: (03) 9839 2801
CS	CSL Limited 45 Poplar Road Parkville Vic 3052 Tel: (03) 9389 1911 Fax: (03) 9388 2351

<i>Code</i>	<i>Manufacturer</i>
CT	Coloplast Pty Ltd 33 Gilby Road Mount Waverley Vic 3149 Tel: 1800 673 317 Fax: (03) 9541 1199
CV	Ciba Vision Australia Pty Ltd Unit 1/42 Carrington Road Castle Hill NSW 2154 Tel: (02) 9680 6655 Fax: (02) 9899 9640
CX	Contact Lens Centre Australia Ltd Unit 1, 5 Seismic Court Rowville Vic 3178 Tel: (03) 9753 3177 Fax: (03) 9753 2399
DG	Douglas Medication Systems A Division of Douglas Pharmaceuticals Australia Ltd 3/10 Inglewood Place Norwest Business Park Baulkham Hills NSW 2153 Tel: (02) 8818 2111 Fax: (02) 8818 2122
DK	Dakota Pharmaceuticals A Division of Sanofi-Synthelabo Australia Pty Limited 16 Byfield Street North Ryde NSW 2113 Tel: (02) 8899 0700 Fax: (02) 8899 0600
DP	Douglas Pharmaceuticals Australia Ltd 3/10 Inglewood Place Norwest Business Park Baulkham Hills NSW 2153 Tel: (02) 8818 2111 Fax: (02) 8818 2122
DR	DiaCare International Pty Ltd Unit 30, 10 Gladstone Road Castle Hill NSW 2154 Tel: (02) 8875 7909 Fax: (02) 8875 7777
DT	DermaTech Laboratories Pty Ltd Unit 17, 167 Prospect Highway Seven Hills NSW 2147 Tel: (02) 9624 5874 Fax: (02) 9624 8822
EG	Eagle Pharmaceuticals Pty Ltd 4/40 Carrington Road Castle Hill NSW 2154 Tel: (02) 9899 9099 Fax: (02) 9899 6564

<i>Code</i>	<i>Manufacturer</i>
EO	Ego Pharmaceuticals Pty Ltd 21-31 Malcolm Road Braeside Vic 3195 Tel: (03) 9587 1088 Fax: (03) 9580 7647
EP	EpiPharm Pty Ltd Level 13, 1 Collins Street Melbourne Vic 3000 Tel: 1800 028 880 Fax: (03) 9660 4999
EX	Essex Laboratories 11 Gibbon Road Baulkham Hills NSW 2153 Tel: (02) 9852 7444 Fax: (02) 9852 7500
FA	F.H. Faulding & Co. Limited Level 6, 390 St Kilda Road Melbourne Vic 3004 Tel: (03) 9868 0700 Fax: (03) 9868 0111
FB	Pierre Fabre Medicament Australia Pty Limited Unit 26B, Parkview Business Centre 1 Maitland Place Baulkham Hills NSW 2153 Tel: (02) 8858 2800 Fax: (02) 8858 2888
FH	Faulding Healthcare Pty Ltd 115 Sherriff Street Underdale SA 5032 Tel: (08) 8408 3200 Fax: (08) 8408 3383
FL	C. B. Fleet Co. (Aust.) Pty Ltd 25 Macbeth Street Braeside Vic 3195 Tel: (03) 9580 2755 Fax: (03) 9580 2899
FM	Fawns and McAllan Pty Ltd A member of Sigma Group of Companies 96 Merrindale Drive Croydon Vic 3136 Tel: (03) 9839 2800 Fax: (03) 9839 2801
FP	Ferring Pharmaceuticals Pty Ltd Suite 2B, Level 2 802 Pacific Highway Gordon NSW 2072 Tel: (02) 9497 2300 Fax: (02) 9497 2399

<i>Code</i>	<i>Manufacturer</i>
FR	Charles E. Frosst Division of Merck Sharp & Dohme (Australia) Pty Ltd 54-68 Ferndell Street South Granville NSW 2142 Tel: (02) 9795 9500 Fax: (02) 9795 9595
GA	Galderma Australia Pty Ltd 9 Rodborough Road Frenchs Forest NSW 2086 Tel: 1800 800 765 Fax: (02) 9975 5374
GC	GlaxoSmithKline Consumer Healthcare 82 Hughes Avenue Ermington NSW 2115 Tel: (02) 9684 0888 Fax: (02) 9684 6958
GD	General Diabetes Services Rear, 137 Mt Alexander Road Flemington Vic 3031 Tel: (03) 9372 2389 Fax: (03) 9376 4158
GI	Gilead Sciences Pty Ltd Unit 2, 41 Stamford Road Oakleigh Vic 3166 Tel: (03) 9563 0433 Fax: (03) 9563 0170
GK	GlaxoSmithKline Australia Pty Ltd 1061 Mountain Highway Boronia Vic 3155 Tel: (03) 9721 6000 Fax: (03) 9729 5319
GP	GP Laboratories A Division of Pfizer Pty Limited 38-42 Wharf Road West Ryde NSW 2114 Tel: (02) 9850 3333 Fax: (02) 9858 1347
GX	GenRx Pty Ltd Suite 2B, Level 3, Building A 11 Talavera Road North Ryde NSW 2113 Tel: 1800 195 055 Fax: 1800 133 300
HA	Hamilton Laboratories Pty Ltd 217 Flinders Street Adelaide SA 5000 Tel: (08) 8223 2957 Fax: (08) 8232 1480

<i>Code</i>	<i>Manufacturer</i>
HO	Hollister Incorporated, U.S.A. Australian Distributor Liberty Medical Pty Ltd 9 Central Boulevard Port Melbourne Vic 3207 Tel: (03) 9646 4033 Fax: (03) 9646 4018
HP	Hoechst Division of Aventis Pharma Pty Limited 27 Sirius Road Lane Cove NSW 2066 Tel: 1800 555 995 Fax: 1800 555 335
HR	Paul Hartmann Pty Ltd 27-28/11-21 Underwood Road Homebush NSW 2140 Tel: 1800 805 839 Fax: (02) 8762 7100
HS	healthsense 115 Sherriff Street Underdale SA 5032 Tel: (08) 8408 3200 Fax: (08) 8408 3383
HX	Hexal Australia Pty Ltd Level 4, Suite 1-6 100 Harris Street Pyrmont NSW 2009 Tel: (02) 9566 1500 Fax: (02) 9566 1458
IQ	loquin A Division of Alcon Laboratories (Australia) Pty Ltd Allambie Grove Park 25 Frenchs Forest Road East Frenchs Forest NSW 2086 Tel: 1800 025 004 Fax: (02) 9452 5209
IS	Ipsen Pty Limited 21 Aristoc Road Glen Waverley Vic 3150 Tel: (03) 9550 1843 Fax: (03) 9562 5152
IT	InterPharma Pty Ltd Suite 3, 14 Sydney Road Manly NSW 2095 Tel: (02) 9976 6876 Fax: (02) 9976 6859
JC	Janssen-Cilag Pty Ltd 1-5 Khartoum Road North Ryde NSW 2113 Tel: (02) 8875 3333 Fax: (02) 8875 3300

<i>Code</i>	<i>Manufacturer</i>
JJ	Johnson & Johnson Medical 1-5 Khartoum Road North Ryde NSW 2113 Tel: (02) 9878 9111 Fax: 1800 808 233
JT	Johnson & Johnson Pacific Pty Limited 45 Jones Street Ultimo NSW 2007 Tel: 13 1565 Fax: (02) 8260 8102
KC	Kimberly-Clark Australia Pty Ltd 52 Alfred Street South Milsons Point NSW 2061 Tel: (02) 9963 8888 Fax: (02) 9957 5687
KE	Kendall Australasia Pty Ltd 22 Giffnock Avenue North Ryde NSW 2113 Tel: 1800 252 467 Fax: (02) 9888 7378
KN	Knoll A Division of Abbott Australasia Pty Ltd Captain Cook Drive Kurnell NSW 2231 Tel: (02) 9668 9711 Fax: (02) 9668 8459
KR	Kenral Division of Pharmacia Australia Pty Limited 59 Kirby Street Rydalmere NSW 2116 Tel: (02) 9848 3000 Fax: (02) 9848 3333
KY	Key Pharmaceuticals Pty Ltd 9-11 Leeds Street Rhodes NSW 2138 Tel: (02) 9736 3811 Fax: (02) 9736 3316
LB	Life Bioscience Pty Ltd 80 Millewa Avenue Malvern East Vic 3145 Tel: 1800 114 610 Fax: (03) 8660 2785
LF	Laboratoires Fournier S.A. Level 4, 35 Spring Street Bondi Junction NSW 2022 Tel: (02) 9389 8611 Fax: (02) 9389 8615
LM	Link Medical Products Pty Ltd Level 1, Bridgepoint Centre 3 Brady Street Mosman NSW 2088 Tel: (02) 9960 0150 Fax: (02) 9960 0149

<i>Code</i>	<i>Manufacturer</i>
LN	Lennon Healthcare A Division of Aspen Pharmacare Australia Pty Ltd First Floor 34-36 Chandos Street St Leonards NSW 2065 Tel: (02) 8436 8300 Fax: (02) 9901 3540
LU	Lundbeck Australia Pty Ltd Unit 1, 10 Inglewood Place Norwest Business Park Baulkham Hills NSW 2153 Tel: (02) 9836 1655 Fax: (02) 9836 1755
LY	Eli Lilly Australia Pty Limited 112 Wharf Road West Ryde NSW 2114 Tel: (02) 9325 4444 Fax: (02) 9325 4410
MD	Macarthur Research Division of Roche Products Pty Ltd 4-10 Inman Road Dee Why NSW 2099 Tel: (02) 9454 9000 Fax: (02) 9981 3229
ME	Menley & James Division of GlaxoSmithKline Australia Pty Ltd 1061 Mountain Highway Boronia Vic 3155 Tel: (03) 9721 6000 Fax: (03) 9729 5319
MF	Mundipharma Pty Ltd Level 26, 6 O'Connell Street Sydney NSW 2000 Tel: (02) 9231 7200 Fax: (02) 9223 0011
MG	McGloins Classic Brands Unit 5, 13 Hoyle Avenue Castle Hill NSW 2154 Tel: (02) 9894 4844 Fax: (02) 9894 4393
MH	Molnlycke Health Care Pty Ltd Building 1, Ground Floor 14 Aquatic Drive Frenchs Forest NSW 2086 Tel: (02) 9453 1144 Fax: (02) 9453 1155
MK	Merck Sharp & Dohme (Australia) Pty Ltd 54-68 Ferndell Street South Granville NSW 2142 Tel: (02) 9795 9500 Fax: (02) 9795 9595

<i>Code</i>	<i>Manufacturer</i>
ML	Marion Division of Aventis Pharma Pty Limited 27 Sirius Road Lane Cove NSW 2066 Tel: 1800 555 995 Fax: 1800 555 335
MM	3M Pharmaceuticals Australia Pty Ltd 9-15 Chilvers Road Thornleigh NSW 2120 Tel: (02) 9875 6333 Fax: (02) 9875 6416
MQ	Merck Pharmaceuticals Chase Building 2 Wentworth Park Road Glebe NSW 2037 Tel: (02) 9298 3999 Fax: (02) 9566 4686
MS	Abbott Diagnostics Division (MediSense Products) 666 Doncaster Road Doncaster Vic 3108 Tel: (03) 9843 7100 Fax: (03) 9855 8020
MT	Mentholatum Australasia Pty Ltd 12-16 Janine Street Scoresby Vic 3179 Tel: (03) 9763 0322 Fax: (03) 9763 2699
MW	McGaw Biomed Australia Pty Ltd c/- B. Braun Australia Pty Ltd Norwest Business Park 17 Lexington Drive Bella Vista NSW 2153 Tel: (02) 9629 0200 Fax: (02) 9629 0299
MX	Mayne Pharma Pty Ltd (David Bull Laboratories, Faulding Pharmaceuticals) Level 6, 390 St Kilda Road Melbourne Vic 3004 Tel: (03) 9868 0700 Fax: (03) 9868 0111
NA	National Diagnostic Products (Aust) Pty Limited 7-9 Merriwa Street Gordon NSW 2072 Tel: (02) 9418 1100 Fax: (02) 9418 1181
NC	Novartis Consumer Health Australasia Pty Ltd 327-333 Police Road Mulgrave Vic 3170 Tel: (03) 9701 2711 Fax: (03) 9701 2911

<i>Code</i>	<i>Manufacturer</i>
NE	Norgine Pty Limited 3/14 Rodborough Road Frenchs Forest NSW 2086 Tel: (02) 9972 7500 Fax: (02) 9972 7522
NF	FlexPen Products of Novo Nordisk Pharmaceuticals Pty Ltd Level 3, 21 Solent Circuit Baulkham Hills NSW 2153 Tel: (02) 8858 3600 Fax: (02) 8858 3799
NI	InnoLet Products of Novo Nordisk Pharmaceuticals Pty Ltd Level 3, 21 Solent Circuit Baulkham Hills NSW 2153 Tel: (02) 8858 3600 Fax: (02) 8858 3799
NL	NovoLet Products of Novo Nordisk Pharmaceuticals Pty Ltd Level 3, 21 Solent Circuit Baulkham Hills NSW 2153 Tel: (02) 8858 3600 Fax: (02) 8858 3799
NO	Novo Nordisk Pharmaceuticals Pty Ltd Level 3, 21 Solent Circuit Baulkham Hills NSW 2153 Tel: (02) 8858 3600 Fax: (02) 8858 3799
NT	Nestlé Australia Ltd 60 Bathurst Street Sydney NSW 2000 Tel: (02) 9931 2345 Fax: (02) 9931 2610
NU	Nutricia Australia Pty Limited Norwest Business Park 14-16 Brookhollow Avenue Baulkham Hills NSW 2153 Tel: (02) 8853 9600 Fax: (02) 9894 6498
NV	Novartis Pharmaceuticals Australia Pty Ltd 54 Waterloo Road North Ryde NSW 2113 Tel: (02) 9805 3555 Fax: (02) 9887 4551
OA	Orphan Australia Pty Ltd 48 Kangan Drive Berwick Vic 3806 Tel: (03) 9769 5744 Fax: (03) 9769 5944

<i>Code</i>	<i>Manufacturer</i>
OB	Oral B Laboratories Pty Ltd Level 3, 90 Mount Street North Sydney NSW 2060 Tel: (02) 9957 6499 Fax: (02) 9957 5383
OL	Owen Laboratories Division of Galderma Australia Pty Ltd 9 Rodborough Road Frenchs Forest NSW 2086 Tel: 1800 800 765 Fax: (02) 9975 5374
OM	Colgate Oral Care 345 George Street Sydney NSW 2000 Tel: (02) 9229 5600 Fax: (02) 9232 8448
OR	Organon (Australia) Pty Limited 31-33 Sirius Road Lane Cove NSW 2066 Tel: (02) 9428 9411 Fax: (02) 9428 1732
PC	Pfizer Consumer Healthcare Pty Ltd 32 Cawarra Road Caringbah NSW 2229 Tel: (02) 9710 6500 Fax: (02) 9710 6644
PD	Parke Davis Pty Ltd 32 Cawarra Road Caringbah NSW 2229 Tel: (02) 9710 6500 Fax: (02) 9710 6400
PE	Pacific EyeCare A Division of Allergan Australia Pty Ltd 77 Ridge Street Gordon NSW 2072 Tel: 1800 252 224 Fax: (02) 9498 0290
PF	Pfizer Pty Limited 38-42 Wharf Road West Ryde NSW 2114 Tel: (02) 9850 3333 Fax: (02) 9858 1347
PH	Pharmacia Australia Pty Limited 59 Kirby Street Rydalmere NSW 2116 Tel: (02) 9848 3000 Fax: (02) 9848 3333
PK	Pharmatel Fresenius Kabi Pty Ltd Unit 6, 6-18 Bridge Road Hornsby NSW 2077 Tel: (02) 9472 2222 Fax: (02) 9472 2255

<i>Code</i>	<i>Manufacturer</i>
PL	Pharmalab 332 Burns Bay Road Lane Cove NSW 2066 Tel: (02) 9420 9199 Fax: (02) 9420 9177
PM	PMC Pharma A Division of AstraZeneca Pty Ltd Alma Road North Ryde NSW 2113 Tel: (02) 9978 3500 Fax: (02) 9978 3700
PP	Petrus Pharmaceuticals Pty Ltd Level 3, IBM Building 1060 Hay Street West Perth WA 6005 Tel: (08) 9368 5954 Fax: (08) 9368 6692
PU	Pharmacia & Upjohn Pty Limited 59 Kirby Street Rydalmere NSW 2116 Tel: (02) 9848 3000 Fax: (02) 9848 3333
PX	Point of Care Diagnostics Australia Pty Ltd Unit 14, 76 Reserve Road Artarmon NSW 2064 Tel: (02) 9437 1355 Fax: (02) 9437 1399
PY	Procter & Gamble Pharmaceuticals Australia Pty Ltd 99 Phillip Street Parramatta NSW 2150 Tel: (02) 9685 4500 Fax: (02) 9685 4777
QM	Qualimed Division of Aventis Pharma Pty Limited 27 Sirius Road Lane Cove NSW 2066 Tel: 1800 555 995 Fax: 1800 555 335
RC	Reckitt Benckiser (Australia) Pty Limited 44 Wharf Road West Ryde NSW 2114 Tel: (02) 9857 2000 Fax: (02) 9857 2004
RD	Roche Diagnostics Australia Pty Ltd 31 Victoria Avenue Castle Hill NSW 2154 Tel: (02) 9899 7999 Fax: (02) 9634 4696

<i>Code</i>	<i>Manufacturer</i>
RE	Real-RL Division of GlaxoSmithKline Australia Pty Ltd 1061 Mountain Highway Boronia Vic 3155 Tel: (03) 9721 6000 Fax: (03) 9721 5319
RO	Roche Products Pty Ltd 4-10 Inman Road Dee Why NSW 2099 Tel: (02) 9454 9000 Fax: (02) 9981 3229
SA	SciGen Pty Limited Level 7, 2 Bligh Street Sydney NSW 2000 Tel: (02) 9234 1700 Fax: (02) 9234 1777
SB	Scientific Hospital Supplies Australia Products Norwest Business Park 14-16 Brookhollow Avenue Baulkham Hills NSW 2153 Tel: (02) 8853 9600 Fax: (02) 9894 6498
SC	Schering Pty Ltd Australian Subsidiary of Schering AG, Berlin 27-31 Doody Street Alexandria NSW 2015 Tel: (02) 9317 8666 Fax: (02) 9317 2138
SE	Servier Laboratories (Aust.) Pty Ltd 8 Cato Street Hawthorn Vic 3122 Tel: (03) 8823 7333 Fax: (03) 9822 9790
SG	Serono Australia Pty Ltd Unit 3-4, 25 Frenchs Forest Road East Frenchs Forest NSW 2086 Tel: (02) 8977 4100 Fax: (02) 9975 1516
SH	Schering-Plough Pty Ltd 11 Gibbon Road Baulkham Hills NSW 2153 Tel: (02) 9852 7444 Fax: (02) 9852 7500
SI	Sigma Pharmaceuticals Pty Ltd 96 Merrindale Drive Croydon Vic 3136 Tel: (03) 9839 2800 Fax: (03) 9839 2801

<i>Code</i>	<i>Manufacturer</i>
SJ	Sharpe Laboratories Pty Ltd 12 Hope Street Ermington NSW 2115 Tel: (02) 9858 5622 Fax: (02) 9858 5957
SL	SBPA A Division of Sandoz Pty Ltd 54 Waterloo Road North Ryde NSW 2113 Tel: (02) 9888 8550 Fax: (02) 9888 8557
SM	Solvay Pharmaceuticals Division of Solvay Biosciences Pty Ltd Level 1, Building 2 Pymble Corporate Centre 20 Bridge Street Pymble NSW 2073 Tel: (02) 9440 0977 Fax: (02) 9440 0910
SN	Smith & Nephew Healthcare 315 Ferntree Gully Road Mount Waverley Vic 3149 Tel: (03) 8540 6777 Fax: 1800 671 000
SS	SSL Australia Pty Ltd 225 Beach Road Mordialloc Vic 3195 Tel: 1800 999 155 Fax: (03) 9587 6870
SU	Sauter Laboratories (Aust.) Pty Ltd 4-10 Inman Road Dee Why NSW 2099 Tel: (02) 9454 9000 Fax: (02) 9981 3229
SW	Sanofi-Synthelabo Australia Pty Limited 16 Byfield Street North Ryde NSW 2113 Tel: (02) 8899 0700 Fax: (02) 8899 0600
SX	Stiefel Laboratories Pty Limited Unit 14, 5 Salisbury Road Castle Hill NSW 2154 Tel: (02) 9894 5088 Fax: (02) 9894 5016
SY	Schering AG 27-31 Doody Street Alexandria NSW 2015 Tel: (02) 9317 8666 Fax: (02) 9317 2138
SZ	Sandoz Pty Ltd 54 Waterloo Road North Ryde NSW 2113 Tel: (02) 9888 8550 Fax: (02) 9888 8557

<i>Code</i>	<i>Manufacturer</i>
TC	Technostic Consulting Pty Ltd 6735 Cutrock Road Lisarow NSW 2250 Tel: 0417 210 398 Fax: (02) 4362 7077
TH	Richard Thomson Pty Ltd Unit 2, 109 Vanessa Street Kingsgrove NSW 2208 Tel: (02) 9502 8400 Fax: (02) 9502 8401
TM	Technipro Marketing Pty Ltd 13 Bourke Street North Parramatta NSW 2151 Tel: (02) 9890 9311 Fax: (02) 9890 7488
TW	Terry White Chemists 115 Sherriff Street Underdale SA 5032 Tel: (08) 8408 3200 Fax: (08) 8408 3383
UC	UCB Pharma A Division of UCB Australia Pty Ltd Level 1, 1155 Malvern Road Malvern Vic 3144 Tel: (03) 9828 1800 Fax: (03) 9828 1860
UM	Unomedical Pty Ltd 11-17 Wilmette Place Mona Vale NSW 2103 Tel: (02) 9997 8033 Fax: (02) 9997 3760
UW	United Works of Pharmaceutical and Dietetic Products, Hungary Australian Distributor Boucher & Muir Pty Limited Tel: (02) 9436 3922 Fax: (02) 9906 7147
VF	Vitaflo Australia Pty Ltd 23-25 Pinto Way Highton Vic 3216 Tel: (03) 5244 5811 Fax: (03) 5244 5822
VT	Valeant Pharmaceuticals Australasia Pty Ltd Level 1, 85 St Hilliers Road Auburn NSW 2144 Tel: (02) 9648 4266 Fax: (02) 9648 4655
WT	Wyeth Consumer Healthcare Pty Ltd 17-19 Solent Circuit Norwest Business Park Baulkham Hills NSW 2153 Tel: 1800 555 057 Fax: (02) 9023 0016

<i>Code</i>	<i>Manufacturer</i>
WW	Wm R. Warner 32 Cawarra Road Caringbah NSW 2229 Tel: (02) 9710 6500 Fax: (02) 9710 6644
WX	Wyeth Australia Pty Limited 17-19 Solent Circuit Norwest Business Park Baulkham Hills NSW 2153 Tel: (02) 9761 8200 Fax: (02) 9023 0000
WY	Wyeth Pharmaceuticals Division of Wyeth Australia Pty Limited 17-19 Solent Circuit Norwest Business Park Baulkham Hills NSW 2153 Tel: (02) 9761 8200 Fax: (02) 9023 0000
ZT	Synthon AU Pty Ltd Level 28, 140 St Georges Terrace Perth WA 6000 Tel: (08) 9409 5200 Fax: (08) 9409 5211

Section 1 — Explanatory Notes

Introduction

These Explanatory Notes are provided to help doctors, dentists and pharmacists work within the Australian Government's Pharmaceutical Benefits Scheme (PBS).

The PBS is a system of subsidising the cost of most prescription medicines. The subsidies are available to all Australian residents and eligible foreign visitors, i.e., people from countries which have Reciprocal Health Care Agreements with Australia. These countries are the United Kingdom, Ireland, New Zealand, Malta, Italy, Sweden, the Netherlands, Finland, and Norway.

The aim of the PBS, which has been in operation since 1948, is to provide reliable and affordable access to a wide range of necessary medicines.

The Schedule of Pharmaceutical Benefits – referred to throughout as the 'Schedule' – lists all of the medicines available under the PBS, and explains how they can be used in order to be subsidised.

The Schedule is produced three times each year by the Australian Department of Health and Ageing (effective 1 April, 1 August and 1 December).

It is vital therefore that doctors, dentists and pharmacists remain up to date with information on which medicines are included in or excluded from the Schedule, whether restrictions apply to the medicines, and how much patients should pay.

Queries relating to the PBS can be made to the Pharmaceutical Branches of Medicare Australia (telephone 132 290 Mondays to Saturdays, during business hours). Queries relating to the Repatriation Pharmaceutical Benefits Scheme (RPBS) can be made to the State offices of the Department of Veterans' Affairs (DVA) (telephone 1800 552 580).

1. The Schedule — Where to Find What

The Schedule of Pharmaceutical Benefits is divided into sections, each with particular page colours for easy reference. At the start of the Schedule, immediately after the table of contents, is a summary of any changes to listed items. This is followed by a list of important information sources, contacts and addresses, then an index of manufacturers' codes.

The last pages of the Schedule provide a generic/proprietary index of PBS and RPBS ready-prepared items.

Section 1 (yellow pages)

Section 1 is what you are reading, the Explanatory Notes. It outlines the correct way to prescribe and supply pharmaceutical benefits; patient charges; who qualifies for concessions; how the Safety Net system works; and, for pharmacists, how to claim reimbursement for PBS items.

Please note that except where indicated, the term '**prescriber**' is used in this section to cover both doctors and dentists who work within the PBS.

And except where stated otherwise, the term '**pharmacist**' means a pharmacist approved to supply medicines under the PBS.

Section 2 (white, lavender, pink and orange pages)

This section lists ready-prepared items, and includes the form, manner of administration, brand and brand equivalents which may be prescribed, and the maximum quantity and number of repeats for each item.

Emergency drug (doctor's bag) supplies are also listed at the beginning of this section.

Any medicines that have restrictions on how they can be prescribed are printed in ***bold italics***. Items appearing in more than one therapeutic group are cross-referenced.

The second page of Section 2 explains symbols used throughout the Schedule.

The use of 'NOTE' in this section is used to clarify how some pharmaceutical benefits should be prescribed.

The use of 'CAUTION' is to warn of known adverse reactions from, or precautions to be taken with, a particular pharmaceutical benefit. (The absence of a cautionary note does not imply reactions may not happen.)

A separate list at the end of Section 2 relates to items that can be prescribed by dentists who work within the PBS (pink pages). This is followed by a list of items that are made available under special arrangements for doctors to prescribe (orange pages).

Section 3 (blue pages)

This section lists container prices, fees related to dispensing, standard packs and prices for ready-prepared preparations.

Section 4 (green pages)

This section deals with extemporaneous preparations. It lists the ingredients which can be used, a table of maximum quantities and number of repeats, container prices, and a list of standard formula preparations and prices (based on formularies in common use and referred to in the Schedule as the Standard Formulae List).

Restrictions applying to the use of a pharmaceutical benefit are indicated against the item.

Repatriation Schedule of Pharmaceutical Benefits (white and buff pages)

After Section 4, the Schedule provides information about pharmaceutical benefits under the RPBS. These may only be prescribed to DVA beneficiaries holding one of the repatriation health cards (see details under '4. Patient Charges').

2. Prescribing Medicines — Information for Doctors and Dentists

Eligible prescribers

Pharmaceutical benefits can only be prescribed by registered doctors and by dentists who are approved to work within the PBS.

PBS Prescription forms

Standard PBS prescription forms are available from Medicare Australia for prescribing pharmaceutical benefits.

For doctors:

- *Personalised forms* – are printed with the doctor's name, qualifications, practice address/es, telephone number and prescriber number (which relates to pharmaceutical benefits). They are only provided to doctors who have a Medicare provider number.
- *Non-personalised (blank) forms* – are distributed as an emergency supply (usually when a doctor has temporarily run out of personalised forms).
- *Locum forms* – have the doctor's name, prescriber number and telephone number (if available) and a space to record the practice where the doctor is working.
- *PBS/RPBS Authority Prescription Forms* – can be in personalised, non-personalised or locum format.

- *Computer PBS prescription forms* — are either continuous or single sheet. On the reverse side they list the name, address and telephone number of the practice, and in the case of a sole doctor practice, the doctor's name.

For dentists:

- *Personalised forms* – have the dentist's name, qualifications, practice address/es, telephone number and prescriber number.
- *Non-personalised (blank) forms* – are distributed for emergency supply only.

PBS prescription forms for both doctors and dentists are supplied free of charge.

The inclusion of the prescriber number on a PBS prescription enables the pharmacist to be sure the prescription is from a legitimate prescriber and satisfies State/Territory legislation.

PBS prescriptions should be provided to the patient in duplicate, as both copies make up a valid PBS prescription. The patient should be reminded to present both the original and the duplicate copy to their local pharmacist.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner, in course of employment with the participating public hospital, may prescribe the subsidised medication. The States of Victoria, Queensland and Western Australia have agreed to implement these arrangements.

Ordering forms

Prescribers are asked not to over order. Getting the right amount of forms helps to reduce the cost to taxpayers and helps to reduce paper wastage. Also, the pads may deteriorate if stored over time.

Order forms for standard and authority PBS prescription forms are available from Medicare Australia stationery officers. Contact details are listed in the front of the Schedule. Order forms for computer PBS prescription form stationery are obtained from Medicare Australia (at the address below). Orders should be sent to:

Prescription Pad Order Clerk
Pharmaceutical Branch
Medicare Australia
GPO Box 9826
Sydney NSW 2001
Telephone (02) 9895 3295

Orders for PBS prescription stationery will only be accepted by application in writing and through the channels mentioned above.

Preparing general PBS prescriptions

Do's and Don't's

A PBS prescription is only valid when it is written by a doctor or dentist.

The PBS prescription must be for the treatment of the person named on the PBS prescription. A PBS prescription may only be written for the treatment of one person.

A prescriber cannot write more than one PBS prescription for the same pharmaceutical benefit for the same person on the same day.

Up to **three** pharmaceutical benefit items may be included on a single PBS prescription form, but pharmaceutical benefits and non-pharmaceutical benefits should not be listed together on the one PBS prescription form.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner, in course of employment with the participating public hospital, may prescribe the subsidised medication. The States of Victoria, Queensland and Western Australia have agreed to implement these arrangements.

If an item has a particular manner of administration it may not, as a pharmaceutical benefit, be administered in any other way, e.g., an ophthalmic preparation may not be prescribed for topical use.

If an item is restricted, but the patient is not suffering from one of the specified conditions, it cannot be prescribed as a pharmaceutical benefit. The prescriber should write the prescription either on a private prescription or on a standard prescription with 'PBS/RPBS' clearly struck out. It should also be endorsed 'non-PBS'.

Prescribers must heed State/Territory laws when prescribing drugs listed as narcotic, specified or restricted in the poisons legislation of the particular State or Territory. Legislative requirements in some States/Territories are such that prescribers may be required to prescribe a drug of addiction on a separate PBS prescription.

A prescriber cannot prescribe a narcotic drug for him/herself.

Prescribers are issued with individual PBS prescription pads by Medicare Australia for their own use – these pads should not be used by other prescribers, as this can cause confusion through incorrect pharmacy records.

Doctors should, and dentists are required to, include their prescriber numbers on non-personalised PBS prescriptions.

The following admixtures are not pharmaceutical benefits:

- the admixture of two or more ready-prepared items listed in the Schedule; or
- the admixture of a ready-prepared item and one or more extemporaneous drugs listed in Section 4 of the Schedule (green pages); or
- the admixture of a non-pharmaceutical benefit item with a pharmaceutical benefit item.

Writing the PBS prescription

The following rules apply for writing PBS prescriptions:

- they must be written in indelible form (i.e., ink or ball-point pen) in the prescriber's own handwriting (exceptions must be approved by Medicare Australia's Chief Executive Officer) either on the standard PBS prescription, or on paper approximately 18 cm x 12 cm, or they can be generated by computer on a form approved by Medicare Australia. For patient safety reasons, both the original and the duplicate must be legible;
- they must record the prescriber's name and address (and, in the case of dentists, the prescriber number), the patient's name, address and entitlement status, and whether the prescription is under the PBS or RPBS;
- they should completely identify the pharmaceutical benefit by detailing the item, dose, form, strength, quantity and instructions for use;
- they should indicate where brand substitution is not permitted. PBS prescriptions must not be prepared using a computer prescribing program that contains a default which would result in all prescriptions being indicated as Brand Substitution Not Permitted;
- where 'solvent required' is included after the form, the volume and number of ampoules must be specified; and
- they must be signed by the prescriber and dated. Forward or back dating is not permitted.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner, in course of employment with the participating public hospital, may prescribe the subsidised medication. The States of Victoria, Queensland and Western Australia have agreed to implement these arrangements.

Restrictions

Pharmaceutical benefits listed in the Schedule fall into three broad categories:

Unrestricted benefits – which have no restrictions on their therapeutic uses;

Restricted benefits – which can only be prescribed for specific therapeutic uses (they are noted as **restricted benefit**); and

Authority required benefits – which are restricted and require prior approval from Medicare Australia or the DVA (they are noted as **authority required**).

Authority PBS prescriptions

Only doctors (not dentists) can write authority PBS prescriptions.

Approval of authority PBS prescriptions by Medicare Australia may be obtained by posting an Authority Prescription Form to Medicare Australia, by calling Medicare Australia's Authority Freecall service (1800 888 333), or by using Medicare Australia's PBS authorities website at www.medicareaustralia.gov.au/providers. Approval of authority prescriptions by the DVA may be obtained either by posting an Authority Prescription Form to the DVA, or by using the DVA's Authority Freecall service (1800 552 580).

An authority PBS prescription cannot have retrospective approval.

An authority PBS prescription is not valid until it has been approved by Medicare Australia or the DVA. Until then, a pharmacist must not supply the item as a PBS benefit.

The Authority PBS Prescription Form includes:

- the original copy, which records doctor, patient and pharmaceutical benefit item details. After approval, Medicare Australia will forward it with the duplicate to the patient or the doctor (if it is to be sent direct to the patient, the doctor should mark the box next to the patient's details).
- the duplicate copy, which is used for repeat supplies and is attached to repeat authorisations until the last supply is made. The duplicate is then retained by the pharmacist. In the new format, the duplicate ("Medicare Australia/DVA copy") is forwarded to Medicare Australia for processing and payment.
- the triplicate ("doctor or Medicare Australia/DVA") copy, which is kept by Medicare Australia or the DVA for record purposes when approval is sought in writing. Where approval is by telephone, the doctor must keep this copy for 12 months. This copy must record the daily dose, details of the disease, the clinical justification for using the item, the patient's age (if the patient is a child) and whether the patient has previously received an authority for this pharmaceutical benefit.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner, in course of employment with the participating public hospital, may prescribe the subsidised medication. The States of Victoria, Queensland and Western Australia have agreed to implement these arrangements.

Writing authority PBS prescriptions

The following rules apply:

- only one item may be prescribed per PBS prescription;
- PBS prescriptions must be completed by doctors in writing, unless otherwise approved by Medicare Australia;

- doctors should include their name, address, telephone number and **prescriber number** (not provider number);
- doctors must include the patient's name, address and entitlement status (i.e. whether they are a 'concessional' or 'general' patient);
- doctors must indicate when brand substitution is not permitted. PBS prescriptions must not be prepared using a computer prescribing program that contains a default which would result in all PBS prescriptions being indicated as Brand Substitution Not Permitted;
- in certain circumstances, the doctor must provide additional information to Medicare Australia with the authority application; and
- the PBS prescription must be signed by the doctor and dated.

Posted applications which lack necessary information, and therefore cannot be approved, will be returned to doctors for correction. If a doctor can clarify the matter via telephone, an Authority to Prescribe Form may be prepared by Medicare Australia or the DVA and sent to the doctor.

In the case of authority PBS prescriptions approved by telephone, the doctor should ensure the approval number is included on the PBS prescription to enable the pharmacist to supply the medication. A prescriber who is granted approval but decides not to continue with the therapy should advise Medicare Australia.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner, in course of employment with the participating public hospital, may prescribe the subsidised medication. The States of Victoria, Queensland and Western Australia have agreed to implement these arrangements.

Maximum quantities and repeats

The maximum quantity and number of repeats allowed for PBS items are recommended by the Pharmaceutical Benefits Advisory Committee (PBAC). In the case of RPBS items, the recommendations come from the Repatriation Pharmaceutical Reference Committee (RPRC).

Only doctors (not dentists) can prescribe repeats.

PBS prescriptions and repeats can be for any quantity up to the maximum. It is not necessary to prescribe the maximum quantity if a lesser quantity is sufficient for the patient's needs. Please clearly indicate the number of tablets, capsules, etc. required and the number of repeats needed, and **do not use** abbreviations such as 'Max. Qty', 'M.Q.', or 'M.R.'.

If a doctor feels the maximum quantity or number of repeats should be increased for a particular patient, he or she must complete an Authority Prescription Form (see procedures above under 'Authority prescriptions'). The provision of increased quantities and repeats on authority PBS prescriptions is intended to provide approximately one month's therapy which may be repeated (if clinically appropriate) to provide 6 months' therapy in total. This situation usually arises where higher than normal dosages are required.

Approval for increased quantities and repeats of authority required and restricted benefit PBS items will be granted only where the reason for the PBS prescription is consistent with the indications published in the Schedule.

Approval for increased quantities and repeats extends only to the provision of a pharmaceutical benefit for the patient and does not imply approval of any aspects of the patient's care, which are the responsibility of the treating doctor.

Regulation 24

Under this regulation, original and repeat supplies of pharmaceutical benefits can be supplied at the one time if a doctor is first satisfied that certain conditions apply, then endorses the PBS prescription 'Regulation 24'. RPBS prescriptions may be endorsed 'hardship conditions apply'.

The doctor must first be satisfied all the following conditions apply:

- the maximum PBS quantity is insufficient for the patient's treatment; **AND**
- the patient has a chronic illness or lives in a remote area where access to PBS supplies is limited; **AND**
- the patient would suffer great hardship trying to get the pharmaceutical benefit on separate occasions.

Urgent cases

In urgent cases and where State/Territory law allows, a prescriber may telephone a pharmacist and ask that a PBS prescription be supplied. He/she must then forward the written PBS prescription and duplicate to the pharmacist within **seven days of the date of supply**.

This also applies to authority PBS prescriptions provided prior approval has been given by Medicare Australia or DVA. The follow-up written PBS prescription must include the approval number provided over the phone by Medicare Australia or DVA.

Drugs of addiction

Prescribers must heed State/Territory laws when prescribing drugs listed as narcotic, specified or restricted and must notify, or receive approval from, the appropriate health authority.

When a PBS/RPBS authority application is for a drug of addiction (other than dexamphetamine sulfate), the following guidelines apply:

- the maximum quantity authorised is generally for one month's therapy (e.g., one week's therapy with three repeats);
- where supply for a longer period is warranted, quantities are usually for up to three months' therapy;
- telephone approvals are limited to one month's therapy.

Doctors should also state the interval of repeat where repeats are called for, and ensure State/Territory health authorities are notified about ongoing treatment.

Emergency drug (doctor's bag) supplies

Certain pharmaceutical benefits are provided without charge to doctors who in turn can supply them free to patients for emergency use.

A doctor must fill out and sign the Emergency Drug (Doctor's Bag) Order Form in triplicate and give the original and duplicate to a pharmacist. Each form is valid for the month indicated on the form.

Doctors can order the maximum quantity of an item provided they do not already have the maximum quantity on hand. Doctors can only get items once a month. They can also ask for a particular brand of a pharmaceutical benefit. If it is unavailable, they must specify another listed brand, and initial the alteration.

A receipt must be signed by the doctor, or by an authorised representative, when supplies are received.

3. Supplying Medicines — What Pharmacists Need to Know

Eligible suppliers

Pharmaceutical benefits are mainly supplied by approved pharmacists – pharmacists who comply with certain conditions. These pharmacists are approved to dispense pharmaceutical benefits from a particular pharmacy.

Other suppliers include approved doctors (usually practising in isolated areas), Friendly Society pharmacies, and approved hospitals. All suppliers are issued with approval numbers by Medicare Australia. They should follow the procedures in these Explanatory Notes.

Unapproved pharmacists *cannot* supply pharmaceutical benefits.

Approval conditions for pharmacists

A pharmacist approved to supply medicines under the PBS:

- can only supply benefits from the pharmacy that he/she is operating;
- will not supply to anyone any pharmaceutical benefit that attracts a Commonwealth contribution for free, or for a price that is less than the relevant patient contribution;
- will clearly advertise that any offer for free or cut-price medicines does not include pharmaceutical benefits which have a Commonwealth contribution;
- will not pay rebates or refunds of patient contributions;
- will publicly display a notice setting out the pharmacy's normal trading hours;
- is obliged to supply pharmaceutical benefits at the pharmacy at any hour if a PBS prescription is marked 'urgent' and initialled by the prescriber;
- will keep adequate stocks for the supply of pharmaceutical benefits;
- may be called on by Medicare Australia to provide details of stocks of pharmaceutical benefits or preparations for pharmaceutical benefits; and
- must keep the duplicates of all old format PBS prescriptions, and the patient/pharmacist copies of all new format PBS prescriptions, with a Commonwealth contribution for at least one year from the date of supply. This includes PBS prescriptions ordering repeats when it is the final supply, and order forms for emergency drug (doctor's bag) supplies. Please note that some State/Territory laws require these copies to be kept for longer periods.

Before supplying pharmaceutical benefits

Several steps must be taken before a pharmaceutical benefit is supplied.

Firstly, a pharmacist must endorse the PBS prescription and duplicate with his/her name and approved supplier number.

Secondly, a PBS prescription identifying number must be given to the PBS prescription item on both the PBS prescription and duplicate. Any recognised series of numbers may be used.

If more than one item is on a PBS prescription, a separate identifying number should be allocated to each item.

In the case of a repeat authorisation, the same PBS prescription identifying number(s) must be carried through for each item. A pharmacist must also allocate his/her own identifying number on the repeat authorisation. It must be written alongside the date and place of supply.

Supplying pharmaceutical benefits

Do's and Don't's

Except in urgent cases (see details under '2. Prescribing Medicines ... Urgent cases'), pharmacists are authorised to supply pharmaceutical benefits only after they receive:

- the original and duplicate of a valid PBS prescription which is not more than 12 months old; or
- the original and duplicate of an approved authority PBS prescription or an authority to prescribe which is not more than 12 months old; or
- a repeat authorisation attached to a duplicate PBS prescription (or in the new format, the "patient/pharmacist" copy) not more than 12 months after the date of the original PBS prescription.

A pharmaceutical benefit cannot be supplied more times than specified in the PBS prescription.

A pharmacist cannot add to, delete from, or alter a PBS prescription in any other way. However, there may be circumstances where after contacting a prescriber, the pharmacist can clarify the prescriber's intentions and endorse the PBS prescription accordingly.

Once a pharmaceutical benefit has been supplied to a patient, it may not be supplied to that patient again:

- on the same day or within the next 20 days, if it is a benefit (other than an eye preparation) that has five or more repeats allowed in the Schedule; or
- on the same day or within the next four days (e.g., if a pharmaceutical benefit is supplied on a Monday, it cannot be supplied again to that patient until the next Saturday) in the case of other benefits.

Exceptions to this are:

- when a PBS prescription is endorsed with the words 'Regulation 24' or 'hardship conditions apply' (see below under 'Regulation 24'); and
- If a pharmacist believes supply is urgently needed to treat the person, or a previous supply has been destroyed, lost or stolen. In this case, the pharmacist can provide another supply but must write 'immediate supply necessary' and sign the PBS prescription.

With the agreement of the patient, a pharmacist can supply an alternative brand of a benefit without reference to the prescriber, provided that:

- the PBS prescription does not indicate 'brand substitution not permitted';
- the Schedule shows the prescribed brand and the substitute brand are bioequivalent; and
- supply of the substitute brand does not contravene relevant State/Territory law.

A pharmacist cannot substitute one medicine for another – only an alternative brand of the medicine, as specified above.

Pharmacists must heed State/Territory laws when supplying drugs listed as narcotic, specified or restricted in the poisons legislation of the particular State or Territory.

What to do if the Schedule changes

If an item or brand is deleted from the Schedule, it *cannot* be supplied as a pharmaceutical benefit from the date the deletion takes effect – regardless of whether the PBS prescription was written before this date. This includes repeat authorisations. (Special conditions applying to RPBS prescriptions are detailed in the RPBS Explanatory Notes.)

However, if restrictions on the prescribing of a pharmaceutical benefit change, or the maximum quantity or number of repeats is altered in the Schedule, valid PBS prescriptions written before the date of effect of the change *may* still be supplied as pharmaceutical benefits, under the conditions applying at the date of prescribing.

Suspected forgery

Pharmacists should take all reasonable steps to satisfy themselves that all items on a PBS prescription were written by a doctor or a dentist.

Regulation 24

This regulation allows pharmacists to supply a pharmaceutical benefit and all of its repeats at the one time. The PBS prescription must be endorsed by the doctor with the words 'Regulation 24' if it is an item under the PBS, or 'hardship conditions apply' if it is being supplied under the RPBS. (For more information see under '2. Prescribing Medicines ... Regulation 24').

Repeat authorisations

When a PBS prescription calls for repeat supplies, the pharmacist shall prepare a Repeat Authorisation Form, except when the PBS prescription is marked 'Regulation 24'.

The repeat may be requested on a standard PBS prescription, an authority PBS prescription or an Authority to Prescribe Form, or on an earlier repeat authorisation. In the latter case, it must come with the duplicate PBS prescription, or in the new format, the "patient/pharmacist copy".

Preparing Repeat Authorisation Forms

A Repeat Authorisation Form must show:

- the category of benefit (concession or general) – by placing a cross (x) in the relevant box;
- the patient's name and full address;
- in the case of repeats authorised on authority PBS prescriptions, the authority number;
- details of the original PBS prescription stating the item, brand, form, strength, quantity and directions;
- if brand substitution has occurred, the name of the brand actually supplied;
- for the first supply, the pharmacy approval number, the date of the original PBS prescription and the allotted PBS prescription identifying number;
- for subsequent supplies, the pharmacy approval number and the date of the original PBS prescription number;
- the number of times the item is to be repeated and the number of times it has been supplied;
- the name and pharmacy approval number of the pharmacist issuing the repeat authorisation; and
- the date of supply.

When a repeat authorisation is prepared for any further repeats or deferred supply, a pharmacist must attach the duplicate copy of an old format PBS prescription, or the patient/pharmacist copy of a new format PBS prescription, and give both to the patient at the time of supply.

Repeat authorisations for injectables and solvents

Where an injectable pharmaceutical benefit requires a solvent, both items should be treated as one pharmaceutical benefit. If repeats are needed, only one repeat authorisation is to be prepared. Details of the injectable and the solvent should appear in the space provided for the 'original prescription transcription'.

Repeat authorisations for deferred supply

When a PBS prescription orders a number of pharmaceutical benefit items, but the patient does not need all of the items at the same time, a separate repeat authorisation for each deferred item must be prepared. The words 'original supply deferred' should be indicated across the relevant item on the original PBS prescription, its duplicate, and on the repeat authorisation.

Deferred items must not be claimed on the original PBS prescription.

The Repeat Authorisation Form when it is used for a deferred supply, is issued in the same way as normal repeat authorisations except that:

- '0' is to be inserted in the space for 'no. of times already dispensed'; and
- if no repeats are ordered, '0' is to be inserted in the space for 'no. of repeats authorised'.

Supplying a benefit on a deferred supply repeat authorisation is to be treated as if it is the first time of supply. If repeats are directed, the normal procedure for repeat authorisations applies. Details of the pharmacy at which the deferred supply was authorised are to be written onto subsequent repeat authorisations.

Authority PBS prescriptions

If a pharmacist is presented with an authority PBS prescription and is not sure if it has been approved, he or she should contact Medicare Australia. Please note that Medicare Australia will not provide clinical information.

Urgent cases

In urgent cases and where State/Territory law allows, pharmacists can supply a pharmaceutical benefit to a person without a PBS prescription, provided details of the prescription are given by the prescriber via telephone or other means. The prescriber must then forward the written PBS prescription and duplicate to the pharmacist within **seven days of the date of supply**.

Where a pharmaceutical benefit needs prior approval from Medicare Australia or the DVA, the prescriber must obtain approval and then advise the pharmacist of the PBS prescription and approval details.

Only an original supply can be provided in this manner – not repeats.

Receipts

A person receiving a pharmaceutical benefit item must sign and date a receipt for it. If the person is not the patient, that person must also endorse the PBS prescription or repeat authorisation with his/her address. A receipt cannot be obtained until supply of the benefit has been made.

If a pharmaceutical benefit has to be sent through the post, by rail, or by other means, and a receipt is not practical, the pharmacist must certify on the PBS prescription or repeat authorisation that the benefit has been supplied, and write the date of supply and details of how it was sent. For example, if a pharmaceutical benefit is mailed to a patient on 1 April 2005, the pharmacist should write: "Certified supplied – mailed to patient 1 April 2005 (name of pharmacist) (signature of pharmacist) (date of certification)".

If an item is supplied in an urgent case (see above), or to a person who cannot read or write, the pharmacist should sign and date a statement on the PBS prescription or repeat authorisation, stating the item has been supplied and the date on which it was supplied, and explaining why there is no receipt. For example, if a pharmaceutical benefit is supplied to a patient with a broken arm on 1 May 2005, the pharmacist should write: "Certified supplied 1 May 2005 – patient has a broken arm and is unable to sign (name of pharmacist) (signature of pharmacist) (date of certification)".

Only the pharmacist approved to supply pharmaceutical benefits can certify supply.

Emergency drug (doctor's bag) supplies

Pharmacists may supply certain pharmaceutical benefit items free of charge to doctors for emergencies if they receive an Emergency Drug (Doctor's Bag) Order Form in duplicate, signed by the doctor.

Pharmacists must be satisfied the form was completed by a doctor and includes the doctor's name and address. If a pharmacist does not know the doctor, he/she should confirm the doctor's registration and endorse this on the back of the form.

For more information about emergency supplies see under '2. Prescribing Medicines ... Emergency drug (doctor's bag) supplies'.

4. Patient Charges

Type of patient

There are two types of PBS beneficiaries – general patients and concessional patients. General patients hold a Medicare card. Concessional patients hold a Medicare card and one of the following cards from Centrelink or the DVA:

- Pensioner Concession Card
- Commonwealth Seniors Health Card
- Health Care Card
- Repatriation Health Card For All Conditions (gold) – concessional patients under RPBS
- Repatriation Health Card For Specific Conditions (white) – only regarded as concessional patients for RPBS prescriptions unless they hold a separate entitlement from Centrelink, otherwise they are general patients
- Repatriation Pharmaceutical Benefits Card (orange) – concessional patients under RPBS
- Safety Net Concession Card or Safety Net Entitlement Card are also issued by Medicare Australia.

The above concessional patients are recognised by public hospitals in all States and Territories apart from South Australia (where DVA beneficiaries are treated as general patients) and New South Wales (where holders of a white DVA card are treated as general patients).

Under the Reciprocal Health Care Agreements (RHCA), visitors from participating countries (see the introduction of this section for the list of countries) are treated as general patients – they do not have concessional entitlements. To receive pharmaceutical benefits, these visitors may need to present a temporary Medicare card or their passport. Pharmacists should contact Medicare Australia if they have queries about these arrangements.

Establishing entitlement

PBS prescription forms supplied by Medicare Australia have spaces provided for details of a patient's entitlement status. Anyone can enter this information, which must include:

- a cross (x) in the appropriate box to indicate the level of patient contribution; and
- the complete Medicare number (including individual reference number) or complete Veteran file number on the card; and
- if applicable, the complete concession number on the card.

The person who signs the receipt for pharmaceutical benefits also accepts responsibility for the validity of the entitlement information on the PBS prescription.

All PBS prescriptions must have a Medicare or Veteran file number. All concessional PBS prescriptions must have a concession number. However, it is not necessary for the Medicare (Veteran file) or the concession number to be endorsed on the PBS prescription if it is included in the electronic prescription details supplied by a pharmacist who is using the Claims Transmission System.

What to charge

Patient contribution

Under the PBS, the maximum cost for a pharmaceutical benefit item at a pharmacy is \$28.60 for general patients and \$4.60 for concessional patients (except where a special patient contribution, a brand premium, or a therapeutic group premium applies). General patients who have reached the safety net threshold (see details under '5. The Safety Net Scheme') receive pharmaceutical benefit items at the concessional rate of \$4.60.

Patients who have a Safety Net Entitlement Card (see details under '5. The Safety Net Scheme') receive PBS items for free, except when a special patient contribution, brand premium, or therapeutic group premium applies.

The contribution rate for general patients as outpatients at public hospitals throughout Australia is \$22.90. The exception is Queensland and hospitals participating in the pharmaceutical reforms where they pay the safety net value of an item when it is listed in the Schedule (see details under '5. The Safety Net Scheme'), or up to \$28.60 for items not listed in the Schedule. The public hospital pharmaceutical reforms enable participating public hospitals to prescribe and supply pharmaceutical medication from the Pharmaceutical Benefits Scheme to outpatients and patients upon discharge. A range of chemotherapy drugs will also be available for day-admitted and non-admitted chemotherapy patients.

The contribution rate for concessional patients in all public hospitals is \$4.60.

The supply of a pharmaceutical benefit or a Repatriation pharmaceutical benefit to a patient is a GST-free supply. Goods and services tax must not be included in the price charged to a patient for the supply of a benefit under the PBS or RPBS.

It is the patient's responsibility to meet any charge lawfully demanded by an approved pharmacist, otherwise supply may be refused.

The patient contribution rates are usually adjusted on 1 January each year in line with inflation.

Patient contributions for early supply of some PBS medicines

From 1 January 2006, prescriptions for some pharmaceutical benefits are not eligible for Safety Net benefits if resupplied within 20 days of a previous supply of the same pharmaceutical benefit for the same person under the PBS or the RPBS. This is known as the 'Safety Net 20 day rule'. (see also details under '5. The Safety Net Scheme' and '7. How Pharmacists Claim Reimbursement')

Where a prescription is subject to the Safety Net 20 day rule, exclusion from Safety Net benefits has the following effects:

- the patient contribution does not count towards the Safety Net
- after the Safety Net threshold is reached, the usual patient payment amount for the corresponding entitlement level (not the Safety Net amount) applies.

For example: The payment for such a prescription for a patient with a Safety Net Entitlement Card would be \$4.60 – not free. For a general patient with a Safety Net Concession Card, the usual general patient amount (up to \$28.60) would apply – not the Safety Net concessional amount of \$4.60.

The Safety Net 20 day rule does not apply to PBS/RPBS prescriptions originating from hospitals or day hospital facilities.

Special patient contributions, brand premiums and therapeutic group premiums

A special patient contribution is payable for a pharmaceutical benefit when there is a disagreement between the manufacturer and the Government over the dispensed price for that benefit item. This extra charge is paid by all patients, together with their usual patient contribution. Other than for bleomycin sulfate, exemptions on medical grounds are available. For RPBS special patient contribution arrangements see the RPBS Explanatory Notes.

Under the brand premium arrangements, Commonwealth reimbursement to pharmacists is based on the lowest-priced brand. Patients pay the difference for higher-priced brands, on top of their usual patient contribution.

Under the therapeutic group premium arrangements, Commonwealth reimbursement to pharmacists is based on the lowest priced benefit items within identified therapeutic groups. Patients pay the difference for higher priced items. Exemptions on medical grounds are available, but must be granted by Medicare Australia.

The Schedule's special patient contributions, brand premiums and therapeutic group premiums apply to maximum quantities. When a quantity is less than, or – on an authority or regulation 24 PBS prescription – more than, the maximum, the contributions or premiums will be a fraction or multiple of the maximum quantity, using standard pricing rules (see details under '9. Pricing Prescriptions').

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner, in course of employment with the participating public hospital, may prescribe the subsidised medication. The States of Victoria, Queensland and Western Australia have agreed to implement these arrangements.

Solvents

Where a solvent is prescribed as part of a pharmaceutical benefit, only one patient contribution is charged.

Increased quantities

Where a doctor has written an authority PBS prescription for a quantity greater than the maximum, the relevant patient contribution should be made for each supply of the increased maximum quantity.

Regulation 24

For 'Regulation 24' PBS prescriptions, a pharmacist should charge the usual patient contribution for the original and for each repeat quantity needed to make up the total supply (plus any special patient contribution, brand premium or therapeutic group premium if applicable).

After hours

A pharmacist may charge an extra fee if he/she supplies a PBS item outside his/her pharmacy's normal trading hours. The charge is paid by the patient and does not count towards the safety net.

Delivery

A charge can be added for delivering pharmaceutical benefits from the pharmacy. This charge does not count towards the safety net. For RPBS delivery arrangements refer to the RPBS Explanatory Notes.

5. The Safety Net Scheme

The Safety Net Scheme is designed to protect those patients and their families who require a large number of PBS and RPBS items, and applies to each calendar year.

For the purposes of the scheme, the family includes:

- the spouse or de facto spouse;
- family members under the age of 16 who are in the care and control of the patient; or
- dependent full-time students under the age of 25.

The scheme requires pharmacists, on request by patients, to record the supply of PBS and RPBS items on Prescription Record Forms (PRF's). When patients reach a certain spending level (or Safety Net threshold) within a calendar year, they qualify to receive PBS or RPBS items at a cheaper price or free of charge for the rest of that year. The reductions do not apply to special patient contributions, brand premiums or therapeutic group premiums – these charges must still be met by patients.

The Safety Net threshold may be reached by accumulating PBS prescriptions through community pharmacies or public hospitals or a combination of both.

Pharmaceutical benefit items (including authority items) can only be counted towards the safety net threshold when prescribed and supplied according to the Schedule's conditions. A medicine supplied by a pharmacist who is not approved to supply pharmaceutical benefits cannot count towards the Safety Net.

From 1 January 2006, prescriptions for some pharmaceutical benefits are not eligible for Safety Net benefits if resupplied within 20 days of a previous supply of the same pharmaceutical Benefit for the same person under the PBS or RPBS. For such prescriptions, the patient contribution cannot count towards the Safety Net. (see also details under '4. Patient Charges' and '7. How Pharmacists Claim Reimbursement') This does not apply to out-patient medications in public hospitals or to any prescriptions originating from a hospital or day hospital facility.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner, in course of employment with the participating public hospital, may prescribe the subsidised medication. The States of Victoria, Queensland and Western Australia have agreed to implement these arrangements.

Safety net thresholds

There are two safety net thresholds – one for general patients and the other for concessional patients.

The general patient safety net threshold is currently \$874.90. When patients and/or their families reach this amount, they can apply for a Safety Net Concession Card and pay only \$4.60 per PBS prescription for the rest of the calendar year.

The concessional safety net threshold is \$239.20 (this also applies to gold, white or orange card holders under the RPBS). Once patients and/or their families reach this amount, they can apply for a Safety Net Entitlement Card and receive items free of charge for the rest of the calendar year.

Brand premiums, therapeutic group premiums and special patient contributions do not count towards the safety net thresholds.

The thresholds may be adjusted on 1 January each year in line with inflation.

Safety net cross-over arrangements

Some patients and/or members of their families will move in and out of concessional status during the calendar year. Patients should apply for the safety net card appropriate to their status at the time they apply.

Concessional patients who were previously general patients can apply for a Safety Net Entitlement Card once they reach the threshold of \$239.20. In this case, any pharmaceutical benefits previously supplied at the general rate in that calendar year will need to be converted to the rate of \$4.60 per item.

General patients who were previously concessional patients can apply for a Safety Net Concession Card when they reach the general threshold of \$874.90. In this case, any pharmaceutical benefits previously supplied at the concessional rate in that calendar year will be counted at the rate of \$4.60 per item.

In the case of families where one parent holds a concession card and other family members are general patients, the family can choose to apply for either a Safety Net Entitlement Card or a Safety Net Concession Card.

To receive a Safety Net Entitlement Card, all pharmaceutical benefits (including general PBS prescriptions) are counted at \$4.60 per item until the \$239.20 threshold is reached. To receive a Safety Net Concession Card, general pharmaceutical benefits are counted at the general rate of up to \$28.60 per item and concessional pharmaceutical benefits at \$4.60 per item, until the threshold of \$874.90 is reached.

White DVA card holders can either be general or concessional patients (depending on their Centrelink entitlements). If they are being treated for a specific disability accepted by the DVA, they are also supplied with specific items under the RPBS at the concessional rate of \$4.60 per item. Therefore, these patients are encouraged to maintain a concessional PRF, plus a general PRF for those items not covered under the RPBS.

White card holders may choose at any time during the year to count contributions made at the general level towards the concessional safety net threshold. The patient receives a credit of \$4.60 for each PBS prescription item purchased. Alternatively, he/she can count contributions at the concessional level towards the general safety net, and receive a credit of \$4.60 for each RPBS prescription item purchased.

Those with gold or orange cards receive all of their prescription items under the RPBS, and only pay \$4.60 for each item.

Dependants of white, gold or orange card holders are treated separately and may be either general patients or concessional patients. Their prescriptions may be included in the cross-over arrangements.

Recording PBS prescriptions

There are two types of PRF's to record PBS prescription items. One (a blue form) is used by general and concessional patients and veterans who pay for items at community pharmacies. It is available from community pharmacies, Medicare offices and Medicare Australia in each State. The other form (maroon) is used by out-patients who pay for items at public hospital pharmacies, and is available through hospital out-patient departments or Medicare Australia State offices.

Patients should record their status (general or concessional) in the appropriate box on the front of the PRF, state their Centrelink, DVA and/or Safety Net Concession/Entitlement Card number, and list family members covered. General patients must also record their Medicare number when applying for a Safety Net Concession Card.

Details to be entered on the form by the pharmacist are:

- date of supply;
- PBS/RPBS code number of the item (for community pharmacies only);
- the safety net value of the item (for community pharmacies only);
- pharmacist's approval number (for community pharmacies only);
- item identification – medicine code, name of medicine or abbreviation (for public hospitals only);
- hospital charge (for public hospitals only);

- hospital safety net number (for public hospitals only); and
- signature of the authorised person making the entry.

Community pharmacists should record in the 'safety net value' column:

- the patient contribution when it is less than the PBS dispensed price; or
- the safety net value shown in this Schedule, or any lesser amount charged, if the PBS dispensed price is less than or equal to the patient contribution. The pharmacist may discount the price for these items.

Some computer software suppliers provide a special label to record this information on the PRF. Some suppliers also provide a computer printout option that is an acceptable replacement for the PRF.

The patient is responsible for maintenance and storage of the PRF. However, it may be kept in the pharmacy. An individual (or family) may have more than one PRF.

Hospital PRF's

Items to be recorded on a hospital PRF must be approved by the hospital's pharmaceutical advisory committee. They can be listed on a hospital's formulary (a list of pharmaceutical items approved by the committee for the treatment of particular illnesses), or authorised on a patient-by-patient basis.

Multi-item forms

If a patient submits a multi-item PBS prescription form, which would take the PRF past the safety net expenditure limit, any items in excess are to be treated as entitled items once the Safety Net Entitlement/Concession Card is issued.

These excess items should be treated as 'deferred supply' items.

For example, if a family has \$840.00 recorded on their PRF and they have a new PBS prescription for three items, the first item should be supplied at the general rate. If the second item would take the family over the threshold of \$874.90, the pharmacist should then issue a Safety Net Concession Card and supply the other two items at the concessional rate. This involves the deferral of two items, entry of the Safety Net Concession Card number into the computer, and the subsequent supply of these items.

Qualifying PBS prescriptions

A PBS prescription should be supplied at either the concessional rate or free of charge, as appropriate, when the safety net value or hospital charge for that PBS prescription takes the PRF over the qualifying amount for a Safety Net Entitlement/Concession Card.

Lost PRF's

If a PRF has been lost, stolen or destroyed, a pharmacist may prepare a duplicate copy, but is under no obligation to do so.

Retrospective entitlement and patient refunds

Responsibility for claiming entitlements rests with the patient. If items accumulated on a PRF have already exceeded the safety net limit, then the cost to the patient of those items in excess of the limit cannot be refunded by a pharmacist.

However, a patient may get a refund in the following circumstances:

- The patient failed to apply for his/her Safety Net Entitlement/Concession Card on reaching the safety net threshold —

The patient should write to Medicare Australia and provide copies of pharmacy accounts or a signed statement from the pharmacist giving the date of supply, description and cost of items supplied and paid for. A copy of the relevant PRF's should also be provided. If they are not available, the patient should give the name of the pharmacy where the card was issued and the number on the card, so Medicare Australia can locate the PRF's through its records. Cash refunds are not available in these circumstances – cheques are issued. Patients can write to the Medicare Australia office in their State. Contact details are provided in the front of the Schedule (pink pages).

- The patient cannot satisfy a pharmacist that he/she has a current entitlement and is charged the general patient price —

The pharmacist should issue the patient with a receipt and a claim form (provided to the pharmacist by Medicare Australia). The patient can then get a refund from Medicare Australia via Medicare offices or PBS processing centres. RPBS prescription refunds are paid at DVA State offices.

Medicare Australia can only pay refunds for PBS items supplied through approved pharmacies. Refunds for hospital supplied items should be referred to the relevant State/Territory hospital or health department. Refunds cannot be made where the patient was charged the general or concessional amount instead of the Safety Net concessional or Safety Net entitlement amount, respectively, as a result of the Safety Net 20 day rule. (see details under '4. Patient Charges') Receipts for prescriptions where the Safety Net 20 day rule has applied must include 'SN20DR' to indicate the reason for the amount charged.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner, in course of employment with the participating public hospital, may prescribe the subsidised medication. The States of Victoria, Queensland and Western Australia have agreed to implement these arrangements.

Applying for a Safety Net Entitlement/Concession Card

Once the safety net threshold has been reached, any adult patient covered by the PRF can complete the application and declaration on the back of the form or the computer-generated application to get a Safety Net Entitlement/Concession Card. Please note that software packages that produce computer generated applications must be approved by Medicare Australia.

If the card is issued to a dependent child or student, it should be in the name of a parent.

When issuing entitlement/concession cards, pharmacists do not have to check all PRF details. However, they should ensure each entry has been signed and that the PRF total qualifies the patient for the relevant safety net card.

In the case of a concession card, pharmacists should check that the patient's Medicare card number is on the PRF.

Issuing a Safety Net Entitlement/Concession Card

When satisfied that the individual or family is entitled, the pharmacist should issue the next blank Safety Net Entitlement/Concession Card with the following details:

- the names of family members covered. If there are more than eight family members, a second card should be issued listing the card holder and family members not listed on the first card. The PRF has space to record that two cards have been issued;
- the two-character code to indicate the relationship to the card holder. Applicable codes are:
 - SP – spouse;
 - DC – child under 16 years; and
 - DS – dependent full-time student under 25 years.

The pharmacist should be satisfied that only family members are listed on the card. The unused space on the card should be ruled through to prevent extra names being added. The sticky label from the Safety Net Entitlement/Concession Card, pre-printed with the card number, should be attached to the PRF. The pharmacist should sign and stamp each PRF with the pharmacy's stamp. He/she must then enter the card issue details on a Safety Net – Claim for Payment Form.

Issuing supplementary cards

A pharmacist can give a card holder a supplementary card for a spouse or dependant only at the time the original card is issued. The duplicate card should be recorded in the additional box on the PRF.

Requests for supplementary cards at a later stage should be referred to Medicare Australia State offices, as should requests to add a new family member to the original card (e.g., a new baby or adopted child).

Notification to Medicare Australia and claim for payment

Payment for issuing a Safety Net Entitlement/Concession Card is made after the Safety Net – Claim for Payment Form is sent to Medicare Australia's State office, no later than one month after a card is issued.

Each form must be accompanied by all supporting documentation (PRF's and cancelled or void Safety Net Entitlement/Concession Cards).

Payment will not be made for void cards.

Lost Safety Net Entitlement/Concession Cards

When a card has been lost, damaged, stolen or destroyed, a pharmacist cannot re-issue a patient with a replacement card. The original card holder (or spouse) must apply to Medicare Australia's office in his/her State.

Pharmacy record of issued cards

A record of all cards issued must be kept at the pharmacy from which the pharmacist is approved to supply pharmaceutical benefits. The duplicate ('bookfast') copy in the Safety Net – Claim for Payment book is provided for this purpose.

6. Medicare Australia Entitlement Checks

General Patients

Medicare Australia validates a patient's entitlement to pharmaceutical benefits by checking Medicare and/or Veteran file numbers in pharmacists' claims. If a number is not recorded correctly, a patient cannot be identified against Medicare Australia's Pharmaceutical Benefits Entitlement File and entitlement cannot be established.

If the Medicare or Veteran file number provided in the pharmacists' claims is incorrect or the number and the name supplied do not match Medicare Australia records to enable patient identification, an appropriate warning or rejection code will be returned to the pharmacy. These notifications of missing or incorrect Medicare or Veteran file numbers are provided to pharmacists in their reconciliation statement produced after the claim period has been paid by Medicare Australia.

Special numbers are available for use in certain circumstances for eligible people who are unable to provide a Medicare number.

Concessional Patients

Medicare Australia routinely validates a patient's entitlement to free or concessional benefits by checking concessional numbers in pharmacists' claims. If a number is not recorded correctly, a patient cannot be identified against Medicare Australia's Pharmaceutical Benefits Entitlement File and entitlement cannot be established.

When a number is found to be from a card which was incorrect, expired at the time of supply or entitlement was withdrawn, warning or rejection codes will be returned to the pharmacy to assist with validation of concessional entitlement in relation to future claims from the same patient.

Entitlement checking procedures

General Patients

Once a pharmacist has been notified by Medicare Australia of an incorrect Medicare or Veteran file number he/she should correct the number for future claims by:

- updating his/her system to reflect the correct number provided by Medicare Australia (if patient consent to do so has been obtained); or
- speaking to the patient; or
- obtaining patient consent and calling Medicare Australia on the Improved Monitoring of Entitlements (IME) hotline (1300 302 122).

If the patient presents a Medicare card that appears correct, but according to Medicare Australia is not a valid number, or not a valid number for that person, a pharmacist may use a special number. A photocopy of the card, or a form must accompany the use of this number. The form is available on Medicare Australia's website or by calling 132 290.

Concessional Patients

Once a pharmacist has been notified by Medicare Australia of an incorrect concessional entitlement number, he/she should view the entitlement card to confirm the entitlement number, and start and end dates, when the patient next presents a PBS prescription.

Step by step

Pharmacists should take the following steps where concession entitlement does not appear to be valid or current:

- Re-confirm entitlement with the cardholder/customer;
- Contact Medicare Australia on 132 290, with consent, to confirm the cardholder/customer concession status;
- If Medicare Australia advises that the cardholder/customer is concessionally entitled to receive the PBS medicines on that day, supply the prescription as a concessional entitlement;
- If Medicare Australia advises that the cardholder/customer is not concessionally entitled to receive the PBS medicines on that day, supply as a general prescription. Provide the customer with the information sheet "Your entitlement card" which explains entitlement checking to the customer and the steps they can follow if they are concessionally entitled.

7. How Pharmacists Claim Reimbursement: Information Required

Medicare Australia uses a computerised system for pricing PBS prescriptions, repeat authorisations and emergency drug (doctor's bag) orders, and for calculating claims.

The payment system is designed to pay pharmacists correctly for the pharmaceutical benefits they supply. It is essential instructions are followed carefully and that each document includes all relevant information. Accurate and complete data ensures claim payment is not delayed.

PBS Prescription identification

Pharmacists must include certain information on each PBS prescription sent in for claim, as specified below. It is important that this information is entered correctly and in the right place on the PBS prescription.

This information will be included in a sticker produced by pharmacy software.

The sticker should be placed on the extreme left front of a PBS prescription, opposite each item being claimed. It must not obscure any details written by the prescriber. Most prescribers use PBS prescriptions, which have space for the sticker. If a sticker is not used, a PBS prescription identification stamp can be used or the information can be written in the same place, and in the same order.

Pharmacists should avoid writing over, or placing the sticker over, the prescriber number pre-printed on PBS/RPBS prescriptions, or the 'DP No.' box on PBS dental prescriptions.

The sticker is not necessary for current repeat authorisation, emergency drug (doctor's bag), or for old style authority PBS prescription and authority to prescribe forms, as they have printed spaces for the necessary details. However, it is required for the new format authority PBS prescription forms.

The following information should be entered next to the appropriate letter on the sticker or stamp:

- 'S' – the serial number for the claim (see below)
- 'A' – (a) the price claimed for pricing elected PBS prescriptions, exceptional PBS prescriptions and RPBS non-scheduled prescriptions (see below under 'Extemporaneously-prepared pharmaceutical benefits not listed in the Standard Formulae List' for explanations of pricing elected PBS prescriptions and exceptional PBS prescriptions); and/or
 - (b) confirmation that the PBS prescription is endorsed 'Regulation 24' or the RPBS prescription is endorsed 'hardship conditions apply'; and/or
 - (c) a claim for a glass dropper bottle where applicable; and/or
 - (d) any clarification of the prescription which will assist Medicare Australia payment processing.
- 'No.' – the PBS prescription identifying number.

Serial numbers

PBS prescription, repeat authorisation, authority PBS prescription, and emergency drug (doctor's bag) forms submitted in each claim must bear consecutive serial numbers starting with:

- 1 – for emergency drug (doctor's bag) supplies;
- 1 – for general benefits;
- C1 – for concessional and Safety Net Concession Card benefits;
- E1 – for Safety Net Entitlement Card benefits; and
- R1 – for RPBS benefits.

Each serial number should also be noted on any document kept by the pharmacist for record purposes.

Each emergency drug (doctor's bag) item should be given a serial number, e.g., if there are five items on the first form in the claim, the first item on the second form in the claim will start with the serial number 6.

For prescriptions subject to the Safety Net 20 day rule, the serial number corresponds to the resulting payment category for the pharmaceutical benefit as supplied, not the patient's entitlement category. (see also '8. How Pharmacists Claim Reimbursement: Documents to be Submitted')

Repeat authorisations for authority PBS prescriptions

When a benefit is supplied on a repeat authorisation which needed an authority PBS prescription, the serial number must be prefixed with the letter 'A' for a general benefit; 'AC' for a concessional benefit or a benefit supplied to a Safety Net Concession Card holder; 'AE' for a Safety Net Entitlement Card holder; or 'AR' for a RPBS benefit.

Repeat authorisations for deferred supply

When a benefit is supplied on a repeat authorisation prepared for deferred supply, the serial number must be prefixed with the letter 'D' for a general benefit; 'DC' for a concessional benefit or a benefit supplied to a Safety Net Concession Card holder; 'DE' for a Safety Net Entitlement Card holder; or 'DR' for a RPBS benefit.

Injectable item ordered with a solvent

When both an injectable item and a solvent are to be supplied, only one serial number is used. This number should be placed on the left hand side of the PBS prescription, opposite the injectable item.

Dropper containers

Dispensed prices for extemporaneously-prepared eye drops, ear drops and nasal instillations include the price of a polythene dropper container. However, if a glass dropper container is supplied, payment should be claimed by writing 'glass bottle' in box 'A' of the stamp.

Extemporaneously-prepared pharmaceutical benefits not listed in the Standard Formulae List

When a formula is not listed on the Standard Formulae List (see section 4 of this Schedule – green pages), the PBS prescription is paid at an average of 10 g/mL rate for the type of preparation, unless the pharmacist elects otherwise. A pharmacist may price an exceptional PBS prescription, or elect to price all non-priced extemporaneous PBS prescriptions (see below).

PBS prescriptions paid on an average price basis

If the PBS prescription is to be claimed as an exceptional PBS prescription, the pharmacist should write details of the formula supplied on the PBS prescription or repeat authorisation form; price the PBS prescription in accordance with the pricing principles (as detailed under '9. Pricing PBS Prescriptions'); and enter the calculated price on the sticker.

An exceptional PBS prescription is for an extemporaneously-prepared pharmaceutical benefit that is not included in the Standard Formulae List and for which the price of the ingredients (based on basic pricing rules) is twice or more than the recovery price of the ingredients calculated on an average price basis. Further information on pricing PBS prescriptions can be accessed from the booklet titled *Explanation of Current Pricing* on the Medicare Australia's website at www.medicareaustralia.gov.au (PBS publications for Health Care Providers).

Pricing non-pre-priced extemporaneous preparations

Pharmacists should notify Medicare Australia when they elect to price non-pre-priced extemporaneous preparations. Each PBS prescription should be priced in accordance with the pricing principles (as detailed under '9. Pricing Prescriptions') and that price entered on the sticker.

RPBS prescriptions for items not included in either the PBS or RPBS Schedule

When a prescription for a RPBS patient is for an item not included in either the PBS or the RPBS Schedule, the price claimed should be entered on the sticker. Full details on pricing and availability of such items under the RPBS are set out in the RPBS Explanatory Notes.

8. How Pharmacists Claim Reimbursement: Documents to be Submitted

A claim for pharmaceutical benefits consists of:

- the original and duplicate of a completed Claim for Payment Form;
- the original orders for emergency drug (doctor's bag) supplies in a separate bundle;
- the originals of all old format PBS prescriptions and authority PBS prescriptions, the Medicare Australia/DVA copies of new format PBS prescriptions and authority PBS prescriptions, and all repeat authorisations, separated into four bundles for benefits supplied to the general public; concessional beneficiaries/Safety Net Concession Card holders; Safety Net Entitlement Card holders and RPBS patients.

PBS prescriptions in each bundle should be in serial number order, with serial number 1 at the top of the bundle.

PBS prescriptions subject to the Safety Net 20 day rule are bundled according to the resulting payment category. For prescription forms with multiple PBS items, where the Safety Net 20 day rule would result in different payment categories for different items, dispensing via 'deferred supply' should be used where necessary to allow all items to be included in the correct bundles.

PBS prescriptions in the wrong bundle may be returned to the pharmacist for clarification. If appropriate, they can be resubmitted in the correct bundle in the next claim period.

Completing the claim form

The claimant's name, address of the pharmacy from which the pharmacist is approved to supply pharmaceutical benefits, approval number, and claim period number should be entered on the Claim for Payment Form. These details should match the latest written information held by Medicare Australia, or payments can be delayed while clarification is sought.

The claim period number should state how many claims have been submitted so far in a calendar year, e.g., the sixth claim submitted by an approved pharmacist in 2005 should have a claim period number of 0506.

The first and last serial numbers given to items in each bundle are to be entered on the Claim for Payment Form.

A total claim amount is not required – this will be calculated by Medicare Australia after the PBS prescriptions have been individually priced.

The declaration must be signed by the pharmacist approved to supply pharmaceutical benefits, unless he/she has made arrangements through Medicare Australia for another pharmacist to sign it.

Lodging claims

A claim may be lodged at any time during the month at the relevant Medicare Australia State office. Unless other arrangements have been made with Medicare Australia, the following conditions apply:

- only one claim period can exist and only one claim can be lodged per month;
- the claim period shall cover pharmaceutical benefits supplied during one month; and
- the claim shall be sent within 30 days from when the benefits were supplied.

Claims for pharmaceutical benefits supplied over 18 months earlier may not be accepted for computer processing. Pharmacists with such claims should contact Medicare Australia.

Reconciliation statements

As mentioned earlier, a pharmacist will receive a PBS reconciliation statement after a claim period has been processed. It provides details of each prescription for each brand of each pharmaceutical benefit item supplied in that claim period.

Reasons for non-payment of any item are coded, with the code numbers explained in the statement.

PBS prescriptions and repeat authorisations not accepted for payment will be returned, with the exception of PBS prescriptions with a dispensed price equal to or less than the patient contribution. Any other items on those PBS prescriptions that have been paid will have been cancelled.

If a PBS prescription was not accepted and can be re-submitted, it must be given a new serial number and included in a subsequent claim period.

If a PBS prescription is finally rejected for payment and a pharmacist is not satisfied with the decision, he/she may apply to the Administrative Appeals Tribunal for a review of that decision.

9. Pricing PBS Prescriptions

Pricing principles

The same pricing principles apply to all PBS prescriptions.

For ready-prepared pharmaceutical benefits, payment is made on the basis of the lowest-priced brand.

For a pharmaceutical benefit not listed as a ready-prepared item, and where a formulation title is stated but no formulary specified, payment is made on the basis of precedence given to formularies by State/Territory legislation.

Prices published in the Schedule do not include any component for goods and services tax (GST).

Further information on pricing PBS prescriptions can be accessed from the booklet titled *Explanation of Current Pricing* on the Medicare Australia's website at www.medicareaustralia.gov.au (PBS publications for Health Care Providers).

Pricing dates

Ready-prepared pharmaceutical benefits are priced on the first day of April, August and December, for items supplied as from each of those days respectively.

Extemporaneously-prepared pharmaceutical benefits and containers are priced on the first day of May each year for items supplied as from the first day of August that year.

Pricing ready-prepared items

For maximum quantities

The price payable for a pharmaceutical benefit is shown in the Schedule against the item. The price is for the maximum quantity available.

If the PBS prescription is for an injectable item and solvent, the price of each is added together, but only one dispensing fee is payable.

The maximum quantity of some pharmaceutical benefits, such as eye drops and oral suspensions, has been determined as a single pack corresponding to the manufacturer's pack. These packs cannot be broken, so if a PBS prescription calls for less, the maximum quantity should be supplied and claimed from Medicare Australia. Packs not to be broken are indicated by a double dagger (‡) in the Schedule.

For lesser quantities

For items where the standard pack is the same as the maximum quantity, and the pack can be broken, the price payable for a lesser quantity is established as follows:

- an amount equal to the dispensing fee, and if applicable the dangerous drug fee, is deducted from the benefit price as shown in the Schedule;
- to this new amount, a wastage percentage is applied, determined from the Wastage Factor Table (see below);
- then the amount equal to the dispensing fee, dangerous drug fee (if applicable), and appropriate container fee, is added.

In no case shall the price for a broken quantity be more than the dispensed price of the Schedule's maximum quantity.

When a standard pack is not the same as the maximum quantity, the price of the pharmaceutical benefit concerned has an asterisk next to it and the standard pack rate is set out in Section 3 of the Schedule (blue pages). The price payable for the quantity supplied is established by:

- applying the appropriate wastage table percentage to the standard pack rate;
- then adding an amount equivalent to the dispensing fee, the dangerous drug fee where applicable, and the appropriate container fee.

In no case shall the supply of a broken quantity, which is less than the item's maximum quantity, cost more than the dispensed price for the maximum quantity.

No container fee is payable when the quantity of pharmaceutical benefit supplied is more than the quantity contained in the standard pack.

Wastage table percentage

The following Wastage Factor Table is used to calculate the price payable for quantities supplied from the standard pack.

Wastage Factor Table

Column A - 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100

Column B - 10, 18, 26, 32, 38, 44, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98, 100

The appropriate wastage table percentage is as follows:

- the percentage of the amount supplied from the amount in the standard pack is determined; and
- where this percentage is the same as a percentage listed in Column A of the table, the percentage used is the figure shown in Column B; or
- where the percentage is not the same as a percentage in Column A, then the nearest upward percentage in Column A applies, and the percentage used is the figure in Column B.

For example, 24 tablets are supplied from a standard pack of 100. Thus 24 per cent of the number contained in the standard pack is supplied. As this percentage does not appear in Column A, the next higher (i.e., 25 per cent) is used. Reading down from 25 per cent to Column B, the wastage table percentage is found to be 38 per cent.

Pricing extemporaneously-prepared items

General

The price payable for supplying the maximum quantity of standard formula preparations is shown in the Standard Formulae List, in Section 4 of this Schedule (green pages).

The following principles apply in determining prices of all pre-priced extemporaneous formulae on the list. They also apply when a pharmacist elects to price extemporaneous PBS prescriptions outside the list, including exceptional PBS prescriptions.

The amount payable is the sum of:

- the recovery price of each ingredient as shown in the Drug Tariff;
- the price of the appropriate container as shown in the price section; and
- a dispensing fee as shown in the price section.

Pricing of ingredients

When the quantity dispensed is not specified in the Drug Tariff, the recovery price is as follows:

(a) determine the basic pricing unit relative to the quantity dispensed by referring to the following table:

<i>Quantity</i>	<i>Basic Pricing Unit</i>
Up to and including 700 mg	100 mg price rate
Over 700 mg and up to and including 1 g	price as if 1 g
Over 1 g and up to and including 7 g	1 g price rate
Over 7 g and up to and including 10 g	price as if 10 g
Over 10 g and up to and including 80 g	10 g price rate
Over 80 g and up to and including 90 g	price as if 80 g
Over 90 g	100 g price rate

(b) find the recovery price of the basic pricing unit by applying the following quantity divisors to the recovery price shown for the ingredient in the Drug Tariff:

- 100 g price is 500 g price divided by 5, or 1 kg price divided by 10
- 10 g price is 100 g price plus 12.5 per cent divided by 10
- 1 g price is 10 g price plus 25 per cent divided by 10
- 100 mg price is 1 g price plus 25 per cent divided by 10

(c) find the recovery price by multiplying the price of the basic pricing unit – as established in (b) – by the fraction that the quantity dispensed bears to the basic pricing unit.

For pricing purposes the quantity is to be taken to the next upward 50 milligrams or 0.05 millilitres.

The minimum recovery price for any ingredient is one cent. In other cases where a fraction of a cent occurs, the price is to be taken to the nearest cent (a half cent being taken up to the next cent).

In no case shall the recovery price for a quantity of an ingredient exceed the recovery price for a greater quantity of that ingredient.

Where liquids are purchased by weight, the recovery price includes the 'Specific Gravity Factor'.

Special pricing provisions apply to drugs marked '(a)' or '(b)' in the Drug Tariff.

For drugs marked '(a)', the pricing rules shown above apply to quantities up to the quantity listed in the Drug Tariff. Greater quantities are priced on a linear basis: the recovery price is ascertained by multiplying the fraction that the quantity dispensed bears to the quantity listed in the Drug Tariff by the price shown for the quantity listed.

Drugs marked '(b)' are packed sterile or are unstable, and all quantities are priced as if whole pack(s) were required. The recovery price is ascertained by multiplying the fraction that the quantity dispensed bears to the quantity listed in the Drug Tariff, taken to the next whole number, by the price shown for the quantity listed.

Pricing PBS prescriptions where extra ingredients are added to a formula

Where the vehicle is liquid and one or more solid ingredients are added, displacement of the liquid by the solid ingredients is disregarded for pricing purposes.

Containers

When a quantity is for more than the container sizes listed in this Schedule, payment will be made as if that quantity had been supplied in the minimum number of containers necessary to supply that quantity.

A double size container is allowed for bulk powders.

Special provisions for extemporaneous PBS prescriptions outside the Standard Formulae List

If a pharmacist elects to price extemporaneous PBS prescriptions outside the Standard Formulae List, there can be no variation for three months. This applies to all extemporaneously-prepared formulae not on the list, and includes both PBS and RPBS prescriptions.

If a pharmacist does not elect to price out these PBS prescriptions, he/she will be paid at an average reimbursement rate.

Under this system, payment is made on the basis of an average 10 g/mL rate applied to the category of preparation concerned, i.e., the price will be determined by multiplying the appropriate 10 g/mL rate by the number of 10 g/mL units supplied and adding container and dispensing fees. For example, an 80 mL mixture would be priced at eight times the average 10 mL rate for mixtures, with container and dispensing fee added.

The average 10 g/mL rate for each type of preparation is calculated monthly. It applies to PBS prescriptions supplied in the following month.

PBS prescriptions ordering a combination of standard formula preparations fall outside the scope of the Standard Formulae List and therefore are subject to this section.

Any variant to a formula included in the list (adding or deleting an ingredient or varying the dose) takes the formula dispensed outside the list.

When an ingredient is added to a standard formula and the recovery price for the standard formula plus additive under the average price system is less than for the standard formula alone, the pharmacist may have the PBS prescription priced as a basic standard formula item.

10. Miscellaneous

References

This Schedule identifies monographs of the British Pharmacopoeia, the British Pharmaceutical Codex, and the Australian Pharmaceutical Formulary and Handbook by the letters BP, BPC and APF respectively. References to all editions of the BPC and to earlier editions of the BP and APF also include the year of publication or the number of the edition.

Standards

Pharmacists can only supply under the PBS medicines which, or whose ingredients, conform to the standards of composition or purity prescribed. These standards are those specified in the *Therapeutic Goods Act 1989*.

Legislation

Copies of the *National Health Act 1953* and the *National Health (Pharmaceutical Benefits) Regulations 1960* are available from Government AusInfo shops in each capital city. The Act and the Regulations may also be accessed through the Attorney-General's Department website at <http://scaleplus.law.gov.au>.

**Therapeutic Index
for
PBS Schedule**

THERAPEUTIC INDEX

ALIMENTARY TRACT AND METABOLISM 77

STOMATOLOGICAL PREPARATIONS 77

Stomatological preparations 77

DRUGS FOR ACID RELATED DISORDERS 77

*Antacids 77**Drugs for peptic ulcer and gastro-oesophageal reflux disease (GORD) 77*

DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS 84

*Belladonna and derivatives, plain 84**Propulsives 84*

ANTIEMETICS AND ANTINAUSEANTS 84

Antiemetics and antinauseants 84

BILE AND LIVER THERAPY 86

Bile therapy 86

LAXATIVES 87

Laxatives 87

ANTIDIARRHEALS, INTESTINAL ANTIINFLAMMATORY/ ANTIINFECTIVE AGENTS 89

*Intestinal antiinfectives 89**Electrolytes with carbohydrates 90**Antipropulsives 90**Intestinal antiinflammatory agents 90*

DIGESTIVES, INCL. ENZYMES 93

Digestives, incl. enzymes 93

DRUGS USED IN DIABETES 93

*Insulins and analogues 93**Oral blood glucose lowering drugs 95*

VITAMINS 99

Vitamin A and D, incl. combinations of the two 99

MINERAL SUPPLEMENTS 99

*Calcium 99**Potassium 99*

ANABOLIC AGENTS FOR SYSTEMIC USE 100

*Anabolic steroids 100***BLOOD AND BLOOD FORMING ORGANS 101**

ANTITHROMBOTIC AGENTS 101

Antithrombotic agents 101

ANTIHEMORRHAGICS 106

Antifibrinolytics 106

ANTIANEMIC PREPARATIONS 106

*Iron preparations 106**Vitamin B₁₂ and folic acid 106*

BLOOD SUBSTITUTES AND PERFUSION SOLUTIONS 107

*Blood and related products 107**I.V. solutions 107***CARDIOVASCULAR SYSTEM 108**

CARDIAC THERAPY 108

*Cardiac glycosides 108**Antiarrhythmics, class I and III 108**Cardiac stimulants excl. cardiac glycosides 110**Vasodilators used in cardiac diseases 110*

ANTIHYPERTENSIVES 112

*Antiadrenergic agents, centrally acting 112**Antiadrenergic agents, peripherally acting 112**Arteriolar smooth muscle, agents acting on 113*

DIURETICS 113

*Low-ceiling diuretics, thiazides 113**Low-ceiling diuretics, excl. thiazides 113**High-ceiling diuretics 114**Potassium-sparing agents 115**Diuretics and potassium-sparing agents in combination 115*

THERAPEUTIC INDEX

- PERIPHERAL VASODILATORS 115
 - Peripheral vasodilators 115*
- BETA BLOCKING AGENTS 116
 - Beta blocking agents 116*
- CALCIUM CHANNEL BLOCKERS 118
 - Selective calcium channel blockers with mainly vascular effects 118*
 - Selective calcium channel blockers with direct cardiac effects 120*
- AGENTS ACTING ON THE RENIN-ANGIOTENSIN SYSTEM 121
 - ACE inhibitors, plain 121*
 - ACE inhibitors, combinations 125*
 - Angiotensin II antagonists, plain 126*
 - Angiotensin II antagonists, combinations 126*
- SERUM LIPID REDUCING AGENTS 127
 - Cholesterol and triglyceride reducers 129*
- DERMATOLOGICALS 134**
 - ANTIFUNGALS FOR DERMATOLOGICAL USE 134
 - Antifungals for systemic use 134*
 - ANTIPSORIATICS 134
 - Antipsoriatics for topical use 134*
 - Antipsoriatics for systemic use 134*
 - ANTIBIOTICS AND CHEMOTHERAPEUTICS FOR DERMATOLOGICAL USE 135
 - Chemotherapeutics for topical use 135*
 - CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS 135
 - Corticosteroids, plain 135*
 - ANTI-ACNE PREPARATIONS 137
 - Anti-acne preparations for systemic use 137*
 - OTHER DERMATOLOGICAL PREPARATIONS 137
 - Other dermatological preparations 137*
- GENITO URINARY SYSTEM AND SEX HORMONES 139**
 - OTHER GYNECOLOGICALS 139
 - Contraceptives for topical use 139*
 - Other gynecologicals 139*
 - SEX HORMONES AND MODULATORS OF THE GENITAL SYSTEM 140
 - Hormonal contraceptives for systemic use 140*
 - Androgens 141*
 - Estrogens 143*
 - Progestogens 145*
 - Progestogens and estrogens in combination 146*
 - Gonadotropins and other ovulation stimulants 148*
 - Antiandrogens 151*
 - Other sex hormones and modulators of the genital system 152*
 - UROLOGICALS 153
 - Other urologicals, incl. antispasmodics 153*
- SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS 154**
 - PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES 154
 - Anterior pituitary lobe hormones and analogues 154*
 - Posterior pituitary lobe hormones 154*
 - Hypothalamic hormones 154*
 - CORTICOSTEROIDS FOR SYSTEMIC USE 155
 - Corticosteroids for systemic use, plain 155*
 - Antiadrenal preparations 157*
 - THYROID THERAPY 157
 - Thyroid preparations 157*
 - Antithyroid preparations 158*
 - PANCREATIC HORMONES 158
 - Glycogenolytic hormones 158*
 - CALCIUM HOMEOSTASIS 158
 - Anti-parathyroid hormones 158*

THERAPEUTIC INDEX**ANTIINFECTIVES FOR SYSTEMIC USE 159****ANTIBACTERIALS FOR SYSTEMIC USE 159**

- Tetracyclines 159*
- Beta-lactam antibacterials, penicillins 161*
- Other beta-lactam antibacterials 166*
- Sulfonamides and trimethoprim 169*
- Macrolides, lincosamides and streptogramins 170*
- Aminoglycoside antibacterials 171*
- Quinolone antibacterials 172*
- Other antibacterials 173*

ANTIMYCOTICS FOR SYSTEMIC USE 175

- Antimycotics for systemic use 175*

ANTIMYCOBACTERIALS 176

- Drugs for treatment of tuberculosis 176*
- Drugs for treatment of lepra 176*

ANTIVIRALS FOR SYSTEMIC USE 177

- Direct acting antivirals 177*

VACCINES 180

- Bacterial vaccines 180*
- Viral vaccines 180*

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS 181**ANTINEOPLASTIC AGENTS 181**

- Alkylating agents 181*
- Antimetabolites 182*
- Plant alkaloids and other natural products 183*
- Cytotoxic antibiotics and related substances 185*
- Other antineoplastic agents 187*

ENDOCRINE THERAPY 189

- Hormones and related agents 189*
- Hormone antagonists and related agents 191*

IMMUNOSTIMULANTS 193

- Cytokines and immunomodulators 193*

IMMUNOSUPPRESSIVE AGENTS 197

- Immunosuppressive agents 197*

MUSCULO-SKELETAL SYSTEM 238**ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS 238**

- Antiinflammatory and antirheumatic products, non-steroids 238*
- Specific antirheumatic agents 242*

MUSCLE RELAXANTS 243

- Muscle relaxants, centrally acting agents 243*
- Muscle relaxants, directly acting agents 243*

ANTIGOUT PREPARATIONS 244

- Antigout preparations 244*

DRUGS FOR TREATMENT OF BONE DISEASES 245

- Drugs affecting bone structure and mineralization 245*

NERVOUS SYSTEM 249**ANALGESICS 249**

- Opioids 249*
- Other analgesics and antipyretics 258*
- Antimigraine preparations 259*

ANTIEPILEPTICS 261

- Antiepileptics 261*

ANTI-PARKINSON DRUGS 266

- Anticholinergic agents 266*
- Dopaminergic agents 266*

THERAPEUTIC INDEX

PSYCHOLEPTICS 268

Antipsychotics 268

Anxiolytics 272

Hypnotics and sedatives 274

PSYCHOANALEPTICS 275

Antidepressants 275

Psychostimulants, agents used for ADHD and nootropics 280

Anti-dementia drugs 282

OTHER NERVOUS SYSTEM DRUGS 286

Parasympathomimetics 286

Drugs used in addictive disorders 287

Other nervous system drugs 288

ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS 289

ANTIPROTOZOALS 289

Agents against amoebiasis and other protozoal diseases 289

Antimalarials 289

ANTHELMINTICS 290

Antinematodal agents 290

ECTOPARASITICIDES, INCL. SCABICIDES, INSECTICIDES AND REPELLENTS 290

Ectoparasiticides, incl. scabicides 290

RESPIRATORY SYSTEM 291

DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES 291

Adrenergics, inhalants 291

Other drugs for obstructive airway diseases, inhalants 294

Adrenergics for systemic use 298

Other systemic drugs for obstructive airway diseases 298

COUGH AND COLD PREPARATIONS 299

Expectorants, excl. combinations with cough suppressants 299

Cough suppressants, excl. combinations with expectorants 299

ANTI-HISTAMINES FOR SYSTEMIC USE 299

Antihistamines for systemic use 299

SENSORY ORGANS 300

OPHTHALMOLOGICALS 300

Anti-infectives 300

Anti-inflammatory agents 301

Antiglaucoma preparations and miotics 302

Mydriatics and cycloplegics 304

Decongestants and antiallergics 304

Other ophthalmologicals 304

OTOLOGICALS 307

Anti-infectives 307

Corticosteroids and anti-infectives in combination 307

OPHTHALMOLOGICAL AND OTOLOGICAL PREPARATIONS 307

Anti-infectives 307

VARIOUS 308

ALLERGENS 308

Allergens 308

ALL OTHER THERAPEUTIC PRODUCTS 308

All other therapeutic products 308

DIAGNOSTIC AGENTS 309

Urine tests 309

Other diagnostic agents 309

GENERAL NUTRIENTS 310

Other nutrients 310

PALLIATIVE CARE BENEFITS**ALIMENTARY TRACT AND METABOLISM 319**

STOMATOLOGICAL PREPARATIONS 319

Stomatological preparations 319

DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS 320

Belladonna and derivatives, plain 320

ANTIEMETICS AND ANTINAUSEANTS 320

Antiemetics and antinauseants 320

LAXATIVES 321

Laxatives 321

MUSCULO-SKELETAL SYSTEM 324

ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS 324

Antiinflammatory and antirheumatic products, non-steroids 324

NERVOUS SYSTEM 329

ANALGESICS 329

Other analgesics and antipyretics 329

ANTIEPILEPTICS 329

Antiepileptics 329

PSYCHOLEPTICS 330

Anxiolytics 330

Hypnotics and sedatives 331

DENTAL BENEFITS**ALIMENTARY TRACT AND METABOLISM 333**

STOMATOLOGICAL PREPARATIONS 333

Stomatological preparations 333

DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS 333

Belladonna and derivatives, plain 333

Propulsives 333

ANTIEMETICS AND ANTINAUSEANTS 333

Antiemetics and antinauseants 333

ANTIDIARRHEALS, INTESTINAL ANTIINFLAMMATORY/ ANTIINFECTIVE AGENTS 334

Intestinal antiinfectives 334

BLOOD AND BLOOD FORMING ORGANS 334

BLOOD SUBSTITUTES AND PERFUSION SOLUTIONS 334

I.V. solutions 334

CARDIOVASCULAR SYSTEM 334

CARDIAC THERAPY 334

Antiarrhythmics, class I and III 334

Cardiac stimulants excl. cardiac glycosides 334

Vasodilators used in cardiac diseases 335

DERMATOLOGICALS 335

CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS 335

Corticosteroids, plain 335

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS 335

CORTICOSTEROIDS FOR SYSTEMIC USE 335

Corticosteroids for systemic use, plain 335

PANCREATIC HORMONES 336

Glycogenolytic hormones 336

THERAPEUTIC INDEX**ANTIINFECTIVES FOR SYSTEMIC USE 336****ANTIBACTERIALS FOR SYSTEMIC USE 336***Tetracyclines 336**Beta-lactam antibacterials, penicillins 337**Other beta-lactam antibacterials 341**Sulfonamides and trimethoprim 344**Macrolides, lincosamides and streptogramins 344**Other antibacterials 345***VACCINES 346***Bacterial vaccines 346***MUSCULO-SKELETAL SYSTEM 346****ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS 346***Antiinflammatory and antirheumatic products, non-steroids 346***NERVOUS SYSTEM 349****ANALGESICS 349***Opioids 349**Other analgesics and antipyretics 354***ANTIEPILEPTICS 355***Antiepileptics 355***ANTI-PARKINSON DRUGS 355***Anticholinergic agents 355***PSYCHOLEPTICS 355***Anxiolytics 355**Hypnotics and sedatives 356***RESPIRATORY SYSTEM 356****DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES 356***Adrenergics for systemic use 356***SENSORY ORGANS 356****OPHTHALMOLOGICALS 356***Antiinfectives 356***VARIOUS 356****ALL OTHER THERAPEUTIC PRODUCTS 356***All other therapeutic products 356*

Section 2

Emergency Drug (Doctor's Bag) Supplies

Special Pharmaceutical Benefits

General Pharmaceutical Benefits

Pharmaceutical Benefits for Palliative Care

Pharmaceutical Benefits for Dental Use

Items Available Under Special Arrangements (s.100)

SYMBOLS USED IN THE SCHEDULE

An arrow (➤) in front of a restriction indicates an additional or amended purpose in this issue.

An asterisk (*) against the dispensed price of a benefit indicates that the manufacturer's pack does not coincide with the maximum quantity.

A double dagger (‡) in the maximum quantity column indicates an item for which the maximum quantity has been specially determined to correspond to the manufacturer's pack and the manufacturer's standard pack should be prescribed and supplied. For any item where a maximum quantity greater than 1 is marked with a double dagger (‡), that maximum quantity should be prescribed and supplied.

A gauge sign (#) against the dispensed price of a benefit indicates that the product is not preconstituted and that an extemporaneously-prepared dispensing fee is included in the dispensed price and, where appropriate, an amount for purified water.

Where a STATE is indicated after a manufacturer's code, that brand may be available only in the State indicated. NSW—(N); Vic—(V); Qld—(Q); SA—(S); WA—(W); Tas—(T).

RESTRICTED BENEFITS

All restricted items are printed in ***bold italics***, with separate headings for authority and non-authority items. In each case these items may be prescribed as pharmaceutical benefits only for use for one of the specified indications. Where more than one indication is specified for an authority required or restricted pharmaceutical benefit, each indication is separated from the preceding indication by a semi-colon and commences on the next line. The drug may be prescribed as a pharmaceutical benefit for a patient who qualifies under any of the specified indications.

A straight line is drawn between entries for different forms and strengths of an item to indicate clearly the different restrictions which apply to these various forms and strengths.

The maximum quantity and/or number of repeats in respect of an item shown in the Schedule may be varied by the Chief Executive Officer of Medicare Australia when approving an Authority Prescription or an Authority to Prescribe. The quantity and number of repeats shown on the authority shall be supplied. (See Explanatory Notes (yellow pages). Payment will be made on the basis of the price shown for that item in the Schedule.

BRAND EQUIVALENCE

'a' located immediately before brand names of a particular strength of an item indicates that the sponsors of these brands have submitted evidence that they have been demonstrated to be bioequivalent or therapeutically equivalent, or that justification for not needing bioequivalence or therapeutic equivalence data has been provided to and accepted by the Department. It would thus be expected that these brands may be interchanged without differences in clinical effect.

For other brands of an item, i.e., those not indicated as above, it is either unknown whether or not they are equivalent, or else the sponsors of these brands have requested that an indication of equivalence NOT be shown.

There may be several reasons for the first situation, such as bioequivalence data not being considered necessary when the products were approved for marketing, or that advice or data have not been forthcoming from sponsors. This does not necessarily suggest a lack of safety or efficacy, but in these circumstances caution should be taken if brands are interchanged.

In the second case, requests not to include an indication of equivalence are meant to prevent brand substitution (even though the products are equivalent) as legislation permitting substitution applies only to brands which are known to be equivalent. This situation applies only to co-marketed products where there is no brand premium and thus no implication for payment by patients.

'b' attached to brand names indicates that these brands are also equivalent, but that it is not known if there is equivalence between brands marked 'a' and brands marked 'b'.

BRAND PREMIUM POLICY

The Brand Premium Policy was introduced on 1 December 1990 to increase price competition by allowing pharmaceutical manufacturers to set their own price on multi-branded items listed on the Pharmaceutical Benefits Scheme and to encourage the development of the generic pharmaceutical industry in Australia.

The policy does this by increasing prescribers' and patients' consciousness about the price of drugs. In effect, it makes both groups question whether it is necessary for the patient to pay more for the drugs when a cheaper brand is available. The policy also allows companies to establish prices taking into account competition and consumer acceptance.

The policy operates where there is more than one brand of a particular drug available through the Pharmaceutical Benefits Scheme and where the brands are therapeutically interchangeable. Due to this, the policy mainly applies to out of patent drugs.

Basically the policy operates by:

- the Australian Government subsidising a drug to the level of the lowest priced brand (except in those instances where the lowest priced brand has, as part of its price, a therapeutic group premium);
- suppliers of other brands of that drug being able to set a price above the price charged by the supplier(s) of the lowest priced brand(s); and
- the patient paying the brand premium which is the price difference between the lowest price brand and the brand prescribed.

If a prescription is written generically or for the lowest priced brand, and the lowest priced brand is supplied, there is no brand premium payable.

'B' located immediately before an amount in the premium column indicates a brand premium which applies to that particular brand of the item.

If a brand of a drug which is subject to a therapeutic group premium also has a brand premium, there will be two amounts shown on separate lines in the premium column, prefixed by 'T' and 'B' respectively.

If a brand of a drug which is subject to a special patient contribution also has a brand premium, there will be two amounts shown on separate lines in the premium column, prefixed by 'S' and 'B' respectively.

THERAPEUTIC GROUP PREMIUM POLICY

The Therapeutic Group Premium Policy was introduced on 1 February 1998 as an extension of the Brand Premium Policy to encourage greater competition between manufacturers of drugs and to make doctors and patients more aware of the costs of medicines.

The Therapeutic Group Premium policy applies within narrowly defined therapeutic sub-groups where the drugs concerned are of similar safety, efficacy and health outcomes.

Basically the policy operates by:

- the Australian Government subsidising drugs within a defined therapeutic sub-group to the level of the lowest priced drug in the sub-group;
- suppliers of other drugs within that sub-group being able to set prices above the price charged by the supplier(s) of the lowest priced drug; and
- the patient paying the therapeutic group premium which is the price difference between the lowest price drug and the drug prescribed.

'T' located immediately before an amount in the premium column indicates a therapeutic group premium which applies to that particular item.

If a brand of a drug which is subject to a therapeutic group premium also has a brand premium, there will be two amounts shown on separate lines in the premium column, prefixed by 'T' and 'B' respectively.

The success of the Government in controlling prices of products supplied through the Pharmaceutical Benefits Scheme has often been criticised by the pharmaceutical industry. Under both the Brand Premium Policy and the Therapeutic Group Premium Policy, suppliers of multi-branded items and therapeutically similar drugs are able to set their own prices at a level that they think the market will bear. At the same time, the prescriber and the patient can decide whether it is necessary to pay more for a particular brand or drug when a cheaper one is available and is therapeutically interchangeable.

The brand premium or therapeutic group premium does not count toward the patient's safety net.

It should be noted that the brand premium or therapeutic group premium is not a Government charge or revenue. The premium arises from the manufacturer's price and the majority goes to the manufacturer with wholesalers and pharmacists receiving a small percentage.

EMERGENCY DRUG (DOCTOR'S BAG) SUPPLIES

Code	Name, Manner of Administration and Form	Max. Qty	Dispensed Price for Max.Qty \$	Proprietary Name and Manufacturer	
3451P	ADRENALINE Injection 1 mg in 1 mL (1 in 1,000)	5	14.76	AP	
3453R	ATROPINE SULFATE Injection 600 micrograms in 1 mL	10	12.20	AP	
3457Y	BENZTROPINE MESYLATE Injection 2 mg in 2 mL	5	20.99	Cogentin	MK
3486L	BENZYLPENICILLIN Powder for injection 600 mg	10	*22.25	BenPen	CS
3485K	<i>or</i> PROCAINE PENICILLIN Injection 1.5 g	5	52.49	Cilicaine	SI
3487M	BENZYLPENICILLIN Powder for injection 3 g	1	11.11	BenPen	CS
3455W	CHLORPROMAZINE HYDROCHLORIDE Injection 50 mg in 2 mL	10	13.99	Largactil	AV
3456X	<i>or</i> HALOPERIDOL Injection 5 mg in 1 mL	10	15.86	Serenace	SI
3472R	DEXAMETHASONE SODIUM PHOSPHATE Injection equivalent to 4 mg dexamethasone phosphate in 1 mL	5	13.85	MX	
3470P	<i>or</i> HYDROCORTISONE SODIUM SUCCINATE Injection equivalent to 100 mg hydrocortisone with 2 mL solvent	2	*15.55	Solu-Cortef	PH
3471Q	<i>or</i> HYDROCORTISONE SODIUM SUCCINATE Injection equivalent to 250 mg hydrocortisone with 2 mL solvent	1	12.27	Solu-Cortef	PH
3458B	DIAZEPAM Injection 10 mg in 2 mL	5	10.55	MX	
3460D	DIHYDROERGOTAMINE MESYLATE Injection 1 mg in 1 mL	5	15.27	Dihydergot	NV
3463G	DIPHThERIA and TETANUS VACCINE, ADSORBED, DILUTED FOR ADULT USE Injection 0.5 mL in pre-filled syringe	15	*90.70	ADT	CS
3466K	FRUSEMIDE Injection 20 mg in 2 mL	5	10.11	^a Frusehexal ^a Lasix	HX AV
3467L	GLUCAGON HYDROCHLORIDE Injection set containing 1 mg (1 i.u.) and 1 mL solvent in disposable syringe	1	40.12	GlucaGen Hypokit	NO
3475X	GLYCERYL TRINITRATE Buccal/sublingual spray (pump pack) 400 micrograms per dose (200 doses)	‡1	18.30	Nitrolingual Pumpspray	AV

EMERGENCY DRUG (DOCTOR'S BAG) SUPPLIES

Code	Name, Manner of administration and form	Max. Qty	Dispensed Price for Max.Qty \$	Proprietary Name and Manufacturer	
3474W	LIGNOCAINE HYDROCHLORIDE Injection 100 mg in 5 mL	5	35.31	PF	
3476Y <i>or</i>	METOCLOPRAMIDE HYDROCHLORIDE Injection 10 mg in 2 mL <i>or</i>	10	11.24	Maxolon	VT
3477B	PROCHLORPERAZINE Injection 12.5 mg in 1 mL	10	14.17	Stemetil	AV
3479D <i>or</i>	MORPHINE SULFATE Injection 15 mg in 1 mL <i>or</i>	5	11.94	MX SI	
3480E	MORPHINE SULFATE Injection 30 mg in 1 mL	5	13.92	MX	
3481F <i>or</i>	NALOXONE HYDROCHLORIDE Injection 800 micrograms in 2 mL <i>or</i>	5	*96.70	Naloxone Min-I-Jet	CS
3482G	NALOXONE HYDROCHLORIDE Injection 2 mg in 5 mL	2	*63.55	Naloxone Min-I-Jet	CS
3483H <i>or</i>	PETHIDINE HYDROCHLORIDE Injection 100 mg in 2 mL <i>or</i>	5	11.31	MX	
3484J	TRAMADOL HYDROCHLORIDE Injection 100 mg in 2 mL	5	11.11	Tramal 100	CS
3488N	PROMETHAZINE HYDROCHLORIDE Injection 50 mg in 2 mL	10	*18.29	MX	
3496B	SALBUTAMOL SULFATE Nebuliser solution single dose units 2.5 mg (base) in 2.5 mL, 30	‡1	14.18	^a Asmol 2.5 uni-dose ^a Butamol 2.5 ^a Chem mart Salbutamol ^a GenRx Salbutamol ^a Terry White Chemists Salbutamol ^a PU 15.28 ^a Ventolin Nebules	AF AW CH GX TW GK
<i>or</i>	<i>or</i>				
3495Y	SALBUTAMOL SULFATE Oral pressurised inhalation 100 micrograms (base) per dose (200 doses), CFC-free formulation	‡1	10.56 10.87 11.06	^a Asmol CFC-free ^a Epaq ^a Airomir ^a Ventolin CFC-free	AL AW MM GK

EMERGENCY DRUG (DOCTOR'S BAG) SUPPLIES

Code	Name, Manner of administration and form	Max. Qty	Dispensed Price for Max.Qty \$	Proprietary Name and Manufacturer	
3497C	SALBUTAMOL SULFATE Nebuliser solution single dose units 5 mg (base) in 2.5 mL, 30	‡1	14.72	^a Asmol 5 uni-dose	AF
				^a Butamol 5	AW
				^a Chem mart Salbutamol	CH
				^a GenRx Salbutamol	GX
				^a Terry White Chemists Salbutamol	TW
				^a Ventolin Nebules	GK
			15.82		
3491R	TERBUTALINE SULFATE Injection 500 micrograms in 1 mL	5	22.54	Bricanyl	AP
3492T	TETANUS VACCINE, ADSORBED Injection 0.5 mL in pre-filled syringe	5	*25.15	Tet-Tox	CS
3494X	VERAPAMIL HYDROCHLORIDE Injection 5 mg in 2 mL	5	10.64	Isoptin	AB

SPECIAL PHARMACEUTICAL BENEFITS

The special patient contribution is payable by all patients in addition to the relevant patient contribution for concessional and general patients. Other than for bleomycin sulfate, exemptions on medical grounds are available. For eligible veterans under RPBS provisions, see RPBS EXPLANATORY NOTES, paragraph 28.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Reimburse- ment Price for Max. Qty \$	Total Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
<u>BLEOMYCIN SULFATE</u>								
<u>Restricted benefit</u>								
<i>Germ cell neoplasms; Lymphoma.</i>								
2315W	<i>Powder for injection 15,000 i.u.</i>	10	..	<i>\$460.80</i>	<i>*516.95</i>	<i>*977.75</i>	<i>28.60</i>	<i>MX</i>
				<i>\$460.80</i>	<i>516.95</i>	<i>977.75</i>	<i>28.60</i>	<i>a Blenamax SI</i>
				<i>\$460.80</i>	<i>516.95</i>	<i>1057.23</i>	<i>28.60</i>	<i>a Blenoxane BQ</i>
				<i>B79.48</i>				
<u>ESCITALOPRAM OXALATE</u>								
<u>Restricted benefit</u>								
<i>Major depressive disorders.</i>								
8700X	<i>Tablet 10 mg (base)</i>	28	5	<i>\$3.31</i>	<i>27.94</i>	<i>31.25</i>	<i>28.60</i>	<i>Lexapro LU</i>
8701Y	<i>Tablet 20 mg (base)</i>	28	5	<i>\$5.53</i>	<i>43.39</i>	<i>48.92</i>	<i>28.60</i>	<i>Lexapro LU</i>
8849R	<i>Oral solution 10 mg (base) per mL, 28 mL</i>	1	5	<i>\$4.46</i>	<i>36.06</i>	<i>40.52</i>	<i>28.60</i>	<i>Lexapro LU</i>
<u>Authority required</u>								
<i>Major depressive disorders, where:</i>								
<i>(1) adverse events have occurred with other suitable PBS-listed drugs; or</i>								
<i>(2) drug interactions have occurred with other suitable PBS-listed drugs; or</i>								
<i>(3) drug interactions are expected to occur with other suitable PBS-listed drugs; or</i>								
<i>(4) transfer to another suitable PBS-listed drug would cause patient confusion resulting in problems with compliance; or</i>								
<i>(5) transfer to another suitable PBS-listed drug is likely to result in adverse clinical consequences.</i>								
9711D	<i>Tablet 10 mg (base)</i>	28	5	<i>..</i>	<i>31.25</i>	<i>31.25</i>	<i>28.60</i>	<i>Lexapro LU</i>
9712E	<i>Tablet 20 mg (base)</i>	28	5	<i>..</i>	<i>48.92</i>	<i>48.92</i>	<i>28.60</i>	<i>Lexapro LU</i>
9700M	<i>Oral solution 10 mg (base) per mL, 28 mL</i>	1	5	<i>..</i>	<i>40.52</i>	<i>40.52</i>	<i>28.60</i>	<i>Lexapro LU</i>

SPECIAL PHARMACEUTICAL BENEFITS

The special patient contribution is payable by all patients in addition to the relevant patient contribution for concessional and general patients. Other than for bleomycin sulfate, exemptions on medical grounds are available. For eligible veterans under RPBS provisions, see RPBS EXPLANATORY NOTES, paragraph 28.

KPBS EXPLANATORY NOTES, paragraph 26.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Reimbursement Price for Max. Qty \$	Total Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
LEVETIRACETAM									
<u>Authority required</u>									
<i>Treatment of partial epileptic seizures which are not controlled satisfactorily by other anti-epileptic drugs.</i>									
8654L	Tablet 250 mg	60	5	\$7.38	56.42	63.80	28.60	Keppra	UC
8655M	Tablet 500 mg	60	5	\$12.31	90.91	103.22	28.60	Keppra	UC
8656N	Tablet 1 g	60	5	\$20.50	148.30	168.80	28.60	Keppra	UC
<u>Authority required</u>									
<i>Treatment of partial epileptic seizures which are not controlled satisfactorily by other anti-epileptic drugs, and where:</i>									
<i>(1) adverse events have occurred with other suitable PBS-listed drugs; or</i>									
<i>(2) drug interactions have occurred with other suitable PBS-listed drugs; or</i>									
<i>(3) drug interactions are expected to occur with other suitable PBS-listed drugs; or</i>									
<i>(4) transfer to another suitable PBS-listed drug would cause patient confusion resulting in problems with compliance; or</i>									
<i>(5) transfer to another suitable PBS-listed drug is likely to result in adverse clinical consequences.</i>									
9708Y	Tablet 250 mg	60	5	..	63.80	63.80	28.60	Keppra	UC
9709B	Tablet 500 mg	60	5	..	103.22	103.22	28.60	Keppra	UC
9710C	Tablet 1 g	60	5	..	168.80	168.80	28.60	Keppra	UC
PEMETREXED DISODIUM									
<u>Authority required</u>									
<i>Locally advanced or metastatic non-small cell lung cancer, after prior platinum-based chemotherapy.</i>									
8809P	Powder for I.V. infusion 500 mg (base)	2	2	\$460.76	*3230.11	*3690.87	28.60	Alimta	LY
<u>Authority required</u>									
<i>Locally advanced or metastatic non-small cell lung cancer, after prior platinum-based chemotherapy, where:</i>									
<i>(1) treatment with paclitaxel or docetaxel is contraindicated; or</i>									
<i>(2) intolerance to treatment with either docetaxel or paclitaxel has developed; or</i>									
<i>(3) treatment with either docetaxel or paclitaxel has been unsuccessful; or</i>									
<i>(4) transfer to docetaxel or paclitaxel is likely to result in adverse clinical consequences.</i>									
9713F	Powder for I.V. infusion 500 mg (base)	2	2	..	*3690.87	*3690.87	28.60	Alimta	LY

SPECIAL PHARMACEUTICAL BENEFITS

The special patient contribution is payable by all patients in addition to the relevant patient contribution for concessional and general patients. Other than for bleomycin sulfate, exemptions on medical grounds are available. For eligible veterans under RPBS provisions, see RPBS EXPLANATORY NOTES, paragraph 28.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium \$	Reimburse- ment Price for Max.Qty \$	Total Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
TOPIRAMATE									
<u>Authority required</u>									
<i>Treatment of partial epileptic seizures, primary generalised tonic-clonic epileptic seizures and seizures of the Lennox-Gastaut syndrome, which are not controlled satisfactorily by other anti-epileptic drugs.</i>									
8163P	Tablet 25 mg	60	5	\$2.40	39.20	41.60	28.60	Topamax	JC
8164Q	Tablet 50 mg	60	5	\$2.40	62.19	64.59	28.60	Topamax	JC
8165R	Tablet 100 mg	60	5	\$2.70	100.45	103.15	28.60	Topamax	JC
8166T	Tablet 200 mg	60	5	\$2.69	165.53	168.22	28.60	Topamax	JC
8371N	Capsule 15 mg	60	5	\$2.40	29.56	31.96	28.60	Topamax Sprinkle	JC
8372P	Capsule 25 mg	60	5	\$2.40	39.20	41.60	28.60	Topamax Sprinkle	JC
8520K	Capsule 50 mg	60	5	\$2.40	62.19	64.59	28.60	Topamax Sprinkle	JC
<u>Authority required</u>									
<i>Treatment of partial epileptic seizures, primary generalised tonic-clonic epileptic seizures and seizures of the Lennox-Gastaut syndrome, which are not controlled satisfactorily by other anti-epileptic drugs, and where:</i>									
<i>(1) adverse events have occurred with other suitable PBS-listed drugs; or</i>									
<i>(2) drug interactions have occurred with other suitable PBS-listed drugs; or</i>									
<i>(3) drug interactions are expected to occur with other suitable PBS-listed drugs; or</i>									
<i>(4) transfer to another suitable PBS-listed drug would cause patient confusion resulting in problems with compliance; or</i>									
<i>(5) transfer to another suitable PBS-listed drug is likely to result in adverse clinical consequences.</i>									
9704R	Tablet 25 mg	60	5	..	41.60	41.60	28.60	Topamax	JC
9705T	Tablet 50 mg	60	5	..	64.59	64.59	28.60	Topamax	JC
9706W	Tablet 100 mg	60	5	..	103.15	103.15	28.60	Topamax	JC
9707X	Tablet 200 mg	60	5	..	168.22	168.22	28.60	Topamax	JC
9701N	Capsule 15 mg	60	5	..	31.96	31.96	28.60	Topamax Sprinkle	JC
9702P	Capsule 25 mg	60	5	..	41.60	41.60	28.60	Topamax Sprinkle	JC
9703Q	Capsule 50 mg	60	5	..	64.59	64.59	28.60	Topamax Sprinkle	JC

ALIMENTARY TRACT AND METABOLISM

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
STOMATOLOGICAL PREPARATIONS								
Stomatological preparations								
• Antiinfectives and antiseptics for local oral treatment								
AMPHOTERICIN								
2931G	Lozenge 10 mg	20	1	..	8.60	9.57	Fungilin	BQ
NYSTATIN								
3033P	Oral suspension 100,000 units per mL, 24 mL	‡1	1	..	9.13	10.10	Mycostatin Nilstat	BQ SI
• Other agents for local oral treatment								
BENZYLAMINE HYDROCHLORIDE								
Restricted benefit								
Radiation induced mucositis.								
1121B	Mouth and throat rinse 22.5 mg per 15 mL, 500 mL	‡1	1	..	18.54	19.51	Difflam	MM
DRUGS FOR ACID RELATED DISORDERS								
Antacids								
• Combinations and complexes of aluminium, calcium and magnesium compounds								
ALUMINIUM HYDROXIDE with MAGNESIUM HYDROXIDE								
2576N	Tablet 200 mg-200 mg	200	5	.. B0.40	*12.69 *13.09	13.66 13.66	^a Gelusil ^a Mylanta P	WW PC
2157M	Oral suspension 200 mg-200 mg per 5 mL, 500 mL	2	5	.. B0.32	*12.69 *13.01	13.66 13.66	^a Gelusil ^a Mylanta P	WW PC
ALUMINIUM HYDROXIDE with MAGNESIUM TRISILICATE and MAGNESIUM HYDROXIDE								
2159P	Oral suspension 250 mg-120 mg- 120 mg per 5 mL, 500 mL	2	5	..	*12.69	13.66	Gastrogel	FM
Drugs for peptic ulcer and gastro-oesophageal reflux disease (GORD)								
• H₂-receptor antagonists								
NOTE:								
The base-priced drugs in this therapeutic group are cimetidine (except cimetidine effervescent tablet 800 mg (as hydrochloride)), famotidine, nizatidine and ranitidine hydrochloride (except ranitidine hydrochloride effervescent tablet 150 mg (base) and syrup 150 mg (base) per 10 mL, 300 mL).								
CIMETIDINE								
NOTE:								
<i>Helicobacter pylori</i> eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.								
1157X	Tablet 200 mg	120	5	.. B2.49	27.29 29.78	28.26 28.26	^a Cimehexal ^a Magicul 200 ^a Tagamet	HX AF GK

continued ➡

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CIMETIDINE—cont.								
1158Y	Tablet 400 mg	60	5	..	27.29	28.26	^a Cimehexal ^a GenRx Cimetidine	HX GX
				^B 2.49	29.78	28.26	^a Magicul 400 ^a Tagamet	AF GK
1159B	Tablet 800 mg	30	5	..	27.29	28.26	^a Cimehexal ^a GenRx Cimetidine	HX GX
				^B 4.96	32.25	28.26	^a Magicul 800 ^a Tagamet	AF GK
1156W	Effervescent tablet 800 mg (as hydrochloride)	30	5	^T 7.01	34.30	28.26	Tagamet 800 Express	GK

CIMETIDINE**NOTE:**

Helicobacter pylori eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.

Authority required

Adverse effects occurring with all of the base-priced drugs;

Drug interactions occurring with all of the base-priced drugs;

Drug interactions expected to occur with all of the base-priced drugs;

Transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.

8901L	Effervescent tablet 800 mg (as hydrochloride)	30	5	..	34.30	28.60	Tagamet 800 Express	GK

FAMOTIDINE**NOTE:**

Helicobacter pylori eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.

2487X	Tablet 20 mg	60	5	..	21.32	22.29	^a Ausfam 20 ^a Chem mart Famotidine	AW CH
							^a Famohexal ^a GenRx Famotidine	HX GX
							^a Pamacid 20 ^a Pepzan ^a Terry White Chemists Famotidine	AF DP TW
				^B 5.41	26.73	22.29	^a Pepcidine M	MK

continued ➡

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
2488Y	FAMOTIDINE—cont. Tablet 40 mg	30	5	..	21.32	22.29	^a Ausfam 40 ^a Chem mart Famotidine ^a Famohexal ^a GenRx Famotidine ^a Pamacid 40 ^a Pepzan ^a Terry White Chemists Famotidine ^a Pepcidine	AW CH HX GX AF DP TW MK
NIZATIDINE								
<u>NOTE:</u> <i>Helicobacter pylori</i> eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.								
1505F	Capsule 150 mg	60	5	.. ^B 2.60	23.05 25.65	24.02 24.02	^a Nizac ^a Tazac	LN AS
1504E	Capsule 300 mg	30	5	.. ^B 2.60	23.05 25.65	24.02 24.02	^a Nizac ^a Tazac	LN AS
RANITIDINE HYDROCHLORIDE								
<u>NOTE:</u> <i>Helicobacter pylori</i> eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.								
1978D	Tablet 150 mg (base)	60	5	..	22.01	22.98	^a Ausran ^a Chem mart Ranitidine ^a GenRx Ranitidine ^a Rani 2 ^a Ranihexal ^a Ranoxyl ^a Terry White Chemists Ranitidine ^a Zantac	SI CH GX AF HX DP TW GK
1937Y	Effervescent tablet 150 mg (base)	60	5	^T 2.14	[*] 24.17	23.00	Zantac	GK

continued ➡

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
RANITIDINE HYDROCHLORIDE—cont.								
1977C	Tablet 300 mg (base)	30	5	..	22.01	22.98	^a Ausran ^a Chem mart Ranitidine ^a GenRx Ranitidine ^a Rani 2 ^a Ranihexal ^a Ranoxyl ^a Terry White Chemists Ranitidine	SI CH GX AF HX DP TW GK
8162N	Syrup 150 mg (base) per 10 mL, 300 mL	2	5	^B 2.17 ^T 2.14	24.18 *24.17	22.98 23.00	^a Zantac Zantac Syrup	GK GK

RANITIDINE HYDROCHLORIDE**NOTE:**

Helicobacter pylori eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.

Authority required

Adverse effects occurring with all of the base-priced drugs;

Drug interactions occurring with all of the base-priced drugs;

Drug interactions expected to occur with all of the base-priced drugs;

Transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.

8903N	Effervescent tablet 150 mg (base)	60	5	..	*24.17	25.14	Zantac	GK
8905Q	Syrup 150 mg (base) per 10 mL, 300 mL	2	5	..	*24.17	25.14	Zantac Syrup	GK

• **Prostaglandins****MISOPROSTOL****CAUTION:**

Misoprostol is a prostaglandin analogue. It should not be used in pregnant women.

Authority required

Reduction in the incidence of gastrointestinal complications in patients who have a history of peptic ulcer disease and in whom NSAID therapy is essential;

Duodenal ulcer (including pyloric and stomal ulcers), proven by current or prior x-ray, endoscopy or surgery. The authority application must state the date and the method by which the ulcer was proven;

Gastric ulcer, proven by x-ray, endoscopy or surgery within the previous 2 years. The authority application must state the date and the method by which the ulcer was proven.

1648R	Tablet 200 micrograms	120	2	..	51.59	28.60	Cytotec	PH

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Proton pump inhibitors ESOMEPRAZOLE MAGNESIUM TRIHYDRATE Restricted benefit <i>Healing of gastro-oesophageal reflux disease.</i> NOTE: <i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8601Q	Tablet (enteric coated), equivalent to 40 mg esomeprazole	30	1	..	75.35	28.60	Nexium	AP
Restricted benefit <i>Maintenance of healed gastro-oesophageal reflux disease.</i> NOTE: <i>No applications for increased maximum quantities will be authorised.</i>								
8600P	Tablet (enteric coated), equivalent to 20 mg esomeprazole	30	5	..	46.28	28.60	Nexium	AP
LANSOPRAZOLE Restricted benefit <i>Initial treatment of peptic ulcer.</i> NOTE: <i>Helicobacter pylori eradication therapy should be considered.</i>								
2240X	Capsule 30 mg	30	1	..	45.19	28.60	Zoton	WY
8528W	Sachet containing granules for oral suspension, 30 mg per sachet	28	1	..	42.50	28.60	Zoton	WY
NOTE: <i>No applications for increased repeats will be authorised.</i>								
Restricted benefit <i>Gastro-oesophageal reflux disease; Scleroderma oesophagus.</i>								
8198L	Capsule 15 mg	30	5	..	28.58	28.60	Zoton	WY
2241Y	Capsule 30 mg	30	5	..	45.19	28.60	Zoton	WY
8529X	Sachet containing granules for oral suspension, 30 mg per sachet	28	5	..	42.50	28.60	Zoton	WY

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
OMEPRAZOLE								
<u>Restricted benefit</u>								
<i>Initial treatment of peptic ulcer.</i>								
<u>NOTE:</u>								
<i>Helicobacter pylori eradication therapy should be considered.</i>								
8331L	Tablet 20 mg	30	1	..	39.26	28.60	^a Acimax Tablets	AL
							^a Meprazol	HX
				^B 2.75	42.01	28.60	^a Losec Tablets	AP
1326T	Capsule 20 mg	30	1	..	39.26	28.60	Probitor	SZ
<u>NOTE:</u>								
<i>No applications for increased repeats will be authorised.</i>								
<u>Restricted benefit</u>								
<i>Gastro-oesophageal reflux disease;</i>								
<i>Scleroderma oesophagus;</i>								
<i>Zollinger-Ellison syndrome.</i>								
8332M	Tablet 10 mg	30	5	..	29.09	28.60	Losec Tablets	AP
8333N	Tablet 20 mg	30	5	..	39.26	28.60	^a Acimax Tablets	AL
							^a Meprazol	HX
				^B 2.75	42.01	28.60	^a Losec Tablets	AP
1327W	Capsule 20 mg	30	5	..	39.26	28.60	Probitor	SZ
PANTOPRAZOLE SODIUM SESQUIHYDRATE								
<u>Restricted benefit</u>								
<i>Initial treatment of peptic ulcer.</i>								
<u>NOTE:</u>								
<i>Helicobacter pylori eradication therapy should be considered.</i>								
8007K	Tablet (enteric coated), equivalent to 40 mg pantoprazole	30	2	..	46.51	28.60	Somac	PH
<u>NOTE:</u>								
<i>No applications for increased repeats will be authorised.</i>								
<u>Restricted benefit</u>								
<i>Gastro-oesophageal reflux disease.</i>								
8399C	Tablet (enteric coated), equivalent to 20 mg pantoprazole	30	5	..	27.26	28.23	Somac	PH
8008L	Tablet (enteric coated), equivalent to 40 mg pantoprazole	30	5	..	46.51	28.60	Somac	PH

continued ➡

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PANTOPRAZOLE SODIUM SESQUIHYDRATE—cont.								
<u>Restricted benefit</u>								
<i>Scleroderma oesophagus; Zollinger-Ellison syndrome.</i>								
8008L	Tablet (enteric coated), equivalent to 40 mg pantoprazole	30	5	..	46.51	28.60	Somac	PH
RABEPRAZOLE SODIUM								
<u>Restricted benefit</u>								
<i>Initial treatment of peptic ulcer.</i>								
NOTE:								
<i>Helicobacter pylori eradication therapy should be considered.</i>								
8509W	Tablet 20 mg (enteric coated)	30	2	..	46.50	28.60	Pariet	JC
NOTE:								
<i>No applications for increased repeats will be authorised.</i>								
<u>Restricted benefit</u>								
<i>Gastro-oesophageal reflux disease; Scleroderma oesophagus.</i>								
8507R	Tablet 10 mg (enteric coated)	28	5	..	27.69	28.60	Pariet	JC
8508T	Tablet 20 mg (enteric coated)	30	5	..	46.50	28.60	Pariet	JC
• Combinations for eradication of Helicobacter pylori								
ESOMEPRAZOLE MAGNESIUM TRIHYDRATE and CLARITHROMYCIN and AMOXYCILLIN								
<u>Restricted benefit</u>								
<i>Eradication of Helicobacter pylori associated with peptic ulcer disease.</i>								
8738X	Pack containing 14 tablets (enteric coated) equivalent to 20 mg esomeprazole, 14 tablets clarithromycin 500 mg and 28 capsules amoxicillin 500 mg	£1	..	..	97.88	28.60	Nexium Hp7	AP
OMEPRAZOLE and CLARITHROMYCIN and AMOXYCILLIN								
<u>Restricted benefit</u>								
<i>Eradication of Helicobacter pylori associated with peptic ulcer disease.</i>								
8272J	Pack containing 14 capsules omeprazole 20 mg, 14 tablets clarithromycin 500 mg and 28 capsules amoxicillin 500 mg	£1	..	..	98.12	28.60	Klacid Hp 7	AB
• Other drugs for peptic ulcer and gastro-oesophageal reflux disease (GORD)								
SODIUM ALGINATE with CALCIUM CARBONATE and SODIUM BICARBONATE								
2014B	Oral liquid 1 g-320 mg-534 mg in 20 mL, 500 mL	2	5	..	*12.91	13.88	Gaviscon P	RC

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
SUCRALFATE								
2055E	Tablet equivalent to 1 g anhydrous sucralfate	120	2	..	22.89	23.86	^a Ulcyte	AF
				^B 2.21	25.10	23.86	^a Carafate	AS

DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS

Belladonna and derivatives, plain• **Belladonna alkaloids, tertiary amines**

ATROPINE SULFATE

1089H	Injection 600 micrograms in 1 mL	10	1	..	12.20	13.17	AP	

Propulsives• **Propulsives**

DOMPERIDONE

1347X	Tablet 10 mg	25	..	..	7.19	8.16	Motilium	JC

METOCLOPRAMIDE HYDROCHLORIDE

1207M	Tablet 10 mg	25	..	..	6.36	7.33	Pramin	AF
				^B 2.98	9.34	7.33	Maxolon	VT
1205K	Syrup 5 mg per 5 mL, 100 mL	‡1	..	..	7.35	8.32	Maxolon	VT
1206L	Injection 10 mg in 2 mL	10	..	..	11.24	12.21	Maxolon	VT

ANTIEMETICS AND ANTINAUSEANTS

Antiemetics and antinauseants• **Serotonin (5HT₃) antagonists**

DOLASETRON MESYLATE

Restricted benefit

Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy.

8191D	Tablet 200 mg	2	..	..	56.45	28.60	Anzemet	AV
8192E	I.V. injection 100 mg in 5 mL	1	..	..	30.60	28.60	Anzemet	AV

GRANISETRON HYDROCHLORIDE

Restricted benefit

Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy.

8728J	Tablet 2 mg (base)	2	..	..	*66.83	28.60	Kytril	MX
8729K	Concentrated injection 3 mg (base) in 3 mL	1	..	..	40.25	28.60	Kytril	MX

Authority required

Management of nausea and vomiting associated with radiotherapy being used to treat malignancy.

8873B	Tablet 2 mg (base)	5	1	..	159.96	28.60	Kytril	MX
8730L	Concentrated injection 3 mg (base) in 3 mL	1	..	..	40.25	28.60	Kytril	MX

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ONDANSETRON								
<u>Restricted benefit</u>								
<i>Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy.</i>								
8224W	Tablet 4 mg	4	..	..	44.38	28.60	Zofran	GK
8225X	Tablet 8 mg	4	..	..	66.83	28.60	Zofran	GK
8410P	Wafer 4 mg	4	..	..	44.38	28.60	Zofran Zydys	GK
8411Q	Wafer 8 mg	4	..	..	66.83	28.60	Zofran Zydys	GK
8226Y	I.V. injection 4 mg in 2 mL	1	..	..	27.10	28.07	Zofran	GK
8227B	I.V. injection 8 mg in 4 mL	1	..	..	40.25	28.60	Zofran	GK
<u>Authority required</u>								
<i>Management of nausea and vomiting associated with radiotherapy being used to treat malignancy.</i>								
1594X	Tablet 4 mg	10	1	..	103.83	28.60	Zofran	GK
1595Y	Tablet 8 mg	10	1	..	159.96	28.60	Zofran	GK
8412R	Wafer 4 mg	10	1	..	103.83	28.60	Zofran Zydys	GK
8413T	Wafer 8 mg	10	1	..	159.96	28.60	Zofran Zydys	GK
8233H	Syrup 4 mg per 5 mL, 50 mL	‡1	1	..	103.83	28.60	Zofran syrup 50 mL	GK
1596B	I.V. injection 4 mg in 2 mL	1	..	..	27.10	28.07	Zofran	GK
1597C	I.V. injection 8 mg in 4 mL	1	..	..	40.25	28.60	Zofran	GK
TROPISETRON HYDROCHLORIDE								
<u>Restricted benefit</u>								
<i>Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy.</i>								
2745L	Capsule 5 mg (base)	2	..	..	56.45	28.60	Navoban	NV
2746M	I.V. injection 5 mg (base) in 5 mL	1	..	..	30.60	28.60	Navoban	NV

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
• Other antiemetics APREPITANT NOTE: <i>Aprepitant is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.</i> Authority required <i>Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy, in combination with a 5HT3 antagonist and dexamethasone, where any 1 of the following chemotherapy agents are to be administered:</i> <i>(a) altretamine;</i> <i>(b) carmustine;</i> <i>(c) cisplatin when a single dose constitutes a cycle of chemotherapy;</i> <i>(d) cyclophosphamide at a dose of 1500 mg per square metre per day or greater;</i> <i>(e) dacarbazine;</i> <i>(f) procarbazine when a single dose constitutes a cycle of chemotherapy;</i> <i>(g) streptozocin.</i> <i>No more than 1 pack containing 1 x 125 mg capsule and 2 x 80 mg capsules will be authorised per cycle of cytotoxic chemotherapy.</i> NOTE: <i>No applications for increased maximum quantities and/or repeats will be authorised.</i>							
8808N	Pack containing 1 capsule 125 mg and 2 capsules 80 mg	‡1	..	..	151.41	28.60	Emend MK
PROCHLORPERAZINE CAUTION: Prochlorperazine may be associated with parkinsonism and tardive dyskinesia and should be used for short-term treatment only.							
2893G	Tablet 5 mg	25	..	..	7.35	8.32	^a Stemizine HP
				B2.15	9.50	8.32	^a Stemetil AV
2369Q	Injection 12.5 mg in 1 mL	10	..	..	14.17	15.14	Stemetil AV
2894H	Suppositories 5 mg, 5	‡1	2	..	15.37	16.34	Stemetil AV
2895J	Suppositories 25 mg, 5	‡1	2	..	16.91	17.88	Stemetil AV
NOTE: As prochlorperazine may be associated with parkinsonism and tardive dyskinesia it should be used for short-term treatment only. However, authorities for increased maximum quantities and/or repeats of prochlorperazine tablets will be granted for the treatment of emesis associated with malignant disease.							
BILE AND LIVER THERAPY Bile therapy • Bile acid preparations URSODEOXYCHOLIC ACID Authority required <i>Primary biliary cirrhosis.</i> NOTE: <i>Not for use in the treatment of sclerosing cholangitis or cholelithiasis.</i>							
8448P	Capsule 250 mg	100	2	..	180.75	28.60	Ursofalk OA

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
LAXATIVES								
Laxatives								
• Contact laxatives								
BISACODYL								
Restricted benefit								
<i>Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;</i>								
<i>Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;</i>								
<i>For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;</i>								
<i>Patients receiving palliative care;</i>								
<i>Terminal malignant neoplasia;</i>								
<i>Anorectal congenital abnormalities;</i>								
<i>Megacolon.</i>								
1259G	Tablet 5 mg	200	2	..	13.00	13.97	Bisalax Lax-Tab	AS AE
1260H	Suppositories 10 mg, 10	3	5	..	*21.52	22.49	Durolax Petrus Bisacodyl Suppositories	BY PP
1258F	Suppositories 10 mg, 12	3	4	..	*17.50	18.47	Fleet Laxative Suppositories Petrus Bisacodyl Suppositories	FL PP
DOCUSATE SODIUM with BISACODYL								
Restricted benefit								
<i>Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;</i>								
<i>Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;</i>								
<i>For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;</i>								
<i>Patients receiving palliative care;</i>								
<i>Terminal malignant neoplasia;</i>								
<i>Anorectal congenital abnormalities;</i>								
<i>Megacolon.</i>								
1125F	Suppositories 100 mg-10 mg, 5	6	5	..	*21.49	22.46	Coloxyl	FM

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Bulk producers STERCULIA with FRANGULA BARK Restricted benefit <i>Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;</i> <i>Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;</i> <i>For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;</i> <i>Patients receiving palliative care;</i> <i>Terminal malignant neoplasia;</i> <i>Anorectal congenital abnormalities;</i> <i>Megacolon.</i>								
1104D	Granules 620 mg-80 mg per g (62%-8%), 500 g	‡1	1	..	23.05	24.02	Normacol Plus	NE
• Osmotically acting laxatives LACTULOSE Restricted benefit <i>Hepatic coma or precoma (chronic porto-systemic encephalopathy);</i> <i>Constipation in patients with malignant neoplasia.</i>								
3064G	Mixture 3.34 g per 5 mL, 500 mL	‡1	5	..	15.05	16.02	^a Actilax ^a Genlac ^a GenRx Lactulose ^a Lac-Dol ^a Lactocur ^a Duphalac	AF AW GX DP HX SM
				B2.20	17.25	16.02		
MACROGOL 3350 Restricted benefit <i>Constipation in patients with malignant neoplasia.</i>								
8612G	Sachets containing powder for solution 13.125 g with electrolytes, 30	‡1	5	..	22.90	23.87	Movicol	NE
• Enemas BISACODYL Restricted benefit <i>Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;</i> <i>Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;</i> <i>For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;</i> <i>Patients receiving palliative care;</i> <i>Terminal malignant neoplasia;</i> <i>Anorectal congenital abnormalities;</i> <i>Megacolon.</i>								
1263L	Enemas 10 mg in 5 mL, 25	‡1	2	..	35.77	28.60	Bisalax	AS

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
SORBITOL with SODIUM CITRATE and SODIUM LAURYL SULFOACETATE								
<u>Restricted benefit</u>								
<i>Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;</i>								
<i>Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;</i>								
<i>For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;</i>								
<i>Patients receiving palliative care;</i>								
<i>Terminal malignant neoplasia;</i>								
<i>Anorectal congenital abnormalities;</i>								
<i>Megacolon.</i>								
2091C	Enemas 3.125 g-450 mg-45 mg in 5 mL, 12	2	2	..	*34.53	28.60	Microlax	PH
• Other laxatives								
GLYCEROL								
<u>Restricted benefit</u>								
<i>Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;</i>								
<i>Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;</i>								
<i>For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;</i>								
<i>Patients receiving palliative care;</i>								
<i>Terminal malignant neoplasia;</i>								
<i>Anorectal congenital abnormalities;</i>								
<i>Megacolon.</i>								
2555L	Suppositories 700 mg (for infants), 12	3	5	..	*14.77	15.74	PP	
2556M	Suppositories 1.4 g (for children), 12	3	5	..	*15.16	16.13	PP	
2557N	Suppositories 2.8 g (for adults), 12	3	5	..	*15.52	16.49	PP	
ANTIDIARRHEALS, INTESTINAL ANTIINFLAMMATORY/ ANTIINFECTIVE AGENTS								
Intestinal antiinfectives								
• Antibiotics								
NEOMYCIN SULFATE								
2325J	Tablet 500 mg	25	1	..	13.29	14.26	Neosulf	AF
NYSTATIN								
1696G	Tablet 500,000 units	50	..	..	16.18	17.15	Nilstat	SI
1699K	Capsule 500,000 units	50	..	..	16.18	17.15	Nilstat	SI

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
VANCOMYCIN								
Authority required								
<i>Antibiotic associated pseudomembranous colitis due to Clostridium difficile which is unresponsive to metronidazole;</i>								
<i>Antibiotic associated pseudomembranous colitis due to Clostridium difficile where there is intolerance to metronidazole.</i>								
NOTE:								
<i>Metronidazole has similar efficacy to vancomycin but may have less selective pressure to vancomycin resistant enterococci and is therefore the preferred treatment.</i>								
3113W	Capsule 125 mg (125,000 i.u.) vancomycin activity	40	..	..	*255.59	28.60	Vancocin	LY
3114X	Capsule 250 mg (250,000 i.u.) vancomycin activity	40	..	..	*489.05	28.60	Vancocin	LY
Electrolytes with carbohydrates								
• Oral rehydration salt formulations								
ELECTROLYTE REPLACEMENT (ORAL)								
3196F	Sachets containing powder for oral solution 4.9 g, 10	‡1	..	..	11.71	12.68	Chem mart Oral Rehydration Salts O.R.S. Repalyte New Formulation Terry White Chemists Oral Rehydration Salts	CH AS AV TW
NOTE:								
Each sachet contains sodium chloride 470 mg, potassium chloride 300 mg, sodium acid citrate 530 mg and glucose 3.56 g.								
Antipropulsives								
• Antipropulsives								
DIPHENOXYLATE HYDROCHLORIDE with ATROPINE SULFATE								
2501P	Tablet 2.5 mg-25 micrograms	20	..	..	6.70	7.67	^a Lofenoxal	KR
				^B 1.83	8.53	7.67	^a Lomotil	PH
LOPERAMIDE HYDROCHLORIDE								
1571Q	Capsule 2 mg	12	..	..	6.70	7.67	^a Gastro-Stop Loperamide	AS
				^B 0.92	7.62	7.67	^a Imodium	JC
Intestinal antiinflammatory agents								
• Corticosteroids acting locally								
HYDROCORTISONE ACETATE								
Restricted benefit								
<i>Proctitis;</i>								
<i>Ulcerative colitis.</i>								
1502C	Rectal foam 90 mg per applicatorful, 14 applications, aerosol 21.1 g	2	3	..	*36.11	28.60	Colifoam	GC

[For other listings for this drug see Generic/Proprietary Index]

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
1920C	PREDNISOLONE SODIUM PHOSPHATE Retention enema equivalent to 20 mg prednisolone in 100 mL	28	3	..	*93.67	28.60	Predsol	SI
PREDNISOLONE SODIUM PHOSPHATE								
<u>Restricted benefit</u>								
<i>Proctitis;</i>								
<i>Ulcerative colitis.</i>								
2554K	Suppositories equivalent to 5 mg prednisolone, 10	3	3	..	*27.94	28.60	Predsol	SI
• Aminosalicylic acid and similar agents								
BALSALAZIDE SODIUM								
<u>Authority required</u>								
<i>Mild to moderate ulcerative colitis where hypersensitivity to sulfonamides exists;</i>								
<i>Mild to moderate ulcerative colitis where intolerance to sulfasalazine exists.</i>								
<u>NOTE:</u>								
<i>Not for the treatment of Crohn's disease.</i>								
8845M	Capsule 750 mg	180	5	..	127.13	28.60	Colazide	PK
MESALAZINE								
<u>Restricted benefit</u>								
<i>Acute episode of mild to moderate ulcerative proctitis.</i>								
<u>NOTE:</u>								
<i>Not for the treatment of Crohn's disease.</i>								
8752P	Suppositories 1 g, 28	‡1	..	..	130.11	28.60	Pentasa	FP
<u>NOTE:</u>								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
<u>Authority required</u>								
<i>Colitis (including ulcerative colitis and Crohn's disease) where hypersensitivity to sulfonamides exists;</i>								
<i>Colitis (including ulcerative colitis and Crohn's disease) where intolerance to sulfasalazine exists.</i>								
1611T	Tablet 250 mg	100	5	..	73.91	28.60	Mesasal	GK
<u>Authority required</u>								
<i>Colitis and Crohn's disease where hypersensitivity to sulfonamides exists;</i>								
<i>Colitis and Crohn's disease where intolerance to sulfasalazine exists.</i>								
8731M	Tablet 500 mg (enteric coated)	100	5	..	139.19	28.60	Salofalk	OA

continued ➡

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MESALAZINE—cont.								
Authority required								
<i>Mild to moderate ulcerative colitis where hypersensitivity to sulfonamides exists;</i>								
<i>Mild to moderate ulcerative colitis where intolerance to sulfasalazine exists.</i>								
NOTE:								
<i>Not for the treatment of Crohn's disease.</i>								
8598M	Sachet containing granules, 500 mg per sachet	100	5	..	143.06	28.60	Salofalk	OA
8599N	Sachet containing granules, 1 g per sachet	100	2	..	259.14	28.60	Salofalk	OA
Authority required								
<i>Acute episode of mild to moderate ulcerative colitis.</i>								
NOTE:								
<i>Not for the treatment of Crohn's disease.</i>								
8616L	Enemas 2 g in 60 mL, 7	4	..	..	*344.99	28.60	Salofalk	OA
8617M	Enemas 4 g in 60 mL, 7	4	..	..	*458.31	28.60	Salofalk	OA
8768L	Rectal foam 1 g per applicatorful, 14 applications, aerosol 80 g	4	..	..	*344.99	28.60	Salofalk	OA
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
Authority required								
<i>Acute episode of mild to moderate ulcerative proctosigmoiditis.</i>								
NOTE:								
<i>Not for the treatment of Crohn's disease.</i>								
8753Q	Enemas 1 g in 100 mL, 7	4	..	..	*344.99	28.60	Pentasa	FP
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
OLSALAZINE SODIUM								
Authority required								
<i>Ulcerative colitis where hypersensitivity to sulfonamides exists;</i>								
<i>Ulcerative colitis where intolerance to sulfasalazine exists.</i>								
NOTE:								
<i>Not for the treatment of Crohn's disease.</i>								
1728Y	Capsule 250 mg	100	5	..	61.58	28.60	Dipentum	UC
8086N	Tablet 500 mg	100	5	..	104.85	28.60	Dipentum	UC
SULFASALAZINE								
2093E	Tablet 500 mg	200	5	..	*53.29	28.60	Salazopyrin	PH
2096H	Tablet 500 mg (enteric coated)	200	5	..	*58.15	28.60	^a Pyralin EN	KR
				^B 1.64	*59.79	28.60	^a Salazopyrin-EN	PH

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DIGESTIVES, INCL. ENZYMES								
Digestives, incl. enzymes								
• Enzyme preparations								
PANCREATIC EXTRACT								
8556H	Capsule (containing enteric coated minimicrospheres) providing not less than 5,000 BP units of lipase activity	500	10	..	*121.35	28.60	Creon 5000	SM
8020D	Capsule (containing enteric coated minimicrospheres) providing not less than 10,000 BP units of lipase activity	500	10	..	*174.60	28.60	Creon	SM
8021E	Capsule (containing enteric coated minimicrospheres) providing not less than 25,000 BP units of lipase activity	200	10	..	*140.63	28.60	Creon Forte	SM
PANCRELIPASE								
2495H	Capsule (containing enteric coated microspheres) providing not less than 10,000 BP units of lipase activity	500	10	..	*177.49	28.60	Cotazym-S Forte	OR
8366H	Capsule (containing enteric coated microtablets) providing not less than 25,000 BP units of lipase activity	200	10	..	*140.63	28.60	Panzytrat 25000	TM
DRUGS USED IN DIABETES								
Insulins and analogues								
• Insulins and analogues, fast-acting								
INSULIN ASPART								
8571D	Injection (human analogue) 100 units per mL, 10 mL	5	2	..	*162.70	28.60	NovoRapid	NO
8435Y	Injections (human analogue) 100 units per mL, 3 mL, 5	5	1	..	*270.60	28.60	NovoRapid FlexPen NovoRapid Penfill 3 mL	NF NO
INSULIN LISPRO								
8084L	Injection (human analogue) 100 units per mL, 10 mL	5	2	..	*162.70	28.60	Humalog	LY
8212F	Injections (human analogue) 100 units per mL, 3 mL, 5	5	1	..	*270.60	28.60	Humalog	LY
INSULIN NEUTRAL								
1713E	Injection (bovine) 100 units per mL, 10 mL	5	2	..	*136.35	28.60	Hypurin Neutral	AS
1531N	Injection (human) 100 units per mL, 10 mL	5	2	..	*136.35	28.60	Actrapid Humulin R	NO LY

continued ➡

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
INSULIN NEUTRAL—cont.								
1762R	Injections (human) 100 units per mL, 3 mL, 5	5	1	..	*229.30	28.60	Actrapid Penfill 3 mL Humulin R	NO LY
• <i>Insulins and analogues, intermediate-acting</i>								
INSULIN ISOPHANE (N.P.H.)								
1711C	Injection (bovine) 100 units per mL, 10 mL	5	2	..	*136.35	28.60	Hypurin Isophane	AS
1533Q	Injection (human) 100 units per mL, 10 mL	5	2	..	*136.35	28.60	Humulin NPH Protaphane	LY NO
1761Q	Injections (human) 100 units per mL, 3 mL, 5	5	1	..	*229.30	28.60	Humulin NPH Protaphane InnoLet Protaphane NovoLet 3 mL Protaphane Penfill 3 mL	LY NI NL NO
• <i>Insulins and analogues, intermediate-acting combined with fast-acting</i>								
INSULIN ASPART—INSULIN ASPART PROTAMINE SUSPENSION								
8609D	Injections (human analogue) 100 units (30 units-70 units) per mL, 3 mL, 5	5	1	..	*270.60	28.60	NovoMix 30 FlexPen NovoMix 30 Penfill 3 mL	NF NO
INSULIN LISPRO—INSULIN LISPRO PROTAMINE SUSPENSION								
8390N	Injections (human analogue) 100 units (25 units-75 units) per mL, 3 mL, 5	5	1	..	*270.60	28.60	Humalog Mix25	LY
INSULIN NEUTRAL—INSULIN ISOPHANE (N.P.H.), (MIXED) (Biphasic Isophane)								
1426C	Injection (human) 100 units (30 units-70 units) per mL, 10 mL	5	2	..	*136.35	28.60	Humulin 30/70 Mixtard 30/70	LY NO
1425B	Injection (human) 100 units (50 units-50 units) per mL, 10 mL	5	2	..	*136.35	28.60	Mixtard 50/50	NO
8006J	Injections (human) 100 units (20 units-80 units) per mL, 3 mL, 5	5	1	..	*229.30	28.60	Mixtard 20/80 Penfill 3 mL	NO
1763T	Injections (human) 100 units (30 units-70 units) per mL, 3 mL, 5	5	1	..	*229.30	28.60	Humulin 30/70 Mixtard 30/70 InnoLet Mixtard 30/70 Penfill 3 mL	LY NI NO
2062M	Injections (human) 100 units (50 units-50 units) per mL, 3 mL, 5	5	1	..	*229.30	28.60	Mixtard 50/50 Penfill 3 mL	NO

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
Oral blood glucose lowering drugs							
• Biguanides							
METFORMIN HYDROCHLORIDE							
2430X	Tablet 500 mg	100	5	..	13.54	14.51	^a Chem mart Metformin CH ^a Diaformin AF ^a Formet 500 AW ^a GenRx GX Metformin ^a Glucohexal HX ^a Glucomet DP 500 mg ^a Terry White Chemists TW Metformin ^B 1.41 14.95 14.51 ^a Glucophage MQ ^B 1.80 15.34 14.51 ^a Diabex AL
1801T	Tablet 850 mg	60	5	..	13.54	14.51	^a Chem mart Metformin CH ^a Diaformin 850 AF ^a Formet 850 AW ^a GenRx GX Metformin ^a Glucohexal HX ^a Glucomet DP 850 mg ^a Terry White Chemists TW Metformin ^B 1.41 14.95 14.51 ^a Glucophage MQ ^B 1.80 15.34 14.51 ^a Diabex 850 AL
8607B	Tablet 1 g	90	5	..	20.05	21.02	^a Diaformin 1000 AF ^a Formet 1000 AW ^B 1.81 21.86 21.02 ^a Diabex 1000 AL
• Sulfonamides, urea derivatives							
GLIBENCLAMIDE							
CAUTION: Sulfonylureas may cause hypoglycaemia, particularly in the elderly.							
2939Q	Tablet 5 mg	100	5	..	10.06	11.03	^a Glimel AF ^B 1.25 11.31 11.03 ^a Daonil AV
GLICLAZIDE							
CAUTION: Sulfonylureas may cause hypoglycaemia, particularly in the elderly.							
8535F	Tablet 30 mg (modified release)	100	5	..	15.76	16.73	Diamicron MR SE

continued ➡

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
GLICLAZIDE—cont.								
2449X	Tablet 80 mg	100	5	..	15.43	16.40	^a Chem mart Gliclazide	CH
							^a GenRx Gliclazide	GX
							^a Glyade	AF
							^a Mellihexal	HX
							^a Nidem	AW
							^a Terry White Chemists Gliclazide	TW
				^B 1.35	16.78	16.40	^a Diamicon	SE
GLIMEPIRIDE								
CAUTION: Sulfonylureas may cause hypoglycaemia, particularly in the elderly.								
8450R	Tablet 1 mg	30	5	..	8.78	9.75	^a Dimirel	ML
				^B 2.50	11.28	9.75	^a Amaryl	AV
8451T	Tablet 2 mg	30	5	..	12.45	13.42	^a Dimirel	ML
				^B 2.50	14.95	13.42	^a Amaryl	AV
8533D	Tablet 3 mg	30	5	..	14.65	15.62	^a Dimirel	ML
				^B 2.50	17.15	15.62	^a Amaryl	AV
8452W	Tablet 4 mg	30	5	..	16.85	17.82	^a Dimirel	ML
				^B 2.50	19.35	17.82	^a Amaryl	AV
GLIPIZIDE								
CAUTION: Sulfonylureas may cause hypoglycaemia, particularly in the elderly.								
2440K	Tablet 5 mg	100	5	..	10.17	11.14	^a Melizide	AF
				^B 4.10	14.27	11.14	^a Minidiab	PH
• Combinations of oral blood glucose lowering drugs METFORMIN HYDROCHLORIDE with GLIBENCLAMIDE CAUTION: Sulfonylureas may cause hypoglycaemia, particularly in the elderly.								
8838E	Tablet 250 mg-1.25 mg	90	5	..	13.37	14.34	Glucovance 250mg/1.25mg	AL
8810Q	Tablet 500 mg-2.5 mg	90	5	..	16.25	17.22	Glucovance 500mg/2.5mg	AL
8811R	Tablet 500 mg-5 mg	90	5	..	17.44	18.41	Glucovance 500mg/5mg	AL
• Alpha glucosidase inhibitors ACARBOSE								
8188Y	Tablet 50 mg	90	5	..	28.95	28.60	Glucobay 50	BN
8189B	Tablet 100 mg	90	5	..	38.85	28.60	Glucobay 100	BN

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Thiazolidinediones PIOGLITAZONE HYDROCHLORIDE Authority required ➤ <i>Initiation of therapy, in combination with either metformin or a sulfonylurea, in type 2 diabetic patients whose blood glucose concentrations are inadequately controlled, and:</i> <i>(a) in whom combination therapy with metformin and a sulfonylurea is contraindicated or not tolerated; or</i> <i>(b) where combination therapy with metformin and a sulfonylurea is being continued despite the development of intolerance or a contraindication to either agent.</i> <i>Inadequate control is defined as Hb A1c greater than 7% despite diet, exercise and maximally tolerated doses of metformin or a sulfonylurea.</i> <i>The patient's Hb A1c level and the date of measurement, which must have occurred no earlier than 4 months prior to the date of application, must be provided at the time authority approval for initial treatment is sought.</i> <i>Pioglitazone hydrochloride is not PBS-subsidised as initial treatment when used in combination with both metformin and a sulfonylurea.</i> Authority required <i>Continuation of therapy, in combination with either metformin or a sulfonylurea, in type 2 diabetic patients where the patient has previously been issued with an authority prescription for pioglitazone hydrochloride or rosiglitazone maleate.</i> <i>Pioglitazone hydrochloride is not PBS-subsidised as continuing treatment when used in combination with both metformin and a sulfonylurea.</i> Authority required ➤ <i>Initiation of therapy, in combination with insulin, in insulin-treated type 2 diabetic patients whose blood glucose concentrations are inadequately controlled.</i> <i>Inadequate control is defined as Hb A1c greater than 7% despite concomitant use of insulin and 1 or more oral anti-diabetic agents or despite use of insulin alone, where metformin would have been added to the patient's treatment regimen but where metformin is contraindicated.</i> <i>The patient's Hb A1c level and the date of measurement, which must have occurred no earlier than 4 months prior to the date of application, must be provided at the time authority approval for initial treatment is sought.</i> Authority required <i>Continuation of therapy, in combination with insulin, in type 2 diabetic patients where the patient has previously been issued with an authority prescription for pioglitazone hydrochloride or rosiglitazone maleate.</i> <i>Patients who commenced pioglitazone hydrochloride therapy in combination with insulin prior to 1 November 2003 may continue to receive PBS-subsidised treatment on the same basis as originally approved.</i>								
8694N	Tablet 15 mg (base)	28	5	..	61.70	28.60	Actos	LY
8695P	Tablet 30 mg (base)	28	5	..	92.34	28.60	Actos	LY
8696Q	Tablet 45 mg (base)	28	5	..	118.64	28.60	Actos	LY

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ROSIGLITAZONE MALEATE							
<u>Authority required</u>							
➤ Initiation of therapy, in combination with 1 or more oral hypoglycaemic agents, in type 2 diabetic patients whose blood glucose concentrations are inadequately controlled, and: (a) in whom combination therapy with metformin and a sulfonylurea is contraindicated or not tolerated; or (b) where combination therapy with metformin and a sulfonylurea is being continued despite the development of intolerance or a contraindication to either agent; or (c) who are taking maximally tolerated doses of metformin and a sulfonylurea. Inadequate control is defined as Hb A1c greater than 7% despite diet, exercise and maximally tolerated doses of 1 or more oral hypoglycaemic agents. Applications for initial PBS-subsidised therapy for patients who were taking maximally tolerated doses of metformin and a sulfonylurea and for whom rosiglitazone maleate was added to therapy prior to 1 April 2005 will be approved providing the patient's Hb A1c level at the time of commencing rosiglitazone maleate therapy was above 7%. The patient's Hb A1c level and the date of measurement, which must have occurred no earlier than 4 months prior to the date of application or prior to commencing rosiglitazone maleate therapy, must be provided at the time authority approval for initial treatment is sought.							
<u>Authority required</u>							
Continuing PBS-subsidised therapy, in combination with 1 or more oral hypoglycaemic agents, in type 2 diabetic patients where the patient has previously been issued with an authority prescription for rosiglitazone maleate or pioglitazone hydrochloride.							
<u>Authority required</u>							
➤ Initiation of therapy, in combination with insulin, in insulin-treated type 2 diabetic patients whose blood glucose concentrations are inadequately controlled. Inadequate control is defined as Hb A1c greater than 7% despite concomitant use of insulin and 1 or more oral anti-diabetic agents or despite use of insulin alone, where metformin would have been added to the patient's treatment regimen but where metformin is contraindicated. The patient's Hb A1c level and the date of measurement, which must have occurred no earlier than 4 months prior to the date of application, must be provided at the time authority approval for initial treatment is sought.							
<u>Authority required</u>							
Continuation of therapy, in combination with insulin, in type 2 diabetic patients where the patient has previously been issued with an authority prescription for rosiglitazone maleate or pioglitazone hydrochloride.							
8689H	Tablet 4 mg (base)	28	5	..	61.70	28.60	Avandia GK
8690J	Tablet 8 mg (base)	28	5	..	92.34	28.60	Avandia GK

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
VITAMINS								
Vitamin A and D, incl. combinations of the two								
• Vitamin D and analogues								
CALCITRIOL								
Authority required								
<i>Hypocalcaemia due to renal disease;</i>								
<i>Hypoparathyroidism;</i>								
<i>Hypophosphataemic rickets;</i>								
<i>Vitamin D-resistant rickets;</i>								
<i>Initial treatment for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be included in the authority application.</i>								
<i>A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body;</i>								
<i>Continuing treatment for established osteoporosis in patients with fracture due to minimal trauma, where the patient has previously been issued with an authority prescription for this drug.</i>								
2502Q	Capsule 0.25 microgram	100	3	..	53.74	28.60	^a Calcitriol-DP	DP
							^a Citrihexal	HX
							^a GenRx	GX
							Calcitriol	
							^a Kosteol	AW
							^a Rocaltrol	RO
							^a Sitriol	AF
MINERAL SUPPLEMENTS								
Calcium								
• Calcium								
CALCIUM								
Authority required								
➤ <i>Hyperphosphataemia associated with chronic renal failure.</i>								
8560M	Tablet 250 mg (as citrate)	120	1	..	12.55	13.52	Citracal	KY
3116B	Tablet (chewable) 500 mg (as carbonate)	120	1	..	*13.49	14.46	Cal-Sup	MM
3117C	Tablet 600 mg (as carbonate)	120	1	..	12.55	13.52	Caltrate	WT
Potassium								
• Potassium								
POTASSIUM CHLORIDE								
2642C	Tablet 600 mg (sustained release)	200	1	..	*10.71	11.68	^a Duro-K	SZ
				..	10.71	11.68	Span-K	AS
				^B 2.66	*13.37	11.68	^a Slow-K	NV
3012M	Effervescent tablet 14 mmol K ⁺ and 8 mmol Cl ⁻	60	1	..	10.84	11.81	^a K-Sol	LN
				^B 2.63	13.47	11.81	^a Chlorvescent	AS

ALIMENTARY TRACT AND METABOLISM —cont.

Code	Name, Restriction, Manner of administration and form	Max. No.of			Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
		Qty	Rpts	Premium				
ANABOLIC AGENTS FOR SYSTEMIC USE								
Anabolic steroids								
• Androstan derivatives								
OXANDROLONE								
<u>Authority required</u>								
<i>Promotion of growth in girls with Turner syndrome (date of confirmation of diagnosis to be stated);</i>								
<i>Promotion of growth in boys of short stature with delayed bone maturation (date of confirmation of diagnosis to be stated).</i>								
<u>NOTE:</u>								
<i>Oxandrolone should not be prescribed for promotion of growth in boys aged less than 11 years or boys who do not have a bone age of at least 9 years.</i>								
2545Y	Tablet 2.5 mg	100	1	..	1011.37	28.60	Oxandrin	CS
• Estren derivatives								
NANDROLONE DECANOATE								
<u>Authority required</u>								
<i>Monotherapy for osteoporosis, where other treatment has failed and where specialist advice confirms that this is the only suitable treatment option for the patient. Specialist advice need only be obtained for the first authority approval;</i>								
<i>Monotherapy for osteoporosis, where other treatment is not tolerated and where specialist advice confirms that this is the only suitable treatment option for the patient. Specialist advice need only be obtained for the first authority approval;</i>								
<i>Monotherapy for osteoporosis, where other treatment is contraindicated and where specialist advice confirms that this is the only suitable treatment option for the patient. Specialist advice need only be obtained for the first authority approval;</i>								
<i>Patients receiving PBS-subsidised therapy with this drug for osteoporosis prior to 1 February 2004;</i>								
<i>Patients on long-term treatment with corticosteroids.</i>								
<u>NOTE:</u>								
<i>Monotherapy for the treatment of osteoporosis does not exclude calcium supplementation.</i>								
1671Y	Injection 50 mg in 1 mL, disposable syringe	1	7	..	19.36	20.33	Deca-Durabolin	OR

BLOOD AND BLOOD FORMING ORGANS

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTITHROMBOTIC AGENTS								
Antithrombotic agents								
• Vitamin K antagonists								
WARFARIN SODIUM								
CAUTION:								
The listed brands have NOT been shown to be bioequivalent and should not be interchanged.								
2843P	Tablet 1 mg	50	2	..	7.59	8.56	Coumadin Marevan	SI FM
2209G	Tablet 2 mg	50	2	..	7.76	8.73	Coumadin	SI
2844Q	Tablet 3 mg	50	2	..	8.00	8.97	Marevan	FM
2211J	Tablet 5 mg	50	2	..	8.49	9.46	Coumadin Marevan	SI FM
• Heparin group								
DALTEPARIN SODIUM								
(Low Molecular Weight Heparin Sodium—porcine mucous)								
8603T	Injection 2,500 units (anti-Xa) in 0.2 mL single dose pre-filled syringe	10	1	..	55.55	28.60	Fragmin	PH
2816F	Injection 5,000 units (anti-Xa) in 0.2 mL single dose pre-filled syringe	10	1	..	57.68	28.60	Fragmin	PH
8271H	Injection 7,500 units (anti-Xa) in 0.75 mL single dose pre-filled syringe	10	1	..	84.56	28.60	Fragmin	PH
8269F	Injection 10,000 units (anti-Xa) in 1 mL single dose pre-filled syringe	10	1	..	111.14	28.60	Fragmin	PH
DALTEPARIN SODIUM								
(Low Molecular Weight Heparin Sodium—porcine mucous)								
Restricted benefit								
Haemodialysis.								
8641T	Injection 2,500 units (anti-Xa) in 0.2 mL single dose pre- filled syringe	20	3	..	*106.35	28.60	Fragmin	PH
8642W	Injection 5,000 units (anti-Xa) in 0.2 mL single dose pre- filled syringe	20	3	..	*110.61	28.60	Fragmin	PH
8643X	Injection 7,500 units (anti-Xa) in 0.75 mL single dose pre- filled syringe	20	3	..	*164.37	28.60	Fragmin	PH

BLOOD AND BLOOD FORMING ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ENOXAPARIN SODIUM								
8558K	Injection 20 mg (2,000 i.u. anti-Xa) in 0.2 mL pre-filled syringe	10	1	..	55.55	28.60	Clexane	AV
8510X	Injection 40 mg (4,000 i.u. anti-Xa) in 0.4 mL pre-filled syringe	10	1	..	57.68	28.60	Clexane	AV
8262W	Injection 60 mg (6,000 i.u. anti-Xa) in 0.6 mL pre-filled syringe	10	1	..	80.45	28.60	Clexane	AV
8263X	Injection 80 mg (8,000 i.u. anti-Xa) in 0.8 mL pre-filled syringe	10	1	..	91.85	28.60	Clexane	AV
8264Y	Injection 100 mg (10,000 i.u. anti-Xa) in 1 mL pre-filled syringe	10	1	..	110.83	28.60	Clexane	AV
<u>ENOXAPARIN SODIUM</u>								
<u>Restricted benefit</u>								
<u>Haemodialysis.</u>								
8716R	Injection 20 mg (2,000 i.u. anti-Xa) in 0.2 mL pre-filled syringe	20	3	..	*106.35	28.60	Clexane	AV
8639Q	Injection 40 mg (4,000 i.u. anti-Xa) in 0.4 mL pre-filled syringe	20	3	..	*110.61	28.60	Clexane	AV
8640R	Injection 60 mg (6,000 i.u. anti-Xa) in 0.6 mL pre-filled syringe	20	3	..	*156.15	28.60	Clexane	AV
HEPARIN SODIUM								
1466E	Injection 5,000 units in 0.2 mL	5	5	..	11.28	12.25	MX	
1467F	Injection 5,000 units in 1 mL	5	5	..	11.28	12.25	MX	
1463B	Injection (preservative-free) 5,000 units in 5 mL	50	5	..	52.50	28.60	PU	
1076P	Injection 35,000 units in 35 mL	12	5	..	*148.51	28.60	MX	
• Platelet aggregation inhibitors excl. heparin								
<u>ABCIXIMAB</u>								
<u>Authority required</u>								
<i>Patients undergoing percutaneous coronary balloon angioplasty;</i>								
<i>Patients undergoing percutaneous coronary atherectomy;</i>								
<i>Patients undergoing percutaneous coronary stent placement.</i>								
8048N	I.V. injection 10 mg in 5 mL	3	..	..	*1531.36	28.60	ReoPro	LY
ASPIRIN								
8202Q	Tablet 100 mg	112	1	..	6.36	7.33	^a DBL Aspirin 100 mg	FA
				^B 1.36	7.72	7.33	^a Astrix	MX
1010E	Tablet 300 mg (dispersible)	96	1	..	6.98	7.95	Solprin	RC

BLOOD AND BLOOD FORMING ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CLOPIDOGREL HYDROGEN SULFATE								
Authority required								
<i>Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events in patients:</i>								
<i>(1) with a history of symptomatic cerebrovascular ischaemic episodes while on therapy with low-dose aspirin; or</i>								
<i>(2) where low-dose aspirin poses an unacceptable risk of gastrointestinal bleeding; or</i>								
<i>(3) where there is a history of anaphylaxis, urticaria or asthma within 4 hours of ingestion of aspirin, other salicylates, or NSAIDs;</i>								
<i>Prevention of recurrence of myocardial infarction or unstable angina in patients:</i>								
<i>(1) with a history of symptomatic cardiac ischaemic events while on therapy with low-dose aspirin; or</i>								
<i>(2) where low-dose aspirin poses an unacceptable risk of gastrointestinal bleeding; or</i>								
<i>(3) where there is a history of anaphylaxis, urticaria or asthma within 4 hours of ingestion of aspirin, other salicylates, or NSAIDs.</i>								
NOTE:								
<i>Not for prophylaxis of DVT or peripheral arterial disease.</i>								
8358X	Tablet 75 mg (base)	28	5	..	84.13	28.60	Iscover Plavix	BQ SW
DIPYRIDAMOLE								
Restricted benefit								
<i>Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events:</i>								
<i>(1) as adjunctive therapy with low-dose aspirin; or</i>								
<i>(2) where low-dose aspirin poses an unacceptable risk of gastrointestinal bleeding; or</i>								
<i>(3) where there is a history of anaphylaxis, urticaria or asthma within 4 hours of ingestion of aspirin, other salicylates, or NSAIDs.</i>								
8335Q	Capsule 200 mg (sustained release)	60	5	..	33.78	28.60	Persantin SR	BY
DIPYRIDAMOLE with ASPIRIN								
Restricted benefit								
<i>Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events.</i>								
8382E	Capsule 200 mg (sustained release)-25 mg	60	5	..	32.51	28.60	Asasantin SR	BY
EPTIFIBATIDE ACETATE								
Authority required								
<i>Patients undergoing non-urgent percutaneous intervention with intracoronary stenting.</i>								
8683B	Solution for I.V. injection 20 mg (base) in 10 mL	2	..	..	*268.81	28.60	Integrilin	SH
8684C	Solution for I.V. infusion 75 mg (base) in 100 mL	3	..	..	*1052.53	28.60	Integrilin	SH
TICLOPIDINE HYDROCHLORIDE								
CAUTION:								
<i>Severe neutropenia is common in the early months of therapy. Haematological monitoring should be undertaken at commencement and every two weeks in the first four months of therapy.</i>								

continued ➡

BLOOD AND BLOOD FORMING ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
TICLOPIDINE HYDROCHLORIDE—cont.								
Authority required								
<i>Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events in patients:</i>								
<i>(1) with a history of symptomatic cerebrovascular ischaemic episodes while on therapy with low-dose aspirin; or</i>								
<i>(2) where low-dose aspirin poses an unacceptable risk of gastrointestinal bleeding; or</i>								
<i>(3) where there is a history of anaphylaxis, urticaria or asthma within 4 hours of ingestion of aspirin, other salicylates, or NSAIDs;</i>								
<i>Patients established on this drug as a pharmaceutical benefit prior to 1 November 1999.</i>								
2095G	Tablet 250 mg	60	5	..	152.45	28.60	^a Ticlopidine Hexal	HX
							^a Tilodene	AF
				^B 2.26	154.71	28.60	^a Ticlid	RO
TIROFIBAN HYDROCHLORIDE								
Authority required								
<i>Patients with high risk unstable angina who have new transient or persistent ST-T ischaemic changes and anginal pain lasting longer than 20 minutes;</i>								
<i>Patients with high risk unstable angina who have new transient or persistent ST-T ischaemic changes and repetitive episodes of angina at rest or during minimal exercise in the previous 12 hours;</i>								
<i>Patients with non-Q-wave myocardial infarction.</i>								
8350L	Solution concentrate for I.V. infusion 12.5 mg (base) in 50 mL	1	2	..	372.75	28.60	Aggrastat	MK
• Enzymes								
ALTEPLASE								
<i>(Recombinant tissue-type plasminogen activator)</i>								
Restricted benefit								
<i>Treatment of acute myocardial infarction within 12 hours of onset of attack.</i>								
1029E	Injection set containing 1 vial 50 mg in 2333 mg dry powder, 1 vial sterile water for injection 50 mL and 1 transfer cannula	2	..	..	*2157.53	28.60	Actilyse	BY
DROTRECIGIN ALFA (ACTIVATED)								
Authority required								
<i>Adult patients with severe sepsis who have a high risk of death as determined by acute dysfunction in at least 2 organs or modified Apache II score of at least 25.</i>								
<i>Acute organ dysfunction is defined as follows:</i>								
<i>(1) For cardiovascular-system dysfunction, an arterial systolic blood pressure of less than or equal to 90 mmHg or mean arterial pressure of less than or equal to 70 mmHg for at least 1 hour despite adequate fluid resuscitation, adequate intravascular volume status or the use of vasopressors in an attempt to maintain a systolic blood pressure of greater than or equal to 90 mmHg or a mean arterial pressure of greater than or equal to 70 mmHg;</i>								
<i>(2) For kidney dysfunction, urine output of less than 0.5 mL per kg of body weight per hour for 1 hour despite adequate fluid resuscitation;</i>								
<i>(3) For respiratory-system dysfunction, a ratio of PaO₂ to FiO₂ of less than or equal to 250;</i>								

continued ➤

BLOOD AND BLOOD FORMING ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DROTRECOGIN ALFA (ACTIVATED)—cont.								
(4) For haematologic dysfunction, a platelet count of less than 80,000 per cubic millimetre or which has decreased by 50 percent in the previous 3 days;								
(5) In the case of unexplained metabolic acidosis, a pH of less than or equal to 7.30 or a base deficit of greater than or equal to 5.0 mmol per L in association with a plasma lactate level of greater than 1.5 times the upper limit of the normal value for the reporting laboratory.								
NOTE: Medical practitioners should request the appropriate quantity of vials at the time of the authority application, according to the weight of the patient, to achieve a dose of 24 micrograms per kg per hour over a maximum of 96 hours.								
8614J	Powder for I.V. infusion 5 mg	1	..	..	480.56	28.60	Xigris	LY
8615K	Powder for I.V. infusion 20 mg	1	..	..	1908.02	28.60	Xigris	LY
RETEPLASE (Recombinant plasminogen activator)								
Restricted benefit Treatment of acute myocardial infarction within 6 hours of onset of attack.								
8253J	Pack containing 2 vials powder for injection 10 units, 2 single use pre-filled syringes with solvent, 2 reconstitution spikes and 2 needles	1	..	..	2224.11	28.60	Rapilysin 10 U	RO
TENECTEPLASE Restricted benefit Treatment of acute myocardial infarction within 12 hours of onset of attack.								
8526R	Powder for injection 40 mg with solvent	1	..	..	2101.39	28.60	Metalyse	BY
8527T	Powder for injection 50 mg with solvent	1	..	..	2212.67	28.60	Metalyse	BY
• Direct thrombin inhibitors BIVALIRUDIN TRIFLUOROACETATE Authority required Patients undergoing non-emergency percutaneous coronary intervention.								
NOTE: Bivalirudin trifluoroacetate has not been evaluated in patients undergoing emergency percutaneous coronary intervention, i.e. as primary or salvage therapy for myocardial infarction.								
8844L	Powder for I.V. injection 250 mg (base)	1	..	..	692.30	28.60	Angiomax	CS
• Other antithrombotic agents FONDAPARINUX SODIUM Authority required Prevention of venous thromboembolic events in patients undergoing major hip surgery; Prevention of venous thromboembolic events in patients undergoing total knee replacement.								
NOTE: No applications for increased maximum quantities and/or repeats will be authorised.								
8775W	Injection 2.5 mg in 0.5 mL single dose pre-filled syringe	7	..	..	120.25	28.60	Arixtra	GK

BLOOD AND BLOOD FORMING ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
BLOOD SUBSTITUTES AND PERFUSION SOLUTIONS								
Blood and related products								
• Blood substitutes and plasma protein fractions								
GELATIN - SUCCINYLATED								
8444K	I.V. infusion 20 g per 500 mL, 500 mL	3	..	..	*44.35	28.60	Gelofusine	BR
POLYGELINE								
2334W	I.V. infusion 17.5 g per 500 mL (3.5%) with electrolytes, 500 mL	3	..	..	*44.35	28.60	Haemaccel	AE
I.V. solutions								
• Solutions for parenteral nutrition								
GLUCOSE								
2245E	I.V. infusion 278 mmol (anhydrous) per L (5%), 1 L	5	1	..	*24.00	24.97	BX	
• Solutions affecting the electrolyte balance								
ELECTROLYTE REPLACEMENT SOLUTION								
3199J	I.V. infusion 1 L	2	1	..	*20.13	21.10	Plasma-Lyte 148	BX
SODIUM CHLORIDE								
2264E	I.V. infusion 154 mmol per L (0.9%), 1 L	5	1	..	*24.00	24.97	BX	
2260Y	I.V. infusion 513 mmol per L (3%), 1 L	2	1	..	*16.41	17.38	BX	
SODIUM CHLORIDE COMPOUND								
2266G	I.V. infusion 1 L	4	1	..	*28.07	28.60	BX	
SODIUM CHLORIDE with GLUCOSE								
2281C	I.V. infusion 31 mmol-222 mmol (anhydrous) per L (0.18%-4%), 1 L	5	1	..	*24.00	24.97	BX	
2279Y	I.V. infusion 19 mmol-104 mmol (anhydrous) per 500 mL (0.225%-3.75%), 500 mL	5	1	..	*30.05	28.60	BX	
2278X	I.V. infusion 39 mmol-69 mmol (anhydrous) per 500 mL (0.45%- 2.5%), 500 mL	5	1	..	*30.05	28.60	BX	
SODIUM LACTATE COMPOUND								
2286H	I.V. infusion 1 L	5	1	..	*24.00	24.97	BX	

CARDIOVASCULAR SYSTEM

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CARDIAC THERAPY								
Cardiac glycosides								
• Digitalis glycosides								
DIGOXIN								
2605D	Tablet 62.5 micrograms	200	1	..	8.51	9.48	^a Sigmaxin-PG	FM
				^B 1.30	9.81	9.48	^a Lanoxin-PG	SI
1322N	Tablet 250 micrograms	100	1	..	8.80	9.77	^a Sigmaxin	FM
				^B 1.30	10.10	9.77	^a Lanoxin	SI
3164M	Oral solution for children 50 micrograms per mL, 60 mL	2	3	..	*26.75	27.72	Lanoxin	SI
Antiarrhythmics, class I and III								
• Antiarrhythmics, class IA								
DISOPYRAMIDE								
2923W	Capsule 100 mg	100	5	..	25.24	26.21	Rythmodan	AV
2924X	Capsule 150 mg	100	5	..	34.16	28.60	Rythmodan	AV
QUINIDINE BISULFATE								
CAUTION: Severe thrombocytopenia has been reported with this drug.								
2623C	Tablet 250 mg (sustained release)	100	5	..	31.08	28.60	Kinidin Durule	AP
• Antiarrhythmics, class IB								
LIGNOCAINE HYDROCHLORIDE								
2875H	Injection 100 mg in 5 mL	5	..	..	35.31	28.60	PF	
2876J	Infusion 500 mg in 5 mL	10	..	..	27.65	28.60	Xylocard 500	AP
MEXILETINE HYDROCHLORIDE								
1682M	Capsule 50 mg	100	5	..	27.85	28.60	Mexitil	BY
1683N	Capsule 200 mg	100	5	..	56.45	28.60	Mexitil	BY
• Antiarrhythmics, class IC								
FLECAINIDE ACETATE								
CAUTION: <i>Flecainide acetate should be avoided in patients with poor cardiac function.</i>								
Restricted benefit <i>Serious supra-ventricular cardiac arrhythmias; Serious ventricular cardiac arrhythmias where treatment is initiated in a hospital (in-patient or out-patient).</i>								
1088G	Tablet 50 mg	60	5	..	38.32	28.60	Tambocor	MM
1090J	Tablet 100 mg	60	5	..	46.37	28.60	^a Flecatag	AF
				^B 2.86	49.23	28.60	^a Tambocor	MM

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
• Antiarrhythmics, class III AMIODARONE HYDROCHLORIDE CAUTION: <i>Amiodarone hydrochloride has been reported to cause frequent and potentially serious toxicity.</i> <i>Regular monitoring of hepatic and thyroid function is recommended.</i> Restricted benefit <i>Severe cardiac arrhythmias.</i>							
2344J	Tablet 100 mg	30	5	..	16.11	17.08	^a Aratac 100 AF ^a Cardinorm HX ^a GenRx GX <i>Amiodarone</i> ^a Rithmik 100 AW ^B 1.00 17.11 17.08 ^a Cordarone X SW 100
2343H	Tablet 200 mg	30	5	..	24.95	25.92	^a Aratac 200 AF ^a Cardinorm HX ^a Chem mart CH <i>Amiodarone</i> ^a GenRx GX <i>Amiodarone</i> ^a Rithmik 200 AW ^a Terry White TW Chemists <i>Amiodarone</i> ^B 1.00 25.95 25.92 ^a Cordarone X SW 200
SOTALOL HYDROCHLORIDE Restricted benefit <i>Severe cardiac arrhythmias.</i>							
8398B	Tablet 80 mg	60	5	..	17.08	18.05	^a GenRx Sotalol GX ^a Solavert AW ^a Sotahexal HX ^B 1.66 18.74 18.05 ^a Sotacor BQ
2043M	Tablet 160 mg	60	5	..	31.21	28.60	^a Cardol AF ^a Chem mart CH <i>Sotalol</i> ^a GenRx Sotalol GX ^a Solavert AW ^a Sotab DP ^a Sotahexal HX ^a Terry White TW Chemists <i>Sotalol</i> ^B 1.80 33.01 28.60 ^a Sotacor BQ

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
Cardiac stimulants excl. cardiac glycosides							
• Adrenergic and dopaminergic agents							
ADRENALINE							
1016L	Injection 1 mg in 1 mL (1 in 1,000)	5	1	..	14.76	15.73	AP
ADRENALINE							
Authority required							
<i>Initial supply for anticipated emergency treatment of acute allergic reactions with anaphylaxis in a patient who has been assessed to be at significant risk of anaphylaxis by, or in consultation with, a clinical immunologist, allergist, paediatrician or respiratory physician. The name of the specialist consulted must be provided at the time of application for initial supply;</i>							
<i>Continuing supply for anticipated emergency treatment of acute allergic reactions with anaphylaxis, where the patient has previously been issued with an authority prescription for this drug.</i>							
NOTE:							
<i>The auto-injector should be provided in the framework of a comprehensive anaphylaxis prevention program and an emergency action plan including training in recognition of the symptoms of anaphylaxis and the use of the auto-injector device. (For further information see the Australasian Society of Clinical Immunology and Allergy website at www.allergy.org.au.)</i>							
8697R	I.M. injection 150 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	96.41	28.60	EpiPen Jr. CS
8698T	I.M. injection 300 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	96.41	28.60	EpiPen CS
NOTE:							
<i>Authorities for increased maximum quantities, up to a maximum of 2, may be authorised for children aged less than 17 years where 2 auto-injectors are necessary to ensure 1 is on hand at all times. No increased maximum quantities will be authorised for patients aged 17 years or older.</i>							
No repeats will be issued.							
Vasodilators used in cardiac diseases							
• Organic nitrates							
GLYCERYL TRINITRATE							
1459T	Tablets 600 micrograms, 100	‡1	5	..	8.40	9.37	^a Lycinate FM
				^B 1.30	9.70	9.37	^a Anginine SI
				..	30.67	28.60	Stabilised LM
8171C	Buccal/sublingual spray (pump pack) 400 micrograms per dose (200 doses)	‡1	5	..	18.30	19.27	Nitrolingual AV
							Pumpspray
NOTE:							
The spray should not be inhaled.							

continued ➡

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
GLYCERYL TRINITRATE—cont.								
8027L	Transdermal patch releasing approximately 5 mg per 24 hours	30	5	..	27.14	28.11	Minitran 5	MM
1515R	Transdermal patch releasing approximately 5 mg per 24 hours	30	5	..	27.14	28.11	Transiderm- Nitro 25	NV
8010N	Transdermal patch releasing approximately 5 mg per 24 hours	30	5	..	27.14	28.11	Nitro-Dur 5	SH
8028M	Transdermal patch releasing approximately 10 mg per 24 hours	30	5	..	34.12	28.60	Minitran 10	MM
1516T	Transdermal patch releasing approximately 10 mg per 24 hours	30	5	..	34.12	28.60	Transiderm- Nitro 50	NV
8011P	Transdermal patch releasing approximately 10 mg per 24 hours	30	5	..	34.12	28.60	Nitro-Dur 10	SH
8119H	Transdermal patch releasing approximately 15 mg per 24 hours	30	5	..	34.12	28.60	Minitran 15	MM
8026K	Transdermal patch releasing approximately 15 mg per 24 hours	30	5	..	34.12	28.60	Nitro-Dur 15	SH
ISOSORBIDE DINITRATE								
2587E	Tablet 10 mg	200	2	..	*12.23	13.20	^a Sorbidin	AF
				^B 3.70	*15.93	13.20	^a Isordil	SI
2588F	Sublingual tablet 5 mg	200	2	..	*13.45	14.42	Isordil Sublingual	SI
ISOSORBIDE MONONITRATE								
1558B	Tablet 60 mg (sustained release)	30	5	..	13.26	14.23	^a Arso r b 60	AW
							^a Chem mart	CH
							Isosorbide	
							Mononitrate	
							^a Duride	AF
							^a GenRx	GX
							Isosorbide	
							Mononitrate	
							^a Imtrate 60 mg	DP
							^a Isomonit	HX
							^a Monodur 60 mg	PM
							^a Terry White	TW
							Chemists	
							Isosorbide	
							Mononitrate	
				^B 2.85	16.11	14.23	^a Imdur Durule	AP
8273K	Tablet 120 mg (sustained release)	30	5	..	24.94	25.91	^a Monodur	PM
							120 mg	
				^B 3.00	27.94	25.91	^a Imdur 120 mg	AP

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Other vasodilators used in cardiac diseases								
NICORANDIL								
8228C	Tablets 10 mg, 60	‡1	5	..	20.87	21.84	Ikorel	AV
8229D	Tablets 20 mg, 60	‡1	5	..	27.58	28.55	Ikorel	AV
PERHEXILINE MALEATE								
CAUTION:								
Regular monitoring of drug serum levels is recommended.								
Authority required								
Angina not responding to other therapy.								
1822X	Tablet 100 mg	100	5	..	53.15	28.60	Pexsig	SI
ANTIHYPERTENSIVES								
Antiadrenergic agents, centrally acting								
• Methyldopa								
METHYLDOPA								
1629R	Tablet 250 mg	100	5	..	12.36	13.33	^a Hydopa	AF
				^B 2.95	15.31	13.33	^a Aldomet	MK
• Imidazoline receptor agonists								
CLONIDINE								
3145M	Tablet 100 micrograms	100	5	..	26.95	27.92	Catapres 100	BY
3141H	Tablet 150 micrograms	100	5	..	35.41	28.60	Catapres	BY
Antiadrenergic agents, peripherally acting								
• Alpha-adrenoceptor antagonists								
PRAZOSIN HYDROCHLORIDE								
1479W	Tablet 1 mg (base)	100	5	..	12.92	13.89	^a Chem mart Prazosin	CH
							^a GenRx Prazosin	GX
							^a Prazohexal	HX
							^a Pressin 1	AF
							^a Terry White Chemists Prazosin	TW
				^B 3.05	15.97	13.89	^a Minipress	PF
1480X	Tablet 2 mg (base)	100	5	..	15.96	16.93	^a Chem mart Prazosin	CH
							^a GenRx Prazosin	GX
							^a Prazohexal	HX
							^a Pressin 2	AF
							^a Terry White Chemists Prazosin	TW
				^B 3.15	19.11	16.93	^a Minipress	PF

continued ➡

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PRAZOSIN HYDROCHLORIDE—cont.								
1478T	Tablet 5 mg (base)	100	5	..	23.77	24.74	^a Chem mart Prazosin	CH
							^a GenRx Prazosin	GX
							^a Prazohexal	HX
							^a Pressin 5	AF
							^a Terry White Chemists Prazosin	TW
				^B 3.40	27.17	24.74	^a Minipress	PF
Arteriolar smooth muscle, agents acting on								
• Hydrazinophthalazine derivatives								
HYDRALAZINE HYDROCHLORIDE								
1640H	Tablet 25 mg	200	2	..	*13.73	14.70	Alphapress 25	AF
1639G	Tablet 50 mg	200	2	..	*15.61	16.58	Alphapress 50	AF
• Pyrimidine derivatives								
MINOXIDIL								
Authority required								
<i>Severe refractory hypertension where treatment with minoxidil was initiated in a hospital (in-patient or out-patient).</i>								
2313R	Tablet 10 mg	100	5	..	52.39	28.60	Loniten	PH
DIURETICS								
Low-ceiling diuretics, thiazides								
• Thiazides, plain								
HYDROCHLOROTHIAZIDE								
1484D	Tablet 25 mg	100	1	..	19.40	20.37	Dithiazide	PL
Low-ceiling diuretics, excl. thiazides								
• Sulfonamides, plain								
CHLORTHALIDONE								
1585K	Tablet 25 mg	100	1	..	*11.87	12.84	Hygroton 25	NV
INDAPAMIDE HEMIHYDRATE								
8532C	Tablet 1.5 mg (sustained release)	90	1	..	19.78	20.75	Natrilix SR	SE
2436F	Tablet 2.5 mg	90	1	..	19.33	20.30	^a Chem mart Indapamide	CH
							^a Dapa-Tabs	AF
							^a GenRx Indapamide	GX
							^a Indahexal	HX
							^a Insig	SI
							^a Napamide 2.5 mg	DP
							^a Terry White Chemists Indapamide	TW
				^B 3.37	22.70	20.30	^a Natrilix	SE

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
High-ceiling diuretics								
• Sulfonamides, plain								
FRUSEMIDE								
2414C	Tablet 20 mg	100	1	..	8.18	9.15	^a Chem mart Frusemide	CH
							^a GenRx Frusemide	GX
							^a Terry White Chemists Frusemide	TW
				..	*8.19	9.16	^a Frusid Urex-M	DP FM
				^B 1.12	*9.31	9.16	^a Lasix-M	AV
2412Y	Tablet 40 mg	100	1	..	7.86	8.83	^a Chem mart Frusemide	CH
							^a Frusehexal 40 mg	HX
							^a Frusid	DP
							^a GenRx Frusemide	GX
							^a Terry White Chemists Frusemide	TW
							^a Uremide	AF
							^a Urex	FM
				^B 1.11	8.97	8.83	^a Lasix	AV
2415D	Tablet 500 mg	50	3	..	20.99	21.96	Urex-Forte	FM
2411X	Oral solution 10 mg per mL, 30 mL	‡1	3	..	13.95	14.92	Lasix	AV
2413B	Injection 20 mg in 2 mL	5	..	..	10.11	11.08	^a Frusehexal	HX
							^a Lasix	AV
• Aryloxyacetic acid derivatives								
ETHACRYNIC ACID								
<u>Restricted benefit</u>								
<i>Patients hypersensitive to other oral diuretics.</i>								
8748K	Tablet 25 mg	200	1	..	*57.73	28.60	Edecrin	MK

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Potassium-sparing agents								
• Aldosterone antagonists								
SPIRONOLACTONE								
CAUTION: <i>Serum electrolytes should be checked regularly.</i>								
Restricted benefit <i>Hyperaldosteronism, including refractory cardiac failure; Female hirsutism.</i>								
CAUTION: <i>Treatment of hirsutism with spironolactone should be considered only after all avenues of non-drug therapy have been explored.</i>								
<i>Appropriate contraceptive measures should be taken by women of child-bearing age in whom spironolactone therapy has been initiated.</i>								
2339D	Tablet 25 mg	100	5	.. B1.89	12.76 14.65	13.73 ^a 13.73 ^a	Spiractin 25 Aldactone	AF PH
2340E	Tablet 100 mg	100	5	.. B2.62	36.24 38.86	28.60 ^a 28.60 ^a	Spiractin 100 Aldactone	AF PH
• Other potassium-sparing agents								
AMILORIDE HYDROCHLORIDE								
CAUTION: <i>Serum electrolytes should be checked regularly.</i>								
3109P	Tablet 5 mg	100	1	..	*8.39	9.36	Kaluril	AF
Diuretics and potassium-sparing agents in combination								
• Low-ceiling diuretics and potassium-sparing agents								
HYDROCHLOROTHIAZIDE with AMILORIDE HYDROCHLORIDE								
CAUTION: <i>Serum electrolytes should be checked regularly.</i>								
1486F	Tablet 50 mg-5 mg	100	1	.. B3.44	*11.15 *14.59	12.12 12.12	^a Amizide ^a Moduretic	AF MK
HYDROCHLOROTHIAZIDE with TRIAMTERENE								
CAUTION: <i>Serum electrolytes should be checked regularly.</i>								
1280J	Tablet 25 mg-50 mg	100	1	..	11.15	12.12	Hydrene 25/50	AF
PERIPHERAL VASODILATORS								
Peripheral vasodilators								
• Other peripheral vasodilators								
PHENOXYBENZAMINE HYDROCHLORIDE								
Restricted benefit <i>Phaeochromocytoma; Neurogenic urinary retention.</i>								
1862B	Capsule 10 mg	100	5	..	52.27	28.60	Dibenyline	LM

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
BETA BLOCKING AGENTS								
Beta blocking agents								
• Beta blocking agents, non-selective								
OXPRENOLOL HYDROCHLORIDE								
2942W	Tablet 20 mg	100	5	..	8.28	9.25	Corbeton 20	AF
2961W	Tablet 40 mg	100	5	..	10.04	11.01	Corbeton 40	AF
PINDOLOL								
3062E	Tablet 5 mg	100	5	..	11.43	12.40	^a Barbloc 5	AF
				^B 2.80	14.23	12.40	^a Visken 5	NV
3065H	Tablet 15 mg	50	5	..	14.35	15.32	^a Barbloc 15	AF
				^B 2.80	17.15	15.32	^a Visken 15	NV
PROPRANOLOL HYDROCHLORIDE								
2565B	Tablet 10 mg	100	5	..	7.72	8.69	Deralin 10	AF
				^B 3.00	10.72	8.69	Inderal	AP
2566C	Tablet 40 mg	100	5	..	7.85	8.82	Deralin 40	AF
				^B 3.01	10.86	8.82	Inderal	AP
2899N	Tablet 160 mg	50	5	..	8.85	9.82	Deralin 160	AF
SOTALOL HYDROCHLORIDE								
For listings see Generic/Proprietary Index								
• Beta blocking agents, selective								
ATENOLOL								
1081X	Tablet 50 mg	30	5	..	9.82	10.79	^a Anselol 50 mg	DP
							^a Atehexal	HX
							^a Chem mart	CH
							Atenolol	
							^a GenRx Atenolol	GX
							^a Noten	AF
							^a Tensig	SI
							^a Terry White	TW
							Chemists	
							Atenolol	
				^B 3.55	13.37	10.79	^a Tenormin	AP
BISOPROLOL FUMARATE								
Authority required								
Moderate to severe heart failure in patients stabilised on conventional therapy which must include an ACE-inhibitor if tolerated.								
8604W	Tablet 2.5 mg	28	5	..	55.44	28.60	Bicor	AL
8605X	Tablet 5 mg	28	5	..	68.12	28.60	Bicor	AL
8606Y	Tablet 10 mg	28	5	..	83.95	28.60	Bicor	AL

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
METOPROLOL SUCCINATE								
Authority required								
<i>Moderate to severe heart failure in patients stabilised on conventional therapy which must include an ACE-inhibitor if tolerated.</i>								
8818D	Pack containing 15 tablets 23.75 mg (controlled release), 15 tablets 47.5 mg (controlled release) and 15 tablets 95 mg (controlled release)	1	..	..	95.96	28.60	Toprol-XL Titration Pack	AP
8732N	Tablet 23.75 mg (controlled release)	15	..	..	19.19	20.16	Toprol-XL 23.75	AP
8733P	Tablet 47.5 mg (controlled release)	30	5	..	72.99	28.60	Toprol-XL 47.5	AP
8734Q	Tablet 95 mg (controlled release)	30	5	..	90.04	28.60	Toprol-XL 95	AP
8735R	Tablet 190 mg (controlled release)	30	5	..	111.37	28.60	Toprol-XL 190	AP
METOPROLOL TARTRATE								
1324Q	Tablet 50 mg	100	5	..	10.57	11.54	^a Chem mart Metoprolol	CH
							^a GenRx Metoprolol	GX
							^a Metohexal Metoprolol	HX
							^a Metolol	DP
							^a Metrol 50	AW
							^a Minax 50	AF
							^a Terry White Chemists Metoprolol	TW
					^B 2.85 13.42	11.54	Lopresor 50	NV
					^B 3.26 13.83	11.54	^a Betaloc	AP
1325R	Tablet 100 mg	60	5	..	12.14	13.11	^a Chem mart Metoprolol	CH
							^a GenRx Metoprolol	GX
							^a Metohexal Metoprolol	HX
							^a Metolol	DP
							^a Metrol 100	AW
							^a Minax 100	AF
							^a Terry White Chemists Metoprolol	TW
					^B 2.85 14.99	13.11	Lopresor 100	NV
					^B 3.25 15.39	13.11	^a Betaloc	AP

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Alpha and beta blocking agents CARVEDILOL Authority required <i>Moderate to severe heart failure in patients stabilised on conventional therapy which must include an ACE-inhibitor if tolerated.</i>								
8742D	Pack containing 30 tablets 3.125 mg, 30 tablets 6.25 mg and 10 tablets 12.5 mg	1	..	..	66.36	28.60	Dilatrend Titration Pack	RO
Authority required <i>Moderate to severe heart failure in patients stabilised on conventional therapy which must include an ACE-inhibitor if tolerated; Patients receiving this drug as a pharmaceutical benefit prior to 1 August 2002.</i>								
8255L	Tablet 3.125 mg	30	..	..	18.92	19.89	^a Dilatrend 3.125 ^a Kredex	RO MD
8256M	Tablet 6.25 mg	60	5	..	71.73	28.60	^a Dilatrend 6.25 ^a Kredex	RO MD
8257N	Tablet 12.5 mg	60	5	..	88.46	28.60	^a Dilatrend 12.5 ^a Kredex	RO MD
8258P	Tablet 25 mg	60	5	..	109.39	28.60	^a Dilatrend 25 ^a Kredex	RO MD
LABETALOL HYDROCHLORIDE								
1566K	Tablet 100 mg	100	5	.. ^B 2.84	13.67 16.51	14.64	^a Presolol 100 ^a Trandate	AF SI
1567L	Tablet 200 mg	100	5	.. ^B 2.97	20.26 23.23	21.23	^a Presolol 200 ^a Trandate	AF SI

CALCIUM CHANNEL BLOCKERS

Selective calcium channel blockers with mainly vascular effects

• Dihydropyridine derivatives

NOTE:

The base-priced drugs in this therapeutic group are felodipine and nifedipine (except nifedipine controlled release tablet 20 mg).

AMLODIPINE BESYLATE								
2751T	Tablet 5 mg (base)	30	5	^T 4.62	25.12	21.47	Norvasc	PF
2752W	Tablet 10 mg (base)	30	5	^T 6.70	39.17	28.60	Norvasc	PF

AMLODIPINE BESYLATE

Authority required

Adverse effects occurring with all of the base-priced drugs;

Drug interactions occurring with all of the base-priced drugs;

Drug interactions expected to occur with all of the base-priced drugs;

Transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.

8923P	Tablet 5 mg (base)	30	5	..	25.12	26.09	Norvasc	PF
8924Q	Tablet 10 mg (base)	30	5	..	39.17	28.60	Norvasc	PF

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
FELODIPINE								
2361G	Tablet 2.5 mg (extended release)	30	5	..	15.39	16.36	^a Felodur ER 2.5 mg	AL
				^B 3.00	18.39	16.36	^a Plendil ER	AP
2366M	Tablet 5 mg (extended release)	30	5	..	19.38	20.35	^a Felodur ER 5 mg	AL
				^B 3.36	22.74	20.35	^a Plendil ER	AP
2367N	Tablet 10 mg (extended release)	30	5	..	30.94	28.60	^a Felodur ER 10 mg	AL
				^B 4.65	35.59	28.60	^a Plendil ER	AP
LERCANIDIPINE HYDROCHLORIDE								
8534E	Tablet 10 mg	30	5	^T 2.15	26.55	25.37	Zanidip	SM
8679T	Tablet 20 mg	30	5	^T 3.60	40.57	28.60	Zanidip	SM

LERCANIDIPINE HYDROCHLORIDE**Authority required***Adverse effects occurring with all of the base-priced drugs;**Drug interactions occurring with all of the base-priced drugs;**Drug interactions expected to occur with all of the base-priced drugs;**Transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.*

8939L	Tablet 10 mg	30	5	..	26.55	27.52	Zanidip	SM
8940M	Tablet 20 mg	30	5	..	40.57	28.60	Zanidip	SM
NIFEDIPINE								
1694E	Tablet 10 mg	60	5	..	18.96	19.93	^a Adefin 10	AF
							^a Nypine 10	AW
				^B 1.78	20.74	19.93	^a Adalat 10	BN
1695F	Tablet 20 mg	60	5	..	22.46	23.43	^a Adefin 20	AF
							^a Chem mart Nifedipine	CH
							^a GenRx Nifedipine	GX
							^a Nifehexal	HX
							^a Nyefax 20 mg	DP
							^a Nypine 20	AW
							^a Terry White Chemists Nifedipine	TW
				^B 3.31	25.77	23.43	^a Adalat 20	BN
8610E	Tablet 20 mg (controlled release)	30	5	^T 2.20	24.59	23.36	Adalat Oros 20mg	BN
1906H	Tablet 30 mg (controlled release)	30	5	..	23.79	24.76	^a Addos XR 30	AW
							^a Adefin XL 30	AF
				^B 3.10	26.89	24.76	^a Adalat Oros 30	BN
1907J	Tablet 60 mg (controlled release)	30	5	..	28.69	28.60	^a Addos XR 60	AW
							^a Adefin XL 60	AF
				^B 3.70	32.39	28.60	^a Adalat Oros 60	BN

continued ➤

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
<i>NIFEDIPINE—cont.</i>								
<u>Authority required</u>								
<i>Adverse effects occurring with all of the base-priced drugs;</i>								
<i>Drug interactions occurring with all of the base-priced drugs;</i>								
<i>Drug interactions expected to occur with all of the base-priced drugs;</i>								
<i>Transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.</i>								
8938K	Tablet 20 mg (controlled release)	30	5	..	24.59	25.56	Adalat Oros 20mg	BN
Selective calcium channel blockers with direct cardiac effects								
• Phenylalkylamine derivatives								
VERAPAMIL HYDROCHLORIDE								
<u>CAUTION:</u>								
The myocardial depressant effects of this drug and of beta-blocking drugs are additive.								
1248Q	Tablet 40 mg	100	5	..	11.46	12.43	^a Anpec 40	AF
							^a Verahexal	HX
				^B 1.00	12.46	12.43	^a Isoptin	AB
1250T	Tablet 80 mg	100	5	..	16.66	17.63	^a Anpec 80	AF
							^a Verahexal	HX
				^B 1.00	17.66	17.63	^a Isoptin	AB
1254B	Tablet 120 mg	100	5	..	22.02	22.99	^a Verahexal	HX
				^B 1.00	23.02	22.99	^a Isoptin	AB
1253Y	Tablet 160 mg	60	5	..	20.59	21.56	Isoptin	AB
2208F	Tablet 180 mg (sustained release)	30	5	..	14.36	15.33	^a Cordilox 180 SR	KN
				^B 3.00	17.36	15.33	^a Isoptin 180 SR	AB
1241H	Tablet 240 mg (sustained release)	30	5	..	17.57	18.54	^a Anpec SR	AF
							^a Cordilox SR	KN
				^B 3.00	20.57	18.54	^a Isoptin SR	AB
2206D	Capsule 160 mg (sustained release)	30	5	..	12.68	13.65	Veracaps SR	SI
2207E	Capsule 240 mg (sustained release)	30	5	..	17.70	18.67	Veracaps SR	SI
1060T	Injection 5 mg in 2 mL	5	..	..	10.64	11.61	Isoptin	AB
• Benzothiazepine derivatives								
DILTIAZEM HYDROCHLORIDE								
<u>CAUTION:</u>								
The myocardial depressant effects of this drug and of beta-blocking drugs are additive.								
1335G	Tablet 60 mg	90	5	..	19.50	20.47	^a Chem mart Diltiazem	CH
							^a Coras	AF
							^a Diltahexal	HX
							^a Dilzem 60 mg	DP
							^a GenRx Diltiazem	GX
							^a Terry White Chemists Diltiazem	TW
							^a Vasocardol	HP
				^B 2.64	22.14	20.47	^a Cardizem	AV

continued ➡

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
DILTIAZEM HYDROCHLORIDE—cont.							
1312C	Capsule 180 mg (controlled delivery)	30	5	..	21.12	22.09	^a Chem mart CH Diltiazem CD ^a Dilzem CD DP ^a GenRx Diltiazem CD GX ^a Terry White Chemists TW Diltiazem CD ^a Vasocardol CD HP Cardizem CD AV
				^B 2.64	23.76	22.09	
1313D	Capsule 240 mg (controlled delivery)	30	5	..	26.85	27.82	^a Chem mart CH Diltiazem CD ^a Dilzem CD DP ^a GenRx Diltiazem CD GX ^a Terry White Chemists TW Diltiazem CD ^a Vasocardol CD HP Cardizem CD AV
				^B 2.64	29.49	27.82	
8480H	Capsule 360 mg (controlled delivery)	30	5	..	32.95	28.60	^a Vasocardol CD HP Cardizem CD AV
				^B 3.21	36.16	28.60	

AGENTS ACTING ON THE RENIN-ANGIOTENSIN SYSTEM

ACE inhibitors, plain• **ACE inhibitors, plain****CAUTION:**

Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

NOTE:

The base-priced drugs in this therapeutic group are captopril, enalapril maleate, fosinopril sodium, lisinopril, perindopril erbumine, quinapril hydrochloride, ramipril (except ramipril capsule 10 mg) andtrandolapril.

CAPTOPRIL							
1147J	Tablet 12.5 mg	90	5	..	20.19	21.16	^a Acenorm AF 12.5 mg ^a Captohexal HX ^a Chem mart CH Captopril ^a GenRx Captopril GX ^a Terry White Chemists TW Captopril ^a Topace FM ^a DP ^B 2.46 22.65 21.16 ^a Capoten BQ

continued ➤

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
1148K	CAPTOPRIL—cont. Tablet 25 mg	90	5	..	27.10	28.07	^a Acenorm 25 mg AF ^a Captohexal HX ^a Chem mart CH Captopril ^a GenRx Captopril GX ^a Terry White TW Chemists Captopril ^a Topace FM ^a DP ^B 2.46 29.56 28.07 ^a Capoten BQ
1149L	Tablet 50 mg	90	5	..	48.86	28.60	^a Acenorm 50 mg AF ^a Captohexal HX ^a Chem mart CH Captopril ^a GenRx Captopril GX ^a Terry White TW Chemists Captopril ^a Topace FM ^a DP ^B 2.45 51.31 28.60 ^a Capoten BQ

CAPTOPRIL**Restricted benefit***For patients unable to take a solid dose form of an ACE inhibitor.*

8760C	Oral solution 5 mg per mL, 95 mL	‡1	5	..	61.61	28.60	Capoten	BQ
1370D	ENALAPRIL MALEATE Tablet 5 mg	30	5	..	14.86	15.83	^a Alphapril ^a Auspril ^a Chem mart Enalapril ^a Enahexal ^a Enalabell ^a Enalapril-DP 5mg ^a GenRx Enalapril ^a Terry White Chemists Enalapril ^B 2.45 ^B 3.15	AF SI CH HX BF DP GX TW AD MK

continued ➤

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ENALAPRIL MALEATE—cont.							
1368B	Tablet 10 mg	30	5	..	21.35	22.32	a Alphapril AF a Auspril SI a Chem mart CH Enalapril a Enahexal HX a Enalabell BF a Enalapril-DP DP 10mg a GenRx Enalapril GX a Terry White TW Chemists Enalapril B2.45 23.80 22.32 a Amprace 10 AD B3.15 24.50 22.32 a Renitec MK
1369C	Tablet 20 mg	30	5	..	25.89	26.86	a Alphapril AF a Auspril SI a Chem mart CH Enalapril a Enahexal HX a Enalabell BF a Enalapril-DP DP 20mg a GenRx Enalapril GX a Terry White TW Chemists Enalapril B2.46 28.35 26.86 a Amprace 20 AD B3.15 29.04 26.86 a Renitec 20 MK
FOSINOPRIL SODIUM							
1182F	Tablet 10 mg	30	5	..	19.02	19.99	Monopril BQ
1183G	Tablet 20 mg	30	5	..	26.10	27.07	Monopril BQ
LISINOPRIL							
2456G	Tablet 5 mg	30	5	..	16.34	17.31	a Chem mart CH Lisinopril a Fibsol 5 AW a GenRx Lisinopril GX a Liprace DP a Lisinobell BF a Lisinopril Hexal HX a Lisodur AF a Terry White TW Chemists Lisinopril B1.95 18.29 17.31 a Prinivil 5 MK B2.15 18.49 17.31 a Zestril AP

continued ➡

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
LISINOPRIL—cont.								
2457H	Tablet 10 mg	30	5	..	22.17	23.14	^a Chem mart Lisinopril	CH
							^a Fibsol 10	AW
							^a GenRx Lisinopril	GX
							^a Liprace	DP
							^a Lisinobell	BF
							^a Lisinopril Hexal	HX
							^a Lisodur	AF
							^a Terry White Chemists Lisinopril	TW
				^B 1.95	24.12	23.14	^a Prinivil 10	MK
				^B 2.15	24.32	23.14	^a Zestril	AP
2458J	Tablet 20 mg	30	5	..	26.68	27.65	^a Chem mart Lisinopril	CH
							^a Fibsol 20	AW
							^a GenRx Lisinopril	GX
							^a Liprace	DP
							^a Lisinobell	BF
							^a Lisinopril Hexal	HX
							^a Lisodur	AF
							^a Terry White Chemists Lisinopril	TW
				^B 1.95	28.63	27.65	^a Prinivil 20	MK
				^B 2.15	28.83	27.65	^a Zestril	AP
PERINDOPRIL ERBUMINE								
3050M	Tablet 2 mg	30	5	..	19.52	20.49	Coversyl	SE
3051N	Tablet 4 mg	30	5	..	24.81	25.78	Coversyl	SE
8704D	Tablet 8 mg	30	5	..	42.89	28.60	Coversyl	SE
QUINAPRIL HYDROCHLORIDE								
1968N	Tablet 5 mg (base)	30	5	..	17.13	18.10	^a Accupril	PF
							^a Asig	SI
1969P	Tablet 10 mg (base)	30	5	..	22.21	23.18	^a Accupril	PF
							^a Asig	SI
1970Q	Tablet 20 mg (base)	30	5	..	26.26	27.23	^a Accupril	PF
							^a Asig	SI
RAMIPRIL								
1944H	Tablet 1.25 mg	30	5	..	13.61	14.58	^a Ramace 1.25 mg	ML
				^B 2.00	15.61	14.58	^a Tritace 1.25 mg	AV
1945J	Tablet 2.5 mg	30	5	..	17.60	18.57	^a Ramace 2.5 mg	ML
				^B 2.01	19.61	18.57	^a Tritace 2.5 mg	AV
1946K	Tablet 5 mg	30	5	..	20.78	21.75	^a Ramace 5 mg	ML
				^B 2.00	22.78	21.75	^a Tritace 5 mg	AV
8470T	Capsule 10 mg	30	5	^T 1.35	34.23	28.60	Tritace 10 mg	AV

continued ➡

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
8668F	RAMIPRIL—cont. Pack containing 7 tablets 2.5 mg, 21 tablets 5 mg and 10 capsules 10 mg	‡1	..	..	28.24	28.60	Tritace Titration Pack	AV

RAMIPRIL**Authority required***Adverse effects occurring with all of the base-priced drugs;**Drug interactions occurring with all of the base-priced drugs;**Drug interactions expected to occur with all of the base-priced drugs;**Transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.*

8937J	Capsule 10 mg	30	5	..	34.23	28.60	Tritace 10 mg	AV
TRANSDOLAPRIL								
2791X	Capsule 500 micrograms	28	5	..	14.60	15.57	^a Gopten	AB
				^B 1.00	15.60	15.57	^a Odrik	KN
2792Y	Capsule 1 mg	28	5	..	20.41	21.38	^a Gopten	AB
				^B 1.01	21.42	21.38	^a Odrik	KN
2793B	Capsule 2 mg	28	5	..	22.17	23.14	^a Gopten	AB
				^B 0.99	23.16	23.14	^a Odrik	KN
8758Y	Capsule 4 mg	28	5	..	36.53	28.60	Gopten	AB

ACE inhibitors, combinations**CAUTION:**

Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

• **ACE inhibitors and diuretics****ENALAPRIL MALEATE with HYDROCHLOROTHIAZIDE****Restricted benefit***Hypertension in patients who are not adequately controlled with 20 mg enalapril maleate.*

8477E	Tablet 20 mg-6 mg	30	5	..	30.18	28.60	Renitec Plus 20/6	MK

FOSINOPRIL SODIUM with HYDROCHLOROTHIAZIDE**Restricted benefit***Hypertension in patients who are not adequately controlled with either hydrochlorothiazide or fosinopril sodium monotherapy.*

8400D	Tablet 10 mg-12.5 mg	30	5	..	23.01	23.98	Monoplus 10/12.5	BQ
8401E	Tablet 20 mg-12.5 mg	30	5	..	31.55	28.60	Monoplus 20/12.5	BQ

PERINDOPRIL ERBUMINE with INDAPAMIDE HEMIHYDRATE**Restricted benefit***Hypertension in patients who are not adequately controlled with either indapamide hemihydrate or perindopril erbumine monotherapy.*

8449Q	Tablet 4 mg-1.25 mg	30	5	..	32.56	28.60	Coversyl Plus 4/1.25	SE

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
QUINAPRIL HYDROCHLORIDE with HYDROCHLOROTHIAZIDE								
<u>Restricted benefit</u>								
<i>Hypertension in patients who are not adequately controlled with either hydrochlorothiazide or quinapril hydrochloride monotherapy.</i>								
8589C	Tablet 10 mg (base)-12.5 mg	30	5	..	24.41	25.38	Accuretic 10/12.5mg	PF
8590D	Tablet 20 mg (base)-12.5 mg	30	5	..	28.46	28.60	Accuretic 20/12.5mg	PF
Angiotensin II antagonists, plain								
• Angiotensin II antagonists, plain								
CANDESARTAN CILEXETIL								
8295N	Tablet 4 mg	30	5	..	19.19	20.16	Atacand	AP
8296P	Tablet 8 mg	30	5	..	22.99	23.96	Atacand	AP
8297Q	Tablet 16 mg	30	5	..	28.77	28.60	Atacand	AP
EPROSARTAN MESYLATE								
8397Y	Tablet 400 mg (base)	56	5	..	*28.31	28.60	Teveten	SM
8447N	Tablet 600 mg (base)	28	5	..	28.30	28.60	Teveten	SM
IRBESARTAN								
8246B	Tablet 75 mg	30	5	..	19.27	20.24	Avapro Karvea	BQ SW
8247C	Tablet 150 mg	30	5	..	23.26	24.23	Avapro Karvea	BQ SW
8248D	Tablet 300 mg	30	5	..	28.29	28.60	Avapro Karvea	BQ SW
TELMISARTAN								
8355R	Tablet 40 mg	28	5	..	19.60	20.57	Micardis	BY
8356T	Tablet 80 mg	28	5	..	26.67	27.64	Micardis	BY
Angiotensin II antagonists, combinations								
• Angiotensin II antagonists and diuretics								
CANDESARTAN CILEXETIL with HYDROCHLOROTHIAZIDE								
<u>Restricted benefit</u>								
<i>Hypertension in patients who are not adequately controlled with 16 mg candesartan cilexetil.</i>								
8504N	Tablet 16 mg-12.5 mg	30	5	..	30.97	28.60	Atacand Plus 16/12.5	AP
EPROSARTAN MESYLATE with HYDROCHLOROTHIAZIDE								
<u>Restricted benefit</u>								
<i>Hypertension in patients who are not adequately controlled with either hydrochlorothiazide or eprosartan mesylate monotherapy.</i>								
8624X	Tablet 600 mg (base)-12.5 mg	28	5	..	30.35	28.60	Teveten Plus 600/12.5	SM

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
<i>IRBESARTAN with HYDROCHLOROTHIAZIDE</i>								
<u>Restricted benefit</u>								
<i>Hypertension in patients who are not adequately controlled with either hydrochlorothiazide or irbesartan monotherapy.</i>								
8404H	Tablet 150 mg-12.5 mg	30	5	..	25.46	26.43	Avapro HCT 150/12.5 Karvezide 150/12.5	BQ SW
8405J	Tablet 300 mg-12.5 mg	30	5	..	30.49	28.60	Avapro HCT 300/12.5 Karvezide 300/12.5	BQ SW
<i>TELMISARTAN with HYDROCHLOROTHIAZIDE</i>								
<u>Restricted benefit</u>								
<i>Hypertension in patients who are not adequately controlled with either hydrochlorothiazide or telmisartan monotherapy.</i>								
8622T	Tablet 40 mg-12.5 mg	28	5	..	21.65	22.62	Micardis Plus 40/12.5 mg	BY
8623W	Tablet 80 mg-12.5 mg	28	5	..	28.72	28.60	Micardis Plus 80/12.5 mg	BY

SERUM LIPID REDUCING AGENTS

GENERAL STATEMENT FOR LIPID-LOWERING DRUGS
PRESCRIBED AS PHARMACEUTICAL BENEFITS


Use the following criteria to determine patient eligibility for subsidisation under the PBS for the following drugs:

- atorvastatin calcium
- fluvastatin sodium
- pravastatin sodium
- simvastatin
- fenofibrate
- gemfibrozil

By writing a PBS prescription, the prescriber is certifying the patient satisfies the qualifying criteria set out below and the use is in accordance with the registered indications which differ between agents in this class. Refer to the current Product Information. Patients already established on a particular lipid-lowering drug, where use satisfies the PBS qualifying criteria, but is outside the registered indications for that drug, are not required to switch to another drug in the class to retain PBS eligibility.

Use the flow-chart and table below to determine whether your patient satisfies criteria for subsidisation under the PBS. Document how the patient meets each of these steps in the patient record. Lipid levels must be measured at an accredited laboratory.

continued ➞

CARDIOVASCULAR SYSTEM —cont.**QUALIFYING CRITERIA**

PATIENT CATEGORY	LIPID LEVEL FOR PBS SUBSIDY
<i>Patients with existing coronary heart disease</i>	cholesterol > 4 mmol/L
<i>Other patients at high risk with one or more of the following:</i> • diabetes mellitus • familial hypercholesterolaemia • family history of coronary heart disease (first degree relative less than 60 years of age) • hypertension • peripheral vascular disease	cholesterol > 6.5 mmol/L or cholesterol > 5.5 mmol/L and HDL < 1 mmol/L
<i>Patients with HDL < 1 mmol/L</i>	cholesterol > 6.5 mmol/L
<i>Patients not eligible under the above:</i> • men 35 to 75 years • post-menopausal women up to 75 years	cholesterol > 7.5 mmol/L or triglyceride > 4 mmol/L
<i>Other patients not included in the above</i>	cholesterol > 9 mmol/L or triglyceride > 8 mmol/L

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Cholesterol and triglyceride reducers								
• HMG CoA reductase inhibitors								
ATORVASTATIN CALCIUM								
<u>Restricted benefit</u>								
<i>For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.</i>								
8213G	Tablet 10 mg (atorvastatin)	30	5	..	42.00	28.60	Lipitor	PF
8214H	Tablet 20 mg (atorvastatin)	30	5	..	59.70	28.60	Lipitor	PF
8215J	Tablet 40 mg (atorvastatin)	30	5	..	82.12	28.60	Lipitor	PF
8521L	Tablet 80 mg (atorvastatin)	30	5	..	115.36	28.60	Lipitor	PF
FLUVASTATIN SODIUM								
<u>Restricted benefit</u>								
<i>For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.</i>								
8023G	Capsule 20 mg (fluvastatin)	28	5	..	25.78	26.75	Lescol Vastin	NV AP
8024H	Capsule 40 mg (fluvastatin)	28	5	..	30.52	28.60	Lescol Vastin	NV AP
PRAVASTATIN SODIUM								
<u>Restricted benefit</u>								
<i>For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.</i>								
2833D	Tablet 10 mg	30	5	..	29.85	28.60	^a Lipostat 10	AW
				^B 1.75	31.60	28.60	^a Pravachol	BQ
2834E	Tablet 20 mg	30	5	..	44.63	28.60	^a Lipostat 20	AW
				^B 1.74	46.37	28.60	^a Pravachol	BQ
8197K	Tablet 40 mg	30	5	..	66.87	28.60	^a Lipostat 40	AW
				^B 1.75	68.62	28.60	^a Pravachol	BQ
8829Q	Tablet 80 mg	30	5	..	99.97	28.60	^a Lipostat 80	AW
				^B 1.75	101.72	28.60	^a Pravachol	BQ
SIMVASTATIN								
<u>Restricted benefit</u>								
<i>For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.</i>								
2013Y	Tablet 5 mg	30	5	..	26.98	27.95	^a Simvabell	BF
							^a Simvahexal	HX
							^a Simvastatin-RL	RE
							^a Zimstat	AF
				^B 0.71	27.69	27.95	^a Lipex 5	AD
							^a Zocor	MK

continued ➤

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
SIMVASTATIN—cont.							
2011W	Tablet 10 mg	30	5	..	35.48	28.60	^a Chem mart CH Simvastatin ^a GenRx GX Simvastatin ^a Simvabell BF ^a Simvahexal HX ^a Simvar 10 AW ^a Simvastatin-DP DP ^a Simvastatin-RL RE ^a Terry White TW Chemists Simvastatin ^a Zimstat AF ^a Lipex 10 AD ^a Zocor MK
2012X	Tablet 20 mg	30	5	..	48.93	28.60	^a Chem mart CH Simvastatin ^a GenRx GX Simvastatin ^a Simvabell BF ^a Simvahexal HX ^a Simvar 20 AW ^a Simvastatin-DP DP ^a Simvastatin-RL RE ^a Terry White TW Chemists Simvastatin ^a Zimstat AF ^a Lipex 20 AD ^a Zocor MK
8173E	Tablet 40 mg	30	5	..	68.33	28.60	^a Chem mart CH Simvastatin ^a GenRx GX Simvastatin ^a Simvabell BF ^a Simvahexal HX ^a Simvar 40 AW ^a Simvastatin-DP DP ^a Simvastatin-RL RE ^a Terry White TW Chemists Simvastatin ^a Zimstat AF ^a Lipex 40 AD ^a Zocor MK
				^B 0.71	36.19	28.60	
				^B 0.70	49.63	28.60	
				^B 0.70	69.03	28.60	

continued ↗

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
SIMVASTATIN—cont.							
8313M	Tablet 80 mg	30	5	..	96.11	28.60	^a Chem mart CH ^a Simvastatin ^a GenRx GX ^a Simvastatin ^a Simvabell BF ^a Simvahexal HX ^a Simvar 80 AW ^a Simvastatin-DP DP ^a Simvastatin-RL RE ^a Terry White Chemists TW ^a Simvastatin ^a Zimstat AF ^a Lipex 80 AD ^a Zocor MK
					^B 0.70	96.81	28.60

• **Fibrates****FENOFIBRATE****NOTE:**

The risk of serious muscle toxicity is increased if fenofibrate is used concomitantly with HMG CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution in patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and patients monitored closely for chronic signs of muscle toxicity.

Restricted benefit

For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.

8721B	Capsule 67 mg	60	5	..	28.10	28.60	Lipidil LF
8722C	Tablet 160 mg	30	5	..	39.79	28.60	Lipidil LF

NOTE:

Three of the 67 mg capsules are considered to be bioequivalent to 1 of the 160 mg tablets.

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
GEMFIBROZIL								
NOTE:								
<i>The risk of serious muscle toxicity is increased if gemfibrozil is used concomitantly with HMG CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution in patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and patients monitored closely for chronic signs of muscle toxicity.</i>								
Restricted benefit								
<i>For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.</i>								
1453L	Tablet 600 mg	60	5	..	39.79	28.60	^a Ausgem	SI
							^a Chem mart	CH
							Gemfibrozil	
							^a Gemhexal	HX
							^a GenRx	GX
							Gemfibrozil	
							^a Jezil	AF
							^a Lipazil 600 mg	DP
							^a Terry White	TW
							Chemists	
							Gemfibrozil	
				^B 3.49	43.28	28.60	^a Lopid	PF
• Bile acid sequestrants								
CHOLESTYRAMINE								
2967E	Sachets 4.7 g (equivalent to 4 g cholestyramine), 50	2	5	..	*57.51	28.60	Questran Lite	BQ
2978R	Sachets 9.4 g (equivalent to 8 g cholestyramine), 50	‡1	5	..	54.37	28.60	Questran Lite	BQ
COLESTIPOL HYDROCHLORIDE								
1224K	Sachets 5 g, 120	‡1	5	..	68.06	28.60	Colestid	PH

CARDIOVASCULAR SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of Qty Rpts Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
• Other cholesterol and triglyceride reducers EZETIMIBE Authority required <i>Patients eligible for PBS-subsidised lipid-lowering medication (according to the criteria set out in the General Statement for Lipid-Lowering Drugs):</i> <i>(a) where treatment with an HMG CoA reductase inhibitor (statin) is contraindicated; or</i> <i>(b) where treatment with an HMG CoA reductase inhibitor (statin) is unsuitable because the patient developed a clinically important product-related adverse event during treatment with a statin, and required discontinuation of all statin treatment.</i> <i>A clinically important product-related adverse event is defined as follows:</i> <i>(i) Severe myalgia (muscle symptoms without CK elevation) which is proven to be temporally associated with statin treatment; or</i> <i>(ii) Myositis (clinically important CK elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or</i> <i>(iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.</i> Authority required <i>Homozygous sitosterolaemia;</i> <i>Patients with homozygous familial hypercholesterolaemia who are eligible for PBS-subsidised lipid-lowering medication (according to the criteria set out in the General Statement for Lipid-Lowering Drugs), in combination with an HMG CoA reductase inhibitor (statin).</i> Authority required <i>Initial treatment for co-administration with an HMG CoA reductase inhibitor (statin) in patients whose cholesterol levels are inadequately controlled with a statin and who have:</i> <i>(a) coronary heart disease; or</i> <i>(b) diabetes mellitus.</i> <i>Inadequate control with a statin is defined as a cholesterol level in excess of the initial threshold for PBS-subsidy according to the General Statement for Lipid-Lowering Drugs after at least 3 months of treatment at a daily dose of 40 mg or greater of a statin.</i> <i>The cholesterol level after 3 months of treatment with a statin and the dose of the statin must be provided at the time of application. The cholesterol level results provided must be no more than 1 month old at the time of application;</i> <i>Continuing treatment for co-administration with HMG CoA reductase inhibitors (statins) in patients with coronary heart disease or diabetes mellitus whose cholesterol levels were inadequately controlled with a statin, where the patient has previously been issued with an authority prescription for this drug.</i>					
8757X	Tablet 10 mg	30 5 ..	71.45	28.60	Ezetrol MK

DERMATOLOGICALS

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTIFUNGALS FOR DERMATOLOGICAL USE								
Antifungals for systemic use								
• Antifungals for systemic use								
GRISEOFULVIN								
1460W	Tablet 125 mg	100	2	..	18.67	19.64	Grisovin	SI
2982Y	Tablet 500 mg	28	2	..	19.16	20.13	Grisovin 500	SI
TERBINAFINE HYDROCHLORIDE								
Authority required								
➤ <i>Proximal or extensive (greater than 80% nail involvement) onychomycosis due to dermatophyte infection where topical treatment has failed. This infection must be proven by microscopy or culture and confirmed by an Approved Pathology Authority. The date of the pathology report must be provided at the time of application and must not be more than 12 months old.</i>								
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
2804N	Tablet 250 mg (base)	42	1	..	137.58	28.60	^a GenRx Terbinafine	GX
							^a Tamsil	AW
							^a Terbihexal	HX
							^a Terbinafine-DP	DP
							^a Zabel	AF
				B2.00	139.58	28.60	^a Lamisil	NV
ANTIPSORIATICS								
Antipsoriatics for topical use								
• Tars								
COAL TAR - PREPARED								
8864M	Gel 10 mg per g (1%), 100 mL	‡1	2	..	31.10	28.60	Exorex	EP
• Other antipsoriatics for topical use								
CALCIPOTRIOL								
Restricted benefit								
<i>Chronic stable plaque type psoriasis vulgaris.</i>								
8291J	Ointment 50 micrograms per g (0.005%), 30 g	‡1	1	..	22.00	22.97	Daivonex	CS
Antipsoriatics for systemic use								
• Retinoids for treatment of psoriasis								
ACITRETIN								
CAUTION:								
<i>This drug is a potent teratogen—pregnancy should be avoided for at least two years after cessation of therapy.</i>								
NOTE:								
<i>Care must be taken to comply with the provisions of State/Territory law when prescribing acitretin.</i>								
Authority required								
<i>Severe intractable psoriasis;</i>								
<i>Severe forms of disorders of keratinisation.</i>								
2019G	Capsule 10 mg	100	2	..	210.16	28.60	Neotigason	RO

continued ➤

DERMATOLOGICALS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
2020H	ACITRETIN—cont. Capsule 25 mg	100	2	..	403.86	28.60	Neotigason	RO

ANTIBIOTICS AND CHEMOTHERAPEUTICS FOR DERMATOLOGICAL USE

Chemotherapeutics for topical use• **Sulfonamides**

**SILVER SULFADIAZINE with
CHLORHEXIDINE GLUCONATE**

Restricted benefit

*Prevention and treatment of infection in partial or full skin thickness loss due to burns;
Prevention and treatment of infection in partial or full skin thickness loss due to
epidermolysis bullosa;
Stasis ulcers.*

1996C	Cream 10 mg-2 mg per g (1%- 0.2%), 50 g	‡1	..	..	15.84	16.81	Silvazine	SN
1997D	Cream 10 mg-2 mg per g (1%- 0.2%), 100 g	‡1	..	..	18.60	19.57	Silvazine	SN

CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS

Corticosteroids, plain• **Corticosteroids, weak (group I)**

HYDROCORTISONE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

1495Q	Cream 10 mg per g (1%), 50 g	‡1	1	..	7.05	8.02	Egocort Cream 1%	EO

HYDROCORTISONE ACETATE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

2887Y	Cream 10 mg per g (1%), 30 g	‡1	1	..	6.59	7.56	a Cortic-DS 1%	FM
				b 1.30	7.89	7.56	a Sigmacort	SI
2881P	Cream 10 mg per g (1%), 50 g	‡1	1	..	7.05	8.02	Cortef	DT
				b 1.30	8.35	8.02	a Cortic-DS 1%	FM
							a Sigmacort	SI
2888B	Topical ointment 10 mg per g (1%), 30 g	‡1	1	..	6.59	7.56	a Cortic-DS 1%	FM
				b 1.30	7.89	7.56	a Sigmacort	SI
2882Q	Topical ointment 10 mg per g (1%), 50 g	‡1	1	..	7.05	8.02	a Cortic-DS 1%	FM
				b 1.30	8.35	8.02	a Sigmacort	SI

[For other listings for this drug see Generic/Proprietary Index]

DERMATOLOGICALS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Corticosteroids, moderately potent (group II)								
TRIAMCINOLONE ACETONIDE								
<u>Restricted benefit</u>								
<i>Treatment of corticosteroid-responsive dermatoses.</i>								
2117K	Cream 200 micrograms per g (0.02%), 100 g	2	..	..	*13.31 B2.60 *15.91	14.28 14.28	a Tricortone a Aristocort 0.02%	FM SI
2118L	Ointment 200 micrograms per g (0.02%), 100 g	2	..	..	*13.31 B2.60 *15.91	14.28 14.28	a Tricortone a Aristocort 0.02%	FM SI
• Corticosteroids, potent (group III)								
BETAMETHASONE DIPROPIONATE								
<u>Restricted benefit</u>								
<i>Treatment of corticosteroid-responsive dermatoses.</i>								
1115Q	Cream 500 micrograms (base) per g (0.05%), 15 g	‡1	1	..	7.12	8.09	a Diprosone a Eleuphrat	SH EX
1119X	Ointment 500 micrograms (base) per g (0.05%), 15 g	‡1	1	..	7.12	8.09	a Diprosone a Eleuphrat	SH EX
BETAMETHASONE VALERATE								
<u>Restricted benefit</u>								
<i>Treatment of corticosteroid-responsive dermatoses.</i>								
2812B	Cream 200 micrograms (base) per g (0.02%), 100 g	2	..	..	*13.31 B2.60 *15.91	14.28 14.28	a Antroquoril a Celestone-M b Cortival 1/5 b Betnovate 1/5	EX SH FM SI
2813C	Cream 500 micrograms (base) per g (0.05%), 15 g	‡1	1	..	6.91 B1.29 8.20	7.88 7.88	b Cortival 1/2 b Betnovate 1/2	FM SI
2820K	Ointment 200 micrograms (base) per g (0.02%), 100 g	2	..	..	*13.31	14.28	a Antroquoril a Celestone-M	EX SH
2815E	Ointment 500 micrograms (base) per g (0.05%), 15 g	‡1	1	..	6.91 B1.29 8.20	7.88 7.88	b Cortival 1/2 b Betnovate 1/2	FM SI
METHYLPREDNISOLONE ACEPONATE								
<u>Restricted benefit</u>								
<i>Treatment of corticosteroid-responsive dermatoses.</i>								
8054X	Cream 1 mg per g (0.1%), 15 g	‡1	..	..	10.40	11.37	Advantan	SC
8055Y	Ointment 1 mg per g (0.1%), 15 g	‡1	..	..	10.40	11.37	Advantan	SC
8128T	Fatty ointment 1 mg per g (0.1%), 15 g	‡1	..	..	10.40	11.37	Advantan	SC
<u>Restricted benefit</u>								
<i>Eczema.</i>								
8618N	Lotion 1 mg per g (0.1%), 20 g	‡1	..	..	11.16	12.13	Advantan	SC

DERMATOLOGICALS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MOMETASONE FUROATE								
Restricted benefit								
<i>Treatment of corticosteroid-responsive dermatoses.</i>								
1913Q	Cream 1 mg per g (0.1%), 15 g	‡1	..	..	10.40	11.37	^a Elocon ^a Novasone	SH EX
1915T	Ointment 1 mg per g (0.1%), 15 g	‡1	..	..	10.40	11.37	^a Elocon ^a Novasone	SH EX
8043H	Lotion 1 mg per g (0.1% w/w), 30 mL	‡1	..	..	14.43	15.40	^a Elocon ^a Novasone	SH EX

ANTI-ACNE PREPARATIONS

Anti-acne preparations for systemic use

- Retinoids for treatment of acne

ISOTRETINOIN**CAUTION:**

This drug causes birth defects. Isotretinoin has been reported to cause other frequent and potentially serious toxicity.

NOTE:

Care must be taken to comply with the provisions of State/Territory law when prescribing isotretinoin.

Authority required

Severe cystic acne not responsive to other therapy.

NOTE:

No applications for increased maximum quantities and/or repeats will be authorised.

2591J	Capsule 10 mg	60	3	..	97.68	28.60	^a GenRx Isotretinoin ^a Isohexal ^a Oratane ^a Roaccutane	GX HX DP RO
2592K	Capsule 20 mg	60	3	..	149.61	28.60	^a Chem mart Isotretinoin ^a GenRx Isotretinoin ^a Isohexal ^a Oratane ^a Terry White Chemists Isotretinoin ^a Roaccutane	CH GX HX DP TW RO
				^B 2.50	152.11	28.60		

OTHER DERMATOLOGICAL PREPARATIONS

Other dermatological preparations

- Other dermatologicals

DAPSONE

8801F	Tablet 25 mg	100	1	..	102.06	28.60	LM
1272Y	Tablet 100 mg	100	1	..	115.75	28.60	LM

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
	<u>PIMECROLIMUS</u>						
	<u>Authority required</u>						
	<i>Treatment of facial or eyelid atopic dermatitis in patients aged between 3 months and 18 years with 1 or more of the following contraindications to topical corticosteroids:</i>						
	<i>(i) perioral dermatitis;</i>						
	<i>(ii) periorbital dermatitis;</i>						
	<i>(iii) rosacea;</i>						
	<i>(iv) epidermal atrophy;</i>						
	<i>(v) dermal atrophy;</i>						
	<i>(vi) allergy to topical corticosteroids;</i>						
	<i>(vii) cataracts;</i>						
	<i>(viii) glaucoma;</i>						
	<i>(ix) raised intraocular pressure.</i>						
	<i>The following information must be provided at the time of application:</i>						
	<i>(i) the patient's date of birth; and</i>						
	<i>(ii) a description of the contraindication to topical corticosteroids; and</i>						
	<i>(iii) where subsidy is requested on the basis of topical corticosteroid allergy, the date and outcome of epicutaneous patch testing.</i>						
	<u>Authority required</u>						
	<i>Short-term (up to 3 weeks) intermittent treatment of atopic dermatitis of the face or eyelids in patients aged between 3 months and 18 years who fail to achieve satisfactory disease control with intermittent topical corticosteroid therapy, and where more than 3 months have passed since the initial diagnosis of atopic dermatitis.</i>						
	<i>Failure to achieve satisfactory disease control with intermittent topical corticosteroid therapy is manifest by:</i>						
	<i>(i) failure of the facial skin to clear despite at least 2 weeks of topical hydrocortisone 1% applied every day; or</i>						
	<i>(ii) failure of the facial skin to clear despite at least 1 week of a moderate or potent topical corticosteroid applied every day; or</i>						
	<i>(iii) clearing of the facial skin with at least 2 weeks of topical hydrocortisone 1% applied every day, but almost immediate and significant flare in facial disease (within 48 hours) upon stopping topical corticosteroids, occurring on at least 2 consecutive occasions; or</i>						
	<i>(iv) clearing of the facial skin with at least 1 week of a moderate or potent topical corticosteroid applied every day, but almost immediate and significant flare in facial disease (within 48 hours) upon stopping topical corticosteroids, occurring on at least 2 consecutive occasions.</i>						
	<i>The following information must be provided at the time of application:</i>						
	<i>(i) the patient's date of birth; and</i>						
	<i>(ii) the duration of atopic dermatitis since initial diagnosis; and</i>						
	<i>(iii) the name, quantity and duration of topical corticosteroid therapy used on the face or eyelids in the last month.</i>						
8802G	Cream 10 mg per g (1%), 15 g	₹1	1	..	31.81	28.60	Elidel NV
	<u>NOTE:</u>						
	<i>No applications for increased maximum quantities and/or repeats will be authorised. Only 1 authority application per 6 months, per patient, will be authorised.</i>						

GENITO URINARY SYSTEM AND SEX HORMONES

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
OTHER GYNECOLOGICALS								
Contraceptives for topical use								
• Intrauterine contraceptives								
LEVONORGESTREL								
Restricted benefit								
Contraception.								
8633J	Intrauterine drug delivery system 52 mg (releasing approximately 20 micrograms per 24 hours)	1	..	..	220.53	28.60	Mirena	SC
Other gynecologicals								
• Prolactine inhibitors								
BROMOCRIPTINE MESYLATE								
Restricted benefit								
Prevention of the onset of lactation in the puerperium for medical reasons.								
1444B	Tablet 2.5 mg (base)	30	..	..	19.15	20.12	^a Bromohexal	HX
							^a Krypton 2.5	AF
				^B 2.94	22.09	20.12	^a Parlodel	NV
Restricted benefit								
Acromegaly;								
Parkinson's disease;								
Pathological hyperprolactinaemia where surgery is not indicated;								
Pathological hyperprolactinaemia where surgery has already been used with incomplete resolution;								
Pathological hyperprolactinaemia where radiotherapy is not indicated;								
Pathological hyperprolactinaemia where radiotherapy has already been used with incomplete resolution.								
1443Y	Tablet 2.5 mg (base)	60	5	..	33.54	28.60	^a Bromohexal	HX
							^a Krypton 2.5	AF
				^B 3.00	36.54	28.60	^a Parlodel	NV
1446D	Capsule 5 mg (base)	60	5	..	61.36	28.60	^a Bromohexal	HX
							^a Krypton 5	AF
				^B 3.00	64.36	28.60	^a Parlodel	NV
1445C	Capsule 10 mg (base)	100	5	..	207.04	28.60	^a Krypton 10	AF
				^B 3.00	210.04	28.60	^a Parlodel	NV
CABERGOLINE								
Restricted benefit								
Prevention of the onset of lactation in the puerperium for medical reasons.								
8115D	Tablet 500 micrograms	2	..	..	25.10	26.07	Dostinex	PH

continued ➡

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CABERGOLINE—cont.								
Authority required								
<i>Pathological hyperprolactinaemia where surgery is not indicated;</i>								
<i>Pathological hyperprolactinaemia where surgery has already been used with incomplete resolution;</i>								
<i>Pathological hyperprolactinaemia where radiotherapy is not indicated;</i>								
<i>Pathological hyperprolactinaemia where radiotherapy has already been used with incomplete resolution.</i>								
8114C	Tablet 500 micrograms	8	5	..	76.88	28.60	Dostinex	PH
QUINAGOLIDE HYDROCHLORIDE								
Authority required								
<i>Pathological hyperprolactinaemia where surgery is not indicated;</i>								
<i>Pathological hyperprolactinaemia where surgery has already been used with incomplete resolution;</i>								
<i>Pathological hyperprolactinaemia where radiotherapy is not indicated;</i>								
<i>Pathological hyperprolactinaemia where radiotherapy has already been used with incomplete resolution.</i>								
8860H	Pack containing 3 tablets 25 micrograms (base) and 3 tablets 50 micrograms (base)	‡1	..	..	9.73	10.70	Norprolac	FP
8822H	Tablet 75 micrograms (base)	30	5	..	54.61	28.60	Norprolac	FP

SEX HORMONES AND MODULATORS OF THE GENITAL SYSTEM

Hormonal contraceptives for systemic use• **Progestogens and estrogens, fixed combinations**

LEVONORGESTREL with ETHINYLOESTRADIOL								
1456P	Pack containing 21 tablets 125 micrograms-50 micrograms and 7 inert tablets	4	2	..	*15.31	16.28	Microgynon 50 ED	SC
1393H	Tablets 150 micrograms- 30 micrograms, 21	4	2	^B 9.76	*25.07	16.28	^a Microgynon 30	SC
1394J	Pack containing 21 tablets 150 micrograms-30 micrograms and 7 inert tablets	4	2	..	*15.31	16.28	^a Levlen ED	SY
				^B 9.76	*25.07	16.28	^b Monofeme 28	WX
							^a Microgynon 30 ED	SC
							^b Nordette 28	WY
NORETHISTERONE with ETHINYLOESTRADIOL								
2772X	Tablets 500 micrograms- 35 micrograms, 21	4	2	^B 7.96	*23.27	16.28	^a Brevinor	PH
2774B	Pack containing 21 tablets 500 micrograms-35 micrograms and 7 inert tablets	4	2	..	*15.31	16.28	^a Norimin 28 Day	KR
				^B 7.96	*23.27	16.28	^a Brevinor	PH
2773Y	Tablets 1 mg-35 micrograms, 21	4	2	^B 7.96	*23.27	16.28	^a Brevinor-1	PH
2775C	Pack containing 21 tablets 1 mg- 35 micrograms and 7 inert tablets	4	2	..	*15.31	16.28	^a Norimin-1 28 Day	KR
				^B 7.96	*23.27	16.28	^a Brevinor-1	PH

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
	NORETHISTERONE with MESTRANOL							
3176E	Tablets 1 mg-50 micrograms, 21	4	2	..	*15.31	16.28	Norinyl-1	PH
3179H	Pack containing 21 tablets 1 mg- 50 micrograms and 7 inert tablets	4	2	..	*15.31	16.28	Norinyl-1/28	PH
• Progestogens and estrogens, sequential preparations								
	LEVONORGESTREL with ETHINYLOESTRADIOL							
1392G	Pack containing 6 tablets	4	2	..	*15.31	16.28	^a Logynon ED	SY
	50 micrograms-30 micrograms,						^b Trifeme 28	WX
	5 tablets 75 micrograms-			^B 9.76	*25.07	16.28	^b Triphasil 28	WY
	40 micrograms, 10 tablets						^a Triquilar ED	SC
	125 micrograms-30 micrograms and 7 inert tablets							
	NORETHISTERONE with ETHINYLOESTRADIOL							
2776D	Pack containing 12 tablets	4	2	..	*15.31	16.28	^a Improvil 28 Day	KR
	500 micrograms-35 micrograms,			^B 7.96	*23.27	16.28	^a Synphasic	PH
	9 tablets 1 mg-35 micrograms and 7 inert tablets							
• Progestogens								
	ETONOGESTREL							
8487Q	Subcutaneous implant 68 mg	1	..	..	220.53	28.60	Implanon	OR
	LEVONORGESTREL							
2913H	Tablets 30 micrograms, 28	4	2	..	*15.31	16.28	Microlut 28 Microval 28	SC WY
	MEDROXYPROGESTERONE ACETATE							
3118D	Injection 150 mg in 1 mL	1	1	..	13.03	14.00	^a Depo-Ralovera	KR
				^B 3.36	16.39	14.00	^a Depo-Provera	PH
<i>[For other listings for this drug see Generic/Proprietary Index]</i>								
	NORETHISTERONE							
1967M	Tablets 350 micrograms, 28	4	2	..	*15.31	16.28	^a Locilan 28 Day Micronor	KR JC
				^B 4.00	*19.31	16.28	^a Noriday 28 Day	PH

Androgens**• 3-oxoandrostens (4) derivatives****TESTOSTERONE****Authority required**

**Androgen deficiency in males with established pituitary or testicular disorders;
Androgen deficiency in males 40 years and older who do not have established pituitary or
testicular disorders other than aging, confirmed by at least 2 morning blood samples taken
on different mornings. Androgen deficiency is confirmed by testosterone less than 8 nmol
per L, or 8-15 nmol per L with high LH (greater than 1.5 times the upper limit of the
eugonadal reference range for young men);
Micropenis, pubertal induction, or constitutional delay of growth or puberty, in males
under 18 years of age.**

8098F	Subcutaneous implant 100 mg	6	..	..	*200.59	28.60	OR	
8099G	Subcutaneous implant 200 mg	3	..	..	*200.56	28.60	OR	

continued ➡

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
TESTOSTERONE—cont.								
8830R	Transdermal gel 50 mg in 5 g sachet, 30	‡1	5	..	96.41	28.60	Testogel	SC
8460G	Transdermal patches 12.2 mg (releasing approximately 2.5 mg per 24 hours), 60	‡1	5	..	97.15	28.60	Androderm	MX
8619P	Transdermal patches 24.3 mg (releasing approximately 5 mg per 24 hours), 30	‡1	5	..	97.15	28.60	Androderm	MX
TESTOSTERONE ENANTHATE								
Authority required								
Androgen deficiency in males with established pituitary or testicular disorders; Androgen deficiency in males 40 years and older who do not have established pituitary or testicular disorders other than aging, confirmed by at least 2 morning blood samples taken on different mornings. Androgen deficiency is confirmed by testosterone less than 8 nmol per L, or 8-15 nmol per L with high LH (greater than 1.5 times the upper limit of the eugonadal reference range for young men); Micropenis, pubertal induction, or constitutional delay of growth or puberty, in males under 18 years of age.								
2114G	Injection 250 mg in 1 mL	3	3	..	31.49	28.60	Primoteston Depot	SC
TESTOSTERONE ESTERS								
Authority required								
Androgen deficiency in males with established pituitary or testicular disorders; Androgen deficiency in males 40 years and older who do not have established pituitary or testicular disorders other than aging, confirmed by at least 2 morning blood samples taken on different mornings. Androgen deficiency is confirmed by testosterone less than 8 nmol per L, or 8-15 nmol per L with high LH (greater than 1.5 times the upper limit of the eugonadal reference range for young men); Micropenis, pubertal induction, or constitutional delay of growth or puberty, in males under 18 years of age.								
2670M	Injection 100 mg	3	3	..	*18.85	19.82	Sustanon 100	OR
2101N	Injection 250 mg	3	3	..	*31.48	28.60	Sustanon 250	OR
TESTOSTERONE UNDECANOATE								
Authority required								
Androgen deficiency in males with established pituitary or testicular disorders; Androgen deficiency in males 40 years and older who do not have established pituitary or testicular disorders other than aging, confirmed by at least 2 morning blood samples taken on different mornings. Androgen deficiency is confirmed by testosterone less than 8 nmol per L, or 8-15 nmol per L with high LH (greater than 1.5 times the upper limit of the eugonadal reference range for young men); Micropenis, pubertal induction, or constitutional delay of growth or puberty, in males under 18 years of age.								
2115H	Capsule 40 mg	60	5	..	35.51	28.60	Andriol Testocaps	OR

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Estrogens								
• Natural and semisynthetic estrogens, plain								
OESTRADIOL								
8274L	Tablet 2 mg	56	2	..	12.38	13.35	Zumenon	SM
1742Q	Vaginal tablets 25 micrograms, 15	‡1	2	..	19.41	20.38	Vagifem	NO
OESTRADIOL								
Restricted benefit								
<i>For use for post-menopausal symptoms where a trial of peri- or post-menopausal low-dose oestrogen therapy has demonstrated intolerance to oral oestrogens.</i>								
NOTE:								
<i>Oestradiol should be used in conjunction with an oral progestogen in women with an intact uterus.</i>								
8286D	Transdermal gel 1 mg in 1 g sachet, 28	‡1	5	..	16.19	17.16	Sandrena	OR
8761D	Transdermal patches 390 micrograms (releasing approximately 25 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Estradot 25	NV
8311K	Transdermal patches 750 micrograms (releasing approximately 25 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Estraderm MX 25	NV
8485N	Transdermal patches 2 mg (releasing approximately 25 micrograms per 24 hours), 4	‡1	5	..	16.19	17.16	^a Climara 25 ^a Femtran 25	SC MM
8194G	Transdermal patches 2 mg (releasing approximately 25 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Dermestril 25	MX
1743R	Transdermal patches 2 mg (releasing approximately 25 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Estraderm 25	NV
8762E	Transdermal patches 585 micrograms (releasing approximately 37.5 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Estradot 37.5	NV
8012Q	Transdermal patches 3.28 mg (releasing approximately 37.5 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Menorest 37.5	NV
8763F	Transdermal patches 780 micrograms (releasing approximately 50 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Estradot 50	NV

continued ➞

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
OESTRADIOL—cont.								
8140K	Transdermal patches 1.5 mg (releasing approximately 50 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Estraderm MX 50	NV
8125P	Transdermal patches 3.8 mg (releasing approximately 50 micrograms per 24 hours), 4	‡1	5	..	16.19	17.16	^a Climara 50 ^a Femtran 50	SC MM
8082J	Transdermal patches 4 mg (releasing approximately 50 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Dermestril 50	MX
1744T	Transdermal patches 4 mg (releasing approximately 50 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Estraderm 50	NV
8013R	Transdermal patches 4.33 mg (releasing approximately 50 micrograms per 24 hours), 8	‡1	5	..	16.19	17.16	Menorest 50	NV
8764G	Transdermal patches 1.17 mg (releasing approximately 75 micrograms per 24 hours), 8	‡1	5	..	18.37	19.34	Estradot 75	NV
8486P	Transdermal patches 5.7 mg (releasing approximately 75 micrograms per 24 hours), 4	‡1	5	..	18.37	19.34	Climara 75	SC
8014T	Transdermal patches 6.57 mg (releasing approximately 75 micrograms per 24 hours), 8	‡1	5	..	18.37	19.34	Menorest 75	NV
8765H	Transdermal patches 1.56 mg (releasing approximately 100 micrograms per 24 hours), 8	‡1	5	..	18.37	19.34	Estradot 100	NV
8312L	Transdermal patches 3 mg (releasing approximately 100 micrograms per 24 hours), 8	‡1	5	..	18.37	19.34	Estraderm MX 100	NV
8126Q	Transdermal patches 7.6 mg (releasing approximately 100 micrograms per 24 hours), 4	‡1	5	..	18.37	19.34	^a Climara 100 ^a Femtran 100	SC MM
8195H	Transdermal patches 8 mg (releasing approximately 100 micrograms per 24 hours), 8	‡1	5	..	18.37	19.34	Dermestril 100	MX

continued ↗

GENITO URINARY SYSTEM AND SEX HORMONES—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
OESTRADIOL—cont.								
1745W	Transdermal patches 8 mg (releasing approximately 100 micrograms per 24 hours), 8	‡1	5	..	18.37	19.34	Estraderm 100	NV
8041F	Transdermal patches 8.66 mg (releasing approximately 100 micrograms per 24 hours), 8	‡1	5	..	18.37	19.34	Menorest 100	NV
OESTRADIOL HEMIHYDRATE								
Restricted benefit								
<i>For use for post-menopausal symptoms where a trial of peri- or post-menopausal low-dose oestrogen therapy has demonstrated intolerance to oral oestrogens.</i>								
NOTE:								
<i>Oestradiol should be used in conjunction with an oral progestogen in women with an intact uterus.</i>								
8645B	Nasal spray 150 micrograms per actuation, 60 actuations, 4.2 mL	‡1	5	..	19.55	20.52	Aerodiol	SE
OESTRADIOL VALERATE								
1663M	Tablet 1 mg	56	2	..	10.37	11.34	Progynova	SC
1664N	Tablet 2 mg	56	2	..	12.38	13.35	Progynova	SC
1061W	Injection 10 mg in 1 mL	3	..	..	62.48	28.60	Primogyn Depot	SC
OESTRIOL								
1776L	Tablet 1 mg	60	2	..	*10.77	11.74	Ovestin	OR
1771F	Pessaries 500 micrograms, 15	‡1	2	..	19.41	20.38	Ovestin Ovula	OR
1781R	Vaginal cream 1 mg per g (0.1%), 15 g	‡1	1	..	16.14	17.11	Ovestin	OR
OESTROGENS—CONJUGATED								
1733F	Tablet 300 micrograms	56	2	..	*10.37	11.34	Premarin	WY
1734G	Tablet 625 micrograms	56	2	..	*12.39	13.36	Premarin	WY
PIPERAZINE OESTRONE SULFATE								
1777M	Tablet 730 micrograms (equivalent to 625 micrograms sodium oestrone sulfate)	56	2	.. B1.48	10.37 11.85	11.34 11.34	^a Genoral 0.625 ^a Ogen .625	KR PH
1778N	Tablet 1.46 mg (equivalent to 1.25 mg sodium oestrone sulfate)	56	2	.. B1.49	12.38 13.87	13.35 13.35	^a Genoral 1.25 ^a Ogen 1.25	KR PH
Progestogens								
• Pregnen (4) derivatives								
MEDROXYPROGESTERONE ACETATE								
2323G	Tablet 5 mg	56	2	.. B1.77	13.62 15.39	14.59 14.59	^a Ralovera ^a Provera	KR PH

continued ➡

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MEDROXYPROGESTERONE ACETATE—cont.								
2321E	Tablet 10 mg	30	2	..	14.25	15.22	^a Medroxyhexal	HX
				^B 1.78	16.03	15.22	^a Ralovera	KR
							^a Provera	PH
2319C	Injection 50 mg in 1 mL	1	1	..	10.28	11.25	Depo-Provera	PH
MEDROXYPROGESTERONE ACETATE								
<u>Restricted benefit</u>								
<i>Endometriosis.</i>								
2722G	Tablet 10 mg	100	2	..	30.77	28.60	^a Ralovera	KR
				^B 1.65	32.42	28.60	^a Provera	PH
[For other listings for this drug see Generic/Proprietary Index]								
• Pregnadien derivatives								
DYDROGESTERONE								
1350C	Tablet 10 mg	28	2	..	13.62	14.59	Duphaston	SM
• Estren derivatives								
NORETHISTERONE								
2993M	Tablet 5 mg	30	2	..	30.82	28.60	Primolut N	SC
Progestogens and estrogens in combination								
• Progestogens and estrogens, combinations								
OESTRADIOL with NORETHISTERONE ACETATE								
8353P	Tablets 1 mg-500 micrograms, 28	‡1	5	..	16.11	17.08	Kliovance	NO
8081H	Tablets 2 mg-1 mg, 28	‡1	5	..	16.11	17.08	Kliogest	NO
OESTRADIOL with NORETHISTERONE ACETATE								
<u>Restricted benefit</u>								
<i>For use for post-menopausal symptoms where a trial of peri- or post-menopausal low-dose oestrogen therapy has demonstrated intolerance to oral oestrogens.</i>								
8427M	Transdermal patches 620 micrograms-2.7 mg (releasing approximately 50 micrograms- 140 micrograms per 24 hours), 8	‡1	5	..	18.37	19.34	Estalis continuous 50/140	NV
8428N	Transdermal patches 510 micrograms-4.8 mg (releasing approximately 50 micrograms- 250 micrograms per 24 hours), 8	‡1	5	..	18.37	19.34	Estalis continuous 50/250	NV
OESTROGENS—CONJUGATED with MEDROXYPROGESTERONE ACETATE								
8168X	Tablets 625 micrograms-2.5 mg, 28	‡1	5	..	16.11	17.08	Premia 2.5 Continuous	WY
8169Y	Tablets 625 micrograms-5 mg, 28	‡1	5	..	16.11	17.08	Premia 5 Continuous	WY

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Progestogens and estrogens, sequential preparations								
OESTRADIOL and OESTRADIOL with DYDROGESTERONE								
8244X	Pack containing 14 tablets oestradiol 2 mg and 14 tablets oestradiol 2 mg with dydrogesterone 10 mg	‡1	5	..	16.11	17.08	Femoston 2/10	SM
OESTRADIOL and OESTRADIOL with NORETHISTERONE ACETATE								
1764W	Pack containing 12 tablets oestradiol 2 mg, 10 tablets oestradiol 2 mg with norethisterone acetate 1 mg and 6 tablets oestradiol 1 mg	‡1	5	..	16.11	17.08	Trisequens	NO
OESTRADIOL and OESTRADIOL with NORETHISTERONE ACETATE								
Restricted benefit								
<i>For use for post-menopausal symptoms where a trial of peri- or post-menopausal low-dose oestrogen therapy has demonstrated intolerance to oral oestrogens.</i>								
8425K	Pack containing 4 transdermal patches oestradiol 4.33 mg (releasing approximately 50 micrograms per 24 hours) and 4 transdermal patches oestradiol with norethisterone acetate 620 micrograms-2.7 mg (releasing approximately 50 micrograms- 140 micrograms per 24 hours)	‡1	5	..	18.37	19.34	Estalis sequi 50/140	NV
8426L	Pack containing 4 transdermal patches oestradiol 4.33 mg (releasing approximately 50 micrograms per 24 hours) and 4 transdermal patches oestradiol with norethisterone acetate 510 micrograms-4.8 mg (releasing approximately 50 micrograms- 250 micrograms per 24 hours)	‡1	5	..	18.37	19.34	Estalis sequi 50/250	NV
8029N	Pack containing 4 transdermal patches oestradiol 4 mg (releasing approximately 50 micrograms per 24 hours) and 4 transdermal patches oestradiol with norethisterone acetate 10 mg- 30 mg (releasing approximately 50 micrograms- 250 micrograms per 24 hours)	‡1	5	..	18.37	19.34	Estracombi	NV

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
8062H	OESTRADIOL VALERATE and OESTRADIOL VALERATE with CYPROTERONE ACETATE Pack containing 11 tablets oestradiol valerate 2 mg and 10 tablets oestradiol valerate 2 mg with cyproterone acetate 1 mg	1	5	..	16.11	17.08	Climen	SC
Gonadotropins and other ovulation stimulants								
• Gonadotropins								
FOLLITROPIN ALFA								
<u>Restricted benefit</u>								
<i>Anovulatory infertility.</i>								
<u>NOTE:</u>								
<i>Except in cases of hypopituitarism or primary amenorrhoea, the patient should have been adequately treated with clomiphene citrate and/or gonadorelin and failed to have conceived.</i>								
<i>Women who have had apparent ovulation induced by other agents and have failed to conceive should have laparoscopic evidence that there is no other impediment to conception.</i>								
<i>Oligomenorrhoea should have been present for at least twelve months or amenorrhoea for at least six months prior to treatment.</i>								
<i>Patients with hyperprolactinaemia should have had appropriate surgical or medical treatment prior to treatment.</i>								
8672K	Injection set containing 1 vial powder for injection 75 i.u. and 1 pre-filled syringe solvent 1 mL	5	5	..	*253.35	28.60	Gonal-f 75	SG
8673L	Injection set containing 10 vials powder for injection 75 i.u. and 10 pre-filled syringes solvent 1 mL	1	5	..	484.37	28.60	Gonal-f 75	SG
8713N	Injection 300 i.u. in 0.5 mL multi-dose cartridge	3	5	..	*580.36	28.60	Gonal-f Pen	SG
8675N	Injection set containing 1 vial powder for injection 450 i.u. and 1 pre-filled syringe solvent 1 mL	3	5	..	*868.12	28.60	Gonal-f	SG
8714P	Injection 450 i.u. in 0.75 mL multi-dose cartridge	3	5	..	*868.15	28.60	Gonal-f Pen	SG
8715Q	Injection 900 i.u. in 1.5 mL multi-dose cartridge	2	5	..	*1155.97	28.60	Gonal-f Pen	SG
8674M	Injection set containing 1 vial powder for injection 1,050 i.u. and 1 pre-filled syringe solvent 2 mL	1	5	..	676.22	28.60	Gonal-f	SG

continued ➡

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
FOLLITROPIN ALFA—cont.								
<u>Restricted benefit</u>								
<i>For the treatment of infertility in males due to hypogonadotrophic hypogonadism, following failure of 6 months' treatment with human chorionic gonadotrophin to achieve adequate spermatogenesis. Combined treatment with HCG must be given.</i>								
8673L	Injection set containing 10 vials powder for injection 75 i.u. and 10 pre-filled syringes solvent 1 mL	1	5	..	484.37	28.60	Gonal-f 75	SG
8713N	Injection 300 i.u. in 0.5 mL multi-dose cartridge	3	5	..	*580.36	28.60	Gonal-f Pen	SG
8675N	Injection set containing 1 vial powder for injection 450 i.u. and 1 pre-filled syringe solvent 1 mL	3	5	..	*868.12	28.60	Gonal-f	SG
8714P	Injection 450 i.u. in 0.75 mL multi-dose cartridge	3	5	..	*868.15	28.60	Gonal-f Pen	SG
8715Q	Injection 900 i.u. in 1.5 mL multi-dose cartridge	2	5	..	*1155.97	28.60	Gonal-f Pen	SG
FOLLITROPIN BETA								
<u>Restricted benefit</u>								
<i>Anovulatory infertility.</i>								
<u>NOTE:</u>								
<i>Except in cases of hypopituitarism or primary amenorrhoea, the patient should have been adequately treated with clomiphene citrate and/or gonadorelin and failed to have conceived.</i>								
<i>Women who have had apparent ovulation induced by other agents and have failed to conceive should have laparoscopic evidence that there is no other impediment to conception.</i>								
<i>Oligomenorrhoea should have been present for at least twelve months or amenorrhoea for at least six months prior to treatment.</i>								
<i>Patients with hyperprolactinaemia should have had appropriate surgical or medical treatment prior to treatment.</i>								
8437C	Solution for injection 100 i.u. in 0.5 mL	10	5	..	643.41	28.60	Puregon 100 IU/0.5 mL	OR
8438D	Solution for injection 150 i.u. in 0.5 mL	10	5	..	962.75	28.60	Puregon 150 IU/0.5 mL	OR
8565T	Solution for injection 300 i.u. in 0.36 mL multi-dose cartridge	3	5	..	*579.55	28.60	Puregon 300 IU/0.36 mL	OR
8566W	Solution for injection 600 i.u. in 0.72 mL multi-dose cartridge	2	5	..	*771.15	28.60	Puregon 600 IU/0.72 mL	OR
8871X	Solution for injection 900 i.u. in 1.08 mL multi-dose cartridge	2	5	..	*1154.35	28.60	Puregon 900 IU/1.08 mL	OR

continued ➡

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
FOLLITROPIN BETA—cont.								
<u>Restricted benefit</u>								
<i>For the treatment of infertility in males due to hypogonadotrophic hypogonadism, following failure of 6 months' treatment with human chorionic gonadotrophin to achieve adequate spermatogenesis. Combined treatment with HCG must be given.</i>								
8437C	Solution for injection 100 i.u. in 0.5 mL	10	5	..	643.41	28.60	Puregon 100 IU/0.5 mL	OR
8438D	Solution for injection 150 i.u. in 0.5 mL	10	5	..	962.75	28.60	Puregon 150 IU/0.5 mL	OR
8835B	Solution for injection 200 i.u. in 0.5 mL	10	5	..	643.41	28.60	Puregon 200 IU/0.5 mL	OR
8565T	Solution for injection 300 i.u. in 0.36 mL multi-dose cartridge	3	5	..	*579.55	28.60	Puregon 300 IU/0.36 mL	OR
8566W	Solution for injection 600 i.u. in 0.72 mL multi-dose cartridge	2	5	..	*771.15	28.60	Puregon 600 IU/0.72 mL	OR
8871X	Solution for injection 900 i.u. in 1.08 mL multi-dose cartridge	2	5	..	*1154.35	28.60	Puregon 900 IU/1.08 mL	OR

HUMAN CHORIONIC GONADOTROPHIN**Restricted benefit***Anovulatory infertility.***NOTE:***Except in cases of hypopituitarism or primary amenorrhoea, the patient should have been adequately treated with clomiphene citrate and/or gonadorelin and failed to have conceived.**Women who have had apparent ovulation induced by other agents and have failed to conceive should have laparoscopic evidence that there is no other impediment to conception.**Oligomenorrhoea should have been present for at least twelve months or amenorrhoea for at least six months prior to treatment.**Patients with hyperprolactinaemia should have had appropriate surgical or medical treatment prior to treatment.***Restricted benefit***For the treatment of infertility in males due to hypogonadotrophic hypogonadism;**For the treatment of infertility in males associated with isolated luteinising hormone deficiency;**For the treatment of males who have combined deficiency of human growth hormone and gonadotrophins and in whom the absence of secondary sexual characteristics indicates a lag in maturation.***Restricted benefit***For the treatment of boys over the age of 16 years who show clinical evidence of hypogonadism or delayed puberty. Treatment must not extend beyond 6 months.*

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
HUMAN CHORIONIC GONADOTROPHIN—cont.								
1579D	Injection set containing 3 ampoules powder for injection 500 units and 3 ampoules solvent 1 mL	1	5	..	28.92	28.60	Pregnyl	OR
1581F	Injection set containing 3 ampoules powder for injection 1,500 units and 3 ampoules solvent 1 mL	1	5	..	38.48	28.60	Pregnyl	OR
Restricted benefit <i>Cryptorchism not due to organic obstruction in boys over 12 months of age.</i>								
1583H	Injection set containing 3 ampoules powder for injection 500 units and 3 ampoules solvent 1 mL	2	1	..	*53.09	28.60	Pregnyl	OR
• Ovulation stimulants, synthetic CLOMIPHENE CITRATE NOTE: <i>Care must be taken to comply with the provisions of State/Territory law when prescribing clomiphene citrate.</i>								
Restricted benefit <i>Anovulatory infertility; Patients undergoing in-vitro fertilisation.</i>								
1211R	Tablet 50 mg	10	5	..	*39.91	28.60	^a Clomhexal	HX
							^a Fermil	AW
							^a GenRx	GX
							Clomiphene	
				^B 0.18	40.09	28.60	^a Serophene	SG
				^B 3.74	43.65	28.60	^a Clomid	AV
Antiandrogens								
• Antiandrogens, plain preparations CYPROTERONE ACETATE Authority required <i>Moderate to severe androgenisation in non-pregnant women (acne alone is not a sufficient indication of androgenisation).</i> CAUTION: <i>This drug should not be used during pregnancy as it may result in feminisation of the male foetus.</i>								
1269T	Tablet 50 mg	20	5	..	68.13	28.60	^a Cyprohexal	HX
							^a Cyprone	AF
							^a Cyprostat	SY
							^a GenRx	GX
							Cyproterone Acetate	
				^B 1.50	69.63	28.60	^a Procur	DP
							^a Androcur	SC

continued ➡

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
CYPROTERONE ACETATE—cont.							
Authority required							
<i>Advanced carcinoma of the prostate; To reduce drive in sexual deviations in males.</i>							
1270W	Tablet 50 mg	100	5	..	*275.31	28.60	^a Cyprohexal HX ^a Cyprone AF ^a Cyprostat SY ^a GenRx GX Cyproterone Acetate ^a Procur DP ^a Androcur SC
					^B 3.28 *278.59	28.60	
8019C	Tablet 100 mg	50	5	..	227.34	28.60	^a Cyprohexal HX ^a Cyprostat-100 SY ^a GenRx GX Cyproterone Acetate ^a Procur 100 DP ^B 1.00 228.34 28.60 ^a Androcur-100 SC

Other sex hormones and modulators of the genital system

- Antigonadotropins and similar agents

DANAZOL**CAUTION:**

Pregnancy must be excluded prior to administration of this drug.

Authority required

Endometriosis, visually proven;

Hereditary angio-oedema;

Short-term treatment (up to 6 months) of intractable primary menorrhagia (not more than 6 months' therapy will be authorised);

Short-term treatment (up to 6 months) of severe benign (fibrocystic) breast disease or mastalgia associated with severe symptomatic benign breast disease in patients refractory to other treatments (not more than 6 months' therapy will be authorised).

1285P	Capsule 100 mg	100	5	..	58.65	28.60	Azol 100 AF
1287R	Capsule 200 mg	100	5	..	88.00	28.60	Azol 200 AF

GESTRINONE**Authority required**

Short-term treatment (up to 6 months) of visually proven endometriosis (only 1 course of not more than 6 months' therapy will be authorised).

8015W	Capsule 2.5 mg	8	5	..	78.67	28.60	Dimetriose AV

- Selective estrogen receptor modulators

RALOXIFENE HYDROCHLORIDE

For listings see Generic/Proprietary Index

GENITO URINARY SYSTEM AND SEX HORMONES —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
UROLOGICALS								
Other urologicals, incl. antispasmodics								
• Urinary antispasmodics								
OXYBUTYNIN HYDROCHLORIDE								
<u>Restricted benefit</u>								
<i>Detrusor overactivity where propantheline bromide has failed.</i>								
8039D	Tablet 5 mg	100	5	..	15.75	16.72	Ditropan	AV
PROPANTHELINE BROMIDE								
<u>Restricted benefit</u>								
<i>Detrusor overactivity.</i>								
1953T	Tablet 15 mg	200	5	..	*24.55	25.52	Pro-Banthine	SI
• Other urologicals								
PHENOXYBENZAMINE HYDROCHLORIDE								
<u>Restricted benefit</u>								
<i>Phaeochromocytoma; Neurogenic urinary retention.</i>								
1862B	Capsule 10 mg	100	5	..	52.27	28.60	Dibenylene	LM

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES								
Anterior pituitary lobe hormones and analogues								
• ACTH								
TETRACOSACTRIN								
2832C	Injection 1 mg in 1 mL	5	5	..	*71.80	28.60	Synacthen Depot 1 mg/1 mL	NV
Posterior pituitary lobe hormones								
• Vasopressin and analogues								
DESMOPRESSIN ACETATE								
Authority required								
<i>Cranial diabetes insipidus.</i>								
8662X	Tablet 200 micrograms	90	5	..	*154.54	28.60	Minirin	FP
2129C	Intranasal solution 100 micrograms per mL, 2.5 mL	5	5	..	*164.70	28.60	Minirin	FP
8711L	Nasal spray (pump pack) 10 micrograms per actuation, 60 actuations, 6 mL	2	5	..	*164.51	28.60	Minirin Nasal Spray	FP
Authority required								
<i>Primary nocturnal enuresis in patients aged 6 years or older:</i>								
<i>(1) who are refractory to an enuresis alarm; or</i>								
<i>(2) for whom an enuresis alarm is contraindicated. The reason that an alarm is</i>								
<i>contraindicated must be included in the authority application.</i>								
NOTE:								
<i>Not to be used in preference to enuresis alarms.</i>								
8663Y	Tablet 200 micrograms	30	5	..	54.68	28.60	Minirin	FP
NOTE:								
<i>Only one application per six months with no more than twice the maximum quantity will be</i>								
<i>authorised for the tablets.</i>								
8712M	Nasal spray (pump pack) 10 micrograms per actuation, 60 actuations, 6 mL	1	5	..	84.63	28.60	Minirin Nasal Spray	FP
Hypothalamic hormones								
• Gonadotropin-releasing hormones								
NAFARELIN ACETATE								
Authority required								
<i>Initial treatment (up to 6 months) of visually proven endometriosis;</i>								
<i>Subsequent treatment (up to 6 months) of visually proven endometriosis, where 2 years or</i>								
<i>more have elapsed since the end of the previous course and where a recent bone density</i>								
<i>assessment has been made. The date of the assessment must be provided.</i>								
2962X	Nasal spray (pump pack) 200 micrograms (base) per dose (60 doses)	1	5	..	96.82	28.60	Synarel	PH

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CORTICOSTEROIDS FOR SYSTEMIC USE								
Corticosteroids for systemic use, plain								
• Mineralocorticoids								
1433K	FLUDROCORTISONE ACETATE Tablet 100 micrograms	200	1	..	*16.83	17.80	Florinef	BQ
• Glucocorticoids								
BETAMETHASONE ACETATE with BETAMETHASONE SODIUM PHOSPHATE								
<u>Restricted benefit</u>								
<i>Alopecia areata; For local intra-articular or peri-articular infiltration; Granulomata, dermal; Keloid; Lichen planus hypertrophic; Lichen simplex chronicus; Lupus erythematosus, chronic discoid; Necrobiosis lipoidica; Uveitis.</i>								
2694T	Injection 3 mg-3.9 mg (equivalent to 5.7 mg betamethasone) in 1 mL	5	..	..	24.66	25.63	Celestone Chronodose	SH
CORTISONE ACETATE								
1246N	Tablet 5 mg	50	4	..	12.59	13.56	Cortate	AS
1247P	Tablet 25 mg	60	4	..	14.27	15.24	Cortate	AS
DEXAMETHASONE								
1292B	Tablet 500 micrograms	30	4	..	6.98	7.95	Dexmethsone	AS
2507Y	Tablet 4 mg	30	4	..	10.44	11.41	Dexmethsone	AS
DEXAMETHASONE SODIUM PHOSPHATE								
2509C	Injection equivalent to 4 mg dexamethasone phosphate in 1 mL	5	..	..	13.85	14.82	MX	
1291Y	Injection equivalent to 8 mg dexamethasone phosphate in 2 mL	5	1	..	21.14	22.11	MX	
2508B	Injection equivalent to 120 mg dexamethasone phosphate in 5 mL	1	..	..	61.24	28.60	MX	
HYDROCORTISONE								
1499X	Tablet 4 mg	50	4	..	9.56	10.53	Hysone 4	AF
1500Y	Tablet 20 mg	60	4	..	13.23	14.20	Hysone 20	AF

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
HYDROCORTISONE SODIUM SUCCINATE								
1501B	Injection equivalent to 100 mg hydrocortisone with 2 mL solvent	2	..	..	*15.55	16.52	Solu-Cortef	PH
3096Y	Injection equivalent to 250 mg hydrocortisone with 2 mL solvent	1	..	..	12.27	13.24	Solu-Cortef	PH
HYDROCORTISONE SODIUM SUCCINATE								
<u>Restricted benefit</u>								
<i>For use in a hospital.</i>								
1510L	Injection equivalent to 100 mg hydrocortisone with 2 mL solvent	6	..	..	*37.15	28.60	Solu-Cortef	PH
1511M	Injection equivalent to 250 mg hydrocortisone with 2 mL solvent	6	..	..	*49.87	28.60	Solu-Cortef	PH
METHYLPREDNISOLONE ACETATE								
<u>Restricted benefit</u>								
<i>For local intra-articular or peri-articular infiltration.</i>								
1928L	Injection 40 mg in 1 mL	5	..	..	23.31	24.28	^a Depo-Nisolone	KR
				^B0.75	24.06	24.28	^a Depo-Medrol	PH
METHYLPREDNISOLONE SODIUM SUCCINATE								
2981X	Powder for injection equivalent to 40 mg methylprednisolone with diluent	5	..	..	32.14	28.60	Solu-Medrol	PF
8834Y	Powder for injection equivalent to 1 g methylprednisolone with diluent	1	..	..	102.47	28.60	Solu-Medrol	PF
PREDNISOLONE								
3152X	Tablet 1 mg	100	4	..	7.12	8.09	^a Predsolone	LN
				^B 0.62	7.74	8.09	^a Panafcortelone	AS
1917X	Tablet 5 mg	60	4	..	7.60	8.57	Panafcortelone Solone	AS FM
1916W	Tablet 25 mg	30	4	..	9.90	10.87	Panafcortelone Solone	AS FM
PREDNISOLONE SODIUM PHOSPHATE								
8285C	Oral solution equivalent to 5 mg prednisolone per mL, 30 mL	‡1	5	..	12.78	13.75	^a PredMix	LN
				^B 1.53	14.31	13.75	^a Redipred	AS

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
1934T	PREDNISONE Tablet 1 mg	100	4	.. B0.62	7.12 7.74	8.09 8.09	^a Predsone ^a Panafcort	LN AS
1935W	Tablet 5 mg	60	4	..	7.60	8.57	Panafcort Sone	AS FM
1936X	Tablet 25 mg	30	4	..	9.90	10.87	Panafcort Sone	AS FM
TRIAMCINOLONE ACETONIDE								
<u>Restricted benefit</u>								
<i>Alopecia areata; For local intra-articular or peri-articular infiltration; Granulomata, dermal; Keloid; Lichen planus hypertrophic; Lichen simplex chronicus; Lupus erythematosus, chronic discoid; Necrobiosis lipoidica; Psoriasis.</i>								
2990J	Injection 10 mg in 1 mL	5	..	..	24.66	25.63	Kenacort-A10	BQ
Antiadrenal preparations								
• Anticorticosteroids								
AMINOGLUTETHIMIDE								
1036M	Tablet 250 mg	100	5	..	163.04	28.60	Cytadren 250	NV
THYROID THERAPY								
Thyroid preparations								
• Thyroid hormones								
LIOETHYRONINE SODIUM								
<u>Authority required</u>								
<i>Management of patients with thyroid cancer; Replacement therapy for hypothyroid patients who have documented intolerance to thyroxine sodium; Replacement therapy for hypothyroid patients who have documented resistance to thyroxine sodium; Initiation of thyroid therapy in severely hypothyroid patients.</i>								
2318B	Tablet 20 micrograms	100	2	..	73.50	28.60	Tertroxin	SI
THYROXINE SODIUM								
2174K	Tablet equivalent to 50 micrograms anhydrous thyroxine sodium	200	1	.. B1.30	13.37 14.67	14.34 14.34	^a Eutroxsig ^a Oroxine	FM SI
2175L	Tablet equivalent to 100 micrograms anhydrous thyroxine sodium	200	1	.. B1.30	14.51 15.81	15.48 15.48	^a Eutroxsig ^a Oroxine	FM SI
2173J	Tablet equivalent to 200 micrograms anhydrous thyroxine sodium	200	1	.. B1.30	16.73 18.03	17.70 17.70	^a Eutroxsig ^a Oroxine	FM SI

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Antithyroid preparations								
• Thiouracils								
1955X	PROPYLTHIOURACIL Tablet 50 mg	200	2	..	*31.73	28.60	PL	
• Sulfur-containing imidazole derivatives								
1153Q	CARBIMAZOLE Tablet 5 mg	200	2	..	*29.07	28.60	Neo-Mercazole	RO
PANCREATIC HORMONES								
Glycogenolytic hormones								
• Glycogenolytic hormones								
1449G	GLUCAGON HYDROCHLORIDE Injection set containing 1 mg (1 i.u.) and 1 mL solvent in disposable syringe	1	1	..	40.12	28.60	GlucaGen Hypokit	NO
CALCIUM HOMEOSTASIS								
Anti-parathyroid hormones								
• Calcitonin preparations								
SALCATONIN								
NOTE:								
<i>The maximum quantities for salcatonin shown represent the number of individual ampoules and NOT multiples of the manufacturer's packs. The pack size for both strengths is five ampoules.</i>								
Authority required								
<i>Symptomatic Paget's disease of bone;</i>								
<i>Treatment initiated in a hospital (in-patient or out-patient) of hypercalcaemia.</i>								
2995P	Injection 50 i.u. in 1 mL	30	5	..	*212.11	28.60	Miacalcic 50	NV
2997R	Injection 100 i.u. in 1 mL	15	5	..	*164.59	28.60	Miacalcic 100	NV

ANTIINFECTIVES FOR SYSTEMIC USE

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ANTIBACTERIALS FOR SYSTEMIC USE							
Tetracyclines							
• Tetracyclines							
DOXYCYCLINE							
2709N	Tablet 100 mg	7	1	..	7.43	8.40	^a Chem mart CH Doxycycline ^a Doxsig SI ^a Doxy-100 DP ^a Doxyhexal HX ^a Doxylin 100 AF ^a GenRx GX Doxycycline ^a Terry White TW Chemists Doxycycline ^a Vibramycin PF
2708M	Capsule 100 mg	7	1	..	7.43	8.40	^a DBL Doxycycline FA ^a Doryx MX
				^B 1.60	9.03	8.40	
				^B 1.52	8.95	8.40	
DOXYCYCLINE							
Restricted benefit							
<i>Bronchiectasis in patients aged 8 years or older; Chronic bronchitis in patients aged 8 years or older; Severe acne.</i>							
2711Q	Tablet 50 mg	25	5	..	9.56	10.53	^a Chem mart CH Doxycycline ^a Doxy-50 DP ^a Doxyhexal HX ^a Doxylin 50 AF ^a Frakas AW ^a GenRx GX Doxycycline ^a Terry White TW Chemists Doxycycline ^a Vibra-Tabs PF
2707L	Capsule 50 mg	25	5	..	9.56	10.53	^a DBL FA Doxycycline ^a Doryx MX
				^B 1.65	11.21	10.53	
				^B 1.71	11.27	10.53	

continued ↗

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
<u>DOXYCYCLINE—cont.</u>								
<u>Restricted benefit</u>								
<i>Pelvic inflammatory disease.</i>								
2702F	Tablet 100 mg	28	..	..	*15.47	16.44	^a Chem mart Doxycycline	CH
							^a Doxsig	SI
							^a Doxy-100	DP
							^a Doxyhexal	HX
							^a Doxylin 100	AF
							^a GenRx	GX
							^a Doxycycline	
							^a Terry White Chemists	TW
							^a Doxycycline	
				^B 6.40	*21.87	16.44	^a Vibramycin	PF
2703G	Capsule 100 mg	28	..	..	*15.47	16.44	^a DBL	FA
							^a Doxycycline	
				^B 6.08	*21.55	16.44	^a Doryx	MX
<u>Restricted benefit</u>								
<i>Urethritis.</i>								
2714W	Tablet 100 mg	21	..	..	*12.79	13.76	^a Chem mart Doxycycline	CH
							^a Doxsig	SI
							^a Doxy-100	DP
							^a Doxyhexal	HX
							^a Doxylin 100	AF
							^a Terry White Chemists	TW
							^a Doxycycline	
				..	12.79	13.76	^a GenRx	GX
							^a Doxycycline	
				^B 4.80	*17.59	13.76	^a Vibramycin	PF
2715X	Capsule 100 mg	21	..	..	12.79	13.76	^a DBL	FA
							^a Doxycycline	
				^B 2.75	15.54	13.76	^a Doryx	MX

MINOCYCLINE

CAUTION:

There are concerns about the incidence of benign intracranial hypertension associated with this drug.

3037W Capsule 100 mg 11 9.02 9.99 Akamin 100 AF

NOTE:

No applications for increased maximum quantities and/or repeats will be authorised.

continued ↗

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MINOCYCLINE—cont.								
CAUTION:								
<i>There are concerns about the incidence of benign intracranial hypertension associated with this drug.</i>								
Restricted benefit								
<i>Severe acne not responding to other tetracyclines.</i>								
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
1616C	Tablet 50 mg	60	5	..	16.71	17.68	^a Akamin 50	AF
				^B 1.12	17.83	17.68	^a Minomycin-50	SI
Beta-lactam antibacterials, penicillins								
• Penicillins with extended spectrum								
AMOXYCILLIN								
1883D	Chewable tablet 250 mg	20	1	..	8.96	9.93	Amoxil	GK
1884E	Capsule 250 mg	20	1	..	7.95	8.92	^a Alphamox 250	AF
							^a Amohexal	HX
							^a Amoxycillin-DP	DG
							^a Bgramin	DP
							^a Chem mart	CH
							Amoxycillin	
							^a Cilamox	SI
							^a GenRx	GX
							Amoxycillin	
							^a Moxacin	CS
							^a Terry White	TW
							Chemists	
							Amoxycillin	
				^B 1.00	8.95	8.92	^a Amoxil	GK
1889K	Capsule 500 mg	20	1	..	11.13	12.10	^a Alphamox 500	AF
							^a Amohexal	HX
							^a Amoxycillin-DP	DG
							^a Bgramin	DP
							^a Chem mart	CH
							Amoxycillin	
							^a Cilamox	SI
							^a GenRx	GX
							Amoxycillin	
							^a Moxacin	CS
							^a Terry White	TW
							Chemists	
							Amoxycillin	
				^B 1.00	12.13	12.10	^a Amoxil	GK
1878W	Sachet containing oral powder 3 g	1	..	..	8.81	9.78	Amoxil	GK
1888J	Powder for paediatric oral drops 100 mg per mL, 20 mL	‡1	1	..	#11.96	13.34	Amoxil	GK

continued ➡

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
AMOXYCILLIN—cont.								
1886G	Powder for syrup 125 mg per 5 mL, 100 mL	‡1	1	..	#10.02	11.40	^a Alphamox 125 ^a Amohexal ^a Bgramin ^a Chem mart Amoxycillin ^a Cilamox ^a GenRx Amoxycillin ^a Moxacin ^a Terry White Chemists Amoxycillin	AF HX DP CH SI GX CS TW
				^B 1.00	#11.02	11.40	^a Amoxil	GK
1887H	Powder for syrup 250 mg per 5 mL, 100 mL	‡1	1	..	#11.28	12.66	^a Alphamox 250 ^a Amohexal ^a Bgramin ^a Chem mart Amoxycillin ^a Cilamox ^a GenRx Amoxycillin ^a Moxacin 250 ^a Terry White Chemists Amoxycillin	AF HX DP CH SI GX CS TW
				^B 1.01	#12.29	12.66	^a Amoxil Forte	GK
8705E	Powder for oral suspension 500 mg per 5 mL, 100 mL	‡1	1	..	#14.49	15.87	Maxamox	SZ
AMOXYCILLIN								
<u>Restricted benefit</u>								
<i>Acute exacerbations of chronic bronchitis.</i>								
8581P	Tablet 1 g	14	1	..	11.33	12.30	^a Amoxycillin Sandoz	BG
				^B 1.49	12.82	12.30	^a Amoxil Duo ^a Maxamox	GK SZ
AMPICILLIN								
1048E	Capsule 250 mg	24	1	..	8.59	9.56	Alphacin 250	AF
2671N	Capsule 500 mg	24	..	..	11.96	12.93	Alphacin 500	AF
2390T	Powder for injection 500 mg	5	1	..	10.89	11.86	^a Austrapen ^a Ibimicyn	CS DP
2977Q	Powder for injection 1 g	5	1	..	14.82	15.79	^a Aspen Ampicyn ^a Austrapen ^a Ibimicyn	AS CS DP

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Beta-lactamase sensitive penicillins								
BENZATHINE PENICILLIN								
8167W	Injection 900 mg in 2 mL cartridge-needle unit (for use with Tubex Injector)	1	..	..	24.95	25.92	Bicillin L-A Tubex	AS
<u>BENZATHINE PENICILLIN</u>								
<u>Restricted benefit</u>								
<u>Syphilis.</u>								
8743E	Injection 900 mg in 2 mL cartridge-needle unit (for use with Tubex Injector)	2	..	..	*45.15	28.60	Bicillin L-A Tubex	AS
BENZYL PENICILLIN								
1775K	Powder for injection 600 mg	10	1	..	*22.25	23.22	BenPen	CS
2647H	Powder for injection 3 g	10	..	..	*50.51	28.60	BenPen	CS
PHENOXYMETHYLPENICILLIN								
1787C	Tablet 250 mg	50	..	..	*11.53	12.50	Abbocillin-VK Filmtab	SI
3028J	Tablet 500 mg	50	..	..	*14.83	15.80	Abbocillin-VK Filmtab	SI
1789E	Capsule 250 mg	50	..	..	11.33	12.30	^a Cilicaine VK ^a Cilopen VK LPV ^a Penhexal VK	FM DP CS HX
2965C	Capsule 500 mg	50	..	..	14.53	15.50	^a Cilicaine VK ^a Cilopen VK LPV ^a Penhexal VK	FM DP CS HX
2356B	Paediatric oral suspension 125 mg per 5 mL, 100 mL	2	..	.. ^B 1.44	*12.35 *13.79	13.32 13.32	^a Cilicaine V ^a Abbocillin-V	FM SI
2354X	Oral suspension 250 mg per 5 mL, 100 mL	2	..	.. ^B 1.46	*15.11 *16.57	16.08 16.08	^a Cilicaine V ^a Abbocillin-V	FM SI
<u>PHENOXYMETHYLPENICILLIN</u>								
<u>Restricted benefit</u>								
<u>Prophylaxis of recurrent streptococcal infections (including rheumatic fever).</u>								
1703P	Tablet 250 mg	50	5	..	*11.53	12.50	Abbocillin-VK Filmtab	SI
1705R	Capsule 250 mg	50	5	..	11.33	12.30	^a Cilicaine VK ^a Cilopen VK LPV ^a Penhexal VK	FM DP CS HX
PROCAINE PENICILLIN								
1794K	Injection 1.5 g	5	..	..	52.49	28.60	Cilicaine	SI

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Beta-lactamase resistant penicillins								
DICLOXACILLIN								
8123M	Powder for injection 500 mg	5	..	..	16.83	17.80	Diclocil	BQ
8124N	Powder for injection 1 g	5	1	..	23.69	24.66	Diclocil	BQ
DICLOXACILLIN								
<u>Restricted benefit</u>								
<i>Serious staphylococcal infections.</i>								
8121K	Capsule 250 mg	24	..	..	11.37	12.34	^a Diclocil	BQ
							^a Dicloxsig	SI
							^a Distaph 250	AF
8122L	Capsule 500 mg	24	..	..	18.62	19.59	^a Diclocil	BQ
							^a Dicloxsig	SI
							^a Distaph 500	AF
FLUCLOXACILLIN								
<u>CAUTION:</u>								
Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.								
1524F	Powder for injection 500 mg	5	..	..	16.71	17.68	^a Flopen	CS
							^a Flubiclox	DP
1525G	Powder for injection 1 g	5	1	..	23.51	24.48	^a Aspen Flucil	AS
							^a Flopen	CS
							^a Flubiclox	DP
							^a MX	
FLUCLOXACILLIN								
<u>CAUTION:</u>								
Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.								
<u>Restricted benefit</u>								
<i>Serious staphylococcal infections.</i>								
1526H	Capsule 250 mg	24	..	..	11.37	12.34	^a Flophen	CS
							^a Floxsig	SI
							^a Staphylex 250	AF
				^B 0.46	11.83	12.34	^a Floxapen	GK
1527J	Capsule 500 mg	24	..	..	18.62	19.59	^a Flophen	CS
							^a Floxsig	SI
							^a Staphylex 500	AF
				^B 0.58	19.20	19.59	^a Floxapen	GK
1528K	Powder for syrup 125 mg per 5 mL, 100 mL	‡1	..	..	#13.13	14.51	^a Flophen	CS
					^B 0.06	#13.19	^a Floxapen	GK
1529L	Powder for syrup 250 mg per 5 mL, 100 mL	‡1	..	..	#16.86	18.24	^a Flophen	CS
				^B 0.10	#16.96	18.24	^a Floxapen	GK

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
• Combinations of penicillins, incl. beta-lactamase inhibitors AMOXYCILLIN with CLAVULANIC ACID CAUTION: <i>Hepatotoxicity has been reported with this drug.</i> Restricted benefit <i>Infections where resistance to amoxycillin is suspected;</i> <i>Infections where resistance to amoxycillin is proven.</i>							
1891M	Tablet 500 mg-125 mg	10	1	..	13.39	14.36	^a Clamohexal Duo 500mg/125mg HX ^a Clamoxyl Duo AL ^a Clavulin Duo AW ^a Curam 500/125 SZ ^a Muric 500/125 SL ^a Augmentin Duo GK
					^B 0.99 14.38	14.36	
8254K	Tablet 875 mg-125 mg	10	1	..	17.06	18.03	^a Chem mart Amoxycillin and Clavulanic Acid CH ^a Clamohexal Duo Forte 875mg/125mg HX ^a Clamoxyl Duo forte AL ^a Clavulin Duo Forte AW ^a Curam 875/125 SZ ^a GenRx GX Amoxycillin and Clavulanic Acid ^a Muric 875/125 SL ^a Terry White Chemists Amoxycillin and Clavulanic Acid TW ^B 1.30 18.36 18.03 ^a Augmentin Duo forte GK
1892N	Powder for syrup 125 mg-31.25 mg per 5 mL, 75 mL	‡1	1	..	#12.47	13.85	^a Clamohexal 125mg/31.25mg/5mL HX ^a Clamoxyl AL ^a Clavulin AW ^B 0.96 #13.43 13.85 ^a Augmentin GK

continued ➞

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
AMOXYCILLIN with CLAVULANIC ACID—cont.								
8319W	Powder for syrup 400 mg- 57 mg per 5 mL, 60 mL	#1	1	..	#14.75	16.13	^a Clamohexal Duo 400mg/ 57mg/5mL	HX
							^a Clamoxyl Duo 400	AL
							^a Clavulin Duo 400	AW
				^B 0.98	#15.73	16.13	^a Augmentin Duo 400	GK
TICARCILLIN with CLAVULANIC ACID								
Restricted benefit								
<i>Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent; Septicaemia, suspected; Septicaemia, proven.</i>								
2179Q	Powder for injection 3 g- 100 mg	10	..	..	152.15	28.60	Timentin	GK
Other beta-lactam antibacterials								
• First-generation cephalosporins								
CEPHALEXIN								
3058Y	Capsule 250 mg	20	1	..	8.40	9.37	^a Chem mart Cephalexin	CH
							^a Cilex	DP
							^a GenRx Cephalexin	GX
							^a Ialex	LN
							^a Ibilex 250	AF
							^a Sporaheal	HX
							^a Terry White Chemists Cephalexin	TW
				^B 1.76	10.16	9.37	^a Keflex	AS
3119E	Capsule 500 mg	20	1	..	11.31	12.28	^a Chem mart Cephalexin	CH
							^a Cilex	DP
							^a GenRx Cephalexin	GX
							^a Ialex	LN
							^a Ibilex 500	AF
							^a Sporaheal	HX
							^a Terry White Chemists Cephalexin	TW
				^B 1.87	13.18	12.28	^a Keflex	AS

continued ↻

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CEPHALEXIN—cont.								
3094W	Granules for syrup 125 mg per 5 mL, 100 mL	‡1	1	..	#11.52	12.90	^a Chem mart Cephalexin	CH
							^a Cilex	DP
							^a GenRx	GX
							Cephalexin	
							^a Ialex	LN
							^a Ibilex 125	AF
							^a Terry White Chemists	TW
							Cephalexin	
				^B 1.78	#13.30	12.90	^a Keflex	AS
3095X	Granules for syrup 250 mg per 5 mL, 100 mL	‡1	1	..	#13.63	15.01	^a Chem mart Cephalexin	CH
							^a Cilex	DP
							^a GenRx	GX
							Cephalexin	
							^a Ialex	LN
							^a Ibilex 250	AF
							^a Terry White Chemists	TW
							Cephalexin	
				^B 1.87	#15.50	15.01	^a Keflex	AS
CEPHALOTHIN								
2964B	Powder for injection 1 g	10	1	..	43.57	28.60	^a MX	
				^B 0.30	43.87	28.60	^a Keflin Neutral	LY
CEPHAZOLIN								
<u>Restricted benefit</u>								
<i>Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent;</i>								
<i>Septicaemia, suspected;</i>								
<i>Septicaemia, proven.</i>								
1256D	Powder for injection 500 mg	10	..	..	*44.91	28.60	MX	
1257E	Powder for injection 1 g	10	..	..	*68.13	28.60	^a Cefazolin Sandoz	SZ
							^a MX	
				^B 2.51	70.64	28.60	^a Kefzol	LY

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Second-generation cephalosporins								
CEFACLOR								
CAUTION:								
Serum sickness-like reactions have been reported with this drug, especially in children.								
1169M	Tablet 375 mg (sustained release)	10	1	..	14.51	15.48	^a Cefkor CD ^a Chem mart Cefaclor CD ^a Douglas Cefaclor-CD ^a GenRx Cefaclor CD ^a Karlor CD ^a Keflor CD ^a Terry White Chemists Cefaclor CD	DP CH DG GX LN AF TW
				^B 1.54	16.05	15.48	^a Ceclor CD	AS
2460L	Powder for oral suspension 125 mg per 5 mL, 100 mL	‡1	1	..	#14.02	15.40	^a Aclor 125 ^a Chem mart Cefaclor ^a GenRx Cefaclor ^a Keflor ^a Terry White Chemists Cefaclor	AW CH GX AF TW
				^B 1.53	#15.55	15.40	^a Ceclor	AS
2461M	Powder for oral suspension 250 mg per 5 mL, 75 mL	‡1	1	..	#14.52	15.90	^a Aclor 250 ^a Chem mart Cefaclor ^a GenRx Cefaclor ^a Keflor ^a Terry White Chemists Cefaclor	AW CH GX AF TW
				^B 1.55	#16.07	15.90	^a Ceclor	AS
CEFUROXIME AXETIL								
8292K	Tablet 250 mg (base)	14	1	..	14.79	15.76	Zinnat	GK
• Third-generation cephalosporins								
CEFOTAXIME								
Restricted benefit								
<i>Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent; Septicaemia, suspected; Septicaemia, proven.</i>								
1085D	Powder for injection 1 g	10	..	..	*56.35	28.60	^a Cefotaxime Sandoz	SZ
				..	56.35	28.60	^a MX	
1086E	Powder for injection 2 g	10	..	..	*100.15	28.60	^a Cefotaxime Sandoz	SZ
				..	100.15	28.60	^a MX	

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CEFTRIAXONE								
<u>Restricted benefit</u>								
<i>Gonorrhoea.</i>								
1782T	Powder for injection 250 mg	1	..	..	11.57	12.54	Rocephin	RO
<u>Restricted benefit</u>								
<i>Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent;</i>								
<i>Septicaemia, suspected;</i>								
<i>Septicaemia, proven.</i>								
1790F	Powder for injection 250 mg	5	..	..	*38.85	28.60	Rocephin	RO
1783W	Powder for injection 500 mg	5	..	..	*54.10	28.60	Rocephin	RO
1784X	Powder for injection 1 g	5	..	..	*82.20	28.60	^a Ceftriaxone Sandoz	SZ
				..	82.20	28.60	^a Rocephin	RO
							^a DBL	MX
							Ceftriaxone	
1785Y	Powder for injection 2 g	5	..	..	*148.10	28.60	^a Ceftriaxone Sandoz	SZ
							^a DBL	MX
							Ceftriaxone	
							^a Rocephin	RO
• Fourth-generation cephalosporins								
CEFEPIME								
<u>Authority required</u>								
<i>Treatment of febrile neutropenia.</i>								
8315P	Powder for injection 1 g	10	..	..	*188.15	28.60	Maxipime	BQ
8316Q	Powder for injection 2 g	10	..	..	*340.15	28.60	Maxipime	BQ
Sulfonamides and trimethoprim								
• Trimethoprim and derivatives								
TRIMETHOPRIM								
2922T	Tablet 300 mg	7	1	..	7.47	8.44	^a Alprim	AF
				^B 1.36	8.83	8.44	^a Triprim	SI
• Combinations of sulfonamides and trimethoprim, incl. derivatives								
TRIMETHOPRIM with SULFAMETHOXAZOLE								
<u>CAUTION:</u>								
There is an increased risk of severe adverse reactions with this combination in the elderly.								
2949F	Tablet 80 mg-400 mg	10	1	..	7.72	8.69	^a Resprim	AF
				^B 1.54	9.26	8.69	^a Seprim	SI

continued ➡

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
TRIMETHOPRIM with SULFAMETHOXAZOLE—cont.								
2951H	Tablet 160 mg-800 mg	10	1	..	8.66	9.63	^a Bactrim DS ^a Chem mart Trimethoprim with Sulfame- thoxazole DS ^a GenRx Trimethoprim with Sulfame- thoxazole DS ^a Resprim Forte ^a Terry White Chemists Trimethoprim with Sulfame- thoxazole DS	RO CH GX AF TW SI
				^B 1.46	10.12	9.63	^a Septrin Forte	SI
3103H	Oral suspension 40 mg-200 mg per 5 mL, 100 mL	‡1	1	..	8.23	9.20	Bactrim ^a Resprim	RO AF
				^B 1.91	10.14	9.20	^a Septrin	SI

Macrolides, lincosamides and streptogramins• **Macrolides****AZITHROMYCIN****Restricted benefit***Uncomplicated urethritis due to Chlamydia trachomatis;**Uncomplicated cervicitis due to Chlamydia trachomatis.***NOTE:***No applications for increased maximum quantities and/or repeats will be authorised.*

8200N	Tablet 500 mg	2	..	..	22.02	22.99	Zithromax	PF

Restricted benefit*Trachoma.***NOTE:***No applications for increased maximum quantities and/or repeats will be authorised.*

8336R	Tablet 500 mg	2	2	..	22.02	22.99	Zithromax	PF
8201P	Powder for oral suspension 200 mg per 5 mL, 15 mL	‡1	..	..	#21.46	22.84	Zithromax	PF

CLARITHROMYCIN								
8318T	Tablet 250 mg	14	1	..	14.55	15.52	^a Chem mart Clarithromycin ^a Clarac ^a Clarihexal ^a GenRx Clarithromycin ^a Kalixocin ^a Terry White Chemists Clarithromycin	CH DP HX GX AF TW AB
				^B 2.00	16.55	15.52	^a Klacid	AB

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
1404X	ERYTHROMYCIN Capsule 250 mg	25	1	..	8.72	9.69	^a DBL Erythromycin	FA
				^B 1.77	10.49	9.69	^a Eryc	MX
2750R	ERYTHROMYCIN ETHYL SUCCINATE Tablet 400 mg (base)	25	1	..	8.39	9.36	^a E-Mycin	AF
				^B 2.89	11.28	9.36	^a E.E.S. 400 Filmtab	AB
2424N	Powder for oral liquid 200 mg (base) per 5 mL, 100 mL	‡1	1	..	#10.45	11.83	^a E-Mycin 200	AF
				^B 2.19	#12.64	11.83	^a E.E.S. 200	AB
2428T	Powder for oral liquid 400 mg (base) per 5 mL, 100 mL	‡1	1	..	#12.08	13.46	^a E-Mycin 400	AF
				^B 1.75	#13.83	13.46	^a E.E.S. Granules	AB
1397M	ERYTHROMYCIN LACTOBIONATE Powder for I.V. infusion 1 g (base)	5	..	..	*50.75	28.60	Erythrocin-I.V.	AB
8129W	ROXITHROMYCIN Tablet for oral suspension 50 mg	10	1	..	10.22	11.19	Rulide D	AV
1760P	Tablet 150 mg	10	1	..	11.78	12.75	^a Biaxsig	HP
				^B 2.50	14.28	12.75	^a Rulide	AV
8016X	Tablet 300 mg	5	1	..	11.78	12.75	^a Biaxsig	HP
				^B 2.50	14.28	12.75	^a Rulide	AV
• Lincosamides								
CLINDAMYCIN								
Restricted benefit								
<i>Gram-positive coccal infections where these cannot be safely and effectively treated with a penicillin.</i>								
3138E	Capsule 150 mg	25	..	..	18.47	19.44	^a Cleocin	KR
				^B 1.47	19.94	19.44	^a Dalacin C	PH
2530E	LINCOMYCIN Injection 600 mg in 2 mL	5	..	..	26.96	27.93	Lincocin	PH
Aminoglycoside antibacterials								
• Other aminoglycosides								
2824P	GENTAMICIN SULFATE Injection 80 mg (base) in 2 mL	10	1	..	*18.91	19.88	^a MX	
				..	18.92	19.89	^a PU	
TOBRAMYCIN SULFATE								
Restricted benefit								
<i>Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent; Septicaemia, suspected; Septicaemia, proven.</i>								
1356J	Injection 80 mg (base) in 2 mL (with preservative)	10	1	..	*65.25	28.60	Nebcin MX	AS
8872Y	Injection 80 mg (base) in 2 mL (without preservative)	10	1	..	*65.25	28.60	PU	

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
Quinolone antibacterials							
• Fluoroquinolones							
CIPROFLOXACIN							
Restricted benefit							
Gonorrhoea.							
NOTE:							
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>							
1311B	Tablet 250 mg	2	..	..	12.23	13.20	Ciproxin 250 BN
Authority required							
<i>Respiratory tract infection proven or suspected to be caused by Pseudomonas aeruginosa in severely immunocompromised patients;</i>							
<i>Bacterial gastroenteritis in severely immunocompromised patients;</i>							
<i>Treatment of infections proven to be due to Pseudomonas aeruginosa or other gram-negative bacteria resistant to all other oral antimicrobials;</i>							
<i>Treatment of joint and bone infections, epididymo-orchitis, prostatitis or perichondritis of the pinna, suspected or proven to be caused by gram-negative bacteria or gram-positive bacteria resistant to all other appropriate antimicrobials.</i>							
1208N	Tablet 250 mg	14	..	..	34.75	28.60	^a C-Flox 250 AL ^a Ciprofloxacin-BC BG ^a Ciprol 250 AW ^a GenRx GX Ciprofloxacin ^a Profloxin HX ^a Ciproxin 250 BN
				^B 1.90	36.65	28.60	
1209P	Tablet 500 mg	14	..	..	63.35	28.60	^a C-Flox 500 AL ^a Ciprofloxacin-BC BG ^a Ciprol 500 AW ^a GenRx GX Ciprofloxacin ^a Profloxin HX ^a Proquin DP ^a Ciproxin 500 BN
				^B 1.90	65.25	28.60	
1210Q	Tablet 750 mg	14	..	..	93.05	28.60	^a C-Flox 750 AL ^a Ciprol 750 AW ^a GenRx GX Ciprofloxacin ^a Profloxin HX ^a Proquin DP ^a Ciproxin 750 BN
				^B 1.90	94.95	28.60	

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer		
MOXIFLOXACIN HYDROCHLORIDE									
Authority required									
<i>Initial treatment of radiologically-confirmed, severe community-acquired pneumonia, requiring admission to an intensive care or high dependency unit, in patients greater than 12 years old with a history of hypersensitivity to penicillin.</i>									
8635L	Solution for I.V. infusion 400 mg (base) in 250 mL	3	..	..	*199.15	28.60	Avelox	BN	
Authority required									
<i>Oral treatment, following initial intravenous treatment with moxifloxacin hydrochloride, of radiologically-confirmed, severe community-acquired pneumonia, requiring admission to an intensive care or high dependency unit, in patients greater than 12 years old with a history of hypersensitivity to penicillin;</i>									
<i>Radiologically-confirmed, severe community-acquired pneumonia requiring hospitalisation, in patients greater than 12 years old with a history of hypersensitivity to penicillin;</i>									
<i>Radiologically-confirmed, community-acquired pneumonia in patients greater than 12 years old with a history of immediate hypersensitivity to penicillin (as defined by urticaria, angioedema, bronchospasm or anaphylaxis within 1 hour of drug administration).</i>									
8636M	Tablet 400 mg (base)	5	1	..	46.62	28.60	Avelox	BN	
NORFLOXACIN									
Authority required									
<i>Acute bacterial enterocolitis;</i>									
<i>Complicated urinary tract infection.</i>									
3010K	Tablet 400 mg	14	1	..	19.64	20.61	^a Chem mart Norfloxacin	CH	
							^a GenRx Norfloxacin	GX	
							^a Norflohexal	HX	
							^a Nufloxib	AF	
							^a Roxin	AW	
							^a Terry White Chemists Norfloxacin	TW	
							^a DP		
					^B 1.73	21.37	20.61	^a Insensye	FR
					^B 3.73	23.37	20.61	^a Noroxin	MK
Other antibacterials									
• Glycopeptide antibacterials									
VANCOMYCIN									
Restricted benefit									
<i>Prophylaxis of endocarditis in patients hypersensitive to penicillin.</i>									
3130R	Powder for injection 500 mg (500,000 i.u.) vancomycin activity	2	..	..	*51.91	28.60	^a MX		
					^B 0.36	*52.27	28.60	^a Vancocin	LY

continued ➡

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
VANCOMYCIN—cont.								
Restricted benefit								
Endophthalmitis;								
Use initiated in a hospital for infections where vancomycin is an appropriate antibiotic.								
3131T	Powder for injection 500 mg (500,000 i.u.) vancomycin activity	5	..	..	*122.65 B0.90 *123.55	28.60 28.60	a MX a Vancocin	LY
• Steroid antibacterials								
FUSIDIC ACID								
Restricted benefit								
For use in combination with another antibiotic in the treatment of proven serious staphylococcal infections.								
2312Q	Tablet (sodium salt) 250 mg	36	1	..	75.49	28.60	Fucidin	CS
• Imidazole derivatives								
METRONIDAZOLE								
1636D	Tablet 200 mg	21	1	..	6.77	7.74	a Metrogyl 200	AF
					B1.98	8.75	a Metronide 200	HP
							a Flagyl	AV
1626N	Tablet 400 mg	5	2	..	6.66	7.63	Metrogyl 400	AF
1642K	Suppositories 500 mg, 10	‡1	..	..	20.26	21.23	Flagyl	AV
METRONIDAZOLE								
Restricted benefit								
Treatment of anaerobic infections.								
1621H	Tablet 400 mg	21	1	..	9.50	10.47	a Metrogyl 400	AF
					B2.07	11.57	a Metronide 400	HP
							a Flagyl	AV
Restricted benefit								
Prophylaxis in large bowel surgery;								
Treatment, in a hospital, of acute anaerobic sepsis.								
1638F	I.V. infusion 500 mg in 100 mL	5	1	..	*43.20	28.60	BX	
METRONIDAZOLE BENZOATE								
1630T	Oral suspension 320 mg per 5 mL (equivalent to 200 mg metronidazole in 5 mL), 100 mL	‡1	..	..	14.39	15.36	Flagyl S	AV
TINIDAZOLE								
1465D	Tablet 500 mg	4	..	..	7.57	8.54	a Simplotan	GP
					B2.50	10.07	a Fasigyn	PF
• Nitrofurantoin derivatives								
NITROFURANTOIN								
CAUTION:								
Nitrofurantoin may cause peripheral neuritis and severe pulmonary reactions.								
1692C	Capsule 50 mg	30	1	..	14.65	15.62	Macrochantin	PU
1693D	Capsule 100 mg	30	1	..	22.31	23.28	Macrochantin	PU

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Other antibacterials								
3124K	HEXAMINE HIPPURATE Tablet 1 g	100	5	..	32.50	28.60	Hiprex	MM
ANTIMYCOTICS FOR SYSTEMIC USE								
Antimycotics for systemic use								
• Antibiotics								
1047D	AMPHOTERICIN Powder for injection 50 mg	1	..	..	27.67	28.60	Fungizone	BQ
• Imidazole derivatives								
KETOCONAZOLE								
<u>Authority required</u>								
<i>Symptomatic genital candidiasis recurring after treatment of at least 2 episodes with topical therapy.</i>								
<u>CAUTION:</u>								
<i>Hepatotoxicity has been reported with ketoconazole.</i>								
1573T	Tablet 200 mg	10	..	..	17.35	18.32	Nizoral	JC
<u>Authority required</u>								
<i>Oral candidiasis in severely immunocompromised persons where topical therapy has failed;</i>								
<i>Systemic or deep mycoses where other forms of therapy have failed.</i>								
<u>CAUTION:</u>								
<i>Hepatotoxicity has been reported with ketoconazole.</i>								
1572R	Tablet 200 mg	30	5	..	38.63	28.60	Nizoral	JC
• Triazole derivatives								
FLUCONAZOLE								
<u>Authority required</u>								
<i>Treatment of cryptococcal meningitis in patients unable to take or tolerate amphotericin;</i>								
<i>Maintenance therapy in patients with cryptococcal meningitis and immunosuppression;</i>								
<i>Treatment of oropharyngeal candidiasis in immunosuppressed patients;</i>								
<i>Treatment of oesophageal candidiasis in immunosuppressed patients;</i>								
<i>Secondary prophylaxis of oropharyngeal candidiasis in immunosuppressed patients;</i>								
<i>Treatment of serious and life-threatening candida infections in patients unable to tolerate amphotericin.</i>								
1471K	Capsule 50 mg	28	5	..	177.51	28.60	Diflucan	PF
1472L	Capsule 100 mg	28	5	..	328.71	28.60	Diflucan	PF
1475P	Capsule 200 mg	28	5	..	624.59	28.60	Diflucan	PF
1473M	Solution for I.V. infusion 100 mg in 50 mL	7	..	..	*196.55	28.60	Diflucan	PF
1474N	Solution for I.V. infusion 200 mg in 100 mL	7	..	..	*350.90	28.60	Diflucan	PF

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ITRACONAZOLE								
Authority required								
<i>Systemic aspergillosis;</i>								
<i>Systemic sporotrichosis;</i>								
<i>Systemic histoplasmosis;</i>								
<i>Treatment and maintenance therapy in patients with AIDS who have disseminated pulmonary histoplasmosis infection;</i>								
<i>Treatment and maintenance therapy in patients with AIDS who have chronic pulmonary histoplasmosis infection;</i>								
<i>Treatment of oropharyngeal candidiasis in immunosuppressed patients;</i>								
<i>Treatment of oesophageal candidiasis in immunosuppressed patients.</i>								
8196J	Capsule 100 mg	60	5	..	210.57	28.60	Sporanox	JC
ANTIMYCOBACTERIALS								
Drugs for treatment of tuberculosis								
• Hydrazides								
ISONIAZID								
1554T	Tablet 100 mg	100	2	..	9.43	10.40	FM	
Drugs for treatment of lepra								
• Drugs for treatment of lepra								
DAPSONE								
8801F	Tablet 25 mg	100	1	..	102.06	28.60	LM	
1272Y	Tablet 100 mg	100	1	..	115.75	28.60	LM	
RIFAMPICIN								
Restricted benefit								
<i>Prophylaxis of meningococcal disease in close contacts and carriers;</i>								
<i>Prophylactic treatment of contacts of patients with Haemophilus influenzae type B.</i>								
1981G	Capsule 150 mg	10	..	..	8.79	9.76	Rimycin 150	AF
1984K	Capsule 300 mg	10	..	..	11.80	12.77	Rimycin 300	AF
8025J	Syrup 100 mg per 5 mL, 60 mL	‡1	..	..	23.58	24.55	Rifadin	AV
Authority required								
<i>Leprosy in adults.</i>								
1982H	Capsule 150 mg	100	..	..	34.99	28.60	Rimycin 150	AF
1983J	Capsule 300 mg	100	..	..	65.23	28.60	Rimycin 300	AF

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ANTIVIRALS FOR SYSTEMIC USE							
Direct acting antivirals							
• Nucleosides and nucleotides excl. reverse transcriptase inhibitors							
ACICLOVIR							
Authority required							
<i>Moderate to severe initial genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is desirable but need not delay treatment.</i>							
NOTE:							
<i>Aciclovir 200 mg is not PBS-subsidised for chickenpox, herpes zoster or herpes simplex infections other than genital herpes.</i>							
1003T	Tablet 200 mg	50	..	..	*104.13	28.60	^a Acihexal HX
							^a Acyclo-V 200 AF
							^a Lovir DP
				..	104.14	28.60	^a GenRx Aciclovir GX
				^B 5.78	*109.91	28.60	^a Zovirax 200 mg GK
NOTE:							
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>							
Authority required							
➤ <i>Episodic treatment or suppressive therapy of moderate to severe recurrent genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.</i>							
NOTE:							
<i>Aciclovir 200 mg is not PBS-subsidised for chickenpox, herpes zoster or herpes simplex infections other than genital herpes.</i>							
1007B	Tablet 200 mg	90	5	..	186.59	28.60	^a Acihexal HX
							^a Acyclo-V 200 AF
							^a Chem mart CH
							Aciclovir
							^a GenRx Aciclovir GX
							^a Lovir DP
							^a Terry White TW
							Chemists
							Aciclovir
							^a Zyclir 200 AW
				^B 4.28	190.87	28.60	^a Zovirax 200 mg GK

continued ↗

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ACICLOVIR—cont.							
Authority required							
<i>Treatment of patients with herpes zoster within 72 hours of the onset of the rash; Herpes zoster ophthalmicus.</i>							
NOTE:							
<i>Aciclovir is effective only if commenced within 72 hours of onset of rash.</i>							
<i>Aciclovir 800 mg is not PBS-subsidised for herpes simplex or chickenpox.</i>							
1052J	Tablet 800 mg	35	..	..	223.02	28.60	^a Acihexal HX ^a Acyclo-V 800 AF ^a GenRx Aciclovir GX ^a Lovir DP ^a Zyclir 800 AW ^B 2.09 225.11 28.60 ^a Zovirax 800 mg GK
NOTE:							
<i>No applications for repeats will be authorised.</i>							
Authority required							
<i>Patients with advanced HIV disease (CD4 cell counts of less than 150 million per litre).</i>							
8234J	Tablet 800 mg	120	5	..	695.71	28.60	^a Acihexal HX ^a Acyclo-V 800 AF ^a Lovir DP ^B 7.19 702.90 28.60 ^a Zovirax 800 mg GK
FAMCICLOVIR							
Authority required							
➤	<i>Episodic treatment of moderate to severe recurrent genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.</i>						
NOTE:							
<i>Famciclovir 125 mg is not PBS-subsidised for chickenpox, herpes zoster or herpes simplex infections other than genital herpes.</i>							
8092X	Tablet 125 mg	40	1	..	159.03	28.60	Famvir NV
Authority required							
<i>Treatment of patients with herpes zoster within 72 hours of the onset of the rash.</i>							
NOTE:							
<i>Famciclovir is effective only if commenced within 72 hours of onset of rash.</i>							
<i>Famciclovir 250 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.</i>							
8002E	Tablet 250 mg	21	..	..	166.74	28.60	Famvir NV
NOTE:							
<i>No applications for repeats will be authorised.</i>							

continued ➤

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
	<u>FAMCICLOVIR—cont.</u>							
	<u>Authority required</u>							
➤	<i>Suppressive therapy of moderate to severe recurrent genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.</i>							
	<u>NOTE:</u>							
	<i>Famciclovir 250 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.</i>							
8217L	Tablet 250 mg	56	5	..	415.45	28.60	Famvir	NV
	<u>VALACICLOVIR HYDROCHLORIDE</u>							
	<u>Authority required</u>							
	<i>Moderate to severe initial genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is desirable but need not delay treatment.</i>							
	<u>NOTE:</u>							
	<i>Valaciclovir 500 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.</i>							
8133C	Tablet 500 mg (base)	20	..	..	*122.07	28.60	Valtrex	GK
	<u>NOTE:</u>							
	<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>							
	<u>Authority required</u>							
➤	<i>Episodic treatment or suppressive therapy of moderate to severe recurrent genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.</i>							
	<u>NOTE:</u>							
	<i>Valaciclovir 500 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.</i>							
8134D	Tablet 500 mg (base)	30	5	..	180.73	28.60	Valtrex	GK
	<u>Authority required</u>							
	<i>Treatment of patients with herpes zoster within 72 hours of the onset of the rash; Herpes zoster ophthalmicus.</i>							
	<u>NOTE:</u>							
	<i>Valaciclovir is effective only if commenced within 72 hours of onset of rash.</i>							
	<i>Valaciclovir 500 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.</i>							
8064K	Tablet 500 mg (base)	42	..	..	246.72	28.60	Valtrex	GK
	<u>NOTE:</u>							
	<i>No applications for repeats will be authorised.</i>							

ANTIINFECTIVES FOR SYSTEMIC USE —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
VACCINES								
Bacterial vaccines								
• Pneumococcal vaccines								
PNEUMOCOCCAL VACCINE, POLYVALENT								
Restricted benefit								
<i>Splenectomised persons over 2 years of age;</i>								
<i>Persons with Hodgkin's disease;</i>								
<i>Persons at high risk of pneumococcal infections.</i>								
1903E	Injection 0.5 mL (23 valent)	1	..	..	40.54	28.60	Pneumovax 23	CS
• Tetanus vaccines								
DIPHTHERIA and TETANUS VACCINE, ADSORBED								
NOTE:								
For immunisation of children up to the age of eight years.								
1341N	Injection 0.5 mL	3	..	..	*18.64	19.61	CS	
DIPHTHERIA and TETANUS VACCINE, ADSORBED, DILUTED FOR ADULT USE								
NOTE:								
For immunisation of adults and children over the age of eight years.								
8783G	Injection 0.5 mL in pre-filled syringe	3	..	..	*21.94	22.91	ADT	CS
TETANUS VACCINE, ADSORBED								
8813W	Injection 0.5 mL in pre-filled syringe	3	..	..	*16.99	17.96	Tet-Tox	CS
Viral vaccines								
• Influenza vaccines								
INFLUENZA VACCINE								
Restricted benefit								
<i>Persons at special risk of adverse consequences from infections of the lower respiratory tract.</i>								
2852D	Injection (trivalent) 0.5 mL (containing A/New Caledonia/20/99, A/Wellington/1/2004 and B/Shanghai/361/2002 like strains)	1	..	..	17.82	18.79	Fluarix Fluvax Influvac Vaxigrip	GK CS SM AX

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTINEOPLASTIC AGENTS								
Alkylating agents								
• Nitrogen mustard analogues								
CHLORAMBUCIL								
1163F	Tablet 2 mg	100	2	..	*82.43	28.60	Leukeran	GK
CYCLOPHOSPHAMIDE								
1266P	Tablet 50 mg	50	2	..	29.35	28.60	Cycloblastin	PH
1079T	Powder for injection 500 mg	2	..	..	*27.83	28.60	Endoxan	BX
1080W	Powder for injection 1 g	1	..	..	25.11	26.08	Endoxan	BX
1031G	Powder for injection 2 g	1	..	..	41.02	28.60	Endoxan	BX
IFOSFAMIDE								
Restricted benefit								
<i>Relapsed or refractory germ cell tumours following first-line chemotherapy;</i>								
<i>Relapsed or refractory sarcomas following first-line chemotherapy.</i>								
8076C	Powder for I.V. injection 1 g	5	5	..	*267.80	28.60	Holoxan	BX
8077D	Powder for I.V. injection 2 g	5	5	..	*491.25	28.60	Holoxan	BX
MELPHALAN								
2547C	Tablet 2 mg	25	1	..	27.45	28.42	Alkeran	GK
• Alkyl sulphonates								
BUSULFAN								
1128J	Tablet 2 mg	100	..	..	46.00	28.60	Myleran	GK
• Ethylene imines								
THIOTEPA								
2345K	Powder for injection 15 mg	2	1	..	*42.17	28.60	SI	
• Nitrosoureas								
FOTEMUSTINE								
Authority required								
<i>Metastatic malignant melanoma.</i>								
8786K	Powder for injection 208 mg with solvent	1	4	..	1034.35	28.60	Muphoran	SE
• Other alkylating agents								
TEMOZOLOMIDE								
Authority required								
<i>Glioblastoma multiforme concomitantly with radiotherapy.</i>								
NOTE:								
<i>Applications for doses above 150 mg per day will not be authorised. No applications for increased repeats will be authorised.</i>								
8819E	Capsule 5 mg	15	2	..	*222.49	28.60	Temodal	SH
8820F	Capsule 20 mg	15	2	..	*615.55	28.60	Temodal	SH
8821G	Capsule 100 mg	15	2	..	*2621.32	28.60	Temodal	SH

continued ➡

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
TEMOZOLOMIDE—cont.								
Authority required								
<i>Recurrence of anaplastic astrocytoma following standard therapy;</i>								
<i>Recurrence of glioblastoma multiforme following standard therapy;</i>								
<i>Glioblastoma multiforme following radiotherapy.</i>								
8378Y	Capsule 5 mg	5	5	..	77.99	28.60	Temodal	SH
8379B	Capsule 20 mg	5	5	..	218.52	28.60	Temodal	SH
8380C	Capsule 100 mg	5	5	..	876.94	28.60	Temodal	SH
8381D	Capsule 250 mg	5	5	..	2093.21	28.60	Temodal	SH
Antimetabolites								
• Folic acid analogues								
METHOTREXATE								
1622J	Tablet 2.5 mg	30	5	..	11.93	12.90	^a Methoblastin	PH
							^a MX	
				^B 0.08	12.01	12.90	^a Ledertrexate	WY
1623K	Tablet 10 mg	50	2	..	47.10	28.60	Methoblastin	PH
2396D	Injection 5 mg in 2 mL	5	..	..	36.67	28.60	MX	
2395C	Injection 50 mg in 2 mL	5	..	..	35.96	28.60	MX PU	
8850T	Solution concentrate for I.V. infusion 500 mg in 5 mL	1	..	..	67.16	28.60	Methotrexate Ebewe	IT
8863L	Solution concentrate for I.V. infusion 500 mg in 20 mL	1	..	..	67.16	28.60	MX	
8851W	Solution concentrate for I.V. infusion 1000 mg in 10 mL	1	..	..	129.59	28.60	^a Methotrexate Ebewe	IT
							^a MX	
8852X	Solution concentrate for I.V. infusion 5000 mg in 50 mL	1	..	..	594.89	28.60	Methotrexate Ebewe	IT
PEMETREXED DISODIUM								
<i>See Special Pharmaceutical Benefits</i>								
RALTITREXED								
Authority required								
<i>For use as a single agent in the treatment of advanced colorectal cancer.</i>								
8284B	Powder for I.V. infusion 2 mg	3	2	..	*883.00	28.60	Tomudex	AP
• Purine analogues								
CLADRIBINE								
Authority required								
<i>Hairy cell leukaemia.</i>								
8800E	Injection 10 mg in 5 mL	7	..	..	*5122.59	28.60	Litak	OA
1811H	Solution for I.V. infusion 10 mg in 10 mL	7	..	..	5122.74	28.60	Leustatin	JC
MERCAPTOPURINE								
1598D	Tablet 50 mg	100	2	..	*103.23	28.60	Purinethol	GK

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
1233X	THIOGUANINE Tablet 40 mg	25	1	..	104.08	28.60	Lanvis	GK
• Pyrimidine analogues CAPECITABINE Authority required <i>Advanced breast cancer after failure of prior therapy which includes a taxane and an anthracycline;</i> <i>Advanced breast cancer where therapy with a taxane and/or an anthracycline is contraindicated;</i> <i>Advanced breast cancer in combination with docetaxel after failure of prior anthracycline-containing chemotherapy;</i> <i>Treatment of advanced or metastatic colorectal cancer.</i>								
8361C	Tablet 150 mg	60	2	..	122.26	28.60	Xeloda	RO
8362D	Tablet 500 mg	120	2	..	693.50	28.60	Xeloda	RO
CYTARABINE								
2884T	Injection 100 mg in 5 mL	10	1	..	*59.83	28.60	PU	
8034W	Injection 500 mg in 5 mL	2	1	..	*68.31	28.60	MX	
FLUOROURACIL								
2528C	Injection 500 mg in 10 mL	10	..	..	*66.45	28.60	MX	
GEMCITABINE HYDROCHLORIDE Authority required <i>Advanced breast cancer in combination with paclitaxel after failure of prior therapy which includes an anthracycline;</i> <i>Advanced epithelial ovarian cancer, in combination with carboplatin, in patients who relapse more than 6 months after platinum-based therapy;</i> <i>Locally advanced or metastatic non-small cell lung cancer;</i> <i>Locally advanced or metastatic adenocarcinoma of the pancreas;</i> <i>Locally advanced or metastatic bladder cancer, in combination with cisplatin.</i>								
8049P	Powder for I.V. infusion 200 mg (base)	4	2	..	*241.31	28.60	Gemzar	LY
8050Q	Powder for I.V. infusion 1 g (base)	2	2	..	*566.01	28.60	Gemzar	LY
Plant alkaloids and other natural products								
• Vinca alkaloids and analogues								
VINBLASTINE SULFATE								
2198Q	Powder for injection 10 mg	5	..	..	*98.85	28.60	Velbe	AS
2199R	Solution for I.V. injection 10 mg in 10 mL	5	..	..	107.49	28.60	MX	
VINCRISTINE SULFATE								
2374Y	I.V. injection 1 mg in 1 mL	10	..	..	*168.11	28.60	MX PU	

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
VINORELBINE TARTRATE								
Authority required								
Advanced breast cancer after failure of prior therapy which includes an anthracycline; Locally advanced or metastatic non-small cell lung cancer.								
8280T	Solution for I.V. infusion 10 mg (base) in 1 mL	16	2	..	*1258.67	28.60	^a Navelbine ^a MX	FB
8281W	Solution for I.V. infusion 50 mg (base) in 5 mL	4	2	..	*1316.47	28.60	^a Navelbine ^a MX	FB
• Podophyllotoxin derivatives								
ETOPOSIDE								
1396L	Capsule 50 mg	20	..	..	494.74	28.60	Vepesid	BQ
1389D	Capsule 100 mg	10	..	..	433.15	28.60	Vepesid	BQ
1390E	Solution for I.V. infusion 100 mg in 5 mL	5	..	..	*180.70	28.60	^a Etoposide Ebewe ^a MX	IT
ETOPOSIDE PHOSPHATE								
8120J	Powder for I.V. infusion 113.6 mg (equivalent to 100 mg etoposide)	5	..	..	*180.70	28.60	Etopophos	BQ
8515E	Powder for I.V. infusion 1136 mg (equivalent to 1 g etoposide)	1	..	..	342.64	28.60	Etopophos	BQ
• Taxanes								
DOCETAXEL								
Authority required								
Advanced breast cancer after failure of prior therapy which includes an anthracycline; Advanced metastatic ovarian cancer after failure of prior therapy which includes a platinum compound; Locally advanced or metastatic non-small cell lung cancer.								
8071T	Injection set containing 1 single use vial concentrate for I.V. infusion 20 mg (anhydrous) in 0.5 mL and 1 single use vial solvent 1.5 mL	2	..	..	*681.29	28.60	Taxotere	AV
8074Y	Injection set containing 1 single use vial concentrate for I.V. infusion 80 mg (anhydrous) in 2 mL and 1 single use vial solvent 6 mL	1	..	..	1361.62	28.60	Taxotere	AV

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PACLITAXEL								
Authority required								
<i>Adjuvant treatment of node-positive, oestrogen receptor negative, breast cancer administered sequentially to doxorubicin hydrochloride and cyclophosphamide;</i>								
<i>Advanced breast cancer after failure of prior therapy which includes an anthracycline;</i>								
<i>Advanced metastatic ovarian cancer after failure of prior therapy which includes a platinum compound;</i>								
<i>Primary treatment of ovarian cancer in combination with a platinum compound;</i>								
<i>Locally advanced or metastatic non-small cell lung cancer.</i>								
3026G	Solution concentrate for I.V. infusion 30 mg in 5 mL	5	..	..	*1020.90	28.60	^a Anzatax ^a Paclitaxel Ebewe ^a Taxol	MX IT BQ
8018B	Solution concentrate for I.V. infusion 100 mg in 16.7 mL	2	..	..	*1373.55	28.60	^a Anzatax ^a Paclitaxel Ebewe ^a Taxol	MX IT BQ
3017T	Solution concentrate for I.V. infusion 150 mg in 25 mL	2	..	..	*2036.97	28.60	^a Anzatax ^a Paclitaxel Ebewe	MX IT
8360B	Solution concentrate for I.V. infusion 300 mg in 50 mL	1	..	..	2057.49	28.60	^a Anzatax ^a Paclitaxel Ebewe ^a Taxol	MX IT BQ
Cytotoxic antibiotics and related substances								
• Anthracyclines and related substances								
DOXORUBICIN HYDROCHLORIDE								
1336H	Solution for I.V. injection or intravesical administration 10 mg in 5 mL	4	..	..	*148.63	28.60	^a Adriamycin Solution ^a Doxorubicin Ebewe ^a MX	PH IT
1340M	Solution for I.V. injection or intravesical administration 20 mg in 10 mL	4	..	..	*256.67	28.60	Adriamycin Solution	PH
1342P	Solution for I.V. injection or intravesical administration 50 mg in 25 mL	3	..	..	*437.50	28.60	^a Adriamycin Solution ^a Doxorubicin Ebewe ^a MX	PH IT
8827N	Solution for I.V. injection or intravesical administration 100 mg in 50 mL	1	..	..	299.25	28.60	Doxorubicin Ebewe	IT
8828P	Solution for I.V. injection or intravesical administration 200 mg in 100 mL	1	..	..	579.87	28.60	Doxorubicin Ebewe	IT

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DOXORUBICIN HYDROCHLORIDE, PEGYLATED LIPOSOMAL								
Authority required								
<i>Advanced epithelial ovarian cancer in women who have failed a first-line platinum-based chemotherapy regimen;</i>								
<i>Metastatic breast cancer, as monotherapy, after failure of prior therapy which includes capecitabine and a taxane;</i>								
<i>Metastatic breast cancer, as monotherapy, where therapy with capecitabine and/or a taxane is contraindicated.</i>								
8569B	Suspension for I.V. infusion 20 mg in 10 mL	1	..	..	750.76	28.60	Caelyx	SH
8570C	Suspension for I.V. infusion 50 mg in 25 mL	1	..	..	1780.96	28.60	Caelyx	SH
1375J	EPIRUBICIN HYDROCHLORIDE Solution for injection 10 mg in 5 mL	4	..	..	*228.75	28.60	^a Epirubicin Ebewe	IT
							^a Pharmorubicin Solution	PH
1376K	Solution for injection 20 mg in 10 mL	4	..	..	*420.35	28.60	Pharmorubicin Solution	PH
1377L	Solution for injection 50 mg in 25 mL	3	..	..	*771.64	28.60	^a Epirubicin Ebewe	IT
							^a Pharmorubicin Solution	PH
8817C	Solution for injection 100 mg in 50 mL	2	..	..	*1013.55	28.60	Epirubicin Ebewe	IT
8858F	Solution for injection 200 mg in 100 mL	1	..	..	998.42	28.60	Epirubicin Ebewe	IT
IDARUBICIN HYDROCHLORIDE								
Restricted benefit								
<i>Acute myelogenous leukaemia.</i>								
2446R	Capsule 5 mg	3	..	..	*246.31	28.60	Zavedos	PH
2448W	Capsule 10 mg	3	..	..	*457.96	28.60	Zavedos	PH
8530Y	Solution for I.V. injection 5 mg in 5 mL	3	..	..	586.00	28.60	Zavedos Solution	PH
2452C	Powder for I.V. injection 5 mg	3	..	..	*586.00	28.60	Zavedos	PH
8531B	Solution for I.V. injection 10 mg in 10 mL	6	..	..	2267.83	28.60	Zavedos Solution	PH
2453D	Powder for I.V. injection 10 mg	6	..	..	*2267.83	28.60	Zavedos	PH
1932Q	MITOZANTRONE HYDROCHLORIDE Injection 10 mg (base) in 5 mL	1	..	..	187.81	28.60	PU	
1929M	Injection 20 mg (base) in 10 mL	1	..	..	355.59	28.60	^a Novantrone ^a Onkotrone ^a PU	SI BX
1930N	Injection 25 mg (base) in 12.5 mL	1	..	..	438.55	28.60	^a Novantrone ^a Onkotrone ^a PU	SI BX

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
• Other cytotoxic antibiotics BLEOMYCIN SULFATE See Special Pharmaceutical Benefits							
Other antineoplastic agents • Platinum compounds							
CARBOPLATIN							
1160C	Solution for I.V. injection 50 mg in 5 mL	2	..	..	*79.33	28.60 ^a	MX PU
1161D	Solution for I.V. injection 150 mg in 15 mL	6	..	..	*515.47	28.60 ^a	MX PU
1162E	Solution for I.V. injection 450 mg in 45 mL	2	..	..	*331.19	28.60 ^a	MX PU
CISPLATIN							
2578Q	I.V. injection 10 mg in 10 mL	1	..	..	14.46	15.43	PU
2579R	I.V. injection 50 mg in 50 mL	1	..	..	30.81	28.60 ^a	MX PU
2580T	I.V. injection 100 mg in 100 mL	1	..	..	70.42	28.60	MX
OXALIPLATIN							
Authority required							
<i>Metastatic colorectal cancer in patients with a WHO performance status of 2 or less, to be used in combination with 5-fluorouracil and folinic acid;</i>							
➤ <i>Adjuvant treatment of stage III (Dukes C) colon cancer, in combination with 5-fluorouracil and folinic acid, following complete resection of the primary tumour.</i>							
NOTE:							
<i>Oxaliplatin is not PBS-subsidised for the treatment of patients with stage II (Dukes B) colon cancer.</i>							
<i>Oxaliplatin is not PBS-subsidised for the adjuvant treatment of patients with rectal cancer.</i>							
8539K	Powder for I.V. infusion 50 mg	1	2	..	440.05	28.60	Eloxatin SW
8847P	Solution concentrate for I.V. infusion 50 mg in 10 mL	1	2	..	440.05	28.60	Eloxatin SW
8540L	Powder for I.V. infusion 100 mg	1	2	..	843.04	28.60	Eloxatin SW
8848Q	Solution concentrate for I.V. infusion 100 mg in 20 mL	1	2	..	843.04	28.60	Eloxatin SW
• Monoclonal antibodies RITUXIMAB Authority required <i>Relapsed or refractory low-grade B-cell non-Hodgkin's lymphoma;</i> <i>Relapsed or refractory follicular B-cell non-Hodgkin's lymphoma.</i>							
8293L	Solution for I.V. infusion 100 mg in 10 mL	2	3	..	946.40	28.60	Mabthera RO
8294M	Solution for I.V. infusion 500 mg in 50 mL	1	3	..	2358.86	28.60	Mabthera RO

continued ➤

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
<u>RITUXIMAB—cont.</u>								
<u>Authority required</u>								
<i>Treatment of previously untreated, CD20 positive, diffuse large B-cell non-Hodgkin's lymphoma, in combination with chemotherapy.</i>								
8665C	<i>Solution for I.V. infusion 100 mg in 10 mL</i>	2	7	..	946.40	28.60	<i>Mabthera</i>	<i>RO</i>
8666D	<i>Solution for I.V. infusion 500 mg in 50 mL</i>	1	7	..	2358.86	28.60	<i>Mabthera</i>	<i>RO</i>

• Other antineoplastic agents

GEFITINIB**NOTE:**

Any queries concerning the arrangements to prescribe gefitinib may be directed to Medicare Australia on 1800 242 679.

Written applications for authority to prescribe gefitinib should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Authority required

Initial PBS-subsidised treatment, as monotherapy, of locally advanced or metastatic non-small cell lung cancer in patients with a WHO performance status of 2 or less, where:

- (1) disease progression has occurred following treatment with at least 1 chemotherapy agent; and
- (2) there is evidence that the patient has an activating mutation(s) of the epidermal growth factor receptor (EGFR) gene in tumour material. The mutation(s) must be demonstrated by analysis of the DNA sequence of the EGFR gene.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Gefitinib (Iressa) PBS Authority Application for Use in the Treatment of Locally Advanced or Metastatic Non-Small Cell Lung Cancer - Supporting Information Form [may be downloaded from the Medicare Australia website at www.medicareaustralia.gov.au]; and
- (3) details of the prior chemotherapy including the name(s) of drug(s) and date of the most recent treatment cycle; and
- (4) details of the patient's WHO performance status; and
- (5) a copy of the pathology report providing evidence of the presence of activating mutation(s) of the EGFR gene from an Approved Pathology Authority.

Patients commenced on gefitinib therapy between 1 July 2004 and 1 December 2004 must demonstrate that they would have met the criteria for initial PBS-subsidised treatment at the time treatment with gefitinib was commenced.

Medical practitioners who wish to apply for authority to prescribe gefitinib for patients who commenced on gefitinib therapy prior to 1 July 2004 should contact Medicare Australia on 1800 242 679.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
GEFITINIB—cont.								
Authority required								
<i>Continuing PBS-subsidised treatment, as monotherapy, of locally advanced or metastatic non-small cell lung cancer in patients with a WHO performance status of 2 or less, where the patient has previously been issued with an authority prescription for gefitinib.</i>								
<i>Applications for continuing treatment may be made in writing or on the telephone by contacting Medicare Australia on 1800 242 679.</i>								
8769M	Tablet 250 mg	30	1	..	4286.09	28.60	Iressa	AP
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
3093T	HYDROXYUREA Capsule 500 mg	100	..	..	77.13	28.60	Hydrea	BQ
IRINOTECAN HYDROCHLORIDE TRIHYDRATE								
Authority required								
<i>Metastatic colorectal cancer in patients with a WHO performance status of 2 or less.</i>								
NOTE:								
<i>In first-line usage, effectiveness and tolerance may be improved when irinotecan is combined with an infusional 5-fluorouracil regimen.</i>								
8414W	I.V. injection 40 mg in 2 mL	1	3	..	169.75	28.60	^a Camptosar ^a MX	PU
8415X	I.V. injection 100 mg in 5 mL	2	3	..	*784.75	28.60	^a Camptosar ^a MX	PU
TOPOTECAN HYDROCHLORIDE								
Authority required								
<i>Advanced metastatic ovarian cancer after failure of prior therapy which includes a platinum compound.</i>								
8199M	Powder for I.V. infusion 4 mg (base)	5	1	..	2292.75	28.60	Hycamtin	GK
ENDOCRINE THERAPY								
Hormones and related agents								
• Progestogens								
MEDROXYPROGESTERONE ACETATE								
Restricted benefit								
<i>Hormone-dependent advanced breast cancer.</i>								
2728N	Tablet 500 mg	30	2	..	136.53	28.60	Provera	PH
Restricted benefit								
<i>Hormone-dependent breast cancer; Endometrial cancer.</i>								
2725K	Tablet 100 mg	100	2	..	98.82	28.60	Provera	PH
2316X	Tablet 200 mg	60	2	..	111.60	28.60	Provera	PH
2727M	Tablet 250 mg	60	2	..	138.32	28.60	Provera	PH
<i>[For other listings for this drug see Generic/Proprietary Index]</i>								

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MEGESTROL ACETATE								
<u>Restricted benefit</u>								
<i>Hormone-dependent advanced breast cancer.</i>								
2734X	Tablet 160 mg	30	2	..	73.50	28.60	Megace	BQ
• Gonadotropin releasing hormone analogues								
GOSERELIN ACETATE								
<u>Authority required</u>								
<i>Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate;</i>								
<i>Hormone-dependent locally advanced (equivalent to stage III) or metastatic (equivalent to stage IV) breast cancer in pre-menopausal women;</i>								
<i>Short-term treatment (up to 6 months) of visually proven endometriosis (only 1 course of not more than 6 months' therapy will be authorised).</i>								
1454M	Subcutaneous implant 3.6 mg (base) in pre-filled injection syringe	1	5	..	341.64	28.60	Zoladex Implant	AP
<u>Authority required</u>								
<i>Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate.</i>								
8093Y	Subcutaneous implant (long acting) 10.8 mg (base) in pre-filled injection syringe	1	1	..	1148.75	28.60	Zoladex 10.8 Implant	AP
LEUPRORELIN ACETATE								
<u>Authority required</u>								
<i>Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate.</i>								
1565J	I.M. injection (modified release), set containing 1 vial powder for injection 7.5 mg, 1 ampoule diluent 2 mL and 1 syringe with 2 needles	1	5	..	431.75	28.60	Lucrin Depot	AB
8707G	Suspension for subcutaneous injection (modified release), 7.5 mg injection set	1	5	..	431.75	28.60	Eligard 1 month	MX
8211E	I.M. injection (modified release), set containing 1 vial powder for injection 22.5 mg, 1 ampoule diluent 2 mL and 1 syringe with 2 needles	1	1	..	1148.75	28.60	Lucrin Depot 3 Month Injection	AB
8708H	Suspension for subcutaneous injection (modified release), 22.5 mg injection set	1	1	..	1148.75	28.60	Eligard 3 month	MX
8484M	I.M. injection (modified release), set containing 1 vial powder for injection 30 mg, 1 ampoule diluent 2 mL and 1 syringe with 2 needles	1	1	..	1529.39	28.60	Lucrin Depot 4 Month Injection	AB

continued ➡

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
LEUPRORELIN ACETATE—cont.								
8709J	Suspension for subcutaneous injection (modified release), 30 mg injection set	1	1	..	1529.39	28.60	Eligard 4 month	MX
8859G	Suspension for subcutaneous injection (modified release), 45 mg injection set	1	..	..	2289.99	28.60	Eligard 6 month	MX
Hormone antagonists and related agents								
• Anti-estrogens								
TAMOXIFEN CITRATE								
Restricted benefit								
<i>Treatment of hormone-dependent breast cancer.</i>								
NOTE:								
<i>This drug is not PBS-subsidised for primary prevention of breast cancer.</i>								
2109B	Tablet 10 mg (base)	60	5	..	47.54	28.60	^a Genox 10 ^a Tamoxen 10 mg ^a Tamoxifen Hexal	AF DP HX
				^B 2.16	[*] 49.71	28.60	^a Nolvadex	AP
2110C	Tablet 20 mg (base)	60	5	..	78.91	28.60	^a Chem mart Tamoxifen ^a Genox 20 ^a GenRx Tamoxifen ^a Tamosin ^a Tamoxen 20 mg ^a Tamoxifen Hexal ^a Terry White Chemists Tamoxifen	CH AF GX SI DP HX TW
				^B 3.94	[*] 82.85	28.60	^a Nolvadex-D	AP
TOREMIFENE CITRATE								
Restricted benefit								
<i>Treatment of hormone-dependent metastatic breast cancer in post-menopausal patients.</i>								
NOTE:								
<i>This drug is not PBS-subsidised for primary prevention of breast cancer.</i>								
8216K	Tablet 60 mg (base)	30	5	..	69.53	28.60	Fareston	SH
• Anti-androgens								
BICALUTAMIDE								
Authority required								
<i>Metastatic (equivalent to stage D) prostatic carcinoma in combination with GnRH (LH-RH) agonist therapy.</i>								
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8094B	Tablet 50 mg	28	5	..	225.67	28.60	Cosudex	AP

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CYPROTERONE ACETATE								
Authority required								
<i>Advanced carcinoma of the prostate; To reduce drive in sexual deviations in males.</i>								
1270W	Tablet 50 mg	100	5	..	*275.31	28.60	^a Cyprohexal	HX
							^a Cyprone	AF
							^a Cyprostat	SY
							^a GenRx	GX
							Cyproterone Acetate	
							^a Procur	DP
					^B 3.28 *278.59	28.60	^a Androcur	SC
8019C	Tablet 100 mg	50	5	..	227.34	28.60	^a Cyprohexal	HX
							^a Cyprostat-100	SY
							^a GenRx	GX
							Cyproterone Acetate	
							^a Procur 100	DP
					^B 1.00 228.34	28.60	^a Androcur-100	SC

[For other listings for this drug see Generic/Proprietary Index]

FLUTAMIDE**Authority required**

Metastatic (equivalent to stage D) prostatic carcinoma in combination with GnRH (LH-RH) agonist therapy.

NOTE:

No applications for increased maximum quantities and/or repeats will be authorised.

1417N	Tablet 250 mg	100	5	..	221.64	28.60	^a Eulexin	SH
							^a Flutamin	AF

NILUTAMIDE**Authority required**

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) prostatic carcinoma, in combination with GnRH (LH-RH) agonist therapy;

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) prostatic carcinoma, in conjunction with surgical orchidectomy.

8131Y	Tablet 150 mg	30	5	..	231.41	28.60	Anandron	AV

• **Enzyme inhibitors**

1036M	AMINOGLUTETHIMIDE Tablet 250 mg	100	5	..	163.04	28.60	Cytadren 250	NV

ANASTROZOLE**Restricted benefit**

*Treatment of hormone-dependent advanced breast cancer in post-menopausal women;
Treatment of hormone-dependent early breast cancer in post-menopausal women in whom tamoxifen citrate therapy is contraindicated;
Treatment of hormone-dependent early breast cancer in post-menopausal women who are intolerant of tamoxifen citrate.*

NOTE:

This drug is not PBS-subsidised for primary prevention of breast cancer.

8179L	Tablet 1 mg	30	2	..	217.14	28.60	Arimidex	AP

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
EXEMESTANE								
Restricted benefit								
<i>Treatment of hormone-dependent advanced breast cancer in post-menopausal women with disease progression following treatment with tamoxifen citrate.</i>								
NOTE:								
<i>This drug is not PBS-subsidised for primary prevention of breast cancer.</i>								
8506Q	Tablet 25 mg	30	2	..	224.65	28.60	Aromasin	PH
LETROZOLE								
Restricted benefit								
<i>Treatment of hormone-dependent advanced breast cancer in post-menopausal women.</i>								
NOTE:								
<i>This drug is not PBS-subsidised for primary prevention of breast cancer.</i>								
8245Y	Tablet 2.5 mg	30	2	..	217.14	28.60	Femara 2.5 mg	NV
IMMUNOSTIMULANTS								
Cytokines and immunomodulators								
• Interferons								
INTERFERON ALFA-2a								
CAUTION:								
<i>Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.</i>								
Authority required								
<i>Hairy cell leukaemia;</i>								
<i>Myeloproliferative disease with excessive thrombocytosis.</i>								
8180M	Injection 3,000,000 i.u. in 0.5 mL single dose pre-filled syringe	15	4	..	*521.20	28.60	Roferon-A	RO
Authority required								
<i>Myeloproliferative disease with excessive thrombocytosis.</i>								
8551C	Injection 4,500,000 i.u. in 0.5 mL single dose pre-filled syringe	5	4	..	*271.10	28.60	Roferon-A	RO
8552D	Injection 6,000,000 i.u. in 0.5 mL single dose pre-filled syringe	5	4	..	*353.75	28.60	Roferon-A	RO
8553E	Injection 9,000,000 i.u. in 0.5 mL single dose pre-filled syringe	5	4	..	*521.15	28.60	Roferon-A	RO

continued ➡

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
INTERFERON ALFA-2a—cont.								
Authority required								
<i>Low grade non-Hodgkin's lymphoma with clinical features suggestive of a poor prognosis, in combination with anthracycline-based chemotherapy.</i>								
8181N	Injection 3,000,000 i.u. in 0.5 mL single dose pre-filled syringe	15	5	..	*521.20	28.60	Roferon-A	RO
8182P	Injection 4,500,000 i.u. in 0.5 mL single dose pre-filled syringe	5	5	..	*271.10	28.60	Roferon-A	RO
8183Q	Injection 6,000,000 i.u. in 0.5 mL single dose pre-filled syringe	5	5	..	*353.75	28.60	Roferon-A	RO
8184R	Injection 9,000,000 i.u. in 0.5 mL single dose pre-filled syringe	5	5	..	*521.15	28.60	Roferon-A	RO
INTERFERON ALFA-2b								
CAUTION:								
<i>Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.</i>								
Authority required								
<i>Hairy cell leukaemia.</i>								
8572E	Solution for injection 18,000,000 i.u. in 1.2 mL multi-dose injection pen	3	4	..	*624.37	28.60	Intron A Redipen	SH
Authority required								
<i>Maintenance treatment of multiple myeloma once remission has been achieved with chemotherapy;</i>								
<i>Low grade non-Hodgkin's lymphoma with clinical features suggestive of a poor prognosis, in combination with anthracycline-based chemotherapy.</i>								
8348J	Solution for injection 18,000,000 i.u. in 1.2 mL multi-dose injection pen	3	5	..	*624.37	28.60	Intron A Redipen	SH
8476D	Solution for injection 30,000,000 i.u. in 1.2 mL multi-dose injection pen	3	5	..	*1037.47	28.60	Intron A Redipen	SH

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
INTERFERON BETA-1a								
Authority required								
<i>Initial treatment of clinically definite relapsing-remitting multiple sclerosis in ambulatory (without assistance or support) patients who have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to the multiple sclerosis, in the preceding 2 years. The diagnosis must be confirmed by magnetic resonance imaging of the brain and/or spinal cord and the date of the scan included in the authority application, unless the authority application is accompanied by written certification provided by a radiologist that an MRI scan is contraindicated because of the risk of physical (not psychological) injury to the patient. The authority will be limited to the maximum quantity and number of repeats indicated in the schedule;</i>								
<i>Continuing treatment of clinically definite relapsing-remitting multiple sclerosis in patients previously issued with an authority prescription for this drug who do not show continuing progression of disability while on treatment with this drug and who have demonstrated compliance with, and an ability to tolerate, this therapy. Authorities will be limited to the maximum quantity and number of repeats indicated in the schedule.</i>								
8289G	Injection set comprising 1 vial powder for injection 30 micrograms (6,000,000 i.u.) with diluent	4	5	..	1090.98	28.60	Avonex	BD
8805K	Injection 30 micrograms (6,000,000 i.u.) in 0.5 mL single dose pre-filled syringe	4	5	..	1090.98	28.60	Avonex	BD
8403G	Injection 44 micrograms (12,000,000 i.u.) in 0.5 mL single dose pre-filled syringe	12	5	..	1090.98	28.60	Rebif 44	SG
INTERFERON BETA-1b								
Authority required								
<i>Initial treatment of clinically definite relapsing-remitting multiple sclerosis in ambulatory (without assistance or support) patients who have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to the multiple sclerosis, in the preceding 2 years. The diagnosis must be confirmed by magnetic resonance imaging of the brain and/or spinal cord and the date of the scan included in the authority application, unless the authority application is accompanied by written certification provided by a radiologist that an MRI scan is contraindicated because of the risk of physical (not psychological) injury to the patient. The authority will be limited to the maximum quantity and number of repeats indicated in the schedule;</i>								
<i>Continuing treatment of clinically definite relapsing-remitting multiple sclerosis in patients previously issued with an authority prescription for this drug who do not show continuing progression of disability while on treatment with this drug and who have demonstrated compliance with, and an ability to tolerate, this therapy. Authorities will be limited to the maximum quantity and number of repeats indicated in the schedule.</i>								
8101J	Injection set comprising 1 vial powder for injection 8,000,000 i.u. (250 micrograms) and solvent	15	5	..	1178.49	28.60	Betaferon	SC

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Other cytokines and immunomodulators BCG IMMUNOTHERAPEUTIC <i>(Bacillus Calmette-Guérin/ Connaught strain)</i> <u>Restricted benefit</u> <i>Treatment of carcinoma in situ of the urinary bladder.</i>								
1140B	Single dose set comprising 1 vial powder for intravesical administration containing 6.6 to 19.2 x 10 ⁸ CFU and 1 vial diluent 3 mL	3	1	..	*472.75	28.60	ImmuCyst	AV
BCG-TICE <i>(Bacillus Calmette-Guérin/ Tice strain)</i> <u>Restricted benefit</u> <i>Primary and relapsing superficial urothelial carcinoma of the bladder.</i>								
1131M	Vial containing powder for intravesical administration approximately 5 x 10 ⁸ CFU	3	1	..	548.97	28.60	OncoTICE	OR
GLATIRAMER ACETATE <u>Authority required</u> <i>Initial treatment of clinically definite relapsing-remitting multiple sclerosis in ambulatory (without assistance or support) patients who have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to the multiple sclerosis, in the preceding 2 years. The diagnosis must be confirmed by magnetic resonance imaging of the brain and/or spinal cord and the date of the scan included in the authority application, unless the authority application is accompanied by written certification provided by a radiologist that an MRI scan is contraindicated because of the risk of physical (not psychological) injury to the patient. The authority will be limited to the maximum quantity and number of repeats indicated in the schedule;</i> <i>Continuing treatment of clinically definite relapsing-remitting multiple sclerosis in patients previously issued with an authority prescription for this drug who do not show continuing progression of disability while on treatment with this drug and who have demonstrated compliance with, and an ability to tolerate, this therapy. Authorities will be limited to the maximum quantity and number of repeats indicated in the schedule.</i>								
8726G	Injection 20 mg in 1 mL single dose pre-filled syringe	28	5	..	1090.98	28.60	Copaxone	AV

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum Recordable Value for Safety Net	Proprietary Name and Manufacturer
					for Max. Qty \$	\$	

IMMUNOSUPPRESSIVE AGENTS

Immunosuppressive agents

• Selective immunosuppressive agents

ADALIMUMAB

NOTE:

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia

Prior Written Approval of Specialised Drugs

Reply Paid 9826

GPO Box 9826

HOBART TAS 7001

Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au.

NOTE:

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the tumour necrosis factor (TNF) alfa antagonists (adalimumab, etanercept and infliximab) and the interleukin-1 inhibitor (anakinra) for adult patients with severe rheumatoid arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs (bDMARDs) at any 1 time.

1. Patients who have received no prior PBS-subsidised bDMARD treatment at 1 December 2004.

From 1 December 2004, the arrangements for prescribing the bDMARDs on the PBS have been amended to allow patients to commence a single cycle of bDMARD treatment that allows them to trial any number of bDMARDs without having to experience a disease flare when swapping between alternate bDMARDs. Within a single treatment cycle, patients may continue to receive long-term treatment with a bDMARD while they continue to show a response to therapy.

Once patients have either failed, or ceased to respond to, treatment with a maximum of 3 bDMARDs, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised bDMARD therapy before they are eligible to commence the next cycle. The 5-year period will be measured from the date the last prescription for PBS-subsidised bDMARD treatment was approved in the most recent cycle to the date of the application for initial treatment with a bDMARD under the new cycle.

Patients who have failed treatment with fewer than 3 bDMARDs within a particular treatment cycle, and where a period of less than 5 years duration has elapsed since the patient's previous course of PBS-subsidised bDMARD treatment in that cycle, may commence a further course of bDMARD treatment within that same treatment cycle.

Patients who have failed treatment with fewer than 3 bDMARDs within a particular treatment cycle, and who have had a break in PBS-subsidised therapy of 5 years or more, are eligible to commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

continued ➞

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ADALIMUMAB—cont.

If patients fail to respond to a particular bDMARD within a single treatment cycle, they are not eligible to receive further PBS-subsidised treatment with that drug until they commence the next cycle.

2. Patients who have received PBS-subsidised TNF-alfa antagonist treatment prior to 1 December 2004.

Patients who commenced PBS-subsidised TNF-alfa antagonist therapy prior to 1 December 2004 are considered to be in their first cycle of bDMARD treatment on 1 December 2004.

Patients who have failed to respond to prior PBS-subsidised treatment with fewer than 3 TNF-alfa antagonists at the time the first application for treatment is made on or after 1 December 2004, will be subject to the same conditions applying to new patients detailed above.

Therefore, patients who have failed:

(a) 1 TNF-alfa antagonist will be eligible to trial further PBS-subsidised treatment with a bDMARD they have not failed in their first treatment cycle, until they fail to demonstrate a response to no more than another 2 bDMARDs;

(b) 2 TNF-alfa antagonists will be eligible to trial further PBS-subsidised treatment with a bDMARD they have not failed in their first treatment cycle, until they fail to demonstrate a response to no more than 1 other bDMARD.

Patients who have failed PBS-subsidised treatment with 3 TNF-alfa antagonists prior to 1 December 2004 or at the first assessment required after this date, will be eligible to trial PBS-subsidised treatment with anakinra if they wish. However, if they fail to demonstrate a response to anakinra, they will not be able to trial any further PBS-subsidised bDMARD treatment until a minimum of 5 years has elapsed from the date that the prescription for the last course of anakinra therapy was approved. Arrangements to allow these patients to fail 4 bDMARDs will only be in place for the first treatment cycle. For subsequent cycles, patients will cease to be eligible to receive PBS-subsidised bDMARD treatment once they have failed to demonstrate a response to a maximum of 3 bDMARDs.

Any queries on these arrangements should be forwarded to Medicare Australia.

3. Information relevant to all patients.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) patients have received no prior PBS-subsidised bDMARD treatment and wish to commence such therapy; or

(ii) patients have received prior PBS-subsidised (initial or continuing) bDMARD therapy and wish to trial an alternate agent [further details are under 'Swapping therapy' below]; or

(iii) patients wish to re-commence treatment with a specific bDMARD following a break in PBS-subsidised therapy with that specific agent.

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted bDMARD supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that bDMARD.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed	Maximum	Proprietary Name and Manufacturer
					Price for Max. Qty \$	Recordable Value for Safety Net \$	

ADALIMUMAB—cont.**(b) Continuing treatment.**

Following the completion of an initial treatment course with a specific bDMARD, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that bDMARD.

(c) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised bDMARD is approved, patients may swap to an alternate bDMARD without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-bDMARD therapy requirements. However, the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialed.

Patients may swap to an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application or not.

Patients may alternate between therapy with any bDMARD of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular bDMARD previously; or*
- (ii) they have demonstrated an adequate response to that particular bDMARD if they have previously trialed it on the PBS.*

Therefore, to maximise the choice of bDMARD patients may alternate between, it is important that patients are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(d) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to a reduction in the total number of active joints.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ADALIMUMAB—cont.

(e) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy. Patients who wish to trial a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with at least 1 non-biological DMARD, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Authority required

Application for initial PBS-subsidised treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and*
- (b) have received no prior PBS-subsidised treatment with a bDMARD for this condition in this treatment cycle; and*
- (c) have failed to achieve an adequate response to the following treatments:*
 - (i) methotrexate at a dose of at least 20 mg weekly; and*
 - (ii) methotrexate (at a minimum dose of 7.5 mg weekly), in combination with 2 other non-biological disease modifying anti-rheumatic drugs (DMARDs), for a minimum of 3 months; and*
 - (iii) a minimum of 3 months' treatment with:*
 - leflunomide alone; or*
 - leflunomide in combination with methotrexate; or*
 - cyclosporin.*

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or intolerance of a severity necessitating permanent treatment withdrawal develops during the relevant period of use, the patient is exempted from demonstrating an inadequate response to that particular agent(s) only. Details of the contraindications or intolerance, including the degree of toxicity, must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrable in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L;

AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or*
- (ii) at least 4 active joints from the following list of major joints:*
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or*
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).*

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ADALIMUMAB—cont.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and*
 - (2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes details of the patient's ESR and CRP measurements and the patient's active joint count which must have been assessed no earlier than 1 month prior to the date of application; and*
 - (3) a copy of the signed patient acknowledgement form which may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm).*
- Completion of this form declares that the patient understands and acknowledges that, within a single treatment cycle, PBS-subsidised treatment with any biological DMARD will cease if they do not demonstrate the response to treatment required to support continuation of PBS-subsidised treatment at any assessment where a response must be demonstrated.*

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Authority required

Application for an initial course of PBS-subsidised treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and*
- (b) have received prior PBS-subsidised bDMARD treatment for this condition in this treatment cycle and are eligible to receive further bDMARD therapy.*

Applications for patients who have received PBS-subsidised treatment with adalimumab within this treatment cycle and who wish to re-commence therapy with this drug within this same cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised adalimumab treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised adalimumab treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised bDMARD therapy or, under this restriction, for patients who have received previous PBS-subsidised bDMARD therapy), patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised adalimumab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum Recordable Value for	Proprietary Name and Manufacturer
					for Max. Qty \$	Safety Net \$	
ADALIMUMAB—cont.							
<i>The authority application must be made in writing and must include:</i>							
<i>(1) a completed authority prescription form; and</i>							
<i>(2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website</i>							
<i>(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].</i>							
<i>Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).</i>							
<i>Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.</i>							
<i>Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.</i>							
8737W	Injection 40 mg in 0.8 mL pre-filled syringe	2	3	..	1888.30	28.60	Humira AB
NOTE:							
<i>No applications for increased maximum quantities and/or repeats will be authorised. Applications for treatment with adalimumab where the dosing frequency exceeds 40 mg per fortnight will not be approved.</i>							

Authority required

Initial PBS-subsidised supply for continuing treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and*
- (b) were receiving treatment with adalimumab prior to 1 November 2004; and*
- (c) failed to qualify for PBS-subsidised therapy after 1 May 2004 due to an inability to receive concomitant methotrexate; and*
- (d) have demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with adalimumab.*

Medical practitioners who wish to apply for authority to prescribe a bDMARD for patients who commenced treatment with adalimumab prior to 1 November 2004 and who have not demonstrated a response to treatment should contact Medicare Australia on 1800 005 750.

The authority application must be in writing and must include sufficient information to determine the patient's eligibility. The date of assessment must be provided.

Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ADALIMUMAB—cont.

Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Authority required

Initial PBS-subsidised supply for continuing treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and*
- (b) were receiving treatment with adalimumab prior to 1 March 2005; and*
- (c) failed to qualify for PBS-subsidised therapy after 1 May 2004 due to testing negative for rheumatoid factor; and*
- (d) have demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with adalimumab.*

Medical practitioners who wish to apply for authority to prescribe a bDMARD for patients who commenced treatment with adalimumab prior to 1 March 2005 and who have not demonstrated a response to treatment should contact Medicare Australia on 1800 005 750.

The authority application must be in writing and must include sufficient information to determine the patient's eligibility. The date of assessment must be provided.

Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Authority required

Continuing PBS-subsidised treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have severe active rheumatoid arthritis; and*
- (b) who have demonstrated an adequate response to treatment with adalimumab; and*
- (c) whose most recent course of PBS-subsidised bDMARD treatment in this treatment cycle was with adalimumab.*

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ADALIMUMAB—cont.							
	An adequate response to treatment is defined as:						
	an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;						
	AND either of the following:						
	(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or						
	(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:						
	— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or						
	— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).						
	The authority application must be made in writing and must include:						
	(1) a completed authority prescription form; and						
	(2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].						
	All applications for continuing treatment with adalimumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with adalimumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.						
	Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).						
	Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.						
	Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.						
8741C	Injection 40 mg in 0.8 mL pre-filled syringe	2	5	..	1888.30	28.60	Humira AB
	<u>NOTE:</u>						
	No applications for increased maximum quantities and/or repeats will be authorised. Applications for treatment with adalimumab where the dosing frequency exceeds 40 mg per fortnight will not be approved.						

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed	Maximum	Proprietary Name and Manufacturer
					Price for Max. Qty \$	Recordable Value for Safety Net \$	

ANAKINRA**NOTE:**

Any queries concerning the arrangements to prescribe anakinra may be directed to Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe anakinra should be forwarded to:

*Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001*

Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au.

NOTE:

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the tumour necrosis factor (TNF) alfa antagonists (adalimumab, etanercept and infliximab) and the interleukin-1 inhibitor (anakinra) for adult patients with severe rheumatoid arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs (bDMARDs) at any 1 time.

1. Patients who have received no prior PBS-subsidised bDMARD treatment at 1 December 2004.

From 1 December 2004, the arrangements for prescribing the bDMARDs on the PBS have been amended to allow patients to commence a single cycle of bDMARD treatment that allows them to trial any number of bDMARDs without having to experience a disease flare when swapping between alternate bDMARDs. Within a single treatment cycle, patients may continue to receive long-term treatment with a bDMARD while they continue to show a response to therapy.

Once patients have either failed, or ceased to respond to, treatment with a maximum of 3 bDMARDs, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised bDMARD therapy before they are eligible to commence the next cycle. The 5-year period will be measured from the date the last prescription for PBS-subsidised bDMARD treatment was approved in the most recent cycle to the date of the application for initial treatment with a bDMARD under the new cycle.

Patients who have failed treatment with fewer than 3 bDMARDs within a particular treatment cycle, and where a period of less than 5 years duration has elapsed since the patient's previous course of PBS-subsidised bDMARD treatment in that cycle, may commence a further course of bDMARD treatment within that same treatment cycle.

Patients who have failed treatment with fewer than 3 bDMARDs within a particular treatment cycle, and who have had a break in PBS-subsidised therapy of 5 years or more, are eligible to commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

If patients fail to respond to a particular bDMARD within a single treatment cycle, they are not eligible to receive further PBS-subsidised treatment with that drug until they commence the next cycle.

continued ➞

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ANAKINRA—cont.

2. Patients who have received PBS-subsidised TNF-alfa antagonist treatment prior to 1 December 2004.

Patients who commenced PBS-subsidised TNF-alfa antagonist therapy prior to 1 December 2004 are considered to be in their first cycle of bDMARD treatment on 1 December 2004.

Patients who have failed to respond to prior PBS-subsidised treatment with fewer than 3 TNF-alfa antagonists at the time the first application for treatment is made on or after 1 December 2004, will be subject to the same conditions applying to new patients detailed above.

Therefore, patients who have failed:

- (a) 1 TNF-alfa antagonist will be eligible to trial further PBS-subsidised treatment with a bDMARD they have not failed in their first treatment cycle, until they fail to demonstrate a response to no more than another 2 bDMARDs;*
- (b) 2 TNF-alfa antagonists will be eligible to trial further PBS-subsidised treatment with a bDMARD they have not failed in their first treatment cycle, until they fail to demonstrate a response to no more than 1 other bDMARD.*

Patients who have failed PBS-subsidised treatment with 3 TNF-alfa antagonists prior to 1 December 2004 or at the first assessment required after this date, will be eligible to trial PBS-subsidised treatment with anakinra if they wish. However, if they fail to demonstrate a response to anakinra, they will not be able to trial any further PBS-subsidised bDMARD treatment until a minimum of 5 years has elapsed from the date that the prescription for the last course of anakinra therapy was approved. Arrangements to allow these patients to fail 4 bDMARDs will only be in place for the first treatment cycle. For subsequent cycles, patients will cease to be eligible to receive PBS-subsidised bDMARD treatment once they have failed to demonstrate a response to a maximum of 3 bDMARDs.

Any queries on these arrangements should be forwarded to Medicare Australia.

3. Information relevant to all patients.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised bDMARD treatment and wish to commence such therapy; or*
- (ii) patients have received prior PBS-subsidised (initial or continuing) bDMARD therapy and wish to trial an alternate agent [further details are under 'Swapping therapy' below]; or*
- (iii) patients wish to re-commence treatment with a specific bDMARD following a break in PBS-subsidised therapy with that specific agent.*

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted bDMARD supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that bDMARD.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed	Maximum	Proprietary Name and Manufacturer
					Price for Max. Qty \$	Recordable Value for Safety Net \$	

ANAKINRA—cont.**(b) Continuing treatment.**

Following the completion of an initial treatment course with a specific bDMARD, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that bDMARD.

(c) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised bDMARD is approved, patients may swap to an alternate bDMARD without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-bDMARD therapy requirements. However, the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialed.

Patients may swap to an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application or not.

Patients may alternate between therapy with any bDMARD of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular bDMARD previously; or**
- (ii) they have demonstrated an adequate response to that particular bDMARD if they have previously trialed it on the PBS.**

Therefore, to maximise the choice of bDMARD patients may alternate between, it is important that patients are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(d) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to a reduction in the total number of active joints.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer

ANAKINRA—cont.

(e) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy. Patients who wish to trial a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with at least 1 non-biological DMARD, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Authority required

Application for initial PBS-subsidised treatment with anakinra, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and*
- (b) have received no prior PBS-subsidised treatment with a bDMARD for this condition in this treatment cycle; and*
- (c) have failed to achieve an adequate response to the following treatments:*
 - (i) methotrexate at a dose of at least 20 mg weekly; and*
 - (ii) methotrexate (at a minimum dose of 7.5 mg weekly), in combination with 2 other non-biological disease modifying anti-rheumatic drugs (DMARDs), for a minimum of 3 months; and*
 - (iii) a minimum of 3 months' treatment with:*
 - leflunomide alone; or*
 - leflunomide in combination with methotrexate; or*
 - cyclosporin.*

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or intolerance of a severity necessitating permanent treatment withdrawal develops during the relevant period of use, the patient is exempted from demonstrating an inadequate response to that particular agent(s) only. Details of the contraindications or intolerance, including the degree of toxicity, must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrable in all patients at the time of the initial application:

- an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L;*
- AND either*
 - (i) a total active joint count of at least 20 active (swollen and tender) joints; or*
 - (ii) at least 4 active joints from the following list of major joints:*
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or*
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).*

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and*
- (2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes details of the patient's ESR and CRP measurements and the patient's active joint count which must have been assessed no earlier than 1 month prior to the date of application; and*

continued ➤

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ANAKINRA—cont.

(3) a copy of the signed patient acknowledgement form which may be downloaded from the Medicare Australia website

(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm).

Completion of this form declares that the patient understands and acknowledges that, within a single treatment cycle, PBS-subsidised treatment with any biological DMARD will cease if they do not demonstrate the response to treatment required to support continuation of PBS-subsidised treatment at any assessment where a response must be demonstrated.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients who fail to demonstrate a response to treatment with anakinra under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial anakinra after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Authority required

Application for an initial course of PBS-subsidised treatment with anakinra, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

(a) have severe active rheumatoid arthritis; and

(b) have received prior PBS-subsidised bDMARD treatment for this condition in this treatment cycle and are eligible to receive further bDMARD therapy.

Applications for patients who have received PBS-subsidised treatment with anakinra within this treatment cycle and who wish to re-commence therapy with this drug within this same cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised anakinra treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised anakinra treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised bDMARD therapy or, under this restriction, for patients who have received previous PBS-subsidised bDMARD therapy), patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised anakinra treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

For the first treatment cycle only, applications for anakinra will also be approved under this restriction for patients who have failed to demonstrate a response to treatment with 3 PBS-subsidised TNF-alfa antagonists prior to 1 December 2004. Once these patients fail to demonstrate a response to anakinra, they will not be able to trial further PBS-subsidised bDMARDs until a 5-year period has elapsed from the date that the last course of anakinra therapy was approved.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ANAKINRA—cont.							
<i>The authority application must be made in writing and must include:</i> <i>(1) a completed authority prescription form; and</i> <i>(2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].</i> <i>Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).</i> <i>Patients who fail to demonstrate a response to treatment with anakinra under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial anakinra after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.</i> <i>Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.</i>							
8773R	Injection 100 mg in 0.67 mL single use pre-filled syringe	28	3	..	1433.33	28.60	Kineret AN
NOTE: No applications for increased maximum quantities and/or repeats will be authorised.							

Authority required

Initial PBS-subsidised supply for continuing treatment with anakinra, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

(a) have severe active rheumatoid arthritis; and

(b) were receiving treatment with anakinra prior to 1 July 2004; and

(c) have demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with anakinra.

Medical practitioners who wish to apply for authority to prescribe a bDMARD for patients who commenced treatment with anakinra prior to 1 July 2004 and who have not demonstrated a response to treatment should contact Medicare Australia on 1800 005 750.

The authority application must be in writing and must include sufficient information to determine the patient's eligibility. The date of assessment must be provided.

Patients who fail to demonstrate a response to treatment with anakinra under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial anakinra after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ANAKINRA—cont.

Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Authority required

Initial PBS-subsidised supply for continuing treatment with anakinra, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and*
- (b) were receiving treatment with anakinra prior to 1 March 2005; and*
- (c) failed to qualify for PBS-subsidised therapy after 1 December 2004 due to testing negative for rheumatoid factor; and*
- (d) have demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with anakinra.*

Medical practitioners who wish to apply for authority to prescribe a bDMARD for patients who commenced treatment with anakinra prior to 1 March 2005 and who have not demonstrated a response to treatment should contact Medicare Australia on 1800 005 750.

The authority application must be in writing and must include sufficient information to determine the patient's eligibility. The date of assessment must be provided.

Patients who fail to demonstrate a response to treatment with anakinra under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial anakinra after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Authority required

Continuing PBS-subsidised treatment with anakinra, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have severe active rheumatoid arthritis; and*
- (b) who have demonstrated an adequate response to treatment with anakinra; and*
- (c) whose most recent course of PBS-subsidised bDMARD treatment in this treatment cycle was with anakinra.*

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ANAKINRA—cont.							
An adequate response to treatment is defined as: an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; AND either of the following: (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%: — elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or — shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth). The authority application must be made in writing and must include: (1) a completed authority prescription form; and (2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm). For the first treatment cycle only, applications for anakinra will also be approved under this restriction for patients who have failed to demonstrate a response to treatment with 3 PBS-subsidised TNF-alfa antagonists prior to 1 December 2004. Once these patients fail to demonstrate a response to anakinra, they will not be able to trial further PBS-subsidised bDMARDs until a 5-year period has elapsed from the date that the last course of anakinra therapy was approved. All applications for continuing treatment with anakinra must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with anakinra, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course. Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Patients who fail to demonstrate a response to treatment with anakinra under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial anakinra after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle. Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.							
8774T	Injection 100 mg in 0.67 mL single use pre-filled syringe	28	5	..	1433.33	28.60	Kineret AN

NOTE:

No applications for increased maximum quantities and/or repeats will be authorised.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CYCLOSPORIN								
CAUTION:								
<i>Careful monitoring of patients is mandatory.</i>								
Authority required								
<i>Maintenance therapy, following initiation and stabilisation of treatment with cyclosporin, of:</i>								
<i>(a) patients with organ or tissue transplants. Therapy must remain under the supervision and direction of the transplant unit reviewing the patient. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application;</i>								
<i>(b) patients with severe atopic dermatitis for whom other systemic therapies are ineffective or inappropriate. Therapy must remain under the supervision and direction of a dermatologist, clinical immunologist or specialised unit reviewing the patient. The name of the dermatologist, clinical immunologist or specialised unit reviewing treatment and the date of the latest review must be included in the authority application;</i>								
<i>(c) patients with severe psoriasis for whom other systemic therapies are ineffective or inappropriate and in whom the disease has caused significant interference with quality of life. Therapy must remain under the supervision and direction of a dermatologist or specialised unit reviewing the patient. The name of the dermatologist or specialised unit reviewing treatment and the date of the latest review must be included in the authority application;</i>								
<i>(d) patients with nephrotic syndrome in whom steroids and cytostatic drugs have failed or are not tolerated or are considered inappropriate and in whom renal function is unimpaired. Therapy must remain under the supervision and direction of a nephrologist or specialised unit reviewing the patient. The name of the nephrologist or specialised unit reviewing treatment and the date of the latest review must be included in the authority application;</i>								
<i>(e) patients with severe active rheumatoid arthritis for whom classical slow-acting anti-rheumatic agents (including methotrexate) are ineffective or inappropriate. Therapy must remain under the supervision and direction of a rheumatologist, clinical immunologist or specialised unit reviewing the patient. The name of the rheumatologist, clinical immunologist or specialised unit reviewing treatment and the date of the latest review must be included in the authority application.</i>								
8657P	Capsule 10 mg	120	3	..	*95.67	28.60	Neoral 10	NV
8658Q	Capsule 25 mg	60	3	..	*104.13	28.60	^a Cysporin	MX
				^B 2.38	*106.51	28.60	^a Cicloral	HX
				^B 4.80	*108.93	28.60	^a Neoral 25	NV
8659R	Capsule 50 mg	60	3	..	*212.83	28.60	^a Cysporin	MX
				^B 2.38	*215.21	28.60	^a Cicloral	HX
				^B 4.76	*217.59	28.60	^a Neoral 50	NV
8660T	Capsule 100 mg	60	3	..	*412.49	28.60	^a Cysporin	MX
				^B 2.40	*414.89	28.60	^a Cicloral	HX
				^B 4.80	*417.29	28.60	^a Neoral 100	NV
8661W	Oral liquid 100 mg per mL, 50 mL	2	3	..	*734.59	28.60	Neoral	NV

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ETANERCEPT**NOTE:**

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe etanercept should be forwarded to:

*Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001*

Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au.

NOTE:

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of etanercept and infliximab for adult patients with HLA-B27 positive, active ankylosing spondylitis only. It does not apply to the prescribing of adalimumab. Therefore, where the term 'tumour necrosis factor (TNF) alfa antagonists' appears in the following NOTES and restrictions, it only refers to etanercept and infliximab.

Patients are eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alfa antagonists above at any 1 time.

From 1 August 2005, under the PBS, all patients will be able to commence a single treatment cycle where they may trial both of the PBS-subsidised TNF-alfa antagonists without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, patients may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

Once patients have either failed or ceased to respond to treatment up to 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. Within the same treatment cycle, patients are not allowed to trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than twice. Patients who have failed the same PBS-subsidised TNF-alfa antagonist twice must change to the alternate PBS-subsidised agent if they wish to continue PBS-subsidised TNF-alfa antagonist treatment.

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised TNF-alfa antagonist treatment was granted in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new cycle.

Where a period of less than 5 years duration has elapsed since the patient's previous course of PBS-subsidised TNF-alfa antagonist treatment, patients who have failed TNF-alfa antagonist therapy less than 3 times within a particular treatment cycle, as defined in the restriction, may commence a further course of treatment within that cycle.

Where a break in PBS-subsidised therapy of 5 years or more has occurred, patients who have failed TNF-alfa antagonist therapy less than 3 times within a particular treatment cycle, as defined in the restriction, are eligible to commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ETANERCEPT—cont.**1. Patients 'grandfathered' onto PBS-subsidised treatment with infliximab and etanercept.**

Both the infliximab and etanercept restrictions allow patients who commenced treatment prior to 1 March 2004 and 1 July 2004 respectively to commence treatment under a different set of criteria than those patients who commenced treatment after these dates (i.e. new patients). This means such patients were 'grandfathered' onto PBS-subsidised treatment. Consequently, the criteria to enable 'grandfather' patients to continue on PBS-subsidised treatment are also different to those for new patients.

Where patients who were 'grandfathered' onto PBS-subsidised treatment with 1 TNF- α antagonist wish to swap to the alternate agent, they can only do so using the restrictions that apply to 'grandfather' patients.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, 'grandfather' patients must requalify for initial treatment under the criteria that apply to new patients. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below for further details. Patients who commenced on PBS-subsidised etanercept or infliximab treatment prior to 1 August 2005 are deemed to be in their first cycle of therapy.

The first course of initial therapy for patients 'grandfathered' onto PBS-subsidised treatment with etanercept or infliximab will be limited to provide for 24 weeks of therapy. All subsequent initial treatment courses made under the 'grandfather' restriction will be limited to provide 6 weeks of therapy.

2. How to prescribe TNF- α antagonist therapy after 1 August 2005.**(a) Initial treatment.**

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised TNF- α antagonist treatment and wish to commence such therapy; or*
- (ii) patients have received prior PBS-subsidised (initial or continuing) TNF- α antagonist therapy and wish to trial an alternate agent [further details are under 'Swapping therapy' below]; or*
- (iii) patients have failed their most recent course of PBS-subsidised TNF- α antagonist therapy but wish to trial a further course of treatment with the same TNF- α antagonist (providing they have not failed that TNF- α antagonist more than once); or*
- (iv) patients wish to re-commence treatment with a specific TNF- α antagonist following a break in PBS-subsidised therapy with that specific agent.*

All applications for initial treatment will be limited to provide for a maximum of 6 weeks of therapy except for the first application for 'grandfather' patients approved for PBS-subsidised treatment with etanercept and infliximab where up to 24 weeks of treatment will be approved.

In order to demonstrate a response to therapy, patients who have received an initial PBS-subsidised course of up to 6 weeks of treatment must be assessed no earlier than 4 weeks from the commencement of the initial therapy course. In order to demonstrate a response to therapy, patients who have received an initial PBS-subsidised course of up to 24 weeks must be assessed in the 4 weeks prior to treatment being ceased.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ETANERCEPT—cont.							
Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that TNF-α antagonist.							
Prescribers should ensure that applications for second and subsequent courses of PBS-subsidised TNF-α antagonist treatment are submitted to Medicare Australia before patients complete their previous treatment course to ensure uninterrupted treatment. Applications where no provisions to fax the prescription and application are made in the restriction should be posted to Medicare Australia no later than 2 weeks prior to the patient completing their previous treatment course.							
(b) Continuing treatment.							
Following the completion of an initial treatment course with a specific TNF-α antagonist, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing TNF-α antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.							
Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.							
Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that TNF-α antagonist.							
(c) How to apply for TNF-α antagonist treatment on or after 1 August 2005.							
Patients for whom the first application for PBS-subsidised treatment with either etanercept or infliximab is made on or after 1 August 2005, will fall into 1 of the following categories:							
(I) Patients who have not received any prior PBS-subsidised infliximab or PBS-subsidised etanercept treatment.							
This includes the following:							
(i) Patients who have not received any prior treatment with infliximab or etanercept for ankylosing spondylitis.							
(ii) Patients who have received non-PBS-subsidised treatment with infliximab after 1 March 2004.							
(iii) Patients who have received non-PBS-subsidised treatment with etanercept after 1 July 2004.							
(iv) Patients who have received non-PBS-subsidised infliximab treatment prior to 1 March 2004.							
(v) Patients who have received non-PBS-subsidised etanercept treatment prior to 1 July 2004.							
To commence PBS-subsidised treatment with etanercept or infliximab, all patients who fall into groups (i)-(iii) must qualify under the first initial restriction for new patients (i.e. the patient must meet the criteria with respect to the indices of severity and prior treatment regimens specified in the 'new patient' restriction).							
To commence PBS-subsidised treatment with infliximab, all patients who fall into group (iv) must qualify under the first initial restriction for 'grandfather' patients (i.e. the patient must meet the criteria with respect to the indices of severity and prior treatment regimens specified in the 'grandfather' restriction).							

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ETANERCEPT—cont.

To commence PBS-subsidised treatment with etanercept, all patients who fall into group (v) must qualify under the first initial restriction for 'grandfather' patients (i.e. the patient must meet the criteria with respect to the indices of severity and prior treatment regimens specified in the 'grandfather' restriction).

Patients in groups (iv) and (v) can not be 'grandfathered' onto treatment with the alternate agent. If they wish to commence PBS-subsidised therapy with the alternate agent, they must qualify under the first initial restriction for new patients (i.e. the patient must meet the criteria with respect to the indices of severity and prior treatment regimens specified in the 'new patient' restriction).

(II) Patients who have received PBS-subsidised etanercept only.

(III) Patients who have received PBS-subsidised infliximab only.

(IV) Patients who have received PBS-subsidised etanercept and non-PBS-subsidised infliximab.

(V) Patients who have received PBS-subsidised infliximab and non-PBS-subsidised etanercept.

To receive PBS-subsidised treatment with etanercept or infliximab, the patients in groups (II)-(V) do not need to requalify with respect to the indices of severity or prior treatment requirements. They use the second initial restriction for either 'new' or 'grandfather' patients (i.e. whichever was applicable for their first course of PBS-subsidised treatment).

NOTE:

Patients who have responded to their most recent course of etanercept therapy as specified in the restriction may:

- continue on etanercept treatment under the 'grandfather' or new patient restriction, whichever is applicable [see 'Patients 'grandfathered' onto PBS-subsidised treatment with infliximab and etanercept']; or*
- swap to an initial course of infliximab treatment of up to 6 weeks under the 'grandfather' or new patient restriction, whichever is applicable [see 'Patients 'grandfathered' onto PBS-subsidised treatment with infliximab and etanercept'].*

Patients who have failed to respond to their most recent course of etanercept treatment may:

- trial a further course of initial therapy of up to 6 weeks with etanercept; or*
- swap to an initial course of infliximab treatment of up to 6 weeks.*

Patients who have responded to their most recent course of infliximab therapy as specified in the restriction may:

- continue on infliximab treatment under the 'grandfather' or new patient restriction, whichever is applicable [see 'Patients 'grandfathered' onto PBS-subsidised treatment with infliximab and etanercept']; or*
- swap to an initial course of etanercept treatment of up to 6 weeks under the 'grandfather' or new patient restriction, whichever is applicable [see 'Patients 'grandfathered' onto PBS-subsidised treatment with infliximab and etanercept'].*

Patients who have failed to respond to their most recent course of infliximab treatment may:

- trial a further course of initial therapy of up to 6 weeks with infliximab; or*
- swap to an initial course of etanercept treatment of up to 6 weeks.*

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ETANERCEPT—cont.**3. Swapping therapy.**

Once an authority for initial treatment with the first PBS-subsidised TNF- α antagonist is approved, patients may swap to the alternate TNF- α antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements. This also applies to patients who fail to achieve or sustain a response to the first PBS-subsidised TNF- α antagonist approved and who wish to trial a further course of treatment with the same TNF- α antagonist.

Patients may trial an alternate TNF- α antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF- α antagonist at the time of the application or not. However, they cannot swap to a particular TNF- α antagonist if they have failed to respond to prior treatment with that particular drug more than twice in the same cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate TNF- α antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF- α antagonist the patient is ceasing.

4. Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a TNF- α antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

For new patients, the first BASDAI result used to determine baseline must be provided while the patient is still receiving NSAID therapy. However, this is not required for any subsequent BASDAI results provided for these patients nor for patients who were 'grandfathered' onto TNF- α antagonist treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response.

5. Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF- α antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with at least 1 NSAID, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the BASDAI, ESR and/or CRP levels are measured.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ETANERCEPT—cont.							
Authority required							
Initial 1 (new patients)							
<i>Application for the first course of initial PBS-subsidised treatment with etanercept, by a rheumatologist, of adults with active ankylosing spondylitis who have radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, who have not received any PBS-subsidised treatment with either etanercept or infliximab;</i>							
AND							
<i>(a) have at least 2 of the following:</i>							
<i>(i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or</i>							
<i>(ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI) [for further information on the BASMI please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; or</i>							
<i>(iii) limitation of chest expansion relative to normal values for age and gender [for chest expansion normal values please refer to the Medicare Australia website at www.medicareaustralia.gov.au];</i>							
AND							
<i>(b) who have documented confirmation of HLA-B27 positive status;</i>							
AND							
<i>(c) who have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs) in combination with an appropriate exercise program for a total period of 3 months.</i>							
<i>The application must include details of the NSAIDs trialled, their doses and duration of treatment. If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reasons why a higher dose cannot be used.</i>							
<i>For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Medicare Australia website at www.medicareaustralia.gov.au.</i>							
<i>Patients who received treatment with etanercept prior to 1 July 2004 and wish to commence PBS-subsidised treatment with etanercept must qualify under the initial treatment restriction for 'grandfather' patients.</i>							
<i>The following initiation criteria indicate failure to achieve an adequate response and must be demonstrable in all patients at the time of the initial application:</i>							
<i>(a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale;</i>							
AND							
<i>(b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.</i>							

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ETANERCEPT—cont.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the patient is exempted from demonstrating an inadequate response to the above treatment regimen. Where appropriate, evidence to support a contraindication must be provided. If adverse events of a severity necessitating permanent withdrawal develop during the relevant period of use of 2 NSAIDs, the patient may be exempted from demonstrating an inadequate response to the above treatment regimen. For details of the adverse events, including the severity, which will be accepted for the purposes of administering this restriction, please refer to the Medicare Australia website at www.medicareaustralia.gov.au.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and*
 - (b) a completed TNF-alfa antagonist PBS Authority Application for Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes the following:*
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and*
 - (ii) a copy of the pathology report from an Approved Pathology Authority confirming the presence of HLA-B27; and*
 - (iii) a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)];*
 - (iv) a copy of the signed patient acknowledgment form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)]*
- Completion of this form declares that the patient understands and acknowledges that PBS-subsidised treatment with the TNF-alfa antagonists (etanercept or infliximab) for ankylosing spondylitis will cease if they do not demonstrate the response to treatment required to support continuation of PBS-subsidised treatment at any assessment where a response must be demonstrated.*

The assessment of the patient's response to the initial course of treatment must be made no earlier than 4 weeks from the commencement of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A maximum of 6 weeks of treatment with etanercept will be approved under this criterion.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ETANERCEPT—cont.**Authority required****Initial 2 (new patients)**

Application for an initial course of PBS-subsidised treatment with etanercept, by a rheumatologist, of adults with active ankylosing spondylitis who, in this treatment cycle, have received prior PBS-subsidised treatment with either etanercept or infliximab for this condition and have not failed PBS-subsidised therapy with etanercept more than once.

All applications for treatment must be accompanied by the results of the most recent course of TNF- α antagonist therapy within the timeframes specified below.

Where the most recent course of PBS-subsidised TNF- α antagonist treatment was approved under an initial treatment restriction for new patients, patients must have been assessed for response to that course no earlier than 4 weeks from the commencement of that course. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised TNF- α antagonist treatment was approved under the continuing treatment criteria for new patients, patients must have been assessed for response to that course, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessments to the previous course of TNF- α antagonist treatment are not submitted as detailed above, patients will be deemed to have failed therapy with that particular course of TNF- α antagonist.

Authority applications must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed TNF- α antagonist PBS Authority Application for Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website

(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website

(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].

A maximum of 6 weeks of treatment with etanercept will be approved under this criterion.

Authority required**Initial 2 ('grandfather' patients)**

Application for an initial course of PBS-subsidised treatment with etanercept, by a rheumatologist, of adults with active ankylosing spondylitis who have radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, who were 'grandfathered' onto PBS-subsidised treatment with etanercept or infliximab and who have not failed PBS-subsidised therapy with etanercept more than once.

Where the most recent course of PBS-subsidised TNF- α antagonist treatment was approved under an initial treatment restriction for 'grandfather' patients, patients must have been assessed for response to that course no earlier than 4 weeks from the commencement of that course. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised TNF- α antagonist treatment was approved under the continuing treatment criteria for 'grandfather' patients, patients must have been assessed for response to that course, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

continued ➤

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ETANERCEPT—cont.							
<i>If the response assessments to the previous course of TNF-alfa antagonist treatment are not submitted as detailed above, patients will be deemed to have failed therapy with that particular course of TNF-alfa antagonist.</i>							
<i>Authority applications must be made in writing and must include:</i>							
<i>(a) a completed authority prescription form; and</i>							
<i>(b) a completed TNF-alfa antagonist PBS Authority Application for Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website</i>							
<i>(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website</i>							
<i>(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].</i>							
<i>A maximum of 6 weeks of treatment with etanercept will be approved under this criterion.</i>							
8778B	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	3	..	..	*2830.09	28.60	Enbrel WY
NOTE:							
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>							

Authority required*Continuing (new patients)**Continuing PBS-subsidised treatment, by a rheumatologist, of adults with active ankylosing spondylitis who:*

- (a) have received 6 weeks or more of PBS-subsidised treatment with etanercept; and*
- (b) have demonstrated a response to treatment with etanercept; and*
- (c) whose most recent course of PBS-subsidised therapy in this treatment cycle was with etanercept.*

Response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or*
- (b) a CRP measurement no greater than 10 mg per L; or*
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.*

The same acute phase reactant measured at the relevant baseline must be measured in all subsequent continuing treatment applications.

The first application for continuing treatment following an initial treatment course must be made no earlier than 4 weeks from the commencement of the most recent initial treatment course with etanercept. This first authority application and a copy of the authority prescription may be faxed to Medicare Australia on (03) 6215 5640 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) in order to seek approval for a maximum of 4 weeks' supply. Medicare Australia will then contact the prescriber by telephone. The original document must then be posted to Medicare Australia with a second authority prescription for the balance of 24 weeks of treatment.

Second and subsequent applications for continuing treatment must be made in writing and should be posted to Medicare Australia no less than 2 weeks prior to the completion of the current treatment course.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ETANERCEPT—cont.

Written applications for authorisation must include:

(a) a completed authority prescription form; and

(b) a completed TNF- α antagonist PBS Authority Application for Continuing Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website

(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes the following:

(i) a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website

(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] including certification by the prescriber and the patient that the patient did not have access to their baseline BASDAI at the time of their continuing treatment assessment.

All measurements provided must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with etanercept will be authorised under this criterion.

Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.

Authority required

Initial 1 ('grandfather' patients)

Application for an initial PBS-subsidised course of continuing treatment with etanercept, by a rheumatologist, of adults with active ankylosing spondylitis who have radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, who were receiving treatment with etanercept prior to 1 July 2004;

AND

(a) are receiving treatment with etanercept at the time of application;

AND

(b) who have not received prior PBS-subsidised treatment with infliximab;

AND

(c) who have documented confirmation of HLA-B27 positive status;

AND

(d) whose Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score is less than or equal to 5 on a 0-10 scale;

AND

(e) who have:

(i) an ESR measurement no greater than 25 mm per hour; or

(ii) a CRP measurement no greater than 10 mg per L; or

(iii) an ESR or CRP measurement reduced by at least 20% from pre-treatment baseline.

The BASDAI assessment and ESR and/or CRP measurements must be no more than 1 month old at the time of application. The same acute phase reactant measured in the first application for PBS-subsidised treatment must be measured in all subsequent continuing treatment applications.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ETANERCEPT—cont.							
Authority applications must be made in writing and must include:							
(a) a completed authority prescription form; and							
(b) a completed TNF-alfa antagonist PBS Authority Application for Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes the following:							
(i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and							
(ii) a copy of the pathology report from an Approved Pathology Authority confirming the presence of HLA-B27; and							
(iii) a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)]]; and							
(iv) a copy of the signed patient acknowledgment form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)]].							
Completion of this form declares that the patient understands and acknowledges that PBS-subsidised treatment with the TNF-alfa antagonists (etanercept or infliximab) for ankylosing spondylitis will cease if they do not demonstrate the response to treatment required to support continuation of PBS-subsidised treatment at any assessment where a response must be demonstrated.							
The assessment of the patient's response to this initial course of therapy must be made within the 4 weeks prior to completion of the course in order to ensure continuity of treatment.							
Those patients ceasing treatment or swapping to an alternate agent and wishing to demonstrate a response to treatment, must be assessed no earlier than 4 weeks from the commencement of PBS-subsidised treatment. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.							
A maximum of 24 weeks of treatment with etanercept will be authorised under this criterion.							
Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.							
Patients may only qualify for PBS-subsidised treatment under this criterion once.							
<u>Authority required</u>							
Continuing ('grandfather' patients)							
Continuing PBS-subsidised treatment, by a rheumatologist, of adults with active ankylosing spondylitis who were 'grandfathered' onto PBS-subsidised treatment with a TNF-alfa antagonist and who:							
(a) have received 6 weeks or more of PBS-subsidised treatment with etanercept; and							
(b) have demonstrated a response to treatment with etanercept; and							
(c) whose most recent course of PBS-subsidised therapy in this treatment cycle was with etanercept.							

continued ➤

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ETANERCEPT—cont.							
Response to treatment is defined as a BASDAI score no more than 20% greater than the score included in the initial application for PBS-subsidised treatment; AND (a) an ESR measurement no greater than 25 mm per hour; or (b) a CRP measurement no greater than 10 mg per L; or (c) an ESR or CRP measurement reduced by at least 20% from pre-treatment baseline. The same acute phase reactant measured at the relevant baseline must be measured in all subsequent continuing treatment applications. The first application for continuing treatment following an initial treatment course must be made no earlier than 4 weeks from the commencement of the most recent initial treatment course with etanercept. This first authority application and a copy of the authority prescription may be faxed to Medicare Australia on (03) 6215 5640 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) in order to seek approval for a maximum of 4 weeks' supply. Medicare Australia will then contact the prescriber by telephone. The original document must then be posted to Medicare Australia with a second authority prescription for the balance of 24 weeks of treatment. Second and subsequent applications for continuing treatment must be made in writing and should be posted to Medicare Australia no less than 2 weeks prior to the completion of the current treatment course. In order to demonstrate a response to treatment where the patient is ceasing or swapping to an alternate agent, the assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased. Written applications for authorisation must include: (a) a completed authority prescription form; and (b) a completed TNF-α antagonist PBS Authority Application for Continuing Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes the following: (i) a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] including certification by the prescriber and the patient that the patient did not have access to their baseline BASDAI at the time of their continuing treatment assessment. All measurements provided must be no more than 1 month old at the time of application. A maximum of 24 weeks of treatment with etanercept will be authorised under this criterion. Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.							
8779C	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	5	..	*1888.31	28.60	Enbrel WY
NOTE: No applications for increased maximum quantities and/or repeats will be authorised.							

continued 

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ETANERCEPT—cont.**NOTE:**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the tumour necrosis factor (TNF) alfa antagonists (adalimumab, etanercept and infliximab) and the interleukin-1 inhibitor (anakinra) for adult patients with severe rheumatoid arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs (bDMARDs) at any 1 time.

1. Patients who have received no prior PBS-subsidised bDMARD treatment at 1 December 2004.

From 1 December 2004, the arrangements for prescribing the bDMARDs on the PBS have been amended to allow patients to commence a single cycle of bDMARD treatment that allows them to trial any number of bDMARDs without having to experience a disease flare when swapping between alternate bDMARDs. Within a single treatment cycle, patients may continue to receive long-term treatment with a bDMARD while they continue to show a response to therapy.

Once patients have either failed, or ceased to respond to, treatment with a maximum of 3 bDMARDs, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised bDMARD therapy before they are eligible to commence the next cycle. The 5-year period will be measured from the date the last prescription for PBS-subsidised bDMARD treatment was approved in the most recent cycle to the date of the application for initial treatment with a bDMARD under the new cycle.

Patients who have failed treatment with fewer than 3 bDMARDs within a particular treatment cycle, and where a period of less than 5 years duration has elapsed since the patient's previous course of PBS-subsidised bDMARD treatment in that cycle, may commence a further course of bDMARD treatment within that same treatment cycle.

Patients who have failed treatment with fewer than 3 bDMARDs within a particular treatment cycle, and who have had a break in PBS-subsidised therapy of 5 years or more, are eligible to commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

If patients fail to respond to a particular bDMARD within a single treatment cycle, they are not eligible to receive further PBS-subsidised treatment with that drug until they commence the next cycle.

2. Patients who have received PBS-subsidised TNF-alfa antagonist treatment prior to 1 December 2004.

Patients who commenced PBS-subsidised TNF-alfa antagonist therapy prior to 1 December 2004 are considered to be in their first cycle of bDMARD treatment on 1 December 2004.

Patients who have failed to respond to prior PBS-subsidised treatment with fewer than 3 TNF-alfa antagonists at the time the first application for treatment is made on or after 1 December 2004, will be subject to the same conditions applying to new patients detailed above.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum Recordable Value for Safety Net	Proprietary Name and Manufacturer
					for Max. Qty \$	\$	

ETANERCEPT—cont.

Therefore, patients who have failed:

(a) 1 TNF-alfa antagonist will be eligible to trial further PBS-subsidised treatment with a bDMARD they have not failed in their first treatment cycle, until they fail to demonstrate a response to no more than another 2 bDMARDs;

(b) 2 TNF-alfa antagonists will be eligible to trial further PBS-subsidised treatment with a bDMARD they have not failed in their first treatment cycle, until they fail to demonstrate a response to no more than 1 other bDMARD.

Patients who have failed PBS-subsidised treatment with 3 TNF-alfa antagonists prior to 1 December 2004 or at the first assessment required after this date, will be eligible to trial PBS-subsidised treatment with anakinra if they wish. However, if they fail to demonstrate a response to anakinra, they will not be able to trial any further PBS-subsidised bDMARD treatment until a minimum of 5 years has elapsed from the date that the prescription for the last course of anakinra therapy was approved. Arrangements to allow these patients to fail 4 bDMARDs will only be in place for the first treatment cycle. For subsequent cycles, patients will cease to be eligible to receive PBS-subsidised bDMARD treatment once they have failed to demonstrate a response to a maximum of 3 bDMARDs.

Any queries on these arrangements should be forwarded to Medicare Australia.

3. Information relevant to all patients.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) patients have received no prior PBS-subsidised bDMARD treatment and wish to commence such therapy; or

(ii) patients have received prior PBS-subsidised (initial or continuing) bDMARD therapy and wish to trial an alternate agent [further details are under 'Swapping therapy' below]; or

(iii) patients wish to re-commence treatment with a specific bDMARD following a break in PBS-subsidised therapy with that specific agent.

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted bDMARD supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that bDMARD.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that bDMARD.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed	Maximum	Proprietary Name and Manufacturer
					Price for Max. Qty \$	Recordable Value for Safety Net \$	
ETANERCEPT—cont.							
(c) Swapping therapy.							
Once an authority for initial treatment with the first PBS-subsidised bDMARD is approved, patients may swap to an alternate bDMARD without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-bDMARD therapy requirements. However, the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialed.							
Patients may swap to an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application or not.							
Patients may alternate between therapy with any bDMARD of their choice (1 at a time) providing:							
(i) they have not received PBS-subsidised treatment with that particular bDMARD previously; or							
(ii) they have demonstrated an adequate response to that particular bDMARD if they have previously trialed it on the PBS.							
Therefore, to maximise the choice of bDMARD patients may alternate between, it is important that patients are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.							
To avoid confusion, applications for patients who wish to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.							
(d) Baseline measurements to determine response.							
Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.							
To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to a reduction in the total number of active joints.							
(e) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.							
Patients who wish to trial a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with at least 1 non-biological DMARD, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.							

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of Qty Rpts Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ETANERCEPT—cont.					
Authority required					
<i>Application for initial PBS-subsidised treatment with etanercept, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:</i>					
<i>(a) have severe active rheumatoid arthritis; and</i>					
<i>(b) have received no prior PBS-subsidised treatment with a bDMARD for this condition in this treatment cycle; and</i>					
<i>(c) have failed to achieve an adequate response to the following treatments:</i>					
<i>(i) methotrexate at a dose of at least 20 mg weekly; and</i>					
<i>(ii) methotrexate (at a minimum dose of 7.5 mg weekly), in combination with 2 other non-biological disease modifying anti-rheumatic drugs (DMARDs), for a minimum of 3 months; and</i>					
<i>(iii) a minimum of 3 months' treatment with:</i>					
<i>— leflunomide alone; or</i>					
<i>— leflunomide in combination with methotrexate; or</i>					
<i>— cyclosporin.</i>					
<i>If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or intolerance of a severity necessitating permanent treatment withdrawal develops during the relevant period of use, the patient is exempted from demonstrating an inadequate response to that particular agent(s) only. Details of the contraindications or intolerance, including the degree of toxicity, must be provided at the time of application.</i>					
<i>The following initiation criteria indicate failure to achieve an adequate response and must be demonstrable in all patients at the time of the initial application:</i>					
<i>an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L;</i>					
<i>AND either</i>					
<i>(i) a total active joint count of at least 20 active (swollen and tender) joints; or</i>					
<i>(ii) at least 4 active joints from the following list of major joints:</i>					
<i>— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or</i>					
<i>— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).</i>					
<i>If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.</i>					
<i>The authority application must be made in writing and must include:</i>					
<i>(1) a completed authority prescription form; and</i>					
<i>(2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes details of the patient's ESR and CRP measurements and the patient's active joint count which must have been assessed no earlier than 1 month prior to the date of application; and</i>					
<i>(3) a copy of the signed patient acknowledgement form which may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm).</i>					

continued ➞

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ETANERCEPT—cont.

Completion of this form declares that the patient understands and acknowledges that, within a single treatment cycle, PBS-subsidised treatment with any biological DMARD will cease if they do not demonstrate the response to treatment required to support continuation of PBS-subsidised treatment at any assessment where a response must be demonstrated.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial etanercept after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Authority required

Application for an initial course of PBS-subsidised treatment with etanercept, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and*
- (b) have received prior PBS-subsidised bDMARD treatment for this condition in this treatment cycle and are eligible to receive further bDMARD therapy.*

Applications for patients who have received PBS-subsidised treatment with etanercept within this treatment cycle and who wish to re-commence therapy with this drug within this same cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised etanercept treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised etanercept treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised bDMARD therapy or, under this restriction, for patients who have received previous PBS-subsidised bDMARD therapy), patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised etanercept treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and*
- (2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].*

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of Qty Rpts Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer

ETANERCEPT—cont.

Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial etanercept after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Authority required

Initial treatment, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of patients aged 18 years or older with a documented history of severe active polyarticular course juvenile chronic arthritis with onset prior to the age of 18 years;

AND

(a) who have signed a patient agreement form indicating that they understand and acknowledge that PBS-subsidised treatment will cease if the predetermined response criteria do not support continuation of PBS-subsidised treatment;

AND

(b) who have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly;

AND

(c) who have failed to achieve an adequate response to methotrexate, in combination with 2 other disease modifying anti-rheumatic drugs (DMARDs), for a minimum of 3 months;

AND

(d) who have subsequently failed to achieve an adequate response following a minimum of 3 months' treatment with:

(i) leflunomide alone; or

(ii) leflunomide in combination with methotrexate; or

(iii) cyclosporin.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or intolerance of a severity necessitating permanent treatment withdrawal develops during the relevant period of use, the patient is exempted from demonstrating an inadequate response to the above treatment regimens. Details of the contraindication or intolerance, including the degree of toxicity, must be provided at the time of application.

The following criteria must be met in order to demonstrate failure to achieve an adequate response:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L;

AND either

(i) an active joint count of at least 20 active (swollen and tender) joints; or

(ii) at least 4 active joints from the following list:

— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

— shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

continued ➞

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer		
ETANERCEPT—cont.									
<i>If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.</i>									
<i>The authority application must be in writing and must include sufficient information to determine the patient's eligibility according to the above criteria. The date of joint assessment must be provided.</i>									
<i>Where fewer than 3 repeats are requested at the time of the initial authority application, authority approvals for sufficient repeats to complete a maximum of 4 months of treatment may be requested by telephone. Under no circumstances will telephone approvals be granted for initial or continuing authority applications, or for treatment that would otherwise extend the initial treatment period beyond 4 months.</i>									
<i>The assessment of the patient's response to the initial course of treatment should be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. Applications for continuing treatment with etanercept should be made prior to the completion of 16 weeks of treatment to ensure continuity for those patients who meet the criteria.</i>									
8637N	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	3	..	*1888.31	28.60	Enbrel		WY
8861J	Injection set containing 4 vials powder for injection 50 mg and 4 pre-filled syringes solvent 1 mL	1	3	..	1888.31	28.60	Enbrel		WY

NOTE:

No applications for increased maximum quantities and/or repeats will be authorised.

Authority required

Initial PBS-subsidised supply for continuing treatment with etanercept, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) were receiving treatment with etanercept prior to 1 March 2005; and
- (c) failed to qualify for PBS-subsidised therapy after 1 August 2003 due to testing negative for rheumatoid factor; and
- (d) have demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with etanercept.

Medical practitioners who wish to apply for authority to prescribe a bDMARD for patients who commenced treatment with etanercept prior to 1 March 2005 and who have not demonstrated a response to treatment should contact Medicare Australia on 1800 005 750.

The authority application must be in writing and must include sufficient information to determine the patient's eligibility. The date of assessment must be provided.

Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial etanercept after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed	Maximum	Proprietary Name and Manufacturer
					Price for Max. Qty \$	Recordable Value for Safety Net \$	

ETANERCEPT—cont.

Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Authority required

Continuing PBS-subsidised treatment with etanercept, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have severe active rheumatoid arthritis; and*
- (b) who have demonstrated an adequate response to treatment with etanercept; and*
- (c) whose most recent course of PBS-subsidised bDMARD treatment in this treatment cycle was with etanercept.*

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or*
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:*

- elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or*
- shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).*

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and*
- (2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].*

All applications for continuing treatment with etanercept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with etanercept, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial etanercept after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

ETANERCEPT—cont.

Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Authority required

Initial PBS-subsidised supply for continuing treatment, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of patients aged 18 years or older with a documented history of severe active polyarticular course juvenile chronic arthritis with onset prior to the age of 18 years, and who were receiving treatment with etanercept prior to 1 December 2002;

AND

(a) who have signed a patient agreement form indicating that they understand and acknowledge that PBS-subsidised treatment will cease if the predetermined response criteria do not support continuation of PBS-subsidised treatment;

AND

(b) who have demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with etanercept.

The authority application must be in writing and must include sufficient information to determine the patient's eligibility. The date of assessment of the patient must be provided.

Authority required

Continuing PBS-subsidised treatment, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of patients aged 18 years or older with a documented history of severe active polyarticular course juvenile chronic arthritis with onset prior to the age of 18 years, who, at the time of application, demonstrate an adequate response to treatment with etanercept as manifested by:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND 1 or more of the following:

(i) an active joint count of fewer than 10 active (swollen and tender) joints; or

(ii) a reduction in the active (swollen and tender) joint count by at least 50% from baseline;

or

(iii) a reduction in the number of the following active joints, from at least 4, by at least 50%:

— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

— shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All authority applications for continuing treatment with etanercept must be in writing and must include sufficient information to determine the patient's response according to the above criteria. The date of assessment of the patient must be provided.

Patients who fail to demonstrate an adequate response, as specified in the criteria for continuing treatment with etanercept, will not be eligible to recommence treatment with etanercept within 12 months of the date on which treatment was ceased.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ETANERCEPT—cont.								
<i>Where re-treatment with etanercept after a break in PBS-subsidised treatment with the drug is being sought, the reason for and date of cessation of the previous treatment course with etanercept must be included in the application.</i>								
8638P	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	5	..	*1888.31	28.60	Enbrel	WY
8862K	Injection set containing 4 vials powder for injection 50 mg and 4 pre-filled syringes solvent 1 mL	1	5	..	1888.31	28.60	Enbrel	WY
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
EVEROLIMUS								
CAUTION:								
<i>Careful monitoring of patients is mandatory.</i>								
Authority required								
<i>Maintenance therapy, following initiation and stabilisation of treatment with everolimus and where therapy remains under the supervision and direction of the transplant unit reviewing that patient, of patients with:</i>								
<i>(a) renal transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application;</i>								
<i>(b) cardiac transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application.</i>								
8840G	Tablet 0.25 mg	60	3	..	289.75	28.60	Certican	NV
8841H	Tablet 0.5 mg	60	3	..	560.11	28.60	Certican	NV
8842J	Tablet 0.75 mg	120	3	..	*1670.83	28.60	Certican	NV
LEFLUNOMIDE								
CAUTION:								
<i>Leflunomide is a category X drug and must not be given to pregnant women. Pregnancy should be avoided for two years after cessation of therapy, unless special wash-out procedures are carried out.</i>								
Authority required								
<i>Initiation by consultant physicians for the treatment of severe active rheumatoid arthritis in patients for whom other disease modifying anti-rheumatic drugs (including methotrexate) are inappropriate and/or ineffective.</i>								
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8373Q	Pack containing 3 tablets leflunomide 100 mg and 30 tablets leflunomide 20 mg	‡1	..	..	250.11	28.60	Arava	AV

continued ➡

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
LEFLUNOMIDE—cont.								
Authority required								
<i>Initiation by consultant physicians for the treatment of severe active rheumatoid arthritis in patients for whom other disease modifying anti-rheumatic drugs (including methotrexate) are inappropriate and/or ineffective;</i>								
<i>Ongoing leflunomide therapy for severe active rheumatoid arthritis in patients for whom other disease modifying anti-rheumatic drugs (including methotrexate) are inappropriate and/or ineffective.</i>								
8374R	Tablet 10 mg	30	5	..	108.78	28.60	^a Arabloc	HP
				^B 1.22	110.00	28.60	^a Arava	AV
8375T	Tablet 20 mg	30	5	..	163.15	28.60	^a Arabloc	HP
				^B 1.22	164.37	28.60	^a Arava	AV
MYCOPHENOLATE MOFETIL								
CAUTION:								
<i>Careful monitoring of patients is mandatory.</i>								
Authority required								
<i>Maintenance therapy, following initiation and stabilisation of treatment with mycophenolate mofetil and where therapy remains under the supervision and direction of the transplant unit reviewing that patient, of patients with:</i>								
<i>(a) renal transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application;</i>								
<i>(b) cardiac transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application.</i>								
8649F	Capsule 250 mg	300	3	..	646.89	28.60	CellCept	RO
8650G	Tablet 500 mg	150	3	..	646.89	28.60	CellCept	RO
8651H	Powder for oral suspension 1 g per 5 mL, 165 mL	‡1	3	..	#296.85	28.60	CellCept	RO
MYCOPHENOLATE SODIUM								
CAUTION:								
<i>Careful monitoring of patients is mandatory.</i>								
Authority required								
<i>Maintenance therapy, following initiation and stabilisation of treatment with mycophenolate sodium and where therapy remains under the supervision and direction of the transplant unit reviewing that patient, of patients with renal transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application.</i>								
8652J	Tablet (enteric coated) 180 mg (mycophenolic acid)	120	3	..	269.73	28.60	Myfortic	NV
8653K	Tablet (enteric coated) 360 mg (mycophenolic acid)	120	3	..	518.47	28.60	Myfortic	NV

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
SIROLIMUS								
CAUTION:								
<i>Careful monitoring of patients is mandatory.</i>								
Authority required								
<i>Maintenance therapy, following initiation and stabilisation of treatment with sirolimus and where therapy remains under the supervision and direction of the transplant unit reviewing that patient, of patients with renal transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application.</i>								
8724E	Tablet 1 mg	100	3	..	778.97	28.60	Rapamune	WY
8833X	Tablet 2 mg	100	3	..	1553.20	28.60	Rapamune	WY
8725F	Oral solution 1 mg per mL, 60 mL	£1	3	..	469.42	28.60	Rapamune	WY
TACROLIMUS								
CAUTION:								
<i>Careful monitoring of patients is mandatory.</i>								
Authority required								
<i>Maintenance therapy, following initiation and stabilisation of treatment with tacrolimus and where therapy remains under the supervision and direction of the transplant unit reviewing that patient, of patients with:</i>								
<i>(a) liver transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application;</i>								
<i>(b) renal transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application.</i>								
8646C	Capsule 500 micrograms	100	3	..	230.90	28.60	Prograf	JC
8647D	Capsule 1 mg	100	3	..	439.05	28.60	Prograf	JC
8648E	Capsule 5 mg	50	3	..	1087.14	28.60	Prograf	JC
• Other immunosuppressive agents								
AZATHIOPRINE								
2688L	Tablet 25 mg	100	2	..	45.29	28.60	^a Azahexal	HX
				^B 1.57	46.86	28.60	^a Imuran	GK
2687K	Tablet 50 mg	100	2	..	73.75	28.60	^a Azahexal	HX
							^a Azamun	DP
							^a Azapin	AW
							^a GenRx	GX
							Azathioprine	
				^B 1.50	75.25	28.60	^a Thioprine	AF
							^a Imuran	GK
METHOTREXATE								
1622J	Tablet 2.5 mg	30	5	..	11.93	12.90	^a Methoblastin	PH
				^B 0.08	12.01	12.90	^a MX	
							^a Ledertrexate	WY
1623K	Tablet 10 mg	50	2	..	47.10	28.60	Methoblastin	PH

MUSCULO-SKELETAL SYSTEM

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS								
Antiinflammatory and antirheumatic products, non-steroids								
• Acetic acid derivatives and related substances								
DICLOFENAC SODIUM								
1302M	Suppository 100 mg	40	3	..	*23.03	24.00	Voltaren 100	NV
DICLOFENAC SODIUM								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
1299J	Tablet 25 mg (enteric coated)	100	3	..	*13.51	14.48	^a Chem mart Diclofenac	CH
							^a Clonac 25	AW
							^a Diclofenac-BC	BG
							^a Diclohexal	HX
							^a Dinac	DP
							^a GenRx	GX
							Diclofenac	
							^a Terry White Chemists	TW
							Diclofenac	
				..	13.51	14.48	^a Fenac 25	AF
				^B 3.30	*16.81	14.48	^a Voltaren 25	NV
1300K	Tablet 50 mg (enteric coated)	50	3	..	10.84	11.81	^a Chem mart Diclofenac	CH
							^a Clonac 50	AW
							^a Diclohexal	HX
							^a Dinac	DP
							^a Fenac	AF
							^a GenRx	GX
							Diclofenac	
							^a Terry White Chemists	TW
							Diclofenac	
				^B 3.21	14.05	11.81	^a Voltaren 50	NV
INDOMETHACIN								
2757D	Suppository 100 mg	40	3	..	*20.63	21.60	Indocid	MK
INDOMETHACIN								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
2454E	Capsule 25 mg	100	3	..	*8.73	9.70	^a Arthrexin	AF
				^B 3.06	*11.79	9.70	^a Indocid	MK

MUSCULO-SKELETAL SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
SULINDAC								
Restricted benefit								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
2047R	Tablet 100 mg	100	3	..	*14.57	15.54	Aclin	AF
2048T	Tablet 200 mg	50	3	..	13.51	14.48	Aclin 200	AF
• Oxicams								
MELOXICAM								
NOTE:								
<i>The use of meloxicam for the treatment of the following conditions is not subsidised through the PBS:</i>								
<i>(a) acute pain;</i>								
<i>(b) soft tissue injury;</i>								
<i>(c) arthrosis without an inflammatory component.</i>								
Restricted benefit								
<i>Symptomatic treatment of osteoarthritis.</i>								
8561N	Tablet 7.5 mg	30	3	..	22.83	23.80	^a Movalis 7.5	AW
				^B 1.85	24.68	23.80	^a Mobic	BY
8562P	Tablet 15 mg	30	3	..	31.13	28.60	^a Movalis 15	AW
				^B 1.85	32.98	28.60	^a Mobic	BY
PIROXICAM								
Restricted benefit								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component.</i>								
1895R	Dispersible tablet 10 mg	50	3	..	12.77	13.74	^a GenRx Piroxicam Dispersible	GX
							^a Mobilis D-10	AF
							^a Pirohexal-D	HX
				^B 2.75	15.52	13.74	^a Feldene-D	PF
1896T	Dispersible tablet 20 mg	25	3	..	12.36	13.33	^a Chem mart Piroxicam Dispersible	CH
							^a GenRx Piroxicam Dispersible	GX
							^a Mobilis D-20	AF
							^a Pirohexal-D	HX
							^a Terry White Chemists Piroxicam Dispersible	TW
				^B 2.73	15.09	13.33	^a Feldene-D	PF

continued ➡

MUSCULO-SKELETAL SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PIROXICAM—cont.								
1897W	Capsule 10 mg	50	3	..	12.77	13.74	^a Chem mart Piroxicam	CH
							^a GenRx Piroxicam	GX
							^a Mobilis 10	AF
							^a Terry White Chemists Piroxicam	TW
				^B 2.75	15.52	13.74	^a Feldene	PF
1898X	Capsule 20 mg	25	3	..	12.36	13.33	^a Chem mart Piroxicam	CH
							^a GenRx Piroxicam	GX
							^a Mobilis 20	AF
							^a Terry White Chemists Piroxicam	TW
				^B 2.73	15.09	13.33	^a Feldene	PF
• Propionic acid derivatives								
IBUPROFEN								
3192B	Tablets 400 mg, 20	‡1	..	..	7.65	8.62	Brufen	AB
IBUPROFEN								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
3198H	Tablet 200 mg	100	3	..	*9.11	10.08	Rafen 200	AF
3190X	Tablet 400 mg	100	3	..	12.49	13.46	Brufen	AB
KETOPROFEN								
1588N	Suppository 100 mg	40	3	..	*21.39	22.36	Orudis	AV
KETOPROFEN								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component.</i>								
1590Q	Capsule 200 mg (sustained release)	28	3	..	15.34	16.31	^a Oruvail SR	HP
				^B 2.01	17.35	16.31	^a Orudis SR 200	AV
NAPROXEN								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
1674D	Tablet 250 mg	100	3	..	*14.37	15.34	^a Inza 250	AF
				^B 3.10	*17.47	15.34	^a Naprosyn	RO
1659H	Tablet 500 mg	50	3	..	13.30	14.27	^a Inza 500	AF
				^B 1.80	15.10	14.27	^a Naprosyn	RO

continued ↻

MUSCULO-SKELETAL SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
NAPROXEN—cont.								
1614Y	Tablet 750 mg (sustained release)	28	3	.. B1.68	12.60 14.28	13.57 ^a 13.57 ^a	Proxen SR 750 Naprosyn SR750	MD RO
1615B	Tablet 1 g (sustained release)	28	3	.. B1.77	15.22 16.99	16.19 ^a 16.19 ^a	Proxen SR 1000 Naprosyn SR1000	MD RO
NAPROXEN SODIUM								
Restricted benefit								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
1795L	Tablet 550 mg	50	3	.. B3.00	13.56 16.56	14.53 ^a 14.53 ^a	Crysanal Anaprox 550	MD RO
NOTE: <i>Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.</i>								
TIAPROFENIC ACID								
CAUTION: <i>Cystitis and other urinary disorders have been reported with this drug.</i>								
NOTE: <i>The recommended maximum dose is 600 mg per day.</i>								
Restricted benefit								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component.</i>								
2103Q	Tablet 300 mg	60	3	..	15.77	16.74	Surgam	AV
• Fenamates								
MEFENAMIC ACID								
Restricted benefit								
<i>Dysmenorrhoea; Menorrhagia.</i>								
1824B	Capsule 250 mg	50	2	..	16.36	17.33 ^a 17.33 ^a	Mefic Ponstan	PC PD
• Coxibs								
CELECOXIB								
NOTE: <i>The use of celecoxib for the treatment of the following conditions is not subsidised through the PBS:</i> <i>(a) acute pain;</i> <i>(b) soft tissue injury;</i> <i>(c) arthrosis without an inflammatory component.</i>								
Restricted benefit								
<i>Symptomatic treatment of osteoarthritis; Symptomatic treatment of rheumatoid arthritis.</i>								
8439E	Capsule 100 mg	60	3	..	30.35	28.60	Celebrex	PH
8440F	Capsule 200 mg	30	3	..	30.35	28.60	Celebrex	PH

MUSCULO-SKELETAL SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MUSCLE RELAXANTS								
Muscle relaxants, centrally acting agents								
• Other centrally acting agents								
BACLOFEN								
2729P	Tablet 10 mg	100	5	..	38.55	28.60	a Baclo a Baclohexal a Chem mart Baclofen a Clofen 10 a GenRx Baclofen a Stelax 10 a Terry White Chemists Baclofen	DP HX CH AF GX AW TW NV
				B3.00	41.55	28.60	a Lioresal 10	NV
2730Q	Tablet 25 mg	100	5	..	78.69	28.60	a Baclo a Baclohexal a Chem mart Baclofen a Clofen 25 a GenRx Baclofen a Stelax 25 a Terry White Chemists Baclofen	DP HX CH AF GX AW TW NV
				B3.01	81.70	28.60	a Lioresal 25	NV
Muscle relaxants, directly acting agents								
• Dantrolene and derivatives								
DANTROLENE SODIUM								
<u>Restricted benefit</u>								
<i>Treatment of chronic spasticity.</i>								
1779P	Capsule 25 mg	100	2	..	35.00	28.60	Dantrium	PU
1780Q	Capsule 50 mg	100	2	..	50.95	28.60	Dantrium	PU

MUSCULO-SKELETAL SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTIGOUT PREPARATIONS								
Antigout preparations								
• <i>Preparations inhibiting uric acid production</i>								
ALLOPURINOL								
NOTE:								
The dose should be adjusted in accordance with renal function.								
2600W	Tablet 100 mg	200	2	..	*13.67	14.64	^a Progout 100	AF
				..	13.67	14.64	^a Allohexal	HX
							^a Allosig	FM
							^a Chem mart	CH
							Allopurinol	
							^a GenRx	GX
							Allopurinol	
							^a Terry White	TW
							Chemists	
							Allopurinol	
				^B 2.43	16.10	14.64	^a Zyloprim	SI
2604C	Tablet 300 mg	60	2	..	10.03	11.00	^a Allohexal	HX
							^a Allosig	FM
							^a Chem mart	CH
							Allopurinol	
							^a GenRx	GX
							Allopurinol	
							^a Progout 300	AF
							^a Terry White	TW
							Chemists	
							Allopurinol	
				^B 2.28	12.31	11.00	^a Zyloprim	SI
• <i>Preparations increasing uric acid excretion</i>								
PROBENECID								
1940D	Tablet 500 mg	100	5	..	70.68	28.60	Pro-Cid	PL
• <i>Preparations with no effect on uric acid metabolism</i>								
COLCHICINE								
1227N	Tablet 500 micrograms	100	2	..	9.08	10.05	^a Lengout	LN
				^B 0.93	10.01	10.05	^a Colgout	AS

MUSCULO-SKELETAL SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DRUGS FOR TREATMENT OF BONE DISEASES								
Drugs affecting bone structure and mineralization								
• Bisphosphonates								
ALENDRONATE SODIUM								
Authority required								
<i>Initial treatment for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be included in the authority application.</i>								
<i>A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body;</i>								
<i>Continuing treatment for established osteoporosis in patients with fracture due to minimal trauma, where the patient has previously been issued with an authority prescription for this drug.</i>								
8511Y	Tablet equivalent to 70 mg alendronic acid	4	5	..	54.46	28.60	^a Alendro Once Weekly	AW
				^b 0.24	54.70	28.60	^a Fosamax Once Weekly	MK
Authority required								
<i>Symptomatic Paget's disease of bone.</i>								
8090T	Tablet equivalent to 40 mg alendronic acid	30	5	..	164.01	28.60	Fosamax 40 mg	MK
DISODIUM ETIDRONATE								
Authority required								
<i>Symptomatic Paget's disease of bone when calcitonin has been found to be unsatisfactory due to:</i>								
<i>(a) lack of efficacy; or</i>								
<i>(b) unacceptable side effects;</i>								
<i>Heterotopic ossification.</i>								
2920Q	Tablet 200 mg	60	5	..	117.14	28.60	Didronel	PU
DISODIUM PAMIDRONATE								
Authority required								
<i>Symptomatic Paget's disease of bone.</i>								
8461H	Concentrated injection 15 mg in 5 mL	4	..	..	*275.59	28.60	Pamisol	MX
8208B	Injection set containing 4 vials powder for I.V. infusion 15 mg and 4 ampoules solvent 5 mL	1	..	..	275.61	28.60	Aredia 15 mg	NV
8462J	Concentrated injection 30 mg in 10 mL	2	..	..	*275.61	28.60	Pamisol	MX
8209C	Injection set containing 2 vials powder for I.V. infusion 30 mg and 2 ampoules solvent 10 mL	1	..	..	275.61	28.60	Aredia 30 mg	NV
8463K	Concentrated injection 60 mg in 10 mL	1	..	..	275.61	28.60	Pamisol	MX

MUSCULO-SKELETAL SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
RISEDRONATE SODIUM								
Authority required								
<i>Initial treatment for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be included in the authority application.</i>								
<i>A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body;</i>								
<i>Continuing treatment for established osteoporosis in patients with fracture due to minimal trauma, where the patient has previously been issued with an authority prescription for this drug.</i>								
8481J	Tablet 5 mg	28	5	..	54.46	28.60	Actonel	AV
8621R	Tablet 35 mg	4	5	..	54.46	28.60	Actonel Once-a-Week	AV
Authority required								
<i>Symptomatic Paget's disease of bone.</i>								
8482K	Tablet 30 mg	28	1	..	312.31	28.60	Actonel	AV
SODIUM CLODRONATE TETRAHYDRATE								
Restricted benefit								
<i>Maintenance treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy;</i>								
<i>Multiple myeloma;</i>								
<i>Bone metastases from breast cancer.</i>								
8132B	Capsule equivalent to 400 mg sodium clodronate	100	2	..	342.75	28.60	Bonefos	SC
8265B	Tablet equivalent to 800 mg sodium clodronate	60	2	..	393.75	28.60	Bonefos 800 mg	SC
TILUDRONATE DISODIUM								
Authority required								
<i>Symptomatic Paget's disease of bone.</i>								
8267D	Tablet equivalent to 200 mg tiludronic acid	56	2	..	312.31	28.60	Skelid	MX

MUSCULO-SKELETAL SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Bisphosphonates and calcium, sequential preparations DISODIUM ETIDRONATE and CALCIUM CARBONATE Authority required <i>Initial treatment for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be included in the authority application.</i> <i>A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body;</i> <i>Continuing treatment for established osteoporosis in patients with fracture due to minimal trauma, where the patient has previously been issued with an authority prescription for this drug.</i>								
8056B	Pack containing 28 tablets disodium etidronate 200 mg and 76 tablets calcium carbonate 1.25 g (equivalent to 500 mg calcium)	1	1	..	71.17	28.60	Didrocal	PU
NOTE: No applications for increased maximum quantities and/or repeats will be authorised. • Other drugs affecting bone structure and mineralization CALCITRIOL Authority required <i>Hypocalcaemia due to renal disease;</i> <i>Hypoparathyroidism;</i> <i>Hypophosphataemic rickets;</i> <i>Vitamin D-resistant rickets;</i> <i>Initial treatment for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be included in the authority application.</i> <i>A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body;</i> <i>Continuing treatment for established osteoporosis in patients with fracture due to minimal trauma, where the patient has previously been issued with an authority prescription for this drug.</i>								
2502Q	Capsule 0.25 microgram	100	3	..	53.74	28.60	^a Calcitriol-DP ^a Citrihexal ^a GenRx Calcitriol ^a Kosteo ^a Rocaltrol ^a Sitriol	DP HX GX AW RO AF

MUSCULO-SKELETAL SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. No.of			Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
		Qty	Rpts	Premium				
<u>RALOXIFENE HYDROCHLORIDE</u>								
<u>Authority required</u>								
<i>Initial treatment for established post-menopausal osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be included in the authority application. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body;</i>								
<i>Continuing treatment for established post-menopausal osteoporosis in patients with fracture due to minimal trauma, where the patient has previously been issued with an authority prescription for this drug.</i>								
8363E	Tablet 60 mg	28	5	..	59.00	28.60	Evista	LY

NERVOUS SYSTEM

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
ANALGESICS							
Opioids							
• Natural opium alkaloids							
CODEINE PHOSPHATE							
1214X	Tablet 30 mg	20	..	..	10.53	11.50	FM
CODEINE PHOSPHATE with PARACETAMOL							
1215Y	Tablet 30 mg-500 mg	20	..	..	6.94	7.91	^a Codalgin Forte FM ^a Codapane Forte AL ^a Dolaforte CO ^a Dymadon Forte GK ^a Prodeine Forte DK ^a Panadeine Forte SW
				^B 1.89	8.83	7.91	
NOTE: Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of codeine phosphate with paracetamol below.							
CODEINE PHOSPHATE with PARACETAMOL							
Authority required							
Severe disabling pain not responding to non-narcotic analgesics.							
NOTE: Each authority approval will be limited to no more than 240 tablets per month for no more than 6 months.							
8785J	Tablet 30 mg-500 mg	60	..	..	*11.32	12.29	^a Codalgin Forte FM ^a Codapane Forte AL ^a Dolaforte CO ^a Dymadon Forte GK ^a Prodeine Forte DK ^a Panadeine Forte SW
					^B 5.67	*16.99	12.29
HYDROMORPHONE HYDROCHLORIDE							
CAUTION: The risk of drug dependence is high.							
8420E	Injection 2 mg in 1 mL	5	..	..	11.65	12.62	Dilaudid AB
8421F	Injection 10 mg in 1 mL	5	..	..	17.13	18.10	Dilaudid-HP AB
8422G	Injection 50 mg in 5 mL	5	..	..	45.86	28.60	Dilaudid-HP AB
8423H	Injection 500 mg in 50 mL	1	..	..	80.02	28.60	Dilaudid-HP AB

continued ➡

NERVOUS SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
HYDROMORPHONE HYDROCHLORIDE—cont.								
CAUTION: <i>The risk of drug dependence is high.</i>								
Restricted benefit <i>Severe disabling pain not responding to non-narcotic analgesics.</i>								
NOTE: <i>Authorities for increased maximum quantities and/or repeats will be granted only for:</i>								
<i>(i) severe disabling pain associated with proven malignant neoplasia; or</i>								
<i>(ii) chronic severe disabling pain not responding to non-narcotic analgesics where treatment was initiated in a hospital (in-patient or out-patient) prior to 1 April 2005. This criterion will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006; or</i>								
<i>(iii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or</i>								
<i>(iv) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or</i>								
<i>(v) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.</i>								
8541M	Tablet 2 mg	20	..	..	12.40	13.37	Dilaudid	AB
8542N	Tablet 4 mg	20	..	..	16.07	17.04	Dilaudid	AB
8543P	Tablet 8 mg	20	..	..	22.55	23.52	Dilaudid	AB
8424J	Oral liquid 1 mg per mL, 473 mL	1	..	..	24.37	25.34	Dilaudid	AB

MORPHINE HYDROCHLORIDE**CAUTION:***The risk of drug dependence is high.***Restricted benefit***Severe disabling pain not responding to non-narcotic analgesics.***NOTE:***Authorities for increased maximum quantities and/or repeats will be granted only for:**(i) severe disabling pain associated with proven malignant neoplasia; or**(ii) chronic severe disabling pain not responding to non-narcotic analgesics where treatment was initiated in a hospital (in-patient or out-patient) prior to 1 April 2005. This criterion will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006; or**(iii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or**(iv) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or*

continued ➤

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MORPHINE HYDROCHLORIDE—cont.								
<i>(v) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.</i>								
2122Q	Oral solution 2 mg per mL, 200 mL	1	..	..	17.27	18.24	Ordine 2	MF
2123R	Oral solution 5 mg per mL, 200 mL	1	..	..	20.14	21.11	Ordine 5	MF
2124T	Oral solution 10 mg per mL, 200 mL	1	..	..	24.70	25.67	Ordine 10	MF

MORPHINE SULFATE

CAUTION:

The risk of drug dependence is high.

1644M	Injection 10 mg in 1 mL	5	..	..	11.61	12.58	MX SI	
1645N	Injection 15 mg in 1 mL	5	..	..	11.94	12.91	MX SI	
1647Q	Injection 30 mg in 1 mL	5	..	..	13.92	14.89	MX	

MORPHINE SULFATE

CAUTION:

The risk of drug dependence is high.

Restricted benefit*Severe disabling pain due to cancer not responding to non-narcotic analgesics.*

8669G	Tablet 10 mg	20	..	..	11.49	12.46	Sevredol	MF
8670H	Tablet 20 mg	20	..	..	12.71	13.68	Sevredol	MF

Restricted benefit*Severe disabling pain not responding to non-narcotic analgesics.***NOTE:***Authorities for increased maximum quantities and/or repeats will be granted only for:*

- (i) severe disabling pain associated with proven malignant neoplasia; or*
- (ii) chronic severe disabling pain not responding to non-narcotic analgesics where treatment was initiated in a hospital (in-patient or out-patient) prior to 1 April 2005. This criterion will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006; or*
- (iii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or*
- (iv) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or*

continued ➞

NERVOUS SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MORPHINE SULFATE—cont.								
<i>(v) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.</i>								
1646P	Tablet 30 mg	20	..	..	13.12	14.09	Anamorph	FM
Restricted benefit								
<i>Chronic severe disabling pain not responding to non-narcotic analgesics.</i>								
NOTE:								
<i>Authorities for increased maximum quantities and/or repeats will be granted only for:</i>								
<i>(i) chronic severe disabling pain associated with proven malignant neoplasia; or</i>								
<i>(ii) chronic severe disabling pain not responding to non-narcotic analgesics where treatment was initiated in a hospital (in-patient or out-patient) prior to 1 April 2005. This criterion will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006; or</i>								
<i>(iii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or</i>								
<i>(iv) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or</i>								
<i>(v) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.</i>								
8035X	Tablet 5 mg (controlled release)	20	..	..	14.11	15.08	MS Contin	MF
1653B	Tablet 10 mg (controlled release)	20	..	..	15.63	16.60	MS Contin	MF
8489T	Tablet 15 mg (controlled release)	20	..	..	19.37	20.34	MS Contin	MF
1654C	Tablet 30 mg (controlled release)	20	..	..	28.45	28.60	MS Contin	MF
1655D	Tablet 60 mg (controlled release)	20	..	..	44.46	28.60	MS Contin	MF
1656E	Tablet 100 mg (controlled release)	20	..	..	61.24	28.60	MS Contin	MF
8349K	Capsule 10 mg (containing sustained release pellets)	20	..	..	15.63	16.60	Kapanol	GK
2839K	Capsule 20 mg (containing sustained release pellets)	20	..	..	20.70	21.67	Kapanol	GK
8491X	Capsule 30 mg (controlled release)	10	..	..	19.37	20.34	MS Mono	MF
2840L	Capsule 50 mg (containing sustained release pellets)	20	..	..	36.07	28.60	Kapanol	GK

continued ➡

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MORPHINE SULFATE—cont.								
8492Y	Capsule 60 mg (controlled release)	10	..	..	28.45	28.60	MS Mono	MF
8493B	Capsule 90 mg (controlled release)	10	..	..	34.50	28.60	MS Mono	MF
2841M	Capsule 100 mg (containing sustained release pellets)	20	..	..	61.24	28.60	Kapanol	GK
8494C	Capsule 120 mg (controlled release)	10	..	..	44.46	28.60	MS Mono	MF
8490W	Sachet containing controlled release granules for oral suspension, 20 mg per sachet	20	..	..	20.70	21.67	MS Contin Suspension 20 mg	MF
8146R	Sachet containing controlled release granules for oral suspension, 30 mg per sachet	20	..	..	28.45	28.60	MS Contin Suspension 30 mg	MF
8305D	Sachet containing controlled release granules for oral suspension, 60 mg per sachet	20	..	..	44.46	28.60	MS Contin Suspension 60 mg	MF
8306E	Sachet containing controlled release granules for oral suspension, 100 mg per sachet	20	..	..	61.24	28.60	MS Contin Suspension 100 mg	MF

Authority required*Chronic severe disabling pain due to cancer.*

8453X	Tablet 200 mg (controlled release)	20	..	..	106.36	28.60	MS Contin	MF
8454Y	Sachet containing controlled release granules for oral suspension, 200 mg per sachet	20	..	..	106.36	28.60	MS Contin Suspension 200 mg	MF

MORPHINE TARTRATE

CAUTION:

The risk of drug dependence is high.

1607N	Injection 120 mg in 1.5 mL	5	..	..	21.30	22.27	MX	

OXYCODONE

CAUTION:*The risk of drug dependence is high.***Restricted benefit***Severe disabling pain not responding to non-narcotic analgesics.***NOTE:***Authorities for increased maximum quantities and/or repeats will be granted only for:**(i) severe disabling pain associated with proven malignant neoplasia; or*

continued ➞

NERVOUS SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
OXYCODONE—cont.							
	(ii) <i>chronic severe disabling pain not responding to non-narcotic analgesics where treatment was initiated in a hospital (in-patient or out-patient) prior to 1 April 2005. This criterion will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006; or</i>						
	(iii) <i>chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or</i>						
	(iv) <i>first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or</i>						
	(v) <i>subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.</i>						
2481N	Suppository 30 mg	12	..	..	29.89	28.60	Proladone PL

OXYCODONE HYDROCHLORIDE**CAUTION:**

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

NOTE:

Authorities for increased maximum quantities and/or repeats will be granted only for:

- (i) *severe disabling pain associated with proven malignant neoplasia; or*
- (ii) *chronic severe disabling pain not responding to non-narcotic analgesics where treatment was initiated in a hospital (in-patient or out-patient) prior to 1 April 2005. This criterion will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006; or*
- (iii) *chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or*
- (iv) *first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or*
- (v) *subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.*

2622B	Tablet 5 mg	20	..	..	10.51	11.48	Endone	SI
8464L	Capsule 5 mg	20	..	..	10.51	11.48	OxyNorm	MF
8501K	Capsule 10 mg	20	..	..	13.58	14.55	OxyNorm	MF
8502L	Capsule 20 mg	20	..	..	18.25	19.22	OxyNorm	MF
8644Y	Oral solution 5 mg per 5 mL, 250 mL	1	..	..	15.09	16.06	OxyNorm Liquid 5mg/5mL	MF

continued ➡

NERVOUS SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
OXYCODONE HYDROCHLORIDE—cont.								
Restricted benefit								
<i>Chronic severe disabling pain not responding to non-narcotic analgesics.</i>								
NOTE:								
<i>Authorities for increased maximum quantities and/or repeats will be granted only for:</i>								
<i>(i) chronic severe disabling pain associated with proven malignant neoplasia; or</i>								
<i>(ii) chronic severe disabling pain not responding to non-narcotic analgesics where treatment was initiated in a hospital (in-patient or out-patient) prior to 1 April 2005. This criterion will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006; or</i>								
<i>(iii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or</i>								
<i>(iv) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or</i>								
<i>(v) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.</i>								
8681X	Tablet 5 mg (controlled release)	20	..	..	15.73	16.70	OxyContin	MF
8385H	Tablet 10 mg (controlled release)	20	..	..	19.77	20.74	OxyContin	MF
8386J	Tablet 20 mg (controlled release)	20	..	..	28.45	28.60	OxyContin	MF
8387K	Tablet 40 mg (controlled release)	20	..	..	44.46	28.60	OxyContin	MF
8388L	Tablet 80 mg (controlled release)	20	..	..	72.02	28.60	OxyContin	MF
• Phenylpiperidine derivatives								
FENTANYL								
CAUTION:								
<i>The risk of drug dependence is high.</i>								
Restricted benefit								
<i>Chronic severe disabling pain which is associated with proven malignant neoplasia and which is unresponsive to non-narcotic analgesics.</i>								
8337T	Transdermal patch 2.5 mg (releasing approximately 25 micrograms per hour)	5	..	..	60.03	28.60	Durogesic 25	JC
8338W	Transdermal patch 5 mg (releasing approximately 50 micrograms per hour)	5	..	..	103.60	28.60	Durogesic 50	JC
8339X	Transdermal patch 7.5 mg (releasing approximately 75 micrograms per hour)	5	..	..	140.78	28.60	Durogesic 75	JC

continued ➞

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
FENTANYL—cont.								
8340Y	Transdermal patch 10 mg (releasing approximately 100 micrograms per hour)	5	..	..	171.44	28.60	Durogesic 100	JC
PETHIDINE HYDROCHLORIDE								
CAUTION: The risk of drug dependence is high.								
Restricted benefit Short-term treatment of acute pain.								
NOTE: No applications for increased maximum quantities and/or repeats will be authorised.								
1828F	Injection 50 mg in 1 mL	5	..	..	10.88	11.85	MX	
1829G	Injection 100 mg in 2 mL	5	..	..	11.31	12.28	MX	
• Diphenylpropylamine derivatives								
METHADONE HYDROCHLORIDE								
CAUTION: The risk of drug dependence is high.								
Restricted benefit Severe disabling pain not responding to non-narcotic analgesics.								
NOTE: Authorities for increased maximum quantities and/or repeats will be granted only for:								
(i) severe disabling pain associated with proven malignant neoplasia; or								
(ii) chronic severe disabling pain not responding to non-narcotic analgesics where treatment was initiated in a hospital (in-patient or out-patient) prior to 1 April 2005. This criterion will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006; or								
(iii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or								
(iv) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or								
(v) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.								
1609Q	Tablet 10 mg	20	..	..	13.08	14.05	Physeptone	GK
1606M	Injection 10 mg in 1 mL	5	..	..	28.80	28.60	Physeptone	GK

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Oripavine derivatives BUPRENORPHINE CAUTION: <i>The risk of drug dependence is high.</i> Restricted benefit <i>Chronic severe disabling pain not responding to non-narcotic analgesics.</i> NOTE: <i>Authorities for increased maximum quantities and/or repeats will be granted only for:</i> <i>(i) chronic severe disabling pain associated with proven malignant neoplasia; or</i> <i>(ii) chronic severe disabling pain not responding to non-narcotic analgesics where treatment was initiated in a hospital (in-patient or out-patient) prior to 1 April 2005. This criterion will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2006; or</i> <i>(iii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or</i> <i>(iv) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or</i> <i>(v) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.</i>								
8865N	Transdermal patch 5 mg (releasing approximately 5 micrograms per hour)	2	..	..	24.73	25.70	Norspan	MF
8866P	Transdermal patch 10 mg (releasing approximately 10 micrograms per hour)	2	..	..	35.53	28.60	Norspan	MF
8867Q	Transdermal patch 20 mg (releasing approximately 20 micrograms per hour)	2	..	..	55.62	28.60	Norspan	MF
• Other opioids TRAMADOL HYDROCHLORIDE Restricted benefit <i>For acute pain where aspirin and/or paracetamol alone are inappropriate or have failed.</i> NOTE: <i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8455B	Capsule 50 mg	20	..	..	8.36	9.33	^a Chem mart Tramadol	CH
							^a GenRx Tramadol	GX
							^a Terry White Chemists Tramadol	TW
							^a Zydol	AW
				^B 0.60	8.96	9.33	^a Tramal	CS

continued ➡

NERVOUS SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
TRAMADOL HYDROCHLORIDE—cont.								
<u>Restricted benefit</u>								
<i>For dosage titration in chronic pain where aspirin and/or paracetamol alone are inappropriate or have failed.</i>								
<u>NOTE:</u>								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8611F	Capsule 50 mg	20	2	..	8.36	9.33	^a Chem mart Tramadol	CH
							^a GenRx Tramadol	GX
							^a Terry White Chemists Tramadol	TW
							^a Zydol	AW
				^B 0.60	8.96	9.33	^a Tramal	CS
<u>Restricted benefit</u>								
<i>For pain where aspirin and/or paracetamol alone are inappropriate or have failed.</i>								
<u>NOTE:</u>								
<i>Authorities for increased maximum quantities and/or repeats will be granted only for severe disabling pain not responding to non-narcotic analgesics.</i>								
8523N	Tablet 100 mg (sustained release)	20	..	..	14.56	15.53	^a Tramahexal SR	HX
							^a Zydol SR 100	AW
				^B 1.04	15.60	15.53	^a Tramal SR 100	CS
8524P	Tablet 150 mg (sustained release)	20	..	..	17.97	18.94	^a Tramahexal SR	HX
							^a Zydol SR 150	AW
				^B 1.04	19.01	18.94	^a Tramal SR 150	CS
8525Q	Tablet 200 mg (sustained release)	20	..	..	20.84	21.81	^a Tramahexal SR	HX
							^a Zydol SR 200	AW
				^B 1.05	21.89	21.81	^a Tramal SR 200	CS
8843K	Oral drops 100 mg per mL, 10 mL	‡1	..	..	8.36	9.33	Tramal	CS
<u>Restricted benefit</u>								
<i>Short-term treatment of acute pain.</i>								
<u>NOTE:</u>								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8582Q	Injection 100 mg in 2 mL	5	..	..	11.11	12.08	Tramal 100	CS
Other analgesics and antipyretics								
• Salicylic acid and derivatives								
ASPIRIN								
1010E	Tablet 300 mg (dispersible)	96	1	..	6.98	7.95	Solprin	RC

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Anilides								
PARACETAMOL								
1746X	Tablet 500 mg	100	1	..	7.39	8.36	Dymadon P	PC
							^a Febridol	DG
							^a Panamax	SW
							Parahexal	HX
							^a Paralgin	FM
							Parmol	AW
							^a Terry White Chemists Paracetamol	TW
							Tylenol	JT
1747Y	Oral liquid 120 mg per 5 mL, 100 mL	‡1	2	..	7.68	8.65	Panamax	SW
1770E	Oral liquid 240 mg per 5 mL, 200 mL	‡1	2	..	8.95	9.92	Panamax 240 Elixir	SW
PARACETAMOL								
<u>Restricted benefit</u>								
<i>Chronic arthropathies.</i>								
8784H	Tablet 500 mg	300	4	..	*12.67	13.64	Dymadon P	PC
							^a Febridol	DG
							^a Panamax	SW
							Parahexal	HX
							^a Paralgin	FM
							Parmol	AW
							^a Terry White Chemists Paracetamol	TW
							Tylenol	JT
<u>Restricted benefit</u>								
<i>Relief of persistent pain associated with osteoarthritis.</i>								
8814X	Tablet 665 mg (modified release)	192	5	..	*11.51	12.48	^a Duatrol SR	ME
				B4.88	*16.39	12.48	^a Panadol Osteo	GC
Antimigraine preparations								
• Ergot alkaloids								
DIHYDROERGOTAMINE MESYLATE								
1323P	Injection 1 mg in 1 mL	5	..	..	15.27	16.24	Dihydergot	NV
METHYSERGIDE								
2826R	Tablet 1 mg	100	2	..	*43.45	28.60	Deseril	NV

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Selective 5HT₁-receptor agonists NARATRIPTAN HYDROCHLORIDE CAUTION: <i>Naratriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.</i> Authority required <i>Migraine attacks in patients receiving, or who have failed a reasonable trial of, prophylactic medication and where attacks in the past have usually failed to respond to oral therapy with ergotamine and other appropriate agents, or in whom these agents are contraindicated.</i> NOTE: <i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8298R	Tablet 2.5 mg (base)	4	5	..	*26.75	27.72	Naramig	GK
SUMATRIPTAN CAUTION: <i>Sumatriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.</i> Authority required <i>Migraine attacks in patients receiving, or who have failed a reasonable trial of, prophylactic medication and where attacks in the past have usually failed to respond to oral therapy with ergotamine and other appropriate agents, or in whom these agents are contraindicated.</i> NOTE: <i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8341B	Nasal spray 20 mg in 0.1 mL single dose unit	2	5	..	15.75	16.72	Imigran	GK
SUMATRIPTAN SUCCINATE CAUTION: <i>Sumatriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.</i> Authority required <i>Migraine attacks in patients receiving, or who have failed a reasonable trial of, prophylactic medication and where attacks in the past have usually failed to respond to oral therapy with ergotamine and other appropriate agents, or in whom these agents are contraindicated.</i> NOTE: <i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8144P	Tablet 50 mg (base)	4	5	..	*26.75	27.72	^a Imigran ^a Suvalan 50	GK AW

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ZOLMITRIPTAN								
CAUTION: <i>Zolmitriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.</i>								
Authority required <i>Migraine attacks in patients receiving, or who have failed a reasonable trial of, prophylactic medication and where attacks in the past have usually failed to respond to oral therapy with ergotamine and other appropriate agents, or in whom these agents are contraindicated.</i>								
NOTE: <i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8266C	Tablet 2.5 mg	4	5	..	*26.69	27.66	Zomig	AP
• Other antimigraine preparations								
CYPROHEPTADINE HYDROCHLORIDE								
Restricted benefit <i>Prevention of migraine.</i>								
NOTE: <i>Cyproheptadine hydrochloride is not PBS-subsidised for use in hay fever or atopy.</i>								
1798P	Tablet 4 mg	100	2	..	*11.83	12.80	Periactin	FR
PIZOTIFEN MALATE								
3074T	Tablet 500 micrograms (base)	100	2	..	19.91	20.88	Sandomigran 0.5	NV
ANTIEPILEPTICS								
Antiepileptics								
• Barbiturates and derivatives								
PHENOBARBITONE								
Restricted benefit <i>Epilepsy.</i>								
1850J	Tablet 30 mg	200	4	..	9.25	10.22	SI	
PHENOBARBITONE SODIUM								
Restricted benefit <i>Epilepsy.</i>								
1853M	Injection 200 mg in 1 mL	5	..	..	17.59	18.56	FM	
PRIMIDONE								
1939C	Tablet 250 mg	200	2	..	52.07	28.60	Mysoline	LM
• Hydantoin derivatives								
PHENYTOIN								
1249R	Tablet 50 mg	200	2	..	28.18	28.60	Dilantin Infatabs	PF
2692Q	Paediatric oral suspension 30 mg per 5 mL, 500 mL	‡1	3	..	19.77	20.74	Dilantin	PF
PHENYTOIN SODIUM								
1873N	Capsule 30 mg	200	2	..	27.26	28.23	Dilantin Sodium	PF
1874P	Capsule 100 mg	200	2	..	28.18	28.60	Dilantin Sodium	PF

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Succinimide derivatives								
ETHOSUXIMIDE								
1413J	Capsule 250 mg	200	2	..	53.83	28.60	Zarontin	PF
1414K	Paediatric syrup 250 mg per 5 mL, 200 mL	‡1	5	..	23.41	24.38	Zarontin	PF
• Benzodiazepine derivatives								
CLONAZEPAM								
<u>Restricted benefit</u>								
<u>Epilepsy.</u>								
1807D	Injection 1 mg in 2 mL (set containing solution 1 mg in 1 mL and 1 mL diluent)	5	..	..	16.76	17.73	Rivotril	RO
<u>Authority required</u>								
<u>Neurologically proven epilepsy.</u>								
<u>CAUTION:</u>								
<u>Abuse of clonazepam has been reported. Refer to the current product information.</u>								
1805B	Tablet 500 micrograms	200	2	..	*22.91	23.88	^a Paxam 0.5	AF
				^B 4.74	*27.65	23.88	^a Rivotril	RO
1806C	Tablet 2 mg	200	2	..	*38.89	28.60	^a Paxam 2	AF
				^B 5.36	*44.25	28.60	^a Rivotril	RO
1808E	Oral liquid 2.5 mg per mL, 10 mL	2	..	..	*13.29	14.26	Rivotril	RO
NITRAZEPAM								
<u>Authority required</u>								
<u>Myoclonic epilepsy;</u>								
<u>Malignant neoplasia (late stage);</u>								
<u>For use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal;</u>								
<u>For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.</u>								
2732T	Tablet 5 mg	50	5	..	*8.63	9.60	^a Alodorm	AF
				^B 4.04	*12.67	9.60	^a Mogadon	VT
[For other listings for this drug see Generic/Proprietary Index]								
• Carboxamide derivatives								
CARBAMAZEPINE								
2422L	Tablet 100 mg	200	2	..	21.51	22.48	^a Carbamazepine- BC	BG
							^a Carbamazepine Sandoz	SZ
				^B 2.50	24.01	22.48	^a Tegretol 100	NV

continued ➡

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CARBAMAZEPINE—cont.								
2419H	Tablet 200 mg	200	2	..	36.09	28.60	^a Carbamazepine- BC	BG
							^a Carbamazepine Sandoz	SZ
				^B 3.00	39.09	28.60	^a Teril	AF
							^a Tegretol 200	NV
2426Q	Tablet 200 mg (controlled release)	200	2	..	36.73	28.60	Tegretol CR 200	NV
2431Y	Tablet 400 mg (controlled release)	200	2	..	65.51	28.60	Tegretol CR 400	NV
2427R	Oral suspension 100 mg per 5 mL, 300 mL	‡1	5	..	19.50	20.47	Tegretol Liquid	NV

OXCARBAZEPINE**Authority required**

Treatment of partial epileptic seizures and primary generalised tonic-clonic seizures, which are not controlled satisfactorily by other anti-epileptic drugs.

8584T	Tablet 150 mg	100	5	..	76.98	28.60	Trileptal	NV
8585W	Tablet 300 mg	100	5	..	123.96	28.60	Trileptal	NV
8586X	Tablet 600 mg	100	5	..	203.82	28.60	Trileptal	NV
8588B	Oral suspension 60 mg per mL, 250 mL	2	5	..	*149.21	28.60	Trileptal	NV

• **Fatty acid derivatives**

SODIUM VALPROATE

CAUTION:

There are reports of

- (i) fatal hepatotoxicity, particularly in children;
- (ii) teratogenesis (spina bifida in humans).

2294R	Crushable tablet 100 mg	200	2	..	*29.67	28.60	Epilim	SW
2289L	Tablet 200 mg (enteric coated)	200	2	..	*36.39	28.60	^a Valpro 200	AF
				^B 1.32	*37.71	28.60	^a Epilim EC	SW
2290M	Tablet 500 mg (enteric coated)	200	2	..	*67.35	28.60	^a Valpro 500	AF
				^B 1.64	*68.99	28.60	^a Epilim EC	SW
2293Q	Oral liquid 200 mg per 5 mL, 300 mL	2	2	..	*32.91	28.60	Epilim Liquid	SW
2295T	Syrup 200 mg per 5 mL, 300 mL	2	2	..	*32.91	28.60	Epilim Syrup	SW

TIAGABINE HYDROCHLORIDE**Authority required**

Treatment of partial epileptic seizures which are not controlled satisfactorily by other anti-epileptic drugs.

8221Q	Tablet 5 mg (base)	100	5	..	77.39	28.60	Gabitril	MX
8222R	Tablet 10 mg (base)	100	5	..	150.02	28.60	Gabitril	MX
8223T	Tablet 15 mg (base)	100	5	..	212.68	28.60	Gabitril	MX

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
VIGABATRIN								
CAUTION:								
<i>Visual field defects have been reported with this drug.</i>								
Authority required								
<i>Treatment of epileptic seizures which are not controlled satisfactorily by other anti-epileptic drugs.</i>								
2667J	Tablet 500 mg	100	5	..	97.88	28.60	Sabril	AV
2668K	Oral powder, sachet 500 mg	60	5	..	64.91	28.60	Sabril	AV
• Other antiepileptics								
GABAPENTIN								
Authority required								
<i>Treatment of partial epileptic seizures which are not controlled satisfactorily by other anti-epileptic drugs.</i>								
8505P	Capsule 100 mg	100	5	..	32.66	28.60	^a Gantin	AW
							^a Neurontin	PF
							^a Nupentin 100	AF
1834M	Capsule 300 mg	100	5	..	86.34	28.60	^a DBL	MX
							^a Gabapentin	
							^a Douglas	DP
							^a Gabapentin	
							^a Gabahexal	HX
							^a 300mg	
							^a Gantin	AW
							^a GenRx	GX
							^a Gabapentin	
							^a Neurontin	PF
							^a Nupentin 300	AF
							^a Pendine 300	AL
1835N	Capsule 400 mg	100	5	..	114.22	28.60	^a DBL	MX
							^a Gabapentin	
							^a Douglas	DP
							^a Gabapentin	
							^a 400mg	
							^a Gabahexal	HX
							^a 400mg	
							^a Gantin	AW
							^a GenRx	GX
							^a Gabapentin	
							^a Neurontin	PF
							^a Nupentin 400	AF
							^a Pendine 400	AL
8559L	Tablet 600 mg	100	5	..	180.78	28.60	Neurontin	PF
8389M	Tablet 800 mg	100	5	..	228.71	28.60	^a Gantin	AW
							^a Neurontin	PF
							^a Pendine 800	AF

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
LAMOTRIGINE							
Authority required							
Treatment of epileptic seizures which are not controlled satisfactorily by other anti-epileptic drugs.							
8063J	Tablet 5 mg	56	5	..	19.19	20.16	a Elmendos ME a Lamitrin HX a Lamogine AF a Seaze 5 AW a Lamictal GK
				B1.00	20.19	20.16	
2848X	Tablet 25 mg	56	5	..	36.90	28.60	a Elmendos ME a GenRx GX Lamotrigine a Lamitrin HX a Lamogine AF a Lamotrigine-DP DP a Seaze 25 AW a Lamictal GK
				B1.00	37.90	28.60	
2849Y	Tablet 50 mg	56	5	..	58.36	28.60	a Elmendos ME a GenRx GX Lamotrigine a Lamitrin HX a Lamogine AF a Lamotrigine-DP DP a Seaze 50 AW a Lamictal GK
				B0.99	59.35	28.60	
2850B	Tablet 100 mg	56	5	..	94.07	28.60	a Elmendos ME a GenRx GX Lamotrigine a Lamitrin HX a Lamogine AF a Lamotrigine-DP DP a Seaze 100 AW a Lamictal GK
				B1.00	95.07	28.60	
2851C	Tablet 200 mg	56	5	..	154.80	28.60	a Elmendos ME a GenRx GX Lamotrigine a Lamitrin HX a Lamogine AF a Lamotrigine-DP DP a Seaze 200 AW a Lamictal GK
				B1.00	155.80	28.60	
LEVETIRACETAM							
See Special Pharmaceutical Benefits							
SULTHIAME							
2099L	Tablet 50 mg	200	2	..	38.44	28.60	Ospolot PL
2100M	Tablet 200 mg	200	2	..	87.88	28.60	Ospolot PL
TOPIRAMATE							
See Special Pharmaceutical Benefits							

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTI-PARKINSON DRUGS								
Anticholinergic agents								
• Tertiary amines								
BENZHEXOL HYDROCHLORIDE								
1109J	Tablet 2 mg	200	2	..	11.89	12.86	Artane	SI
1110K	Tablet 5 mg	200	1	..	13.99	14.96	Artane	SI
BIPERIDEN HYDROCHLORIDE								
2544X	Tablet 2 mg	200	2	..	*19.05	20.02	Akineton	AB
• Ethers of tropine or tropine derivatives								
BENZTROPINE MESYLATE								
2362H	Tablet 2 mg	60	2	..	11.01	11.98	Benztrop	PL
3038X	Injection 2 mg in 2 mL	5	..	..	20.99	21.96	Cogentin	MK
Dopaminergic agents								
• Dopa and dopa derivatives								
LEVODOPA with BENSERAZIDE								
8218M	Dispersible tablet 50 mg-12.5 mg	100	5	..	21.14	22.11	Madopar Rapid 62.5	RO
8219N	Dispersible tablet 100 mg-25 mg	100	5	..	36.87	28.60	Madopar Rapid 125	RO
2229H	Tablet 100 mg-25 mg	100	5	..	36.87	28.60	Madopar 125	RO
2228G	Tablet 200 mg-50 mg	100	5	..	49.19	28.60	Madopar	RO
2227F	Capsule 50 mg-12.5 mg	100	5	..	21.14	22.11	Madopar 62.5	RO
2225D	Capsule 100 mg-25 mg	100	5	..	36.87	28.60	Madopar 125	RO
2231K	Capsule 100 mg-25 mg (sustained release)	100	5	..	40.08	28.60	Madopar HBS	RO
2226E	Capsule 200 mg-50 mg	100	5	..	49.19	28.60	Madopar	RO
LEVODOPA with CARBIDOPA								
1242J	Tablet 100 mg-25 mg	100	5	.. B5.41	38.91 44.32	28.60	^a Kinson ^a Sinemet 100/25	AF MK
1245M	Tablet 250 mg-25 mg	100	5	.. B3.20	46.87 50.07	28.60	^a Levohexal ^a Sinemet	HX MK
LEVODOPA with CARBIDOPA								
Authority required								
Parkinson's disease where fluctuations in motor function are not adequately controlled by frequent dosing with conventional formulations of levodopa with decarboxylase inhibitor.								
1255C	Tablet 200 mg-50 mg (modified release)	100	5	..	65.25	28.60	Sinemet CR	MK

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
LEVODOPA with CARBIDOPA and ENTACAPONE								
Authority required								
<i>Parkinson's disease in patients being treated with levodopa—decarboxylase inhibitor combinations who are experiencing fluctuations in motor function due to end-of-dose effect; Parkinson's disease in patients stabilised on concomitant treatment with levodopa—decarboxylase inhibitor combinations and entacapone.</i>								
8797B	Tablet 50 mg-12.5 mg-200 mg	200	4	..	*319.81	28.60	Stalevo 50/12.5/ 200mg	NV
8798C	Tablet 100 mg-25 mg-200 mg	200	4	..	*350.87	28.60	Stalevo 100/25/ 200mg	NV
8799D	Tablet 150 mg-37.5 mg-200 mg	200	4	..	*381.91	28.60	Stalevo 150/37.5/ 200mg	NV
• Adamantane derivatives								
AMANTADINE HYDROCHLORIDE								
Restricted benefit								
<i>Parkinson's disease which is not drug induced.</i>								
3016R	Capsule 100 mg	100	5	..	42.70	28.60	Symmetrel 100	NV
• Dopamine agonists								
BROMOCRIPTINE MESYLATE								
Restricted benefit								
<i>Acromegaly; Parkinson's disease; Pathological hyperprolactinaemia where surgery is not indicated; Pathological hyperprolactinaemia where surgery has already been used with incomplete resolution; Pathological hyperprolactinaemia where radiotherapy is not indicated; Pathological hyperprolactinaemia where radiotherapy has already been used with incomplete resolution.</i>								
1443Y	Tablet 2.5 mg (base)	60	5	..	33.54	28.60	^a Bromohexal	HX
							^a Krypton 2.5	AF
					^B 3.00	36.54	^a Parlodel	NV
1446D	Capsule 5 mg (base)	60	5	..	61.36	28.60	^a Bromohexal	HX
							^a Krypton 5	AF
					^B 3.00	64.36	^a Parlodel	NV
1445C	Capsule 10 mg (base)	100	5	..	207.04	28.60	^a Krypton 10	AF
					^B 3.00	210.04	^a Parlodel	NV
[For other listings for this drug see Generic/Proprietary Index]								
CABERGOLINE								
Restricted benefit								
<i>Parkinson's disease.</i>								
8393R	Tablet 1 mg	30	5	..	70.38	28.60	Cabaser	PU
8394T	Tablet 2 mg	30	5	..	92.70	28.60	Cabaser	PU
8395W	Tablet 4 mg	30	5	..	111.78	28.60	Cabaser	PU

NERVOUS SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PERGOLIDE MESYLATE								
<u>Restricted benefit</u>								
<i>Parkinson's disease as adjunctive therapy in patients being treated with levodopa—decarboxylase inhibitor combinations.</i>								
2808T	Tablet 50 micrograms (base)	100	..	..	52.50	28.60	Permax	AS
2809W	Tablet 250 micrograms (base)	100	5	..	66.50	28.60	Permax	AS
2810X	Tablet 1 mg (base)	100	5	..	247.27	28.60	Permax	AS
• Monoamine oxidase type B inhibitors								
SELEGILINE HYDROCHLORIDE								
<u>Restricted benefit</u>								
<i>Late stage Parkinson's disease as adjunctive therapy in patients being treated with levodopa—decarboxylase inhibitor combinations.</i>								
1973W	Tablet 5 mg	100	5	..	56.56	28.60	^a Eldepryl ^a Selgene	DP AF
• Other dopaminergic agents								
ENTACAPONE								
<u>Authority required</u>								
<i>Parkinson's disease as adjunctive therapy in patients being treated with levodopa—decarboxylase inhibitor combinations who are experiencing fluctuations in motor function due to end-of-dose effect.</i>								
8367J	Tablet 200 mg	200	4	..	*288.75	28.60	Comtan	NV
PSYCHOLEPTICS								
Antipsychotics								
• Phenothiazine with aliphatic side-chain								
CHLORPROMAZINE								
HYDROCHLORIDE								
1196Y	Tablet 10 mg	100	5	..	7.87	8.84	Largactil	AV
1197B	Tablet 25 mg	100	5	..	9.37	10.34	Largactil	AV
1199D	Tablet 100 mg	100	5	..	13.19	14.16	Largactil	AV
1201F	Mixture 25 mg per 5 mL, 100 mL	‡1	5	..	9.15	10.12	Largactil	AV
1195X	Injection 50 mg in 2 mL	10	..	..	13.99	14.96	Largactil	AV
• Phenothiazine with piperazine structure								
FLUPHENAZINE DECANOATE								
1046C	Injection 12.5 mg in 0.5 mL	5	..	..	17.40	18.37	Modecate	BQ
3098C	Injection 25 mg in 1 mL	5	..	..	24.48	25.45	Modecate	BQ
1001Q	Injection 50 mg in 2 mL	5	..	..	35.63	28.60	Modecate	BQ
TRIFLUOPERAZINE HYDROCHLORIDE								
2185B	Tablet 1 mg (base)	100	5	..	7.94	8.91	Stelazine	LM
2386N	Tablet 2 mg (base)	100	5	..	8.82	9.79	Stelazine	LM
2186C	Tablet 5 mg (base)	100	5	..	9.66	10.63	Stelazine	LM

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Phenothiazines with piperidine structure								
PERICYAZINE								
3052P	Tablet 2.5 mg	100	5	..	8.49	9.46	Neulactil	AV
3053Q	Tablet 10 mg	100	5	..	12.69	13.66	Neulactil	AV
THIORIDAZINE HYDROCHLORIDE								
CAUTION:								
<i>Thioridazine may cause serious cardiac arrhythmias.</i>								
Authority required								
<i>Management of patients with schizophrenia who have failed to respond adequately to treatment with appropriate courses of at least 2 other antipsychotic drugs, at an adequate dose and for an adequate duration, because of:</i>								
<i>(a) insufficient effectiveness; or</i>								
<i>(b) the inability to achieve an effective dose due to intolerable adverse effects from those drugs.</i>								
2163W	Tablet 10 mg	100	5	..	8.60	9.57	Aldazine 10	AF
2359E	Tablet 25 mg	100	5	..	10.61	11.58	Aldazine 25	AF
2164X	Tablet 50 mg	100	5	..	13.96	14.93	Aldazine 50	AF
2165Y	Tablet 100 mg	100	5	..	20.90	21.87	Aldazine 100	AF
• Butyrophenone derivatives								
HALOPERIDOL								
2761H	Tablet 500 micrograms	100	5	..	8.35	9.32	Serenace	SI
2767P	Tablet 1.5 mg	100	5	..	8.81	9.78	Serenace	SI
2770T	Tablet 5 mg	50	5	..	8.82	9.79	Serenace	SI
2763K	Oral liquid 2 mg per mL, 100 mL	‡1	5	..	14.58	15.55	Serenace	SI
2768Q	Injection 5 mg in 1 mL	10	..	..	15.86	16.83	Serenace	SI
HALOPERIDOL DECANOATE								
2765M	I.M. injection equivalent to 50 mg haloperidol in 1 mL	5	..	..	24.48	25.45	Haldol decanoate	JC
2766N	I.M. injection equivalent to 150 mg haloperidol in 3 mL	5	..	..	44.33	28.60	Haldol decanoate	JC
• Thioxanthene derivatives								
FLUPENTHIXOL DECANOATE								
2255Q	Oily I.M. injection 20 mg in 1 mL	5	..	..	17.40	18.37	Fluanxol Depot	LU
2256R	Oily I.M. injection 40 mg in 2 mL	5	..	..	24.48	25.45	Fluanxol Depot	LU
2257T	Oily I.M. injection 100 mg in 1 mL	5	..	..	43.35	28.60	Fluanxol Concentrated Depot	LU
ZUCLOPENTHIXOL DECANOATE								
8097E	Oily I.M. injection 200 mg in 1 mL	5	..	..	23.45	24.42	Clopixol Depot	LU

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Diazepines, oxazepines and thiazepines OLANZAPINE <u>Authority required</u> Schizophrenia; Maintenance treatment of bipolar I disorder.								
8170B	Tablet 2.5 mg	30	5	..	58.17	28.60	Zyprexa	LY
8185T	Tablet 5 mg	30	5	..	110.53	28.60	Zyprexa	LY
8186W	Tablet 7.5 mg	30	5	..	165.01	28.60	Zyprexa	LY
8187X	Tablet 10 mg	30	5	..	217.01	28.60	Zyprexa	LY
8433W	Wafer 5 mg	28	5	..	103.46	28.60	Zyprexa Zydys	LY
8434X	Wafer 10 mg	28	5	..	204.05	28.60	Zyprexa Zydys	LY
QUETIAPINE FUMARATE <u>Authority required</u> Schizophrenia.								
8456C	Tablet 25 mg (base)	60	5	..	53.68	28.60	Seroquel	AP
8457D	Tablet 100 mg (base)	90	5	..	148.26	28.60	Seroquel	AP
8458E	Tablet 200 mg (base)	60	5	..	201.24	28.60	Seroquel	AP
8580N	Tablet 300 mg (base)	60	5	..	283.75	28.60	Seroquel	AP
• Neuroleptics, in tardive dyskinesia TETRABENAZINE <u>Authority required</u> Hyperkinetic extrapyramidal disorders.								
1330B	Tablet 25 mg	112	5	..	320.86	28.60	OA	
• Benzamides AMISULPRIDE <u>Authority required</u> Schizophrenia.								
8594H	Tablet 100 mg	30	5	..	36.85	28.60	Solian 100	SW
8595J	Tablet 200 mg	60	5	..	143.05	28.60	Solian 200	SW
8596K	Tablet 400 mg	60	5	..	263.05	28.60	Solian 400	SW
8736T	Oral solution 100 mg per mL, 60 mL	2	5	..	*143.05	28.60	Solian Solution	SW
• Lithium LITHIUM CARBONATE For listings see Generic/Proprietary Index								
• Other antipsychotics ARIPIPRAZOLE <u>Authority required</u> Schizophrenia.								
8717T	Tablet 10 mg	30	5	..	159.78	28.60	Abilify	BQ
8718W	Tablet 15 mg	30	5	..	222.77	28.60	Abilify	BQ

continued ↗

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ARIPIPRAZOLE—cont.								
8719X	Tablet 20 mg	30	5	..	266.23	28.60	Abilify	BQ
8720Y	Tablet 30 mg	30	5	..	319.45	28.60	Abilify	BQ
RISPERIDONE								
Authority required								
<i>Behavioural disturbances characterised by psychotic symptoms and aggression in patients with dementia where non-pharmacological methods have been unsuccessful.</i>								
CAUTION:								
<i>In placebo controlled trials in elderly patients with dementia there was a significantly higher incidence of cerebrovascular adverse events, such as stroke (including fatalities) and transient ischaemic attacks, in patients treated with risperidone compared with patients treated with placebo.</i>								
8787L	Tablet 0.5 mg	60	2	..	*39.16	28.60	Risperdal	JC
8788M	Tablet 0.5 mg (orally disintegrating)	56	2	..	*36.89	28.60	Risperdal Quicklet	JC
8789N	Tablet 1 mg	60	2	..	73.61	28.60	Risperdal	JC
8790P	Tablet 1 mg (orally disintegrating)	56	2	..	*69.01	28.60	Risperdal Quicklet	JC
8791Q	Oral solution 1 mg per mL, 30 mL	‡1	2	..	39.35	28.60	Risperdal	JC
Authority required								
<i>Schizophrenia.</i>								
8869T	Tablet 0.5 mg	60	5	..	*39.16	28.60	Risperdal	JC
8870W	Tablet 0.5 mg (orally disintegrating)	56	5	..	*36.89	28.60	Risperdal Quicklet	JC
3169T	Tablet 1 mg	60	5	..	73.61	28.60	Risperdal	JC
8792R	Tablet 1 mg (orally disintegrating)	56	5	..	*69.01	28.60	Risperdal Quicklet	JC
3170W	Tablet 2 mg	60	5	..	143.13	28.60	Risperdal	JC
8794W	Tablet 2 mg (orally disintegrating)	56	5	..	*133.91	28.60	Risperdal Quicklet	JC
3171X	Tablet 3 mg	60	5	..	209.75	28.60	Risperdal	JC
3172Y	Tablet 4 mg	60	5	..	271.95	28.60	Risperdal	JC
8100H	Oral solution 1 mg per mL, 100 mL	‡1	5	..	120.07	28.60	Risperdal	JC
8780D	Powder for I.M. injection 25 mg (modified release) with 2 mL diluent in pre-filled syringe	2	5	..	*329.37	28.60	Risperdal Consta	JC
8781E	Powder for I.M. injection 37.5 mg (modified release) with 2 mL diluent in pre-filled syringe	2	5	..	*421.61	28.60	Risperdal Consta	JC

continued ➞

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
RISPERIDONE—cont.								
8782F	Powder for I.M. injection 50 mg (modified release) with 2 mL diluent in pre-filled syringe	2	5	..	*514.55	28.60	Risperdal Consta	JC
Anxiolytics								
• Benzodiazepine derivatives								
ALPRAZOLAM								
<u>Authority required</u>								
<i>Panic disorder where other treatments have failed or are inappropriate.</i>								
2130D	Tablet 250 micrograms	50	..	..	8.86	9.83	^a Alprax 0.25 ^a Kalma 0.25 ^a Zamhexal 0.25mg	AW AF HX
				^B 1.08	9.94	9.83	^a Xanax	PH
2131E	Tablet 500 micrograms	50	..	..	11.41	12.38	^a Alprax 0.5 ^a Kalma 0.5 ^a Zamhexal 0.5mg	AW AF HX
				^B 1.16	12.57	12.38	^a Xanax	PH
2132F	Tablet 1 mg	50	2	..	16.37	17.34	^a Alprax 1 ^a Alprazolam-DP ^a Chem mart Alprazolam ^a GenRx Alprazolam ^a Kalma 1 ^a Terry White Chemists Alprazolam ^a Zamhexal 1.0mg	AW DP CH GX AF TW HX
				^B 1.35	17.72	17.34	^a Xanax	PH
8118G	Tablet 2 mg	50	2	..	22.71	23.68	^a Alprax 2 ^a Alprazolam-DP ^a Chem mart Alprazolam ^a GenRx Alprazolam ^a Kalma 2 ^a Terry White Chemists Alprazolam ^a Zamhexal 2mg ^B 1.65 24.36 23.68 ^a Xanax Tri-Score	AW DP CH GX AF TW HX PH
DIAZEPAM								
3161J	Tablet 2 mg	50	..	..	6.79	7.76	^a Antenex 2 ^a Valpam 2	AF AW
				^B 0.91	7.70	7.76	Ducene	SU
				^B 1.30	8.09	7.76	^a Valium	RO

continued ➡

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
DIAZEPAM—cont.							
3162K	Tablet 5 mg	50	..	..	7.03	8.00	^a Antenex 5 AF
							^a Diazepam-DP DP
				^B 0.93	7.96	8.00	^a Valpam 5 AW
				^B 1.32	8.35	8.00	Ducene SU
							^a Valium RO
2558P	Injection 10 mg in 2 mL	5	..	..	10.55	11.52	MX

NOTE:

Authorities for increased maximum quantities and/or repeats for the oral forms of diazepam will be granted only for

- (i) the treatment of disabling spasticity; or
 - (ii) malignant neoplasia (late stage); or
 - (iii) use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past six months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal; or
 - (iv) use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past six months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.
- Up to six months' treatment (i.e. one month's treatment with five repeats) may be requested.

OXAZEPAM							
3132W	Tablet 15 mg	25	..	..	6.04	7.01	^a Alepam 15 AF
				^B 1.44	7.48	7.01	^a Serepax SI
3133X	Tablet 30 mg	25	..	..	6.25	7.22	^a Alepam 30 AF
							^a Murelax FM
				^B 1.45	7.70	7.22	^a Serepax SI

NOTE:

Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of oxazepam below.

OXAZEPAM**Authority required**

Malignant neoplasia (late stage);

For use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.

3134Y	Tablet 15 mg	50	5	..	*7.33	8.30	^a Alepam 15 AF
				^B 2.88	*10.21	8.30	^a Serepax SI
3135B	Tablet 30 mg	50	5	..	*7.75	8.72	^a Alepam 30 AF
							^a Murelax FM
				^B 2.90	*10.65	8.72	^a Serepax SI

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
• Other anxiolytics CLOMIPRAMINE HYDROCHLORIDE Restricted benefit <i>Cataplexy associated with narcolepsy;</i> <i>Obsessive-compulsive disorder;</i> <i>Phobic disorders in adults.</i>							
1561E	Tablet 25 mg	50	2	..	18.48	19.45	^a Chem mart CH ^a Clomipramine ^a GenRx GX ^a Clomipramine ^a Placil AF ^a Terry White TW ^a Chemists ^a Clomipramine ^B 2.94 21.42 19.45 ^a Anafranil 25 NV

Hypnotics and sedatives**• Benzodiazepine derivatives**

NITRAZEPAM							
2723H	Tablet 5 mg	25	..	..	6.69	7.66	^a Alodorm AF
				^B 2.02	8.71	7.66	^a Mogadon VT

NOTE:

Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of nitrazepam below.

NITRAZEPAM**Authority required**

Myoclonic epilepsy;

Malignant neoplasia (late stage);

For use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.

2732T	Tablet 5 mg	50	5	..	*8.63	9.60	^a Alodorm AF
				^B 4.04	*12.67	9.60	^a Mogadon VT
TEMAZEPAM							
2089Y	Tablet 10 mg	25	..	..	6.69	7.66	^a Temaze AF
				^B 1.06	7.75	7.66	^a Temtabs FM
							^a Normison SI

NOTE:

Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of temazepam below.

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
TEMAZEPAM—cont.								
Authority required								
<i>Malignant neoplasia (late stage);</i>								
<i>For use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal;</i>								
<i>For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.</i>								
2088X	Tablet 10 mg	50	5	..	*8.63	9.60	^a Temaze	AF
							^a Temtabs	FM
				^B 2.12	*10.75	9.60	^a Normison	SI
PSYCHOANALEPTICS								
Antidepressants								
• Non-selective monoamine reuptake inhibitors								
AMITRIPTYLINE HYDROCHLORIDE								
2417F	Tablet 10 mg	50	2	..	6.75	7.72	Endep 10	AF
2418G	Tablet 25 mg	50	2	..	6.84	7.81	^a Endep 25	AF
				^B 1.25	8.09	7.81	^a Tryptanol	MK
2429W	Tablet 50 mg	50	2	..	7.48	8.45	Endep 50	AF
CLOMIPRAMINE HYDROCHLORIDE								
Restricted benefit								
<i>Cataplexy associated with narcolepsy;</i>								
<i>Obsessive-compulsive disorder;</i>								
<i>Phobic disorders in adults.</i>								
1561E	Tablet 25 mg	50	2	..	18.48	19.45	^a Chem mart	CH
							^a Clomipramine	
							^a GenRx	GX
							^a Clomipramine	
							^a Placil	AF
							^a Terry White	TW
							^a Chemists	
							^a Clomipramine	
				^B 2.94	21.42	19.45	^a Anafranil 25	NV
DOTHIEPIN HYDROCHLORIDE								
1357K	Capsule 25 mg	50	2	..	7.38	8.35	^a Dothep 25	AF
				^B 1.07	8.45	8.35	^a Prothiaden	AB
1358L	Tablet 75 mg	30	2	..	7.68	8.65	^a Dothep 75	AF
				^B 1.05	8.73	8.65	^a Prothiaden	AB
DOXEPIH HYDROCHLORIDE								
1012G	Tablet 50 mg (base)	50	2	..	8.01	8.98	Deptran 50	AF
1011F	Capsule 10 mg (base)	50	2	..	6.99	7.96	Deptran 10	AF
				^B 1.85	8.84	7.96	Sinequan	PF
1013H	Capsule 25 mg (base)	50	2	..	7.38	8.35	Deptran 25	AF
				^B 1.55	8.93	8.35	Sinequan	PF

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
IMIPRAMINE HYDROCHLORIDE								
2420J	Tablet 10 mg	50	2	..	7.04	8.01	Tofranil 10	NV
2421K	Tablet 25 mg	50	2	..	7.04	8.01	Melipramine Tofranil 25	UW NV
<u>NORTRIPTYLINE HYDROCHLORIDE</u>								
<u>Restricted benefit</u>								
<i>Major depression where other antidepressant therapy has failed;</i>								
<i>Major depression where other antidepressant therapy is contraindicated.</i>								
2522R	Tablet 10 mg (base)	50	2	..	10.66	11.63	Allegron	AS
2523T	Tablet 25 mg (base)	50	2	..	12.37	13.34	Allegron	AS
• Selective serotonin reuptake inhibitors								
<u>CITALOPRAM HYDROBROMIDE</u>								
<u>Restricted benefit</u>								
<i>Major depressive disorders.</i>								
8702B	Tablet 10 mg (base)	28	5	..	19.80	20.77	Celapram	AF
8220P	Tablet 20 mg (base)	28	5	..	27.94	28.60	^a Celapram	AF
							^a Chem mart	CH
							^a Ciazil	DP
							^a GenRx	GX
							^a Citalopram	
							^a Talam	AW
							^a Talohexal	HX
							^a Terry White Chemists	TW
							^a Citalopram	
				^B 3.20	31.14	28.60	^a Cipramil	LU
8703C	Tablet 40 mg (base)	28	5	..	43.39	28.60	Celapram	AF
							GenRx	GX
							^a Citalopram	
							^a Talohexal	HX
<u>ESCITALOPRAM OXALATE</u>								
<i>See Special Pharmaceutical Benefits</i>								
<u>FLUOXETINE HYDROCHLORIDE</u>								
<u>Restricted benefit</u>								
<i>Major depressive disorders;</i>								
<i>Obsessive-compulsive disorder.</i>								
8270G	Tablet 20 mg (base)	28	5	..	27.61	28.58	^a Lovan 20 Tab	AL
	(dispersible)			^B 4.10	31.71	28.58	^a Prozac Tab	LY

continued ➡

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
FLUOXETINE HYDROCHLORIDE—cont.							
1434L	Capsule 20 mg (base)	28	5	..	27.61	28.58	^a Auscap <i>SI</i> ^a Chem mart <i>CH</i> <i>Fluoxetine</i> ^a Fluohexal <i>HX</i> ^a Fluoxebell <i>BF</i> ^a Fluoxetine-DP <i>DP</i> ^a GenRx <i>GX</i> <i>Fluoxetine</i> ^a Lovan <i>AL</i> ^a Terry White Chemists <i>TW</i> <i>Fluoxetine</i> ^a Zactin <i>AF</i> ^a Prozac 20 <i>LY</i>
1809F	Oral solution 20 mg (base) per 5 mL, 140 mL	‡1	5	..	36.79	28.60	Lovan Liquid <i>AL</i>
FLUVOXAMINE MALEATE							
<u>Restricted benefit</u> <i>Major depressive disorders;</i> <i>Obsessive-compulsive disorder.</i>							
8512B	Tablet 50 mg	30	5	..	23.13	24.10	^a Faverin 50 <i>AW</i> ^a Movox 50 <i>AL</i> ^a Voxam <i>HX</i> ^b 3.90 27.03 24.10 <i>SM</i> ^a Luvox
8174F	Tablet 100 mg	30	5	..	32.59	28.60	^a Faverin 100 <i>AW</i> ^a Movox 100 <i>AF</i> ^a Voxam <i>HX</i> ^b 3.90 36.49 28.60 <i>SM</i> ^a Luvox
PAROXETINE HYDROCHLORIDE							
<u>Restricted benefit</u> <i>Major depressive disorders;</i> <i>Obsessive-compulsive disorder;</i> <i>Panic disorder.</i>							
2242B	Tablet 20 mg (base)	30	5	..	30.74	28.60	^a Chem mart <i>CH</i> <i>Paroxetine</i> ^a GenRx <i>GX</i> <i>Paroxetine</i> ^a Oxetine <i>HX</i> ^a Paxtine <i>AF</i> ^a Terry White Chemists <i>TW</i> <i>Paroxetine</i> ^b 1.10 31.84 28.60 ^a Aropax <i>GK</i>

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
SERTRALINE HYDROCHLORIDE								
<u>Restricted benefit</u>								
<i>Major depressive disorders.</i>								
2236Q	Tablet 50 mg (base)	30	5	..	30.64	28.60	^a Chem mart Sertraline	CH
							^a Concorz	HX
							^a Eleva 50	AF
							^a GenRx	GX
							Sertraline	
							^a Sertraline-DP	DP
							^a Terry White Chemists	TW
							Sertraline	
							^a Xydep 50	AW
				^B 0.55	31.19	28.60	^a Zoloft	PF
2237R	Tablet 100 mg (base)	30	5	..	30.64	28.60	^a Chem mart Sertraline	CH
							^a Concorz	HX
							^a Eleva 100	AF
							^a GenRx	GX
							Sertraline	
							^a Sertraline-DP	DP
							^a Terry White Chemists	TW
							Sertraline	
							^a Xydep 100	AW
				^B 0.55	31.19	28.60	^a Zoloft	PF
<u>Restricted benefit</u>								
<i>Obsessive-compulsive disorder;</i>								
<i>Panic disorder where other treatments have failed or are inappropriate.</i>								
8836C	Tablet 50 mg (base)	30	5	..	30.64	28.60	^a Eleva 50	AF
							^a Xydep 50	AW
				^B 0.55	31.19	28.60	^a Zoloft	PF
8837D	Tablet 100 mg (base)	30	5	..	30.64	28.60	^a Eleva 100	AF
							^a Xydep 100	AW
				^B 0.55	31.19	28.60	^a Zoloft	PF
• Monoamine oxidase inhibitors, non-selective								
PHENELZINE SULFATE								
<u>CAUTION:</u>								
<i>This drug is an irreversible monoamine oxidase inhibitor.</i>								
<u>Restricted benefit</u>								
<i>Depression where all other anti-depressant therapy has failed or is inappropriate.</i>								
2856H	Tablet 15 mg (base)	100	1	..	101.55	28.60	Nardil	LM
TRANLYCYPROMINE SULFATE								
<u>CAUTION:</u>								
<i>This drug is an irreversible monoamine oxidase inhibitor.</i>								
2444P	Tablet 10 mg (base)	50	2	..	26.24	27.21	Parnate	LM

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Monoamine oxidase type A inhibitors MOCLOBEMIDE Restricted benefit Major depressive disorders.								
1900B	Tablet 150 mg	60	5	..	23.89	24.86	^a Arima	AF
							^a Chem mart	CH
							^a Moclobemide	
							^a Clobemix	DP
							^a GenRx	GX
							^a Moclobemide	
							^a Mohexal	HX
							^a Terry White	TW
							^a Chemists	
							^a Moclobemide	
				^B 0.95	24.84	24.86	^a Aurorix	RO
8003F	Tablet 300 mg	60	5	..	43.03	28.60	^a Arima 300	AF
							^a Chem mart	CH
							^a Moclobemide	
							^a Clobemix	DP
							^a GenRx	GX
							^a Moclobemide	
							^a Maosig	SI
							^a Mohexal	HX
							^a Terry White	TW
							^a Chemists	
							^a Moclobemide	
				^B 1.93	44.96	28.60	^a Aurorix 300 mg	RO
• Other antidepressants LITHIUM CARBONATE								
3059B	Tablet 250 mg	200	2	..	13.34	14.31	Lithicarb	AS
8290H	Tablet 450 mg (slow release)	200	2	..	*26.79	27.76	Quilonum SR	GK
MIANSERIN HYDROCHLORIDE CAUTION: Neutropenia and agranulocytosis are more frequent in the elderly, especially in the early months of therapy.								
Restricted benefit Severe depression.								
1627P	Tablet 10 mg	50	5	..	14.35	15.32	^a Lumin 10	AF
				^B 1.29	15.64	15.32	^a Tolvon	OR
1628Q	Tablet 20 mg	50	5	..	25.02	25.99	^a Lumin 20	AF
				^B 1.60	26.62	25.99	^a Tolvon	OR

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MIRTAZAPINE								
Restricted benefit								
Major depressive disorders.								
8855C	Tablet 15 mg (orally disintegrating)	30	5	..	23.13	24.10	Avanza SolTab	BP
8513C	Tablet 30 mg	30	5	..	32.32	28.60	^a Axit 30	AF
					^B 1.25	28.60	^a Mirtazon	AW
					^B 26.69	28.60	^a Avanza	BP
						28.60	^a Remeron	OR
8856D	Tablet 30 mg (orally disintegrating)	30	5	..	32.32	28.60	Avanza SolTab	BP
8857E	Tablet 45 mg (orally disintegrating)	30	5	..	50.70	28.60	Avanza SolTab	BP
REBOXETINE MESILATE								
Restricted benefit								
Major depressive disorders.								
8583R	Tablet 4 mg (base)	60	5	..	36.72	28.60	Edronax	PH
VENLAFAXINE HYDROCHLORIDE								
Restricted benefit								
Major depressive disorders.								
8068P	Tablet 37.5 mg (base)	56	5	..	43.42	28.60	Efexor	WY
8069Q	Tablet 75 mg (base)	56	5	..	52.15	28.60	Efexor	WY
8868R	Capsule 37.5 mg (base) (modified release)	28	..	..	24.09	25.06	Efexor-XR	WY
8301X	Capsule 75 mg (base) (modified release)	28	5	..	43.42	28.60	Efexor-XR	WY
8302Y	Capsule 150 mg (base) (modified release)	28	5	..	52.15	28.60	Efexor-XR	WY
Psychostimulants, agents used for ADHD and nootropics								
• Centrally acting sympathomimetics								
DEXAMPHETAMINE SULFATE								
NOTE:								
Care must be taken to comply with the provisions of State/Territory law when prescribing dexamphetamine.								
Authority required								
Use in attention deficit hyperactivity disorder, in accordance with State/Territory law; Narcolepsy.								
1165H	Tablet 5 mg	100	5	..	16.31	17.28	SI	
METHYLPHENIDATE HYDROCHLORIDE								
NOTE:								
Care must be taken to comply with the provisions of State/Territory law when prescribing methylphenidate hydrochloride.								
Authority required								
Use in attention deficit hyperactivity disorder, in accordance with State/Territory law.								
8839F	Tablet 10 mg	100	5	..	16.31	17.28	^a Attenta	AF
					^B 1.28	17.59	^a Ritalin 10	NV

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of Qty Rpts Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer

MODAFINIL**NOTE:**

Any queries concerning the arrangements to prescribe modafinil may be directed to Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe modafinil should be forwarded to:

*Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001*

Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au.

NOTE:

Modafinil is not PBS-subsidised when used in combination with PBS-subsidised dexamphetamine sulfate.

Authority required

Initial treatment, by a qualified sleep medicine practitioner, of patients with narcolepsy where:

- (i) therapy with dexamphetamine sulfate poses an unacceptable medical risk; or*
- (ii) intolerance to dexamphetamine sulfate of a severity necessitating permanent treatment withdrawal develops.*

The presence of any 1 of the following indicates treatment with dexamphetamine sulfate poses an unacceptable medical risk:

- (a) a psychiatric disorder;*
- (b) a cardiac disorder;*
- (c) a history of substance abuse;*
- (d) glaucoma;*
- (e) any other absolute contraindication to dexamphetamine sulfate as specified in the TGA-approved Product Information.*

Patients must meet the following definition of narcolepsy:

Excessive daytime sleepiness, recurrent naps or lapses into sleep occurring almost daily for at least 3 months and either:

- (i) a definite history of cataplexy and a Multiple Sleep Latency Test (MSLT) with a mean sleep latency less than 8 minutes; or*
- (ii) a MSLT with a mean sleep latency less than 8 minutes and 2 or more sleep onset rapid eye movement (REM) periods.*

The MSLT must be preceded by nocturnal polysomnography. Sleep prior to the MSLT must be at least 6 hours.

Polygraph records and sleep reports must be available for audit by Medicare Australia.

The authority application must be made in writing and must include the following:

- (a) a completed authority prescription form; and*
- (b) a completed Modafinil (Modavigil) PBS Authority Application for Use in the Treatment of Narcolepsy - Supporting Information Form [may be downloaded from the Medicare Australia website at www.medicareaustralia.gov.au.]; and*
- (c) the result and date of the polysomnography test conducted by, or under the supervision of, a qualified sleep medicine practitioner; and*
- (d) the results and date of the MSLT conducted by, or under the supervision of, a qualified sleep medicine practitioner; and*

continued ➤

NERVOUS SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
<i>MODAFINIL—cont.</i>							
<i>(e) details of the contraindication or intolerance to dexamphetamine sulfate.</i>							
<i>The results of the polysomnography test and the MSLT must be no more than 5 years old at the time of application.</i>							
<u>Authority required</u>							
<i>Continuing treatment of narcolepsy, where the patient has previously been issued with an authority prescription for this drug.</i>							
8816B	Tablet 100 mg	120	5	..	*356.09	28.60	Modavigil CS

Anti-dementia drugs• **Anticholinesterases****DONEPEZIL HYDROCHLORIDE****NOTE:**

Authority applications for initial treatment and the first application for continuing treatment should be made to Medicare Australia in writing. The second and subsequent authority applications for continuing treatment may be made by telephone. Up to a maximum of 6 months' therapy (1 month plus 5 repeats) will be approved for initial and continuing therapy.

To avoid unnecessary delays in initiating treatment with this drug, prescribers may seek telephone approval from Medicare Australia for up to 2 months' initial therapy. In order to continue the initial course of treatment, a written application must be provided to Medicare Australia following the initial telephone authority approval to confirm the information provided. No more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment will be approved for this follow-up application.

Authority required

Initial treatment of mild to moderately severe Alzheimer's disease. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist). The authority application must include the result of the baseline Mini-Mental State Examination (MMSE). This baseline MMSE must be a score of 10 or more and, if this score is at least 25 points, the result of the baseline Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) must also be specified. An application must be made in writing, but initial supply may be sought on the telephone as indicated in the Note above. For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. For the subsequent written authority application, no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment will be approved. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised; Continuing treatment, following initial therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in cognitive function as measured by an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE), OR a decrease of at least 4 points from baseline on the Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) for patients with an MMSE baseline score of at least 25 points (the initial authority application for continuing treatment must include the relevant result from the MMSE or the ADAS-Cog and must be in writing).

NERVOUS SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
DONEPEZIL HYDROCHLORIDE—cont.							
Authority required							
<i>Initial treatment of mild to moderately severe Alzheimer's disease of patients with a baseline Mini-Mental State Examination (MMSE) score of 9 or less, who are unable to register a score of 10 or more for reasons other than their Alzheimer's disease, as specified below. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist). Such patients will need to be assessed subsequently using the Clinicians Interview Based Impression of Change (CIBIC) scale. The authority application must include the result of the baseline MMSE and specify to which group(s) (see below) the patient belongs. An application must be made in writing, but initial supply may be sought on the telephone as indicated in the Note above. For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. For the subsequent written authority application, no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment will be approved. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.</i>							
<i>Patients who qualify under this criterion are from 1 or more of the following groups:</i>							
<i>(1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;</i>							
<i>(2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;</i>							
<i>(3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an MMSE test;</i>							
<i>(4) Intellectual (developmental or acquired) disability, eg Down's syndrome;</i>							
<i>(5) Significant sensory impairment despite best correction, which precludes completion of an MMSE test;</i>							
<i>(6) Prominent dysphasia, out of proportion to other cognitive and functional impairment;</i>							
<i>Continuing treatment, following initial therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in function, based on a rating of "very much improved" or "much improved" on the CIBIC scale, which must be assessed by the same clinician who initiated treatment (the initial authority application for continuing treatment must state the improvement achieved on the CIBIC scale and must be in writing).</i>							
8495D	Tablet 5 mg	28	5	..	158.75	28.60	Aricept PF
8496E	Tablet 10 mg	28	5	..	158.75	28.60	Aricept PF

GALANTAMINE HYDROBROMIDE**NOTE:**

Authority applications for initial treatment and the first application for continuing treatment should be made to Medicare Australia in writing. The second and subsequent authority applications for continuing treatment may be made by telephone. Up to a maximum of 6 months' therapy (1 month plus 5 repeats) will be approved for initial and continuing therapy.

To avoid unnecessary delays in initiating treatment with this drug, prescribers may seek telephone approval from Medicare Australia for up to 2 months' initial therapy. In order to continue the initial course of treatment, a written application must be provided to Medicare Australia following the initial telephone authority approval to confirm the information provided. No more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment will be approved for this follow-up application.

continued ➞

NERVOUS SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

GALANTAMINE HYDROBROMIDE—cont.**Authority required**

Initial treatment of mild to moderately severe Alzheimer's disease. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist). The authority application must include the result of the baseline Mini-Mental State Examination (MMSE). This baseline MMSE must be a score of 10 or more and, if this score is at least 25 points, the result of the baseline Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) must also be specified. An application must be made in writing, but initial supply may be sought on the telephone as indicated in the Note above. For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. For the subsequent written authority application, no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment will be approved. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised; Continuing treatment, following initial therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in cognitive function as measured by an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE), OR a decrease of at least 4 points from baseline on the Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) for patients with an MMSE baseline score of at least 25 points (the initial authority application for continuing treatment must include the relevant result from the MMSE or the ADAS-Cog and must be in writing).

Authority required

Initial treatment of mild to moderately severe Alzheimer's disease of patients with a baseline Mini-Mental State Examination (MMSE) score of 9 or less, who are unable to register a score of 10 or more for reasons other than their Alzheimer's disease, as specified below. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist). Such patients will need to be assessed subsequently using the Clinicians Interview Based Impression of Change (CIBIC) scale. The authority application must include the result of the baseline MMSE and specify to which group(s) (see below) the patient belongs. An application must be made in writing, but initial supply may be sought on the telephone as indicated in the Note above. For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. For the subsequent written authority application, no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment will be approved. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;*
 - (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;*
 - (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an MMSE test;*
 - (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;*
 - (5) Significant sensory impairment despite best correction, which precludes completion of an MMSE test;*
 - (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment;*
- Continuing treatment, following initial therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in function, based on a rating of "very much improved" or "much improved" on the CIBIC scale, which must be assessed by the same clinician who initiated treatment (the initial authority application for continuing treatment must state the improvement achieved on the CIBIC scale and must be in writing).*

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
GALANTAMINE HYDROBROMIDE—cont.								
8536G	Tablet 4 mg (base)	56	5	..	132.35	28.60	Reminyl	JC
8537H	Tablet 8 mg (base)	56	5	..	158.75	28.60	Reminyl	JC
8756W	Tablet 12 mg (base)	56	5	..	191.75	28.60	Reminyl	JC
8770N	Capsule 8 mg (base) (prolonged release)	28	5	..	132.35	28.60	Reminyl	JC
8771P	Capsule 16 mg (base) (prolonged release)	28	5	..	158.75	28.60	Reminyl	JC
8772Q	Capsule 24 mg (base) (prolonged release)	28	5	..	191.75	28.60	Reminyl	JC

RIVASTIGMINE HYDROGEN TARTRATE**NOTE:**

Authority applications for initial treatment and the first application for continuing treatment should be made to Medicare Australia in writing. The second and subsequent authority applications for continuing treatment may be made by telephone. Up to a maximum of 6 months' therapy (1 month plus 5 repeats) will be approved for initial and continuing therapy.

To avoid unnecessary delays in initiating treatment with this drug, prescribers may seek telephone approval from Medicare Australia for up to 2 months' initial therapy. In order to continue the initial course of treatment, a written application must be provided to Medicare Australia following the initial telephone authority approval to confirm the information provided. No more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment will be approved for this follow-up application.

Authority required

Initial treatment of mild to moderately severe Alzheimer's disease. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist). The authority application must include the result of the baseline Mini-Mental State Examination (MMSE). This baseline MMSE must be a score of 10 or more and, if this score is at least 25 points, the result of the baseline Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) must also be specified. An application must be made in writing, but initial supply may be sought on the telephone as indicated in the Note above. For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. For the subsequent written authority application, no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment will be approved. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised; Continuing treatment, following initial therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in cognitive function as measured by an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE), OR a decrease of at least 4 points from baseline on the Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) for patients with an MMSE baseline score of at least 25 points (the initial authority application for continuing treatment must include the relevant result from the MMSE or the ADAS-Cog and must be in writing).

NERVOUS SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
RIVASTIGMINE HYDROGEN TARTRATE—cont.								
Authority required								
<i>Initial treatment of mild to moderately severe Alzheimer's disease of patients with a baseline Mini-Mental State Examination (MMSE) score of 9 or less, who are unable to register a score of 10 or more for reasons other than their Alzheimer's disease, as specified below. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist). Such patients will need to be assessed subsequently using the Clinicians Interview Based Impression of Change (CIBIC) scale. The authority application must include the result of the baseline MMSE and specify to which group(s) (see below) the patient belongs. An application must be made in writing, but initial supply may be sought on the telephone as indicated in the Note above. For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. For the subsequent written authority application, no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment will be approved. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.</i>								
<i>Patients who qualify under this criterion are from 1 or more of the following groups:</i>								
<i>(1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;</i>								
<i>(2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;</i>								
<i>(3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an MMSE test;</i>								
<i>(4) Intellectual (developmental or acquired) disability, eg Down's syndrome;</i>								
<i>(5) Significant sensory impairment despite best correction, which precludes completion of an MMSE test;</i>								
<i>(6) Prominent dysphasia, out of proportion to other cognitive and functional impairment; Continuing treatment, following initial therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in function, based on a rating of "very much improved" or "much improved" on the CIBIC scale, which must be assessed by the same clinician who initiated treatment (the initial authority application for continuing treatment must state the improvement achieved on the CIBIC scale and must be in writing).</i>								
8497F	Capsule 1.5 mg (base)	56	5	..	158.75	28.60	Exelon	NV
8498G	Capsule 3 mg (base)	56	5	..	158.75	28.60	Exelon	NV
8499H	Capsule 4.5 mg (base)	56	5	..	158.75	28.60	Exelon	NV
8500J	Capsule 6 mg (base)	56	5	..	158.75	28.60	Exelon	NV
8563Q	Oral solution 2 mg (base) per mL, 120 mL	‡1	5	..	158.75	28.60	Exelon	NV

OTHER NERVOUS SYSTEM DRUGS

Parasympathomimetics

• Anticholinesterases

PYRIDOSTIGMINE BROMIDE

2724J	Tablet 10 mg	100	5	..	17.84	18.81	Mestinon	VT
1959D	Tablet 60 mg	150	5	..	55.99	28.60	Mestinon	VT
2608G	Tablet 180 mg (modified release)	100	5	..	115.30	28.60	Mestinon Timespan	VT

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Choline esters								
BETHANECHOL CHLORIDE								
1062X	Tablet 10 mg	100	2	..	16.14	17.11	Uro-Carb	HA
Drugs used in addictive disorders								
• Drugs used in nicotine dependence								
BUPROPION HYDROCHLORIDE								
NOTE:								
<i>Only one treatment course per year with no increased maximum quantities or repeats will be authorised.</i>								
Authority required								
<i>Commencement of treatment as short-term adjunctive therapy for nicotine dependence to facilitate the goal of achieving abstinence in patients who have indicated that they are ready to cease smoking and:</i>								
<i>(a) who have entered a comprehensive support and counselling program; or</i>								
<i>(b) who are entering a comprehensive support and counselling program during the consultation at which this authority is requested.</i>								
<i>Details of the program must be specified in the authority application.</i>								
8465M	Tablet 150 mg (sustained release)	30	..	..	90.11	28.60	Zyban	GK
Authority required								
<i>Completion of treatment as short-term adjunctive therapy for nicotine dependence to facilitate the goal of achieving abstinence in patients who have indicated that they are ready to cease smoking and who have entered a comprehensive support and counselling program.</i>								
8710K	Tablet 150 mg (sustained release)	90	..	..	200.11	28.60	Zyban	GK
• Drugs used in alcohol dependence								
ACAMPROSATE CALCIUM								
Authority required								
<i>For use within a comprehensive treatment program for alcohol dependence with the goal of maintaining abstinence.</i>								
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8357W	Tablet 333 mg (enteric coated)	180	1	..	170.27	28.60	Campral	AF
NALTREXONE HYDROCHLORIDE								
CAUTION:								
<i>Naltrexone hydrochloride is contraindicated in patients receiving opioid drugs.</i>								
Authority required								
<i>For use within a comprehensive treatment program for alcohol dependence with the goal of maintaining abstinence.</i>								
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8370M	Tablet 50 mg	30	1	..	167.45	28.60	ReVia	BQ

NERVOUS SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
Other nervous system drugs							
• Other nervous system drugs							
RILUZOLE							
<u>Authority required</u>							
<i>Initial treatment of amyotrophic lateral sclerosis, as diagnosed by a neurologist, in patients aged 75 years or less, with disease duration of 2 years or less and who have at least 60 percent of predicted forced vital capacity within 2 months prior to commencing riluzole therapy and who:</i>							
<i>(1) are ambulatory, and</i>							
<i>(a) have not undergone tracheostomy, and</i>							
<i>(b) have not experienced respiratory failure;</i>							
OR							
<i>(2) are not ambulatory, and</i>							
<i>(a) have not undergone tracheostomy, and</i>							
<i>(b) have not experienced respiratory failure, and</i>							
<i>(c) are either able to use upper limbs or able to swallow.</i>							
<i>The date of diagnosis and the date and results of spirometry (in terms of percent of predicted forced vital capacity) must be supplied with the initial authority application.</i>							
<u>Authority required</u>							
<i>Initial PBS-subsidised treatment of amyotrophic lateral sclerosis for patients receiving treatment with riluzole prior to 1 March 2003 who would have qualified under the initial treatment criteria for PBS subsidy.</i>							
<i>The date of diagnosis, date of commencement of riluzole treatment and the date and results of spirometry (in terms of percent of predicted forced vital capacity) must be supplied with the initial authority application.</i>							
<u>Authority required</u>							
<i>Continuing treatment of amyotrophic lateral sclerosis in patients who have previously been issued with an authority prescription for this drug and who:</i>							
<i>(1) are ambulatory, and</i>							
<i>(a) have not undergone tracheostomy, and</i>							
<i>(b) have not experienced respiratory failure;</i>							
OR							
<i>(2) are not ambulatory, and</i>							
<i>(a) have not undergone tracheostomy, and</i>							
<i>(b) have not experienced respiratory failure, and</i>							
<i>(c) are either able to use upper limbs or able to swallow.</i>							
8664B	Tablet 50 mg	56	5	..	682.23	28.60	Rilutek AV

ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTIPROTOZOALS								
Agents against amoebiasis and other protozoal diseases								
• Nitroimidazole derivatives								
METRONIDAZOLE								
<i>For listings see Generic/Proprietary Index</i>								
METRONIDAZOLE BENZOATE								
<i>For listings see Generic/Proprietary Index</i>								
TINIDAZOLE								
<i>For listings see Generic/Proprietary Index</i>								
• Other agents against amoebiasis and other protozoal diseases								
ATOVAQUONE								
<u>Authority required</u>								
<i>Treatment of mild to moderate Pneumocystis carinii pneumonia in adult patients who are intolerant of trimethoprim/sulfamethoxazole therapy.</i>								
8300W	Oral suspension 750 mg per 5 mL, 210 mL	‡1	..	..	890.15	28.60	Wellvone	GK
1966L	PYRIMETHAMINE Tablet 25 mg	50	..	..	12.96	13.93	Daraprim	GK
Antimalarials								
• Methanolquinolines								
QUININE BISULFATE								
<u>CAUTION:</u>								
<i>Severe thrombocytopenia has been reported with this drug.</i>								
<u>Authority required</u>								
<i>Malaria.</i>								
1972T	Tablet 300 mg	50	2	..	10.13	11.10	Quinbisul	AS
QUININE SULFATE								
<u>CAUTION:</u>								
<i>Severe thrombocytopenia has been reported with this drug.</i>								
<u>Authority required</u>								
<i>Malaria.</i>								
1975Y	Tablet 300 mg	50	2	.. ^B 2.19	10.13 12.32	11.10 ^a 11.10 ^a	Quinsul Quinate	LN AS

ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTHELMINTICS								
Antinematodal agents								
• Benzimidazole derivatives								
ALBENDAZOLE								
<u>Authority required</u>								
<i>Treatment of tapeworm infestation.</i>								
8503M	Tablet 200 mg	6	1	..	26.73	27.70	Zentel	GK
<u>Authority required</u>								
<i>For the treatment of hydatid disease in conjunction with surgery or when a surgical cure cannot be achieved or where surgery cannot be used.</i>								
8459F	Tablet 400 mg	60	2	..	158.75	28.60	Eskazole	GK
• Tetrahydropyrimidine derivatives								
PYRANTEL EMBONATE								
3047J	Tablet 125 mg (base)	6	..	..	6.36	7.33	Anthel 125	AF
3048K	Tablet 250 mg (base)	6	..	..	7.76	8.73	Anthel 250	AF
• Avermectines								
IVERMECTIN								
<u>Authority required</u>								
<i>Onchocerciasis; Strongyloidiasis.</i>								
8359Y	Tablet 3 mg	4	..	..	29.36	28.60	Stromectol	MK
ECTOPARASITICIDES, INCL. SCABICIDES, INSECTICIDES AND REPELLENTS								
Ectoparasiticides, incl. scabicides								
• Pyrethrines, incl. synthetic compounds								
PERMETHRIN								
3054R	Cream 50 mg per g (5%), 30 g	‡1	1	..	14.98	15.95	Lyclear	PC
• Other ectoparasiticides, incl. scabicides								
BENZYL BENZOATE								
1114P	Application 50 g in 200 mL (25%)	‡1	2	..	7.23	8.20	Benzemul	MG

RESPIRATORY SYSTEM

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES								
Adrenergics, inhalants								
• Selective beta-2-adrenoceptor agonists								
EFORMOTEROL FUMARATE DIHYDRATE								
Restricted benefit								
Patients with frequent episodes of asthma who are currently receiving treatment with oral corticosteroids;								
Patients with frequent episodes of asthma who are currently receiving treatment with optimal doses of inhaled corticosteroids.								
8136F	Capsule containing powder for oral inhalation 12 micrograms (for use in Foradile Aerolizer)	60	5	..	35.31	28.60	Foradile	NV
8239P	Powder for oral inhalation in breath actuated device 6 micrograms per dose (60 doses)	‡1	5	..	24.48	25.45	Oxis Turbuhaler	AP
8240Q	Powder for oral inhalation in breath actuated device 12 micrograms per dose (60 doses)	‡1	5	..	34.42	28.60	Oxis Turbuhaler	AP
SALBUTAMOL SULFATE								
1099W	Capsule containing powder for oral inhalation 200 micrograms (base) (for use in Ventolin Rotahaler)	200	5	..	*20.65	21.62	Ventolin Rotacaps	GK
8288F	Oral pressurised inhalation 100 micrograms (base) per dose (200 doses), CFC-free formulation	2	5	..	*16.37	17.34	^a Asmol CFC-free	AL
					^B 0.62	*16.99	^a Epaq	AW
					^B 1.00	*17.37	^a Airomir	MM
							^a Ventolin CFC-free	GK
SALBUTAMOL SULFATE								
Restricted benefit								
Patients unable to achieve co-ordinated use of other metered dose inhalers containing this drug.								
8354Q	Oral pressurised inhalation in breath actuated device 100 micrograms (base) per dose (200 doses), CFC-free formulation	2	5	..	*36.55	28.60	Airomir Autohaler	MM

continued ↗

RESPIRATORY SYSTEM—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
SALBUTAMOL SULFATE—cont.								
Restricted benefit								
<i>Asthma in patients unable to use this drug delivered from an oral pressurised inhalation device via a spacer;</i>								
<i>Chronic obstructive pulmonary disease in patients unable to use this drug delivered from an oral pressurised inhalation device via a spacer.</i>								
2000G	Nebuliser solution single dose units 2.5 mg (base) in 2.5 mL, 30	2	5	..	*23.61	24.58	a Asmol 2.5 uni-dose a Butamol 2.5 a Chem mart Salbutamol a GenRx Salbutamol a Terry White Chemists Salbutamol a PU	AF AW CH GX TW GK
				B2.20	*25.81	24.58	a Ventolin Nebules	GK
2001H	Nebuliser solution single dose units 5 mg (base) in 2.5 mL, 30	2	5	..	*24.69	25.66	a Asmol 5 uni-dose a Butamol 5 a Chem mart Salbutamol a GenRx Salbutamol a Terry White Chemists Salbutamol a Ventolin Nebules	AF AW CH GX TW GK
				B2.20	*26.89	25.66		
2003K	Nebuliser solution 5 mg (base) per mL (0.5%), 30 mL	2	2	..	*11.65	12.62	PU	
SALMETEROL XINAFOATE								
Restricted benefit								
<i>Patients with frequent episodes of asthma who are currently receiving treatment with oral corticosteroids;</i>								
<i>Patients with frequent episodes of asthma who are currently receiving treatment with optimal doses of inhaled corticosteroids.</i>								
3027H	Oral pressurised inhalation 25 micrograms (base) per dose (120 doses)	‡1	5	..	34.42	28.60	Serevent	GK
8141L	Powder for oral inhalation in breath actuated device 50 micrograms (base) per dose (60 doses)	‡1	5	..	34.42	28.60	Serevent Accuhaler	GK

RESPIRATORY SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
TERBUTALINE SULFATE								
1252X	Powder for oral inhalation in breath actuated device 500 micrograms per dose (200 doses)	‡1	5	..	16.03	17.00	Bricanyl Turbuhaler	AP
TERBUTALINE SULFATE								
<u>Restricted benefit</u>								
<i>Asthma in patients unable to use this drug delivered from a breath actuated device; Chronic obstructive pulmonary disease in patients unable to use this drug delivered from a breath actuated device.</i>								
1251W	Nebuliser solution single dose units 5 mg in 2 mL, 30	2	5	..	*26.87	27.84	Bricanyl Respules	AP
• Adrenergics and other drugs for obstructive airway diseases								
BUDESONIDE with EFORMOTEROL FUMARATE DIHYDRATE								
<u>Restricted benefit</u>								
<i>Patients who previously had frequent episodes of asthma while receiving treatment with oral corticosteroids and who have been stabilised on concomitant inhaled eformoterol fumarate dihydrate and budesonide;</i>								
<i>Patients who previously had frequent episodes of asthma while receiving treatment with optimal doses of inhaled corticosteroids and who have been stabilised on concomitant inhaled eformoterol fumarate dihydrate and budesonide.</i>								
8796Y	Powder for oral inhalation in breath actuated device 100 micrograms- 6 micrograms per dose (120 doses)	‡1	5	..	54.25	28.60	Symbicort Turbuhaler 100/6	AP
8625Y	Powder for oral inhalation in breath actuated device 200 micrograms- 6 micrograms per dose (120 doses)	‡1	5	..	58.85	28.60	Symbicort Turbuhaler 200/6	AP
8750M	Powder for oral inhalation in breath actuated devices 400 micrograms- 12 micrograms per dose (60 doses), 2	1	5	..	87.90	28.60	Symbicort Turbuhaler 400/12	AP

RESPIRATORY SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
FLUTICASONE PROPIONATE with SALMETEROL XINAFOATE								
<u>Restricted benefit</u>								
<i>Patients who previously had frequent episodes of asthma while receiving treatment with oral corticosteroids and who have been stabilised on concomitant inhaled salmeterol xinafoate and fluticasone propionate;</i>								
<i>Patients who previously had frequent episodes of asthma while receiving treatment with optimal doses of inhaled corticosteroids and who have been stabilised on concomitant inhaled salmeterol xinafoate and fluticasone propionate.</i>								
8517G	Oral pressurised inhalation 50 micrograms- 25 micrograms (base) per dose (120 doses), CFC-free formulation	‡1	5	..	45.99	28.60	Seretide MDI 50/25	GK
8518H	Oral pressurised inhalation 125 micrograms- 25 micrograms (base) per dose (120 doses), CFC-free formulation	‡1	5	..	59.41	28.60	Seretide MDI 125/25	GK
8519J	Oral pressurised inhalation 250 micrograms- 25 micrograms (base) per dose (120 doses), CFC-free formulation	‡1	5	..	79.55	28.60	Seretide MDI 250/25	GK
8430Q	Powder for oral inhalation in breath actuated device 100 micrograms- 50 micrograms (base) per dose (60 doses)	‡1	5	..	45.99	28.60	Seretide Accuhaler 100/50	GK
8431R	Powder for oral inhalation in breath actuated device 250 micrograms- 50 micrograms (base) per dose (60 doses)	‡1	5	..	59.41	28.60	Seretide Accuhaler 250/50	GK
8432T	Powder for oral inhalation in breath actuated device 500 micrograms- 50 micrograms (base) per dose (60 doses)	‡1	5	..	79.55	28.60	Seretide Accuhaler 500/50	GK
Other drugs for obstructive airway diseases, inhalants								
• Glucocorticoids								
BECLOMETHASONE DIPROPIONATE								
8406K	Oral pressurised inhalation 50 micrograms per dose (200 doses), CFC-free formulation	‡1	5	..	17.48	18.45	Qvar 50	MM
8407L	Oral pressurised inhalation 100 micrograms per dose (200 doses), CFC-free formulation	‡1	5	..	31.48	28.60	Qvar 100	MM

continued ➤

RESPIRATORY SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
BECLOMETHASONE DIPROPIONATE—cont.								
<u>Restricted benefit</u>								
<i>Patients unable to achieve co-ordinated use of other metered dose inhalers containing this drug.</i>								
8408M	Oral pressurised inhalation in breath actuated device 50 micrograms per dose (200 doses), CFC-free formulation	‡1	5	..	25.96	26.93	Qvar 50 Autohaler	MM
8409N	Oral pressurised inhalation in breath actuated device 100 micrograms per dose (200 doses), CFC-free formulation	‡1	5	..	37.08	28.60	Qvar 100 Autohaler	MM
BUDESONIDE								
2070Y	Powder for oral inhalation in breath actuated device 100 micrograms per dose (200 doses)	‡1	5	..	21.47	22.44	Pulmicort Turbuhaler	AP
2071B	Powder for oral inhalation in breath actuated device 200 micrograms per dose (200 doses)	‡1	5	..	29.13	28.60	Pulmicort Turbuhaler	AP
2072C	Powder for oral inhalation in breath actuated device 400 micrograms per dose (200 doses)	‡1	5	..	44.45	28.60	Pulmicort Turbuhaler	AP
<u>BUDESONIDE</u>								
<u>Authority required</u>								
<i>Severe chronic asthma in patients who require long-term steroid therapy and who are unable to use other forms of inhaled steroid therapy.</i>								
2065Q	Nebuliser suspension single dose units 500 micrograms in 2 mL, 30	‡1	5	..	35.84	28.60	Pulmicort Respules	AP
2066R	Nebuliser suspension single dose units 1 mg in 2 mL, 30	‡1	5	..	48.05	28.60	Pulmicort Respules	AP
CICLESONIDE								
8853Y	Oral pressurised inhalation 80 micrograms per dose (120 doses), CFC-free formulation	‡1	5	..	23.92	24.89	Alvesco 80	AH
8854B	Oral pressurised inhalation 160 micrograms per dose (120 doses), CFC-free formulation	‡1	5	..	40.04	28.60	Alvesco 160	AH
FLUTICASONE PROPIONATE								
8516F	Oral pressurised inhalation 50 micrograms per dose (120 doses), CFC-free formulation	‡1	5	..	15.30	16.27	Flixotide Junior	GK

continued ➡

RESPIRATORY SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
FLUTICASONE PROPIONATE—cont.								
8345F	Oral pressurised inhalation 125 micrograms per dose (120 doses), CFC-free formulation	‡1	5	..	28.72	28.60	Flixotide	GK
8346G	Oral pressurised inhalation 250 micrograms per dose (120 doses), CFC-free formulation	‡1	1	..	48.86	28.60	Flixotide	GK
8147T	Powder for oral inhalation in breath actuated device 100 micrograms per dose (60 doses)	‡1	5	..	15.30	16.27	Flixotide Junior Accuhaler	GK
8148W	Powder for oral inhalation in breath actuated device 250 micrograms per dose (60 doses)	‡1	5	..	28.72	28.60	Flixotide Accuhaler	GK
8149X	Powder for oral inhalation in breath actuated device 500 micrograms per dose (60 doses)	‡1	1	..	48.86	28.60	Flixotide Accuhaler	GK
• Anticholinergics								
IPRATROPIUM BROMIDE								
8671J	Oral pressurised inhalation 21 micrograms per dose (200 doses), CFC-free formulation	2	5	..	*39.59	28.60	Atrovent	BY
<u>IPRATROPIUM BROMIDE</u>								
<u>Restricted benefit</u>								
<i>Asthma in patients unable to use this drug delivered from an oral pressurised inhalation device via a spacer;</i>								
<i>Chronic obstructive pulmonary disease in patients unable to use this drug delivered from an oral pressurised inhalation device via a spacer.</i>								
1542E	Nebuliser solution single dose units 250 micrograms (anhydrous) in 1 mL, 30	2	5	..	*45.43	28.60	^a Aeron 250 ^a Apoven 250 ^a Chem mart Ipratropium ^a DBL Ipratropium ^a GenRx Ipratropium ^a Ipratrin ^a Ipravent ^a Terry White Chemists Ipratropium ^a Atrovent	AW DP CH MX GX AF PU TW BY
				^B 0.94	*46.37	28.60		

continued ↗

RESPIRATORY SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
IPRATROPIUM BROMIDE—cont.								
8238N	Nebuliser solution single dose units 500 micrograms (anhydrous) in 1 mL, 30	2	5	..	*52.83	28.60	^a Aeron 500 ^a Apoven 500 ^a Chem mart Ipratropium ^a DBL Ipratropium ^a GenRx Ipratropium ^a Ipratrin Adult ^a Ipravent ^a Terry White Chemists Ipratropium ^a Atrovent Adult	AW DP CH MX GX AF PU TW BY
				^B 0.94	*53.77	28.60		
1541D	Nebuliser solution 250 micrograms (anhydrous) per mL (0.025%), 20 mL	2	2	..	*16.87	17.84	Atrovent	BY
TIOTROPIUM BROMIDE MONOHYDRATE								
Restricted benefit								
For the long-term maintenance treatment of bronchospasm and dyspnoea associated with chronic obstructive pulmonary disease.								
8626B	Capsule containing powder for oral inhalation 18 micrograms (base) (for use in HandiHaler)	30	5	..	77.57	28.60	Spiriva	BY
• Antiallergic agents, excl. corticosteroids								
NEDOCROMIL SODIUM								
8365G	Oral pressurised inhalation 2 mg per dose (112 doses), CFC-free formulation	‡1	5	..	33.70	28.60	Tilade CFC-Free	AV
SODIUM CROMOGLYCATE								
2878L	Capsule containing powder for oral inhalation 20 mg (for use in Intal Spinhaler or Intal Halermatic)	100	5	..	31.51	28.60	Intal Spincaps	DP
1124E	Solution for inhalation 20 mg in 2 mL ampoule	120	3	..	*60.21	28.60	Intal	AV
2872E	Oral pressurised inhalation 1 mg per dose (200 doses)	‡1	5	..	28.10	28.60	Intal	AV
8767K	Oral pressurised inhalation 1 mg per dose (200 doses), CFC-free formulation	‡1	5	..	28.10	28.60	Intal CFC-Free	AV
8334P	Oral pressurised inhalation 5 mg per dose (112 doses), CFC-free formulation	‡1	5	..	33.70	28.60	Intal Forte CFC-Free	AV

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Adrenergics for systemic use								
• Alpha- and beta-adrenoceptor agonists								
ADRENALINE								
1016L	Injection 1 mg in 1 mL (1 in 1,000)	5	1	..	14.76	15.73	AP	
ADRENALINE								
Authority required								
<i>Initial supply for anticipated emergency treatment of acute allergic reactions with anaphylaxis in a patient who has been assessed to be at significant risk of anaphylaxis by, or in consultation with, a clinical immunologist, allergist, paediatrician or respiratory physician. The name of the specialist consulted must be provided at the time of application for initial supply;</i>								
<i>Continuing supply for anticipated emergency treatment of acute allergic reactions with anaphylaxis, where the patient has previously been issued with an authority prescription for this drug.</i>								
NOTE:								
<i>The auto-injector should be provided in the framework of a comprehensive anaphylaxis prevention program and an emergency action plan including training in recognition of the symptoms of anaphylaxis and the use of the auto-injector device. (For further information see the Australasian Society of Clinical Immunology and Allergy website at www.allergy.org.au.)</i>								
8697R	I.M. injection 150 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	96.41	28.60	EpiPen Jr.	CS
8698T	I.M. injection 300 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	96.41	28.60	EpiPen	CS
NOTE:								
<i>Authorities for increased maximum quantities, up to a maximum of 2, may be authorised for children aged less than 17 years where 2 auto-injectors are necessary to ensure 1 is on hand at all times. No increased maximum quantities will be authorised for patients aged 17 years or older.</i>								
No repeats will be issued.								
• Selective beta-2-adrenoceptor agonists								
SALBUTAMOL SULFATE								
1103C	Syrup 2 mg (base) per 5 mL, 150 mL	2	5	..	*12.53	13.50	Ventolin	GK
TERBUTALINE SULFATE								
1034K	Injection 500 micrograms in 1 mL	5	..	..	22.54	23.51	Bricanyl	AP
Other systemic drugs for obstructive airway diseases								
• Xanthines								
THEOPHYLLINE								
CAUTION:								
<i>Because of variable effects of food on absorption of sustained release theophylline preparations, patients stabilised on one brand should not be changed to another without appropriate monitoring.</i>								
1143E	Tablet 125 mg	100	5	..	8.63	9.60	Nuelin	MM

continued

RESPIRATORY SYSTEM —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
THEOPHYLLINE—cont.							
8230E	Tablet 200 mg (sustained release)	100	5	..	10.43	11.40	Nuelin-SR 200 MM
2634P	Tablet 250 mg (sustained release)	100	5	..	11.57	12.54	Nuelin-SR 250 MM
8231F	Tablet 300 mg (sustained release)	100	5	..	12.93	13.90	Nuelin-SR 300 MM
2614N	Syrup 133.3 mg per 25 mL, 500 mL	‡1	5	..	9.10	10.07	Nuelin MM

- **Leukotriene receptor antagonists**

MONTELUKAST SODIUM**NOTE:**

Montelukast sodium is not PBS-subsidised for use in children with moderate or severe asthma. It is not intended as an alternative for children who require a corticosteroid as a preventer medication. Montelukast sodium is not PBS-subsidised for use in combination with other preventer medications. PBS subsidy for montelukast sodium will therefore cease for patients who require a preventer medication in addition to montelukast sodium.

NOTE:

No applications for increased maximum quantities and/or repeats will be authorised.

Authority required

First-line preventer medication, as the single preventer agent for children aged 2 to 5 years with frequent episodic or mild persistent asthma, as an alternative to sodium cromoglycate or nedocromil sodium.

8627C	Chewable tablet 4 mg (base)	28	5	..	46.81	28.60	Singulair	MK

Authority required

First-line preventer medication, as the single preventer agent for children aged 6 to 14 years with frequent episodic or mild persistent asthma, as an alternative to sodium cromoglycate or nedocromil sodium.

8628D	Chewable tablet 5 mg (base)	28	5	..	46.81	28.60	Singulair	MK

COUGH AND COLD PREPARATIONS

Expectorants, excl. combinations with cough suppressants

- **Mucolytics**

ACETYLCYSTEINE**Restricted benefit**

Bronchiectasis;

Cystic fibrosis.

8747J	Sterile inhalation solution 200 mg per mL (20%), 5 mL	30	3	..	*149.95	28.60	Mucomyst	BQ

Cough suppressants, excl. combinations with expectorants

- **Opium alkaloids and derivatives**

CODEINE PHOSPHATE

1214X	Tablet 30 mg	20	..	..	10.53	11.50	FM

ANTI-HISTAMINES FOR SYSTEMIC USE

Antihistamines for systemic use

- **Phenothiazine derivatives**

PROMETHAZINE HYDROCHLORIDE

1948M	Injection 50 mg in 2 mL	10	..	..	*18.29	19.26	MX

SENSORY ORGANS

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
OPHTHALMOLOGICALS								
Antiinfectives								
• Antibiotics								
AZITHROMYCIN								
<u>Restricted benefit</u>								
<i>Trachoma.</i>								
<u>NOTE:</u>								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8336R	Tablet 500 mg	2	2	..	22.02	22.99	Zithromax	PF
8201P	Powder for oral suspension 200 mg per 5 mL, 15 mL	‡1	..	..	#21.46	22.84	Zithromax	PF
CHLORAMPHENICOL								
2360F	Eye drops 5 mg per mL (0.5%), 10 mL	‡1	2	..	7.60	8.57	Chloromycetin Chlorsig	PF SI
1171P	Eye ointment 10 mg per g (1%), 4 g	‡1	..	..	7.27	8.24	Chloromycetin Chlorsig	PF SI
GENTAMICIN SULFATE								
<u>Restricted benefit</u>								
<i>Invasive ocular infection;</i>								
<i>Perioperative use in ophthalmic surgery;</i>								
<i>Suspected pseudomonal eye infection.</i>								
1441W	Eye drops 3 mg (base) per mL (0.3%), 5 mL	‡1	2	..	16.33	17.30	Genoptic	AG
POLYMYXIN B SULFATE with BACITRACIN and NEOMYCIN SULFATE								
1910M	Eye ointment 5,000 units- 400 units-5 mg per g, 4 g	‡1	..	..	7.46	8.43	Neosporin	PC
TOBRAMYCIN								
<u>Restricted benefit</u>								
<i>Invasive ocular infection;</i>								
<i>Perioperative use in ophthalmic surgery;</i>								
<i>Suspected pseudomonal eye infection.</i>								
2328M	Eye drops 3 mg per mL (0.3%), 5 mL	‡1	2	..	16.33	17.30	Tobrex	AQ
2329N	Eye ointment 3 mg per g (0.3%), 3.5 g	‡1	..	..	19.35	20.32	Tobrex	AQ
• Sulfonamides								
SULFACETAMIDE SODIUM								
2063N	Eye drops 100 mg per mL (10%), 15 mL	‡1	2	..	10.36	11.33	Bleph 10	AG
• Antivirals								
ACICLOVIR								
<u>Restricted benefit</u>								
<i>Herpes simplex keratitis.</i>								
1002R	Eye ointment 30 mg per g (3%), 4.5 g	‡1	..	..	23.01	23.98	Zovirax	GK

SENSORY ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Other antiinfectives CIPROFLOXACIN <u>Authority required</u> Bacterial keratitis.								
1217C	Eye drops 3 mg per mL (0.3%), 5 mL	2	..	.. 2.48	*27.91 *30.39	28.60 ^a 28.60 ^a	CiloQuin Ciloxan	IQ AQ
OFLOXACIN <u>Authority required</u> Bacterial keratitis.								
8383F	Eye drops 3 mg per mL (0.3%), 5 mL	2	..	..	*27.91	28.60	Ocuflox	AG
Antiinflammatory agents • Corticosteroids, plain								
DEXAMETHASONE								
1288T	Eye drops 1 mg per mL (0.1%), 5 mL	‡1	2	..	8.33	9.30	Maxidex	AQ
FLUOROMETHOLONE								
1204J	Eye drops 1 mg per mL (0.1%), 5 mL	‡1	5	..	8.33	9.30	Flucon FML Liquifilm	AQ AG
FLUOROMETHOLONE ACETATE								
1438Q	Eye drops 1 mg per mL (0.1%), 5 mL	‡1	2	..	8.33	9.30	Flarex	AQ
HYDROCORTISONE								
1489J	Eye drops 5 mg per mL (0.5%), 10 mL	‡1	5	..	10.36	11.33	Hycor	SI
1492M	Eye drops 10 mg per mL (1%), 10 mL	‡1	5	..	10.70	11.67	Hycor	SI
HYDROCORTISONE ACETATE								
1497T	Eye ointment 5 mg per g (0.5%), 5 g	‡1	..	..	10.15	11.12	Hycor	SI
2441L	Eye ointment 10 mg per g (1%), 5 g	‡1	..	..	10.35	11.32	Hycor	SI
• Corticosteroids and mydriatics in combination PREDNISOLONE ACETATE with PHENYLEPHRINE HYDROCHLORIDE <u>Restricted benefit</u> Corneal grafts; Uveitis.								
3112T	Eye drops 10 mg-1.2 mg per mL (1%-0.12%), 10 mL	‡1	2	..	19.49	20.46	Prednefrin Forte	AG
• Antiinflammatory agents, non-steroids DICLOFENAC SODIUM <u>Restricted benefit</u> Inhibition of intraoperative miosis.								
8329J	Eye drops 1 mg per mL (0.1%), 5 mL	‡1	..	..	12.27	13.24	Voltaren Ophtha	NV

SENSORY ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
FLURBIPROFEN SODIUM								
8699W	Eye drops 300 micrograms per mL (0.03%), single dose units 0.4 mL, 5	1	..	..	12.27	13.24	Ocufen	AG
Antiglaucoma preparations and miotics								
• Sympathomimetics in glaucoma therapy								
APRACLOPIDINE HYDROCHLORIDE								
Restricted benefit								
<i>Short-term reduction of intra-ocular pressure in patients already on maximally tolerated anti-glaucoma therapy.</i>								
8083K	Eye drops 5 mg (base) per mL (0.5%), 10 mL	‡1	2	..	35.48	28.60	Iopidine 0.5%	AQ
BRIMONIDINE TARTRATE								
8351M	Eye drops 2 mg per mL (0.2%), 5 mL	‡1	5	..	18.83	19.80	^a Enidin	PE
				^B 1.50	20.33	19.80	^a Alphagan	AG
BRIMONIDINE TARTRATE with TIMOLOL MALEATE								
Restricted benefit								
<i>Reduction of elevated intra-ocular pressure in patients with open-angle glaucoma who are not adequately controlled with timolol maleate 5 mg (base) per mL (0.5%) eye drops; Reduction of elevated intra-ocular pressure in patients with ocular hypertension who are not adequately controlled with timolol maleate 5 mg (base) per mL (0.5%) eye drops.</i>								
8826M	Eye drops 2 mg-5 mg (base) per mL (0.2%-0.5%), 5 mL	‡1	5	..	27.00	27.97	Combigan	AG
DIPIVEFRINE HYDROCHLORIDE								
1351D	Eye drops 1 mg per mL (0.1%), 10 mL	‡1	5	..	20.56	21.53	Propine	AG
• Parasympathomimetics								
CARBACHOL								
2535K	Eye drops 15 mg per mL (1.5%), 15 mL	‡1	5	..	23.63	24.60	Isopto Carbachol	AQ
2536L	Eye drops 30 mg per mL (3%), 15 mL	‡1	5	..	24.18	25.15	Isopto Carbachol	AQ
PILOCARPINE HYDROCHLORIDE								
2778F	Eye drops 5 mg per mL (0.5%), 15 mL	‡1	5	..	10.45	11.42	^a Pilopt	PE
				^B 2.00	12.45	11.42	^a P.V. Carpine	AG
2595N	Eye drops 10 mg per mL (1%), 15 mL	‡1	5	..	10.45	11.42	Isopto Carpine	AQ
				^B 2.00	12.45	11.42	^a Pilopt	PE
							^a P.V. Carpine	AG
2596P	Eye drops 20 mg per mL (2%), 15 mL	‡1	5	..	11.60	12.57	Isopto Carpine	AQ
				^B 2.12	13.72	12.57	^a Pilopt	PE
							^a P.V. Carpine	AG
2597Q	Eye drops 30 mg per mL (3%), 15 mL	‡1	5	..	14.25	15.22	Isopto Carpine	AQ
				^B 2.60	16.85	15.22	^a Pilopt	PE
							^a P.V. Carpine	AG

continued ➤

SENSORY ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PILOCARPINE HYDROCHLORIDE—cont.								
2598R	Eye drops 40 mg per mL (4%), 15 mL	‡1	5	..	14.25	15.22	Isopto Carpine	AQ
							^a Pilopt	PE
				^B 2.60	16.85	15.22	^a P.V. Carpine	AG
2779G	Eye drops 60 mg per mL (6%), 15 mL	‡1	5	..	18.45	19.42	^a Pilopt	PE
				^B 1.98	20.43	19.42	^a P.V. Carpine	AG
• Carbonic anhydrase inhibitors								
ACETAZOLAMIDE								
1004W	Tablet 250 mg	100	3	..	21.91	22.88	Diamox	SI
BRINZOLAMIDE								
8483L	Eye drops 10 mg per mL (1%), 5 mL	‡1	5	..	18.83	19.80	^a BrinzoQuin	IQ
				^B 1.00	19.83	19.80	^a Azopt	AQ
DORZOLAMIDE HYDROCHLORIDE								
8488R	Eye drops 20 mg (base) per mL (2%), 5 mL	‡1	5	..	18.83	19.80	Trusopt	MK
DORZOLAMIDE HYDROCHLORIDE with TIMOLOL MALEATE								
<u>Restricted benefit</u>								
<i>Reduction of elevated intra-ocular pressure in patients with open-angle glaucoma who are not adequately controlled with timolol maleate 5 mg (base) per mL (0.5%) eye drops;</i>								
<i>Reduction of elevated intra-ocular pressure in patients with ocular hypertension who are not adequately controlled with timolol maleate 5 mg (base) per mL (0.5%) eye drops.</i>								
8567X	Eye drops 20 mg (base)-5 mg (base) per mL (2%-0.5%), 5 mL	‡1	5	..	27.00	27.97	Cosopt	MK
• Beta blocking agents								
BETAXOLOL HYDROCHLORIDE								
2811Y	Eye drops, suspension, 2.5 mg (base) per mL (0.25%), 5 mL	‡1	5	..	12.92	13.89	Betoptic S	AQ
2825Q	Eye drops, solution, 5 mg (base) per mL (0.5%), 5 mL	‡1	5	..	12.92	13.89	^a BetoQuin	IQ
				^B 2.01	14.93	13.89	^a Betoptic	AQ
LEVOBUNOLOL HYDROCHLORIDE								
1819R	Eye drops 2.5 mg per mL (0.25%), 5 mL	‡1	5	..	11.83	12.80	Betagan	AG
TIMOLOL MALEATE								
8803H	Eye gel 1 mg (base) per g (0.1%), 5 g	‡1	5	..	11.83	12.80	Nyogel	NV
1278G	Eye drops 2.5 mg (base) per mL (0.25%), 5 mL	‡1	5	..	11.83	12.80	^a Tenopt	SI
				^B 1.09	12.92	12.80	^a Timoptol	FR
1279H	Eye drops 5 mg (base) per mL (0.5%), 5 mL	‡1	5	..	12.92	13.89	^a Tenopt	SI
				^B 1.15	14.07	13.89	^a Timoptol	FR
1925H	Eye drops (gellan gum solution) 2.5 mg (base) per mL (0.25%), 2.5 mL	‡1	5	..	11.83	12.80	Timoptol XE	MK
1926J	Eye drops (gellan gum solution) 5 mg (base) per mL (0.5%), 2.5 mL	‡1	5	..	12.92	13.89	Timoptol XE	MK

SENSORY ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Prostaglandin analogues								
BIMATOPROST								
8620Q	Eye drops 300 micrograms per mL (0.03%), 3 mL	‡1	5	..	34.21	28.60	Lumigan	AG
LATANOPROST								
8243W	Eye drops 50 micrograms per mL (0.005%), 2.5 mL	‡1	5	..	34.21	28.60	Xalatan	PU
TRAVOPROST								
8597L	Eye drops 40 micrograms per mL (0.004%), 2.5 mL	‡1	5	..	34.21	28.60	Travatan	AQ
Mydriatics and cycloplegics								
• Anticholinergics								
ATROPINE SULFATE								
1093M	Eye drops 10 mg per mL (1%), 15 mL	‡1	2	..	9.49	10.46	Atropt	SI
HOMATROPINE HYDROBROMIDE								
2541R	Eye drops 20 mg per mL (2%), 15 mL	‡1	2	..	14.20	15.17	Isopto Homatropine	AQ
2542T	Eye drops 50 mg per mL (5%), 15 mL	‡1	2	..	18.35	19.32	Isopto Homatropine	AQ
Decongestants and antiallergics								
• Other antiallergics								
LODOXAMIDE TROMETAMOL								
<u>Authority required</u>								
<i>Vernal kerato-conjunctivitis.</i>								
8268E	Eye drops 1 mg (base) per mL (0.1%), 10 mL	‡1	5	..	14.29	15.26	Lomide	AQ
SODIUM CROMOGLYCATE								
<u>Authority required</u>								
<i>Vernal kerato-conjunctivitis.</i>								
1127H	Eye drops 20 mg per mL (2%), 10 mL	‡1	5	..	14.29	15.26	Opticrom	AV
Other ophthalmologicals								
• Other ophthalmologicals								
CARBOMER 974								
<u>Authority required</u>								
<i>Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.</i>								
8514D	Ocular lubricating gel 3 mg per g (0.3%), single dose units 0.5 g, 30	3	5	..	*34.06	28.60	Poly Gel	AQ

SENSORY ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CARBOMER 980								
<u>Restricted benefit</u>								
<i>Severe dry eye syndrome, including Sjogren's syndrome.</i>								
8384G	Ocular lubricating gel 2 mg per g (0.2%), 10 g	‡1	5	..	8.89	9.86	GelTears	BU
							^a PAA	CV
				^B 1.00	9.89	9.86	^a Viscotears Liquid Gel	NV
<u>Authority required</u>								
<i>Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.</i>								
8578L	Eye drops 2 mg per g (0.2%), single dose units 0.6 mL, 30	3	5	..	*34.06	28.60	Viscotears	NV
CARMELLOSE SODIUM								
<u>Restricted benefit</u>								
<i>Severe dry eye syndrome, including Sjogren's syndrome.</i>								
8548X	Eye drops 5 mg per mL (0.5%), 15 mL	‡1	5	..	8.89	9.86	Refresh Tears Plus	AG
8593G	Eye drops 10 mg per mL (1%), 15 mL	‡1	5	..	8.89	9.86	Refresh Liquigel	AG
<u>Authority required</u>								
<i>Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.</i>								
8823J	Eye drops 2.5 mg per mL (0.25%), single dose units 0.6 mL, 24	4	5	..	*35.99	28.60	TheraTears	CX
2338C	Eye drops 5 mg per mL (0.5%), single dose units 0.4 mL, 30	3	5	..	*34.06	28.60	Cellufresh	AG
2324H	Eye drops 10 mg per mL (1%), single dose units 0.4 mL, 30	3	5	..	*34.06	28.60	Celluvisc	AG
8824K	Ocular lubricating gel 10 mg per mL (1%), single dose units 0.6 mL, 28	3	5	..	*32.11	28.60	TheraTears	CX
HYDROXYPROPYLCELLULOSE								
<u>Restricted benefit</u>								
<i>Severe dry eye syndrome unresponsive to artificial tear solutions.</i>								
1522D	Ophthalmic inserts 5 mg, 60	‡1	5	..	37.75	28.60	Lacrisert	SI

SENSORY ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
<u>HYPROMELLOSE</u>								
<u>Restricted benefit</u>								
<i>Severe dry eye syndrome, including Sjogren's syndrome.</i>								
8287E	Eye drops 3 mg per mL (0.3%), 15 mL (contains sodium perborate as preservative)	‡1	5	..	8.89	9.86	^a In a Wink Moisturising	CV
				^B 1.86	10.75	9.86	^a Genteal	NV
2956N	Eye drops 5 mg per mL (0.5%), 15 mL	‡1	5	..	8.89	9.86	Isopto Tears Methopt	AQ SI
<u>HYPROMELLOSE with CARBOMER 980</u>								
<u>Restricted benefit</u>								
<i>Severe dry eye syndrome, including Sjogren's syndrome.</i>								
8564R	Ocular lubricating gel 3 mg- 2 mg per g (0.3%-0.2%), 10 g	‡1	5	..	8.89	9.86	^a HPMC PAA	CV
				^B 1.86	10.75	9.86	^a Genteal gel	NV
<u>HYPROMELLOSE with DEXTRAN</u>								
<u>Restricted benefit</u>								
<i>Severe dry eye syndrome, including Sjogren's syndrome.</i>								
1509K	Eye drops 3 mg-1 mg per mL (0.3%-0.1%), 15 mL	‡1	5	..	8.89	9.86	^a Poly-Tears	IQ
				^B 1.67	10.56	9.86	^a Tears Naturale	AQ
<u>Authority required</u>								
<i>Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.</i>								
8299T	Eye drops 3 mg-1 mg per mL (0.3%-0.1%), single dose units 0.4 mL, 28	3	5	..	*32.11	28.60	Bion Tears	AQ
<u>PARAFFIN</u>								
1754H	Compound eye ointment 3.5 g	2	5	..	*18.09	19.06	^a Poly Visc	IQ
				^B 2.00	*20.09	19.06	^a Duratears	AQ
1750D	Pack containing 2 tubes compound eye ointment 3.5 g	‡1	5	..	18.09	19.06	^a Ircal	PE
				^B 2.01	20.10	19.06	^a Poly Visc	IQ
						19.06	^a Lacri-Lube	AG
<u>POLYETHYLENE GLYCOL 400 with PROPYLENE GLYCOL</u>								
<u>Restricted benefit</u>								
<i>Severe dry eye syndrome, including Sjogren's syndrome.</i>								
8676P	Eye drops 4 mg-3 mg per mL (0.4%-0.3%), 15 mL	‡1	5	..	8.89	9.86	Systane	AQ

SENSORY ORGANS —cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
POLYVINYL ALCOHOL								
Restricted benefit								
<i>Severe dry eye syndrome, including Sjogren's syndrome.</i>								
2682E	Eye drops 14 mg per mL (1.4%), 15 mL	‡1	5	.. B1.70	8.89 10.59	9.86 ^a 9.86 ^a	PVA Tears Liquifilm Tears	PE AG
8831T	Eye drops 14 mg per mL (1.4%), 15 mL (contains sodium chlorite/hydrogen peroxide as preservative)	‡1	5	..	8.89	9.86	Vistil	AE
2681D	Eye drops 30 mg per mL (3%), 15 mL	‡1	5	.. B5.98	8.89 14.87	9.86 ^a 9.86 ^a	PVA Forte Liquifilm Forte	PE AG
8832W	Eye drops 30 mg per mL (3%), 15 mL (contains sodium chlorite/hydrogen peroxide as preservative)	‡1	5	..	8.89	9.86	Vistil Forte	AE
OTOLOGICALS								
Antiinfectives								
• Antiinfectives								
CHLORAMPHENICOL								
1172Q	Ear drops (aqueous) 5 mg per mL (0.5%), 5 mL	‡1	2	..	9.34	10.31	Chloromycetin	PF
NEOMYCIN UNDECENOATE with BACITRACIN ZINC								
2296W	Ear ointment 12 mg (3.5 mg base)-400 units per g, 10 g	‡1	..	..	6.95	7.92	Nemdyn	HA
Corticosteroids and antiinfectives in combination								
• Corticosteroids and antiinfectives in combination								
DEXAMETHASONE with FRAMYCETIN SULFATE and GRAMICIDIN								
2781J	Ear drops 500 micrograms-5 mg- 50 micrograms per mL, 8 mL	‡1	2	.. B1.80	7.10 8.90	8.07 ^a 8.07 ^a	Otodex Sofradex	QM AV
TRIAMCINOLONE ACETONIDE with NEOMYCIN SULFATE, GRAMICIDIN and NYSTATIN								
2971J	Ear drops 1 mg-2.5 mg (base)- 250 micrograms-100,000 units per g (0.1%-0.25%-0.025%- 100,000 units per g), 7.5 mL	‡1	2	.. B1.10	7.10 8.20	8.07 ^a 8.07 ^a	Otocomb Otic Kenacomb Otic	BC BQ
2974M	Ear ointment 1 mg-2.5 mg (base)- 250 micrograms-100,000 units per g (0.1%-0.25%-0.025%- 100,000 units per g), 5 g	‡1	2	.. B1.10	6.62 7.72	7.59 ^a 7.59 ^a	Otocomb Otic Kenacomb Otic	BC BQ
OPHTHALMOLOGICAL AND OTOLOGICAL PREPARATIONS								
Antiinfectives								
• Antiinfectives								
FRAMYCETIN SULFATE								
1440T	Eye and ear drops 5 mg per mL (0.5%), 8 mL	‡1	2	..	7.63	8.60	Soframycin	AV

VARIOUS

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ALLERGENS								
Allergens								
• Allergen extracts								
INSECT ALLERGEN EXTRACT— HONEY BEE VENOM								
2886X	Injection set containing 550 micrograms	1	..	..	201.65	28.60	Albey Bee Venom	TH
INSECT ALLERGEN EXTRACT— PAPER WASP VENOM								
NOTE: Paper wasp venom is not European wasp venom.								
2918N	Injection set containing 550 micrograms	1	..	..	201.65	28.60	Albey Paper Wasp Venom	TH
INSECT ALLERGEN EXTRACT— YELLOW JACKET VENOM								
2883R	Injection set containing 550 micrograms	1	..	..	201.65	28.60	Albey Yellow Jacket Venom	TH
ALL OTHER THERAPEUTIC PRODUCTS								
All other therapeutic products								
• Antidotes								
NALOXONE HYDROCHLORIDE								
1751E	Injection 400 micrograms in 1 mL	1	..	..	22.04	23.01	Naloxone Min-I-Jet	CS
1752F	Injection 800 micrograms in 2 mL	1	..	..	23.14	24.11	Naloxone Min-I-Jet	CS
1753G	Injection 2 mg in 5 mL	1	..	..	34.15	28.60	Naloxone Min-I-Jet	CS
• Detoxifying agents for antineoplastic treatment								
CALCIUM FOLINATE								
8740B	Injection equivalent to 50 mg folinic acid in 5 mL	5	5	..	*183.50	28.60	^a Leucovorin Calcium	MX
							^a Leucovorin Calcium	PF
8812T	Injection equivalent to 100 mg folinic acid in 10 mL	10	1	..	322.75	28.60	Leucovorin Calcium	PF
CALCIUM FOLINATE								
Restricted benefit								
Antidote to folic acid antagonists.								
2308L	Tablet equivalent to 15 mg folinic acid	10	..	..	98.35	28.60	Leucovorin Calcium	MX

VARIOUS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MESNA								
Restricted benefit								
<i>Adjunctive therapy for use with ifosfamide or high dose cyclophosphamide.</i>								
8078E	Solution for I.V. injection 400 mg in 4 mL	15	5	..	73.36	28.60	Uromitexan	BX
8079F	Solution for I.V. injection 1 g in 10 mL	15	5	..	161.57	28.60	Uromitexan	BX
• Drugs for treatment of hypercalcemia								
SODIUM ACID PHOSPHATE								
Authority required								
<i>Familial hypophosphataemia; Hypercalcaemia; Hypophosphataemic rickets; Vitamin D-resistant rickets.</i>								
2946C	Compound effervescent tablet containing elemental phosphorus 500 mg, sodium 469 mg (20.4 mmol), potassium 123 mg (3.1 mmol)	100	5	..	82.47	28.60	Phosphate Sandoz	NV
DIAGNOSTIC AGENTS								
Urine tests								
• Urine tests								
COPPER SULFATE								
1228P	Diagnostic compound tablets, 36	2	3	..	*71.99	28.60	Clinitest	BN
GLUCOSE INDICATOR—URINE								
2352T	Reagent strips, 50	2	2	..	*17.99	18.96	Clinistix	BN
3104J	Reagent strips, 50	2	2	..	*17.03	18.00	Diastix	BN
GLUCOSE and KETONE INDICATOR—URINE								
3106L	Reagent strips, 50	2	2	..	*15.51	16.48	Keto-Diabur- Test 5000	RD
3107M	Reagent strips, 50	2	2	..	*15.61	16.58	Keto-Diastix	BN
Other diagnostic agents								
• Tests for diabetes								
GLUCOSE INDICATOR—BLOOD								
2891E	Electrode strips, 50	2	5	..	*52.77	28.60	Advantage II	RD
8749L	Electrode strips, 50	2	5	..	*52.77	28.60	GlucoCare	DR
8766J	Electrode strips, 50	2	5	..	*52.77	28.60	GlucoCare Super Sensor	DR
8557J	Electrode strips, 50	2	5	..	*52.77	28.60	GlucoMen Sensor	GD
8723D	Electrode strips, 50	2	5	..	*52.77	28.60	Omnitest EZ	BR
8825L	Electrode strips, 50	2	5	..	*52.77	28.60	TrueTrack	DR

continued ➤

VARIOUS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
GLUCOSE INDICATOR—BLOOD—cont.								
8176H	Discs containing electrode sensors, 10 sensors per disc, 5	2	5	..	*52.77	28.60	Ascensia Glucodisc	BN
8522M	Electrode strips, 100	‡1	5	..	52.77	28.60	Optium glucose	MS
8573F	Electrode strips, 100	‡1	5	..	52.77	28.60	SofTact	MS
8608C	Electrode strips, 100	‡1	5	..	52.77	28.60	TrueSense	MS
8190C	Reagent strips, 50	2	5	..	*52.77	28.60	Accu-Chek Active	RD
8739Y	Reagent strips, 50	2	5	..	*52.77	28.60	Accu-Chek Go	RD
8806L	Reagent strips, 51	2	5	..	*52.77	28.60	Accu-Chek Integra	RD
2890D	Reagent strips, 50	2	5	..	*52.77	28.60	Betachek	NA
8053W	Reagent strips, 50	2	5	..	*52.77	28.60	Betachek MERIDIAN	NA
8759B	Reagent strips, 50	2	5	..	*52.77	28.60	CareSens	LB
2914J	Reagent strips, 50	2	5	..	*44.53	28.60	Glucoflex-R	TC
2917M	Reagent strips, 50	2	5	..	*44.53	28.60	Glucostix	BN
8795X	Reagent strips, 50	2	5	..	*52.77	28.60	SensoCard	PX

GLUCOSE INDICATOR—BLOOD

Authority required*Patients who have previously received this product as a pharmaceutical benefit;**Patients who have purchased a meter to be used with this product prior to 1 August 2003.*

8634K	Electrode strips, 50	2	5	..	*60.43	28.60	Ascensia Elite	BN
2926B	Electrode strips, 100	‡1	5	..	54.47	28.60	Precision Plus	MS

GENERAL NUTRIENTS

Other nutrients• **Other nutrients****TRIGLYCERIDES, MEDIUM CHAIN****NOTE:***No applications for increased maximum quantities and/or repeats will be authorised.***Authority required***Chylous ascites;**Chylothorax;**Fat malabsorption due to liver disease, short gut syndrome, cystic fibrosis and gastrointestinal disorders;**Hyperlipoproteinaemia type 1;**Intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect, requiring a ketogenic diet;**Long chain fatty acid oxidation disorders.*

3128P	Oil 500 mL	2	5	..	*51.87	28.60	MCT Oil	SB

VARIOUS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Fat/carbohydrates/proteins/minerals/vitamins, combinations								
AMINO ACIDS—SYNTHETIC, FORMULA								
<u>Authority required</u>								
<i>Initial treatment, for up to 3 months, for combined intolerance (not infant colic) to cows' milk protein, soy protein and protein hydrolysate formulae in children up to the age of 2 years. Combined intolerance is demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula. The date of birth of the patient must be included in the authority application.</i>								
<u>NOTE:</u>								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8574G	Compound powder 400 g	8	5	..	*370.75	28.60	EleCare	AB
8443J	Compound powder 400 g	8	5	..	*370.75	28.60	Neocate	SB
8754R	Compound powder 400 g	8	5	..	*370.75	28.60	Neocate Advance	SB

Authority required

Continuing treatment for combined intolerance (not infant colic) to cows' milk protein, soy protein and protein hydrolysate formulae in children up to the age of 2 years, where the child has been assessed by a suitably qualified allergist or paediatrician. The date of birth of the patient must be included in the authority application;

Treatment for combined intolerance (not infant colic) to cows' milk protein, soy protein and protein hydrolysate formulae in children aged 2 years and over, where the child is assessed by a suitably qualified allergist or paediatrician at intervals not greater than 6 months. The date of birth of the patient must be included in the authority application;

Severe intestinal malabsorption including short bowel syndrome where protein hydrolysate formulae have failed;

Severe intestinal malabsorption including short bowel syndrome where the patient has been receiving parenteral nutrition.

NOTE:

Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

8575H	Compound powder 400 g	8	5	..	*370.75	28.60	EleCare	AB
3066J	Compound powder 400 g	8	5	..	*370.75	28.60	Neocate	SB
8755T	Compound powder 400 g	8	5	..	*370.75	28.60	Neocate Advance	SB

VARIOUS—cont.

Code	Name, Restriction, Manner of administration and form	Max. No. of			Dispensed	Maximum	Proprietary Name and Manufacturer
		Qty	Rpts	Premium	Price for Max. Qty \$	Recordable Value for Safety Net \$	
PROTEIN HYDROLYSATE FORMULA with MEDIUM CHAIN TRIGLYCERIDES							
NOTE:							
No applications for increased maximum quantities and/or repeats will be authorised.							
Authority required							
<i>Initial treatment, for up to 3 months, for intolerance (not infant colic) to both cows' milk protein and soy protein in children up to the age of 2 years. Intolerance is demonstrated when the child has failed to respond to a strict cows' milk protein free diet with a soy protein as the principal formula. The date of birth of the patient must be included in the authority application;</i>							
<i>Continuing treatment for intolerance (not infant colic) to both cows' milk protein and soy protein in children up to the age of 2 years, where clinical improvement has been demonstrated with the protein hydrolysate formula with medium chain triglycerides. The date of birth of the patient must be included in the authority application;</i>							
<i>Continuing treatment for intolerance (not infant colic) to both cows' milk protein and soy protein in children aged 2 years and over, where the child has been assessed by a suitably qualified allergist or paediatrician. The date of birth of the patient must be included in the authority application;</i>							
<i>Biliary atresia;</i>							
<i>Chronic liver failure with fat malabsorption;</i>							
<i>Chylous ascites;</i>							
<i>Chylothorax;</i>							
<i>Cystic fibrosis;</i>							
<i>Enterokinase deficiency;</i>							
<i>Proven fat malabsorption;</i>							
<i>Severe diarrhoea of greater than 2 weeks' duration in infants under the age of 4 months. The date of birth of the patient must be included in the authority application;</i>							
<i>Severe intestinal malabsorption including short bowel syndrome.</i>							
2676W	Compound powder 400 g	8	5	..	*99.71	28.60	Alfaré NT
Authority required							
<i>Initial treatment, for up to 3 months, for intolerance (not infant colic) to both cows' milk protein and soy protein in children up to the age of 2 years. Intolerance is demonstrated when the child has failed to respond to a strict cows' milk protein free diet with a soy protein as the principal formula. The date of birth of the patient must be included in the authority application;</i>							
<i>Continuing treatment for intolerance (not infant colic) to both cows' milk protein and soy protein in children up to the age of 2 years, where clinical improvement has been demonstrated with the protein hydrolysate formula with medium chain triglycerides. The date of birth of the patient must be included in the authority application;</i>							
<i>Continuing treatment for intolerance (not infant colic) to both cows' milk protein and soy protein in children aged 2 years and over, where the child has been assessed by a suitably qualified allergist or paediatrician. The date of birth of the patient must be included in the authority application;</i>							
<i>Biliary atresia;</i>							
<i>Chronic liver failure with fat malabsorption;</i>							
<i>Chylous ascites;</i>							
<i>Cystic fibrosis;</i>							
<i>Enterokinase deficiency;</i>							
<i>Proven fat malabsorption;</i>							
<i>Severe diarrhoea of greater than 2 weeks' duration in infants under the age of 4 months. The date of birth of the patient must be included in the authority application;</i>							
<i>Severe intestinal malabsorption including short bowel syndrome.</i>							
8259Q	Compound powder 450 g	8	5	..	*111.55	28.60	Pepti-Junior NU

VARIOUS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
TRIGLYCERIDES—MEDIUM CHAIN, FORMULA								
NOTE: <i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
Restricted benefit								
<i>Chylous ascites;</i>								
<i>Chylothorax;</i>								
<i>Fat malabsorption due to liver disease, short gut syndrome, cystic fibrosis and gastrointestinal disorders;</i>								
<i>Hyperlipoproteinaemia type 1;</i>								
<i>Long chain fatty acid oxidation disorders.</i>								
NOTE: <i>Monogen is not indicated for the treatment of intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect requiring a ketogenic diet.</i>								
8478F	Compound powder 400 g	8	5	..	*320.67	28.60	Monogen	SB
Restricted benefit								
<i>Chylous ascites;</i>								
<i>Chylothorax;</i>								
<i>Fat malabsorption due to liver disease, short gut syndrome, cystic fibrosis and gastrointestinal disorders.</i>								
NOTE: <i>Caprilon is not indicated for the treatment of intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect requiring a ketogenic diet, long chain fatty acid oxidation disorders or hyperlipoproteinaemia type 1.</i>								
8629E	Compound powder 420 g	8	5	..	*349.79	28.60	Caprilon	SB
• Milk substitutes								
MILK POWDER—LACTOSE FREE FORMULA								
Authority required <i>Acute lactose intolerance in infants up to the age of 12 months. The date of birth of the patient must be included in the authority application.</i>								
NOTE: <i>No applications for increased maximum quantities and/or repeats will be authorised. No more than 1 application per patient will be authorised.</i>								
8282X	Infant formula powder 900 g	5	..	..	*90.00	28.60	S-26 LF	WY
2350Q	Lactose-predigested powder infant formula 900 g	5	..	..	*90.00	28.60	Karicare De-Lact	NU

continued ➤

VARIOUS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MILK POWDER—LACTOSE FREE FORMULA—cont.								
Authority required								
<i>Proven chronic lactose intolerance in infants up to the age of 12 months. The date of birth of the patient must be included in the authority application. Lactose intolerance must have been proven by either:</i>								
<i>(a) relief of symptoms on supervised withdrawal of lactose from the diet for 3 or 4 days and subsequent re-emergence of symptoms on rechallenge with lactose containing formulae or milk or food; or</i>								
<i>(b) not less than 0.5% reducing substance in stool exudate tested with copper sulfate diagnostic compound tablet.</i>								
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
8283Y	Infant formula powder 900 g	5	5	..	*90.00	28.60	S-26 LF	WY
2349P	Lactose-predigested powder infant formula 900 g	5	5	..	*90.00	28.60	Karicare De-Lact	NU
MILK POWDER—LACTOSE MODIFIED								
Authority required								
<i>Acute lactose intolerance in children aged 1 year and over. The date of birth of the patient must be included in the authority application.</i>								
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised. No more than 1 application per patient will be authorised.</i>								
2358D	Lactose-predigested powder 900 g	3	1	..	*57.46	28.60	Digestelact	SJ
Authority required								
<i>Proven chronic lactose intolerance in children aged 1 year and over who are significantly malnourished. The date of birth of the patient must be included in the authority application. Lactose intolerance must have been proven by either:</i>								
<i>(a) relief of symptoms on supervised withdrawal of lactose from the diet for 3 or 4 days and subsequent re-emergence of symptoms on rechallenge with lactose containing formulae or milk or food; or</i>								
<i>(b) not less than 0.5% reducing substance in stool exudate tested with copper sulfate diagnostic compound tablet.</i>								
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
2357C	Lactose-predigested powder 900 g	3	10	..	*57.46	28.60	Digestelact	SJ
MILK POWDER—SYNTHETIC								
NOTE:								
<i>No applications for increased maximum quantities and/or repeats will be authorised.</i>								
Authority required								
<i>Hypercalcaemia in children under the age of 4 years.</i>								
3092R	Low calcium compound powder 400 g	8	5	..	*295.95	28.60	Locasol	NU

VARIOUS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Other combinations of nutrients AMINO ACID FORMULA without METHIONINE, THREONINE and VALINE and low in ISOLEUCINE <u>Restricted benefit</u> Methylmalonic acidaemia; Propionic acidaemia.								
3079C	Powder 200 g	5	5	..	*1081.65	28.60	XMTVI Asadon	SB
AMINO ACID FORMULA without PHENYLALANINE <u>Restricted benefit</u> Phenylketonuria.								
8554F	Capsules 500 mg, 200	16	5	..	*1334.99	28.60	Phlexy-10	SB
8678R	Tablets 1 g, 75	24	5	..	*1502.35	28.60	Phlexy-10	SB
8706F	Bars 42 g, 20	10	5	..	*1662.75	28.60	Phlexy-10	SB
2347M	Sachets containing powder 20 g, 30	7	5	..	*1542.30	28.60	Phlexy-10 Drink Mix	SB
3072Q	Powder 250 g	8	5	..	*1261.07	28.60	PK AID II	SB
AMINO ACID FORMULA without PHENYLALANINE, TYROSINE and METHIONINE <u>Restricted benefit</u> Tyrosinaemia.								
2379F	Powder 500 g	4	5	..	*1469.07	28.60	XPTM Tyrosidon	SB
AMINO ACID FORMULA with VITAMINS, MINERALS and LONG CHAIN POLYUNSATURATED FATTY ACIDS without PHENYLALANINE <u>Restricted benefit</u> Phenylketonuria.								
8479G	Infant formula, powder 400 g	8	5	..	*725.23	28.60	XP Analog LCP	SB
AMINO ACID FORMULA with VITAMINS and MINERALS without METHIONINE <u>Restricted benefit</u> For infants and very young children with pyridoxine non-responsive homocystinuria.								
8417B	Infant formula, powder 400 g	8	5	..	*718.51	28.60	XMET Analog	SB
<u>Restricted benefit</u> Pyridoxine non-responsive homocystinuria.								
8677Q	Sachets 20 g, 30	4	5	..	*1430.47	28.60	HCU gel	VF
8744F	Sachets 25 g, 30	4	5	..	*1914.19	28.60	HCU express	VF
8328H	Powder 500 g	8	5	..	*1419.15	28.60	XMET Maxamaid	SB
8416Y	Powder 500 g	8	5	..	*1665.39	28.60	XMET Maxamum	SB

VARIOUS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
AMINO ACID FORMULA with VITAMINS and MINERALS without METHIONINE, THREONINE and VALINE and low in ISOLEUCINE								
<u>Restricted benefit</u>								
<i>Methylmalonic acidaemia; Propionic acidaemia.</i>								
8058D	Infant formula, powder 400 g	8	5	..	*648.59	28.60	XMTVI Analog	SB
8059E	Powder 500 g	8	5	..	*1630.99	28.60	XMTVI Maxamaid	SB
8061G	Powder 500 g	8	5	..	*2068.11	28.60	XMTVI Maxamum	SB
AMINO ACID FORMULA with VITAMINS and MINERALS without PHENYLALANINE								
<u>Restricted benefit</u>								
<i>Phenylketonuria.</i>								
8751N	Sachets 19 g, 30	4	5	..	*914.35	28.60	PK Max	SB
8555G	Sachets 20 g, 30	4	5	..	*918.87	28.60	PKU-gel	VF
8591E	Sachets 25 g, 30	4	5	..	*1638.03	28.60	PKU-Express	VF
8804J	Sachets 27.8 g, 30	3	5	..	*1638.40	28.60	Lophlex	SB
8613H	Sachets 29 g, 30	4	5	..	*919.99	28.60	Minaphlex	SB
8727H	Sachets 50 g, 30	3	5	..	*1596.88	28.60	XP Maxamum	SB
8466N	Infant formula, powder 350 g	8	5	..	*512.27	28.60	Phenex-1	AB
8544Q	Infant formula, powder 400 g	8	5	..	*584.75	28.60	Phenex-1	AB
2737C	Infant formula, powder 400 g	8	5	..	*507.47	28.60	XP Analog	SB
8467P	Powder 325 g	10	5	..	*888.75	28.60	Phenex-2	AB
8545R	Powder 400 g	8	5	..	*874.99	28.60	Phenex-2	AB
2738D	Powder 500 g	8	5	..	*911.63	28.60	XP Maxamaid	SB
2739E	Powder 500 g	8	5	..	*1419.47	28.60	XP Maxamum	SB
8746H	Oral liquid 250 mL	90	5	..	*1376.00	28.60	Easiphen	SB
8846N	Oral liquid 130 mL, 30	4	5	..	*1637.15	28.60	PKU Express Liquid	VF
AMINO ACID FORMULA with VITAMINS and MINERALS without PHENYLALANINE and TYROSINE								
<u>Restricted benefit</u>								
<i>Tyrosinaemia.</i>								
8631G	Sachets 20 g, 30	4	5	..	*1723.99	28.60	TYR gel	VF
8667E	Sachets 25 g, 30	4	5	..	*2567.95	28.60	TYR Express	VF
8445L	Infant formula, powder 400 g	8	5	..	*795.15	28.60	XPhen, Tyr Analog	SB
8446M	Powder 500 g	8	5	..	*1710.35	28.60	XPhen, Tyr Maxamaid	SB
3078B	Powder 500 g	8	5	..	*2226.19	28.60	XPhen, Tyr Maxamum	SB

VARIOUS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
AMINO ACID FORMULA with VITAMINS and MINERALS without VALINE, LEUCINE and ISOLEUCINE								
<u>Restricted benefit</u>								
<i>Maple syrup urine disease.</i>								
8592F	Sachets 20 g, 30	4	5	..	*918.87	28.60	MSUD-gel	VF
8632H	Sachets 25 g, 30	4	5	..	*1638.03	28.60	MSUD Express	VF
8745G	Sachets 29 g, 30	4	5	..	*919.99	28.60	Mapleflex	SB
8468Q	Infant formula, powder 350 g	8	5	..	*512.27	28.60	Ketonex-1	AB
8546T	Infant formula, powder 400 g	8	5	..	*584.75	28.60	Ketonex-1	AB
2380G	Infant formula, powder 400 g	8	5	..	*507.47	28.60	MSUD Analog	SB
8469R	Powder 325 g	10	5	..	*888.75	28.60	Ketonex-2	AB
8547W	Powder 400 g	8	5	..	*874.99	28.60	Ketonex-2	AB
8310J	Powder 500 g	4	5	..	*1419.15	28.60	MSUD AID III	SB
8260R	Powder 500 g	8	5	..	*911.63	28.60	MSUD Maxamaid	SB
8057C	Powder 500 g	8	5	..	*1419.47	28.60	MSUD Maxamum	SB
CARBOHYDRATE, FAT, VITAMINS, MINERALS and TRACE ELEMENTS								
<u>Restricted benefit</u>								
<i>Patients with proven inborn errors of protein metabolism who are unable to meet their energy requirements with permitted food and formulae.</i>								
8576J	Powder 400 g	8	5	..	*266.75	28.60	Pro-Phree	AB
8369L	Powder 400 g	8	5	..	*298.75	28.60	Energivit	SB
ESSENTIAL AMINO ACIDS FORMULA with MINERALS and VITAMIN C								
<u>Restricted benefit</u>								
<i>Gyrate atrophy of the choroid and retina; Urea cycle disorders.</i>								
8001D	Powder 200 g	10	5	..	*653.55	28.60	Dialamine	SB
MILK PROTEIN and FAT FORMULA with VITAMINS and MINERALS—CARBOHYDRATE FREE								
<u>Restricted benefit</u>								
<i>Patients with intractable seizures requiring treatment with a ketogenic diet; Glucose transport protein defects; Pyruvate dehydrogenase deficiency; Infants and young children with glucose-galactose intolerance and multiple monosaccharide intolerance.</i>								
8630F	Powder 225 g	24	5	..	*668.11	28.60	Carbohydrate Free Mixture	SB

VARIOUS—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MINERAL MIXTURE								
NOTE:								
<i>For use with Amino Acid Formula without Phenylalanine.</i>								
Restricted benefit								
<i>Metabolic disorders.</i>								
1451J	Powder 250 g	‡1	5	..	50.07	28.60	Metabolic Mineral Mixture	SB
SOY PROTEIN and FAT FORMULA with VITAMINS and MINERALS—CARBOHYDRATE FREE								
Restricted benefit								
<i>Patients with intractable seizures requiring treatment with a ketogenic diet; Glucose transport protein defects; Pyruvate dehydrogenase deficiency; Infants and young children with glucose-galactose intolerance and multiple monosaccharide intolerance.</i>								
8577K	Liquid 384 mL	120	5	..	*691.15	28.60	RCF	AB
TRIGLYCERIDES, MEDIUM CHAIN and LONG CHAIN with GLUCOSE POLYMER								
Restricted benefit								
<i>Patients with proven inborn errors of protein metabolism who are unable to meet their energy requirements with permitted food and formulae.</i>								
3136C	Compound powder 400 g	8	5	..	*256.27	28.60	Duocal	SB
WHEY PROTEIN FORMULA supplemented with AMINO ACIDS, VITAMINS and MINERALS, and low in PROTEIN, PHOSPHATE, POTASSIUM and LACTOSE								
Authority required								
<i>Infants and young children with chronic renal failure requiring treatment with a low protein and a low phosphorus diet, or a low protein, a low phosphorus and a low potassium diet.</i>								
8587Y	Powder 400 g	16	5	..	*786.83	28.60	Kindergen	SB

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE

The prescribing of medications listed in this section is in accordance with the requirements for General Pharmaceutical Benefits in the Schedule unless otherwise detailed in the listing for the item.

In addition, certain additional principles have been applied by the Pharmaceutical Benefits Advisory Committee (PBAC) in recommending for whom these medications may be prescribed, and the number of repeats that may be approved by Medicare Australia. These principles have been encompassed in the listings for the items, and further details are provided below to help doctors prescribing under this section.

For the purposes of this section a patient receiving palliative care is defined as:

A patient with an active, progressive, far-advanced disease for whom the prognosis is limited and the focus of care is the quality of life.

The provision for increased maximum quantities and up to 3 repeats on the *initial* authority prescription is intended to provide up to 4 months' therapy in total. Where *continuing* treatment is required the provision of repeats is subject to confirmation by the prescriber that a palliative care physician or palliative care service has been consulted regarding the care of the patient.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ALIMENTARY TRACT AND METABOLISM								
STOMATOLOGICAL PREPARATIONS								
Stomatological preparations								
• Other agents for local oral treatment								
CARMELLOSE SODIUM								
Authority required								
<i>Initial supply (for up to 4 months) for palliative care patients where dry mouth is a symptom;</i>								
<i>Continuing supply for palliative care patients where dry mouth is a symptom, and where consultation with a palliative care specialist or service has occurred.</i>								
NOTE:								
<i>No applications for increased repeats will be authorised.</i>								
5333Q	Mouth spray 10 mg per mL, 25 mL	1	3	..	8.28	9.25	Aquae	HA
5334R	Mouth spray 10 mg per mL, 100 mL	1	3	..	10.48	11.45	Aquae	HA

continued 

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CARMELLOSE SODIUM—cont.								
<u>Authority required</u>								
<i>Continuing supply for palliative care patients where dry mouth is a symptom.</i>								
<u>NOTE:</u>								
<i>No applications for repeats will be authorised.</i>								
5335T	Mouth spray 10 mg per mL, 25 mL	‡1	..	..	8.28	9.25	Aquae	HA
5336W	Mouth spray 10 mg per mL, 100 mL	‡1	..	..	10.48	11.45	Aquae	HA

DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS

Belladonna and derivatives, plain

- *Belladonna alkaloids semisynthetic, quaternary ammonium compounds*

HYOSCINE BUTYLBROMIDE

Authority required

Initial supply (for up to 4 months) for palliative care patients where colicky pain is a symptom;

Continuing supply for palliative care patients where colicky pain is a symptom, and where consultation with a palliative care specialist or service has occurred.

NOTE:

No applications for increased repeats will be authorised.

5317W	Injection 20 mg in 1 mL	5	3	..	16.97	17.94	Buscopan	BY

Authority required

Continuing supply for palliative care patients where colicky pain is a symptom.

NOTE:

No applications for repeats will be authorised.

5318X	Injection 20 mg in 1 mL	5	..	..	16.97	17.94	Buscopan	BY

ANTIEMETICS AND ANTINAUSEANTS

Antiemetics and antinauseants

- *Other antiemetics*

PROMETHAZINE HYDROCHLORIDE

Authority required

Initial supply (for up to 4 months) for palliative care patients where nausea and/or vomiting is a problem;

Continuing supply for palliative care patients where nausea and/or vomiting is a problem, and where consultation with a palliative care specialist or service has occurred.

NOTE:

No applications for increased repeats will be authorised.

5325G	Tablet 10 mg	50	3	..	12.90	13.87	Phenergan	AV
5326H	Tablet 25 mg	50	3	..	14.97	15.94	Phenergan	AV
5327J	Oral liquid 5 mg per 5 mL, 100 mL	‡1	3	..	12.90	13.87	Phenergan	AV

continued ↻

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PROMETHAZINE HYDROCHLORIDE—cont.								
<u>Authority required</u>								
<i>Continuing supply for palliative care patients where nausea and/or vomiting is a problem.</i>								
<u>NOTE:</u>								
<i>No applications for repeats will be authorised.</i>								
5328K	Tablet 10 mg	50	..	..	12.90	13.87	Phenergan	AV
5329L	Tablet 25 mg	50	..	..	14.97	15.94	Phenergan	AV
5330M	Oral liquid 5 mg per 5 mL, 100 mL	‡1	..	..	12.90	13.87	Phenergan	AV
LAXATIVES								
Laxatives								
• Contact laxatives								
BISACODYL								
<u>Authority required</u>								
<i>Initial supply (for up to 4 months) for palliative care patients where constipation is a problem;</i>								
<i>Continuing supply for palliative care patients where constipation is a problem, and where consultation with a palliative care specialist or service has occurred.</i>								
<u>NOTE:</u>								
<i>No applications for increased repeats will be authorised.</i>								
5301B	Tablet 5 mg	200	3	..	13.00	13.97	Bisalax Lax-Tab	AS AE
5303D	Suppositories 10 mg, 10	3	3	..	*21.52	22.49	Durolax Petrus Bisacodyl Suppositories	BY PP
5304E	Suppositories 10 mg, 12	3	3	..	*17.50	18.47	Fleet Laxative Suppositories Petrus Bisacodyl Suppositories	FL PP
<u>Authority required</u>								
<i>Continuing supply for palliative care patients where constipation is a problem.</i>								
<u>NOTE:</u>								
<i>No applications for repeats will be authorised.</i>								
5305F	Tablet 5 mg	200	..	..	13.00	13.97	Bisalax Lax-Tab	AS AE
5307H	Suppositories 10 mg, 10	3	..	..	*21.52	22.49	Durolax Petrus Bisacodyl Suppositories	BY PP
5308J	Suppositories 10 mg, 12	3	..	..	*17.50	18.47	Fleet Laxative Suppositories Petrus Bisacodyl Suppositories	FL PP

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DOCUSATE SODIUM with BISACODYL								
<u>Authority required</u>								
<i>Initial supply (for up to 4 months) for palliative care patients where constipation is a problem;</i>								
<i>Continuing supply for palliative care patients where constipation is a problem, and where consultation with a palliative care specialist or service has occurred.</i>								
<u>NOTE:</u>								
<i>No applications for increased repeats will be authorised.</i>								
5309K	Suppositories 100 mg-10 mg, 5	6	3	..	*21.49	22.46	Coloxyl	FM
<u>Authority required</u>								
<i>Continuing supply for palliative care patients where constipation is a problem.</i>								
<u>NOTE:</u>								
<i>No applications for repeats will be authorised.</i>								
5310L	Suppositories 100 mg-10 mg, 5	6	..	..	*21.49	22.46	Coloxyl	FM
• Bulk producers								
STERCULIA with FRANGULA BARK								
<u>Authority required</u>								
<i>Initial supply (for up to 4 months) for palliative care patients where constipation is a problem;</i>								
<i>Continuing supply for palliative care patients where constipation is a problem, and where consultation with a palliative care specialist or service has occurred.</i>								
<u>NOTE:</u>								
<i>No applications for increased repeats will be authorised.</i>								
5322D	Granules 620 mg-80 mg per g (62%-8%), 500 g	‡1	3	..	23.05	24.02	Normacol Plus	NE
<u>Authority required</u>								
<i>Continuing supply for palliative care patients where constipation is a problem.</i>								
<u>NOTE:</u>								
<i>No applications for repeats will be authorised.</i>								
5324F	Granules 620 mg-80 mg per g (62%-8%), 500 g	‡1	..	..	23.05	24.02	Normacol Plus	NE

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Enemas BISACODYL Authority required <i>Initial supply (for up to 4 months) for palliative care patients where constipation is a problem;</i> <i>Continuing supply for palliative care patients where constipation is a problem, and where consultation with a palliative care specialist or service has occurred.</i> NOTE: <i>No applications for increased repeats will be authorised.</i>								
5302C	Enemas 10 mg in 5 mL, 25	‡1	3	..	35.77	28.60	Bisalax	AS
Authority required <i>Continuing supply for palliative care patients where constipation is a problem.</i> NOTE: <i>No applications for repeats will be authorised.</i>								
5306G	Enemas 10 mg in 5 mL, 25	‡1	..	..	35.77	28.60	Bisalax	AS
SORBITOL with SODIUM CITRATE and SODIUM LAURYL SULFOACETATE Authority required <i>Initial supply (for up to 4 months) for palliative care patients where constipation is a problem;</i> <i>Continuing supply for palliative care patients where constipation is a problem, and where consultation with a palliative care specialist or service has occurred.</i> NOTE: <i>No applications for increased repeats will be authorised.</i>								
5331N	Enemas 3.125 g-450 mg-45 mg in 5 mL, 12	2	3	..	*34.53	28.60	Microlax	PH
Authority required <i>Continuing supply for palliative care patients where constipation is a problem.</i> NOTE: <i>No applications for repeats will be authorised.</i>								
5332P	Enemas 3.125 g-450 mg-45 mg in 5 mL, 12	2	..	..	*34.53	28.60	Microlax	PH
• Other laxatives GLYCEROL Authority required <i>Initial supply (for up to 4 months) for palliative care patients where constipation is a problem;</i> <i>Continuing supply for palliative care patients where constipation is a problem, and where consultation with a palliative care specialist or service has occurred.</i> NOTE: <i>No applications for increased repeats will be authorised.</i>								
5311M	Suppositories 700 mg (for infants), 12	3	3	..	*14.77	15.74	PP	

continued ➞

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
GLYCEROL—cont.							
5312N	Suppositories 1.4 g (for children), 12	3	3	..	*15.16	16.13	PP
5313P	Suppositories 2.8 g (for adults), 12	3	3	..	*15.52	16.49	PP

Authority required

Continuing supply for palliative care patients where constipation is a problem.

NOTE:

No applications for repeats will be authorised.

5314Q	Suppositories 700 mg (for infants), 12	3	..	..	*14.77	15.74	PP
5315R	Suppositories 1.4 g (for children), 12	3	..	..	*15.16	16.13	PP
5316T	Suppositories 2.8 g (for adults), 12	3	..	..	*15.52	16.49	PP

MUSCULO-SKELETAL SYSTEM

ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS

Antiinflammatory and antirheumatic products, non-steroids

- **Acetic acid derivatives and related substances**

DICLOFENAC SODIUM

Authority required

Initial supply (for up to 4 months) for palliative care patients where severe pain is a problem;

Continuing supply for palliative care patients where severe pain is a problem, and where consultation with a palliative care specialist or service has occurred.

NOTE:

No applications for increased repeats will be authorised.

5361E	Tablet 25 mg (enteric coated)	100	3	..	*13.51	14.48	^a Chem mart	CH
							^a Diclofenac	
							^a Clonac 25	AW
							^a Diclofenac-BC	BG
							^a Diclohexal	HX
							^a Dinac	DP
							^a GenRx	GX
							^a Diclofenac	
							^a Terry White Chemists	TW
							^a Diclofenac	
				..	13.51	14.48	^a Fenac 25	AF
				^B 3.30	*16.81	14.48	^a Voltaren 25	NV

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DICLOFENAC SODIUM—cont.								
5362F	Tablet 50 mg (enteric coated)	50	3	..	10.84	11.81	a Chem mart Diclofenac	CH
							a Clonac 50	AW
							a Diclohexal	HX
							a Dinac	DP
							a Fenac	AF
							a GenRx	GX
							Diclofenac	
							a Terry White Chemists	TW
							Diclofenac	
				^B 3.21	14.05	11.81	a Voltaren 50	NV
5363G	Suppository 100 mg	40	3	..	*23.03	24.00	Voltaren 100	NV
Authority required								
Continuing supply for palliative care patients where severe pain is a problem.								
NOTE:								
No applications for repeats will be authorised.								
5364H	Tablet 25 mg (enteric coated)	100	..	..	*13.51	14.48	a Chem mart Diclofenac	CH
							a Clonac 25	AW
							a Diclofenac-BC	BG
							a Diclohexal	HX
							a Dinac	DP
							a GenRx	GX
							Diclofenac	
							a Terry White Chemists	TW
							Diclofenac	
				..	13.51	14.48	a Fenac 25	AF
				^B 3.30	*16.81	14.48	a Voltaren 25	NV
5365J	Tablet 50 mg (enteric coated)	50	..	..	10.84	11.81	a Chem mart Diclofenac	CH
							a Clonac 50	AW
							a Diclohexal	HX
							a Dinac	DP
							a Fenac	AF
							a GenRx	GX
							Diclofenac	
							a Terry White Chemists	TW
							Diclofenac	
				^B 3.21	14.05	11.81	a Voltaren 50	NV
5366K	Suppository 100 mg	40	..	..	*23.03	24.00	Voltaren 100	NV

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
INDOMETHACIN								
<u>Authority required</u>								
<i>Initial supply (for up to 4 months) for palliative care patients where severe pain is a problem;</i>								
<i>Continuing supply for palliative care patients where severe pain is a problem, and where consultation with a palliative care specialist or service has occurred.</i>								
<u>NOTE:</u>								
<i>No applications for increased repeats will be authorised.</i>								
5377B	Capsule 25 mg	100	3	..	*8.73	9.70	^a Arthrexin	AF
				^B 3.06	*11.79	9.70	^a Indocid	MK
5378C	Suppository 100 mg	40	3	..	*20.63	21.60	Indocid	MK
<u>Authority required</u>								
<i>Continuing supply for palliative care patients where severe pain is a problem.</i>								
<u>NOTE:</u>								
<i>No applications for repeats will be authorised.</i>								
5379D	Capsule 25 mg	100	..	..	*8.73	9.70	^a Arthrexin	AF
				^B 3.06	*11.79	9.70	^a Indocid	MK
5380E	Suppository 100 mg	40	..	..	*20.63	21.60	Indocid	MK
SULINDAC								
<u>Authority required</u>								
<i>Initial supply (for up to 4 months) for palliative care patients where severe pain is a problem;</i>								
<i>Continuing supply for palliative care patients where severe pain is a problem, and where consultation with a palliative care specialist or service has occurred.</i>								
<u>NOTE:</u>								
<i>No applications for increased repeats will be authorised.</i>								
5381F	Tablet 100 mg	100	3	..	*14.57	15.54	Aclin	AF
5382G	Tablet 200 mg	50	3	..	13.51	14.48	Aclin 200	AF
<u>Authority required</u>								
<i>Continuing supply for palliative care patients where severe pain is a problem.</i>								
<u>NOTE:</u>								
<i>No applications for repeats will be authorised.</i>								
5383H	Tablet 100 mg	100	..	..	*14.57	15.54	Aclin	AF
5384J	Tablet 200 mg	50	..	..	13.51	14.48	Aclin 200	AF

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Propionic acid derivatives IBUPROFEN Authority required <i>Initial supply (for up to 4 months) for palliative care patients where severe pain is a problem;</i> <i>Continuing supply for palliative care patients where severe pain is a problem, and where consultation with a palliative care specialist or service has occurred.</i> NOTE: <i>No applications for increased repeats will be authorised.</i>								
5367L	Tablet 200 mg	100	3	..	*9.11	10.08	Rafen 200	AF
5368M	Tablet 400 mg	100	3	..	12.49	13.46	Brufen	AB
Authority required <i>Continuing supply for palliative care patients where severe pain is a problem.</i> NOTE: <i>No applications for repeats will be authorised.</i>								
5369N	Tablet 200 mg	100	..	..	*9.11	10.08	Rafen 200	AF
5370P	Tablet 400 mg	100	..	..	12.49	13.46	Brufen	AB
NAPROXEN Authority required <i>Initial supply (for up to 4 months) for palliative care patients where severe pain is a problem;</i> <i>Continuing supply for palliative care patients where severe pain is a problem, and where consultation with a palliative care specialist or service has occurred.</i> NOTE: <i>No applications for increased repeats will be authorised.</i>								
5345H	Tablet 250 mg	100	3	..	*14.37	15.34	^a Inza 250	AF
				^b 3.10	*17.47	15.34	^a Naprosyn	RO
5346J	Tablet 500 mg	50	3	..	13.30	14.27	^a Inza 500	AF
				^b 1.80	15.10	14.27	^a Naprosyn	RO
5347K	Tablet 750 mg (sustained release)	28	3	..	12.60	13.57	^a Proxen SR 750	MD
				^b 1.68	14.28	13.57	^a Naprosyn SR750	RO
5348L	Tablet 1 g (sustained release)	28	3	..	15.22	16.19	^a Proxen SR 1000	MD
				^b 1.77	16.99	16.19	^a Naprosyn SR1000	RO

continued ↗

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
<u>NAPROXEN—cont.</u>								
<u>Authority required</u>								
<i>Continuing supply for palliative care patients where severe pain is a problem.</i>								
<u>NOTE:</u>								
<i>No applications for repeats will be authorised.</i>								
5349M	Tablet 250 mg	100	..	..	*14.37	15.34	^a Inza 250	AF
				^B 3.10	*17.47	15.34	^a Naprosyn	RO
5350N	Tablet 500 mg	50	..	..	13.30	14.27	^a Inza 500	AF
				^B 1.80	15.10	14.27	^a Naprosyn	RO
5351P	Tablet 750 mg (sustained release)	28	..	..	12.60	13.57	^a Proxen SR 750	MD
				^B 1.68	14.28	13.57	^a Naprosyn SR750	RO
5352Q	Tablet 1 g (sustained release)	28	..	..	15.22	16.19	^a Proxen SR 1000	MD
				^B 1.77	16.99	16.19	^a Naprosyn SR1000	RO

NAPROXEN SODIUM**Authority required**

Initial supply (for up to 4 months) for palliative care patients where severe pain is a problem;

Continuing supply for palliative care patients where severe pain is a problem, and where consultation with a palliative care specialist or service has occurred.

NOTE:

No applications for increased repeats will be authorised.

5353R	Tablet 550 mg	50	3	..	13.56	14.53	^a Crysanal	MD
				^B 3.00	16.56	14.53	^a Anaprox 550	RO

NOTE:

Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.

Authority required

Continuing supply for palliative care patients where severe pain is a problem.

NOTE:

No applications for repeats will be authorised.

5354T	Tablet 550 mg	50	..	..	13.56	14.53	^a Crysanal	MD
				^B 3.00	16.56	14.53	^a Anaprox 550	RO

NOTE:

Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
NERVOUS SYSTEM								
ANALGESICS								
Other analgesics and antipyretics								
• Anilides								
PARACETAMOL								
Authority required								
<i>Initial supply (for up to 4 months) for palliative care patients for analgesia or fever where alternative therapy cannot be tolerated;</i>								
<i>Continuing supply for palliative care patients for analgesia or fever where alternative therapy cannot be tolerated, and where consultation with a palliative care specialist or service has occurred.</i>								
NOTE:								
<i>No applications for increased repeats will be authorised.</i>								
5343F	Tablet 665 mg (modified release)	96	3	.. B2.44	8.13 10.57	9.10 ^a 9.10 ^a	Duatrol SR Panadol Osteo	ME GC
5319Y	Suppositories 500 mg, 24	‡1	3	..	24.91	25.88	Panadol	GC

Authority required

Continuing supply for palliative care patients for analgesia or fever where alternative therapy cannot be tolerated.

NOTE:

No applications for repeats will be authorised.

5344G	Tablet 665 mg (modified release)	96	..	.. B2.44	8.13 10.57	9.10 ^a 9.10 ^a	Duatrol SR Panadol Osteo	ME GC
5320B	Suppositories 500 mg, 24	‡1	..	..	24.91	25.88	Panadol	GC

ANTIEPILEPTICS

Antiepileptics

• **Benzodiazepine derivatives**

CLONAZEPAM

Authority required

Initial supply (for up to 4 months) for palliative care patients for the prevention of epilepsy;

Continuing supply for palliative care patients for the prevention of epilepsy, where consultation with a palliative care specialist or service has occurred.

NOTE:

No applications for increased repeats will be authorised.

5337X	Tablet 500 micrograms	100	3	.. B2.37	13.83 16.20	14.80 ^a 14.80 ^a	Paxam 0.5 Rivotril	AF RO
5338Y	Tablet 2 mg	100	3	.. B2.68	21.82 24.50	22.79 ^a 22.79 ^a	Paxam 2 Rivotril	AF RO
5339B	Oral liquid 2.5 mg per mL, 10 mL	2	3	..	*13.29	14.26	Rivotril	RO

continued ➡

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CLONAZEPAM—cont.								
Authority required								
<i>Continuing supply for palliative care patients for the prevention of epilepsy.</i>								
NOTE:								
<i>No applications for repeats will be authorised.</i>								
5340C	Tablet 500 micrograms	100	..	..	13.83	14.80	^a Paxam 0.5	AF
				^B 2.37	16.20	14.80	^a Rivotril	RO
5341D	Tablet 2 mg	100	..	..	21.82	22.79	^a Paxam 2	AF
				^B 2.68	24.50	22.79	^a Rivotril	RO
5342E	Oral liquid 2.5 mg per mL, 10 mL	2	..	..	*13.29	14.26	Rivotril	RO

PSYCHOLEPTICS

Anxiolytics

• **Benzodiazepine derivatives**

DIAZEPAM

Authority required

Initial supply (for up to 4 months) for palliative care patients where anxiety is a problem;

Continuing supply for palliative care patients where anxiety is a problem, and where consultation with a palliative care specialist or service has occurred.

NOTE:

No applications for increased repeats will be authorised.

5355W	Tablet 2 mg	50	3	..	6.79	7.76	^a Antenex 2	AF
				^B 0.91	7.70	7.76	^a Valpam 2	AW
				^B 1.30	8.09	7.76	^a Ducene	SU
							^a Valium	RO
5356X	Tablet 5 mg	50	3	..	7.03	8.00	^a Antenex 5	AF
							^a Diazepam-DP	DP
							^a Valpam 5	AW
				^B 0.93	7.96	8.00	^a Ducene	SU
				^B 1.32	8.35	8.00	^a Valium	RO

Authority required

Continuing supply for palliative care patients where anxiety is a problem.

NOTE:

No applications for repeats will be authorised.

5357Y	Tablet 2 mg	50	..	..	6.79	7.76	^a Antenex 2	AF
							^a Valpam 2	AW
				^B 0.91	7.70	7.76	^a Ducene	SU
				^B 1.30	8.09	7.76	^a Valium	RO
5358B	Tablet 5 mg	50	..	..	7.03	8.00	^a Antenex 5	AF
							^a Diazepam-DP	DP
							^a Valpam 5	AW
				^B 0.93	7.96	8.00	^a Ducene	SU
				^B 1.32	8.35	8.00	^a Valium	RO

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
OXAZEPAM								
Authority required								
<i>Initial supply (for up to 4 months) for palliative care patients where anxiety is a problem;</i>								
<i>Continuing supply for palliative care patients where anxiety is a problem, and where</i>								
<i>consultation with a palliative care specialist or service has occurred.</i>								
NOTE:								
<i>No applications for increased repeats will be authorised.</i>								
5371Q	Tablet 15 mg	50	3	..	*7.33	8.30	^a Alepam 15	AF
				^B 2.88	*10.21	8.30	^a Serepax	SI
5372R	Tablet 30 mg	50	3	..	*7.75	8.72	^a Alepam 30	AF
				^B 2.90	*10.65	8.72	^a Murelax	FM
							^a Serepax	SI
Authority required								
<i>Continuing supply for palliative care patients where anxiety is a problem.</i>								
NOTE:								
<i>No applications for repeats will be authorised.</i>								
5373T	Tablet 15 mg	50	..	..	*7.33	8.30	^a Alepam 15	AF
				^B 2.88	*10.21	8.30	^a Serepax	SI
5374W	Tablet 30 mg	50	..	..	*7.75	8.72	^a Alepam 30	AF
				^B 2.90	*10.65	8.72	^a Murelax	FM
							^a Serepax	SI
Hypnotics and sedatives								
• Benzodiazepine derivatives								
NITRAZEPAM								
Authority required								
<i>Initial supply (for up to 4 months) for palliative care patients where insomnia is a problem;</i>								
<i>Continuing supply for palliative care patients where insomnia is a problem, and where</i>								
<i>consultation with a palliative care specialist or service has occurred.</i>								
NOTE:								
<i>No applications for increased repeats will be authorised.</i>								
5359C	Tablet 5 mg	50	3	..	*8.63	9.60	^a Alodorm	AF
				^B 4.04	*12.67	9.60	^a Mogadon	VT
Authority required								
<i>Continuing supply for palliative care patients where insomnia is a problem.</i>								
NOTE:								
<i>No applications for repeats will be authorised.</i>								
5360D	Tablet 5 mg	50	..	..	*8.63	9.60	^a Alodorm	AF
				^B 4.04	*12.67	9.60	^a Mogadon	VT

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
TEMAZEPAM								
<u>Authority required</u>								
<i>Initial supply (for up to 4 months) for palliative care patients where insomnia is a problem;</i>								
<i>Continuing supply for palliative care patients where insomnia is a problem, and where</i>								
<i>consultation with a palliative care specialist or service has occurred.</i>								
<u>NOTE:</u>								
<i>No applications for increased repeats will be authorised.</i>								
5375X	Tablet 10 mg	50	3	..	*8.63	9.60	^a Temaze	AF
							^a Temtabs	FM
				^B 2.12	*10.75	9.60	^a Normison	SI
<u>Authority required</u>								
<i>Continuing supply for palliative care patients where insomnia is a problem.</i>								
<u>NOTE:</u>								
<i>No applications for repeats will be authorised.</i>								
5376Y	Tablet 10 mg	50	..	..	*8.63	9.60	^a Temaze	AF
							^a Temtabs	FM
				^B 2.12	*10.75	9.60	^a Normison	SI

PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ALIMENTARY TRACT AND METABOLISM								
STOMATOLOGICAL PREPARATIONS								
Stomatological preparations								
• Antiinfectives and antiseptics for local oral treatment								
AMPHOTERICIN								
3306B	Lozenge 10 mg	20	..	..	8.60	9.57	Fungilin	BQ
NYSTATIN								
3343Y	Oral suspension 100,000 units per mL, 24 mL	‡1	..	..	9.13	10.10	Mycostatin Nilstat	BQ SI
• Other agents for local oral treatment								
BENZYDAMINE HYDROCHLORIDE								
Restricted benefit								
Radiation induced mucositis.								
5032W	Mouth and throat rinse 22.5 mg per 15 mL, 500 mL	‡1	..	..	18.54	19.51	Difflam	MM
DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS								
Belladonna and derivatives, plain								
• Belladonna alkaloids, tertiary amines								
ATROPINE SULFATE								
5022H	Injection 600 micrograms in 1 mL	10	..	..	12.20	13.17	AP	
Propulsives								
• Propulsives								
METOCLOPRAMIDE HYDROCHLORIDE								
5151D	Tablet 10 mg	25	..	..	6.36	7.33	Pramin	AF
				‡2.98	9.34	7.33	Maxolon	VT
5152E	Syrup 5 mg per 5 mL, 100 mL	‡1	..	..	7.35	8.32	Maxolon	VT
5153F	Injection 10 mg in 2 mL	10	..	..	11.24	12.21	Maxolon	VT
ANTIEMETICS AND ANTINAUSEANTS								
Antiemetics and antinauseants								
• Other antiemetics								
PROCHLORPERAZINE								
CAUTION:								
Prochlorperazine may be associated with parkinsonism and tardive dyskinesia and should be used for short-term treatment only.								
5205Y	Tablet 5 mg	25	..	..	7.35	8.32	^a Stemizine	HP
				‡2.15	9.50	8.32	^a Stemetil	AV
5206B	Injection 12.5 mg in 1 mL	10	..	..	14.17	15.14	Stemetil	AV
5207C	Suppositories 5 mg, 5	‡1	..	..	15.37	16.34	Stemetil	AV
5208D	Suppositories 25 mg, 5	‡1	..	..	16.91	17.88	Stemetil	AV

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
3374N	PROMETHAZINE HYDROCHLORIDE Injection 50 mg in 2 mL	10	..	..	*18.29	19.26	MX	
ANTIDIARRHEALS, INTESTINAL ANTIINFLAMMATORY/ ANTIINFECTIVE AGENTS								
Intestinal antiinfectives								
• Antibiotics								
NYSTATIN								
3342X	Tablet 500,000 units	50	..	..	16.18	17.15	Nilstat	SI
3345C	Capsule 500,000 units	50	..	..	16.18	17.15	Nilstat	SI
BLOOD AND BLOOD FORMING ORGANS								
BLOOD SUBSTITUTES AND PERFUSION SOLUTIONS								
I.V. solutions								
• Solutions for parenteral nutrition								
GLUCOSE								
5106R	I.V. infusion 278 mmol (anhydrous) per L (5%), 1 L	5	..	..	*24.00	24.97	BX	
• Solutions affecting the electrolyte balance								
SODIUM CHLORIDE								
5212H	I.V. infusion 154 mmol per L (0.9%), 1 L	5	..	..	*24.00	24.97	BX	
5213J	I.V. infusion 513 mmol per L (3%), 1 L	2	..	..	*16.41	17.38	BX	
SODIUM CHLORIDE with GLUCOSE								
5214K	I.V. infusion 31 mmol-222 mmol (anhydrous) per L (0.18%-4%), 1 L	5	..	..	*24.00	24.97	BX	
5215L	I.V. infusion 19 mmol-104 mmol (anhydrous) per 500 mL (0.225%-3.75%), 500 mL	5	..	..	*30.05	28.60	BX	
5216M	I.V. infusion 39 mmol-69 mmol (anhydrous) per 500 mL (0.45%- 2.5%), 500 mL	5	..	..	*30.05	28.60	BX	
CARDIOVASCULAR SYSTEM								
CARDIAC THERAPY								
Antiarrhythmics, class I and III								
• Antiarrhythmics, class IB								
LIGNOCAINE HYDROCHLORIDE								
5142P	Injection 100 mg in 5 mL	5	..	..	35.31	28.60	PF	
Cardiac stimulants excl. cardiac glycosides								
• Adrenergic and dopaminergic agents								
ADRENALINE								
5004J	Injection 1 mg in 1 mL (1 in 1,000)	5	..	..	14.76	15.73	AP	

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Vasodilators used in cardiac diseases								
• Organic nitrates								
GLYCERYL TRINITRATE								
5108W	Tablets 600 micrograms, 100	‡1	..	..	8.40	9.37	^a Lycinat	FM
				^B 1.30	9.70	9.37	^a Anginine	SI
				..	30.67	28.60	Stabilised Nitrostat	LM

DERMATOLOGICALS

CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS

Corticosteroids, plain

• **Corticosteroids, weak (group I)**

HYDROCORTISONE ACETATE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

5111B	Cream 10 mg per g (1%), 30 g	‡1	..	..	6.59	7.56	^a Cortic-DS 1%	FM
				^B 1.30	7.89	7.56	^a Sigmacort	SI
5113D	Cream 10 mg per g (1%), 50 g	‡1	..	..	7.05	8.02	Cortef	DT
				^B 1.30	8.35	8.02	^a Cortic-DS 1%	FM
							^a Sigmacort	SI
5112C	Topical ointment 10 mg per g (1%), 30 g	‡1	..	..	6.59	7.56	^a Cortic-DS 1%	FM
				^B 1.30	7.89	7.56	^a Sigmacort	SI
5114E	Topical ointment 10 mg per g (1%), 50 g	‡1	..	..	7.05	8.02	^a Cortic-DS 1%	FM
				^B 1.30	8.35	8.02	^a Sigmacort	SI

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

CORTICOSTEROIDS FOR SYSTEMIC USE

Corticosteroids for systemic use, plain

• **Glucocorticoids**

BETAMETHASONE ACETATE with

BETAMETHASONE SODIUM PHOSPHATE

Restricted benefit

For local intra-articular or peri-articular infiltration;

Keloid;

Lichen planus hypertrophic.

5034Y	Injection 3 mg-3.9 mg (equivalent to 5.7 mg betamethasone) in 1 mL	5	..	..	24.66	25.63	Celestone Chronodose	SH

HYDROCORTISONE SODIUM SUCCINATE

Restricted benefit

For use in a hospital.

5118J	Injection equivalent to 100 mg hydrocortisone with 2 mL solvent	6	..	..	*37.15	28.60	Solu-Cortef	PH
5119K	Injection equivalent to 250 mg hydrocortisone with 2 mL solvent	6	..	..	*49.87	28.60	Solu-Cortef	PH

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
METHYLPREDNISOLONE ACETATE								
<u>Restricted benefit</u>								
<i>For local intra-articular or peri-articular infiltration.</i>								
5148Y	Injection 40 mg in 1 mL	5	..	..	23.31	24.28	^a Depo-Nisolone	KR
				^B 0.75	24.06	24.28	^a Depo-Medrol	PH
TRIAMCINOLONE ACETONIDE								
<u>Restricted benefit</u>								
<i>For local intra-articular or peri-articular infiltration; Keloid; Lichen planus hypertrophic.</i>								
5233K	Injection 10 mg in 1 mL	5	..	..	24.66	25.63	Kenacort-A10	BQ
PANCREATIC HORMONES								
Glycogenolytic hormones								
• Glycogenolytic hormones								
5105Q	GLUCAGON HYDROCHLORIDE Injection set containing 1 mg (1 i.u.) and 1 mL solvent in disposable syringe	1	..	..	40.12	28.60	GlucaGen Hypokit	NO
ANTIINFECTIVES FOR SYSTEMIC USE								
ANTIBACTERIALS FOR SYSTEMIC USE								
Tetracyclines								
• Tetracyclines								
3321T	DOXYCYCLINE Tablet 100 mg	7	..	..	7.43	8.40	^a Chem mart Doxycycline	CH
							^a Doxsig	SI
							^a Doxy-100	DP
							^a Doxyhexal	HX
							^a Doxylin 100	AF
							^a GenRx	GX
							Doxycycline	
							^a Terry White Chemists	TW
							Doxycycline	
				^B 1.60	9.03	8.40	^a Vibramycin	PF
3322W	Capsule 100 mg	7	..	..	7.43	8.40	^a DBL Doxycycline	FA
				^B 1.52	8.95	8.40	^a Doryx	MX

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Beta-lactam antibacterials, penicillins								
• Penicillins with extended spectrum								
AMOXYCILLIN								
3303W	Chewable tablet 250 mg	20	..	..	8.96	9.93	Amoxil	GK
3301R	Capsule 250 mg	20	..	..	7.95	8.92	^a Alphamox 250	AF
							^a Amohexal	HX
							^a Amoxycillin-DP	DG
							^a Bgramin	DP
							^a Chem mart	CH
							Amoxycillin	
							^a Cilamox	SI
							^a GenRx	GX
							Amoxycillin	
							^a Moxacin	CS
							^a Terry White	TW
							Chemists	
							Amoxycillin	
				^B 1.00	8.95	8.92	^a Amoxil	GK
3300Q	Capsule 500 mg	20	..	..	11.13	12.10	^a Alphamox 500	AF
							^a Amohexal	HX
							^a Amoxycillin-DP	DG
							^a Bgramin	DP
							^a Chem mart	CH
							Amoxycillin	
							^a Cilamox	SI
							^a GenRx	GX
							Amoxycillin	
							^a Moxacin	CS
							^a Terry White	TW
							Chemists	
							Amoxycillin	
				^B 1.00	12.13	12.10	^a Amoxil	GK
3309E	Sachet containing oral powder 3 g	1	..	..	8.81	9.78	Amoxil	GK
3310F	Powder for paediatric oral drops 100 mg per mL, 20 mL	‡1	..	..	#11.96	13.34	Amoxil	GK
3302T	Powder for syrup 125 mg per 5 mL, 100 mL	‡1	..	..	#10.02	11.40	^a Alphamox 125	AF
							^a Amohexal	HX
							^a Bgramin	DP
							^a Chem mart	CH
							Amoxycillin	
							^a Cilamox	SI
							^a GenRx	GX
							Amoxycillin	
							^a Moxacin	CS
							^a Terry White	TW
							Chemists	
							Amoxycillin	
				^B 1.00	#11.02	11.40	^a Amoxil	GK

continued ➡

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
AMOXYCILLIN—cont.								
3393N	Powder for syrup 250 mg per 5 mL, 100 mL	‡1	..	..	#11.28	12.66	^a Alphamox 250 ^a Amohexal ^a Bgramin ^a Chem mart Amoxycillin ^a Cilamox ^a GenRx Amoxycillin ^a Moxacin 250 ^a Terry White Chemists Amoxycillin	AF HX DP CH SI GX CS TW
				^B 1.01	#12.29	12.66	^a Amoxil Forte	GK
5225B	Powder for oral suspension 500 mg per 5 mL, 100 mL	‡1	..	..	#14.49	15.87	Maxamox	SZ
AMPICILLIN								
5013W	Capsule 250 mg	24	..	..	8.59	9.56	Alphacin 250	AF
5014X	Capsule 500 mg	24	..	..	11.96	12.93	Alphacin 500	AF
3313J	Powder for injection 500 mg	5	..	..	10.89	11.86	^a Austrapen ^a Ibimicyn	CS DP
3314K	Powder for injection 1 g	5	..	..	14.82	15.79	^a Aspen Ampicyn ^a Austrapen ^a Ibimicyn	AS CS DP
• Beta-lactamase sensitive penicillins								
BENZATHINE PENICILLIN								
5025L	Injection 900 mg in 2 mL cartridge-needle unit (for use with Tubex Injector)	1	..	..	24.95	25.92	Bicillin L-A Tubex	AS
BENZYL PENICILLIN								
3398W	Powder for injection 600 mg	10	..	..	*22.25	23.22	BenPen	CS
3399X	Powder for injection 3 g	10	..	..	*50.51	28.60	BenPen	CS
PHENOXYMETHYLPENICILLIN								
3360W	Tablet 250 mg	50	..	..	*11.53	12.50	Abbocillin-VK Filmtab	SI
3361X	Tablet 500 mg	50	..	..	*14.83	15.80	Abbocillin-VK Filmtab	SI
3363B	Capsule 250 mg	50	..	..	11.33	12.30	^a Cilicaine VK ^a Cilopen VK LPV ^a Penhexal VK	FM DP CS HX
3364C	Capsule 500 mg	50	..	..	14.53	15.50	^a Cilicaine VK ^a Cilopen VK LPV ^a Penhexal VK	FM DP CS HX

continued ➡

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PHENOXYMETHYLPENICILLIN—cont.								
3365D	Paediatric oral suspension 125 mg per 5 mL, 100 mL	2	..	.. B1.44	*12.35 *13.79	13.32 13.32	^a Cilicaine V ^a Abbocillin-V	FM SI
3366E	Oral suspension 250 mg per 5 mL, 100 mL	2	..	.. B1.46	*15.11 *16.57	16.08 16.08	^a Cilicaine V ^a Abbocillin-V	FM SI
PROCAINE PENICILLIN								
3371K	Injection 1.5 g	5	..	..	52.49	28.60	Cilicaine	SI
• Beta-lactamase resistant penicillins								
DICLOXACILLIN								
5098H	Powder for injection 500 mg	5	..	..	16.83	17.80	Diclocil	BQ
5099J	Powder for injection 1 g	5	..	..	23.69	24.66	Diclocil	BQ
DICLOXACILLIN								
<u>Restricted benefit</u>								
<i>Serious staphylococcal infections.</i>								
5096F	Capsule 250 mg	24	..	..	11.37	12.34	^a Diclocil ^a Dicloxsig ^a Distaph 250	BQ SI AF
5097G	Capsule 500 mg	24	..	..	18.62	19.59	^a Diclocil ^a Dicloxsig ^a Distaph 500	BQ SI AF
FLUCLOXACILLIN								
<u>CAUTION:</u>								
Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.								
5094D	Powder for injection 500 mg	5	..	..	16.71	17.68	^a Flopen ^a Flubiclox	CS DP
5095E	Powder for injection 1 g	5	..	..	23.51	24.48	^a Aspen Flucil ^a Flopen ^a Flubiclox ^a MX	AS CS DP
FLUCLOXACILLIN								
<u>CAUTION:</u>								
<i>Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.</i>								
<u>Restricted benefit</u>								
<i>Serious staphylococcal infections.</i>								
5090X	Capsule 250 mg	24	..	..	11.37	12.34	^a Flopen ^a Floxsig ^a Staphylex 250 ^a Floxapen	CS SI AF GK
				B0.46	11.83	12.34		

continued ➡

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
FLUCLOXACILLIN—cont.								
5091Y	Capsule 500 mg	24	..	..	18.62	19.59	^a Flopen	CS
							^a Floxsig	SI
				^B 0.58	19.20	19.59	^a Staphylex 500	AF
							^a Floxapen	GK
5092B	Powder for syrup 125 mg per 5 mL, 100 mL	‡1	..	..	#13.13	14.51	^a Flopen	CS
				^B 0.06	#13.19	14.51	^a Floxapen	GK
5093C	Powder for syrup 250 mg per 5 mL, 100 mL	‡1	..	..	#16.86	18.24	^a Flopen	CS
				^B 0.10	#16.96	18.24	^a Floxapen	GK
• Combinations of penicillins, incl. beta-lactamase inhibitors								
AMOXYCILLIN with CLAVULANIC ACID								
CAUTION:								
<i>Hepatotoxicity has been reported with this drug.</i>								
Restricted benefit								
<i>Infections where resistance to amoxycillin is suspected;</i>								
<i>Infections where resistance to amoxycillin is proven.</i>								
5008N	Tablet 500 mg-125 mg	10	..	..	13.39	14.36	^a Clamohexal Duo 500mg/125mg	HX
							^a Clamoxyl Duo	AL
							^a Clavulin Duo	AW
							^a Curam 500/125	SZ
							^a Muric 500/125	SL
				^B 0.99	14.38	14.36	^a Augmentin Duo	GK
5006L	Tablet 875 mg-125 mg	10	..	..	17.06	18.03	^a Chem mart Amoxycillin and Clavu- lanic Acid	CH
							^a Clamohexal Duo Forte 875mg/125mg	HX
							^a Clamoxyl Duo forte	AL
							^a Clavulin Duo Forte	AW
							^a Curam 875/125	SZ
							^a GenRx Amoxycillin and Clavu- lanic Acid	GX
							^a Muric 875/125	SL
							^a Terry White Chemists Amoxycillin and Clavulanic Acid	TW
				^B 1.30	18.36	18.03	^a Augmentin Duo forte	GK

continued ➤

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
AMOXYCILLIN with CLAVULANIC ACID—cont.								
5009P	Powder for syrup 125 mg- 31.25 mg per 5 mL, 75 mL	#1	..	..	#12.47	13.85	^a Clamohexal 125mg/ 31.25mg/5mL	HX
							^a Clamoxyl	AL
							^a Clavulin	AW
				^B 0.96	#13.43	13.85	^a Augmentin	GK
5011R	Powder for syrup 400 mg- 57 mg per 5 mL, 60 mL	#1	..	..	#14.75	16.13	^a Clamohexal Duo 400mg/ 57mg/5mL	HX
							^a Clamoxyl Duo 400	AL
							^a Clavulin Duo 400	AW
				^B 0.98	#15.73	16.13	^a Augmentin Duo 400	GK
TICARCILLIN with CLAVULANIC ACID								
Restricted benefit								
<i>Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent.</i>								
5230G	Powder for injection 3 g- 100 mg	10	..	..	152.15	28.60	Timentin	GK
Other beta-lactam antibacterials								
• First-generation cephalosporins								
CEPHALEXIN								
3317N	Capsule 250 mg	20	..	..	8.40	9.37	^a Chem mart Cephalexin	CH
							^a Cilex	DP
							^a GenRx	GX
							Cephalexin	
							^a Ialex	LN
							^a Ibilex 250	AF
							^a Sporahexal	HX
							^a Terry White Chemists	TW
							Cephalexin	
				^B 1.76	10.16	9.37	^a Keflex	AS
3318P	Capsule 500 mg	20	..	..	11.31	12.28	^a Chem mart Cephalexin	CH
							^a Cilex	DP
							^a GenRx	GX
							Cephalexin	
							^a Ialex	LN
							^a Ibilex 500	AF
							^a Sporahexal	HX
							^a Terry White Chemists	TW
							Cephalexin	
				^B 1.87	13.18	12.28	^a Keflex	AS

continued ➡

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CEPHALEXIN—cont.								
3319Q	Granules for syrup 125 mg per 5 mL, 100 mL	‡1	..	..	#11.52	12.90	^a Chem mart Cephalexin	CH
							^a Cilex	DP
							^a GenRx Cephalexin	GX
							^a Ialex	LN
							^a Ibilex 125	AF
							^a Terry White Chemists Cephalexin	TW
				^B 1.78	#13.30	12.90	^a Keflex	AS
3320R	Granules for syrup 250 mg per 5 mL, 100 mL	‡1	..	..	#13.63	15.01	^a Chem mart Cephalexin	CH
							^a Cilex	DP
							^a GenRx Cephalexin	GX
							^a Ialex	LN
							^a Ibilex 250	AF
							^a Terry White Chemists Cephalexin	TW
				^B 1.87	#15.50	15.01	^a Keflex	AS
CEPHALOTHIN								
3376Q	Powder for injection 1 g	10	..	..	43.57	28.60	^a MX	
				^B 0.30	43.87	28.60	^a Keflin Neutral	LY
• Second-generation cephalosporins								
CEFACLOR								
CAUTION: Serum sickness-like reactions have been reported with this drug, especially in children.								
5045M	Tablet 375 mg (sustained release)	10	..	..	14.51	15.48	^a Cefkor CD	DP
							^a Chem mart Cefaclor CD	CH
							^a Douglas Cefaclor-CD	DG
							^a GenRx Cefaclor CD	GX
							^a Karlol CD	LN
							^a Keflor CD	AF
							^a Terry White Chemists Cefaclor CD	TW
				^B 1.54	16.05	15.48	^a Ceclor CD	AS

continued ➡

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CEFACLOR—cont.								
5046N	Powder for oral suspension 125 mg per 5 mL, 100 mL	‡1	..	..	#14.02	15.40	^a Aclor 125 ^a Chem mart Cefaclor ^a GenRx Cefaclor ^a Keflor ^a Terry White Chemists Cefaclor	AW CH GX AF TW
				^B 1.53	#15.55	15.40	^a Ceclor	AS
5047P	Powder for oral suspension 250 mg per 5 mL, 75 mL	‡1	..	..	#14.52	15.90	^a Aclor 250 ^a Chem mart Cefaclor ^a GenRx Cefaclor ^a Keflor ^a Terry White Chemists Cefaclor	AW CH GX AF TW
				^B 1.55	#16.07	15.90	^a Ceclor	AS
5052X	CEFUROXIME AXETIL Tablet 250 mg (base)	14	..	..	14.79	15.76	Zinnat	GK
• Third-generation cephalosporins CEFOTAXIME Restricted benefit <i>Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent.</i>								
5048Q	Powder for injection 1 g	10	..	..	*56.35	28.60	^a Cefotaxime Sandoz	SZ
				..	56.35	28.60	^a MX	
5049R	Powder for injection 2 g	10	..	..	*100.15	28.60	^a Cefotaxime Sandoz	SZ
				..	100.15	28.60	^a MX	

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Sulfonamides and trimethoprim								
• Combinations of sulfonamides and trimethoprim, incl. derivatives								
TRIMETHOPRIM with SULFAMETHOXAZOLE								
CAUTION:								
There is an increased risk of severe adverse reactions with this combination in the elderly.								
3389J	Tablet 80 mg-400 mg	10	..	..	7.72	8.69	^a Resprim	AF
				^B 1.54	9.26	8.69	^a Septrin	SI
3390K	Tablet 160 mg-800 mg	10	..	..	8.66	9.63	^a Bactrim DS	RO
							^a Chem mart	CH
							Trimethoprim with Sulfame- thoxazole DS	
							^a GenRx	GX
							Trimethoprim with Sulfame- thoxazole DS	
							^a Resprim Forte	AF
							^a Terry White	TW
							Chemists	
							Trimethoprim with Sulfame- thoxazole DS	
				^B 1.46	10.12	9.63	^a Septrin Forte	SI
3391L	Oral suspension 40 mg-200 mg per 5 mL, 100 mL	‡1	..	..	8.23	9.20	^a Bactrim	RO
				^B 1.91	10.14	9.20	^a Resprim	AF
							^a Septrin	SI
Macrolides, lincosamides and streptogramins								
• Macrolides								
ERYTHROMYCIN								
3325B	Capsule 250 mg	25	..	..	8.72	9.69	^a DBL	FA
				^B 1.77	10.49	9.69	^a Erythromycin	MX
ERYTHROMYCIN ETHYL SUCCINATE								
3336N	Tablet 400 mg (base)	25	..	..	8.39	9.36	^a E-Mycin	AF
				^B 2.89	11.28	9.36	^a E.E.S. 400	AB
							Filmtab	
3334L	Powder for oral liquid 200 mg (base) per 5 mL, 100 mL	‡1	..	..	#10.45	11.83	^a E-Mycin 200	AF
				^B 2.19	#12.64	11.83	^a E.E.S. 200	AB
3337P	Powder for oral liquid 400 mg (base) per 5 mL, 100 mL	‡1	..	..	#12.08	13.46	^a E-Mycin 400	AF
				^B 1.75	#13.83	13.46	^a E.E.S. Granules	AB
ERYTHROMYCIN LACTOBIONATE								
5088T	Powder for I.V. infusion 1 g (base)	5	..	..	*50.75	28.60	Erythrocin-I.V.	AB

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Lincosamides CLINDAMYCIN <u>Restricted benefit</u> <i>Gram-positive coccal infections where these cannot be safely and effectively treated with a penicillin.</i>								
5057E	Capsule 150 mg	25	..	.. ^B 1.47	18.47 19.94	19.44 ^a 19.44 ^a	Cleocin Dalacin C	KR PH
5144R	LINCOMYCIN Injection 600 mg in 2 mL	5	..	..	26.96	27.93	Lincocin	PH
Other antibacterials • Glycopeptide antibacterials VANCOMYCIN <u>Restricted benefit</u> <i>Prophylaxis of endocarditis in patients hypersensitive to penicillin.</i>								
3323X	Powder for injection 500 mg (500,000 i.u.) vancomycin activity	2	..	.. ^B 0.36	*51.91 *52.27	28.60 ^a 28.60 ^a	MX Vancocin	LY
• Imidazole derivatives METRONIDAZOLE								
3339R	Tablet 200 mg	21	..	.. ^B 1.98	6.77 8.75	7.74 ^a 7.74 ^a	Metrogyl 200 Metronide 200 Flagyl	AF HP AV
5159M	Tablet 400 mg	5	..	..	6.66	7.63	Metrogyl 400	AF
5157K	Suppositories 500 mg, 10	‡1	..	..	20.26	21.23	Flagyl	AV
METRONIDAZOLE <u>Restricted benefit</u> <i>Treatment of anaerobic infections.</i>								
5155H	Tablet 400 mg	21	..	.. ^B 2.07	9.50 11.57	10.47 ^a 10.47 ^a	Metrogyl 400 Metronide 400 Flagyl	AF HP AV
<u>Restricted benefit</u> <i>Treatment, in a hospital, of acute anaerobic sepsis.</i>								
5154G	I.V. infusion 500 mg in 100 mL	5	..	..	*43.20	28.60	BX	
3341W	METRONIDAZOLE BENZOATE Oral suspension 320 mg per 5 mL (equivalent to 200 mg metronidazole in 5 mL), 100 mL	‡1	..	..	14.39	15.36	Flagyl S	AV

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
VACCINES								
Bacterial vaccines								
• Tetanus vaccines								
5251J	TETANUS VACCINE, ADSORBED Injection 0.5 mL in pre-filled syringe	3	..	..	*16.99	17.96	Tet-Tox	CS
MUSCULO-SKELETAL SYSTEM								
ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS								
Antiinflammatory and antirheumatic products, non-steroids								
• Acetic acid derivatives and related substances								
5079H	DICLOFENAC SODIUM Suppository 100 mg	40	..	..	*23.03	24.00	Voltaren 100	NV
<u>DICLOFENAC SODIUM</u>								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
5076E	Tablet 25 mg (enteric coated)	100	..	..	*13.51	14.48	^a Chem mart Diclofenac	CH
							^a Clonac 25	AW
							^a Diclofenac-BC	BG
							^a Diclohexal	HX
							^a Dinac	DP
							^a GenRx	GX
							Diclofenac	
							^a Terry White Chemists	TW
							Diclofenac	
				..	13.51	14.48	^a Fenac 25	AF
				^B 3.30	*16.81	14.48	^a Voltaren 25	NV
5077F	Tablet 50 mg (enteric coated)	50	..	..	10.84	11.81	^a Chem mart Diclofenac	CH
							^a Clonac 50	AW
							^a Diclohexal	HX
							^a Dinac	DP
							^a Fenac	AF
							^a GenRx	GX
							Diclofenac	
							^a Terry White Chemists	TW
							Diclofenac	
				^B 3.21	14.05	11.81	^a Voltaren 50	NV

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
5128X	INDOMETHACIN Suppository 100 mg	40	..	..	*20.63	21.60	Indocid	MK
INDOMETHACIN								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
5126T	Capsule 25 mg	100	..	..	*8.73	9.70	^a Arthrexin	AF
				^B 3.06	*11.79	9.70	^a Indocid	MK
SULINDAC								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
5217N	Tablet 100 mg	100	..	..	*14.57	15.54	Aclin	AF
5218P	Tablet 200 mg	50	..	..	13.51	14.48	Aclin 200	AF
• Oxicams								
PIROXICAM								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component.</i>								
5201R	Dispersible tablet 10 mg	50	..	..	12.77	13.74	^a GenRx Piroxicam Dispersible	GX
							^a Mobilis D-10	AF
							^a Pirohexal-D	HX
				^B 2.75	15.52	13.74	^a Feldene-D	PF
5202T	Dispersible tablet 20 mg	25	..	..	12.36	13.33	^a Chem mart Piroxicam Dispersible	CH
							^a GenRx Piroxicam Dispersible	GX
							^a Mobilis D-20	AF
							^a Pirohexal-D	HX
							^a Terry White Chemists Piroxicam Dispersible	TW
				^B 2.73	15.09	13.33	^a Feldene-D	PF
5203W	Capsule 10 mg	50	..	..	12.77	13.74	^a Chem mart Piroxicam	CH
							^a GenRx Piroxicam	GX
							^a Mobilis 10	AF
							^a Terry White Chemists Piroxicam	TW
				^B 2.75	15.52	13.74	^a Feldene	PF

continued ➡

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PIROXICAM—cont.								
5204X	Capsule 20 mg	25	..	..	12.36	13.33	^a Chem mart Piroxicam	CH
							^a GenRx Piroxicam	GX
							^a Mobilis 20	AF
							^a Terry White Chemists Piroxicam	TW
				B2.73	15.09	13.33	^a Feldene	PF
• Propionic acid derivatives								
IBUPROFEN								
5124Q	Tablets 400 mg, 20	‡1	..	..	7.65	8.62	Brufen	AB
IBUPROFEN								
Restricted benefit								
Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.								
5121M	Tablet 200 mg	100	..	..	*9.11	10.08	Rafen 200	AF
5123P	Tablet 400 mg	100	..	..	12.49	13.46	Brufen	AB
KETOPROFEN								
5139L	Suppository 100 mg	40	..	..	*21.39	22.36	Orudis	AV
KETOPROFEN								
Restricted benefit								
Chronic arthropathies (including osteoarthritis) with an inflammatory component.								
5136H	Capsule 200 mg (sustained release)	28	..	..	15.34	16.31	^a Oruvail SR	HP
				B2.01	17.35	16.31	^a Orudis SR 200	AV
NAPROXEN								
Restricted benefit								
Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.								
5176K	Tablet 250 mg	100	..	..	*14.37	15.34	^a Inza 250	AF
				B3.10	*17.47	15.34	^a Naprosyn	RO
5177L	Tablet 500 mg	50	..	..	13.30	14.27	^a Inza 500	AF
				B1.80	15.10	14.27	^a Naprosyn	RO
5178M	Tablet 750 mg (sustained release)	28	..	..	12.60	13.57	^a Proxen SR 750	MD
				B1.68	14.28	13.57	^a Naprosyn SR750	RO
5179N	Tablet 1 g (sustained release)	28	..	..	15.22	16.19	^a Proxen SR 1000	MD
				B1.77	16.99	16.19	^a Naprosyn SR1000	RO

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
NAPROXEN SODIUM								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
5186Y	Tablet 550 mg	50	..	..	13.56	14.53 ^a	Crysanal	MD
				^B 3.00	16.56	14.53 ^a	Anaprox 550	RO
<u>NOTE:</u>								
<i>Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.</i>								
• Other antiinflammatory and antirheumatic agents, non-steroids								
DIFLUNISAL								
<u>Restricted benefit</u>								
<i>Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.</i>								
5080J	Tablet 250 mg	100	..	..	*15.29	16.26	Dolobid	MK
5081K	Tablet 500 mg	50	..	..	14.98	15.95	Dolobid	MK
NERVOUS SYSTEM								
ANALGESICS								
Opioids								
• Natural opium alkaloids								
CODEINE PHOSPHATE								
5063L	Tablet 30 mg	20	..	..	10.53	11.50	FM	
<u>NOTE:</u>								
Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.								
CODEINE PHOSPHATE with PARACETAMOL								
3316M	Tablet 30 mg-500 mg	20	..	..	6.94	7.91	^a Codalgin Forte	FM
							^a Codapane Forte	AL
							^a Dolaforte	CO
							^a Dymadon Forte	GK
							^a Prodeine Forte	DK
					^B 1.89	8.83	^a Panadeine Forte	SW
HYDROMORPHONE HYDROCHLORIDE								
<u>CAUTION:</u>								
The risk of drug dependence is high.								
5129Y	Injection 2 mg in 1 mL	5	..	..	11.65	12.62	Dilaudid	AB
5130B	Injection 10 mg in 1 mL	5	..	..	17.13	18.10	Dilaudid-HP	AB
5131C	Injection 50 mg in 5 mL	5	..	..	45.86	28.60	Dilaudid-HP	AB
<u>NOTE:</u>								
Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.								

continued ➡

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
HYDROMORPHONE HYDROCHLORIDE—cont.								
<u>CAUTION:</u> <i>The risk of drug dependence is high.</i>								
<u>Restricted benefit</u> <i>Severe disabling pain not responding to non-narcotic analgesics.</i>								
5115F	Tablet 2 mg	20	..	..	12.40	13.37	Dilaudid	AB
5116G	Tablet 4 mg	20	..	..	16.07	17.04	Dilaudid	AB
5117H	Tablet 8 mg	20	..	..	22.55	23.52	Dilaudid	AB
5132D	Oral liquid 1 mg per mL, 473 mL	1	..	..	24.37	25.34	Dilaudid	AB

NOTE:

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

MORPHINE HYDROCHLORIDE**CAUTION:**

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

5237P	Oral solution 2 mg per mL, 200 mL	1	..	..	17.27	18.24	Ordine 2	MF
5238Q	Oral solution 5 mg per mL, 200 mL	1	..	..	20.14	21.11	Ordine 5	MF
5239R	Oral solution 10 mg per mL, 200 mL	1	..	..	24.70	25.67	Ordine 10	MF

NOTE:

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

MORPHINE SULFATE**CAUTION:**

The risk of drug dependence is high.

5168B	Injection 10 mg in 1 mL	5	..	..	11.61	12.58	MX SI	
5169C	Injection 15 mg in 1 mL	5	..	..	11.94	12.91	MX SI	
5170D	Injection 30 mg in 1 mL	5	..	..	13.92	14.89	MX	

NOTE:

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MORPHINE SULFATE—cont.								
<u>CAUTION:</u>								
<i>The risk of drug dependence is high.</i>								
<u>Restricted benefit</u>								
<i>Severe disabling pain not responding to non-narcotic analgesics.</i>								
5163R	Tablet 30 mg	20	..	..	13.12	14.09	Anamorph	FM
<u>NOTE:</u>								
<i>Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.</i>								
<u>Restricted benefit</u>								
<i>Chronic severe disabling pain not responding to non-narcotic analgesics.</i>								
5162Q	Tablet 5 mg (controlled release)	20	..	..	14.11	15.08	MS Contin	MF
5164T	Tablet 10 mg (controlled release)	20	..	..	15.63	16.60	MS Contin	MF
5161P	Tablet 15 mg (controlled release)	20	..	..	19.37	20.34	MS Contin	MF
5165W	Tablet 30 mg (controlled release)	20	..	..	28.45	28.60	MS Contin	MF
5166X	Tablet 60 mg (controlled release)	20	..	..	44.46	28.60	MS Contin	MF
5167Y	Tablet 100 mg (controlled release)	20	..	..	61.24	28.60	MS Contin	MF
5246D	Capsule 10 mg (containing sustained release pellets)	20	..	..	15.63	16.60	Kapanol	GK
5240T	Capsule 20 mg (containing sustained release pellets)	20	..	..	20.70	21.67	Kapanol	GK
5064M	Capsule 30 mg (controlled release)	10	..	..	19.37	20.34	MS Mono	MF
5241W	Capsule 50 mg (containing sustained release pellets)	20	..	..	36.07	28.60	Kapanol	GK
5065N	Capsule 60 mg (controlled release)	10	..	..	28.45	28.60	MS Mono	MF
5066P	Capsule 90 mg (controlled release)	10	..	..	34.50	28.60	MS Mono	MF
5242X	Capsule 100 mg (containing sustained release pellets)	20	..	..	61.24	28.60	Kapanol	GK
5067Q	Capsule 120 mg (controlled release)	10	..	..	44.46	28.60	MS Mono	MF
5171E	Sachet containing controlled release granules for oral suspension, 20 mg per sachet	20	..	..	20.70	21.67	MS Contin Suspension 20 mg	MF

continued ➞

PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MORPHINE SULFATE—cont.								
5243Y	Sachet containing controlled release granules for oral suspension, 30 mg per sachet	20	..	..	28.45	28.60	MS Contin Suspension 30 mg	MF
5244B	Sachet containing controlled release granules for oral suspension, 60 mg per sachet	20	..	..	44.46	28.60	MS Contin Suspension 60 mg	MF
5245C	Sachet containing controlled release granules for oral suspension, 100 mg per sachet	20	..	..	61.24	28.60	MS Contin Suspension 100 mg	MF

NOTE:

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

OXYCODONE**CAUTION:**

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

5194J	Suppository 30 mg	12	..	..	29.89	28.60	Proladone	PL

NOTE:

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

OXYCODONE HYDROCHLORIDE**CAUTION:**

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

5195K	Tablet 5 mg	20	..	..	10.51	11.48	Endone	SI
5191F	Capsule 5 mg	20	..	..	10.51	11.48	OxyNorm	MF
5197M	Capsule 10 mg	20	..	..	13.58	14.55	OxyNorm	MF
5198N	Capsule 20 mg	20	..	..	18.25	19.22	OxyNorm	MF
5190E	Oral solution 5 mg per 5 mL, 250 mL	1	..	..	15.09	16.06	OxyNorm Liquid 5mg/5mL	MF

NOTE:

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Chronic severe disabling pain not responding to non-narcotic analgesics.

5227D	Tablet 5 mg (controlled release)	20	..	..	15.73	16.70	OxyContin	MF
5247E	Tablet 10 mg (controlled release)	20	..	..	19.77	20.74	OxyContin	MF

continued ➤

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
OXYCODONE HYDROCHLORIDE—cont.								
5248F	Tablet 20 mg (controlled release)	20	..	..	28.45	28.60	OxyContin	MF
5249G	Tablet 40 mg (controlled release)	20	..	..	44.46	28.60	OxyContin	MF
5250H	Tablet 80 mg (controlled release)	20	..	..	72.02	28.60	OxyContin	MF

NOTE:

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

- **Phenylpiperidine derivatives**

PETHIDINE HYDROCHLORIDE**CAUTION:**

The risk of drug dependence is high.

Restricted benefit

Short-term treatment of acute pain.

5199P	Injection 50 mg in 1 mL	5	..	..	10.88	11.85	MX	
5200Q	Injection 100 mg in 2 mL	5	..	..	11.31	12.28	MX	

NOTE:

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

- **Other opioids**

TRAMADOL HYDROCHLORIDE**Restricted benefit**

*For acute pain where aspirin and/or paracetamol alone are inappropriate or have failed;
For dosage titration in chronic pain where aspirin and/or paracetamol alone are
inappropriate or have failed.*

5232J	Capsule 50 mg	20	..	..	8.36	9.33	^a Chem mart Tramadol	CH
							^a GenRx Tramadol	GX
							^a Terry White Chemists Tramadol	TW
							^a Zydol	AW
				^B 0.60	8.96	9.33	^a Tramal	CS

Restricted benefit

For pain where aspirin and/or paracetamol alone are inappropriate or have failed.

5234L	Tablet 100 mg (sustained release)	20	..	..	14.56	15.53	^a TramaHexal SR	HX
							^a Zydol SR 100	AW
				^B 1.04	15.60	15.53	^a Tramal SR 100	CS
5235M	Tablet 150 mg (sustained release)	20	..	..	17.97	18.94	^a TramaHexal SR	HX
							^a Zydol SR 150	AW
				^B 1.04	19.01	18.94	^a Tramal SR 150	CS

continued ➡

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTIEPILEPTICS								
Antiepileptics								
• Carboxamide derivatives								
CARBAMAZEPINE								
5039F	Tablet 100 mg	200	..	..	21.51	22.48	^a Carbamazepine-BC	BG
							^a Carbamazepine Sandoz	SZ
				^B 2.50	24.01	22.48	^a Tegretol 100	NV
5040G	Tablet 200 mg	200	..	..	36.09	28.60	^a Carbamazepine-BC	BG
							^a Carbamazepine Sandoz	SZ
				^B 3.00	39.09	28.60	^a Teril	AF
							^a Tegretol 200	NV
5038E	Tablet 200 mg (controlled release)	200	..	..	36.73	28.60	Tegretol CR 200	NV
5037D	Tablet 400 mg (controlled release)	200	..	..	65.51	28.60	Tegretol CR 400	NV
5041H	Oral suspension 100 mg per 5 mL, 300 mL	‡1	..	..	19.50	20.47	Tegretol Liquid	NV
ANTI-PARKINSON DRUGS								
Anticholinergic agents								
• Ethers of tropine or tropine derivatives								
BENZTROPINE MESYLATE								
5031T	Injection 2 mg in 2 mL	5	..	..	20.99	21.96	Cogentin	MK
PSYCHOLEPTICS								
Anxiolytics								
• Benzodiazepine derivatives								
DIAZEPAM								
5071X	Tablet 2 mg	50	..	..	6.79	7.76	^a Antenex 2	AF
							^a Valpam 2	AW
				^B 0.91	7.70	7.76	Ducene	SU
				^B 1.30	8.09	7.76	^a Valium	RO
5072Y	Tablet 5 mg	50	..	..	7.03	8.00	^a Antenex 5	AF
							^a Diazepam-DP	DP
							^a Valpam 5	AW
				^B 0.93	7.96	8.00	Ducene	SU
				^B 1.32	8.35	8.00	^a Valium	RO
5073B	Injection 10 mg in 2 mL	5	..	..	10.55	11.52	MX	
OXAZEPAM								
5192G	Tablet 15 mg	25	..	..	6.04	7.01	^a Alepam 15	AF
				^B 1.44	7.48	7.01	^a Serepax	SI
5193H	Tablet 30 mg	25	..	..	6.25	7.22	^a Alepam 30	AF
							^a Murelax	FM
				^B 1.45	7.70	7.22	^a Serepax	SI

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL
PRACTITIONERS FOR DENTAL TREATMENT ONLY—cont.**

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Hypnotics and sedatives								
• Benzodiazepine derivatives								
NITRAZEPAM								
5189D	Tablet 5 mg	25	..	.. B2.02	6.69 8.71	7.66 7.66	a Alodorm a Mogadon	AF VT
TEMAZEPAM								
5221T	Tablet 10 mg	25	..	.. B1.06	6.69 7.75	7.66 7.66	a Temaze a Temtabs a Normison	AF FM SI
RESPIRATORY SYSTEM								
DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES								
Adrenergics for systemic use								
• Alpha- and beta-adrenoceptor agonists								
ADRENALINE								
5004J	Injection 1 mg in 1 mL (1 in 1,000)	5	..	..	14.76	15.73	AP	
SENSORY ORGANS								
OPHTHALMOLOGICALS								
Antiinfectives								
• Antibiotics								
CHLORAMPHENICOL								
5055C	Eye drops 5 mg per mL (0.5%), 10 mL	‡1	..	..	7.60	8.57	Chloromycetin Chlorsig	PF SI
VARIOUS								
ALL OTHER THERAPEUTIC PRODUCTS								
All other therapeutic products								
• Antidotes								
NALOXONE HYDROCHLORIDE								
5173G	Injection 400 micrograms in 1 mL	1	..	..	22.04	23.01	Naloxone Min-I-Jet	CS
5174H	Injection 800 micrograms in 2 mL	1	..	..	23.14	24.11	Naloxone Min-I-Jet	CS
5175J	Injection 2 mg in 5 mL	1	..	..	34.15	28.60	Naloxone Min-I-Jet	CS

SECTION 100 ITEMS

In addition to the drugs and medicinal preparations available under normal PBS arrangements listed in this Schedule, a number of drugs are also available as pharmaceutical benefits but are distributed under alternative arrangements where these are considered more appropriate.

These alternative arrangements are provided for under section 100 of the *National Health Act 1953*. Several programs exist for the provision of drugs as pharmaceutical benefits in this way and this section lists those drugs which are available under the following programs:

HIGHLY SPECIALISED DRUGS PROGRAM

BOTULINUM TOXIN PROGRAM

HUMAN GROWTH HORMONE PROGRAM

IVF/GIFT PROGRAM

OPIATE DEPENDENCE TREATMENT PROGRAM

SPECIAL AUTHORITY PROGRAM

Complete details concerning the availability of drugs as benefits under these programs may be obtained by telephoning the relevant contact number(s) shown in each section, or in certain cases, by referring to the telephone number provided for individual drugs listings.

HIGHLY SPECIALISED DRUGS PROGRAM

The Australian Government provides funding for certain specialised medications under the Highly Specialised Drugs Program. Highly Specialised Drugs are medicines for the treatment of chronic conditions which, because of their clinical use or other special features, are restricted to supply through public and private hospitals having access to appropriate specialist facilities. To prescribe these drugs as pharmaceutical benefit items, medical practitioners are required to be affiliated with these specialist hospital units. A general practitioner or non-specialist hospital doctor may only prescribe Highly Specialised Drugs to provide maintenance therapy under the guidance of the treating specialist.

Benefits are available for the listed clinical indications only. There is no facility for individual patient approval for indications outside those listed.

To gain access to a Commonwealth funded drug under this program, a patient must attend a participating hospital and be a day admitted patient, a non-admitted patient or a patient on discharge, be under appropriate specialist medical care, meet the specific medical criteria and be an Australian resident in Australia (or other eligible person).

A patient will be required to pay a contribution for each supply of a highly specialised drug at a similar rate to the Pharmaceutical Benefits Scheme. Commonwealth subsidy is not available for hospital in-patients.

Reciprocal Health Care Agreement – Where a patient is entitled to be treated as an eligible person as a visitor from a country with which Australia has entered into a Reciprocal Health Care Agreement, the supply will be limited to the original prescription only. Repeat prescriptions for these patients are not permitted.

Private Hospitals – **In addition to the above requirements**, for Highly Specialised Drugs prescribed through private hospitals, claiming and approval of authority prescriptions is administered by Medicare Australia. Highly Specialised Drugs are authority required items. Medical practitioners must seek approval to prescribe these items as pharmaceutical benefits prior to their dispensing under the PBS. Approval of authority prescriptions by Medicare Australia may be obtained either by posting an Authority Prescription Form to Medicare Australia, or by using Medicare Australia's Authority Freecall service (1800 888 333). **Prescribers must quote the provider number of the hospital when applying.** Not more than two months' supply (one month's supply in the case of Clozapine), with provision for up to 5 repeats, will be authorised. Prescriptions for Highly Specialised Drugs can be dispensed by an approved private hospital's dispensary or by a community pharmacy.

The remuneration rates for Highly Specialised Drugs prescribed through private hospitals comprise the normal PBS ready-prepared dispensing fee plus a mark-up ascertained as follows:

- 10% for drugs with a price ex-manufacturer not more than \$40; or
- \$4 for drugs with a price ex-manufacturer more than \$40 but not more than \$100; or
- 4% for drugs with a price ex-manufacturer more than \$100.

Public Hospitals – For Highly Specialised Drugs prescribed through public hospitals, claiming and access to the program is administered by the States/Territories Health Departments. Prescriptions for Highly Specialised Drugs can be dispensed by public hospital pharmacies.

If you would like further information about the Highly Specialised Drugs Program, please contact your pharmacy, Medicare Australia (telephone 132 290) or the Australian Government adviser below:

Highly Specialised Drugs Working Party Secretariat (02) 6289 7238

For information on Pharmaceutical Benefits Pricing Authority therapeutic relativity sheets please visit www.health.gov.au and search for 'relativity sheets'.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
ABACAVIR SULFATE					
<u>Private hospital authority required</u>					
<i>Treatment of HIV infection in patients with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6264Q	Tablet 300 mg (base)	60	423.00	Ziagen	GK
6265R	Oral solution 20 mg (base) per mL, 240 mL	1	75.20	Ziagen	GK
<u>NOTE:</u>					
<i>These prices are based on special supply arrangements—see Pharmaceutical Benefits Pricing Authority relativity sheet for full details.</i>					
ABACAVIR SULFATE with LAMIVUDINE					
<u>Private hospital authority required</u>					
<i>Treatment of HIV infection in patients over 12 years of age, weighing 40 kg or more, with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6458X	Tablet 600 mg (base)-300 mg	30	564.00	Kivexa	GK
ABACAVIR SULFATE with LAMIVUDINE and ZIDOVUDINE					
<u>Private hospital authority required</u>					
<i>Treatment of HIV infection in patients over 12 years of age, weighing 40 kg or more, with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6327B	Tablet 300 mg (base)-150 mg-300 mg	60	852.00	Trizivir	GK
ADEFOVIR DIPIVOXIL					
<u>Private hospital authority required</u>					
<i>Patients with chronic hepatitis B who have failed lamivudine therapy and who satisfy all of the following criteria:</i>					
<i>(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);</i>					
<i>(2) Repeatedly elevated (greater than 1.2 times the upper limit of normal) serum ALT levels while on concurrent antihepadnaviral therapy of greater than or equal to 6 months duration in conjunction with documented chronic hepatitis B infection (HBe antigen positive and/or serum HBV DNA positive);</i>					
<i>(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.</i>					
<i>Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.</i>					
<u>NOTE:</u>					
<i>Patients may receive treatment in combination with lamivudine for the initial 3 months only of PBS-subsidised adefovir dipivoxil therapy. Patients who are immunocompromised may receive treatment in combination with lamivudine for the initial 12 months of PBS-subsidised adefovir dipivoxil therapy. Thereafter, PBS-subsidised adefovir dipivoxil must be used as monotherapy.</i>					
6450L	Tablet 10 mg	30	625.00	Hepsera	GI

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
AMPRENAVIR					
<u>Private hospital authority required</u>					
<i>Treatment of HIV infection in patients who have:</i>					
<i>(a) failed treatment with other protease inhibitors; or</i>					
<i>(b) experienced treatment-limiting toxicity with other protease inhibitors.</i>					
6333H	Capsule 150 mg	240	227.50	Agenerase	GK
6334J	Oral solution 15 mg per mL, 240 mL	1	19.50	Agenerase	GK
APOMORPHINE HYDROCHLORIDE					
<u>Private hospital authority required</u>					
<i>Parkinson's disease in patients severely disabled by motor fluctuations which do not respond to other therapy.</i>					
6104G	Injection 10 mg in 1 mL	5	31.87	Apomine	MX
ATAZANAVIR SULFATE					
<u>Private hospital authority required</u>					
<i>Treatment, in combination with 2 or more other antiretroviral drugs, of HIV infection in patients with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6451M	Capsule 150 mg (base)	60	521.91	Reyataz	BQ
6452N	Capsule 200 mg (base)	60	695.88	Reyataz	BQ
AZITHROMYCIN					
<u>Private hospital authority required</u>					
<i>Prophylaxis against Mycobacterium avium complex infections in HIV-positive patients with CD4 cell counts of less than 75 per cubic millimetre.</i>					
6221K	Tablet 600 mg	8	67.82	Zithromax	PF
BACLOFEN					
<u>Private hospital authority required</u>					
<i>Severe chronic spasticity, where oral antispastic agents have failed or have caused unacceptable side effects, in patients with chronic spasticity:</i>					
<i>(1) of cerebral origin; or</i>					
<i>(2) due to multiple sclerosis; or</i>					
<i>(3) due to spinal cord injury; or</i>					
<i>(4) due to spinal cord disease.</i>					
6284R	Intrathecal injection 10 mg in 5 mL	1	148.37	Lioresal Intrathecal	NV
6285T	Intrathecal injection 10 mg in 20 mL	1	148.37	Lioresal Intrathecal	NV

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	BOSENTAN MONOHYDRATE			
	CAUTION:			
	<i>Bosentan monohydrate is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of treatment with this drug.</i>			
	NOTE:			
	<i>Any queries concerning the arrangements to prescribe bosentan monohydrate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).</i>			
	<i>Written applications for authority to prescribe bosentan monohydrate should be forwarded to:</i>			
	<i>Medicare Australia Prior Written Approval of Specialised Drugs Reply Paid 9826 GPO Box 9826 HOBART TAS 7001</i>			
	NOTE:			
	<i>Bosentan monohydrate is not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma, where the total lung capacity is less than 70% of that predicted.</i>			
	<i>Bosentan monohydrate is not PBS-subsidised when used in combination with PBS-subsidised iloprost trometamol.</i>			
	<i>The following provides some explanatory notes regarding the availability of PBS-subsidised treatment with:</i>			
	<i>(a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, in patients with disease of WHO Functional Class III or IV severity; AND</i>			
	<i>(b) iloprost trometamol, of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in adult patients with disease of WHO Functional Class III or IV severity.</i>			
	<i>From 1 April 2005, adult patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma will be able to access, through the PBS, either bosentan monohydrate or iloprost trometamol. Once these patients are approved initial treatment with 1 of these 2 drugs, they may swap between bosentan monohydrate and iloprost trometamol at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary (unless the prescriber wishes to submit new baselines).</i>			
	<i>Patients may only swap to bosentan monohydrate or iloprost trometamol if they have not failed prior PBS-subsidised treatment with that drug.</i>			
	<i>Patients with pulmonary arterial hypertension secondary to connective tissue disease other than scleroderma or with drug-induced pulmonary arterial hypertension are only eligible for treatment with iloprost trometamol. They may not swap to bosentan monohydrate.</i>			
	<i>Patients aged less than 18 years with primary pulmonary hypertension are only eligible to receive treatment with bosentan monohydrate. They may qualify for treatment with iloprost trometamol when they are aged 18 years or older.</i>			
	<i>1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma.</i>			

continued ➞

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	BOSENTAN MONOHYDRATE—cont.			
	<i>Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, are defined as follows:</i>			
	<i>(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or</i>			
	<i>(ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or</i>			
	<i>(iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.</i>			
	2. Definition of WHO Functional Class III or IV disease severity.			
	<i>(a) WHO Functional Class III disease severity is defined as follows:</i>			
	<i>Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.</i>			
	<i>(b) WHO Functional Class IV disease severity is defined as follows:</i>			
	<i>Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.</i>			
	3. Designated hospitals.			
	<i>Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.</i>			
	4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.			
	<i>(a) Initiation of treatment.</i>			
	<i>The first written application for PBS-subsidised treatment with the first of either bosentan monohydrate or iloprost trometamol should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an ECHO composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.</i>			
	<i>Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:</i>			
	<i>(1) RHC plus ECHO composite assessments;</i>			
	<i>(2) RHC composite assessment plus 6MWT;</i>			
	<i>(3) RHC composite assessment only.</i>			
	<i>In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:</i>			
	<i>(1) ECHO composite assessment plus 6MWT;</i>			
	<i>(2) ECHO composite assessment only.</i>			
	<i>Where fewer than 3 tests are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application.</i>			
	<i>Where patients were initiated on PBS-subsidised treatment either with bosentan monohydrate on or after 1 March 2004, or with iloprost trometamol on or after 1 April 2005, the test results provided with the initial application must be no more than 2 months old at the time of application. These results will form the baseline against which response assessments will be made.</i>			

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	BOSENTAN MONOHYDRATE—cont.			
	<i>Where patients received treatment with either bosentan monohydrate or iloprost trometamol prior to being commenced on PBS-subsidised treatment with the first of either bosentan monohydrate or iloprost trometamol, the test requirements above still apply. The results that will form the baseline against which response assessments will be made will be those measured at the time patients commenced non-PBS-subsidised treatment with either bosentan monohydrate or iloprost trometamol, whichever of the 2 drugs the patient received first.</i>			
	(b) Continuation of treatment.			
	<i>The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:</i>			
	<i>(1) RHC plus ECHO composite assessments plus 6MWT;</i>			
	<i>(2) RHC plus ECHO composite assessments;</i>			
	<i>(3) RHC composite assessment plus 6MWT;</i>			
	<i>(4) ECHO composite assessment plus 6MWT;</i>			
	<i>(5) RHC composite assessment only;</i>			
	<i>(6) ECHO composite assessment only.</i>			
	<i>The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a reason why the test(s) could not be conducted must be provided with the application.</i>			
	<i>The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.</i>			
	5. Definition of response to iloprost trometamol, bosentan monohydrate or prior vasodilator treatment.			
	<i>For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.</i>			
	<i>For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.</i>			
	<i>For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.</i>			
	<i>For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.</i>			
	NOTE:			
	6. Authority approval requirements. <i>[The following 2 sections are only relevant to the PBS listing of bosentan monohydrate. The requirements specific to iloprost trometamol are given in parts 6 and 7 of the NOTE included in the iloprost trometamol Schedule entry.]</i>			
	(a) Initiation of PBS-subsidised treatment with bosentan monohydrate, where the patient has not received prior PBS-subsidised treatment with iloprost trometamol.			
	<i>All applications for initial treatment must be made in writing, must include 2 separate authority prescriptions and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months.</i>			

continued ↗

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	BOSENTAN MONOHYDRATE—cont.			
	<i>Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.</i> <i>(b) Continuation of treatment.</i> <i>Written applications for continuing treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.</i> <i>Applications for continuing treatment will only be approved for patients who have demonstrated a response to an initial course of treatment with bosentan monohydrate.</i> <i>The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with bosentan monohydrate should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.</i> <i>(c) Swapping between bosentan monohydrate and iloprost trometamol.</i> <i>For eligible patients, applications to swap between these 2 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.</i> <i>(d) Cessation of treatment.</i> <i>Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the times where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.</i> <i>For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.</i> 7. Re-treatment with bosentan monohydrate. <i>Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with bosentan monohydrate under any circumstances.</i> 8. Further information. <i>A diagrammatical representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.</i>			

continued ➤

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	BOSENTAN MONOHYDRATE—cont. Public and private hospital authority required <i>Application for initial PBS-subsidised treatment with bosentan monohydrate of adult patients who have not received prior PBS-subsidised treatment with iloprost trometamol and who have been assessed by a physician from a designated hospital to have:</i> (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, unless a RHC is contraindicated on clinical grounds; OR (b) WHO Functional Class III pulmonary arterial hypertension secondary to scleroderma and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, unless a RHC is contraindicated on clinical grounds. <i>Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.</i> <i>In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the test results of the ECHO composite assessment plus 6MWT or the ECHO composite assessment only.</i> <i>Applications for authorisation must be in writing and must include:</i> (1) two completed authority prescription forms [see Note for authority approval requirements]; and (2) a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes results from the 3 tests below, where available: (i) RHC composite assessment; and (ii) ECHO composite assessment; and (iii) 6MWT; and (3) a copy of a signed patient acknowledgment form indicating that the patient understands and acknowledges that PBS-subsidised treatment with bosentan monohydrate for primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, OR with iloprost trometamol for primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response]. <i>Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient is commenced on bosentan monohydrate treatment due to an adverse event or a contraindication to vasodilator treatment, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.</i> <i>Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].</i> <i>The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone.</i>			

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	BOSENTAN MONOHYDRATE—cont. Public and private hospital authority required <i>Application for initial PBS-subsidised treatment with bosentan monohydrate of adult patients who have not received prior PBS-subsidised treatment with iloprost trometamol and who have been assessed by a physician from a designated hospital to have:</i> <i>(a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, unless a RHC is contraindicated on clinical grounds;</i> <i>OR</i> <i>(b) WHO Functional Class III pulmonary arterial hypertension secondary to scleroderma and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, unless a RHC is contraindicated on clinical grounds;</i> <i>OR</i> <i>(c) WHO Functional Class IV primary pulmonary hypertension;</i> <i>OR</i> <i>(d) WHO Functional Class IV pulmonary arterial hypertension secondary to scleroderma.</i> <i>In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the test results of the ECHO composite assessment plus 6MWT or the ECHO composite assessment only.</i> <i>Applications for authorisation must be in writing and must include:</i> <i>(1) two completed authority prescription forms [see Note for authority approval requirements]; and</i> <i>(2) a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes results from the 3 tests below, where available:</i> <i>(i) RHC composite assessment; and</i> <i>(ii) ECHO composite assessment; and</i> <i>(iii) 6MWT; and</i> <i>(3) a copy of a signed patient acknowledgment form indicating that the patient understands and acknowledges that PBS-subsidised treatment with bosentan monohydrate for primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, OR with iloprost trometamol for primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].</i> <i>Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].</i> <i>The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone.</i>			

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

BOSENTAN MONOHYDRATE—cont.**Public and private hospital authority required**

Application for initial PBS-subsidised treatment of patients aged less than 18 years who have been assessed by a physician from a designated hospital and who have WHO Functional Class III primary pulmonary hypertension and either a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, normal right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate prior vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

(1) two completed authority prescription forms [see Note for authority approval requirements]; and

(2) a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes results from the 3 tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6MWT; and

(3) a copy of a patient acknowledgment form, signed by the parent or authorised guardian, indicating that they understand and acknowledge that PBS-subsidised treatment with bosentan monohydrate for primary pulmonary hypertension will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient is commenced on bosentan monohydrate treatment due to an adverse event or a contraindication to vasodilator treatment, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone.

Public and private hospital authority required

Application for initial PBS-subsidised treatment of patients aged less than 18 years who have been assessed by a physician from a designated hospital to have:

(a) WHO Functional Class III primary pulmonary hypertension and either a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular dysfunction as assessed by ECHO;

OR

(b) WHO Functional Class IV primary pulmonary hypertension.

Code	Item and Section 100 Restriction	Pack	Price ex	Proprietary
		Size	Manufacturer	Name and
			\$	Manufacturer

BOSENTAN MONOHYDRATE—cont.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and*
- (2) a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes results from the 3 tests below, where available:*
 - (i) RHC composite assessment; and*
 - (ii) ECHO composite assessment; and*
 - (iii) 6MWT; and*
- (3) a copy of a patient acknowledgment form, signed by the parent or authorised guardian, indicating that they understand and acknowledge that PBS-subsidised treatment with bosentan monohydrate for primary pulmonary hypertension will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].*

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone.

Public and private hospital authority required

Application for initial treatment with bosentan monohydrate of adult patients with either of the following:

- (a) primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma who wish to re-commence PBS-subsidised bosentan monohydrate after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with bosentan monohydrate;*

OR

- (b) primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma and whose most recent course of PBS-subsidised treatment was with iloprost trometamol.*

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and*
- (2) a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the results on which approval for the first application for PBS-subsidised bosentan monohydrate or iloprost trometamol, whichever was initiated first, was granted; and*
- (3) the date of the first application for PBS-subsidised treatment with bosentan monohydrate or iloprost trometamol, whichever was initiated first; and*
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised bosentan monohydrate and iloprost trometamol.*

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	BOSENTAN MONOHYDRATE—cont.				
	<i>The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone.</i>				
	Public and private hospital authority required				
	<i>Continuing PBS-subsidised treatment with bosentan monohydrate of patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, who have received approval for initial PBS-subsidised treatment with bosentan monohydrate and have completed the initial treatment course, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of bosentan monohydrate treatment [see Note for definition of response].</i>				
	<i>Applications for authorisation must be in writing and must include:</i>				
	<i>(1) a completed authority prescription form; and</i>				
	<i>(2) a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes results from the 3 tests below, where available:</i>				
	<i>(i) RHC composite assessment; and</i>				
	<i>(ii) ECHO composite assessment; and</i>				
	<i>(iii) 6MWT.</i>				
	<i>Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].</i>				
	<i>The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. Up to 5 repeats will be authorised.</i>				
	<i>Where fewer than 5 repeats are initially requested under this criterion, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone.</i>				
	Public and private hospital authority required				
	<i>Final PBS-subsidised supply for patients with WHO Functional Class III or IV primary pulmonary hypertension or WHO Functional Class III or IV pulmonary arterial hypertension secondary to scleroderma who have not responded to bosentan monohydrate therapy [see Note for definition of response], to allow for gradual cessation of treatment.</i>				
	<i>Applications for authorisation under this criterion should be made on the telephone [see Note on authority approval requirements].</i>				
	<i>Approval will only be granted for the 62.5 mg tablet strength. The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment.</i>				
	<i>Under no circumstances will telephone approvals be granted for treatment that would extend the final treatment period beyond 1 month.</i>				
6429J	Tablet 62.5 mg (base)	60	4035.00	Tracleer	AT
6430K	Tablet 125 mg (base)	60	4035.00	Tracleer	AT

NOTE:

These prices are based on special supply arrangements—see Pharmaceutical Benefits Pricing Authority relativity sheet for full details.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
CIDOFOVIR					
<u>Private hospital authority required</u>					
<i>Treatment of cytomegalovirus retinitis in patients with AIDS.</i>					
6247T	Solution for I.V. infusion 375 mg (anhydrous) in 5 mL single use vial	1	900.00	Vistide	PU
CLARITHROMYCIN					
<u>Private hospital authority required</u>					
<i>Treatment of Mycobacterium avium complex infections.</i>					
6151R	Tablet 250 mg	100	67.50	Klacid	AB
6152T	Tablet 500 mg	100	135.00	Klacid	AB
CLOZAPINE					
<u>Private hospital authority required</u>					
<i>Schizophrenia in patients who are:</i>					
<i>(a) non-responsive to other neuroleptic agents; or</i>					
<i>(b) intolerant of other neuroleptic agents.</i>					
6462D	Tablet 25 mg	28	20.16	Clozaril 25	NV
6101D	Tablet 25 mg	100	72.00	^a Clopine 25	MX
				^a CloSyn	ZT
				^a Clozaril 25	NV
6417R	Tablet 50 mg	100	144.00	Clopine 50	MX
6463E	Tablet 100 mg	28	75.60	Clozaril 100	NV
6102E	Tablet 100 mg	100	270.00	^a Clopine 100	MX
				^a CloSyn	ZT
				^a Clozaril 100	NV
6418T	Tablet 200 mg	100	540.00	Clopine 200	MX
CYCLOSPORIN					
CAUTION:					
<i>Careful monitoring of patients is mandatory.</i>					
<u>Private hospital authority required</u>					
<i>For use by organ or tissue transplant recipients.</i>					
6109M	Solution concentrate for I.V. infusion 50 mg in 1 mL ampoule	10	54.10	Sandimmun	NV
6110N	Solution concentrate for I.V. infusion 250 mg in 5 mL ampoule	10	257.95	Sandimmun	NV

continued ↻

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
CYCLOSPORIN—cont.					
Private hospital authority required					
<i>Management of rejection in patients following organ or tissue transplantation, under the supervision and direction of a transplant unit. Management includes initiation, stabilisation and review of therapy as required;</i>					
<i>Management (which includes initiation, stabilisation and review of therapy) by:</i>					
<i>(1) dermatologists or clinical immunologists of patients with severe atopic dermatitis for whom other systemic therapies are ineffective or inappropriate;</i>					
<i>(2) dermatologists of patients with severe psoriasis for whom other systemic therapies are ineffective or inappropriate and in whom the disease has caused significant interference with quality of life;</i>					
<i>(3) nephrologists of patients with nephrotic syndrome in patients in whom steroids and cytostatic drugs have failed or are not tolerated or are considered inappropriate and in whom renal function is unimpaired;</i>					
<i>(4) rheumatologists or clinical immunologists of patients with severe active rheumatoid arthritis for whom classical slow-acting anti-rheumatic agents (including methotrexate) are ineffective or inappropriate.</i>					
6232B	Capsule 10 mg	60	37.20	Neoral 10	NV
6352H	Capsule 25 mg	30	40.65	^a Cicloral	HX
				^a Cysporin	MX
				^a Neoral 25	NV
NOTE:					
<i>A brand premium of \$0.98 applies to Cicloral brand. A brand premium of \$1.96 applies to Neoral 25 brand. All 3 brands may not be available in all hospitals.</i>					
6353J	Capsule 50 mg	30	85.54	^a Cicloral	HX
				^a Cysporin	MX
				^a Neoral 50	NV
NOTE:					
<i>A brand premium of \$1.07 applies to Cicloral brand. A brand premium of \$2.14 applies to Neoral 50 brand. All 3 brands may not be available in all hospitals.</i>					
6354K	Capsule 100 mg	30	175.38	^a Cicloral	HX
				^a Cysporin	MX
				^a Neoral 100	NV
NOTE:					
<i>A brand premium of \$1.08 applies to Cicloral brand. A brand premium of \$2.16 applies to Neoral 100 brand. All 3 brands may not be available in all hospitals.</i>					
6125J	Oral liquid 100 mg per mL, 50 mL	1	315.79	Neoral	NV
DARBEPOETIN ALFA					
Private hospital authority required					
<i>Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.</i>					
6320P	Injection 10 micrograms in 0.4 mL pre-filled syringe	4	187.41	Aranesp	AN
6321Q	Injection 20 micrograms in 0.5 mL pre-filled syringe	4	352.96	Aranesp	AN
6322R	Injection 30 micrograms in 0.3 mL pre-filled syringe	4	482.87	Aranesp	AN
6323T	Injection 40 micrograms in 0.4 mL pre-filled syringe	4	586.10	Aranesp	AN

continued ↗

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
DARBEPOETIN ALFA—cont.					
6324W	Injection 50 micrograms in 0.5 mL pre-filled syringe	4	724.62	Aranesp	AN
6325X	Injection 60 micrograms in 0.3 mL pre-filled syringe	4	850.87	Aranesp	AN
6438W	Injection 80 micrograms in 0.4 mL pre-filled syringe	4	1120.00	Aranesp	AN
6326Y	Injection 100 micrograms in 0.5 mL pre-filled syringe	4	1379.21	Aranesp	AN
6365B	Injection 150 micrograms in 0.3 mL pre-filled syringe	4	2055.00	Aranesp	AN
DEFERIPRONE					
<u>Private hospital authority required</u>					
Iron overload in patients with thalassaemia major who are unable to take desferrioxamine therapy;					
Iron overload in patients with thalassaemia major in whom desferrioxamine therapy has proven ineffective.					
6416Q	Tablet 500 mg	100	400.00	Ferriprox	OA
DELAVIDINE MESYLATE					
<u>Private hospital authority required</u>					
Treatment of HIV infection in patients with:					
(a) CD4 cell counts of less than 500 per cubic millimetre; or					
(b) viral load of greater than 10,000 copies per mL.					
6243N	Tablet 100 mg	360	271.58	Rescriptor	PF
DEFERRIOXAMINE MESYLATE					
<u>Private hospital authority required</u>					
Disorders of erythropoiesis associated with treatment-related chronic iron overload.					
6113R	Powder for injection 500 mg vial	10	99.00	^a Desferal 500 mg	NV
				^a MX	
<u>NOTE:</u>					
A brand premium of \$8.19 applies to Desferal 500 mg brand. Both brands may not be available in all hospitals.					
6270B	Powder for injection 2 g vial	1	39.60	^a Desferal 2 g	NV
				^a MX	
<u>NOTE:</u>					
A brand premium of \$0.40 applies to Desferal 2 g brand. Both brands may not be available in all hospitals.					
DIDANOSINE					
<u>Private hospital authority required</u>					
Treatment of HIV infection in patients with:					
(a) CD4 cell counts of less than 500 per cubic millimetre; or					
(b) viral load of greater than 10,000 copies per mL.					
6298L	Capsule 125 mg (containing enteric coated beadlets)	30	102.12	Videx EC	BQ
6299M	Capsule 200 mg (containing enteric coated beadlets)	30	163.40	Videx EC	BQ

continued ➤

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	<i>DIDANOSINE—cont.</i>				
6300N	<i>Capsule 250 mg (containing enteric coated beadlets)</i>	30	204.24	Videx EC	BQ
6301P	<i>Capsule 400 mg (containing enteric coated beadlets)</i>	30	326.79	Videx EC	BQ
	<i>DISODIUM PAMIDRONATE</i>				
	<u>Private hospital authority required</u>				
	<i>Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.</i>				
6286W	<i>Concentrated injection 15 mg in 5 mL</i>	1	56.89	Pamisol	MX
6290C	<i>Injection set containing 4 vials powder for I.V. infusion 15 mg and 4 ampoules solvent 5 mL</i>	1	227.57	Aredia 15 mg	NV
6287X	<i>Concentrated injection 30 mg in 10 mL</i>	1	113.79	Pamisol	MX
6279L	<i>Injection set containing 2 vials powder for I.V. infusion 30 mg and 2 ampoules solvent 10 mL</i>	1	227.57	Aredia 30 mg	NV
6288Y	<i>Concentrated injection 60 mg in 10 mL</i>	1	227.57	Pamisol	MX
6289B	<i>Concentrated injection 90 mg in 10 mL</i>	1	341.36	Pamisol	MX
6223M	<i>Injection set containing 1 vial powder for I.V. infusion 90 mg and 1 ampoule solvent 10 mL</i>	1	341.36	Aredia 90 mg	NV
	<u>Private hospital authority required</u>				
	<i>Multiple myeloma; Bone metastases from breast cancer.</i>				
6289B	<i>Concentrated injection 90 mg in 10 mL</i>	1	341.36	Pamisol	MX
6223M	<i>Injection set containing 1 vial powder for I.V. infusion 90 mg and 1 ampoule solvent 10 mL</i>	1	341.36	Aredia 90 mg	NV
	<i>DORNASE ALFA</i>				
	<u>Private hospital authority required</u>				
	<i>Use by cystic fibrosis patients who satisfy all of the following criteria:</i>				
	<i>(1) are 5 years of age or older;</i>				
	<i>(2) have a FVC greater than 40% predicted for age, gender and height;</i>				
	<i>(3) have evidence of chronic suppurative lung disease (cough and sputum most days of the week, or greater than 3 respiratory tract infections of more than 2 weeks' duration in any 12 months, or objective evidence of obstructive airways disease);</i>				
	<i>(4) are participating in a 4 week trial as detailed below or have achieved a 10% or greater improvement in FEV1 (compared to baseline established prior to dornase alfa treatment) after a 4 week trial.</i>				
	<i>In order for patients to be eligible for participation in the HSD program, the following conditions must be met:</i>				
	<i>(1) Patients must be assessed at cystic fibrosis clinics/centres which are under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis and the prescribing of dornase alfa under the HSD program is limited to such physicians. If attendance at such units is not possible because of geographical isolation, management (including prescribing) may be by specialist physician or paediatrician in consultation with such a unit;</i>				
	<i>(2) The measurement of lung function is to be conducted by independent (other than the treating doctor) experienced personnel at established lung function testing laboratories, unless this is not possible because of geographical isolation;</i>				

continued ➞

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	DORNASE ALFA—cont.				
	(3) Prior to dornase alfa therapy, a baseline measurement of FEV1 must be undertaken during a stable period of the disease;				
	(4) Initial therapy is limited to 4 weeks' treatment with dornase alfa at a dose of 2.5 mg daily;				
	(5) At or towards the end of the initial 4 weeks' trial, patients must be reassessed and a further FEV1 measurement be undertaken (single test under conditions as above). Patients who achieve a 10% or greater improvement in FEV1 (compared to baseline established prior to dornase alfa treatment) are eligible for continued subsidy under the HSD program at a dose of 2.5 mg daily;				
	(6) Patients who fail to meet a 10% or greater improvement in FEV1 after the initial 4 weeks' treatment at a dose of 2.5 mg daily, may have 1 further trial in the next 12 months but not before 3 months after the initial trial;				
	(7) Following an initial 6 months' therapy, a global assessment must be undertaken involving the patient, the patient's family (in the case of paediatric patients) and the treating physician(s) to establish that all agree that dornase alfa treatment is continuing to produce worthwhile benefits. (Dornase alfa therapy should cease if there is not general agreement of benefit as there is always the possibility of harm from unnecessary use.) Further reassessments are to be undertaken at six-monthly intervals;				
	(8) Other aspects of treatment, such as physiotherapy, must be continued;				
	(9) Where there is documented evidence that a patient already receiving dornase alfa therapy would have met the criteria for subsidy (i.e. satisfied the criteria for the 4 week trial and achieved a 10% or greater improvement in FEV1) then the patient is eligible to continue treatment under the HSD program. Where such evidence is not available, patients will need to satisfy the initiation and continuation criteria as for new patients. (Four weeks is considered a suitable wash-out period).				
	NOTE: It is highly desirable that all patients be included in the national cystic fibrosis patient data-base.				
6120D	Solution for inhalation 2.5 mg (2,500 units) in 2.5 mL ampoule	30	1180.00	Pulmozyme	RO
	DOXORUBICIN HYDROCHLORIDE, PEGYLATED LIPOSOMAL				
	Private hospital authority required				
	Treatment of AIDS-related Kaposi's sarcoma in patients with CD4 cell counts of less than 200 per cubic millimetre and:				
	(a) extensive mucocutaneous involvement; or				
	(b) extensive visceral involvement.				
6249X	Suspension for I.V. infusion 20 mg in 10 mL vial	1	645.59	Caelyx	SH
	EFAVIRENZ				
	Private hospital authority required				
	Treatment of HIV infection in patients with:				
	(a) CD4 cell counts of less than 500 per cubic millimetre; or				
	(b) viral load of greater than 10,000 copies per mL.				
6259K	Capsule 100 mg	30	75.50	Stocrin	MK
6283Q	Capsule 200 mg	90	452.64	Stocrin	MK
6356M	Tablet 600 mg	30	452.64	Stocrin	MK
6372J	Oral solution 30 mg per mL, 180 mL	1	135.79	Stocrin	MK
	NOTE: These prices are based on special supply arrangements—see Pharmaceutical Benefits Pricing Authority relativity sheet for full details.				

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	<u>EMTRICITABINE</u>				
	<u>Private hospital authority required</u>				
	<i>Treatment of HIV infection in patients with:</i>				
	<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>				
	<i>(b) viral load of greater than 10,000 copies per mL.</i>				
6137B	Capsule 200 mg	30	282.00	Emtriva	GI
	<u>ENFUVIRTIDE</u>				
	<u>Private hospital authority required</u>				
	<i>Treatment, in combination with other antiretroviral agents, of HIV infection in antiretroviral experienced patients with treatment failure characterised by:</i>				
	<i>(a) evidence of HIV replication, despite ongoing therapy; or</i>				
	<i>(b) treatment-limiting toxicity to previous antiretroviral agents.</i>				
	<i>Patients must have failed previous treatment with 3 different antiretroviral regimens. At least 1 of each of the following classes of antiretroviral drugs must have been attempted:</i>				
	<i>(i) at least 1 non-nucleoside reverse transcriptase inhibitor; and</i>				
	<i>(ii) at least 1 nucleoside reverse transcriptase inhibitor; and</i>				
	<i>(iii) at least 1 protease inhibitor.</i>				
6455R	Pack containing 60 vials powder for injection 90 mg with 60 vials water for injections 1.1 mL (with syringes and swabs)	1	2213.00	Fuzeon	RO
	<u>EPOETIN ALFA</u>				
	<u>Private hospital authority required</u>				
	<i>Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.</i>				
6251B	Injection 1,000 units in 0.5 mL pre-filled syringe	6	147.00	Eprex 1000	JC
6204M	Injection 2,000 units in 0.5 mL pre-filled syringe	6	272.00	Eprex 2000	JC
6205N	Injection 3,000 units in 0.3 mL pre-filled syringe	6	351.00	Eprex 3000	JC
6206P	Injection 4,000 units in 0.4 mL pre-filled syringe	6	447.00	Eprex 4000	JC
6302Q	Injection 5,000 units in 0.5 mL pre-filled syringe	6	556.50	Eprex 5000	JC
6303R	Injection 6,000 units in 0.6 mL pre-filled syringe	6	660.60	Eprex 6000	JC
6305W	Injection 8,000 units in 0.8 mL pre-filled syringe	6	856.80	Eprex 8000	JC
6207Q	Injection 10,000 units in 1 mL pre-filled syringe	6	1037.00	Eprex 10000	JC
6434P	Injection 20,000 units in 0.5 mL pre-filled syringe	6	2040.00	Eprex 20,000	JC
6339P	Injection 40,000 units in 1 mL pre-filled syringe	1	660.00	Eprex 40,000	JC

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	ETANERCEPT			
	NOTE:			
	<i>Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).</i>			
	<i>Written applications for authority to prescribe etanercept should be forwarded to:</i>			
	<i>Medicare Australia Prior Written Approval of Specialised Drugs Reply Paid 9826 GPO Box 9826 HOBART TAS 7001</i>			
	<i>Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au.</i>			
	<u>Public and private hospital authority required</u>			
	<i>Initial treatment by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of patients under 18 years who have severe active polyarticular course juvenile chronic arthritis;</i>			
	AND			
	<i>(a) whose parent or authorised guardian has signed a patient agreement form indicating that they understand and acknowledge that PBS-subsidised treatment will cease if the predetermined response criteria do not support continuation of PBS-subsidised treatment;</i>			
	AND			
	<i>(b) who have demonstrated either:</i>			
	<i>(i) severe intolerance of, or toxicity due to, methotrexate (see below for definition of severe intolerance and toxicity); or</i>			
	<i>(ii) failure to achieve an adequate response to 1 or more of the following treatment regimens:</i>			
	<i>— oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or</i>			
	<i>— oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other DMARD, alone or in combination with corticosteroids, for a minimum of 3 months. (Note: use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.)</i>			
	<i>Severe intolerance is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant NSAIDs on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.</i>			
	<i>Toxicity is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.</i>			
	<i>The following criteria must be met in order to demonstrate failure to achieve an adequate response to either of the above treatment regimens:</i>			
	<i>(a) an active joint count of at least 20 active (swollen and tender) joints;</i>			
	OR			
	<i>(b) at least 4 active joints from the following list:</i>			
	<i>(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or</i>			
	<i>(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).</i>			

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

ETANERCEPT—cont.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, or intolerance develops during the period of use such that permanent withdrawal is necessary and a suitably effective treatment regimen cannot be implemented, this exempts the requirement to demonstrate an inadequate response within the time period specified above for these agents.

The authority application must be in writing and must include the information used to determine the patient's eligibility under the criteria above. The date of the joint assessment must be provided.

Only 16 weeks of treatment will be approved. The assessment of the patient's response to initial treatment should be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated.

Public and private hospital authority required

Initial PBS-subsidised supply for continuing treatment by a rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of severe active polyarticular course juvenile chronic arthritis in patients receiving treatment with etanercept prior to 1 December 2002;

AND

(a) whose parent or authorised guardian has signed a patient agreement form indicating that they understand and acknowledge that PBS-subsidised treatment will cease if the predetermined response criteria do not support continuation of PBS-subsidised treatment;

AND

(b) who have demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with etanercept.

The authority application must be in writing and must include sufficient information to determine the patient's eligibility. The date of the joint assessment must be provided.

Only 6 months of treatment will be approved.

Public and private hospital authority required

Continuing PBS-subsidised treatment by a rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of severe active polyarticular course juvenile chronic arthritis in patients who have demonstrated an adequate response to treatment with etanercept as manifested by:

(a) an active joint count of fewer than 10 active (swollen and tender) joints;

OR

(b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline;

OR

(c) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender);

and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All authority applications for continuing treatment with etanercept must be in writing and must include sufficient information to determine the patient's response according to the above criteria. The date of the joint assessment must be provided.

Only 6 months of treatment per application will be approved. Applications for continuing treatment with etanercept should be made prior to the completion of 16 weeks of treatment to ensure continuity for those patients who meet the criteria.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
ETANERCEPT—cont.					
<i>Patients who fail to demonstrate an adequate response, as specified in the criteria for continuing treatment with etanercept, will not be eligible to recommence treatment with etanercept within 12 months of the date on which treatment was ceased.</i>					
<i>Withdrawal of treatment with etanercept should be considered in patients who have achieved and sustained complete remission of disease for 12 months. Subsequent applications for PBS-subsidised re-treatment with etanercept will be subject to the authority conditions applying to initial treatment and will not be authorised within 12 months of the date on which treatment with etanercept was ceased.</i>					
<i>Where re-treatment with etanercept after a break in PBS-subsidised treatment with the drug is being sought, the reason for and date of cessation of the previous treatment course with etanercept must be included in the application.</i>					
6367D	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	1	815.00	Enbrel	WY
EVEROLIMUS					
<u>CAUTION:</u>					
<i>Careful monitoring of patients is mandatory.</i>					
<u>Private hospital authority required</u>					
<i>Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for:</i>					
<i>(a) Prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required;</i>					
<i>(b) Prophylaxis of cardiac allograft rejection. Management includes initiation, stabilisation and review of therapy as required.</i>					
6459Y	Tablet 0.25 mg	60	240.30	Certican	NV
6460B	Tablet 0.5 mg	60	480.60	Certican	NV
6461C	Tablet 0.75 mg	60	720.90	Certican	NV
FILGRASTIM					
<u>Private hospital authority required</u>					
<i>For use in patients undergoing induction and consolidation therapy for acute myeloid leukaemia;</i>					
<i>Mobilisation of peripheral blood progenitor cells to facilitate harvest of such cells for autologous transplantation into patients with non-myeloid malignancies who have had myeloablative or myelosuppressive therapy;</i>					
<i>Mobilisation of peripheral blood progenitor cells, in normal volunteers, for use in allogeneic transplantation;</i>					
<i>Patients receiving marrow-ablative chemotherapy and subsequent bone marrow transplantation;</i>					
<i>Patients with non-myeloid malignancies receiving marrow-ablative chemotherapy and subsequent autologous peripheral blood progenitor cell transplantation;</i>					

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	FILGRASTIM—cont.			
	<i>Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in:</i>			
	<i>Acute lymphoblastic leukaemia;</i>			
	<i>Ewing's sarcoma;</i>			
	<i>Germ cell tumours;</i>			
	<i>Infants and children with CNS tumours;</i>			
	<i>Neuroblastoma;</i>			
	<i>Non-Hodgkin's lymphoma (intermediate or high grade);</i>			
	<i>Osteosarcoma;</i>			
	<i>Relapsed Hodgkin's disease;</i>			
	<i>Rhabdomyosarcoma;</i>			
	<i>Patients with breast cancer receiving standard dose adjuvant chemotherapy who have had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;</i>			
	<i>Patients receiving first-line chemotherapy for Hodgkin's disease who have had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;</i>			
	<i>Patients receiving chemotherapy for myeloma who have had a prior episode of febrile neutropenia, and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;</i>			
	<i>Patients with severe congenital neutropenia (absolute neutrophil count of less than 100 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, and in whom a bone marrow examination has shown evidence of maturational arrest of the neutrophil lineage);</i>			
	<i>Patients with severe chronic neutropenia (absolute neutrophil count of less than 1,000 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, or evidence of neutrophil dysfunction, and, either having experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics in the previous 12 months, or having recurrent clinically significant infections (a minimum of 3 in the previous 12 months));</i>			
	<i>Patients with chronic cyclic neutropenia (absolute neutrophil count of less than 500 million cells per litre lasting for 3 days per cycle, measured over 3 separate cycles, and, either having experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics, or having recurrent clinically significant infections (a minimum of 3 in the previous 12 months)).</i>			
6126K	Injection 300 micrograms in 1 mL vial	10	1504.00	Neupogen AN
6291D	Injection 300 micrograms in 0.5 mL single use pre-filled syringe	10	1504.00	Neupogen AN
6127L	Injection 480 micrograms in 1.6 mL vial	10	2407.00	Neupogen AN
6292E	Injection 480 micrograms in 0.5 mL single use pre-filled syringe	10	2407.00	Neupogen AN

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	FOSAMPRENAVIR CALCIUM				
	<u>Private hospital authority required</u>				
	<i>Treatment, in combination with 2 or more other antiretroviral drugs, of HIV infection in patients with:</i>				
	<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>				
	<i>(b) viral load of greater than 10,000 copies per mL.</i>				
6453P	Tablet 700 mg (base)	60	568.74	Telzir	GK
	<u>NOTE:</u>				
	<i>This price is based on special supply arrangements—see Pharmaceutical Benefits Pricing Authority relativity sheet for full details.</i>				
6454Q	Oral liquid 50 mg (base) per mL, 225 mL	1	101.56	Telzir	GK
	FOSCARNET SODIUM				
	<u>Private hospital authority required</u>				
	<i>Treatment of cytomegalovirus retinitis in patients with AIDS;</i>				
	<i>Treatment of aciclovir-resistant herpes simplex virus infection in immunocompromised patients with HIV infection.</i>				
6134W	I.V. infusion 24 mg per mL, 250 mL bottle	6	395.00	Foscavir	AP
	GANCICLOVIR				
	<u>Private hospital authority required</u>				
	<i>Cytomegalovirus retinitis in severely immunocompromised patients.</i>				
6256G	Intravitreal implant 4.5 mg	1	6000.00	Vitrasert	BU
	GANCICLOVIR SODIUM				
	<u>Private hospital authority required</u>				
	<i>Cytomegalovirus retinitis in severely immunocompromised patients;</i>				
	<i>Prophylaxis of cytomegalovirus disease in bone marrow transplant patients at risk of cytomegalovirus disease;</i>				
	<i>Prophylaxis of cytomegalovirus disease in solid organ transplant patients at risk of cytomegalovirus disease.</i>				
6136Y	Powder for I.V. infusion equivalent to 500 mg ganciclovir, vial	5	280.00	Cymevene	RO

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

ILOPROST TROMETAMOL**NOTE:**

Any queries concerning the arrangements to prescribe iloprost trometamol may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe iloprost trometamol should be forwarded to:

Medicare Australia

Prior Written Approval of Specialised Drugs

Reply Paid 9826

GPO Box 9826

HOBART TAS 7001

NOTE:

Iloprost trometamol is not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma, where the total lung capacity is less than 70% of that predicted.

Iloprost trometamol is not PBS-subsidised when used in combination with PBS-subsidised bosentan monohydrate.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment with:

(a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, in patients with disease of WHO Functional Class III or IV severity; AND

(b) iloprost trometamol, of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in adult patients with disease of WHO Functional Class III or IV severity.

From 1 April 2005, adult patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma will be able to access, through the PBS, either bosentan monohydrate or iloprost trometamol. Once these patients are approved initial treatment with 1 of these 2 drugs, they may swap between bosentan monohydrate and iloprost trometamol at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary (unless the prescriber wishes to submit new baselines).

Patients may only swap to bosentan monohydrate or iloprost trometamol if they have not failed prior PBS-subsidised treatment with that drug.

Patients with pulmonary arterial hypertension secondary to connective tissue disease other than scleroderma or with drug-induced pulmonary arterial hypertension are only eligible for treatment with iloprost trometamol. They may not swap to bosentan monohydrate.

Patients aged less than 18 years with primary pulmonary hypertension are only eligible to receive treatment with bosentan monohydrate. They may qualify for treatment with iloprost trometamol when they are aged 18 years or older.

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	ILOPROST TROMETAMOL—cont.			
	<i>Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, are defined as follows:</i>			
	<i>(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or</i>			
	<i>(ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or</i>			
	<i>(iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.</i>			
	2. Definition of WHO Functional Class III or IV disease severity.			
	<i>(a) WHO Functional Class III disease severity is defined as follows:</i>			
	<i>Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.</i>			
	<i>(b) WHO Functional Class IV disease severity is defined as follows:</i>			
	<i>Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.</i>			
	3. Designated hospitals.			
	<i>Refer to Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.</i>			
	4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.			
	<i>(a) Initiation of treatment.</i>			
	<i>(i) New patients.</i>			
	<i>The first written application for PBS-subsidised treatment with the first of either bosentan monohydrate or iloprost trometamol should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an ECHO composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.</i>			
	<i>Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:</i>			
	<i>(1) RHC plus ECHO composite assessments;</i>			
	<i>(2) RHC composite assessment plus 6MWT;</i>			
	<i>(3) RHC composite assessment only.</i>			
	<i>In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:</i>			
	<i>(1) ECHO composite assessment plus 6MWT;</i>			
	<i>(2) ECHO composite assessment only.</i>			
	<i>Where fewer than 3 tests are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application.</i>			
	<i>Where patients were initiated on PBS-subsidised treatment either with bosentan monohydrate on or after 1 March 2004, or with iloprost trometamol on or after 1 April 2005, the test results provided with the initial application must be no more than 2 months old at the time of application. These results will form the baseline against which response assessments will be made.</i>			

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

ILOPROST TROMETAMOL—cont.

Where patients received treatment with either bosentan monohydrate or iloprost trometamol prior to being commenced on PBS-subsidised treatment with the first of either bosentan monohydrate or iloprost trometamol, the test requirements above still apply. The results that will form the baseline against which response assessments will be made will be those measured at the time patients commenced non-PBS-subsidised treatment with either bosentan monohydrate or iloprost trometamol, whichever of the 2 drugs the patient received first.

(ii) Patients who received non-PBS-subsidised treatment with iloprost trometamol prior to 1 April 2005.

For patients with primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on iloprost trometamol treatment prior to 1 April 2005 and who have received less than 6 months of treatment with iloprost trometamol at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with iloprost trometamol or bosentan monohydrate, whichever was initiated first.

For patients with primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on iloprost trometamol treatment prior to 1 April 2005 and who have received 6 or more months of treatment at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with iloprost trometamol or bosentan monohydrate, whichever was initiated first. The results at the time of application for initial treatment must also be provided and must be no older than 3 months.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;*
- (2) RHC plus ECHO composite assessments;*
- (3) RHC composite assessment plus 6MWT;*
- (4) ECHO composite assessment plus 6MWT;*
- (5) RHC composite assessment only;*
- (6) ECHO composite assessment only.*

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to iloprost trometamol, bosentan monohydrate or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

continued ➤

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
ILOPROST TROMETAMOL—cont.				
<i>For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.</i>				
NOTE:				
6. Authority approval requirements. <i>[The following 2 sections are only relevant to the PBS listing of iloprost trometamol. The requirements specific to bosentan monohydrate are given in parts 6 and 7 of the NOTE included in the bosentan monohydrate Schedule entry.]</i>				
(a) Initiation of PBS-subsidised treatment with iloprost trometamol, where the patient has not received prior PBS-subsidised treatment with bosentan monohydrate.				
<i>All applications for initial treatment must be made in writing.</i>				
<i>Patients who commence PBS-subsidised iloprost trometamol treatment after 1 April 2005 and patients who received 6 or more months of iloprost trometamol treatment prior to 1 April 2005 are eligible to receive up to 6 months of treatment per authority application.</i>				
<i>Patients who commenced treatment with iloprost trometamol prior to 1 April 2005 and who have received less than 6 months of treatment at the time of application are eligible to receive sufficient supply to allow the patient to complete a total of 6 months of combined PBS-subsidised and non-PBS-subsidised treatment.</i>				
<i>All patients with primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease who commenced treatment with iloprost trometamol prior to 1 April 2005 will be eligible to commence PBS-subsidised treatment with iloprost trometamol. Thereafter, to be eligible for further PBS-subsidised supply, these patients must demonstrate a response to iloprost trometamol treatment, as defined above under definition of response.</i>				
(b) Continuation of treatment.				
<i>Written applications for continuing treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment.</i>				
<i>Applications for continuing treatment will only be approved for patients who have demonstrated a response to an initial course of treatment with iloprost trometamol.</i>				
<i>The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with iloprost trometamol should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.</i>				
(c) Swapping between bosentan monohydrate and iloprost trometamol.				
<i>For eligible patients, applications to swap between these 2 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.</i>				

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	ILOPROST TROMETAMOL—cont.			
	(d) Cessation of treatment.			
	<i>Patients who fail to demonstrate a response to PBS-subsidised iloprost trometamol treatment at the times where an assessment is required must cease PBS-subsidised iloprost trometamol therapy.</i>			
	7. Re-treatment with iloprost trometamol.			
	<i>Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with iloprost trometamol under any circumstances.</i>			
	8. Further information.			
	<i>A diagrammatical representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.</i>			
	<u>Public and private hospital authority required</u>			
	<i>Application for initial PBS-subsidised treatment with iloprost trometamol of adult patients who have not received prior PBS-subsidised treatment with bosentan monohydrate and who have been assessed by a physician from a designated hospital to have:</i>			
	<i>(a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, unless a RHC is contraindicated on clinical grounds;</i>			
	OR			
	<i>(b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, unless a RHC is contraindicated on clinical grounds;</i>			
	OR			
	<i>(c) WHO Functional Class III drug-induced pulmonary arterial hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, unless a RHC is contraindicated on clinical grounds.</i>			
	<i>Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.</i>			
	<i>In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the test results of the ECHO composite assessment plus 6MWT or the ECHO composite assessment only.</i>			
	Applications for authorisation must be in writing and must include:			
	(1) a completed authority prescription form [see Note for authority approval requirements]; and			
	(2) a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes results from the 3 tests below, where available:			
	(i) RHC composite assessment; and			
	(ii) ECHO composite assessment; and			
	(iii) 6MWT; and			
	(3) a copy of a signed patient acknowledgment form indicating that the patient understands and acknowledges that PBS-subsidised treatment with iloprost trometamol for primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, OR with bosentan monohydrate for primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].			

continued ➞

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	ILOPROST TROMETAMOL—cont.			
	<i>Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient is commenced on iloprost trometamol treatment due to an adverse event or a contraindication to vasodilator treatment, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.</i>			
	<i>Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].</i>			
	<i>The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone.</i>			
	<u>Public and private hospital authority required</u>			
	<i>Application for initial PBS-subsidised treatment with iloprost trometamol of adult patients who have not received prior PBS-subsidised treatment with bosentan monohydrate and who have been assessed by a physician from a designated hospital to have:</i>			
	<i>(a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, unless a RHC is contraindicated on clinical grounds;</i>			
	OR			
	<i>(b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, unless a RHC is contraindicated on clinical grounds;</i>			
	OR			
	<i>(c) WHO Functional Class III drug-induced pulmonary arterial hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, unless a RHC is contraindicated on clinical grounds;</i>			
	OR			
	<i>(d) WHO Functional Class IV primary pulmonary hypertension;</i>			
	OR			
	<i>(e) WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease;</i>			
	OR			
	<i>(f) WHO Functional Class IV drug-induced pulmonary arterial hypertension.</i>			
	<i>In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the test results of the ECHO composite assessment plus 6MWT or the ECHO composite assessment only.</i>			
	<i>Applications for authorisation must be in writing and must include:</i>			
	<i>(1) a completed authority prescription form [see Note for authority approval requirements]; and</i>			
	<i>(2) a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes results from the 3 tests below, where available:</i>			
	<i>(i) RHC composite assessment; and</i>			
	<i>(ii) ECHO composite assessment; and</i>			
	<i>(iii) 6MWT; and</i>			

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

ILOPROST TROMETAMOL—cont.

(3) a copy of a signed patient acknowledgment form indicating that the patient understands and acknowledges that PBS-subsidised treatment with iloprost trometamol for primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, OR with bosentan monohydrate for primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone.

Public and private hospital authority required

Application for initial PBS-subsidised treatment with iloprost trometamol of adult patients who were receiving treatment with iloprost trometamol prior to 1 April 2005, who have not received prior PBS-subsidised treatment with bosentan monohydrate and who have been assessed by a physician from a designated hospital to have:

(a) WHO Functional Class III or IV primary pulmonary hypertension;

OR

(b) WHO Functional Class III or IV drug-induced pulmonary arterial hypertension;

OR

(c) WHO Functional Class III or IV pulmonary arterial hypertension secondary to connective tissue disease.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form [see Note for authority approval requirements]; and

(2) (a) for patients who have received less than 6 months of iloprost trometamol treatment at the time of application — a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes results of the following 3 tests, where available, at the time treatment with iloprost trometamol was commenced:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6MWT; or

(b) for patients who have received 6 or more months of iloprost trometamol treatment at the time of application — a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes results of the following 3 tests, both at the time treatment with iloprost trometamol was commenced and at the time of application, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6MWT; and

(3) the date of commencement of iloprost trometamol treatment; and

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

ILOPROST TROMETAMOL—cont.

(4) a copy of a signed patient acknowledgment form indicating that the patient understands and acknowledges that PBS-subsidised treatment with iloprost trometamol for primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease, OR with bosentan monohydrate for primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment. The number of repeats authorised will be dependent on the duration of prior iloprost trometamol therapy. Where patients have received less than 6 months of non-PBS-subsidised treatment with iloprost trometamol, sufficient repeats to allow the patient to complete a total of 6 months of combined PBS-subsidised and non-PBS-subsidised therapy may be requested. Where patients have received 6 months or more of non-PBS-subsidised treatment with iloprost trometamol, a maximum of 5 repeats may be requested. Where fewer than the maximum allowable number of repeats are requested at the time of application, authority approvals for the remainder of the allowable repeats may be requested by telephone.

Public and private hospital authority required

Application for initial treatment with iloprost trometamol of adult patients with either of the following:

(a) primary pulmonary hypertension, drug-induced pulmonary arterial hypertension or pulmonary arterial hypertension secondary to connective tissue disease who wish to recommence PBS-subsidised iloprost trometamol after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with iloprost trometamol;

OR

(b) primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma and whose most recent course of PBS-subsidised treatment was with bosentan monohydrate.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form [see Note for authority approval requirements]; and

(2) a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the results on which approval for the first application for PBS-subsidised iloprost trometamol or bosentan monohydrate, whichever was initiated first, was granted; and

(3) the date of the first application for PBS-subsidised treatment with iloprost trometamol or bosentan monohydrate, whichever was initiated first; and

(4) the results of the patient's response to treatment with their last course of PBS-subsidised iloprost trometamol and bosentan monohydrate.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
<i>ILOPROST TROMETAMOL—cont.</i>					
<u>Public and private hospital authority required</u>					
<i>Continuing PBS-subsidised treatment with iloprost trometamol of patients with primary pulmonary hypertension, pulmonary arterial hypertension secondary to connective tissue disease or drug-induced pulmonary arterial hypertension, who have received approval for initial PBS-subsidised treatment with iloprost trometamol, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of iloprost trometamol treatment [see Note for definition of response].</i>					
<i>Applications for authorisation must be in writing and must include:</i>					
<i>(1) a completed authority prescription form; and</i>					
<i>(2) a completed Bosentan Monohydrate (Tracleer) and Iloprost Trometamol (Ventavis) PBS Authority Application - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes results from the 3 tests below, where available:</i>					
<i>(i) RHC composite assessment; and</i>					
<i>(ii) ECHO composite assessment; and</i>					
<i>(iii) 6MWT.</i>					
<i>Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].</i>					
<i>The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone.</i>					
6456T	Solution for inhalation 20 micrograms (base) in 2 mL	30	1076.00	Ventavis	SC
<u>NOTE:</u>					
<i>This price is based on special supply arrangements—see Pharmaceutical Benefits Pricing Authority relativity sheet for full details.</i>					
<u>INDINAVIR SULFATE</u>					
<u>Private hospital authority required</u>					
<i>Treatment of HIV infection in patients with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6344X	Capsule 100 mg (base)	180	113.75	Crixivan 100 mg	MK
6201J	Capsule 200 mg (base)	360	455.00	Crixivan 200 mg	MK
6202K	Capsule 400 mg (base)	180	455.00	Crixivan 400 mg	MK

Code	Item and Section 100 Restriction	Pack Size	Price ex	Proprietary
			Manufacturer \$	Name and Manufacturer

INFLIXIMAB**NOTE:**

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe infliximab should be forwarded to:

*Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001*

Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au.

NOTE:

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of etanercept and infliximab for adult patients with HLA-B27 positive, active ankylosing spondylitis only. It does not apply to the prescribing of adalimumab. Therefore, where the term 'tumour necrosis factor (TNF) alfa antagonists' appears in the following NOTES and restrictions, it only refers to etanercept and infliximab.

Patients are eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alfa antagonists above at any 1 time.

From 1 August 2005, under the PBS, all patients will be able to commence a single treatment cycle where they may trial both of the PBS-subsidised TNF-alfa antagonists without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, patients may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

Once patients have either failed or ceased to respond to treatment up to 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. Within the same treatment cycle, patients are not allowed to trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than twice. Patients who have failed the same PBS-subsidised TNF-alfa antagonist twice must change to the alternate PBS-subsidised agent if they wish to continue PBS-subsidised TNF-alfa antagonist treatment.

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised TNF-alfa antagonist treatment was granted in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new cycle.

Where a period of less than 5 years duration has elapsed since the patient's previous course of PBS-subsidised TNF-alfa antagonist treatment, patients who have failed TNF-alfa antagonist therapy less than 3 times within a particular treatment cycle, as defined in the restriction, may commence a further course of treatment within that cycle.

Where a break in PBS-subsidised therapy of 5 years or more has occurred, patients who have failed TNF-alfa antagonist therapy less than 3 times within a particular treatment cycle, as defined in the restriction, are eligible to commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

INFLIXIMAB—cont.**1. Patients 'grandfathered' onto PBS-subsidised treatment with infliximab and etanercept.**

Both the infliximab and etanercept restrictions allow patients who commenced treatment prior to 1 March 2004 and 1 July 2004 respectively to commence treatment under a different set of criteria than those patients who commenced treatment after these dates (i.e. new patients). This means such patients were 'grandfathered' onto PBS-subsidised treatment. Consequently, the criteria to enable 'grandfather' patients to continue on PBS-subsidised treatment are also different to those for new patients.

Where patients who were 'grandfathered' onto PBS-subsidised treatment with 1 TNF- α antagonist wish to swap to the alternate agent, they can only do so using the restrictions that apply to 'grandfather' patients.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, 'grandfather' patients must requalify for initial treatment under the criteria that apply to new patients. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below for further details. Patients who commenced on PBS-subsidised etanercept or infliximab treatment prior to 1 August 2005 are deemed to be in their first cycle of therapy.

The first course of initial therapy for patients 'grandfathered' onto PBS-subsidised treatment with etanercept or infliximab will be limited to provide for 24 weeks of therapy. All subsequent initial treatment courses made under the 'grandfather' restriction will be limited to provide 6 weeks of therapy.

2. How to prescribe TNF- α antagonist therapy after 1 August 2005.**(a) Initial treatment.**

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised TNF- α antagonist treatment and wish to commence such therapy; or*
- (ii) patients have received prior PBS-subsidised (initial or continuing) TNF- α antagonist therapy and wish to trial an alternate agent [further details are under 'Swapping therapy' below]; or*
- (iii) patients have failed their most recent course of PBS-subsidised TNF- α antagonist therapy but wish to trial a further course of treatment with the same TNF- α antagonist (providing they have not failed that TNF- α antagonist more than once); or*
- (iv) patients wish to re-commence treatment with a specific TNF- α antagonist following a break in PBS-subsidised therapy with that specific agent.*

All applications for initial treatment will be limited to provide for a maximum of 6 weeks of therapy except for the first application for 'grandfather' patients approved for PBS-subsidised treatment with etanercept and infliximab where up to 24 weeks of treatment will be approved.

In order to demonstrate a response to therapy, patients who have received an initial PBS-subsidised course of up to 6 weeks of treatment must be assessed no earlier than 4 weeks from the commencement of the initial therapy course. In order to demonstrate a response to therapy, patients who have received an initial PBS-subsidised course of up to 24 weeks must be assessed in the 4 weeks prior to treatment being ceased.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that TNF- α antagonist.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	INFLIXIMAB—cont.			
	<i>Prescribers should ensure that applications for second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment are submitted to Medicare Australia before patients complete their previous treatment course to ensure uninterrupted treatment. Applications where no provisions to fax the prescription and application are made in the restriction should be posted to Medicare Australia no later than 2 weeks prior to the patient completing their previous treatment course.</i>			
	(b) Continuing treatment.			
	<i>Following the completion of an initial treatment course with a specific TNF-alfa antagonist, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.</i>			
	<i>Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.</i>			
	<i>Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.</i>			
	(c) How to apply for TNF-alfa antagonist treatment on or after 1 August 2005.			
	<i>Patients for whom the first application for PBS-subsidised treatment with either etanercept or infliximab is made on or after 1 August 2005, will fall into 1 of the following categories:</i>			
	(I) Patients who have not received any prior PBS-subsidised infliximab or PBS-subsidised etanercept treatment.			
	<i>This includes the following:</i>			
	(i) Patients who have not received any prior treatment with infliximab or etanercept for ankylosing spondylitis.			
	(ii) Patients who have received non-PBS-subsidised treatment with infliximab after 1 March 2004.			
	(iii) Patients who have received non-PBS-subsidised treatment with etanercept after 1 July 2004.			
	(iv) Patients who have received non-PBS-subsidised infliximab treatment prior to 1 March 2004.			
	(v) Patients who have received non-PBS-subsidised etanercept treatment prior to 1 July 2004.			
	<i>To commence PBS-subsidised treatment with etanercept or infliximab, all patients who fall into groups (i)-(iii) must qualify under the first initial restriction for new patients (i.e. the patient must meet the criteria with respect to the indices of severity and prior treatment regimens specified in the 'new patient' restriction).</i>			
	<i>To commence PBS-subsidised treatment with infliximab, all patients who fall into group (iv) must qualify under the first initial restriction for 'grandfather' patients (i.e. the patient must meet the criteria with respect to the indices of severity and prior treatment regimens specified in the 'grandfather' restriction).</i>			
	<i>To commence PBS-subsidised treatment with etanercept, all patients who fall into group (v) must qualify under the first initial restriction for 'grandfather' patients (i.e. the patient must meet the criteria with respect to the indices of severity and prior treatment regimens specified in the 'grandfather' restriction).</i>			

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

INFLIXIMAB—cont.

Patients in groups (iv) and (v) can not be 'grandfathered' onto treatment with the alternate agent. If they wish to commence PBS-subsidised therapy with the alternate agent, they must qualify under the first initial restriction for new patients (i.e. the patient must meet the criteria with respect to the indices of severity and prior treatment regimens specified in the 'new patient' restriction).

(II) Patients who have received PBS-subsidised etanercept only.

(III) Patients who have received PBS-subsidised infliximab only.

(IV) Patients who have received PBS-subsidised etanercept and non-PBS-subsidised infliximab.

(V) Patients who have received PBS-subsidised infliximab and non-PBS-subsidised etanercept.

To receive PBS-subsidised treatment with etanercept or infliximab, the patients in groups (II)-(V) do not need to requalify with respect to the indices of severity or prior treatment requirements. They use the second initial restriction for either 'new' or 'grandfather' patients (i.e. whichever was applicable for their first course of PBS-subsidised treatment).

NOTE:

Patients who have responded to their most recent course of etanercept therapy as specified in the restriction may:

- continue on etanercept treatment under the 'grandfather' or new patient restriction, whichever is applicable [see 'Patients 'grandfathered' onto PBS-subsidised treatment with infliximab and etanercept']; or*
- swap to an initial course of infliximab treatment of up to 6 weeks under the 'grandfather' or new patient restriction, whichever is applicable [see 'Patients 'grandfathered' onto PBS-subsidised treatment with infliximab and etanercept'].*

Patients who have failed to respond to their most recent course of etanercept treatment may:

- trial a further course of initial therapy of up to 6 weeks with etanercept; or*
- swap to an initial course of infliximab treatment of up to 6 weeks.*

Patients who have responded to their most recent course of infliximab therapy as specified in the restriction may:

- continue on infliximab treatment under the 'grandfather' or new patient restriction, whichever is applicable [see 'Patients 'grandfathered' onto PBS-subsidised treatment with infliximab and etanercept']; or*
- swap to an initial course of etanercept treatment of up to 6 weeks under the 'grandfather' or new patient restriction, whichever is applicable [see 'Patients 'grandfathered' onto PBS-subsidised treatment with infliximab and etanercept'].*

Patients who have failed to respond to their most recent course of infliximab treatment may:

- trial a further course of initial therapy of up to 6 weeks with infliximab; or*
- swap to an initial course of etanercept treatment of up to 6 weeks.*

3. Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised TNF- α antagonist is approved, patients may swap to the alternate TNF- α antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements. This also applies to patients who fail to achieve or sustain a response to the first PBS-subsidised TNF- α antagonist approved and who wish to trial a further course of treatment with the same TNF- α antagonist.

Code	Item and Section 100 Restriction	Pack Size	Price ex	Proprietary
			Manufacturer \$	Name and Manufacturer

INFLIXIMAB—cont.

Patients may trial an alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application or not. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that particular drug more than twice in the same cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

4. Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

For new patients, the first BASDAI result used to determine baseline must be provided while the patient is still receiving NSAID therapy. However, this is not required for any subsequent BASDAI results provided for these patients nor for patients who were 'grandfathered' onto TNF-alfa antagonist treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response.

5. Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with at least 1 NSAID, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the BASDAI, ESR and/or CRP levels are measured.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

INFLIXIMAB—cont.**Public and private hospital authority required*****Initial 1 (new patients)***

Application for the first course of initial PBS-subsidised treatment with infliximab, by a rheumatologist, of adults with active ankylosing spondylitis who have radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, who have not received any PBS-subsidised treatment with either etanercept or infliximab;

AND

(a) have at least 2 of the following:

(i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or

(ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI) [for further information on the BASMI please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; or

(iii) limitation of chest expansion relative to normal values for age and gender [for chest expansion normal values please refer to the Medicare Australia website at www.medicareaustralia.gov.au];

AND

(b) who have documented confirmation of HLA-B27 positive status;

AND

(c) who have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs) in combination with an appropriate exercise program for a total period of 3 months.

The application must include details of the NSAIDs trialled, their doses and duration of treatment. If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reasons why a higher dose cannot be used.

For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Medicare Australia website at www.medicareaustralia.gov.au.

Patients who received treatment with infliximab prior to 1 March 2004 and wish to commence PBS-subsidised treatment with infliximab must qualify under the initial treatment restriction for 'grandfather' patients.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrable in all patients at the time of the initial application:

(a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale;

AND

(b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

INFLIXIMAB—cont.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the patient is exempted from demonstrating an inadequate response to the above treatment regimen. Where appropriate, evidence to support a contraindication must be provided. If adverse events of a severity necessitating permanent withdrawal develop during the relevant period of use of 2 NSAIDs, the patient may be exempted from demonstrating an inadequate response to the above treatment regimen. For details of the adverse events, including the severity, which will be accepted for the purposes of administering this restriction, please refer to the Medicare Australia website at www.medicareaustralia.gov.au.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and*
 - (b) a completed TNF-alfa antagonist PBS Authority Application for Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes the following:*
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and*
 - (ii) a copy of the pathology report from an Approved Pathology Authority confirming the presence of HLA-B27; and*
 - (iii) a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)];*
 - (iv) a copy of the signed patient acknowledgment form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)]*
- Completion of this form declares that the patient understands and acknowledges that PBS-subsidised treatment with the TNF-alfa antagonists (etanercept or infliximab) for ankylosing spondylitis will cease if they do not demonstrate the response to treatment required to support continuation of PBS-subsidised treatment at any assessment where a response must be demonstrated.*

The assessment of the patient's response to the initial course of treatment must be made no earlier than 4 weeks from the commencement of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A maximum of 6 weeks of treatment with infliximab will be approved under this criterion.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for 2 infusions at a dose of 5 mg per kg. No repeats will be authorised.

Public and private hospital authority required**Initial 2 (new patients)**

Application for an initial course of PBS-subsidised treatment with infliximab, by a rheumatologist, of adults with active ankylosing spondylitis who, in this treatment cycle, have received prior PBS-subsidised treatment with either etanercept or infliximab for this condition and have not failed PBS-subsidised therapy with infliximab more than once.

All applications for treatment must be accompanied by the results of the most recent course of TNF-alfa antagonist therapy within the timeframes specified below.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

INFLIXIMAB—cont.

Where the most recent course of PBS-subsidised TNF- α antagonist treatment was approved under an initial treatment restriction for new patients, patients must have been assessed for response to that course no earlier than 4 weeks from the commencement of that course. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised TNF- α antagonist treatment was approved under the continuing treatment criteria for new patients, patients must have been assessed for response to that course, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessments to the previous course of TNF- α antagonist treatment are not submitted as detailed above, patients will be deemed to have failed therapy with that particular course of TNF- α antagonist.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and*
- (b) a completed TNF- α antagonist PBS Authority Application for Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].*

A maximum of 6 weeks of treatment with infliximab will be approved under this criterion.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for 2 infusions at a dose of 5 mg per kg. No repeats will be authorised.

Public and private hospital authority required

Continuing (new patients)

Continuing PBS-subsidised treatment, by a rheumatologist, of adults with active ankylosing spondylitis who:

- (a) have received 6 weeks or more of PBS-subsidised treatment with infliximab; and*
- (b) have demonstrated a response to treatment with infliximab; and*
- (c) whose most recent course of PBS-subsidised therapy in this treatment cycle was with infliximab.*

Response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or*
- (b) a CRP measurement no greater than 10 mg per L; or*
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.*

The same acute phase reactant measured at the relevant baseline must be measured in all subsequent continuing treatment applications.

The first application for continuing treatment following an initial treatment course must be made no earlier than 4 weeks from the commencement of the most recent initial treatment course with infliximab. This first authority application and a copy of the authority prescription may be faxed to Medicare Australia on (03) 6215 5640 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) in order to seek approval for a maximum of 4 weeks' supply. Medicare Australia will then contact the prescriber by telephone. The original document must then be posted to Medicare Australia with a second authority prescription for the balance of 24 weeks of treatment.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

INFLIXIMAB—cont.

Second and subsequent applications for continuing treatment must be made in writing and should be posted to Medicare Australia no less than 2 weeks prior to the completion of the current treatment course.

Written applications for authorisation must include:

- (a) a completed authority prescription form; and*
- (b) a completed TNF-alfa antagonist PBS Authority Application for Continuing Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes the following:*
 - (i) a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] including certification by the prescriber and the patient that the patient did not have access to their baseline BASDAI at the time of their continuing treatment assessment.*

All measurements provided must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with infliximab will be authorised under this criterion.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.

Public and private hospital authority required

Initial 1 ('grandfather' patients)

Application for an initial PBS-subsidised course of continuing treatment with infliximab, by a rheumatologist, of adults with active ankylosing spondylitis who have radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, who were receiving treatment with infliximab prior to 1 March 2004;

AND

(a) are receiving treatment with infliximab at the time of application;

AND

(b) who have not received prior PBS-subsidised treatment with etanercept;

AND

(c) who have documented confirmation of HLA-B27 positive status;

AND

(d) whose Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score is less than or equal to 5 on a 0-10 scale;

AND

(e) who have:

- (i) an ESR measurement no greater than 25 mm per hour; or*
- (ii) a CRP measurement no greater than 10 mg per L; or*
- (iii) an ESR or CRP measurement reduced by at least 20% from pre-treatment baseline.*

The BASDAI assessment and ESR and/or CRP measurements must be no more than 1 month old at the time of application. The same acute phase reactant measured in the first application for PBS-subsidised treatment must be measured in all subsequent continuing treatment applications.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

INFLIXIMAB—cont.

Authority applications must be made in writing and must include:

*(a) a completed authority prescription form; and
(b) a completed TNF- α antagonist PBS Authority Application for Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website*

(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes the following:

(i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and

(ii) a copy of the pathology report from an Approved Pathology Authority confirming the presence of HLA-B27; and

(iii) a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website

(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)]; and

(iv) a copy of the signed patient acknowledgment form [may be downloaded from the Medicare Australia website

(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].

Completion of this form declares that the patient understands and acknowledges that PBS-subsidised treatment with the TNF- α antagonists (etanercept or infliximab) for ankylosing spondylitis will cease if they do not demonstrate the response to treatment required to support continuation of PBS-subsidised treatment at any assessment where a response must be demonstrated.

The assessment of the patient's response to this initial course of therapy must be made within the 4 weeks prior to completion of the course in order to ensure continuity of treatment.

Those patients ceasing treatment or swapping to an alternate agent and wishing to demonstrate a response to treatment, must be assessed no earlier than 4 weeks from the commencement of PBS-subsidised treatment. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A maximum of 24 weeks of treatment with infliximab will be authorised under this criterion.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.

Patients may only qualify for PBS-subsidised treatment under this criterion once.

Public and private hospital authority required

Initial 2 ('grandfather' patients)

Application for an initial course of PBS-subsidised treatment with infliximab, by a rheumatologist, of adults with active ankylosing spondylitis who have radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, who were 'grandfathered' onto PBS-subsidised treatment with etanercept or infliximab and who have not failed PBS-subsidised therapy with infliximab more than once.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

INFLIXIMAB—cont.

Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under an initial treatment restriction for 'grandfather' patients, patients must have been assessed for response to that course no earlier than 4 weeks from the commencement of that course. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under the continuing treatment criteria for 'grandfather' patients, patients must have been assessed for response to that course, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessments to the previous course of TNF-alfa antagonist treatment are not submitted as detailed above, patients will be deemed to have failed therapy with that particular course of TNF-alfa antagonist.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and*
- (b) a completed TNF-alfa antagonist PBS Authority Application for Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].*

A maximum of 6 weeks of treatment with infliximab will be approved under this criterion.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for 2 infusions at a dose of 5 mg per kg. No repeats will be authorised.

Public and private hospital authority required

Continuing ('grandfather' patients)

Continuing PBS-subsidised treatment, by a rheumatologist, of adults with active ankylosing spondylitis who were 'grandfathered' onto PBS-subsidised treatment with a TNF-alfa antagonist and who:

- (a) have received 6 weeks or more of PBS-subsidised treatment with infliximab; and*
- (b) have demonstrated a response to treatment with infliximab; and*
- (c) whose most recent course of PBS-subsidised therapy in this treatment cycle was with infliximab.*

Response to treatment is defined as a BASDAI score no more than 20% greater than the score included in the initial application for PBS-subsidised treatment;

AND

- (a) an ESR measurement no greater than 25 mm per hour; or*
- (b) a CRP measurement no greater than 10 mg per L; or*
- (c) an ESR or CRP measurement reduced by at least 20% from pre-treatment baseline.*

The same acute phase reactant measured at the relevant baseline must be measured in all subsequent continuing treatment applications.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	INFLIXIMAB—cont.			
	<i>The first application for continuing treatment following an initial treatment course must be made no earlier than 4 weeks from the commencement of the most recent initial treatment course with infliximab. This first authority application and a copy of the authority prescription may be faxed to Medicare Australia on (03) 6215 5640 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) in order to seek approval for a maximum of 4 weeks' supply. Medicare Australia will then contact the prescriber by telephone. The original document must then be posted to Medicare Australia with a second authority prescription for the balance of 24 weeks of treatment.</i> <i>Second and subsequent applications for continuing treatment must be made in writing and should be posted to Medicare Australia no less than 2 weeks prior to the completion of the current treatment course.</i> <i>In order to demonstrate a response to treatment where the patient is ceasing or swapping to an alternate agent, the assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.</i> <i>Written applications for authorisation must include:</i> <i>(a) a completed authority prescription form; and</i> <i>(b) a completed TNF-alfa antagonist PBS Authority Application for Continuing Use in the Treatment of Ankylosing Spondylitis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes the following:</i> <i>(i) a copy of the completed BASDAI Assessment Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] including certification by the prescriber and the patient that the patient did not have access to their baseline BASDAI at the time of their continuing treatment assessment.</i> <i>All measurements provided must be no more than 1 month old at the time of application.</i> <i>A maximum of 24 weeks of treatment with infliximab will be authorised under this criterion.</i> <i>At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised. No applications for increased repeats will be authorised.</i> <i>Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.</i>			
6448J	Powder for I.V. infusion 100 mg	1	875.00	Remicade SH

Code	Item and Section 100 Restriction	Price ex		Proprietary Name and Manufacturer
		Pack Size	Manufacturer \$	

INFLIXIMAB—cont.**NOTE:**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the tumour necrosis factor (TNF) alfa antagonists (adalimumab, etanercept and infliximab) and the interleukin-1 inhibitor (anakinra) for adult patients with severe rheumatoid arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs (bDMARDs) at any 1 time.

1. Patients who have received no prior PBS-subsidised bDMARD treatment at 1 December 2004.

From 1 December 2004, the arrangements for prescribing the bDMARDs on the PBS have been amended to allow patients to commence a single cycle of bDMARD treatment that allows them to trial any number of bDMARDs without having to experience a disease flare when swapping between alternate bDMARDs. Within a single treatment cycle, patients may continue to receive long-term treatment with a bDMARD while they continue to show a response to therapy.

Once patients have either failed, or ceased to respond to, treatment with a maximum of 3 bDMARDs, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised bDMARD therapy before they are eligible to commence the next cycle. The 5-year period will be measured from the date the last prescription for PBS-subsidised bDMARD treatment was approved in the most recent cycle to the date of the application for initial treatment with a bDMARD under the new cycle.

Patients who have failed treatment with fewer than 3 bDMARDs within a particular treatment cycle, and where a period of less than 5 years duration has elapsed since the patient's previous course of PBS-subsidised bDMARD treatment in that cycle, may commence a further course of bDMARD treatment within that same treatment cycle.

Patients who have failed treatment with fewer than 3 bDMARDs within a particular treatment cycle, and who have had a break in PBS-subsidised therapy of 5 years or more, are eligible to commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

If patients fail to respond to a particular bDMARD within a single treatment cycle, they are not eligible to receive further PBS-subsidised treatment with that drug until they commence the next cycle.

2. Patients who have received PBS-subsidised TNF-alfa antagonist treatment prior to 1 December 2004.

Patients who commenced PBS-subsidised TNF-alfa antagonist therapy prior to 1 December 2004 are considered to be in their first cycle of bDMARD treatment on 1 December 2004.

Patients who have failed to respond to prior PBS-subsidised treatment with fewer than 3 TNF-alfa antagonists at the time the first application for treatment is made on or after 1 December 2004, will be subject to the same conditions applying to new patients detailed above.

Code	Item and Section 100 Restriction	Price ex		Proprietary Name and Manufacturer
		Pack Size	Manufacturer \$	

INFLIXIMAB—cont.

Therefore, patients who have failed:

(a) 1 TNF-alfa antagonist will be eligible to trial further PBS-subsidised treatment with a bDMARD they have not failed in their first treatment cycle, until they fail to demonstrate a response to no more than another 2 bDMARDs;

(b) 2 TNF-alfa antagonists will be eligible to trial further PBS-subsidised treatment with a bDMARD they have not failed in their first treatment cycle, until they fail to demonstrate a response to no more than 1 other bDMARD.

Patients who have failed PBS-subsidised treatment with 3 TNF-alfa antagonists prior to 1 December 2004 or at the first assessment required after this date, will be eligible to trial PBS-subsidised treatment with anakinra if they wish. However, if they fail to demonstrate a response to anakinra, they will not be able to trial any further PBS-subsidised bDMARD treatment until a minimum of 5 years has elapsed from the date that the prescription for the last course of anakinra therapy was approved. Arrangements to allow these patients to fail 4 bDMARDs will only be in place for the first treatment cycle. For subsequent cycles, patients will cease to be eligible to receive PBS-subsidised bDMARD treatment once they have failed to demonstrate a response to a maximum of 3 bDMARDs.

Any queries on these arrangements should be forwarded to Medicare Australia.

3. Information relevant to all patients.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) patients have received no prior PBS-subsidised bDMARD treatment and wish to commence such therapy; or

(ii) patients have received prior PBS-subsidised (initial or continuing) bDMARD therapy and wish to trial an alternate agent [further details are under 'Swapping therapy' below]; or

(iii) patients wish to re-commence treatment with a specific bDMARD following a break in PBS-subsidised therapy with that specific agent.

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted bDMARD supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that bDMARD.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that bDMARD.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	INFLIXIMAB—cont.			
	(c) Swapping therapy.			
	<i>Once an authority for initial treatment with the first PBS-subsidised bDMARD is approved, patients may swap to an alternate bDMARD without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-bDMARD therapy requirements. However, the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialed.</i>			
	<i>Patients may swap to an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application or not.</i>			
	<i>Patients may alternate between therapy with any bDMARD of their choice (1 at a time) providing:</i>			
	<i>(i) they have not received PBS-subsidised treatment with that particular bDMARD previously; or</i>			
	<i>(ii) they have demonstrated an adequate response to that particular bDMARD if they have previously trialed it on the PBS.</i>			
	<i>Therefore, to maximise the choice of bDMARD patients may alternate between, it is important that patients are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.</i>			
	<i>To avoid confusion, applications for patients who wish to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.</i>			
	(d) Baseline measurements to determine response.			
	<i>Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.</i>			
	<i>To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to a reduction in the total number of active joints.</i>			
	(e) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.			
	<i>Patients who wish to trial a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with at least 1 non-biological DMARD, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.</i>			

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	<i>INFLIXIMAB—cont.</i>			
	<i>Public and private hospital authority required</i>			
	<i>Application for initial PBS-subsidised treatment with infliximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:</i>			
	<i>(a) have severe active rheumatoid arthritis; and</i>			
	<i>(b) have received no prior PBS-subsidised treatment with a bDMARD for this condition in this treatment cycle; and</i>			
	<i>(c) have failed to achieve an adequate response to the following treatments:</i>			
	<i>(i) methotrexate at a dose of at least 20 mg weekly; and</i>			
	<i>(ii) methotrexate (at a minimum dose of 7.5 mg weekly), in combination with 2 other non-biological disease modifying anti-rheumatic drugs (DMARDs), for a minimum of 3 months; and</i>			
	<i>(iii) a minimum of 3 months' treatment with:</i>			
	<i>— leflunomide alone; or</i>			
	<i>— leflunomide in combination with methotrexate; or</i>			
	<i>— cyclosporin.</i>			
	<i>If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or intolerance of a severity necessitating permanent treatment withdrawal develops during the relevant period of use, the patient is exempted from demonstrating an inadequate response to that particular agent(s) only. Details of the contraindications or intolerance, including the degree of toxicity, must be provided at the time of application.</i>			
	<i>The following initiation criteria indicate failure to achieve an adequate response and must be demonstrable in all patients at the time of the initial application:</i>			
	<i>an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L;</i>			
	<i>AND either</i>			
	<i>(i) a total active joint count of at least 20 active (swollen and tender) joints; or</i>			
	<i>(ii) at least 4 active joints from the following list of major joints:</i>			
	<i>— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or</i>			
	<i>— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).</i>			
	<i>If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.</i>			
	<i>The authority application must be made in writing and must include:</i>			
	<i>(1) a completed authority prescription form; and</i>			
	<i>(2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)] which includes details of the patient's ESR and CRP measurements and the patient's active joint count which must have been assessed no earlier than 1 month prior to the date of application; and</i>			
	<i>(3) a copy of the signed patient acknowledgement form which may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm).</i>			
	<i>Completion of this form declares that the patient understands and acknowledges that, within a single treatment cycle, PBS-subsidised treatment with any biological DMARD will cease if they do not demonstrate the response to treatment required to support continuation of PBS-subsidised treatment at any assessment where a response must be demonstrated.</i>			

continued 

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

INFLIXIMAB—cont.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 3 mg per kg. Up to a maximum of 3 repeats may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Public and private hospital authority required

Application for an initial course of PBS-subsidised treatment with infliximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and*
- (b) have received prior PBS-subsidised bDMARD treatment for this condition in this treatment cycle and are eligible to receive further bDMARD therapy.*

Patients who were commenced on treatment with infliximab prior to 1 December 2004 and who have received methotrexate at a dose of less than 7.5 mg per week will be able to continue to receive PBS-subsidised treatment with infliximab in combination with methotrexate at this lower dose for the duration of the first bDMARD treatment cycle. For subsequent treatment cycles, patients must receive concomitant methotrexate at a dose of at least 7.5 mg weekly.

Applications for patients who have received PBS-subsidised treatment with infliximab within this treatment cycle and who wish to re-commence therapy with this drug within this same cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised infliximab treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised infliximab treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised bDMARD therapy or, under this restriction, for patients who have received previous PBS-subsidised bDMARD therapy), patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised infliximab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and*
- (2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].*

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer

INFLIXIMAB—cont.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 3 mg per kg. Up to a maximum of 3 repeats may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Public and private hospital authority required

Initial PBS-subsidised supply for continuing treatment with infliximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and*
- (b) were receiving treatment with infliximab prior to 1 March 2005; and*
- (c) failed to qualify for PBS-subsidised therapy after 1 November 2003 due to testing negative for rheumatoid factor; and*
- (d) have demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with infliximab.*

Medical practitioners who wish to apply for authority to prescribe a bDMARD for patients who commenced treatment with infliximab prior to 1 March 2005 and who have not demonstrated a response to treatment should contact Medicare Australia on 1800 005 750.

The authority application must be in writing and must include sufficient information to determine the patient's eligibility. The date of assessment must be provided.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 3 mg per kg. Up to a maximum of 2 repeats may be authorised.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	<i>INFLIXIMAB—cont.</i>			
	<u>Public and private hospital authority required</u>			
	<i>Continuing PBS-subsidised treatment with infliximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:</i>			
	<i>(a) who have severe active rheumatoid arthritis; and</i>			
	<i>(b) who have demonstrated an adequate response to treatment with infliximab; and</i>			
	<i>(c) whose most recent course of PBS-subsidised bDMARD treatment in this treatment cycle was with infliximab.</i>			
	<i>An adequate response to treatment is defined as:</i>			
	<i>an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;</i>			
	<i>AND either of the following:</i>			
	<i>(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or</i>			
	<i>(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:</i>			
	<i>— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or</i>			
	<i>— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).</i>			
	<i>Patients who were commenced on treatment with infliximab prior to 1 December 2004 and who have received methotrexate at a dose of less than 7.5 mg per week will be able to continue to receive PBS-subsidised treatment with infliximab in combination with methotrexate at this lower dose for the duration of the first bDMARD treatment cycle. For subsequent treatment cycles, patients must receive concomitant methotrexate at a dose of at least 7.5 mg weekly.</i>			
	<i>The authority application must be made in writing and must include:</i>			
	<i>(1) a completed authority prescription form; and</i>			
	<i>(2) a completed Biological DMARD PBS Authority Application for Use in the Treatment of Rheumatoid Arthritis - Supporting Information Form [may be downloaded from the Medicare Australia website</i>			
	<i>(www.medicareaustralia.gov.au/providers/forms/pbs/medical_practitioners.htm)].</i>			
	<i>All applications for continuing treatment with infliximab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with infliximab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.</i>			
	<i>At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 3 mg per kg. Up to a maximum of 2 repeats may be authorised.</i>			
	<i>Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 005 750 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).</i>			

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	INFLIXIMAB—cont.				
	<i>Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this treatment cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.</i>				
	<i>Once patients fail to respond to treatment with 3 bDMARDs, they are deemed to have completed this treatment cycle and must cease PBS-subsidised therapy. These patients may re-commence a new bDMARD treatment cycle after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised bDMARD was approved in this cycle and the date of the first application under the new cycle.</i>				
6397Q	Powder for I.V. infusion 100 mg	1	875.00	Remicade	SH
	INTERFERON ALFA-2a				
	CAUTION:				
	<i>Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.</i>				
	Private hospital authority required				
	<i>Use in the treatment of Philadelphia chromosome positive myelogenous leukaemia in the chronic phase;</i>				
	<i>Patients with chronic hepatitis B who satisfy all of the following criteria:</i>				
	<i>(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);</i>				
	<i>(2) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection (HBe antigen positive and/or HBV DNA positive);</i>				
	<i>(3) Are not persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L);</i>				
	<i>(4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception;</i>				
	<i>Patients with chronic hepatitis C who satisfy all of the following criteria:</i>				
	<i>(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);</i>				
	<i>(2) Abnormal serum ALT levels in conjunction with documented chronic hepatitis C infection (repeatedly anti-HCV positive and/or HCV RNA positive);</i>				
	<i>(3) No other forms of chronic liver disease;</i>				
	<i>(4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.</i>				
	<i>The treatment course is limited to 3 million units subcutaneously 3 times weekly for up to 52 weeks.</i>				
	<i>Treatment is to cease if plasma HCV RNA remains detectable by an HCV RNA qualitative assay after 12 weeks of therapy.</i>				
	<i>The course of treatment must be continuous and excludes retreatment of nonresponders or patients who relapse.</i>				
	NOTE:				
	<i>Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:</i>				
	<i>(a) a nurse educator/counsellor for patients; and</i>				
	<i>(b) 24 hour access by patients to medical advice; and</i>				
	<i>(c) an established liver clinic; and</i>				
	<i>(d) facilities for safe liver biopsy.</i>				
6210W	Injection 3,000,000 i.u. in 0.5 mL single dose pre-filled syringe	1	29.80	Roferon-A	RO

continued ➡

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	INTERFERON ALFA-2a—cont.				
6211X	Injection 4,500,000 i.u. in 0.5 mL single dose pre-filled syringe	1	44.70	Roferon-A	RO
6212Y	Injection 6,000,000 i.u. in 0.5 mL single dose pre-filled syringe	1	59.58	Roferon-A	RO
6213B	Injection 9,000,000 i.u. in 0.5 mL single dose pre-filled syringe	1	89.38	Roferon-A	RO

INTERFERON ALFA-2b**CAUTION:**

Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Private hospital authority required

Adjunctive therapy of malignant melanoma following surgery in patients with nodal involvement;

Use in the treatment of Philadelphia chromosome positive myelogenous leukaemia in the chronic phase;

Patients with chronic hepatitis B who satisfy all of the following criteria:

- (1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);*
- (2) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection (HBe antigen positive and/or HBV DNA positive);*
- (3) Are not persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L);*
- (4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception;*

Patients with chronic hepatitis C who satisfy all of the following criteria:

- (1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);*
- (2) Abnormal serum ALT levels in conjunction with documented chronic hepatitis C infection (repeatedly anti-HCV positive and/or HCV RNA positive);*
- (3) No other forms of chronic liver disease;*
- (4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.*

The treatment course is limited to 3 million units subcutaneously 3 times weekly for up to 52 weeks.

Treatment is to cease if plasma HCV RNA remains detectable by an HCV RNA qualitative assay after 12 weeks of therapy.

The course of treatment must be continuous and excludes retreatment of nonresponders or patients who relapse.

NOTE:

Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:

- (a) a nurse educator/counsellor for patients; and*
- (b) 24 hour access by patients to medical advice; and*
- (c) an established liver clinic; and*
- (d) facilities for safe liver biopsy.*

6246R	Solution for injection 10,000,000 i.u. in 1 mL single dose vial	5	496.50	Intron A	SH

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
INTERFERON ALFA-2b—cont.					
6253D	Solution for injection 18,000,000 i.u. in 1.2 mL multi-dose injection pen	1	178.74	Intron A Redipen	SH
6218G	Solution for injection 18,000,000 i.u. in 3 mL single dose vial	1	178.74	Intron A	SH
6219H	Solution for injection 25,000,000 i.u. in 2.5 mL single dose vial	1	248.25	Intron A	SH
6254E	Solution for injection 30,000,000 i.u. in 1.2 mL multi-dose injection pen	1	297.90	Intron A Redipen	SH
6255F	Solution for injection 60,000,000 i.u. in 1.2 mL multi-dose injection pen	1	595.80	Intron A Redipen	SH
INTERFERON GAMMA-1b					
<u>Private hospital authority required</u>					
<i>Treatment of chronic granulomatous disease in patients with frequent and severe infections despite adequate prophylaxis with antimicrobial agents.</i>					
6148N	Injection 2,000,000 i.u. in 0.5 mL vial	6	1188.76	Imukin	BY
LAMIVUDINE					
<u>Private hospital authority required</u>					
<i>Patients with chronic hepatitis B who satisfy all of the following criteria:</i>					
<i>(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);</i>					
<i>(2) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection (HBe antigen positive and/or HBV DNA positive);</i>					
<i>(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.</i>					
<i>Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.</i>					
6257H	Tablet 100 mg	28	119.50	Zeffix	GK
6271C	Oral solution 5 mg per mL, 240 mL	1	51.21	Zeffix	GK
<u>Private hospital authority required</u>					
<i>Treatment of HIV infection in patients with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6193Y	Tablet 150 mg	60	282.00	3TC	GK
6435Q	Tablet 300 mg	30	282.00	3TC	GK
6194B	Oral solution 10 mg per mL, 240 mL	1	75.20	3TC	GK
LAMIVUDINE with ZIDOVUDINE					
<u>Private hospital authority required</u>					
<i>Treatment of HIV infection in patients with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6234D	Tablet 150 mg-300 mg	60	578.60	Combivir	GK

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	LANREOTIDE ACETATE				
	<u>Private hospital authority required</u>				
	<i>Active acromegaly in patients with persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre AND</i>				
	<i>(a) after failure of other therapy including dopamine agonists; or</i>				
	<i>(b) as interim treatment in patients awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; or</i>				
	<i>(c) where surgery and radiotherapy are contraindicated.</i>				
	<i>Treatment is to cease in patients previously treated with radiotherapy where there is biochemical evidence of remission (normal IGF1) after lanreotide acetate withdrawal for at least 4 weeks (6 weeks after the last dose). Lanreotide acetate should be withdrawn for assessment of remission every 2 years in the 10 years after radiotherapy.</i>				
	<i>Treatment is to cease if there has been failure to lower IGF1 after 3 months treatment.</i>				
6332G	<i>Powder for suspension for injection 30 mg (base) with diluent ampoule</i>	1	750.00	Somatuline LA	IS
	<u>Private hospital authority required</u>				
	<i>Active acromegaly in patients with persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre AND</i>				
	<i>(a) after failure of other therapy including dopamine agonists; or</i>				
	<i>(b) as interim treatment in patients awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; or</i>				
	<i>(c) where surgery and radiotherapy are contraindicated.</i>				
	<i>Treatment is to cease in patients previously treated with radiotherapy where there is biochemical evidence of remission (normal IGF1) after lanreotide acetate withdrawal for at least 4 weeks (8 weeks after the last dose). Lanreotide acetate should be withdrawn for assessment of remission every 2 years in the 10 years after radiotherapy.</i>				
	<i>Treatment is to cease if there has been failure to lower IGF1 after 3 months treatment;</i>				
	➤ <i>Patients with a histologically-confirmed diagnosis of a functional carcinoid tumour, experiencing on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persists despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, and for whom surgery or antineoplastic therapy has failed or is inappropriate.</i>				
	<i>Treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 120 mg every 28 days. Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.</i>				
6423C	<i>Injection 60 mg (base) in single dose pre-filled syringe</i>	1	1345.00	Somatuline Autogel	IS
6424D	<i>Injection 90 mg (base) in single dose pre-filled syringe</i>	1	1790.00	Somatuline Autogel	IS
6425E	<i>Injection 120 mg (base) in single dose pre-filled syringe</i>	1	2240.00	Somatuline Autogel	IS

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
LENOGRASTIM					
<u>Private hospital authority required</u>					
<i>Mobilisation of peripheral blood progenitor cells to facilitate harvest of such cells for reinfusion into patients with non-myeloid malignancies who have had myeloablative or myelosuppressive therapy;</i>					
<i>Mobilisation of peripheral blood progenitor cells, in normal volunteers, for use in allogeneic transplantation to facilitate harvest of such cells in healthy donors;</i>					
<i>Patients with non-myeloid malignancies receiving marrow-ablative chemotherapy and subsequent peripheral blood progenitor cell or bone marrow transplantation;</i>					
<i>Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in:</i>					
<i>Acute lymphoblastic leukaemia;</i>					
<i>Ewing's sarcoma;</i>					
<i>Germ cell tumours;</i>					
<i>Infants and children with CNS tumours;</i>					
<i>Neuroblastoma;</i>					
<i>Non-Hodgkin's lymphoma (intermediate or high grade);</i>					
<i>Osteosarcoma;</i>					
<i>Relapsed Hodgkin's disease;</i>					
<i>Rhabdomyosarcoma;</i>					
<i>Patients with breast cancer receiving standard dose adjuvant chemotherapy who have had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;</i>					
<i>Patients receiving first-line chemotherapy for Hodgkin's disease who have had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned.</i>					
6337M	Powder for injection 13,400,000 i.u. (105 micrograms) vial	10	512.50	Granocyte 13	MX
6338N	Powder for injection 33,600,000 i.u. (263 micrograms) vial	10	1283.60	Granocyte 34	MX
LOPINAVIR with RITONAVIR					
<u>Private hospital authority required</u>					
<i>Treatment, in combination with 2 or more other antiretroviral drugs, of HIV infection in patients with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6340Q	Capsule 133.3 mg-33.3 mg	90	322.50	Kaletra	AB
6341R	Oral liquid 400 mg-100 mg per 5 mL, 60 mL	1	129.00	Kaletra	AB

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
MYCOPHENOLATE MOFETIL					
<u>CAUTION:</u>					
<i>Careful monitoring of patients is mandatory.</i>					
<u>Private hospital authority required</u>					
<i>Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for:</i>					
<i>(a) Prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required;</i>					
<i>(b) Prophylaxis of cardiac allograft rejection. Management includes initiation, stabilisation and review of therapy as required.</i>					
6208R	Capsule 250 mg	300	555.70	CellCept	RO
6209T	Tablet 500 mg	150	555.70	CellCept	RO
6364Y	Powder for oral suspension 1 g per 5 mL, 165 mL	1	244.51	CellCept	RO
MYCOPHENOLATE SODIUM					
<u>CAUTION:</u>					
<i>Careful monitoring of patients is mandatory.</i>					
<u>Private hospital authority required</u>					
<i>Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required.</i>					
6369F	Tablet (enteric coated) 180 mg (mycophenolic acid)	120	222.28	Myfortic	NV
6370G	Tablet (enteric coated) 360 mg (mycophenolic acid)	120	444.56	Myfortic	NV
NELFINAVIR MESYLATE					
<u>Private hospital authority required</u>					
<i>Treatment of HIV infection in patients with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6331F	Tablet 250 mg (base)	300	505.56	Viracept	RO
6231Y	Oral powder 50 mg (base) per g, 144 g	1	48.53	Viracept	RO
NEVIRAPINE					
<u>Private hospital authority required</u>					
<i>Treatment of HIV infection in patients with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6215D	Tablet 200 mg	60	271.58	Viramune	BY

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
OCTREOTIDE					
<u>Private hospital authority required</u>					
<i>Active acromegaly in patients with persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre AND</i> <i>(a) after failure of other therapy including dopamine agonists; or</i> <i>(b) as interim treatment in patients awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; or</i> <i>(c) where surgery and radiotherapy are contraindicated.</i> <i>Treatment is to cease in patients previously treated with radiotherapy where there is biochemical evidence of remission (normal IGF1) after octreotide withdrawal for at least 4 weeks. Octreotide should be withdrawn for assessment of remission every 2 years in the 10 years after radiotherapy.</i> <i>Treatment is to cease if there has been failure to lower IGF1 after 3 months treatment at a dose of 100 micrograms 3 times daily;</i> <i>Patients with a histologically-confirmed diagnosis of a functional carcinoid tumour or vasoactive intestinal peptide secreting tumour (VIPoma), experiencing on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persists despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, and for whom surgery or antineoplastic therapy has failed or is inappropriate.</i> <i>Treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months' therapy. Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.</i>					
6227R	Injection 50 micrograms in 1 mL ampoule	5	41.76	Sandostatin 0.05	NV
6228T	Injection 100 micrograms in 1 mL ampoule	5	83.50	Sandostatin 0.1	NV
6229W	Injection 500 micrograms in 1 mL ampoule	5	417.96	Sandostatin 0.5	NV
OCTREOTIDE ACETATE					
<u>Private hospital authority required</u>					
<i>Patients with acromegaly who are controlled on Sandostatin subcutaneous injections.</i> <i>Treatment is to cease in patients previously treated with radiotherapy where there is biochemical evidence of remission (normal IGF1) after octreotide acetate withdrawal for at least 4 weeks (8 weeks after the last dose). Octreotide acetate should be withdrawn for assessment of remission every 2 years in the 10 years after radiotherapy.</i> <i>Treatment is to cease if there has been failure to lower IGF1 after 3 months of treatment;</i> <i>Patients with a histologically-confirmed diagnosis of a functional carcinoid tumour or vasoactive intestinal peptide secreting tumour (VIPoma) and who are controlled on Sandostatin subcutaneous injections.</i> <i>Treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with Sandostatin subcutaneous injections. Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.</i>					
6426F	Injection (modified release) 10 mg (base) vial and diluent syringe	1	1390.00	Sandostatin LAR	NV
6427G	Injection (modified release) 20 mg (base) vial and diluent syringe	1	1850.00	Sandostatin LAR	NV
6428H	Injection (modified release) 30 mg (base) vial and diluent syringe	1	2315.00	Sandostatin LAR	NV

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	PEGFILGRASTIM <u>Private hospital authority required</u> <i>For use in patients undergoing induction and consolidation therapy for acute myeloid leukaemia;</i> <i>Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in:</i> <i>Acute lymphoblastic leukaemia;</i> <i>Ewing's sarcoma;</i> <i>Germ cell tumours;</i> <i>Infants and children with CNS tumours;</i> <i>Neuroblastoma;</i> <i>Non-Hodgkin's lymphoma (intermediate or high grade);</i> <i>Osteosarcoma;</i> <i>Relapsed Hodgkin's disease;</i> <i>Rhabdomyosarcoma;</i> <i>Patients with breast cancer receiving standard dose adjuvant chemotherapy who have had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;</i> <i>Patients receiving first-line chemotherapy for Hodgkin's disease who have had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;</i> <i>Patients receiving chemotherapy for myeloma who have had a prior episode of febrile neutropenia, and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned.</i>			
6363X	Injection 6 mg in 0.6 mL single use pre-filled syringe	1	1925.00	Neulasta AN

PEGINTERFERON ALFA-2a
CAUTION:

Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Private hospital authority required

Chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa therapy and have a contraindication to ribavirin, who satisfy all of the following criteria:

- (1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);*
- (2) Abnormal serum ALT levels in conjunction with documented chronic hepatitis C infection (repeatedly anti-HCV positive and/or HCV RNA positive);*
- (3) No other forms of chronic liver disease;*
- (4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.*

The treatment course is limited to up to 48 weeks.

Patients may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
PEGINTERFERON ALFA-2a—cont.					
NOTE:					
<i>Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:</i>					
<i>(a) a nurse educator/counsellor for patients; and</i>					
<i>(b) 24 hour access by patients to medical advice; and</i>					
<i>(c) an established liver clinic; and</i>					
<i>(d) facilities for safe liver biopsy.</i>					
6439X	Injection 135 micrograms in 0.5 mL single use pre-filled syringe	4	1165.90	Pegasys	RO
6449K	Injection 180 micrograms in 0.5 mL single use pre-filled syringe	4	1350.23	Pegasys	RO
PEGINTERFERON ALFA-2b					
CAUTION:					
<i>Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.</i>					
Private hospital authority required					
<i>Chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa therapy and have a contraindication to ribavirin, who satisfy all of the following criteria:</i>					
<i>(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);</i>					
<i>(2) Abnormal serum ALT levels in conjunction with documented chronic hepatitis C infection (repeatedly anti-HCV positive and/or HCV RNA positive);</i>					
<i>(3) No other forms of chronic liver disease;</i>					
<i>(4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.</i>					
<i>The treatment course is limited to 0.5 to 1 microgram per kilogram weekly for up to 48 weeks.</i>					
<i>Patients may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop.</i>					
NOTE:					
<i>Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:</i>					
<i>(a) a nurse educator/counsellor for patients; and</i>					
<i>(b) 24 hour access by patients to medical advice; and</i>					
<i>(c) an established liver clinic; and</i>					
<i>(d) facilities for safe liver biopsy.</i>					
6411K	Powder for injection 50 micrograms with diluent in single use injection pen	4	920.00	PEG-Intron Redipen	SH
6412L	Powder for injection 80 micrograms with diluent in single use injection pen	4	1472.00	PEG-Intron Redipen	SH
6413M	Powder for injection 100 micrograms with diluent in single use injection pen	4	1840.00	PEG-Intron Redipen	SH
6414N	Powder for injection 120 micrograms with diluent in single use injection pen	4	2208.00	PEG-Intron Redipen	SH
6415P	Powder for injection 150 micrograms with diluent in single use injection pen	4	2760.00	PEG-Intron Redipen	SH

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
	RIBAVIRIN and PEGINTERFERON ALFA-2a			
	CAUTION: <i>Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.</i>			
	CAUTION: <i>Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.</i>			
	Private hospital authority required <i>Treatment of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa or peginterferon alfa treatment and who satisfy all of the following criteria:</i>			
	<i>(1) (a) Histological evidence of Metavir (or equivalent index) stage 2, 3 or 4 fibrosis or stage 1 with grade A2 or A3 inflammation, i.e. moderate to severe inflammation evident on liver biopsy; or</i>			
	<i>(b) in those patients with coagulation disorders considered severe enough to prevent liver biopsy, evidence of abnormal serum ALT levels;</i>			
	<i>(2) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);</i>			
	<i>(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and both patient and their partner are using effective forms of contraception (one for each partner). Male patients and their partners are using effective forms of contraception (one for each partner). Female partners of male patients are not pregnant.</i>			
	<i>For patients with genotype 2 or 3 hepatitis C without hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 24 weeks. For hepatitis C patients with genotype 1, 4, 5 or 6 and those genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 48 weeks.</i>			
	<i>Patients with genotype 1, 4, 5 or 6 who are eligible for 48 weeks of treatment may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop. (An HCV RNA assay at week 12 is unnecessary for genotype 2 and 3 patients because of the high likelihood of early viral response by week 12).</i>			
	<i>Patients with genotype 1, 4, 5 or 6 who are viral positive at week 12 but have attained at least a 2 log drop in viral load may only continue treatment after the first 24 weeks of treatment if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. Similarly, genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis may only continue treatment after the first 24 weeks if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. An HCV RNA qualitative assay at week 24 is unnecessary for those patients with genotype 1, 4, 5 or 6 who became viral negative at week 12.</i>			
	NOTE: <i>Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:</i>			
	<i>(a) a nurse educator/counsellor for patients; and</i>			
	<i>(b) 24 hour access by patients to medical advice; and</i>			
	<i>(c) an established liver clinic; and</i>			
	<i>(d) facilities for safe liver biopsy.</i>			
6389G	Pack containing 84 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 135 micrograms		£1 1432.08	Pegasys RBV RO

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	RIBAVIRIN and PEGINTERFERON ALFA-2a—cont.				
6390H	Pack containing 112 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 135 micrograms	‡1	1520.81	Pegasys RBV	RO
6391J	Pack containing 140 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 135 micrograms	‡1	1609.53	Pegasys RBV	RO
6392K	Pack containing 168 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 135 micrograms	‡1	1698.26	Pegasys RBV	RO
6393L	Pack containing 84 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 180 micrograms	‡1	1616.41	Pegasys RBV	RO
6394M	Pack containing 112 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 180 micrograms	‡1	1705.14	Pegasys RBV	RO
6395N	Pack containing 140 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 180 micrograms	‡1	1793.86	Pegasys RBV	RO
6396P	Pack containing 168 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 180 micrograms	‡1	1882.59	Pegasys RBV	RO

RIBAVIRIN and PEGINTERFERON ALFA-2b

CAUTION:

Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

CAUTION:

Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.

Private hospital authority required

Treatment of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa or peginterferon alfa treatment and who satisfy all of the following criteria:

- (1) (a) Histological evidence of Metavir (or equivalent index) stage 2, 3 or 4 fibrosis or stage 1 with grade A2 or A3 inflammation, i.e. moderate to severe inflammation evident on liver biopsy; or
 - (b) in those patients with coagulation disorders considered severe enough to prevent liver biopsy, evidence of abnormal serum ALT levels;
 - (2) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);
 - (3) Female patients of child-bearing age are not pregnant, not breast-feeding, and both patient and their partner are using effective forms of contraception (one for each partner). Male patients and their partners are using effective forms of contraception (one for each partner). Female partners of male patients are not pregnant.
- For patients with genotype 2 or 3 hepatitis C without hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 24 weeks. For hepatitis C patients with genotype 1, 4, 5 or 6 and those genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 48 weeks.

continued ➤

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
RIBAVIRIN and PEGINTERFERON ALFA-2b—cont.					
<i>Patients with genotype 1, 4, 5 or 6 who are eligible for 48 weeks of treatment may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop. (An HCV RNA assay at week 12 is unnecessary for genotype 2 and 3 patients because of the high likelihood of early viral response by week 12).</i> <i>Patients with genotype 1, 4, 5 or 6 who are viral positive at week 12 but have attained at least a 2 log drop in viral load may only continue treatment after the first 24 weeks of treatment if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. Similarly, genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis may only continue treatment after the first 24 weeks if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. An HCV RNA qualitative assay at week 24 is unnecessary for those patients with genotype 1, 4, 5 or 6 who became viral negative at week 12.</i>					
NOTE: <i>Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:</i> <i>(a) a nurse educator/counsellor for patients; and</i> <i>(b) 24 hour access by patients to medical advice; and</i> <i>(c) an established liver clinic; and</i> <i>(d) facilities for safe liver biopsy.</i>					
6399T	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 50 micrograms with diluent	‡1	1014.02	Pegatron	SH
6400W	Pack containing 112 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 50 micrograms with diluent	‡1	1171.51	Pegatron	SH
6401X	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 80 micrograms with diluent	‡1	1338.96	Pegatron	SH
6402Y	Pack containing 140 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 80 micrograms with diluent	‡1	1496.44	Pegatron	SH
6403B	Pack containing 168 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 80 micrograms with diluent	‡1	1496.44	Pegatron	SH
6404C	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 100 micrograms with diluent	‡1	1555.58	Pegatron	SH
6405D	Pack containing 112 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 100 micrograms with diluent	‡1	1713.07	Pegatron	SH

continued ➡

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	RIBAVIRIN and PEGINTERFERON ALFA-2b—cont.				
6406E	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 120 micrograms with diluent	‡1	1772.20	Pegatron	SH
6407F	Pack containing 140 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 120 micrograms with diluent	‡1	1929.69	Pegatron	SH
6408G	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	‡1	2097.14	Pegatron	SH
6409H	Pack containing 140 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	‡1	2254.63	Pegatron	SH
6410J	Pack containing 168 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	‡1	2254.63	Pegatron	SH
	RIFABUTIN				
	<u>Private hospital authority required</u>				
	Treatment of <i>Mycobacterium avium</i> complex infections in HIV-positive patients; Prophylaxis against <i>Mycobacterium avium</i> complex infections in HIV-positive patients with CD4 cell counts of less than 75 per cubic millimetre.				
6195C	Capsule 150 mg	30	147.00	Mycobutin	PH
	RITONAVIR				
	<u>Private hospital authority required</u>				
	Treatment of HIV infection in patients with: (a) CD4 cell counts of less than 500 per cubic millimetre; or (b) viral load of greater than 10,000 copies per mL.				
6203L	Capsule 100 mg	84	106.17	Norvir	AB
6235E	Oral solution 600 mg per 7.5 mL (80 mg per mL), 240 mL	1	242.67	Norvir	AB
	SAQUINAVIR				
	<u>Private hospital authority required</u>				
	Treatment of HIV infection in patients with: (a) CD4 cell counts of less than 500 per cubic millimetre; or (b) viral load of greater than 10,000 copies per mL.				
6248W	Soft gelatin capsule 200 mg	180	227.55	Fortovase	RO
	SAQUINAVIR MESYLATE				
	<u>Private hospital authority required</u>				
	Treatment of HIV infection in patients with: (a) CD4 cell counts of less than 500 per cubic millimetre; or (b) viral load of greater than 10,000 copies per mL.				
6199G	Capsule 200 mg (base)	270	455.00	Invirase	RO

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
SIROLIMUS					
CAUTION:					
<i>Careful monitoring of patients is mandatory.</i>					
Private hospital authority required					
<i>Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required.</i>					
6436R	Tablet 1 mg	100	670.00	Rapamune	WY
6457W	Tablet 2 mg	100	1340.00	Rapamune	WY
6437T	Oral solution 1 mg per mL, 60 mL	1	402.00	Rapamune	WY
STAVUDINE					
Private hospital authority required					
<i>Treatment of HIV infection in patients with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6186N	Capsule 20 mg	60	280.00	Zerit	BQ
6189R	Capsule 30 mg	60	333.68	Zerit	BQ
6190T	Capsule 40 mg	60	444.90	Zerit	BQ
6250Y	Powder for oral solution 1 mg per mL, 200 mL	1	55.56	Zerit	BQ
TACROLIMUS					
CAUTION:					
<i>Careful monitoring of patients is mandatory.</i>					
Private hospital authority required					
<i>Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for:</i>					
<i>(a) Prophylaxis and treatment of liver allograft rejection. Management includes initiation, stabilisation and review of therapy as required;</i>					
<i>(b) Prophylaxis and treatment of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required.</i>					
6328C	Capsule 500 micrograms	100	187.34	Prograf	JC
6216E	Capsule 1 mg	100	374.67	Prograf	JC
6217F	Capsule 5 mg	50	936.68	Prograf	JC
TENOFOVIR DISOPROXIL FUMARATE					
Private hospital authority required					
<i>Treatment of HIV infection in patients with:</i>					
<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>					
<i>(b) viral load of greater than 10,000 copies per mL.</i>					
6358P	Tablet 300 mg	30	499.00	Viread	GI
VALACICLOVIR HYDROCHLORIDE					
Private hospital authority required					
<i>Prophylaxis of cytomegalovirus (CMV) infection and disease following renal transplantation in patients at risk of CMV disease.</i>					
6280M	Tablet 500 mg (base)	100	483.63	Valtrex	GK

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	VALGANCICLOVIR HYDROCHLORIDE				
	<u>Private hospital authority required</u>				
	<i>Cytomegalovirus retinitis in patients with acquired immunodeficiency syndrome; Prophylaxis of cytomegalovirus infection and disease in solid organ transplant patients at risk of cytomegalovirus disease.</i>				
6357N	Tablet 450 mg (base)	60	2245.80	Valcyte	RO
	ZALCITABINE				
	<u>Private hospital authority required</u>				
	<i>Treatment of HIV infection in patients with:</i>				
	<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>				
	<i>(b) viral load of greater than 10,000 copies per mL.</i>				
6149P	Tablet 375 micrograms	100	193.75	Hivid	RO
6150Q	Tablet 750 micrograms	100	242.18	Hivid	RO
	ZIDOVUDINE				
	<u>Private hospital authority required</u>				
	<i>Treatment of HIV infection in patients with:</i>				
	<i>(a) CD4 cell counts of less than 500 per cubic millimetre; or</i>				
	<i>(b) viral load of greater than 10,000 copies per mL.</i>				
6153W	Capsule 100 mg	100	205.46	Retrovir	GK
6154X	Capsule 250 mg	60	308.19	Retrovir	GK
6155Y	Syrup 10 mg per mL, 200 mL bottle	1	41.09	Retrovir	GK
	ZOLEDRONIC ACID				
	<u>Private hospital authority required</u>				
	<i>Multiple myeloma;</i>				
	<i>Bone metastases from breast cancer;</i>				
	<i>Bone metastases from hormone-resistant prostate cancer, with demonstration of biochemical progression of disease despite maximal therapy with hormonal treatments;</i>				
	<i>Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.</i>				
6371H	Injection concentrate for I.V. infusion 4 mg in 5 mL	1	450.00	Zometa	NV
	<u>NOTE:</u>				
	<i>This price is based on special supply arrangements—see Pharmaceutical Benefits Pricing Authority relativity sheet for full details.</i>				

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
BOTULINUM TOXIN PROGRAM					
BOTULINUM TOXIN TYPE A PURIFIED NEUROTOXIN COMPLEX					
NOTE: <i>Arrangements to prescribe this item should be made by medical practitioners with Medicare Australia, contact telephone number 1800 819 296.</i>					
<i>Treatment of blepharospasm associated with dystonia, including benign blepharospasm and VIIth nerve disorders (hemifacial spasm) in patients 12 years and older; Treatment of dynamic equinus foot deformity due to spasticity in ambulant paediatric cerebral palsy patients 2 years of age or older; Treatment of spasmodic torticollis, either as monotherapy or as adjunctive therapy to current standard care.</i>					
6103F	<i>Lyophilised powder for I.M. injection 100 units vial</i>	1	415.50	<i>Botox</i>	<i>AG</i>
NOTE: <i>The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.</i>					
CLOSTRIDIUM BOTULINUM TYPE A TOXIN—HAEMAGGLUTININ COMPLEX					
NOTE: <i>Arrangements to prescribe this item should be made by medical practitioners with Medicare Australia, contact telephone number 1800 819 296.</i>					
<i>Treatment of dynamic equinus foot deformity due to spasticity in ambulant paediatric cerebral palsy patients 2 years of age or older; Treatment of spasmodic torticollis, either as monotherapy or as adjunctive therapy to current standard care.</i>					
6293F	<i>Lyophilised powder for I.M. injection 500 units vial</i>	1	650.00	<i>Dysport</i>	<i>IS</i>
NOTE: <i>The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.</i>					
HUMAN GROWTH HORMONE PROGRAM					
SOMATROPIN <i>(Recombinant human growth hormone)</i>					
<i>Short stature in accordance with the 'Guidelines for the Availability of Human Growth Hormone (hGH) as a Pharmaceutical Benefit'.</i>					
NOTE: <i>These guidelines may be obtained from the Department of Health and Ageing's internet site at http://www.health.gov.au/pbs/supply/hghguidelines.htm, or from:</i>					
<i>Growth Hormone Program Pharmaceutical Benefits Branch Department of Health and Ageing GPO Box 9848 CANBERRA ACT 2601 Contact telephone number (02) 6289 7274</i>					
6266T	<i>Injection 4 mg (12 i.u.) vial with 3.5 mL diluent (with preservative)</i>	1	198.00	<i>SciTropin</i>	<i>SA</i>
6295H	<i>Solution for injection 5 mg (15 i.u.) in 1.5 mL cartridge (with preservative)</i>	1	247.50	<i>Norditropin SimpleXx</i>	<i>NO</i>

continued ↻

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	SOMATROPIN (Recombinant human growth hormone)—cont.				
6330E	Injection 5 mg (15 i.u.) in 1 mL cartridge (with preservative)	1	247.50	Genotropin	PH
6465G	Solution for injection 5 mg (15 i.u.) in 1.5 mL cartridge (with preservative)	1	315.50	Norditropin Nordiflex	NO
	NOTE: This price is based on special supply arrangements—see Pharmaceutical Benefits Pricing Authority relativity sheet for full details.				
6169Q	Injection 18 i.u. (6 mg) cartridge with 3.15 mL diluent (with preservative)	1	297.00	Humatrope	LY
6329D	Injection 8 mg (24 i.u.) vial with 1.37 mL diluent cartridge (with preservative) (for use with one.click auto-injector)	1	396.00	Saizen 8 mg click.easy	SG
6296J	Solution for injection 10 mg (30 i.u.) in 1.5 mL cartridge (with preservative)	1	495.00	Norditropin SimpleXx	NO
6466H	Solution for injection 10 mg (30 i.u.) in 1.5 mL cartridge (with preservative)	1	631.00	Norditropin Nordiflex	NO
	NOTE: This price is based on special supply arrangements—see Pharmaceutical Benefits Pricing Authority relativity sheet for full details.				
6170R	Injection 36 i.u. (12 mg) cartridge with 3.15 mL diluent (with preservative)	1	594.00	Humatrope	LY
6312F	Injection 12 mg (36 i.u.) in 1 mL cartridge (with preservative)	1	594.00	Genotropin	PH
6297K	Solution for injection 15 mg (45 i.u.) in 1.5 mL cartridge (with preservative)	1	742.50	Norditropin SimpleXx	NO
6467J	Solution for injection 15 mg (45 i.u.) in 1.5 mL cartridge (with preservative)	1	946.50	Norditropin Nordiflex	NO
	NOTE: This price is based on special supply arrangements—see Pharmaceutical Benefits Pricing Authority relativity sheet for full details.				
6345Y	Injection 72 i.u. (24 mg) cartridge with 3.15 mL diluent (with preservative)	1	1188.00	Humatrope	LY
6313G	Injection 0.8 mg (2.4 i.u.) with diluent in single use syringe (without preservative)	7	277.20	Genotropin MiniQuick	PH
6314H	Injection 1 mg (3 i.u.) with diluent in single use syringe (without preservative)	7	346.50	Genotropin MiniQuick	PH
6315J	Injection 1.2 mg (3.6 i.u.) with diluent in single use syringe (without preservative)	7	415.80	Genotropin MiniQuick	PH
6316K	Injection 1.4 mg (4.2 i.u.) with diluent in single use syringe (without preservative)	7	485.10	Genotropin MiniQuick	PH
6317L	Injection 1.6 mg (4.8 i.u.) with diluent in single use syringe (without preservative)	7	554.40	Genotropin MiniQuick	PH
6318M	Injection 1.8 mg (5.4 i.u.) with diluent in single use syringe (without preservative)	7	623.70	Genotropin MiniQuick	PH
6319N	Injection 2 mg (6 i.u.) with diluent in single use syringe (without preservative)	7	693.00	Genotropin MiniQuick	PH

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
IVF/GIFT PROGRAM					
FOLLITROPIN ALFA					
<i>Patients who are receiving medical treatment as described in items 13200 or 13203 of the Medicare Benefits Schedule.</i>					
NOTE:					
<i>Arrangements to prescribe this item should be made by medical practitioners with Medicare Australia, contact telephone number 1800 700 270.</i>					
6373K	Injection set containing 1 vial powder for injection 75 i.u. and 1 pre-filled syringe solvent 1 mL	1	36.00	Gonal-f 75	SG
6374L	Injection set containing 10 vials powder for injection 75 i.u. and 10 pre-filled syringes solvent 1 mL	1	360.00	Gonal-f 75	SG
6431L	Injection 300 i.u. in 0.5 mL multi-dose cartridge	1	144.00	Gonal-f Pen	SG
6376N	Injection set containing 1 vial powder for injection 450 i.u. and 1 pre-filled syringe solvent 1 mL	1	216.00	Gonal-f	SG
6432M	Injection 450 i.u. in 0.75 mL multi-dose cartridge	1	216.00	Gonal-f Pen	SG
6433N	Injection 900 i.u. in 1.5 mL multi-dose cartridge	1	432.00	Gonal-f Pen	SG
6375M	Injection set containing 1 vial powder for injection 1,050 i.u. and 1 pre-filled syringe solvent 2 mL	1	504.00	Gonal-f	SG
FOLLITROPIN BETA					
<i>Patients who are receiving medical treatment as described in items 13200 or 13203 of the Medicare Benefits Schedule.</i>					
NOTE:					
<i>Arrangements to prescribe this item should be made by medical practitioners with Medicare Australia, contact telephone number 1800 700 270.</i>					
6274F	Solution for injection 100 i.u. in 0.5 mL, single use vial	10	480.00	Puregon 100 IU/0.5 mL	OR
6275G	Solution for injection 150 i.u. in 0.5 mL, single use vial	10	720.00	Puregon 150 IU/0.5 mL	OR
6276H	Solution for injection 200 i.u. in 0.5 mL, single use vial	10	960.00	Puregon 200 IU/0.5 mL	OR
6335K	Solution for injection 300 i.u. in 0.36 mL multi-dose cartridge	1	143.84	Puregon 300 IU/0.36 mL	OR
6336L	Solution for injection 600 i.u. in 0.72 mL multi-dose cartridge	1	287.68	Puregon 600 IU/0.72 mL	OR
6464F	Solution for injection 900 i.u. in 1.08 mL multi-dose cartridge	1	431.52	Puregon 900 IU/1.08 mL	OR

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
HUMAN CHORIONIC GONADOTROPHIN					
<i>Patients who are receiving medical treatment as described in items 13200 or 13203 of the Medicare Benefits Schedule.</i>					
NOTE:					
<i>Arrangements to prescribe this item should be made by medical practitioners with Medicare Australia, contact telephone number 1800 700 270.</i>					
6176C	Injection set containing 3 ampoules powder for injection 500 units and 3 ampoules solvent 1 mL	1	19.77	Pregnyl	OR
6178E	Injection set containing 3 ampoules powder for injection 1,500 units and 3 ampoules solvent 1 mL	1	27.59	Pregnyl	OR
6181H	Injection set containing 3 ampoules powder for injection 5,000 units and 3 ampoules solvent 1 mL	1	32.00	Pregnyl	OR
PROGESTERONE					
<i>For luteal phase support in patients who are receiving medical treatment as described in item 13200 of the Medicare Benefits Schedule. The luteal phase is defined as the time span from embryo transfer until implantation confirmed by positive B-hCG measurement.</i>					
NOTE:					
<i>Arrangements to prescribe this item should be made by medical practitioners with Medicare Australia, contact telephone number 1800 700 270.</i>					
6366C	Vaginal gel (prolonged release) 90 mg in single dose pre-filled applicator	15	148.50	Crinone 8%	SG
NOTE:					
<i>This price is based on special supply arrangements—see Pharmaceutical Benefits Pricing Authority relativity sheet for full details. Relativity sheets are available on the Department of Health and Ageing's internet site (visit www.health.gov.au and search for 'relativity sheets').</i>					
OPIATE DEPENDENCE TREATMENT PROGRAM					
The Australian Government funds the cost of buprenorphine hydrochloride and methadone hydrochloride supplied as a pharmaceutical benefit through clinics and pharmacies approved by State and Territory governments. For further information about this program, please contact your State or Territory Government.					
BUPRENORPHINE HYDROCHLORIDE					
<i>Treatment of opiate dependence, including maintenance and detoxification (withdrawal), within a framework of medical, social and psychological treatment.</i>					
NOTE:					
<i>Treatment must be in accordance with the law of the relevant State or Territory.</i>					
6307Y	Sublingual tablet 400 micrograms (base)	7	6.16	Subutex	RC
6308B	Sublingual tablet 2 mg (base)	7	10.50	Subutex	RC
6309C	Sublingual tablet 8 mg (base)	7	30.10	Subutex	RC

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer
METHADONE HYDROCHLORIDE				
CAUTION: <i>The risk of drug dependence is high.</i>				
<i>Treatment of opiate dependence in accordance with the law of the relevant State or Territory.</i>				
6171T	Oral liquid 25 mg per 5 mL, 200 mL	1	7.40	^a Biodone Forte MW ^a GK
6172W	Oral liquid 25 mg per 5 mL, 1 L	1	36.00	^a Biodone Forte MW ^a GK
6174Y	Powder 1 g (for preparation of other dosage forms)	1	3.50	GK

SPECIAL AUTHORITY PROGRAM

IMATINIB MESYLATE

NOTE:

Any queries concerning the arrangements to prescribe imatinib mesylate may be directed to Medicare Australia on 1800 242 679.

Written applications for authority to prescribe imatinib mesylate should be forwarded to:

*Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001*

NOTE:

Imatinib mesylate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Section 100 authority required

Initial PBS-subsidised treatment, for up to 3 months, of adult patients with a metastatic or unresectable malignant gastrointestinal stromal tumour which has been histologically confirmed by the detection of CD117 on immunohistochemical staining.

Patients who have not previously been treated with imatinib mesylate for a metastatic or unresectable malignant gastrointestinal stromal tumour must commence treatment at a dose not exceeding 400 mg per day for at least 3 months. Authority prescriptions for a higher dose will not be approved during this initial 3 month treatment period.

Patients who have previously been treated with non-PBS-subsidised imatinib mesylate for a metastatic or unresectable malignant gastrointestinal stromal tumour are eligible to receive up to 3 months treatment at a dose of up to 600 mg per day.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and*
- (2) a completed Imatinib Mesylate (Glivec) PBS Authority Application for Use in the Treatment of Gastrointestinal Stromal Tumour - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:*
 - (i) a copy of a pathology report from an Approved Pathology Authority supporting the diagnosis of a gastrointestinal stromal tumour and confirming the presence of CD117 on immunohistochemical staining; and*

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	IMATINIB MESYLATE—cont.				
	(ii) a copy of the most recent (within 2 months of the application) computed tomography (CT) scan, magnetic resonance imaging (MRI) or ultrasound assessment of the tumour(s), including whether or not there is evidence of metastatic disease; and (iii) where the application for authority to prescribe is being sought on the basis of an unresectable tumour, written evidence in support of that claim must be provided; and (iv) for patients who commenced treatment with imatinib mesylate for a metastatic or unresectable malignant gastrointestinal stromal tumour prior to 1 December 2004, the date on which therapy with imatinib mesylate was commenced.				
	Section 100 authority required				
	Continuing PBS-subsidised treatment, at a dose of up to 600 mg per day, of adult patients with a metastatic or unresectable malignant gastrointestinal stromal tumour who have previously been issued with an authority prescription for this drug.				
	Applications for continuing treatment may be made by telephone (1800 242 679, hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).				
	NOTE:				
	Patients who achieve a response to treatment at an imatinib dose of 400 mg per day should be continued at this dose and assessed for response at regular intervals. Patients who fail to achieve a response to 400 mg per day may have their dose increased to 600 mg per day. Authority applications for doses higher than 600 mg per day will not be approved.				
	A response to treatment is defined as a decrease from baseline in the sum of the products of the perpendicular diameters of all measurable lesions of 50% or greater. (Response definition based on the Southwest Oncology Group standard criteria, see Demetri et al. <i>N Engl J Med</i> 2002; 347: 472-80.)				
6440Y	Tablet 100 mg (base)	60	1886.17	Glivec	NV
6441B	Tablet 400 mg (base)	30	3772.33	Glivec	NV

NOTE:

No applications for increased maximum quantities will be authorised. Up to 2 repeats may be authorised for item 6440Y or 6441B.

Section 100 authority required

- Initial treatment of patients in the chronic phase of chronic myeloid leukaemia expressing the Philadelphia chromosome or the transcript, bcr-abl tyrosine kinase, and who have a primary diagnosis of chronic myeloid leukaemia.

Applications under this restriction will be limited to provide patients with a maximum of 18 months of therapy from the date the first application for initial treatment was approved.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Imatinib Mesylate (Glivec) PBS Authority Application for Use in the Treatment of Chronic Myeloid Leukaemia - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of chronic myeloid leukaemia to confirm eligibility for treatment, or a qualitative PCR report documenting the presence of the bcr-abl transcript in either peripheral blood or bone marrow; and

Code	Item and Section 100 Restriction	Price ex		Proprietary Name and Manufacturer
		Pack Size	Manufacturer \$	

IMATINIB MESYLATE—cont.

(4) a copy of a signed patient acknowledgement form indicating that the patient understands and acknowledges that PBS-subsidised treatment with imatinib mesylate for the chronic phase of chronic myeloid leukaemia will cease if subsequent testing demonstrates that:

- (i) the patient has failed to achieve a major cytogenetic response within the initial 18 months of treatment [see Note defining major cytogenetic response]; or*
- (ii) the patient has failed to sustain a major cytogenetic response for 12 months from the date of the last pathology report that indicated that a major cytogenetic response had been achieved [see Note defining major cytogenetic response].*

NOTE:

Imatinib mesylate in the chronic phase of chronic myeloid leukaemia will only be subsidised for patients who are not receiving concomitant PBS-subsidised interferon alfa therapy.

Patients should be commenced on a dose of imatinib mesylate of 400 mg (base) daily and maintained on a minimum dose of imatinib mesylate of 400 mg (base) daily. Prescribing of lower doses should be carefully considered. Continuing therapy is dependent on patients demonstrating a response to imatinib mesylate therapy following the initial 18 months of treatment and at 12 monthly intervals thereafter, irrespective of the daily imatinib mesylate dose received.

Section 100 authority required

Continuing treatment of patients who have received initial treatment with imatinib mesylate as a pharmaceutical benefit for the chronic phase of chronic myeloid leukaemia and who have demonstrated a major cytogenetic response in the preceding 12 months.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and*
- (2) demonstration of continued response to treatment as evidenced by major cytogenetic response [see Note explaining requirements]. Where this has been supplied within the previous 12 months, only the date of the relevant pathology report need be provided.*

NOTE:

Definition of response.

A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells.

Authority approval requirements.

For the purposes of assessing response to PBS-subsidised treatment with imatinib mesylate, cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping must be submitted. Where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with bcr-abl specific probe or quantitative PCR must be submitted. The cytogenetic analyses must be submitted as follows:

- (i) between 10 and 12 months of the commencement of treatment with imatinib mesylate, at which time patients in whom a major cytogenetic response has been demonstrated may receive authorisation for a further 12 months of treatment; and*
- (ii) within 18 months of the commencement of treatment with imatinib mesylate, in patients who have failed to demonstrate a major cytogenetic response at between 10 and 12 months (patients in whom a major cytogenetic response is demonstrable by 18 months may also receive authorisation for a further 12 months of treatment); and*
- (iii) at no greater than 12 month intervals thereafter, to demonstrate that the major cytogenetic response has been sustained.*

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
IMATINIB MESYLATE—cont.					
<i>For each authority application where eligibility for continuing PBS-subsidised treatment is to be demonstrated, a copy of the cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping must be submitted as described in (i) to (iii) above, unless, at the time of the authority application, the standard karyotyping is not informative. Where the standard karyotyping conducted at the time of application is not informative, a copy of a cytogenetic analysis conducted on the bone marrow using either FISH with bcr-abl specific probe or quantitative PCR must be submitted with the authority application. For patients in whom cytogenetic status is determined on the basis of bcr-abl transcript level measured by quantitative PCR performed on the bone marrow, continuation of treatment will be authorised irrespective of changes to bcr-abl transcript levels. A copy of the non-informative standard karyotype analysis must be included with the authority application.</i> <i>For patients in whom failure to obtain a bone marrow sample (dry tap) has occurred on 2 separate consecutive occasions at any point described in (i) to (iii) above and at no greater than 2 months between each attempt, results of a quantitative PCR analysis of peripheral blood providing bcr-abl transcript levels will be accepted as the basis for continuing therapy thereafter. The first application for continuing treatment for patients in whom there has been a failure to obtain bone marrow samples on 2 separate consecutive occasions, must include 2 pathology reports demonstrating failure to obtain an adequate bone marrow sample. In cases where pathology reports are not obtainable, a statement from the treating specialist that no bone marrow sample could be obtained from 2 separate attempts will be accepted. A pathology report from an Approved Pathology Authority providing the results of the quantitative PCR analysis of peripheral blood must also be submitted and must clearly state the level of bcr-abl transcript which reliably corresponds to less than 35% Philadelphia positive bone marrow cells.</i> <i>Where a patient has previously received PBS-subsidised treatment with imatinib mesylate, no approval will be granted for PBS-subsidised re-treatment in the chronic phase of chronic myeloid leukaemia, where that patient has at any time failed to meet the criteria for continuing treatment.</i>					
6444E	Tablet 100 mg (base)	60	1886.17	Glivec	NV
6445F	Tablet 400 mg (base)	30	3772.33	Glivec	NV

NOTE:

Up to 5 repeats may be authorised for item 6444E or 6445F.

Code	Item and Section 100 Restriction	Pack Size	Price ex Manufacturer \$	Proprietary Name and Manufacturer	
	IMATINIB MESYLATE—cont. Section 100 authority required <i>Treatment of patients in the accelerated phase of chronic myeloid leukaemia expressing the Philadelphia chromosome or the transcript, bcr-abl tyrosine kinase, and who have a primary diagnosis of chronic myeloid leukaemia. Progress to the accelerated phase is defined by the presence of 1 or more of the following:</i> <i>(1) Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 15% but less than 30%; or</i> <i>(2) Percentage of blasts plus promyelocytes in the peripheral blood or bone marrow greater than or equal to 30%; or</i> <i>(3) Peripheral basophils greater than or equal to 20%; or</i> <i>(4) Progressive splenomegaly to a size greater than or equal to 10 cm below the left costal margin to be confirmed on 2 occasions at least 4 weeks apart, or a greater than or equal to 50% increase in size below the left costal margin over 4 weeks; or</i> <i>(5) Karyotypic evolution (chromosomal abnormalities in addition to a single Philadelphia chromosome).</i> <i>Applications for authorisation must be in writing and must include:</i> <i>(a) a completed authority prescription form; and</i> <i>(b) a completed Imatinib Mesylate (Glivec) PBS Authority Application for Use in the Treatment of Chronic Myeloid Leukaemia - Supporting Information form, stating which of the above criteria are satisfied by the patient; and</i> <i>(c) a copy of the confirming pathology report from an Approved Pathology Authority in the case of criteria (1), (2), (3) and (5) above, or details of the dates of assessments in the case of progressive splenomegaly.</i> Section 100 authority required <i>Treatment of patients in the blast phase of chronic myeloid leukaemia expressing the Philadelphia chromosome or the transcript, bcr-abl tyrosine kinase, and who have a primary diagnosis of chronic myeloid leukaemia. Progress to myeloid blast crisis is defined as either:</i> <i>(1) Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 30%; or</i> <i>(2) Extramedullary involvement other than spleen and liver.</i> <i>Applications for authorisation must be in writing and must include:</i> <i>(a) a completed authority prescription form; and</i> <i>(b) a completed Imatinib Mesylate (Glivec) PBS Authority Application for Use in the Treatment of Chronic Myeloid Leukaemia - Supporting Information form, stating which of the above criteria are satisfied by the patient; and</i> <i>(c) a copy of the confirming pathology report from an Approved Pathology Authority in the case of criterion (1) above, or details of the date of assessment in the case of extramedullary involvement.</i> Section 100 authority required <i>Continuing treatment of patients with chronic myeloid leukaemia expressing the Philadelphia chromosome or the transcript, bcr-abl tyrosine kinase, where the patient has previously received PBS-subsidised treatment with imatinib mesylate of:</i> <i>(i) the accelerated phase of chronic myeloid leukaemia; or</i> <i>(ii) the blast phase of chronic myeloid leukaemia.</i>				
6446G	Tablet 100 mg (base)	60	1886.17	Glivec	NV
6447H	Tablet 400 mg (base)	30	3772.33	Glivec	NV
	NOTE: <i>Up to 2 repeats may be authorised for item 6446G or 6447H.</i>				

Section 3

Container Prices, Fees, Standard Packs and Prices for Ready Prepared Pharmaceutical Benefits

CONTAINER PRICES FOR QUANTITIES OF READY PREPARED BENEFITS LESS THAN THE STANDARD PACK:

Injectables	150 mL vial	\$0.67
Other Items	25 mL vial	\$0.24

(The 25 mL is the most commonly used size)

FEES:

Dispensing Fee for Ready Prepared Benefits	\$4.75
Dangerous Drug Fee	\$2.61
Additional Fee for Agreed Price Ready Prepared Benefits	\$0.97

NOTE—

Standard packs and prices (including mark-up, but without dispensing fee and dangerous drug fee) are for items against the price of which an asterisk () is shown in Section 2 of the Schedule.*

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
EMERGENCY DRUG (DOCTOR'S BAG) SUPPLIES				
3486L	Benzylpenicillin	600 mg	5 @ 8.75	CS
3463G	Diphtheria and Tetanus Vaccine, Adsorbed, Diluted For Adult Use	0.5 mL	1 @ 5.73	CS
3470P	Hydrocortisone Sodium Succinate	100 mg with 2 mL solvent	1 @ 5.40	PH
3481F	Naloxone Hydrochloride	800 mcg in 2 mL	1 @ 18.39	CS
3482G		2 mg in 5 mL	1 @ 29.40	CS
3488N	Promethazine Hydrochloride	50 mg in 2 mL	5 @ 6.77	MX
3492T	Tetanus Vaccine, Adsorbed	0.5 mL	1 @ 4.08	CS
SPECIAL PHARMACEUTICAL BENEFITS				
2315W	Bleomycin Sulfate	15,000 i.u.	1 @ 51.22 (for reimbursement price)	MX
			1 @ 97.30 (for total dispensed price)	MX
8809P	Pemetrexed Disodium	500 mg (base)	1 @ 1612.68 (for reimbursement price)	LY
			1 @ 1843.06 (for total dispensed price)	LY
9713F		500 mg (base)	1 @ 1843.06 (for reimbursement price)	LY
			1 @ 1843.06 (for total dispensed price)	LY
GENERAL PHARMACEUTICAL BENEFITS				
8048N	Abciximab	10 mg in 5 mL	1 @ 508.87	LY
8747J	Acetylcysteine	200 mg per mL, 5 mL	6 @ 29.04	BQ
1003T	Aciclovir	200 mg	25 @ 49.69	AF,DP,HX
			25 @ 52.58	GK
2600W	Allopurinol	100 mg	100 @ 4.46	AF
1029E	Alteplase (Recombinant tissue-type plasminogen activator)	50 mg set	1 @ 1076.39	BY
2576N	Aluminium Hydroxide with Magnesium Hydroxide	200 mg-200 mg	100 @ 3.97	WW
			100 @ 4.17	PC
2157M		200 mg-200 mg per 5 mL, 500 mL	1 @ 3.97	WW
			1 @ 4.13	PC
2159P	Aluminium Hydroxide with Magnesium Trisilicate and Magnesium Hydroxide	250 mg-120 mg-120 mg per 5 mL, 500 mL	1 @ 3.97	FM
3109P	Amiloride Hydrochloride	5 mg	50 @ 1.82	AF
3079C	Amino Acid Formula without Methionine, Threonine and Valine and low in Isoleucine	200 g	1 @ 215.38	SB
8554F	Amino Acid Formula without Phenylalanine	500 mg, 200	1 @ 83.14	SB
8678R		1 g, 75	1 @ 62.40	SB
8706F		42 g, 20	1 @ 165.80	SB
2347M		20 g, 30	1 @ 219.65	SB
3072Q		250 g	1 @ 157.04	SB
2379F	Amino Acid Formula without Phenylalanine, Tyrosine and Methionine	500 g	1 @ 366.08	SB

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Approved Name	Form/Strength	Pack and Price \$		Manufacturer
8479G	Amino Acid Formula with Vitamins, Minerals and Long Chain Polyunsaturated Fatty Acids without Phenylalanine	400 g	1 @	90.06	SB
8417B	Amino Acid Formula with Vitamins and Minerals without Methionine	400 g	1 @	89.22	SB
8677Q		20 g, 30	1 @	356.43	VF
8744F		25 g, 30	1 @	477.36	VF
8328H		500 g	1 @	176.80	SB
8416Y		500 g	1 @	207.58	SB
8058D	Amino Acid Formula with Vitamins and Minerals without Methionine, Threonine and Valine and low in Isoleucine	400 g	1 @	80.48	SB
8059E		500 g	1 @	203.28	SB
8061G		500 g	1 @	257.92	SB
8751N	Amino Acid Formula with Vitamins and Minerals without Phenylalanine	19 g, 30	1 @	227.40	SB
8555G		20 g, 30	1 @	228.53	VF
8591E		25 g, 30	1 @	408.32	VF
8804J		27.8 g, 30	1 @	544.55	SB
8613H		29 g, 30	1 @	228.81	SB
8727H		50 g, 30	1 @	530.71	SB
8466N		350 g	1 @	63.44	AB
8544Q		400 g	1 @	72.50	AB
2737C		400 g	1 @	62.84	SB
8467P		325 g	1 @	88.40	AB
8545R		400 g	1 @	108.78	AB
2738D		500 g	1 @	113.36	SB
2739E		500 g	1 @	176.84	SB
8746H		250 mL	18 @	274.25	SB
8846N		130 mL, 30	1 @	408.10	VF
8631G	Amino Acid Formula with Vitamins and Minerals without Phenylalanine and Tyrosine	20 g, 30	1 @	429.81	VF
8667E		25 g, 30	1 @	640.80	VF
8445L		400 g	1 @	98.80	SB
8446M		500 g	1 @	213.20	SB
3078B		500 g	1 @	277.68	SB
8592F	Amino Acid Formula with Vitamins and Minerals without Valine, Leucine and Isoleucine	20 g, 30	1 @	228.53	VF
8632H		25 g, 30	1 @	408.32	VF
8745G		29 g, 30	1 @	228.81	SB
8468Q		350 g	1 @	63.44	AB
8546T		400 g	1 @	72.50	AB
2380G		400 g	1 @	62.84	SB
8469R		325 g	1 @	88.40	AB
8547W		400 g	1 @	108.78	AB
8310J		500 g	1 @	353.60	SB
8260R		500 g	1 @	113.36	SB
8057C		500 g	1 @	176.84	SB
8574G	Amino Acids—Synthetic, Formula	400 g	1 @	45.75	AB
8443J		400 g	1 @	45.75	SB
8754R		400 g	1 @	45.75	SB
8575H		400 g	1 @	45.75	AB
3066J		400 g	1 @	45.75	SB

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
8755T	Amino Acids—Synthetic, Formula	400 g	1 @ 45.75	SB
8736T	Amisulpride	100 mg per mL, 60 mL	1 @ 69.15	SW
1140B	BCG Immunotherapeutic (Bacillus Calmette-Guérin/ Connaught strain)	6.6 to 19.2 x 10 ⁸ CFU set	1 @ 156.00	AV
8743E	Benzathine Penicillin	900 mg in 2 mL	1 @ 20.20	AS
1775K	Benzylpenicillin	600 mg	5 @ 8.75	CS
2647H		3 g	5 @ 22.88	CS
2812B	Betamethasone Valerate	200 mcg (base) per g, 100 g	1 @ 4.28	EX,FM,SH
			1 @ 5.58	SI
2820K		200 mcg (base) per g, 100 g	1 @ 4.28	EX,SH
2544X	Biperiden Hydrochloride	2 mg	100 @ 7.15	AB
1260H	Bisacodyl	10 mg, 10	1 @ 5.59	BY,PP
1258F		10 mg, 12	1 @ 4.25	FL,PP
3116B	Calcium	500 mg	60 @ 4.37	MM
8740B	Calcium Folate	equiv. to 50 mg folic acid in 5 mL	1 @ 35.75	MX
			10 @ 279.12	PF
1153Q	Carbimazole	5 mg	100 @ 12.16	RO
8576J	Carbohydrate, Fat, Vitamins, Minerals and Trace Elements	400 g	1 @ 32.75	AB
8369L		400 g	1 @ 36.75	SB
8514D	Carbomer 974	3 mg per g, 0.5 g, 30	1 @ 9.77	AQ
8578L	Carbomer 980	2 mg per g, 0.6 mL, 30	1 @ 9.77	NV
1160C	Carboplatin	50 mg in 5 mL	1 @ 37.29	MX,PU
1161D		150 mg in 15 mL	1 @ 85.12	MX,PU
1162E		450 mg in 45 mL	1 @ 163.22	MX,PU
8823J	Carmellose Sodium	2.5 mg per mL, 0.6 mL, 24	1 @ 7.81	CX
2338C		5 mg per mL, 0.4 mL, 30	1 @ 9.77	AG
2324H		10 mg per mL, 0.4 mL, 30	1 @ 9.77	AG
8824K		10 mg per mL, 0.6 mL, 28	1 @ 9.12	CX
8315P	Cefepime	1 g	1 @ 18.34	BQ
8316Q		2 g	1 @ 33.54	BQ
1085D	Cefotaxime	1 g	1 @ 5.16	SZ
1086E		2 g	1 @ 9.54	SZ
1790F	Ceftriaxone	250 mg	1 @ 6.82	RO
1783W		500 mg	1 @ 9.87	RO
1784X		1 g	1 @ 15.49	RO,SZ
1785Y		2 g	1 @ 28.67	MX,RO,SZ
1256D	Cephazolin	500 mg	5 @ 20.08	MX
1257E		1 g	5 @ 31.69	MX,SZ
1163F	Chlorambucil	2 mg	25 @ 19.42	GK
1585K	Chlorthalidone	25 mg	50 @ 3.56	NV
2967E	Cholestyramine	4.7 g (equiv. to 4 g cholestyramine)	1 @ 26.38	BQ
1217C	Ciprofloxacin	3 mg per mL, 5 mL	1 @ 11.58	IQ
			1 @ 12.82	AQ
8800E	Cladribine	10 mg in 5 mL	1 @ 731.12	OA
1211R	Clomiphene Citrate	50 mg	5 @ 17.58	AW,GX,HX
1805B	Clonazepam	500 mcg	100 @ 9.08	AF
			100 @ 11.45	RO
1806C		2 mg	100 @ 17.07	AF
			100 @ 19.75	RO
1808E		2.5 mg per mL, 10 mL	1 @ 4.27	RO
8785J	Codeine Phosphate with Paracetamol	30 mg-500 mg	20 @ 2.19	AL,CO,DK, FM,GK
			20 @ 4.08	SW
1228P	Copper Sulfate	Tablets, 36	1 @ 33.62	BN

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Approved Name	Form/Strength	Pack and Price \$	Manufacturer
1079T	Cyclophosphamide	500 mg	1 @ 11.54	BX
8657P	Cyclosporin	10 mg	60 @ 45.46	NV
8658Q		25 mg	30 @ 49.69	MX
			30 @ 50.88	HX
			30 @ 52.09	NV
8659R		50 mg	30 @ 104.04	MX
			30 @ 105.23	HX
			30 @ 106.42	NV
8660T		100 mg	30 @ 203.87	MX
			30 @ 205.07	HX
			30 @ 206.27	NV
8661W		100 mg per mL, 50 mL	1 @ 364.92	NV
1798P	Cyproheptadine Hydrochloride	4 mg	50 @ 3.54	FR
1270W	Cyproterone Acetate	50 mg	50 @ 135.28	AF,DP,GX, HX,SY
			50 @ 136.92	SC
2884T	Cytarabine	100 mg in 5 mL	5 @ 27.54	PU
8034W		500 mg in 5 mL	1 @ 31.78	MX
8641T	Dalteparin Sodium (Low Molecular Weight Heparin Sodium—porcine mucous)	2,500 units (anti-Xa) in 0.2 mL	10 @ 50.80	PH
8642W		5,000 units (anti-Xa) in 0.2 mL	10 @ 52.93	PH
8643X		7,500 units (anti-Xa) in 0.75 mL	10 @ 79.81	PH
8662X	Desmopressin Acetate	200 mcg	30 @ 49.93	FP
2129C		100 mcg per mL, 2.5 mL	1 @ 31.99	FP
8711L		10 mcg per actuation, 60 actuations, 6 mL	1 @ 79.88	FP
1299J	Diclofenac Sodium	25 mg (e.c.)	50 @ 4.38	AW,BG,CH, DP,GX,HX, TW
			50 @ 6.03	NV
1302M		100 mg	20 @ 9.14	NV
1319K	Diflunisal	250 mg	50 @ 5.27	MK
3164M	Digoxin	50 mcg per mL, 60 mL	1 @ 11.00	SI
1341N	Diphtheria and Tetanus Vaccine, Adsorbed	0.5 mL	1 @ 4.63	CS
8783G	Diphtheria and Tetanus Vaccine, Adsorbed, Diluted For Adult Use	0.5 mL	1 @ 5.73	CS
8461H	Disodium Pamidronate	15 mg in 5 mL	1 @ 67.71	MX
8462J		30 mg in 10 mL	1 @ 135.43	MX
8071T	Docetaxel	20 mg (anhydrous) set	1 @ 338.27	AV
1125F	Docusate Sodium with Bisacodyl	100 mg-10 mg, 5	1 @ 2.79	FM
1336H	Doxorubicin Hydrochloride	10 mg in 5 mL	1 @ 35.97	IT,MX,PH
1340M		20 mg in 10 mL	1 @ 62.98	PH
1342P		50 mg in 25 mL	1 @ 144.25	IT,MX,PH
2702F	Doxycycline	100 mg	7 @ 2.68	AF,CH,DP, GX,HX,SI, TW
			7 @ 4.28	PF
2703G		100 mg	7 @ 2.68	FA
			7 @ 4.20	MX
2714W		100 mg	7 @ 2.68	AF,CH,DP, HX,SI,TW
			7 @ 4.28	PF
3199J	Electrolyte Replacement Solution	1 L	1 @ 7.69	BX

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
8716R	Enoxaparin Sodium	20 mg (2,000 i.u. anti-Xa) in 0.2 mL	10 @ 50.80	AV
8639Q		40 mg (4,000 i.u. anti-Xa) in 0.4 mL	10 @ 52.93	AV
8640R		60 mg (6,000 i.u. anti-Xa) in 0.6 mL	10 @ 75.70	AV
8367J	Entacapone	200 mg	100 @ 142.00	NV
1375J	Epirubicin Hydrochloride	10 mg in 5 mL	1 @ 56.00	IT,PH
1376K		20 mg in 10 mL	1 @ 103.90	PH
1377L		50 mg in 25 mL	1 @ 255.63	IT,PH
8817C		100 mg in 50 mL	1 @ 504.40	IT
8397Y	Eprosartan Mesylate	400 mg (base)	28 @ 11.78	SM
8683B	Eptifibatide Acetate	20 mg (base) in 10 mL	1 @ 132.03	SH
8684C		75 mg (base) in 100 mL	1 @ 349.26	SH
1397M	Erythromycin Lactobionate	1 g (base)	1 @ 9.20	AB
8001D	Essential Amino Acids Formula with Minerals and Vitamin C	200 g	1 @ 64.88	SB
8778B	Etanercept	25 mg and 1 mL solvent, 4	1 @ 941.78	WY
8779C		25 mg and 1 mL solvent, 4	1 @ 941.78	WY
8637N		25 mg and 1 mL solvent, 4	1 @ 941.78	WY
8638P		25 mg and 1 mL solvent, 4	1 @ 941.78	WY
8748K	Ethacrynic Acid	25 mg	100 @ 26.49	MK
1390E	Etoposide	100 mg in 5 mL	1 @ 35.19	IT,MX
8120J	Etoposide Phosphate	113.6 mg (equiv. to 100 mg etoposide)	1 @ 35.19	BQ
8842J	Everolimus	0.75 mg	60 @ 833.04	NV
1473M	Fluconazole	100 mg in 50 mL	1 @ 27.40	PF
1474N		200 mg in 100 mL	1 @ 49.45	PF
1433K	Fludrocortisone Acetate	100 mcg	100 @ 6.04	BQ
2528C	Fluorouracil	500 mg in 10 mL	1 @ 6.17	MX
2958Q	Folic Acid	500 mcg	100 @ 3.38	AF
1437P		5 mg	100 @ 3.55	AF
8672K	Follitropin Alfa	75 i.u.	1 @ 49.72	SG
8713N		300 i.u.	1 @ 191.87	SG
8675N		450 i.u.	1 @ 287.79	SG
8714P		450 i.u.	1 @ 287.80	SG
8715Q		900 i.u.	1 @ 575.61	SG
8565T	Follitropin Beta	300 i.u. in 0.36 mL	1 @ 191.60	OR
8566W		600 i.u. in 0.72 mL	1 @ 383.20	OR
8871X		900 i.u. in 1.08 mL	1 @ 574.80	OR
2414C	Frusemide	20 mg	50 @ 1.72	DP,FM
			50 @ 2.28	AV
8444K	Gelatin - Succinylated	20 g per 500 mL, 500 mL	1 @ 13.20	BR
8049P	Gemcitabine Hydrochloride	200 mg (base)	1 @ 59.14	LY
8050Q		1 g (base)	1 @ 280.63	LY
2824P	Gentamicin Sulfate	80 mg (base) in 2 mL	5 @ 7.08	MX
2245E	Glucose	278 mmol per L, 1 L	1 @ 3.85	BX
2891E	Glucose Indicator—Blood	Electrode strips, 50	1 @ 24.01	RD
8634K		Electrode strips, 50	1 @ 27.84	BN
8749L		Electrode strips, 50	1 @ 24.01	DR
8766J		Electrode strips, 50	1 @ 24.01	DR
8557J		Electrode strips, 50	1 @ 24.01	GD
8723D		Electrode strips, 50	1 @ 24.01	BR
8825L		Electrode strips, 50	1 @ 24.01	DR
8176H		Discs containing electrode sensors, 10 sensors per disc, 5	1 @ 24.01	BN
8190C		Reagent strips, 50	1 @ 24.01	RD

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Approved Name	Form/Strength	Pack and Price \$	Manufacturer
8739Y	Glucose Indicator—Blood	Reagent strips, 50	1 @ 24.01	RD
8806L		Reagent strips, 51	1 @ 24.01	RD
2890D		Reagent strips, 50	1 @ 24.01	NA
8053W		Reagent strips, 50	1 @ 24.01	NA
8759B		Reagent strips, 50	1 @ 24.01	LB
2914J		Reagent strips, 50	1 @ 19.89	TC
2917M	Glucose Indicator—Urine	Reagent strips, 50	1 @ 19.89	BN
8795X		Reagent strips, 50	1 @ 24.01	PX
2352T		Reagent strips, 50	1 @ 6.62	BN
3104J		Reagent strips, 50	1 @ 6.14	BN
3106L	Glucose and Ketone Indicator—Urine	Reagent strips, 50	1 @ 5.38	RD
3107M		Reagent strips, 50	1 @ 5.43	BN
2555L	Glycerol	700 mg, 12	1 @ 3.34	PP
2556M		1.4 g, 12	1 @ 3.47	PP
2557N		2.8 g, 12	1 @ 3.59	PP
8728J	Granisetron Hydrochloride	2 mg (base)	1 @ 31.04	MX
1076P	Heparin Sodium	35,000 units in 35 mL	1 @ 11.98	MX
1583H	Human Chorionic Gonadotrophin	500 units set	1 @ 24.17	OR
1640H	Hydralazine Hydrochloride	25 mg	100 @ 4.49	AF
1639G	Hydrochlorothiazide with Amiloride Hydrochloride	50 mg	100 @ 5.43	AF
1486F		50 mg-5 mg	50 @ 3.20	AF
			50 @ 4.92	MK
1502C	Hydrocortisone Acetate	21.1 g	1 @ 15.68	GC
1510L	Hydrocortisone Sodium Succinate	100 mg with 2 mL solvent	1 @ 5.40	PH
1511M		250 mg with 2 mL solvent	1 @ 7.52	PH
1501B		100 mg with 2 mL solvent	1 @ 5.40	PH
8299T	Hypromellose with Dextran	3 mg-1 mg per mL, 0.4 mL, 28	1 @ 9.12	AQ
3198H	Ibuprofen	200 mg	50 @ 2.18	AF
2446R	Idarubicin Hydrochloride	5 mg	1 @ 80.52	PH
2448W		10 mg	1 @ 151.07	PH
2452C		5 mg	1 @ 193.75	PH
2453D	Ifosfamide	10 mg	1 @ 377.18	PH
8076C		1 g	1 @ 52.61	BX
8077D		2 g	1 @ 97.30	BX
2454E	Indomethacin	25 mg	50 @ 1.99	AF
			50 @ 3.52	MK
2757D	Insulin Aspart	100 mg	20 @ 7.94	MK
8571D		100 units per mL, 10 mL	1 @ 31.59	NO
8435Y		100 units per mL, 3 mL, 5	1 @ 53.17	NF,NO
8609D	Insulin Aspart—Insulin Aspart Protamine Suspension	100 units (30 units-70 units) per mL, 3 mL, 5	1 @ 53.17	NF,NO
1711C		100 units per mL, 10 mL	1 @ 26.32	AS
1533Q	Insulin Isophane (N.P.H.)	100 units per mL, 10 mL	1 @ 26.32	LY,NO
1761Q		100 units per mL, 3 mL, 5	1 @ 44.91	LY,NI,NL,NO
8084L		100 units per mL, 10 mL	1 @ 31.59	LY
8212F	Insulin Lispro	100 units per mL, 3 mL, 5	1 @ 53.17	LY
8390N		100 units (25 units-75 units) per mL, 3 mL, 5	1 @ 53.17	LY
1713E	Insulin Lispro—Insulin Lispro Protamine Suspension	100 units (25 units-75 units) per mL, 3 mL, 5	1 @ 53.17	LY
1713E		100 units per mL, 10 mL	1 @ 26.32	AS
1531N		100 units per mL, 10 mL	1 @ 26.32	LY,NO
1762R	Insulin Neutral	100 units per mL, 3 mL, 5	1 @ 44.91	LY,NO

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
1426C	Insulin Neutral—Insulin Isophane (N.P.H.), (Mixed) (Biphasic Isophane)	100 units (30 units- 70 units) per mL, 10 mL	1 @ 26.32	LY,NO
1425B		100 units (50 units- 50 units) per mL, 10 mL	1 @ 26.32	NO
8006J		100 units (20 units- 80 units) per mL, 3 mL, 5	1 @ 44.91	NO
1763T		100 units (30 units- 70 units) per mL, 3 mL, 5	1 @ 44.91	LY,NI,NO
2062M	Interferon Alfa-2a	100 units (50 units- 50 units) per mL, 3 mL, 5	1 @ 44.91	NO
8180M		3,000,000 i.u. in 0.5 mL	1 @ 34.43	RO
8551C		4,500,000 i.u. in 0.5 mL	1 @ 53.27	RO
8552D		6,000,000 i.u. in 0.5 mL	1 @ 69.80	RO
8553E		9,000,000 i.u. in 0.5 mL	1 @ 103.28	RO
8181N		3,000,000 i.u. in 0.5 mL	1 @ 34.43	RO
8182P		4,500,000 i.u. in 0.5 mL	1 @ 53.27	RO
8183Q		6,000,000 i.u. in 0.5 mL	1 @ 69.80	RO
8184R		9,000,000 i.u. in 0.5 mL	1 @ 103.28	RO
8572E		18,000,000 i.u. in 1.2 mL	1 @ 206.54	SH
8348J	Interferon Alfa-2b	18,000,000 i.u. in 1.2 mL	1 @ 206.54	SH
8476D		30,000,000 i.u. in 1.2 mL	1 @ 344.24	SH
8671J	Ipratropium Bromide	21 mcg per dose (200 doses)	1 @ 17.42	BY
1542E		250 mcg (anhydrous) in 1 mL, 30	1 @ 20.34	AF,AW,CH, DP,GX,MX, PU,TW
			1 @ 20.81	BY
8238N		500 mcg (anhydrous) in 1 mL, 30	1 @ 24.04	AF,AW,CH, DP,GX,MX, PU,TW
			1 @ 24.51	BY
1541D		250 mcg (anhydrous) per mL, 20 mL	1 @ 6.06	BY
8415X	Irinotecan Hydrochloride Trihydrate	100 mg in 5 mL	1 @ 390.00	MX,PU
2587E	Isosorbide Dinitrate	10 mg	100 @ 3.74	AF
			100 @ 5.59	SI
2588F	Ketoprofen	5 mg	100 @ 4.35	SI
1588N		100 mg	20 @ 8.32	AV
8797B	Levodopa with Carbidopa and Entacapone	50 mg-12.5 mg-200 mg	100 @ 157.53	NV
8798C		100 mg-25 mg-200 mg	100 @ 173.06	NV
8799D	Levonorgestrel	150 mg-37.5 mg-200 mg	100 @ 188.58	NV
2913H		30 mcg	1 @ 2.64	SC,WY
1456P	Levonorgestrel with Ethinylestradiol	Tablet-Pack	1 @ 2.64	SC
1393H		150 mcg-30 mcg	1 @ 5.08	SC
1394J		Tablet-Pack	1 @ 2.64	SY,WX
			1 @ 5.08	SC,WY
1392G		Tablet-Pack	1 @ 2.64	SY,WX
			1 @ 5.08	SC,WY
8290H	Lithium Carbonate	450 mg (s.r.)	100 @ 11.02	GK
1598D	Mercaptopurine	50 mg	25 @ 24.62	GK
8616L	Mesalazine	2 g in 60 mL, 7	1 @ 85.06	OA
8617M		4 g in 60 mL, 7	1 @ 113.39	OA
8768L		80 g	1 @ 85.06	OA
8753Q	Methysergide	1 g in 100 mL, 7	1 @ 85.06	FP
2826R		1 mg	50 @ 19.35	NV

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Approved Name	Form/Strength	Pack and Price \$	Manufacturer
1638F	Metronidazole	500 mg in 100 mL	1 @ 7.69	BX
8282X	Milk Powder—Lactose Free Formula	900 g	1 @ 17.05	WY
2350Q		900 g	1 @ 17.05	NU
8283Y		900 g	1 @ 17.05	WY
2349P		900 g	1 @ 17.05	NU
2358D	Milk Powder—Lactose Modified	900 g	1 @ 17.57	SJ
2357C		900 g	1 @ 17.57	SJ
3092R	Milk Powder—Synthetic	400 g	1 @ 36.40	NU
8630F	Milk Protein and Fat Formula with Vitamins and Minerals— Carbohydrate Free	225 g	1 @ 27.64	SB
8816B	Modafinil	100 mg	60 @ 175.67	CS
8635L	Moxifloxacin Hydrochloride	400 mg (base) in 250 mL	1 @ 64.80	BN
1674D	Naproxen	250 mg	50 @ 4.81	AF
			50 @ 6.36	RO
8298R	Naratriptan Hydrochloride	2.5 mg (base)	2 @ 11.00	GK
2732T	Nitrazepam	5 mg	25 @ 1.94	AF
			25 @ 3.96	VT
1967M	Norethisterone	350 mcg	1 @ 2.64	JC,KR
			1 @ 3.64	PH
2772X	Norethisterone with Ethinylestradiol	500 mcg-35 mcg	1 @ 4.63	PH
2774B		Tablet-Pack	1 @ 2.64	KR
			1 @ 4.63	PH
2773Y		1 mg-35 mcg	1 @ 4.63	PH
2775C		Tablet-Pack	1 @ 2.64	KR
			1 @ 4.63	PH
2776D		Tablet-Pack	1 @ 2.64	KR
			1 @ 4.63	PH
3176E	Norethisterone with Mestranol	1 mg-50 mcg	1 @ 2.64	PH
3179H		Tablet-Pack	1 @ 2.64	PH
1776L	Oestriol	1 mg	30 @ 3.01	OR
1733F	Oestrogens—Conjugated	300 mcg	28 @ 2.81	WY
1734G		625 mcg	28 @ 3.82	WY
8383F	Ofloxacin	3 mg per mL, 5 mL	1 @ 11.58	AG
3134Y	Oxazepam	15 mg	25 @ 1.29	AF
			25 @ 2.73	SI
3135B		30 mg	25 @ 1.50	AF,FM
			25 @ 2.95	SI
8588B	Oxcarbazepine	60 mg per mL, 250 mL	1 @ 72.23	NV
3026G	Paclitaxel	30 mg in 5 mL	1 @ 203.23	BQ,IT,MX
8018B		100 mg in 16.7 mL	1 @ 684.40	BQ,IT,MX
3017T		150 mg in 25 mL	1 @ 1016.11	IT,MX
8556H	Pancreatic Extract	not less than 5,000 BP units lipase activity	100 @ 23.32	SM
8020D		not less than 10,000 BP units lipase activity	100 @ 33.97	SM
8021E		not less than 25,000 BP units lipase activity	100 @ 67.94	SM
2495H	Pancrelipase	not less than 10,000 BP units lipase activity	250 @ 86.37	OR
8366H		not less than 25,000 BP units lipase activity	100 @ 67.94	TM

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
8784H	Paracetamol	500 mg	100 @ 2.64	AW,DG,FM, HX,JT,PC, SW,TW
8814X		665 mg (m.r.)	96 @ 3.38	ME
			96 @ 5.82	GC
1754H	Paraffin	3.5 g	1 @ 6.67	IQ
			1 @ 7.67	AQ
1703P	Phenoxymethylpenicillin	250 mg	25 @ 3.39	SI
1787C		250 mg	25 @ 3.39	SI
3028J		500 mg	25 @ 5.04	SI
2356B		125 mg per 5 mL, 100 mL	1 @ 3.80	FM
			1 @ 4.52	SI
2354X		250 mg per 5 mL, 100 mL	1 @ 5.18	FM
			1 @ 5.91	SI
2334W	Polygeline	17.5 g per 500 mL, 500 mL	1 @ 13.20	AE
2642C	Potassium Chloride	600 mg	100 @ 2.98	SZ
			100 @ 4.31	NV
1920C	Prednisolone Sodium Phosphate	equiv. to 20 mg prednisolone in 100 mL	7 @ 22.23	SI
2554K		equiv. to 5 mg prednisolone, 10	1 @ 7.73	SI
1948M	Promethazine Hydrochloride	50 mg in 2 mL	5 @ 6.77	MX
1953T	Propantheline Bromide	15 mg	100 @ 9.90	SI
1955X	Propylthiouracil	50 mg	100 @ 13.49	PL
2676W	Protein Hydrolysate Formula with Medium Chain Triglycerides	400 g	1 @ 11.87	NT
8259Q		450 g	1 @ 13.35	NU
8284B	Raltitrexed	2 mg	1 @ 292.75	AP
1937Y	Ranitidine Hydrochloride	150 mg (base), effervescent	30 @ 9.71	GK
8903N		150 mg (base), effervescent	30 @ 9.71	GK
8162N		150 mg (base) per 10 mL, 300 mL	1 @ 9.71	GK
8905Q		150 mg (base) per 10 mL, 300 mL	1 @ 9.71	GK
8787L	Risperidone	0.5 mg	20 @ 11.47	JC
8788M		0.5 mg (orally disintegrating)	28 @ 16.07	JC
8790P		1 mg (orally disintegrating)	28 @ 32.13	JC
8869T		0.5 mg	20 @ 11.47	JC
8870W		0.5 mg (orally disintegrating)	28 @ 16.07	JC
8792R		1 mg (orally disintegrating)	28 @ 32.13	JC
8794W		2 mg (orally disintegrating)	28 @ 64.58	JC
8780D		25 mg	1 @ 162.31	JC
8781E		37.5 mg	1 @ 208.43	JC
8782F		50 mg	1 @ 254.90	JC
1103C	Salbutamol Sulfate	2 mg (base) per 5 mL, 150 mL	1 @ 3.89	GK
1099W		200 mcg (base)	100 @ 7.95	GK
8288F		100 mcg (base) per dose (200 doses)	1 @ 5.81	AL,AW
			1 @ 6.12	MM
			1 @ 6.31	GK
8354Q		100 mcg (base) per dose (200 doses)	1 @ 15.90	MM
2000G		2.5 mg (base) in 2.5 mL, 30	1 @ 9.43	AF,AW,CH, GX,PU,TW
			1 @ 10.53	GK
2001H		5 mg (base) in 2.5 mL, 30	1 @ 9.97	AF,AW,CH, GX,TW
			1 @ 11.07	GK

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Approved Name	Form/Strength	Pack and Price \$	Manufacturer
2003K	Salbutamol Sulfate	5 mg (base) per mL, 30 mL	1 @ 3.45	PU
2995P	Salcatonin	50 i.u. in 1 mL	5 @ 34.56	NV
2997R		100 i.u. in 1 mL	5 @ 53.28	NV
2014B	Sodium Alginate with Calcium Carbonate and Sodium Bicarbonate	1 g-320 mg-534 mg in 20 mL, 500 mL	1 @ 4.08	RC
2264E	Sodium Chloride	154 mmol per L, 1 L	1 @ 3.85	BX
2260Y		513 mmol per L, 1 L	1 @ 5.83	BX
2266G	Sodium Chloride Compound	1 L	1 @ 5.83	BX
2281C	Sodium Chloride with Glucose	31 mmol-222 mmol per L, 1 L	1 @ 3.85	BX
2279Y		19 mmol-104 mmol per 500 mL, 500 mL	1 @ 5.06	BX
2278X		39 mmol-69 mmol per 500 mL, 500 mL	1 @ 5.06	BX
1124E	Sodium Cromoglycate	20 mg in 2 mL	60 @ 27.73	AV
2286H	Sodium Lactate Compound	1 L	1 @ 3.85	BX
2294R	Sodium Valproate	100 mg	100 @ 12.46	SW
2289L		200 mg (e.c.)	100 @ 15.82	AF
			100 @ 16.48	SW
2290M		500 mg (e.c.)	100 @ 31.30	AF
			100 @ 32.12	SW
2293Q		200 mg per 5 mL, 300 mL	1 @ 14.08	SW
2295T		200 mg per 5 mL, 300 mL	1 @ 14.08	SW
2091C	Sorbitol with Sodium Citrate and Sodium Lauryl Sulfoacetate	3.125 g-450 mg-45 mg in 5 mL, 12	1 @ 14.89	PH
8577K	Soy Protein and Fat Formula with Vitamins and Minerals—Carbohydrate Free	384 mL	1 @ 5.72	AB
2093E	Sulfasalazine	500 mg	100 @ 24.27	PH
2096H		500 mg (e.c.)	100 @ 26.70	KR
			100 @ 27.52	PH
2047R	Sulindac	100 mg	50 @ 4.91	AF
8144P	Sumatriptan Succinate	50 mg (base)	2 @ 11.00	AW,GK
2109B	Tamoxifen Citrate	10 mg (base)	30 @ 22.48	AP
2110C		20 mg (base)	30 @ 39.05	AP
2088X	Temazepam	10 mg	25 @ 1.94	AF,FM
			25 @ 3.00	SI
8819E	Temozolomide	5 mg	5 @ 72.58	SH
8820F		20 mg	5 @ 203.60	SH
8821G		100 mg	5 @ 872.19	SH
1251W	Terbutaline Sulfate	5 mg in 2 mL, 30	1 @ 11.06	AP
8098F	Testosterone	100 mg	1 @ 32.64	OR
8099G		200 mg	1 @ 65.27	OR
2670M	Testosterone Esters	100 mg	1 @ 4.70	OR
2101N		250 mg	1 @ 8.91	OR
8813W	Tetanus Vaccine, Adsorbed	0.5 mL	1 @ 4.08	CS
2832C	Tetracosactrin	1 mg in 1 mL	1 @ 13.41	NV
2345K	Thiotepa	15 mg	1 @ 18.71	SI
1356J	Tobramycin Sulfate	80 mg (base) in 2 mL (with preservative)	1 @ 6.05	AS
			5 @ 30.25	MX
8872Y		80 mg (base) in 2 mL (without preservative)	5 @ 30.25	PU
2117K	Triamcinolone Acetonide	200 mcg per g, 100 g	1 @ 4.28	FM
			1 @ 5.58	SI
2118L		200 mcg per g, 100 g	1 @ 4.28	FM
			1 @ 5.58	SI
3128P	Triglycerides, Medium Chain	500 mL	1 @ 23.56	SB

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
8478F	Triglycerides—Medium Chain, Formula	400 g	1 @ 39.49	SB
8629E		420 g	1 @ 43.13	SB
3136C	Triglycerides, Medium Chain and Long Chain with Glucose Polymer	400 g	1 @ 31.44	SB
8133C	Valaciclovir Hydrochloride	500 mg (base)	10 @ 58.66	GK
3113W	Vancomycin	125 mg	20 @ 125.42	LY
3114X		250 mg	20 @ 242.15	LY
3130R		500 mg	1 @ 23.58	MX
			1 @ 23.76	LY
3131T		500 mg	1 @ 23.58	MX
			1 @ 23.76	LY
2198Q	Vinblastine Sulfate	10 mg	1 @ 18.82	AS
2374Y	Vincristine Sulfate	1 mg in 1 mL	5 @ 81.68	MX,PU
8280T	Vinorelbine Tartrate	10 mg (base) in 1 mL	1 @ 78.37	FB,MX
8281W		50 mg (base) in 5 mL	1 @ 327.93	FB,MX
8587Y	Whey Protein Formula supplemented with Amino Acids, Vitamins and Minerals, and low in Protein, Phosphate, Potassium and Lactose	400 g	1 @ 48.88	SB
8266C	Zolmitriptan	2.5 mg	2 @ 10.97	AP
PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE				
5303D	Bisacodyl	10 mg, 10	1 @ 5.59	BY,PP
5304E		10 mg, 12	1 @ 4.25	FL,PP
5307H		10 mg, 10	1 @ 5.59	BY,PP
5308J		10 mg, 12	1 @ 4.25	FL,PP
5339B	Clonazepam	2.5 mg per mL, 10 mL	1 @ 4.27	RO
5342E		2.5 mg per mL, 10 mL	1 @ 4.27	RO
5361E	Diclofenac Sodium	25 mg (e.c.)	50 @ 4.38	AW,BG,CH, DP,GX,HX, TW
			50 @ 6.03	NV
5364H		25 mg (e.c.)	50 @ 4.38	AW,BG,CH, DP,GX,HX, TW
			50 @ 6.03	NV
5363G		100 mg	20 @ 9.14	NV
5366K		100 mg	20 @ 9.14	NV
5309K	Docusate Sodium with Bisacodyl	100 mg-10 mg, 5	1 @ 2.79	FM
5310L		100 mg-10 mg, 5	1 @ 2.79	FM
5311M	Glycerol	700 mg, 12	1 @ 3.34	PP
5312N		1.4 g, 12	1 @ 3.47	PP
5313P		2.8 g, 12	1 @ 3.59	PP
5314Q		700 mg, 12	1 @ 3.34	PP
5315R		1.4 g, 12	1 @ 3.47	PP
5316T		2.8 g, 12	1 @ 3.59	PP
5367L	Ibuprofen	200 mg	50 @ 2.18	AF
5369N		200 mg	50 @ 2.18	AF
5377B	Indomethacin	25 mg	50 @ 1.99	AF
			50 @ 3.52	MK
5379D		25 mg	50 @ 1.99	AF
			50 @ 3.52	MK
5378C		100 mg	20 @ 7.94	MK
5380E		100 mg	20 @ 7.94	MK

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Approved Name	Form/Strength	Pack and Price \$		Manufacturer
5345H	Naproxen	250 mg	50 @	4.81	AF
			50 @	6.36	RO
5349M		250 mg	50 @	4.81	AF
			50 @	6.36	RO
5359C	Nitrazepam	5 mg	25 @	1.94	AF
			25 @	3.96	VT
5360D		5 mg	25 @	1.94	AF
			25 @	3.96	VT
5371Q	Oxazepam	15 mg	25 @	1.29	AF
			25 @	2.73	SI
5372R		30 mg	25 @	1.50	AF,FM
			25 @	2.95	SI
5373T		15 mg	25 @	1.29	AF
			25 @	2.73	SI
5374W		30 mg	25 @	1.50	AF,FM
			25 @	2.95	SI
5331N	Sorbitol with Sodium Citrate and Sodium Lauryl Sulfoacetate	3.125 g-450 mg-45 mg in 5 mL, 12	1 @	14.89	PH
5332P		3.125 g-450 mg-45 mg in 5 mL, 12	1 @	14.89	PH
5381F	Sulindac	100 mg	50 @	4.91	AF
5383H		100 mg	50 @	4.91	AF
5375X	Temazepam	10 mg	25 @	1.94	AF,FM
			25 @	3.00	SI
5376Y		10 mg	25 @	1.94	AF,FM
			25 @	3.00	SI

PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING DENTAL PRACTITIONERS
FOR DENTAL TREATMENT ONLY

3398W	Benzylpenicillin	600 mg	5 @	8.75	CS
3399X		3 g	5 @	22.88	CS
5048Q	Cefotaxime	1 g	1 @	5.16	SZ
5049R		2 g	1 @	9.54	SZ
5076E	Diclofenac Sodium	25 mg (e.c.)	50 @	4.38	AW,BG,CH, DP,GX,HX, TW
			50 @	6.03	NV
5079H		100 mg	20 @	9.14	NV
5080J		250 mg	50 @	5.27	MK
5088T	Erythromycin Lactobionate	1 g (base)	1 @	9.20	AB
5106R	Glucose	278 mmol per L, 1 L	1 @	3.85	BX
5118J	Hydrocortisone Sodium Succinate	100 mg with 2 mL solvent	1 @	5.40	PH
5119K		250 mg with 2 mL solvent	1 @	7.52	PH
5121M	Ibuprofen	200 mg	50 @	2.18	AF
5126T	Indomethacin	25 mg	50 @	1.99	AF
			50 @	3.52	MK
5128X		100 mg	20 @	7.94	MK
5139L		100 mg	20 @	8.32	AV
5154G	Metronidazole	500 mg in 100 mL	1 @	7.69	BX
5176K	Naproxen	250 mg	50 @	4.81	AF
			50 @	6.36	RO
5224Y	Paracetamol	500 mg	100 @	2.64	AW,DG,FM, HX,JT,PC, SW,TW

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$		Manufacturer
3360W	Phenoxymethylpenicillin	250 mg	25 @	3.39	SI
3361X		500 mg	25 @	5.04	SI
3365D		125 mg per 5 mL, 100 mL	1 @	3.80	FM
			1 @	4.52	SI
3366E		250 mg per 5 mL, 100 mL	1 @	5.18	FM
			1 @	5.91	SI
3374N	Promethazine Hydrochloride	50 mg in 2 mL	5 @	6.77	MX
5212H	Sodium Chloride	154 mmol per L, 1 L	1 @	3.85	BX
5213J		513 mmol per L, 1 L	1 @	5.83	BX
5214K	Sodium Chloride with Glucose	31 mmol-222 mmol per L, 1 L	1 @	3.85	BX
5215L		19 mmol-104 mmol per 500 mL, 500 mL	1 @	5.06	BX
5216M		39 mmol-69 mmol per 500 mL, 500 mL	1 @	5.06	BX
5217N	Sulindac	100 mg	50 @	4.91	AF
5251J	Tetanus Vaccine, Adsorbed	0.5 mL	1 @	4.08	CS
3323X	Vancomycin	500 mg	1 @	23.58	MX
			1 @	23.76	LY

Section 4

Drug Tariff

Container Prices

Standard Formulae Preparations

Table of Codes, Maximum Quantities and Number of
Repeats for Extemporaneously Prepared
Pharmaceutical Benefits

Special Note:

Purified Water BP is the minimum requirement for water in all PBS extemporaneous preparations.

Drug Tariff

Drug	Standard	Recovery Prices			
		0.1 g/mL	1 g/mL	10 g/mL	100 g/mL
		\$	\$	\$	\$
Acacia Mucilage (by weight)	APF 15	0.01	0.09	0.72	6.44
Acacia, powdered	BP	0.02	0.18	1.40	12.48
Acetic Acid (6 per cent)	BP	0.01	0.01	0.10	0.85
Acetic Acid (33 per cent)	BP	0.01	0.04	0.33	2.90
Acetone	BP	0.01	0.07	0.52	4.63
(use as additive only)					
Alum	BP	0.02	0.14	1.15	10.25
Aluminium Acetate Solution	BP	0.02	0.12	0.95	8.44
Anise Water Concentrated 1 in 40	BP	0.01	0.05	0.36	3.20
(use as additive only)					
Aqueous Cream	APF	0.01	0.02	0.16	1.40
(for use only as a base combined with active ingredients)					
Ascorbic Acid	BP	0.03	0.25	2.03	18.02
(for use only as an ingredient of ferrous sulfate mixtures)					
Aspirin	BP	0.04	0.28	2.22	19.71
Belladonna Tincture	BP	0.02	0.19	1.55	13.78
Benzocaine	BP	0.06	0.51	4.10	36.40
Benzoic Acid	BP	0.03	0.22	1.77	15.77
Benzoic Acid Compound Ointment	APF	0.01	0.08	0.63	5.56
Benzoic Acid Solution	BP	0.01	0.08	0.62	5.55
Benzooin Compound Tincture	BP	0.02	0.15	1.16	10.29
Boric Acid	BP	0.01	0.07	0.59	5.21
(use as additive only)					
Boric Acid, Olive Oil and Zinc Oxide Ointment	QHF	0.01	0.06	0.47	4.18
Calcium Hydroxide	BP	0.04	0.28	2.23	19.81
Calcium Hydroxide Solution	BP	0.01	0.01	0.08	0.67
Castor Oil	BP	0.01	0.05	0.37	3.25
(use as additive only)					
Cetomacrogol Aqueous Cream	APF	0.01	0.02	0.19	1.66
(for use only as a base combined with active ingredients)					
Cetrimide Aqueous Cream	APF	0.01	0.11	0.88	7.79
(for use only as a base combined with active ingredients)					
Chlorhexidine Acetate	BP	0.44	3.53	28.20	250.67
(use as additive only)					
Chlorhexidine Aqueous Cream	APF	0.02	0.19	1.51	13.39
(for use only as a base combined with active ingredients)					
Chloroform	BP	0.02	0.12	0.94	8.35
(use as additive only)					
Chloroform Spirit	BP	0.01	0.03	0.23	2.05
Chloroform Water Concentrated 1 in 40	APF 15	0.01	0.03	0.22	1.98
Citric Acid Monohydrate	BP	0.01	0.05	0.43	3.80
Coal Tar	BP	0.05	0.42	3.36	29.89

Drug	Standard	Recovery Prices			
		0.1 g/mL	1 g/mL	10 g/mL	100 g/mL
		\$	\$	\$	\$
Coal Tar Solution	BP	0.01	0.09	0.72	6.43
Cocaine Hydrochloride	BP	4.43	35.44	283.53	2520.28
Coconut Oil	BP	0.01	0.10	0.78	6.89
Codeine Linctus	APF	0.01	0.06	0.45	4.03
Codeine Phosphate (may only be prescribed in linctuses, mixtures or mixtures for children)	BP	0.50	4.02	32.13	285.64
Collodion Flexible	BP	0.04	0.32	2.57	22.88
Dithranol	BP	3.18	25.47	203.73	1810.97
Emulsifying Ointment (for use only as a base combined with active ingredients)	BP	0.01	0.06	0.45	4.02
Ephedrine Hydrochloride (may only be prescribed in nasal instillations)	BP	0.19	1.54	12.31	109.45
Ethanol (90 per cent) (use as additive only)	BP	0.01	0.02	0.17	1.53
Ethanol (96 per cent) (use as additive only)	BP	0.01	0.03	0.22	1.96
Ether Solvent (use as additive only)	BP	0.01	0.11	0.87	7.71
Eucalyptus Oil (use as additive only)	BP	0.01	0.09	0.75	6.63
Ferrous Sulfate	BP	0.05	0.41	3.31	29.40
Formaldehyde Solution	BP	0.02	0.17	1.32	11.73
Gentian Alkaline Mixture	APF	0.01	0.06	0.50	4.48
Glycerol	BP	0.01	0.04	0.34	3.06
Honey Purified (use as additive only)	BP 1993	0.01	0.02	0.15	1.37
Hydroxybenzoate Compound Solution	APF	0.06	0.44	3.52	31.26
Iodine	BP	0.17	1.33	10.66	94.77
Iodine Alcoholic Solution	BP	0.01	0.11	0.86	7.66
Iodine Aqueous Oral Solution	BP	0.02	0.17	1.35	12.00
Kaolin Mixture	BPC 1968	0.01	0.07	0.59	5.24
Kaolin and Opium Mixture	APF 14	0.01	0.06	0.45	4.00
Lactic Acid	BP	0.05	0.38	3.03	26.96
Lavender Spike Oil	BPC 1968	0.07	0.57	4.52	40.18
Liquorice Liquid Extract	BP	0.03	0.23	1.80	16.02
Magnesium Carbonate Light	BP	0.01	0.10	0.77	6.87
Magnesium Sulfate (may only be prescribed for other than oral use)	BP	0.01	0.01	0.10	0.93
Magnesium Trisilicate	BP	0.02	0.17	1.32	11.73
Menthol, Racemic or Levomenthol	BP	0.17	1.32	10.52	93.51
Methyl Hydroxybenzoate	BP	0.22	1.76	14.07	125.07
Methyl Hydroxybenzoate Solution	APF	0.03	0.20	1.56	13.91

Drug	Standard	Recovery Prices			
		0.1 g/mL	1 g/mL	10 g/mL	100 g/mL
		\$	\$	\$	\$
Methylated Industrial Spirit (use as additive only)	BP	0.01	0.03	0.21	1.90
Olive Oil (use as additive only)	BP	0.01	0.06	0.44	3.91
Paraffin Hard	BP	0.01	0.06	0.45	4.00
Paraffin Liquid (may only be prescribed for other than oral use)	BP	0.01	0.02	0.19	1.66
Paraffin Light Liquid	BP	0.01	0.08	0.66	5.85
Paraffin Soft White	BP	0.01	0.03	0.23	2.06
Paraffin Soft Yellow	BP	0.01	0.04	0.28	2.47
Peppermint Oil (use as additive only)	BP	0.09	0.72	5.78	51.37
Peppermint Water Concentrated 1 in 40 (use as additive only)	APF 16	0.02	0.18	1.41	12.51
Phenobarbitone Sodium (may only be prescribed for the treatment of epilepsy)	BP	10.88	87.00	696.00	6186.67
Phenol Liquefied (not available for ear drops)	BP	0.05	0.37	2.93	26.00
Pholcodine Citrate Syrup (use as additive only)	BPC 1959	0.01	0.09	0.68	6.06
Podophyllum Resin	BP	0.39	3.13	25.03	222.51
Potassium Citrate	BP	0.01	0.08	0.67	5.97
Potassium Iodide	BP	0.05	0.43	3.41	30.33
Potassium Permanganate	BP	0.02	0.14	1.08	9.57
Propyl Hydroxybenzoate	BP	0.14	1.11	8.86	78.72
Propylene Glycol	BP	0.01	0.07	0.58	5.11
Red Syrup	APF 15	0.02	0.13	1.01	8.98
Resorcinol	BP	0.07	0.56	4.51	40.12
Salicylic Acid	BP	0.02	0.19	1.51	13.40
Salicylic Acid Ointment	APF	0.01	0.08	0.64	5.66
Salicylic Acid Ointment	BP	0.01	0.08	0.64	5.66
Simple Ointment (white) (for use only as a base combined with active ingredients)	BP	0.02	0.13	1.01	8.97
Simple Ointment (yellow) (for use only as a base combined with active ingredients)	BP	0.02	0.13	1.01	8.97
Sodium Bicarbonate	BP	0.01	0.04	0.31	2.73
Sodium Chloride	BP	0.01	0.06	0.48	4.29
Sodium Chloride Solution	BP	0.01	0.01	0.05	0.45
Sodium Citrate	BP	0.01	0.08	0.60	5.32
Sodium Thiosulfate (use as additive only)	BP	0.01	0.10	0.80	7.14
Starch	BP	0.01	0.06	0.47	4.21
Sulfur Ointment (for use only as a base combined with active ingredients)	BP 1980	0.02	0.13	1.01	8.94
Sulfur Precipitated	BP 1980	0.01	0.11	0.86	7.66
Syrup	BP	0.01	0.03	0.22	1.95

Drug	Standard	Recovery Prices			
		0.1 g/mL	1 g/mL	10 g/mL	100 g/mL
		\$	\$	\$	\$
Talc Purified, sterilised	BP	0.01	0.10	0.76	6.75
Thymol	BP	0.15	1.20	9.62	85.47
Thymol Compound Mouth Wash	APF 15	0.01	0.07	0.59	5.23
Tragacanth Compound Powder	BP 1980	0.06	0.44	3.50	31.09
Tragacanth Mucilage	APF 13	0.01	0.02	0.18	1.61
Tragacanth Mucilage	BPC 1973	0.01	0.02	0.12	1.09
Tragacanth, powdered	BP	0.06	0.48	3.86	34.32
Trichloroacetic Acid	BP 1980	0.24	1.88	15.03	133.62
Triethanolamine	BP	0.04	0.30	2.36	21.00
Water For Injections, sterilised (b) (extemporaneously prepared eye drops and eye lotions)	BP	..	..	..	11.23
Water Purified	BP	0.01	0.01	0.04	0.39
Wool Alcohols Ointment (white) (for use only as a base combined with active ingredients)	BP	0.02	0.12	0.94	8.39
Wool Alcohols Ointment (yellow) (for use only as a base combined with active ingredients)	BP	0.02	0.12	0.94	8.39
Wool Fat	BP	0.01	0.09	0.74	6.60
Wool Fat Hydrous	BP	0.01	0.08	0.67	5.93
Zinc Compound Paste	BP	0.02	0.13	1.07	9.52
Zinc Cream (for use only as a base combined with active ingredients)	BP	0.01	0.06	0.48	4.22
Zinc Oxide	BP	0.01	0.07	0.57	5.06
Zinc and Salicylic Acid Paste	BP	0.03	0.27	2.18	19.41
Zinc Sulfate	BP	0.01	0.09	0.69	6.16

Container Prices

	\$
DISPENSING BOTTLES—	
25 mL	0.43
50 mL	0.41
100 mL	0.40
200 mL	0.46
500 mL	1.05
POISON BOTTLES—	
25 mL	0.36
50 mL	0.39
100 mL	0.40
200 mL	0.56
500 mL	1.16
SCREW CAP JARS—	
25 g	0.48
50 g	0.59
100 g	0.62
200 g	0.57
500 g	1.07
DROPPER CONTAINERS—	
15 mL polythene	0.61
15 mL glass	0.48

Dispensing Fee for Extemporaneously Prepared Benefits	\$6.78
Additional Fee for Agreed Price Extemporaneously Prepared Benefits	\$1.38

Standard Formula Preparations

The following list is not intended to indicate in any way which particular formula an approved pharmacist should use in filling a prescription.

The prices shown in the column 'Dispensed Price for Max. Qty' are for the ingredients, the container and the dispensing fee. The prices shown in the column 'Maximum Recordable Value for Safety Net' are for the ingredients, the container and the dispensing fee and, where applicable, the additional fee for agreed price benefits.

KEY TO REFERENCES:

APF	Australian Pharmaceutical Formulary
BP	British Pharmacopoeia
BPC	British Pharmaceutical Codex
QHF	Queensland Hospital Formulary

Code	Item	Reference	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$
	CREAMS (Maximum Quantity 100 g and 1 Repeat)			
7502W	Salicylic Acid and Sulfur Aqueous	APF	9.34	10.72
	DUSTING POWDERS (Maximum Quantity 100 g and 1 Repeat)			
7458M	Zinc, Starch and Talc	APF 15 & BPC 1973	13.76	15.14
	EAR DROPS (Maximum Quantity 15 mL and 2 Repeats)			
7642F	Aluminium Acetate	APF	8.38	9.76
7643G	Aluminium Acetate	BP	8.82	10.20
7314Y	Sodium Bicarbonate	APF & BP	7.65	9.03
7313X	Spirit	APF	7.60	8.98

—CONTAINER RATES ARE INCLUDED—

Code	Item	Reference	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$
	INHALATIONS (Maximum Quantity 50 mL and 1 Repeat)			
7484X	Benzoin and Menthol	APF	14.29	15.67
7308P	Menthol	APF	9.34	10.72
7310R	Menthol and Eucalyptus	BP 1980	9.47	10.85
	LINCTUSES CONTAINING CODEINE PHOSPHATE (Maximum Quantity 100 mL and no Repeats)			
7530H	Codeine	APF	11.21	12.59
	LOTIONS (Maximum Quantity 200 mL and 2 Repeats)			
7709R	Aluminium Acetate Aqueous	APF	9.03	10.41
	MIXTURES, OTHER (Maximum Quantity 200 mL and 4 Repeats)			
7604F	Gentian Alkaline	APF	16.20	17.58
7348R	Kaolin	BPC 1968	17.72	19.10
7301G	Kaolin and Opium	APF 14	15.24	16.62
7342K	Magnesium Trisilicate	BPC 1968	11.30	12.68
7343L	Magnesium Trisilicate and Belladonna	BPC 1968	12.77	14.15
	MOUTH WASHES (Maximum Quantity 200 mL and 1 Repeat)			
7457L	Thymol Compound	APF 15	17.80	19.18
	OINTMENTS (Maximum Quantity 100 g and 1 Repeat)			
7914M	Benzoic Acid Compound	APF	12.96	14.34
7914M	Benzoic Acid Compound (extemporaneous formula)	BP	12.96	14.34
7902X	Boric Acid, Olive Oil and Zinc Oxide	QHF	11.58	12.96
7926E	Salicylic Acid	APF	13.06	14.44
7928G	Salicylic Acid (extemporaneous formula)	BP	13.06	14.44
	PAINTS (Maximum Quantity 25 mL and 1 Repeat)			
7567G	Podophyllin Compound	APF 16 & BP	21.78	23.16
7568H	Salicylic Acid	APF	14.05	15.43
	PASTES, OTHER (Maximum Quantity 100 g and 1 Repeat)			
7558T	Zinc	APF	16.92	18.30
7558T	Zinc Compound (extemporaneous formula)	BP	16.92	18.30
	POWDER FOR INTERNAL USE (Maximum Quantity 100 g and 2 Repeats)			
7545D	Magnesium Trisilicate	BP	19.08	20.46

Table of Codes, Maximum Quantities, and Number of Repeats for Extemporaneously Prepared Benefits

Code	Preparation	Maximum Quantity	Number of Repeats
13Q	Creams	100 g	1
48M	Dusting Powders	100 g	1
15T	Ear Drops	15 mL	2
19B	Eye Drops containing Cocaine Hydrochloride	15 mL	..
22E	Eye Drops, Other	15 mL	5
23F	Eye Lotions	200 mL	2
29M	Inhalations	50 mL	1
64J	Linctuses containing Codeine Phosphate	100 mL	..
34T	Linctuses, Other	100 mL	2
39C	Lotions	200 mL	2
65K	Mixtures containing Codeine Phosphate	200 mL	..
40D	Mixtures, Other	200 mL	4
66L	Mixtures for Children containing Codeine Phosphate	100 mL	..
41E	Mixtures for Children, Other	100 mL	4
30N	Mouth Washes	200 mL	1
42F	Nasal Instillations	15 mL	2
43G	Ointments, Waxes	100 g	1
44H	Paints	25 mL	1
63H	Pastes containing Cocaine Hydrochloride	25 g	..
45J	Pastes, Other	100 g	1
49N	Powders for Internal Use	100 g	2
52R	Solutions	200 mL	2



Australian Government

Department of Veterans' Affairs

REPATRIATION SCHEDULE OF PHARMACEUTICAL BENEFITS

1 DECEMBER 2005


The benefits listed in this Schedule may only be prescribed to Department of Veterans' Affairs beneficiaries holding a:

- Repatriation Health Card For All Conditions (gold); or
- Repatriation Health Card For Specific Conditions (white); or
- Repatriation Pharmaceutical Benefits Card (orange);

BENEFICIARIES' ENTITLEMENT CARDS AND ELIGIBILITY

The diagram below outlines the drug eligibility of Department of Veterans' Affairs

REPATRIATION PHARMACEUTICAL BENEFITS



FOR REPATRIATION PHARMACEUTICAL BENEFITS

beneficiaries in accordance with their treatment entitlement.

PBS Schedule listings written on the common PBS/RPBS prescription form, ticked as 'RPBS' (restrictions apply)	+	Repatriation Schedule listings written on the common PBS/RPBS prescription form, ticked as 'RPBS'	+	Items for which prescribing approval has been authorised on the common PBS/RPBS authority prescription form for: (i) PBS and Repatriation Schedules 'Authority required' items; (ii) greater quantities/repeats of drugs listed in PBS and Repatriation Schedules; or (iii) items not listed in PBS or Repatriation Schedules.

Only for disabilities that have been accepted for treatment by the Department of Veterans' Affairs

PBS Schedule listings written on the common PBS/RPBS prescription form, ticked as 'RPBS' (restrictions apply)	+	Repatriation Schedule listings written on the common PBS/RPBS prescription form, ticked as 'RPBS'	+	Items for which prescribing approval has been authorised on a common PBS/RPBS authority prescription form for: (i) PBS and Repatriation Schedules 'Authority required' items; (ii) greater quantities/repeats of drugs listed in PBS and Repatriation Schedules; or (iii) items not listed in PBS or Repatriation Schedules.

RPBS Explanatory Notes

Introduction

The Australian Repatriation System

1. The Australian Repatriation system is based primarily on the principle of compensation to veterans and eligible dependants for injury or death related to war service. In certain cases, treatment is also provided for accepted injuries or conditions that are not service-related or have occurred as a result of other than war service.
2. Through the *Veterans' Entitlements Act 1986* the Department of Veterans' Affairs provides programs of compensation, income support and treatment for eligible veterans and their dependants. One of the defined benefits for eligible veterans is the Repatriation Pharmaceutical Benefits Scheme. This range of medications and dressings is more comprehensive than is available through the Pharmaceutical Benefits Scheme.

RPBS prescribing provisions

3. Unless otherwise stated, Repatriation Pharmaceutical Benefits Scheme (RPBS) prescriptions must conform with the requirements of Pharmaceutical Benefits Scheme (PBS) prescriptions, as detailed in Section 1 - Explanatory Notes in the *Schedule of Pharmaceutical Benefits* book. The prescriber shall ensure that a prescription contains the following details:
 - the category of benefit, i.e., RPBS, by placing a cross in the relevant box;
 - the patient's full name and address;
 - the prescription date;
 - the DVA file number of the patient as evidence of entitlement;
 - in the case of authority prescriptions, the Authority approval number;
 - the item, form, strength, quantity and directions;
 - the number of repeats, if applicable;
 - indicate when brand substitution is not permitted; and
 - the name, signature, the prescriber number and address of the prescriber.

Prior Approval Arrangements

4. The prior approval of the Department is required to prescribe the following:
 - 'Authority required' items listed in either the PBS or RPBS Schedule;
 - increased quantities and/or repeats of items listed in either the PBS or RPBS Schedule;
 - items listed under section 100 of the *National Health Act 1953*; and
 - other items not listed in either Schedule (non-Schedule items).
5. The above items are to be prescribed on the common PBS/RPBS authority prescription form in accordance with the directions stated in Section 1 - Explanatory Notes in the *Schedule of Pharmaceutical Benefits* book.

6. All authority prescriptions and requests for non-Schedule items must receive prior approval from the Department. This can be achieved by either:
 - using the Department's national free call number 1800 552 580; or
 - by mailing the written authority prescription to the Veterans' Affairs Pharmaceutical Approvals Centre (VAPAC) at the reply paid address shown at the end of these RPBS Explanatory Notes.
7. Some requests for prior approval (including some non-Schedule items) need to be referred by VAPAC to the Repatriation Pharmaceutical Reference Committee for consideration. In such cases a VAPAC pharmacist will advise the prescriber to submit a request in writing that provides the following information:
 - A current clinical report on the patient's condition (such as age, co-morbidities, renal, liver failure) and clinical reports including pathology, biochemistry, diagnostic and other investigations if appropriate.
 - Details of past and current therapy for the condition. Include details of PBS, RPBS and non-Schedule items utilised, and the results of those therapies.
 - Details of the proposed treatment regimen. Include intended dose and duration of treatment and objectives measures of response.
 - When the proposed use of the item is outside the TGA-approved indications for use in Australia, provide copies of articles from peer reviewed publications supporting the proposed treatment.
 - Signed, informed patient consent where the item is to be used for a non-TGA-approved indication.
 - For items without Australian marketing approval, a copy of the TGA Special Access Scheme approval to prescribe the drug.
8. Requests for prior approval to prescribe a non-Schedule (PBS or RPBS) item that is of the same therapeutic class (ATC level 3) as an item that is listed on the Schedule, will not be approved unless unequivocal clinical evidence is presented to demonstrate that the requested item is essential for effective treatment of the nominated patient.
9. A pharmacist should not supply an item prescribed on an RPBS Authority Prescription Form unless the form has been approved and stamped by VAPAC, or has been endorsed by the prescriber with a telephone Authority approval number provided by VAPAC. Medicare Australia will not accept RPBS Authority prescriptions that have not been approved by the Department of Veterans' Affairs for payment.

Palliative Care Drugs

10. The following medications may be available, or made available in increased quantities or doses under prior approval arrangements for use only in the palliative care of terminal disease:

• clonazepam • cyclizine • dexamethasone • disodium pamidronate • fentanyl • glycopyrrolate	• hyoscine butylbromide • hyoscine hydrobromide • ketamine • midazolam • octreotide

For further information telephone VAPAC on 1800 552 580.

Dental Prescribing

11. Under Department of Veterans' Affairs arrangements, financial responsibility for pharmaceutical benefits prescribed by a Local Dental Officer (LDO) is limited to treatment to which holders of the following cards are entitled:

- a Gold Repatriation Health Card - For All Conditions; or
- a White Repatriation Health Card - For Specific Conditions; or
- an Orange Repatriation Pharmaceutical Benefits Card.

Where possible the LDO shall prescribe in accordance with the provisions governing dental prescribing under the Pharmaceutical Benefits Scheme (PBS).

12. Prescriptions for PBS Dental Schedule items for Gold, White and Orange Card holders are to be dispensed at the PBS concessional rate. Claims for payment by the dispensing pharmacist are to be included with other Repatriation prescriptions. The card holder is required to meet the cost of any applicable brand premium.
13. When a non-PBS Dental Schedule item is prescribed for an eligible card holder, the LDO's private prescription form should be used. The dispensing pharmacist may charge the patient the full cost of the prescription. The patient may claim a refund for the full cost of a non-Schedule item from the Department if an itemised receipt (not a cash register receipt) and a copy of the prescription are provided.

Provisions governing pricing and payment for RPBS benefits**Introduction**

14. Unless otherwise stated, the pricing and payment principles and arrangements for approved pharmacists supplying pharmaceutical benefits under the RPBS will be the same as those arrangements applying under the PBS.
15. Where a pharmaceutical benefit that is not listed on the PBS or RPBS Schedule is dispensed on an RPBS Authority prescription, a pharmacist will price the benefit and enter the serial number, prescription identifying number and price on the sticker or stamp imprint affixed to the prescription.

Pricing of Schedule Items

16. Items supplied under the RPBS from the PBS Schedule, both ready-prepared and extemporaneously-prepared, will be paid on the same basis as benefits supplied under the PBS. Items supplied under the RPBS from the Repatriation Schedule, including wound dressings, will be paid on the basis of the price as given in the Repatriation Pharmaceutical Benefits section (Section 1 – RPBS Schedule, Drugs, Medicines and Dressings) of the *Schedule of Pharmaceutical Benefits* book.

Pricing of Non-Schedule Ready Prepared Items

17. Non-Schedule ready-prepared items are to be priced on the basis of the invoiced, GST-exclusive wholesale price to pharmacists plus the appropriate PBS mark-up and the PBS dispensing fee. Where the item price to pharmacists is greater than \$100.00, a copy of the invoice pertaining to the supply of that item is to be submitted together with the appropriate copy of the authority prescription as part of the claim for payment.

Pricing of Non-Schedule Extemporaneously Prepared Items

18. When an ingredient drug is not listed in the PBS Drug Tariff (Section 4, PBS Schedule), the recovery price will be based on the invoiced wholesale price to pharmacists, increased by a mark-up of 100%, calculated in accordance with the directions contained in the pricing instructions for pricing of PBS extemporaneously-prepared benefits in this Schedule (Section 1 –, PBS Schedule Part 9, Pricing Prescriptions). The price paid by the pharmacist for the commercial pack from which the ingredient is used shall be endorsed on the prescription form.

Miscellaneous Pricing Rules

19. The price to pharmacists used as the basis of pricing will be the invoiced, GST-exclusive price from the wholesaler.
20. If multiple quantities of a manufacturer's original pack are supplied, the PBS mark-up is applied to the price to pharmacist of each pack and then totalled. The PBS dispensing fee, and the PBS dangerous drug fee if applicable, are then added to the total of the marked-up prices.
21. When the quantity prescribed corresponds with the quantity of a manufacturer's original pack, in no circumstances will the price payable for one pack exceed that payable for multiples or combinations of packs to supply the quantity prescribed.
22. The list of ingredient drugs and prices included in the PBS Drug Tariff are common to both the PBS and RPBS. Certain restrictions apply regarding the prescribing and dispensing of some of these ingredient drugs as pharmaceutical benefits, e.g., use as additive only.
23. For items prescribed generically, including non-Schedule and wound dressings, the pharmacist should indicate on the prescription the quantity and brand supplied. If prescriptions are not endorsed, the Department will pay the lowest priced acceptable product available.

General**Packaging Material, Postage or Freight**

24. Payment to a pharmacist for the costs of packaging materials, postage or freight required to supply a pharmaceutical benefit is to be paid by the patient, who may then claim reimbursement from the Department through the provision of a pharmacist's itemised receipt.

Payment for Items Supplied at Short Intervals

25. For all items dispensed at specific short intervals of time, the Department will pay a separate PBS dispensing fee for each occasion that the drug is supplied and which is acknowledged on receipt by the patient or agent.
26. The price payable on the items supplied will be based on the individual dose quantity supplied. Where applicable, a PBS dangerous drug fee and a minimum container charge will be payable for each supply.

Receipts for Patient Charges

27. Where a charge is paid by a patient in any of the circumstances of paragraphs 13 or 24, the pharmacist is required to provide a printed receipt to the patient with the details of the items or services provided, the amount paid, date of supply and the patient's name and address. The patient may apply for reimbursement from the Department.

Special Patient Contributions

28. The Special Patient Contribution for items listed as Special Pharmaceutical Benefits in the PBS Schedule is not payable by veterans entitled to pharmaceutical benefits under the RPBS. Eligible veterans receiving Special Pharmaceutical Benefits under the RPBS are required to pay only the concessional patient contribution and any applicable brand premium. If a Safety Net Entitlement card is held, the veteran should receive a Special Pharmaceutical Benefit free of charge, subject to any brand premium applicable. Medicare Australia will reimburse the dispensing pharmacist the total dispensed price, less the concessional patient contribution and/or brand premium if applicable.

Therapeutic Group Premiums – Authority Processing

29. Items attracting a therapeutic group premium are dual listed. Dispensing pharmacists are therefore required to select the appropriate code for those items that are dual listed as authority and non-authority items, in order to correctly charge the patient and claim from Medicare Australia. Those authority prescriptions that grant exemption from a therapeutic group premium will have the letters 'TPX' at the beginning of the telephone Authority approval number, or, in the case of a written approval, will be stamped with the words "This prescription does not attract a therapeutic group premium".

DEPARTMENT OF VETERANS' AFFAIRS
Authority Prescription Applications

Applications for authority to prescribe under the Repatriation Pharmaceutical Benefits Scheme (RPBS) should be sent to the Veterans' Affairs Pharmaceutical Approvals Centre (VAPAC) using the free postal service:

REPLY PAID No. 372
VAPAC (Veterans' Affairs Pharmaceutical Approvals Centre)
GPO Box 9998
BRISBANE QLD 4001

For RPBS enquiries and telephone approvals 24 hours a day the Freecall number is:

1800 552 580

Departmental pharmacists answer applications for prior approval for non-Schedule items and Authority application calls.

REPATRIATION PHARMACEUTICAL BENEFITS

This Schedule will take effect on 1 December 2005 and all previous issues are cancelled. New Schedules will take effect on 1 April, 1 August and 1 December each year.

SUMMARY OF CHANGES

ADDITIONS

Additions — Items

4806Y **Dressing—Hydrogel—Sheet**, dressings 11 cm x 10 cm, 5 (*Aquaclear*)

BIOEQUIVALENCE INDICATORS

The bioequivalence indicator (a) has been added to the following brands:

4238C *Xergic, AF; Telfast 120, AV* — **Fexofenadine Hydrochloride**, tablet 120 mg (effective 1 October 2005)

DELETIONS

Deletions — Items

4910K **Dressing—Hydrogel—Sheet**, dressing 10 cm x 10 cm (*Clearsite Borderless 409240*)
 4087D **Beclomethasone Dipropionate**, aqueous nasal spray 50 micrograms per dose, 400 doses set containing 1 pump pack (200 doses) and 1 refill (200 doses) (*Aldecin Aqueous Set*)

Deletions — Brands

4074K *Senagar, SI* — **Senega and Ammonia**, mixture 200 mL

ALTERATIONS

Alterations — Restriction Changes

The following item has changed from “authority required ” to “restricted benefit”:

4092J **Budesonide**, aqueous nasal spray (pump pack) 64 micrograms per dose (120 doses) (*Budamax Aqueous*)

Other restriction changes:

4092J **Budesonide**, aqueous nasal spray (pump pack) 64 micrograms per dose (120 doses) (*Budamax Aqueous*)

4570M **Orlistat**, capsule 120 mg (*Xenical*)

Alterations — Proprietary Name

4945G	Dressing—Hydrocolloid (Superficial Wound—Moderate Exudate) , dressings 10 cm x 10 cm, 10	From: <i>Hydrocoll Thick 900938/1</i> To: <i>Hydrocoll 900938/1</i>
4946H	Dressing—Hydrocolloid (Superficial Wound—Moderate Exudate) , dressings 15 cm x 15 cm, 10	From: <i>Hydrocoll Thick 900939/1</i> To: <i>Hydrocoll 900939/1</i>
4767X	Gauze and Cotton Tissue (Combine Roll) , wrapped pack 9 cm x 10 m	From: <i>BV 36121054</i> To: <i>BSN 2902165</i>

Alterations — Manufacturer's Code

		<i>From</i>	<i>To</i>
4117Q	Aluminium Hydroxide with Magnesium Hydroxide and Simethicone , tablet 400 mg-400 mg-30 mg (<i>Mylanta Double Strength</i>)	WR	PC
4118R	Aluminium Hydroxide with Magnesium Hydroxide and Simethicone , oral suspension 400 mg-400 mg-30 mg per 5 mL, 500 mL (<i>Mylanta Double Strength</i>)	WR	PC
4175R	Cetirizine Hydrochloride , tablet 10 mg (<i>Zyrtec</i>)	WR	PC
4222F	Fluorouracil , cream 50 mg per g (5%), 20 g (<i>Efudix</i>)	ID	VT
4420P	Pseudoephedrine Hydrochloride , tablet 60 mg (<i>Sudafed Sinus & Nasal Decongestant</i>)	WR	PC
4039N	Zinc Oxide , compound ointment 50 g (<i>Anusol</i>)	WW	PC
4040P	Zinc Oxide , compound suppositories, 12 (<i>Anusol</i>)	WW	PC

**Therapeutic Index
for
RPBS Schedule**

ALIMENTARY TRACT AND METABOLISM 472

STOMATOLOGICAL PREPARATIONS 472

Stomatological preparations 472

DRUGS FOR ACID RELATED DISORDERS 472

*Antacids 472**Drugs for peptic ulcer and gastro-oesophageal reflux disease (GORD) 472*

DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS 473

*Drugs for functional bowel disorders 473**Belladonna and derivatives, plain 473*

LAXATIVES 473

Laxatives 473

ANTIOBESITY PREPARATIONS, EXCL. DIET PRODUCTS 474

Antibesity preparations, excl. diet products 474

VITAMINS 475

*Vitamin B₁, plain and in combination with vitamin B₆ and vitamin B₁₂ 475**Vitamin B-complex, incl. combinations 475**Ascorbic acid (vitamin C), incl. combinations 475*

MINERAL SUPPLEMENTS 475

*Other mineral supplements 475***BLOOD AND BLOOD FORMING ORGANS 475**

ANTITHROMBOTIC AGENTS 475

Antithrombotic agents 475

BLOOD SUBSTITUTES AND PERFUSION SOLUTIONS 476

*Irrigating solutions 476***CARDIOVASCULAR SYSTEM 476**

VASOPROTECTIVES 476

Antihemorrhoidals for topical use 476

CALCIUM CHANNEL BLOCKERS 476

Selective calcium channel blockers with mainly vascular effects 476

AGENTS ACTING ON THE RENIN-ANGIOTENSIN SYSTEM 477

*ACE inhibitors, plain 477***DERMATOLOGICALS 477**

ANTIFUNGALS FOR DERMATOLOGICAL USE 477

*Antifungals for topical use 477**Antifungals for systemic use 478*

EMOLLIENTS AND PROTECTIVES 479

*Emollients and protectives 479**Protectives against UV-radiation 479*

ANTIPRURITICS, INCL. ANTIHISTAMINES, ANESTHETICS, ETC. 480

Antipruritics, incl. antihistamines, anesthetics, etc. 480

ANTIPSORIATICS 480

Antipsoriatics for topical use 480

ANTIBIOTICS AND CHEMOTHERAPEUTICS FOR DERMATOLOGICAL USE 480

*Antibiotics for topical use 480**Chemotherapeutics for topical use 481*

CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS 481

*Corticosteroids, plain 481**Corticosteroids, combinations with antibiotics 481*

ANTISEPTICS AND DISINFECTANTS 482

Antiseptics and disinfectants 482

OTHER DERMATOLOGICAL PREPARATIONS 482

Other dermatological preparations 482

THERAPEUTIC INDEX

GENITO URINARY SYSTEM AND SEX HORMONES 483

- GYNECOLOGICAL ANTIINFECTIVES AND ANTISEPTICS 483
 - Antiinfectives and antiseptics, excl. comb. with corticosteroids 483*
- OTHER GYNECOLOGICALS 484
 - Other gynecologicals 484*
- SEX HORMONES AND MODULATORS OF THE GENITAL SYSTEM 485
 - Estrogens 485*
- UROLOGICALS 485
 - Other urologicals, incl. antispasmodics 485*
 - Drugs used in benign prostatic hypertrophy 486*

ANTIINFECTIVES FOR SYSTEMIC USE 487

- ANTIBACTERIALS FOR SYSTEMIC USE 487
 - Macrolides, lincosamides and streptogramins 487*
 - Quinolone antibacterials 487*

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS 487

- ANTINEOPLASTIC AGENTS 487
 - Antimetabolites 487*
- IMMUNOSUPPRESSIVE AGENTS 488
 - Immunosuppressive agents 488*

MUSCULO-SKELETAL SYSTEM 491

- ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS 491
 - Antiinflammatory and antirheumatic products, non-steroids 491*
- TOPICAL PRODUCTS FOR JOINT AND MUSCULAR PAIN 491
 - Topical products for joint and muscular pain 491*
- DRUGS FOR TREATMENT OF BONE DISEASES 491
 - Drugs affecting bone structure and mineralization 491*

NERVOUS SYSTEM 492

- ANALGESICS 492
 - Opioids 492*
 - Other analgesics and antipyretics 492*
- PSYCHOLEPTICS 493
 - Anxiolytics 493*
 - Hypnotics and sedatives 494*
- OTHER NERVOUS SYSTEM DRUGS 494
 - Drugs used in addictive disorders 494*

ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS 495

- ANTHELMINTICS 495
 - Antinematodal agents 495*

RESPIRATORY SYSTEM 495

- NASAL PREPARATIONS 495
 - Decongestants and other nasal preparations for topical use 495*
 - Nasal decongestants for systemic use 496*
- DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES 496
 - Adrenergics, inhalants 496*
- COUGH AND COLD PREPARATIONS 496
 - Expectorants, excl. combinations with cough suppressants 496*
 - Cough suppressants, excl. combinations with expectorants 496*
- ANTIHISTAMINES FOR SYSTEMIC USE 496
 - Antihistamines for systemic use 496*

SENSORY ORGANS 497

- OPHTHALMOLOGICALS 497
 - Antiglaucoma preparations and miotics 497*
 - Decongestants and antiallergics 497*
- OTOLOGICALS 498
 - Corticosteroids and antiinfectives in combination 498*
 - Other otologicals 498*

THERAPEUTIC INDEX

VARIOUS 498

ALL OTHER THERAPEUTIC PRODUCTS 498

All other therapeutic products 498

ALL OTHER NON-THERAPEUTIC PRODUCTS 499

All other non-therapeutic products 499

Section 1

Drugs, Medicines and Dressings

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ALIMENTARY TRACT AND METABOLISM								
STOMATOLOGICAL PREPARATIONS								
Stomatological preparations								
• Antiinfectives and antiseptics for local oral treatment								
CHLORHEXIDINE GLUCONATE								
4161B	Mouth wash 2 mg per mL (0.2%), 250 mL	‡1	..	..	9.99 10.48	4.60 4.60	Plaqacide Savacol Mouth and Throat Rinse	OB OM
4160Y	Mouth wash 2 mg per mL (0.2%), 250 mL	‡1	..	..	11.78	4.60	Periogard (Chlorohex) Mouth Rinse	OM
• Other agents for local oral treatment								
CARMELLOSE SODIUM								
4568K	Mouth spray 10 mg per mL, 25 mL	‡1	1	..	8.28	4.60	Aquae	HA
4569L	Mouth spray 10 mg per mL, 100 mL	‡1	..	..	10.48	4.60	Aquae	HA
DRUGS FOR ACID RELATED DISORDERS								
Antacids								
• Calcium compounds								
CALCIUM CARBONATE with GLYCINE								
NOTE: For patients with chronic renal failure.								
4055K	Tablet 420 mg-180 mg	200	5	..	*19.69	4.60	Titralac	MM
• Combinations and complexes of aluminium, calcium and magnesium compounds								
ALUMINIUM HYDROXIDE with MAGNESIUM HYDROXIDE and SIMETHICONE								
4117Q	Tablet 400 mg-400 mg-30 mg	200	5	..	*24.11	4.60	Mylanta Double Strength	PC
4118R	Oral suspension 400 mg-400 mg- 30 mg per 5 mL, 500 mL	2	5	..	*20.27	4.60	Mylanta Double Strength	PC
Drugs for peptic ulcer and gastro-oesophageal reflux disease (GORD)								
• H₂-receptor antagonists								
NOTE: The base-priced drugs in this therapeutic group are cimetidine (except cimetidine effervescent tablet 800 mg (as hydrochloride)), famotidine, nizatidine and ranitidine hydrochloride (except ranitidine hydrochloride effervescent tablet 150 mg (base) and syrup 150 mg (base) per 10 mL, 300 mL).								
CIMETIDINE								
NOTE: <i>Helicobacter pylori eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.</i>								
Authority required <i>To be approved where other base-priced (benchmark) drug treatment is inappropriate.</i>								
4976X	Effervescent tablet 800 mg (as hydrochloride)	30	5	..	34.30	4.60	Tagamet 800 Express	GK

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
RANITIDINE HYDROCHLORIDE								
NOTE: <i>Helicobacter pylori eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.</i>								
Authority required <i>To be approved where other base-priced (benchmark) drug treatment is inappropriate.</i>								
4978B	Effervescent tablet 150 mg (base)	60	5	..	*24.17	4.60	Zantac	GK
4980D	Syrup 150 mg (base) per 10 mL, 300 mL	2	5	..	*24.17	4.60	Zantac Syrup	GK
DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS								
Drugs for functional bowel disorders								
• Synthetic anticholinergics, esters with tertiary amino group								
MEBEVERINE HYDROCHLORIDE								
4328T	Tablet 135 mg	90	..	..	25.51 28.65	4.60 4.60	^a Colese ^a Colofac	AF SM
Belladonna and derivatives, plain								
• Belladonna alkaloids semisynthetic, quaternary ammonium compounds								
HYOSCINE BUTYLBROMIDE								
4279F	Injection 20 mg in 1 mL	5	..	..	16.97	4.60	Buscopan	BY
LAXATIVES								
Laxatives								
• Softeners, emollients								
DOCUSATE SODIUM								
4200C	Tablet 50 mg	100	2	..	10.79	4.60	Coloxyl 50	FM
• Contact laxatives								
DOCUSATE SODIUM with SENNA								
4198Y	Tablet 50 mg-8 mg	90	2	..	12.40	4.60	Coloxyl with Senna	FM
SENNA STANDARDISED								
4455L	Tablet 7.5 mg	100	1	..	10.69	4.60	Senokot	RC
• Bulk producers								
ISPAGHULA HUSK								
4285M	Sachets 3.5 g, 30	‡1	1	..	13.70	4.60	Fybogel	RC
PSYLLIUM HYDROPHILIC MUCILLOID								
4419N	Oral powder (orange-flavoured, sugar-free) 283 g	‡1	1	..	19.34	4.60	Metamucil Smooth Texture Orange	PY
4422R	Oral powder (non-flavoured) 336 g	‡1	1	..	19.34	4.60	Metamucil Regular	PY
PSYLLIUM HYDROPHILIC MUCILLOID with HIGH AMYLOSE MAIZE STARCH								
4416K	Oral powder 2.7 g-0.7 g per 7.5 g, 440 g	‡1	1	..	18.95	4.60	Nucolox	SI

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
4558X	STERCULIA with FRANGULA BARK Granules 620 mg-80 mg per g (62%-8%), 500 g	‡1	1	..	23.05	4.60	Normacol Plus	NE
• Enemas								
4462W	SORBITOL with SODIUM CITRATE and SODIUM LAURYL SULFOACETATE Enemas 3.125 g-450 mg-45 mg in 5 mL, 4	‡1	..	..	10.18	4.60	MicroLax	PH
• Other laxatives								
GLYCEROL								
<u>Restricted benefit</u>								
<i>Short-term use when oral laxative therapy has failed or is inappropriate.</i>								
4246L	Suppositories 2.8 g (for adults), 12	3	..	..	*15.52	4.60	PP	

ANTI-OBESITY PREPARATIONS, EXCL. DIET PRODUCTS

Antiobesity preparations, excl. diet products• **Peripherally acting antiobesity products****ORLISTAT****Authority required**➤ ***For the treatment of obese patients.******Total treatment will not exceed 12 months from initial application.******Patients are eligible for 1 continuous treatment in a lifetime.******The patient must be receiving, or enrolled to receive, professional dietetic and weight management advice (where this is available).******Initial treatment for patients who meet the following criteria to qualify:******(a) Body Mass Index (BMI) greater than or equal to 35 with no known co-morbidities;******or******(b) BMI greater than or equal to 30 with 1 or more of the following co-morbidities:******(i) diabetes;******(ii) ischaemic heart disease;******(iii) psychiatric conditions;******(iv) hypertension.******The prescriber must provide the following:******(a) initial body weight; and******(b) BMI.******Continuing treatment for patients who have previously been issued with an authority prescription for orlistat. After 3 months and up to 6 months following commencement of orlistat treatment, patient's initial body weight must have been reduced by 2.5 kg or 2.5% (whichever is the lesser).******Continuing treatment for patients who have previously been issued with an authority prescription for orlistat. After 6 months and up to 12 months following commencement of orlistat treatment, patient's initial body weight must have been reduced by 5 kg or 5% (whichever is the lesser).*****NOTE:*****The patient should be ideally enrolled in an exercise program and be receiving supplemental vitamins.***

4570M	Capsule 120 mg	84	2	..	126.97	4.60	Xenical	RO

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
VITAMINS								
Vitamin B₁, plain and in combination with vitamin B6 and vitamin B12								
• Vitamin B₁, plain								
THIAMINE HYDROCHLORIDE								
4043T	Tablet 100 mg	100	2	..	9.10	4.60	Betamin	AV
Vitamin B-complex, incl. combinations								
• Vitamin B-complex, plain								
VITAMIN B GROUP COMPLEX								
4493L	Oral liquid 200 mL	‡1	2	..	11.37	4.60	Accomin Adult Tonic	WT
Ascorbic acid (vitamin C), incl. combinations								
• Ascorbic acid (vitamin C), plain								
ASCORBIC ACID								
<u>Authority required</u>								
<i>For management of wound healing.</i>								
4565G	Tablet 250 mg	100	2	..	7.39	4.60	Vitelle Vitamin C	FH
MINERAL SUPPLEMENTS								
Other mineral supplements								
• Magnesium								
MAGNESIUM ASPARTATE								
<u>Restricted benefit</u>								
<i>Patients with documented hypomagnesaemia.</i>								
4321K	Tablet 500 mg	50	..	..	12.37	4.60	Magmin	BB
BLOOD AND BLOOD FORMING ORGANS								
ANTITHROMBOTIC AGENTS								
Antithrombotic agents								
• Platelet aggregation inhibitors excl. heparin								
ASPIRIN								
4076M	Tablet 100 mg (with glycine)	90	1	..	12.18	4.60	Cardiprin 100	RC
4077N	Tablet 100 mg (enteric coated)	84	1	..	11.54	4.60	Cartia	GK
4078P	Capsule 100 mg (containing enteric coated pellets)	84	1	..	12.59	4.60	Astrix	MX
NOTE:								
The enteric coated preparations are for patients with a significant risk of gastrointestinal bleeding.								
CLOPIDOGREL HYDROGEN SULFATE								
<u>Authority required</u>								
<i>For use in patients pre- and post-angioplasty.</i>								
4179Y	Tablet 75 mg (base)	28	3	..	84.13	4.60	Iscover Plavix	BQ SW

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
BLOOD SUBSTITUTES AND PERFUSION SOLUTIONS								
Irrigating solutions								
• Salt solutions								
SODIUM CHLORIDE								
4460R	Irrigation solution 9 mg per mL (0.9%), 500 mL	‡1	2	..	8.49	4.60	BX	
4461T	Irrigation solution 9 mg per mL (0.9%), 1 L	‡1	2	..	8.80	4.60	BX	
CARDIOVASCULAR SYSTEM								
VASOPROTECTIVES								
Antihemorrhoidals for topical use								
• Products containing corticosteroids								
HYDROCORTISONE with CINCHOCAINE HYDROCHLORIDE								
CAUTION:								
Long-term use may lead to skin atrophy.								
4036K	Ointment 5 mg-5 mg per g (0.5%- 0.5%), 30 g	‡1	..	..	17.77	4.60	Proctosedyl	AV
4038M	Suppositories 5 mg-5 mg, 12	‡1	..	..	16.73	4.60	Proctosedyl	AV
• Other antihemorrhoidals for topical use								
ZINC OXIDE								
4039N	Compound ointment 50 g	‡1	1	..	12.42	4.60	Anusol	PC
4040P	Compound suppositories, 12	‡1	1	..	11.38	4.60	Anusol	PC
CALCIUM CHANNEL BLOCKERS								
Selective calcium channel blockers with mainly vascular effects								
• Dihydropyridine derivatives								
NOTE:								
The base-priced drugs in this therapeutic group are felodipine and nifedipine (except nifedipine controlled release tablet 20 mg).								
AMLODIPINE BESYLATE								
Authority required								
<i>To be approved where other base-priced (benchmark) drug treatment is inappropriate.</i>								
4985J	Tablet 5 mg (base)	30	5	..	25.12	4.60	Norvasc	PF
4986K	Tablet 10 mg (base)	30	5	..	39.17	4.60	Norvasc	PF
LERCANIDIPINE HYDROCHLORIDE								
Authority required								
<i>To be approved where other base-priced (benchmark) drug treatment is inappropriate.</i>								
4960C	Tablet 10 mg	30	5	..	26.55	4.60	Zanidip	SM
4959B	Tablet 20 mg	30	5	..	40.57	4.60	Zanidip	SM

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
<u>NIFEDIPINE</u>								
<u>Authority required</u>								
<i>To be approved where other base-priced (benchmark) drug treatment is inappropriate.</i>								
4961D	Tablet 20 mg (controlled release)	30	5	..	24.59	4.60	Adalat Oros 20mg	BN

AGENTS ACTING ON THE RENIN-ANGIOTENSIN SYSTEM

ACE inhibitors, plain• **ACE inhibitors, plain****CAUTION:**

Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

NOTE:

The base-priced drugs in this therapeutic group are captopril, enalapril maleate, fosinopril sodium, lisinopril, perindopril erbumine, quinapril hydrochloride, ramipril (except ramipril capsule 10 mg) and trandolapril.

RAMIPRIL**Authority required**

To be approved where other base-priced (benchmark) drug treatment is inappropriate.

4962E	Capsule 10 mg	30	5	..	34.23	4.60	Tritace 10 mg	AV

DERMATOLOGICALS

ANTIFUNGALS FOR DERMATOLOGICAL USE

Antifungals for topical use• **Antibiotics**

NYSTATIN

4001N	Cream 100,000 units per g, 15 g	‡1	1	..	9.23	4.60	Mycostatin	BQ

• **Imidazole and triazole derivatives**

BIFONAZOLE

4003Q	Cream 10 mg per g (1%), 15 g	‡1	..	..	14.03	4.60	Mycospor	BN

CLOTRIMAZOLE

4004R	Cream 10 mg per g (1%), 20 g	‡1	1	..	7.19	4.60	Clonea	AF

4005T	Lotion 10 mg per mL (1%), 20 mL	‡1	1	..	10.46	4.60	Canesten	BN

KETOCONAZOLE**Restricted benefit**

Severe seborrhoeic dermatitis.

4008Y	Shampoo 20 mg per g (2%), 60 mL	‡1	..	..	17.20	4.60	Nizoral	JC

4007X	Shampoo 20 mg per g (2%), 100 mL	‡1	..	..	17.14	4.60	Sebizole	DP
				..	17.16	4.60	Hexal Konazol 2% Shampoo	HX

MICONAZOLE

4341L	Tincture 20 mg per mL (2%), 30 mL	‡1	1	..	18.41	4.60	Daktarin	JC

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
4009B	MICONAZOLE NITRATE Cream 20 mg per g (2%), 20 g	‡1	1	..	9.32	4.60	Monistat Derm	JC
	• Other antifungals for topical use AMOROLFINE HYDROCHLORIDE <u>Restricted benefit</u> <i>Onychomycosis.</i>							
4010C	Nail treatment kit containing nail lacquer 50 mg (base) per mL (5%), 5 mL, 60 isopropyl alcohol cleaning pads, 10 spatulas and 30 nail files	‡1	1	..	90.99	4.60	Loceryl	GA
	CICLOPIROX OLAMINE <u>Restricted benefit</u> <i>Severe seborrhoeic dermatitis.</i>							
4106D	Shampoo 15 mg per g (1.5%), 60 mL	‡1	..	..	14.45	4.60	Stieprox Liquid	SX
	TERBINAFINE <u>Restricted benefit</u> <i>Tinea pedis.</i>							
4463X	Gel 10 mg per g (1%), 15 g	‡1	..	..	19.29	4.60	Lamisil DermGel	NC
	TERBINAFINE HYDROCHLORIDE <u>Restricted benefit</u> <i>Tinea pedis.</i>							
4473K	Cream 10 mg per g (1%), 15 g	‡1	1	..	19.29	4.60	Lamisil	NC
4481W	TOLNAFTATE Spray aerosol 10 mg per g (1%), 100 g	‡1	..	..	12.65	4.60	Tinaderm	SH
	Antifungals for systemic use • Antifungals for systemic use TERBINAFINE HYDROCHLORIDE <u>Authority required</u> <i>Onychomycosis due to dermatophyte infection proven by microscopy or culture and confirmed by an approved pathology provider.</i>							
4011D	Tablet 250 mg (base)	42	1	..	137.58	4.60	^a Zabel	AF
				..	139.58	4.60	^a Lamisil	NV

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
EMOLLIENTS AND PROTECTIVES								
Emollients and protectives								
• Silicone products								
DIMETHICONE with GLYCEROL								
Restricted benefit								
<i>For colostomy and ileostomy use;</i>								
<i>For use by paraplegic and quadriplegic patients;</i>								
<i>For use with surgical appliances.</i>								
4556T	Cream 150 mg-20 mg per g (15%-2%), 75 g	‡1	..	..	9.57	4.60	Silic 15	EO
4551M	Cream 150 mg-20 mg per g (15%-2%), 500 g	‡1	..	..	20.51	4.60	Silic 15	EO
• Soft paraffin and fat products								
WOOL ALCOHOLS								
4041Q	Ointment 100 g	‡1	1	..	10.86	4.60	Eucerin	BE
• Carbamide products								
UREA								
4042R	Cream 100 mg per g (10%), 100 g	‡1	2	..	9.89	4.60	Urederm	HA
				..	10.16	4.60	Aquacare H.P.	AG
				..	10.25	4.60	Calmurid	OL
							Nutraplus	GA
• Other emollients and protectives								
CARMELLOSE SODIUM with PECTIN and GELATIN								
4518T	Paste 167 mg-167 mg-167 mg per g (16.7%-16.7%- 16.7%), 5 g	‡1	..	..	9.02	4.60	Orabase	BQ
SKIN EMOLLIENT								
4122Y	Bath oil 500 mL	‡1	2	..	15.20	4.60	Alpha Keri Bath Oil	MT
				..	15.93	4.60	QV Bath Oil	EO
				..	17.27	4.60	Hamilton Bath Oil	HA
4107E	Lotion 500 mL	‡1	2	..	15.20	4.60	Alpha Keri Lotion	MT
Protectives against UV-radiation								
• Protectives against UV-radiation for topical use								
SUNSCREENS								
4543D	Solid stick 4.5 g	‡1	2	..	9.95	4.60	Hamilton Solastick 30+	HA
4544E	Cream 100 g	‡1	2	..	13.06	4.60	Hamilton Sunscreen Broad Spectrum Cream 15+	HA
				..	14.61	4.60	SunSense Cream SPF 30+	EO

continued ➡

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
SUNSCREENS—cont.								
4546G	Lotion (non-alcoholic) 125 mL	‡1	2	..	13.06	4.60	Hamilton Broad Spectrum Milky Lotion 15+	HA
				..	13.89	4.60	Aquasun Lotion SPF 18	PF
				..	14.86	4.60	SunSense Ultra SPF 30+	EO

ANTIPRURITICS, INCL. ANTIHISTAMINES, ANESTHETICS, ETC.

Antipruritics, incl. antihistamines, anesthetics, etc.

• **Anesthetics for topical use**

LIGNOCAINE HYDROCHLORIDE with CARBOXYMETHYLCELLULOSE								
4308R	Mucilage 20 mg-25 mg per mL (2%-2.5%), 200 mL	‡1	..	..	67.65	4.60	Xylocaine Viscous	AP

• **Other antipruritics**

PINE TAR with TRIETHANOLAMINE LAURYL SULFATE

NOTE:

For patients who have failed to respond to simple moisturising agents.

4408B	Solution 23 mg-60 mg per mL (2.3%-6%), 500 mL	‡1	2	..	17.41	4.60	Hamilton Pine Tar Solution	HA
				..	17.55	4.60	Pinetarsol	EO

ANTIPSORIATICS

Antipsoriatics for topical use

• **Tars**

ALLANTOIN with SULFUR, PHENOL, COAL TAR SOLUTION and MENTHOL								
4506E	Gel 25 mg-5 mg-5 mg-0.05 mL-7.5 mg per g (2.5%-0.5%-0.5%-5%-0.75%), 75 g	‡1	2	..	16.85	4.60	Egopsoryl-TA	EO

ANTIBIOTICS AND CHEMOTHERAPEUTICS FOR DERMATOLOGICAL USE

Antibiotics for topical use

• **Other antibiotics for topical use**

MUPIROCIN

Restricted benefit

For the topical treatment of secondarily infected traumatic skin lesions.

4348W	Cream 20 mg (as calcium) per g (2%), 15 g	‡1	..	..	14.18	4.60	Bactroban	GK
4350Y	Ointment 20 mg per g (2%), 15 g	‡1	..	..	14.18	4.60	Bactroban	GK

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Chemotherapeutics for topical use								
• Antivirals								
PODOPHYLLOTOXIN								
<u>Authority required</u>								
<i>For the treatment of ano-genital warts.</i>								
4566H	Paint 5 mg per mL (0.5%), 3.5 mL (with 30 swabs)	‡1	..	..	35.24	4.60	Condyline Paint	HA
• Other chemotherapeutics								
METRONIDAZOLE								
4030D	Gel 7.5 mg per g (0.75%), 50 g	‡1	..	..	26.79	4.60	Rozex	GA
4340K	Cream 7.5 mg per g (0.75%), 30 g	‡1	1	..	18.83	4.60	Rozex	GA
CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS								
Corticosteroids, plain								
• Corticosteroids, potent (group III)								
BETAMETHASONE VALERATE								
4511K	Cream 500 micrograms (base) per g (0.05%), 30 g	‡1	2	..	10.33	4.60	Betnovate 1/2	SI
4131K	Cream 1 mg (base) per g (0.1%), 30 g	‡1	2	..	20.06	4.60	Betnovate	SI
4132L	Ointment 1 mg (base) per g (0.1%), 30 g	‡1	2	..	20.06	4.60	Betnovate	SI
4133M	Scalp application 1 mg (base) per g (0.1%), 30 g	‡1	2	..	16.73	4.60	Betnovate	SI
MOMETASONE FUROATE								
4342M	Cream 1 mg per g (0.1%), 45 g	‡1	..	..	24.25	4.60	Elocon	SH
4343N	Ointment 1 mg per g (0.1%), 45 g	‡1	..	..	24.25	4.60	Elocon	SH
<u>NOTE:</u>								
Application to large areas of skin for longer than four weeks is not recommended.								
Corticosteroids, combinations with antibiotics								
• Corticosteroids, moderately potent, combinations with antibiotics								
TRIAMCINOLONE ACETONIDE with NEOMYCIN SULFATE, GRAMICIDIN and NYSTATIN								
4482X	Ointment 1 mg-2.5 mg (base)- 250 micrograms- 100,000 units per g (0.1%-0.25% (base)- 0.025%- 100,000 units in 1 g), 15 g	‡1	..	..	14.71	4.60	Kenacomb	BQ
<u>CAUTION:</u>								
For the short-term treatment of localised infective eczema only.								

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTISEPTICS AND DISINFECTANTS								
Antiseptics and disinfectants								
• Iodine products								
POVIDONE-IODINE								
4411E	Solution 100 mg per mL (10%), 100 mL	‡1	..	..	18.52	4.60	Betadine Antiseptic Liquid	FH
OTHER DERMATOLOGICAL PREPARATIONS								
Other dermatological preparations								
• Antihidrotics								
DIPHEMANIL METHYLSULFATE								
4191N	Dusting powder 20 mg per g (2%), 50 g	‡1	1	..	15.01	4.60	Prantal	SH
• Medicated shampoos								
PINE TAR with CADE OIL, COAL TAR SOLUTION, ARACHIS OIL EXTRACT OF CRUDE COAL TAR and OLEYL ALCOHOL								
4409C	Scalp cleanser 3 mg-3 mg-1 mg- 3 mg-10 mg per mL (0.3%-0.3%- 0.1%-0.3%-1%), 350 mL	‡1	2	..	19.13	4.60	Polytar	SX
SALICYLIC ACID with BENZALKONIUM CHLORIDE, ALCOHOL and POLYOXYETHYLENE ETHERS								
4445Y	Scalp cleanser 20 mg-2 mg- 130 mg-216 mg per mL (2%- 0.2%-13%-21.6%), 250 mL	‡1	2	..	16.19	4.60	Ionil	GA
SALICYLIC ACID with BENZALKONIUM CHLORIDE, ALCOHOL, COAL TAR and POLYOXYETHYLENE ETHERS								
4560B	Scalp cleanser 20 mg-2 mg- 130 mg-50 mg-216 mg per mL (2%-0.2%-13%-5%-21.6%), 250 mL	‡1	2	..	16.73	4.60	Ionil-T	GA
SALICYLIC ACID with COAL TAR SOLUTION, PINE TAR and UNDECYLENAMIDE								
4447C	Scalp cleanser 20 mg-10 mg- 10 mg-10 mg per mL (2%- 1%- 1%-1%), 250 mL	‡1	2	..	15.55	4.60	Sebitar	EO
SELENIUM SULFIDE								
4452H	Shampoo 25 mg per mL (2.5%), 125 mL	‡1	..	..	12.13	4.60	Selsun	AB
ZINC PYRITHIONE								
4498R	Shampoo 10 mg per mL (1%), 200 mL	‡1	1	..	10.29	4.60	Dan Gard	FH
• Wart and anti-corn preparations								
SALICYLIC ACID with PODOPHYLLIN RESIN								
4450F	Paint 100 mg-200 mg per mL (10%-20%), 6 mL	‡1	1	..	13.25	4.60	Posalfilin	NE

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Other dermatologicals								
ALLANTOIN with GLYCEROL and ICHTHAMMOL								
NOTE: For patients who have failed to respond to simple moisturising agents.								
4281H	Cream 5 mg-10 mg-10 mg per g (0.5%-1%-1%), 50 g	‡1	2	..	13.97	4.60	Egoderm Cream	EO
4280G	Ointment 5 mg-10 mg-10 mg per g (0.5%-1%-1%), 50 g	‡1	2	..	13.97	4.60	Egoderm Ointment	EO
CATIONIC CONDITIONER with PANTHENOL								
NOTE: To be used in conjunction with the scalp cleanser salicylic acid with coal tar solution, pine tar and undecylenamide (code 4447C).								
4519W	Solution 250 mL	‡1	2	..	11.56	4.60	SebiRinse Conditioner	EO
HYDROLYZED COLLAGEN PROTEINS								
NOTE: To be used in conjunction with the two scalp cleansers: salicylic acid, benzalkonium chloride, alcohol and polyoxyethylene ethers (code 4445Y); and salicylic acid, benzalkonium chloride, alcohol, coal tar and polyoxyethylene ethers (code 4560B).								
4271T	Hair conditioner 250 mL	‡1	2	..	11.14	4.60	Ionil Rinse	GA
SKIN CLEANSER								
4549K	Lotion 500 mL	‡1	2	..	18.08	4.60	Hamilton Body Wash	HA
ZINC OXIDE with STARCH and CHLORPHENESIN								
4497Q	Dusting powder 100 g	‡1	1	..	10.34	4.60	Z.S.C.	SI
GENITO URINARY SYSTEM AND SEX HORMONES								
GYNECOLOGICAL ANTIINFECTIVES AND ANTISEPTICS								
Antiinfectives and antiseptics, excl. comb. with corticosteroids								
• Antibiotics								
NYSTATIN								
4012E	Cream pessaries 100,000 units, 15	‡1	1	..	11.80	4.60	Nilstat	SI
4013F	Vaginal cream 100,000 units per dose, 15 doses, 75 g	‡1	1	..	11.80	4.60	Nilstat	SI
• Imidazole derivatives								
CLOTRIMAZOLE								
4014G	Pessaries 100 mg, 6	‡1	..	..	13.03 14.75	4.60 4.60	Clofeme Canesten	HX BN
4015H	Pessary 500 mg	1	..	..	13.61 15.88	4.60 4.60	Clofeme Canesten 1	HX BN

continued ➡

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
CLOTRIMAZOLE—cont.								
4016J	Vaginal cream 50 mg per 5 g (1%), 35 g	‡1	..	..	13.03	4.60	Chem mart Clotrimazole 6 Day Cream GenRx Clotrimazole 6 Day Cream healthsense Clotrimazole 6 Day Cream Terry White Chemists Clotrimazole 6 Day Cream	CH GX HS TW
				..	13.14	4.60	Hexal Clofeme 6 Day Cream	HX
				..	14.38	4.60	Canesten	BN
4017K	Vaginal cream 100 mg per 5 g (2%), 20 g	‡1	..	..	13.03	4.60	Chem mart Clotrimazole 3 Day Cream GenRx Clotrimazole 3 Day Cream healthsense Clotrimazole 3 Day Cream Terry White Chemists Clotrimazole 3 Day Cream	CH GX HS TW
				..	13.14	4.60	Hexal Clofeme 3 Day Cream	HX
				..	15.31	4.60	Canesten 3	BN
MICONAZOLE NITRATE								
4020N	Pessaries 100 mg, 7	‡1	..	..	12.81	4.60	Monistat 7	JC
4021P	Vaginal cream 100 mg per 5 g (2%), 40 g	‡1	..	..	12.81	4.60	Monistat 7	JC
OTHER GYNECOLOGICALS								
Other gynecologicals								
• Other gynecologicals								
RICINOLEIC ACID with ACETIC ACID and HYDROXYQUINOLINE SULFATE								
4434J	Vaginal jelly 7 mg-9.4 mg- 250 micrograms per g (0.7%- 0.94%-0.025%), 100 g	‡1	..	..	30.08	4.60	Aci-Jel	JC

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
SEX HORMONES AND MODULATORS OF THE GENITAL SYSTEM								
Estrogens								
• <i>Natural and semisynthetic estrogens, plain</i>								
OESTRADIOL								
<u>Restricted benefit</u>								
<i>Post-menopausal symptoms in women who have failed to respond using oral or topical estrogens.</i>								
4365R	Implant 50 mg	1	..	..	71.36	4.60	OR	
4366T	Implant 100 mg	1	..	..	109.20	4.60	OR	
UROLOGICALS								
Other urologicals, incl. antispasmodics								
• <i>Drugs used in erectile dysfunction</i>								
ALPROSTADIL								
<u>Authority required</u>								
<i>Specific accepted war-caused or service-related disabilities for males with vasculogenic, psychogenic or neurogenic erectile dysfunction.</i>								
<i>Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.</i>								
4579B	Intracavernosal injection 10 micrograms with diluent in single use syringe	6	3	..	*80.95	4.60	Caverject Impulse	PH
4580C	Intracavernosal injection 20 micrograms with diluent in single use syringe	6	3	..	*101.95	4.60	Caverject Impulse	PH
SILDENAFIL CITRATE								
<u>Authority required</u>								
<i>Specific accepted war-caused or service-related disabilities for males with vasculogenic, psychogenic or neurogenic erectile dysfunction.</i>								
<i>Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.</i>								
4584G	Tablet 25 mg (base)	4	5	..	51.10	4.60	Viagra	PF
4585H	Tablet 50 mg (base)	4	5	..	63.75	4.60	Viagra	PF
4586J	Tablet 100 mg (base)	4	5	..	68.56	4.60	Viagra	PF
TADALAFIL								
<u>Authority required</u>								
<i>Specific accepted war-caused or service-related disabilities for males with vasculogenic, psychogenic or neurogenic erectile dysfunction.</i>								
<i>Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.</i>								
4596X	Tablet 10 mg	4	5	..	66.71	4.60	Cialis	LY
4597Y	Tablet 20 mg	4	5	..	71.75	4.60	Cialis	LY

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Other urologicals								
4458P	SODIUM BICARBONATE Capsule 840 mg	100	2	..	12.00	4.60	Sodibic	AS
CAUTION: For use in the treatment of renal disease.								
SODIUM CITRO-TARTRATE								
Restricted benefit <i>For relief of urinary symptoms when antibiotic or other therapy alone is inappropriate.</i>								
4047B	Sachets containing oral effervescent powder 3.7 g, 28	‡1	4	..	11.48	4.60	Citralite	MM
4048C	Sachets containing oral effervescent powder 4 g, 28	‡1	4	..	11.48	4.60	Citravescent Sachets	MM
4049D	Sachets containing oral effervescent powder 4 g, 28	‡1	4	..	11.57	4.60	Ural Sachets	SI
Drugs used in benign prostatic hypertrophy								
• Alpha-adrenoreceptor antagonists								
TAMSULOSIN HYDROCHLORIDE								
Authority required <i>Treatment of benign prostatic hyperplasia where surgery is inappropriate, or where other drug treatment has failed or is contraindicated.</i>								
4464Y	Capsule 400 micrograms (modified release)	30	5	..	58.13	4.60	Flomax	CS
TERAZOSIN HYDROCHLORIDE								
Authority required <i>Treatment of benign prostatic hyperplasia where surgery is inappropriate, or where other drug treatment has failed or is contraindicated.</i>								
4396J	Starter pack containing 7 tablets 1 mg and 7 tablets 2 mg	‡1	..	..	17.79	4.60	Hytrin	AB
4397K	Tablet 2 mg	28	5	..	38.60	4.60	Hytrin	AB
4398L	Tablet 5 mg	28	5	..	56.52	4.60	Hytrin	AB
4399M	Tablet 10 mg	28	5	..	84.39	4.60	Hytrin	AB
• Testosterone-5-alpha reductase inhibitors								
FINASTERIDE								
Authority required <i>Treatment of benign prostatic hyperplasia where surgery is inappropriate, or where other drug treatment has failed or is contraindicated.</i>								
4233T	Tablet 5 mg	30	5	..	97.10	4.60	Proscar	MK

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTIINFECTIVES FOR SYSTEMIC USE								
ANTIBACTERIALS FOR SYSTEMIC USE								
Macrolides, lincosamides and streptogramins								
• Macrolides								
AZITHROMYCIN								
<u>Restricted benefit</u>								
<i>Upper and lower respiratory tract infections.</i>								
4115N	Tablet 500 mg	3	..	..	32.18	4.60	Zithromax	PF
Quinolone antibacterials								
• Fluoroquinolones								
GATIFLOXACIN								
<u>Authority required</u>								
<i>For treatment, where other therapies have failed or are inappropriate, of:</i>								
<i>Community-acquired pneumonia;</i>								
<i>Acute bacterial exacerbations of chronic bronchitis;</i>								
<i>Acute bacterial sinusitis.</i>								
4297E	Tablet 400 mg	7	1	..	79.35	4.60	Tequin	BQ
4298F	Solution concentrate for I.V. infusion 400 mg in 40 mL	3	..	..	*244.72	4.60	Tequin	BQ
4299G	I.V. infusion 400 mg in 200 mL	3	..	..	*244.72	4.60	Tequin	BQ
MOXIFLOXACIN HYDROCHLORIDE								
<u>Authority required</u>								
<i>For treatment, where other therapies have failed or are inappropriate, of:</i>								
<i>Community-acquired pneumonia;</i>								
<i>Acute bacterial exacerbations of chronic bronchitis;</i>								
<i>Acute bacterial sinusitis.</i>								
4329W	Tablet 400 mg (base)	5	1	..	46.62	4.60	Avelox	BN
4330X	Solution for I.V. infusion 400 mg (base) in 250 mL	3	..	..	*199.15	4.60	Avelox	BN
ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS								
ANTINEOPLASTIC AGENTS								
Antimetabolites								
• Pyrimidine analogues								
FLUOROURACIL								
4222F	Cream 50 mg per g (5%), 20 g	‡1	..	..	35.72	4.60	Efudix	VT

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. No. of			Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
		Qty	Rpts	Premium			

IMMUNOSUPPRESSIVE AGENTS

Immunosuppressive agents

• *Selective immunosuppressive agents***ALEFACEPT****NOTE:**

Any queries concerning the arrangements to prescribe alefacept may be directed to the Veterans' Affairs Pharmaceutical Approvals Centre (VAPAC) on 1800 552 580.

Written applications for authority to prescribe alefacept should be forwarded to:

Reply Paid 372

Veterans' Affairs Pharmaceutical Approvals Centre (VAPAC)

GPO Box 9998

BRISBANE QLD 4001

No telephone approvals will be granted for initial or continuing authority applications.

No application for increased maximum quantities and/or repeats will be authorised. A maximum of 2 courses of either intravenous or intramuscular alefacept (Amevive) treatment will be authorised in a 12-month period.

The assessment of the patient's response to the initial course of treatment should be made up to 24 weeks following commencement of the treatment course so that there is adequate time for a response to be demonstrated. Measurement of response should be made prior to the completion of the 24-week follow up to ensure continuity for those patients who meet the criteria. Use of alefacept in combination with other systemic therapies must be in line with the Therapeutic Goods Administration (TGA)-approved Product Information.

Authority required

Initial treatment by a dermatologist for patients who have severe chronic plaque psoriasis, in whom other systemic therapies are ineffective or inappropriate, with the following criteria:

(1) CD4+ lymphocyte counts above the lower limit of normal within 2 weeks prior to dosing;

AND

(2) Baseline Psoriasis Area and Severity Index (PASI) score of 15 or more (the date of PASI assessment must be provided);

AND

(3) who have signed a patient agreement form indicating that they understand and acknowledge that RPBS-subsidised treatment will cease if the predetermined response criteria do not support continuation of RPBS-subsidised treatment.

If the above requirements cannot be met, the application must state the reasons why these criteria cannot be met.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form;

(2) a completed Alefacept (Amevive) RPBS Authority Application - Supporting Information Form (contact the VAPAC on 1800 552 580 for a copy of this form);

and

(3) a signed patient agreement form.

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ALEFACEPT—cont.								
Authority required								
<i>Initial RPBS-subsidised supply for continuing treatment by a dermatologist for patients who have severe chronic plaque psoriasis and were receiving treatment with alefacept prior to July 2004, who:</i>								
<i>(1) have demonstrated a response as specified in the criteria for continuing RPBS-subsidised treatment with alefacept (the date of assessment of the patient must be provided); and</i>								
<i>(2) have signed a patient agreement form indicating that they understand and acknowledge that RPBS-subsidised treatment will cease if the predetermined response criteria do not support continuation of RPBS-subsidised treatment.</i>								
<i>Applications for authorisation must be in writing and must include:</i>								
<i>(1) a completed authority prescription form;</i>								
<i>(2) a completed Alefacept (Amevive) RPBS Authority Application - Supporting Information Form (contact the VAPAC on 1800 552 580 for a copy of this form); and</i>								
<i>(3) a signed patient agreement form.</i>								
Authority required								
<i>Continuing RPBS-subsidised treatment by a dermatologist for patients who have severe chronic plaque psoriasis and who:</i>								
<i>(1) at the time of application, have demonstrated a reduction in PASI score of 75% or more from baseline at any time up to 24 weeks following commencement of initial treatment. Re-treatment may be initiated as needed, provided that CD4+ T-cell count is above 250 cells per microlitre and a minimum of 12 weeks has passed between courses of treatment;</i>								
<i>OR</i>								
<i>(2) at the time of application, have demonstrated a reduction in PASI score of from at least 50% to up to 75% from baseline at any time up to 24 weeks following initiation of treatment. These patients may receive a second course of alefacept, provided that their CD4+ T-cell count is within the normal range and a minimum of 12 weeks has passed between courses of treatment. Patients must demonstrate a reduction of 75% or more from baseline PASI score at any time up to 24 weeks following the second course of treatment to be eligible to continue RPBS-subsidised treatment.</i>								
<i>Patients who fail to demonstrate an adequate response, as specified in the criteria for continuing treatment with alefacept, will not be eligible to recommence treatment. Patients unable to complete the course of treatment for reasons other than safety will be eligible to recommence treatment with alefacept within 12 months of the date on which treatment was ceased. Where re-treatment with alefacept after a break in RPBS-subsidised treatment with the drug is sought, the reason for and date of cessation of the previous treatment course with alefacept must be included in the application.</i>								
<i>Applications for authorisation must be in writing and must include:</i>								
<i>(1) a completed authority prescription form; and</i>								
<i>(2) a completed Alefacept (Amevive) RPBS Authority Application - Supporting Information Form (contact the VAPAC on 1800 552 580 for a copy of this form).</i>								
4534P	Pack containing 4 vials powder for I.V. injection 7.5 mg with 4 vials solvent	1	2	..	5998.16	4.60	Amevive	BD
4535Q	Pack containing 4 vials powder for I.M. injection 15 mg with 4 vials solvent	1	2	..	5998.16	4.60	Amevive	BD

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
INFLIXIMAB							
NOTE:							
<i>Any queries concerning the arrangements to prescribe infliximab may be directed to the Veterans' Affairs Pharmaceutical Approvals Centre (VAPAC) on 1800 552 580.</i>							
<i>Written applications for authority to prescribe infliximab should be forwarded to:</i>							
<i>Reply Paid 372 Veterans' Affairs Pharmaceutical Approvals Centre (VAPAC) GPO Box 9998 BRISBANE QLD 4001</i>							
Authority required							
<i>Initial treatment, in combination with methotrexate, of specific accepted war-caused or service-related disability of refractory rheumatoid arthritis. Initial treatment may be prescribed by rheumatologists or consultant physicians for the reduction of signs and symptoms and prevention of structural joint damage in adult patients with active rheumatoid arthritis who satisfy all of the following criteria:</i>							
<i>(1) (a) Proven raised erythrocyte sedimentation rate (ESR) and/or C-reactive protein (CRP); and</i>							
<i>(1) (b) Proven erosive rheumatoid arthritis without end-stage disease;</i>							
<i>(2) Failure of an adequate trial of methotrexate and 2 other disease modifying anti-rheumatic drugs (such as sulfasalazine, hydroxychloroquine, leflunomide or cyclosporin) — unless these drugs were contraindicated or intolerance had developed;</i>							
<i>(3) No history of active tuberculosis requiring treatment in the last 3 years;</i>							
<i>(4) No history of opportunistic infection in the last 2 months;</i>							
<i>(5) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.</i>							
<i>Applications for authorisation must be in writing and must include:</i>							
<i>(1) a completed authority prescription form; and</i>							
<i>(2) a completed Infliximab (Remicade) RPBS Authority Application - Supporting Information form (contact the VAPAC on 1800 552 580 for a copy of the form).</i>							
Authority required							
<i>Continuing treatment, in combination with methotrexate, of specific accepted war-caused or service-related disability of refractory rheumatoid arthritis. Continuing treatment may be prescribed by rheumatologists or consultant physicians, following initial therapy of 3 doses, in patients who satisfy the following criteria:</i>							
<i>(1) There is improvement in ESR and/or CRP; and</i>							
<i>(2) An ACR20 (American College of Rheumatology) response is achieved by 14 weeks after the commencement of therapy.</i>							
<i>Applications for authorisation must be in writing and must include:</i>							
<i>(1) a completed authority prescription form; and</i>							
<i>(2) a completed Infliximab (Remicade) RPBS Authority Application - Supporting Information form (contact the VAPAC on 1800 552 580 for a copy of the form).</i>							
4284L	Powder for I.V. infusion 100 mg	1	2	..	1015.86	4.60	Remicade SH

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
MUSCULO-SKELETAL SYSTEM								
ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS								
Antiinflammatory and antirheumatic products, non-steroids								
• Acetic acid derivatives and related substances								
DICLOFENAC SODIUM with MISOPROSTOL								
Authority required								
<i>Patients requiring an NSAID in whom a risk of upper gastrointestinal complications is high or with a history of peptic ulcer disease.</i>								
4190M	Tablet 50 mg-200 micrograms	60	2	..	34.75	4.60	Arthrotec 50	PH
TOPICAL PRODUCTS FOR JOINT AND MUSCULAR PAIN								
Topical products for joint and muscular pain								
• Preparations with salicylic acid derivatives								
METHYL SALICYLATE								
4022Q	Compound cream APF, 100 g	‡1	1	..	11.93	4.60	Gold Cross	BI
4023R	Ointment BP, 100 g	‡1	1	..	10.18	4.60	Gold Cross	BI
4025W	Compound ointment APF 1934, 100 g	‡1	1	..	8.95	4.60	Gold Cross	BI
4026X	Liniment APF, 100 mL	‡1	1	..	7.92	4.60	Gold Cross	BI
4027Y	Compound liniment APF, 100 mL	‡1	1	..	9.62	4.60	Gold Cross	BI
DRUGS FOR TREATMENT OF BONE DISEASES								
Drugs affecting bone structure and mineralization								
• Bisphosphonates								
RISEDRONATE SODIUM								
Authority required								
<i>For preservation of bone mineral density in patients on long-term glucocorticoid therapy where patients are undergoing continuous treatment with a dose equal to or greater than 7.5 mg of prednisone or equivalent per day. Prescribers need to demonstrate that the patient has been on continuous therapy for 3 months or more and demonstrate that the patient is osteopenic (bone mineral density t-score of less than -1.0).</i>								
4443W	Tablet 5 mg	28	5	..	54.46	4.60	Actonel	AV
4444X	Tablet 35 mg	4	5	..	54.46	4.60	Actonel Once-a-Week	AV

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
NERVOUS SYSTEM								
ANALGESICS								
Opioids								
• Natural opium alkaloids								
MORPHINE SULFATE								
CAUTION:								
<i>The risk of drug dependence is high.</i>								
Restricted benefit								
<i>Chronic severe disabling pain not responding to non-narcotic analgesics.</i>								
NOTE:								
<i>Authorities for increased maximum quantities and/or repeats will be granted only for</i>								
<i>(i) chronic severe disabling pain associated with proven malignant neoplasia; or</i>								
<i>(ii) chronic severe disabling pain where treatment has been initiated by a specialist with</i>								
<i>appropriate expertise in pain management.</i>								
4349X	Tablet 200 mg (controlled release)	20	..	..	106.36	4.60	MS Contin	MF
• Diphenylpropylamine derivatives								
DEXTROPROPOXYPHENE NAPSYLATE								
CAUTION:								
<i>Chronic use of this preparation is likely to cause drug dependence.</i>								
4081T	Capsule 100 mg	50	..	..	*16.40	4.60	Doloxene	AS
Other analgesics and antipyretics								
• Salicylic acid and derivatives								
CODEINE PHOSPHATE with ASPIRIN								
4061R	Tablet soluble 8 mg-300 mg	50	2	..	11.59	4.60	Aspalgin	FM
• Anilides								
CODEINE PHOSPHATE with PARACETAMOL								
4171M	Tablet 8 mg-500 mg	50	2	..	9.84	4.60	Panamax Co.	SW
				..	10.91	4.60	Codalgin	FM
4170L	Tablet 15 mg-500 mg	20	2	..	7.43	4.60	Prodeine 15	SW
• Other analgesics and antipyretics								
GABAPENTIN								
Authority required								
<i>To be approved for the treatment of refractory neuropathic pain not controlled by other drugs.</i>								
4591P	Capsule 100 mg	100	5	..	32.66	4.60	^a Gantin	AW
							^a Neurontin	PF
							^a Nupentin 100	AF

continued ➡

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer
GABAPENTIN—cont.							
4592Q	Capsule 300 mg	100	5	..	86.34	4.60	^a DBL MX ^a <i>Gabapentin</i> ^a Douglas DP <i>Gabapentin</i> 300mg ^a Gantin AW ^a GenRx GX <i>Gabapentin</i> ^a Neurontin PF ^a Nupentin 300 AF
4593R	Capsule 400 mg	100	5	..	114.22	4.60	^a DBL MX ^a <i>Gabapentin</i> ^a Douglas DP <i>Gabapentin</i> 400mg ^a Gantin AW ^a GenRx GX <i>Gabapentin</i> ^a Neurontin PF ^a Nupentin 400 AF
4594T	Tablet 600 mg	100	5	..	180.78	4.60	Neurontin PF
4595W	Tablet 800 mg	100	5	..	228.71	4.60	^a Gantin AW ^a Neurontin PF

PSYCHOLEPTICS

Anxiolytics

• Benzodiazepine derivatives

BROMAZEPAMAuthority required*Patients with terminal disease;**Patients with refractory phobic or anxiety states.*NOTE:

For short-term use and palliative care. This drug should not be used as the first line of treatment. Other PBS-listed benzodiazepines should have been adequately tried and found to be ineffective or inappropriate. Authorities for increased quantities and/or repeats may be granted to patients with terminal disease, and other patients who have been shown to be dependent on this item by an unsuccessful attempt at gradual withdrawal.

4150K	Tablet 3 mg	60	..	..	*24.79	4.60	Lexotan RO
4151L	Tablet 6 mg	60	..	..	*30.55	4.60	Lexotan RO

• Azaspirodecanedione derivatives

BUSPIRONE HYDROCHLORIDEAuthority required*For the short-term treatment of anxiety.*

4144D	Tablet 5 mg	50	..	..	29.56	4.60	Buspar BQ
4145E	Tablet 10 mg	50	..	..	44.44	4.60	Buspar BQ

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Hypnotics and sedatives								
• Benzodiazepine derivatives								
FLUNITRAZEPAM								
<u>Authority required</u>								
<i>Patients with terminal disease;</i>								
<i>Patients with refractory phobic or anxiety states.</i>								
<u>NOTE:</u>								
<i>For short-term use and palliative care. This drug should not be used as the first line of treatment. Other PBS-listed benzodiazepines should have been adequately tried and found to be ineffective or inappropriate. Authorities for increased quantities and/or repeats may be granted to patients with terminal disease, and other patients who have been shown to be dependent on this item by an unsuccessful attempt at gradual withdrawal.</i>								
4216X	Tablet 1 mg	30	..	..	11.55	4.60	Hypnodorm	AF
• Benzodiazepine related drugs								
ZOPICLONE								
<u>Restricted benefit</u>								
<i>For the short-term treatment of insomnia.</i>								
4522B	Tablet 7.5 mg	30	..	..	20.93	4.60	Imovane	AV
OTHER NERVOUS SYSTEM DRUGS								
Drugs used in addictive disorders								
• Drugs used in nicotine dependence								
NICOTINE								
<u>Authority required</u>								
<i>Patients who have indicated that they are ready to cease smoking and who have entered a support and counselling program.</i>								
<u>NOTE:</u>								
<i>Studies have shown that successful therapy with this drug is enhanced by patient participation in a support and counselling program.</i>								
4576W	Transdermal patches releasing approximately 5 mg per 16 hours, 7	2	..	..	*48.65	4.60	Nicorette Patch	PH
4571N	Transdermal patches releasing approximately 7 mg per 24 hours, 7	2	..	..	*43.55 *72.65	4.60 4.60	QuitX Nicabate CQ 7	AF GK
4577X	Transdermal patches releasing approximately 10 mg per 16 hours, 7	2	..	..	*52.99	4.60	Nicorette Patch	PH
4572P	Transdermal patches releasing approximately 14 mg per 24 hours, 7	2	..	..	*47.21 *72.65	4.60 4.60	QuitX Nicabate CQ 14	AF GK
4578Y	Transdermal patches releasing approximately 15 mg per 16 hours, 7	2	2	..	*58.29	4.60	Nicorette Patch	PH
4573Q	Transdermal patches releasing approximately 21 mg per 24 hours, 7	2	2	..	*51.61 *72.65	4.60 4.60	QuitX Nicabate CQ 21	AF GK

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS								
ANTHELMINTICS								
Antinematodal agents								
• Benzimidazole derivatives								
	MEBENDAZOLE							
4325P	Tablet 100 mg	6	..	..	17.79	4.60	Vermox	JC
RESPIRATORY SYSTEM								
NASAL PREPARATIONS								
Decongestants and other nasal preparations for topical use								
• Sympathomimetics, plain								
	OXYMETAZOLINE HYDROCHLORIDE							
4377J	Nasal drops 500 micrograms per mL (0.05%), 15 mL	‡1	..	..	14.52	4.60	Drixine	SH
4378K	Nasal spray 500 micrograms per mL (0.05%), 15 mL	‡1	..	..	14.52	4.60	Drixine	SH
4379L	Nasal spray 500 micrograms per mL (0.05%), 18 mL	‡1	..	..	14.64	4.60	Logicin Rapid Relief	SI
• Antiallergic agents, excl. corticosteroids								
	LEVOCABASTINE HYDROCHLORIDE							
4311X	Nasal spray 500 micrograms per mL (0.05%), 10 mL (100 doses)	‡1	2	..	16.06	4.60	Livostin	JC
	SODIUM CROMOGLYCATE							
4468E	Nasal spray metered dose pump 20 mg per mL (2%), 26 mL	‡1	5	..	18.08	4.60	Rynacrom	AV
• Corticosteroids								
	BUDESONIDE							
	Restricted benefit							
	➤ Severe intractable rhinitis.							
4092J	Aqueous nasal spray (pump pack) 64 micrograms per dose (120 doses)	‡1	..	..	25.63	4.60	Budamax Aqueous	PM
• Other nasal preparations								
	IPRATROPIUM BROMIDE							
	Restricted benefit							
	Severe intractable rhinorrhoea, associated with perennial rhinitis, unresponsive to insufflated nasal steroids.							
4089F	Aqueous nasal spray (pump pack) 21 micrograms (anhydrous) per dose (180 doses)	‡1	5	..	19.09	4.60	Atrovent Nasal Aqueous	BY
4090G	Aqueous nasal spray (pump pack) 42 micrograms (anhydrous) per dose (180 doses)	‡1	5	..	24.83	4.60	Atrovent Nasal Forte	BY

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
Nasal decongestants for systemic use								
• Sympathomimetics								
4420P	PSEUDOEPHEDRINE HYDROCHLORIDE Tablet 60 mg	30	..	..	14.01	4.60	Logicin Sinus Sudafed Sinus & Nasal Decongestant	SI PC
DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES								
Adrenergics, inhalants								
• Adrenergics and other drugs for obstructive airway diseases								
IPRATROPIUM BROMIDE with SALBUTAMOL SULFATE								
Restricted benefit								
<i>Chronic obstructive pulmonary disease where treatment with a beta-agonist and ipratropium is indicated.</i>								
4283K	Oral pressurised inhalation 20 micrograms (anhydrous)- 100 micrograms (base) per dose (200 doses)	2	5	..	*47.15	4.60	Combivent	BY
COUGH AND COLD PREPARATIONS								
Expectorants, excl. combinations with cough suppressants								
• Expectorants								
4074K	SENEGA and AMMONIA Mixture 200 mL	‡1	4	..	7.31	4.60	Gold Cross	BI
Cough suppressants, excl. combinations with expectorants								
• Opium alkaloids and derivatives								
4071G	PHOLCODINE Linctus 1 mg per mL (0.1%), 100 mL	‡1	2	..	7.16 11.25	4.60 4.60	Gold Cross Duro-Tuss	BI MM
ANTI-HISTAMINES FOR SYSTEMIC USE								
Antihistamines for systemic use								
• Phenothiazine derivatives								
4072H	PROMETHAZINE HYDROCHLORIDE Tablet 10 mg	50	2	..	12.90	4.60	Phenergan	AV
4073J	Tablet 25 mg	50	2	..	14.97	4.60	Phenergan	AV
CAUTION: Significant side effects may occur.								
• Piperazine derivatives								
4175R	CETIRIZINE HYDROCHLORIDE Tablet 10 mg	30	..	..	36.34	4.60	Zyrtec	PC

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
• Other antihistamines for systemic use								
FEXOFENADINE HYDROCHLORIDE								
4237B	Tablet 60 mg	60	..	..	*45.73	4.60	Telfast	AV
4238C	Tablet 120 mg	30	..	..	31.81	4.60	^a Xergic	AF
				..	38.41	4.60	^a Telfast 120	AV
LORATADINE								
4313B	Tablet 10 mg	30	..	..	41.95	4.60	Claratyne	SH

SENSORY ORGANS

OPHTHALMOLOGICALS

Antiglaucoma preparations and miotics

• Prostaglandin analogues

LATANOPROST with TIMOLOL MALEATE

Restricted benefit

For the reduction of elevated intra-ocular pressure in patients with open-angle glaucoma who are insufficiently responsive to beta-blockers, prostaglandins and other intra-ocular pressure lowering medications, or where combination therapy is required to aid compliance.

4300H	Eye drops 50 micrograms-5 mg (base) per mL (0.005%-0.5%), 2.5 mL	₹1	5	..	44.79	4.60	Xalacom	PU

Decongestants and antiallergics

• Sympathomimetics used as decongestants

ANTAZOLINE with NAPHAZOLINE

4031E	Eye drops 5 mg (sulfate)-250 micrograms (nitrate) per mL (0.5%-0.025%), 10 mL	₹1	1	..	11.87	4.60	Antistine-Privine	NV

4032F	Eye drops 5 mg (phosphate)-500 micrograms (hydrochloride) per mL (0.5%-0.05%), 15 mL	₹1	1	..	11.82	4.60	Albalon-A	AG

NAPHAZOLINE HYDROCHLORIDE

4035J	Eye drops 1 mg per mL (0.1%), 15 mL	₹1	1	..	11.06	4.60	Naphcon Forte	AQ
				..	11.88	4.60	Albalon Liquifilm	AG

NAPHAZOLINE HYDROCHLORIDE with PHENIRAMINE MALEATE

4355F	Eye drops 250 micrograms-3 mg per mL (0.025%-0.3%), 15 mL	₹1	1	..	12.13	4.60	Naphcon-A	AQ

ZINC SULFATE with PHENYLEPHRINE HYDROCHLORIDE

4034H	Eye drops 2.5 mg-1.2 mg per mL (0.25%-0.12%), 15 mL	₹1	5	..	11.06	4.60	Zincfrin	AQ

• Other antiallergics

LEVOCABASTINE HYDROCHLORIDE

4310W	Eye drops 500 micrograms per mL (0.05%), 4 mL (120 doses)	₹1	1	..	16.06	4.60	Livostin	JC

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
OTOLOGICALS								
Corticosteroids and antiinfectives in combination								
• Corticosteroids and antiinfectives in combination								
CIPROFLOXACIN HYDROCHLORIDE with HYDROCORTISONE								
Authority required								
<i>Indicated where first-line treatment has not been successful or is inappropriate.</i>								
4528H	Ear drops 2 mg (base)-10 mg per mL (0.2%-1%), 10 mL	‡1	2	..	27.11	4.60	Ciproxin HC	AQ
Other otologicals								
• Indifferent preparations								
CARBAMIDE PEROXIDE								
4176T	Ear drops 65 mg per mL (6.5%), 12 mL	‡1	..	..	12.38	4.60	Ear Clear for Ear Wax Removal	KY
DICHLOROBENZENE with CHLORBUTOL and TURPENTINE OIL								
4180B	Ear drops 20 mg-50 mg-0.1 mL per mL (2%-5%-10%), 11 mL	‡1	..	..	12.08	4.60	Cerumol	AC
DOCUSATE SODIUM								
4199B	Ear drops 5 mg per mL (0.5%), 10 mL	‡1	..	..	12.45	4.60	Waxsol	NE
VARIOUS								
ALL OTHER THERAPEUTIC PRODUCTS								
All other therapeutic products								
• Drugs for treatment of hyperkalemia and hyperphosphatemia								
SODIUM POLYSTYRENE SULFONATE								
4470G	Oral powder 454 g	‡1	2	..	63.89	4.60	Resonium-A	SW

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. No. of Qty Rpts Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer

ALL OTHER NON-THERAPEUTIC PRODUCTS

All other non-therapeutic products

- *Other non-therapeutic auxiliary products*

REPATRIATION PHARMACEUTICAL BENEFITS SCHEME (RPBS) WOUND ASSESSMENT AND DRESSING IDENTIFICATION

It is essential to define the aetiology of the wound before selecting a dressing. Recommendations are based on wound type, colour of wound base, depth of wound, and amount of exudate.

This wound chart adheres to the MOIST WOUND concept of healing and wound dressings are described below as ABSORBING or MOISTURE DONATING.

Most wound healing products are designed to remain in situ for several days, with the exception of those for infected wounds which should be changed daily. The quantities and repeats listed in the Repatriation Schedule are considered to be adequate to manage the treatment of a wound for two weeks to one month, when an assessment of the wound's healing process should be undertaken.

DRESSINGS

PINK EPITHELIALISING WOUND

Aim: To protect and promote epithelialisation. Epithelialising wounds normally are superficial and only produce a light exudate.

(A) Covering	• Film; • Film Island;	• Gauze—Paraffin; • Non-adherent
(B) Absorbing	• Foam (Light Exudate); • Hydroactive (Superficial Wound—Light Exudate);	• Hydrocolloid (Superficial Wound—Light Exudate)

RED GRANULATING WOUND

Aims: (1) to protect the granulating tissue; (2) to encourage epithelialisation; (3) to absorb excess exudate.

LIGHT EXUDATE:	Superficial	Cavity
(A) Absorbing	• Foam (Light Exudate); • Hydroactive (Superficial Wound—Light Exudate); • Hydrocolloid (Superficial Wound—Light Exudate)	• Hydrocolloid (Cavity Wound)
(B) Moisture donating	• Hydrogel—Amorphous • Hydrogel—Sheet	• Hydrogel—Amorphous

HIGH EXUDATE:	Superficial	Cavity
(A) Absorbing	• Alginate (Superficial Wound); • Foam—Heavy Exudate; • Hydroactive (Superficial Wound—Moderate Exudate); • Hydrocolloid (Superficial Wound—Moderate/High Exudate)	• Alginate (Cavity Wound); • Foam—Moderate Exudate (see “cavity conforming” product); • Hydroactive (Cavity Wound); • Hydrocolloid (Cavity Wound)
(B) Moisture donating	NOT APPROPRIATE	

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. No.of Qty Rpts Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer

YELLOW SLOUGHY WOUND

Aims: (1) to remove slough; (2) to encourage granulation; (3) to absorb excess exudate.

LIGHT EXUDATE:	Superficial	Cavity
(A) Absorbing	• Cadexomer Iodine • Foam—Light Exudate; • Foam with Charcoal; • Hydroactive (Superficial Wound—Moderate Exudate); • Hydrocolloid (Superficial Wound—Moderate Exudate)	• Cadexomer Iodine • Hydrocolloid (Cavity Wound)
(B) Moisture Donating	• Hydrogel—Amorphous; • Hydrogel—Sheet	• Hydrogel—Amorphous

HIGH EXUDATE:	Superficial	Cavity
(A) Absorbing	• Alginate (Superficial Wound); • Cadexomer Iodine • Foam—Heavy Exudate; • Hydroactive Superficial Wound—Moderate/High Exudate); • Hydrocolloid (Superficial Wound—Moderate/High Exudate)	• Alginate (Cavity Wound); • Cadexomer Iodine • Hydrocolloid (Cavity Wound)
(B) Moisture donating	NOT APPROPRIATE	

BLACK NECROTIC WOUND

Aim: To remove eschar by — (1) sharp debridement, e.g., scissor/scalpel and/or (2) rehydration and autolytic debridement. (These wounds usually produce a LIGHT EXUDATE.)

DRY / LIGHT EXUDATE:	Superficial	Cavity
(A) Absorbing	• Hydroactive Superficial Wound—Light Exudate); • Hydrocolloid (Superficial Wound—Light/Moderate Exudate)	• Hydrocolloid (Cavity Wound)
(B) Moisture donating	• Hydrogel—Amorphous: • Hydrogel—Sheet	• Hydrogel—Amorphous: • Hydrogel—Sheet

INFECTED WOUNDS

Aims: (1) to clear the infection with systemic antibiotics; (2) to absorb excess exudate; (3) to remove slough if present.

MALODOROUS WOUNDS

Aims: (1) to clear infection if present; (2) to remove slough if present; (3) to clear colonising odour-producing bacteria **in** slough — by applying metronidazole gel; (4) to absorb excess exudate.

Products: Activated Charcoal; Alginate with Charcoal; Foam with Charcoal.

MINOR SKIN TRAUMA

Aims: (1) to stop bleeding; (2) to prevent infection; (3) to minimise the surface defect; (4) to promote epithelialisation.

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
ORDERING HARTMANN PRODUCTS								
Hartmann wound dressings are available through the Hartmann Alliance only. To order please telephone the Hartmann Alliance distributor in your State/Territory as below.								
	NSW & ACT	Paul Hartmann Pty Ltd				1800 805 839		
	VIC & TAS	Sutherland Medical				1300 664 027		
	QLD	Medical and Surgical Requisites Pty Ltd				1800 801 226		
	WA	Perth Surgical Supply Co				(08) 9344 3111		
	SA	McNeil's Surgical				(08) 8363 0888		
	NT	McNeil's Surgical				(08) 8948 4047		
BANDAGE—ABSORBENT WOOL								
4651T	Bandage (natural, non-sterile) 10 cm x 2.7 m	6	..	..	*31.57	4.60	Soffban 7224	BV
4653X	Bandage 10 cm x 3 m	6	..	..	18.05	4.60	Surepress 650948	CC
BANDAGE—CALICO								
4717G	Bandage, triangular, large	‡1	..	..	11.52	4.60	Handy 5608	BV
BANDAGE—COMPRESSION								
NOTE: Treatment of varices and oedema associated with venous disease and lymphoedema; contraindicated in arterial disease.								
4654Y	Bandage, short stretch, 8 cm x 5 m	5	..	..	*69.05	4.60	Comprilan 1027	BV
4736G	Bandage, high stretch, 7.5 cm x 3 m	5	..	..	*82.45	4.60	Tensopress 66004347	BV
4656C	Bandage, high stretch, 7.5 cm x 3.5 m	5	..	..	*66.70	4.60	Setopress 3504	SS
4748X	Bandage, high stretch, 10 cm x 3 m	5	..	..	*71.25	4.60	Surepress 650947	CC
				..	*107.60	4.60	Tensopress 66004348	BV
4657D	Bandage, high stretch, 10 cm x 3.5 m	5	..	..	*54.00	4.60	Eloflex 2480	BV
				..	*76.90	4.60	Setopress 3505	SS
4658E	Bandage, four layer	5	..	..	*196.30	4.60	Profore 66050016	SN
4598B	Bandage, four layer	5	..	..	*132.10	4.60	Profore Lite 66050415	SN

continued ➡

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
BANDAGE—COMPRESSION—cont.								
NOTE: <i>Treatment of varices and oedema associated with venous disease and lymphoedema; contraindicated in arterial disease.</i>								
Restricted benefit <i>Initial treatment of venous ulcers.</i>								
4938X	Bandage, two layer, 18 cm- 22 cm (red)	1	..	..	41.97	4.60	ProGuide 66000780	SN
4939Y	Bandage, two layer, 22 cm- 28 cm (yellow)	1	..	..	41.97	4.60	ProGuide 66000781	SN
4940B	Bandage, two layer, 28 cm- 32 cm (green)	1	..	..	41.97	4.60	ProGuide 66000782	SN
Restricted benefit <i>Continuation of treatment of venous ulcers where patient's ability to tolerate dressing has been demonstrated.</i>								
4941C	Bandage, two layer, 18 cm- 22 cm (red)	4	..	..	*153.63	4.60	ProGuide 66000780	SN
4942D	Bandage, two layer, 22 cm- 28 cm (yellow)	4	..	..	*153.63	4.60	ProGuide 66000781	SN
4943E	Bandage, two layer, 28 cm- 32 cm (green)	4	..	..	*153.63	4.60	ProGuide 66000782	SN
BANDAGE—RETENTION—COHESIVE—HEAVY								
4811F	Bandage 5 cm x 1.3 m	2	..	..	*12.01	4.60	Peg 7420	BK
4812G	Bandage 7.5 cm x 1.3 m	2	..	..	*15.15	4.60	Peg 7422	BK
4659F	Bandage 7.5 cm x 3 m	2	..	..	*17.79	4.60	Coplus 3629	BV
4813H	Bandage 10 cm x 1.3 m	2	..	..	*18.75	4.60	Peg 7423	BK
4660G	Bandage 10 cm x 2 m	2	..	..	*17.13	4.60	Coban 1584	MM
4814J	Bandage 15 cm x 1.3 m	2	..	..	*25.57	4.60	Peg 7425	BK
BANDAGE—RETENTION—COHESIVE—LIGHT								
4718H	Bandages 2.5 cm x 4 m, 2	‡1	..	..	10.58	4.60	Handygauze Cohesive 8631	BV
4719J	Bandage 6 cm x 4 m	2	..	..	*12.67	4.60	Handygauze Cohesive 8633	BV
4662J	Bandage 10 cm x 4 m	2	..	..	*15.01	4.60	Handygauze Cohesive 8635	BV

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
BANDAGE—RETENTION—COTTON CREPE								
4727T	Bandage 5 cm x 2.3 m	2	..	..	*15.29 *16.11	4.60 4.60	Telfa 8252F Elastocrepe 36102520	KE BV
4728W	Bandage 7.5 cm x 2.3 m	2	..	..	*19.87 *20.05	4.60 4.60	Telfa 8253F Elastocrepe 36102420	KE BV
4729X	Bandage 10 cm x 2.3 m	2	..	..	*22.87 *25.21	4.60 4.60	Telfa 8254F Elastocrepe 36102320	KE BV
BANDAGE—TUBULAR								
4855M	Bandage 6.25 cm x 1 m	‡1	..	..	15.99	4.60	Tubigrip B 1520	SS
4856N	Bandage 6.75 cm x 1 m	‡1	..	..	15.99	4.60	Tubigrip C 1545	SS
4857P	Bandage 7.5 cm x 1 m	‡1	..	..	15.99	4.60	Tubigrip D 1546	SS
4858Q	Bandage 8.75 cm x 1 m	‡1	..	..	15.99	4.60	Tubigrip E 1547	SS
4859R	Bandage 10 cm x 1 m	‡1	..	..	15.99	4.60	Tubigrip F 1548	SS
4663K	Bandage, straight, size C	‡1	..	..	13.07	4.60	Elastoplast 2225	BE
4664L	Bandage, straight, size D	‡1	..	..	13.07	4.60	Elastoplast 2226	BE
4665M	Bandage, straight, size E	‡1	..	..	13.07	4.60	Elastoplast 2227	BE
4667P	Bandage, lightweight, 8.75 cm x 1 m	‡1	..	..	14.94	4.60	Tensogrip 36361259	BV
BANDAGE—TUBULAR (FINGER)								
4798M	Complete pack including applicator	‡1	..	..	15.61	4.60	Tubegauz 0501633	SS
4726R	Refill	‡1	..	..	11.75	4.60	Tubegauz 0501658	SS
BANDAGE—TUBULAR (LIGHTWEIGHT)								
4671W	Bandage, small limb size (red), 10 m	‡1	..	..	25.74	4.60	Tubifast 2434	SS
4672X	Bandage, medium limb size (green), 10 m	‡1	..	..	29.24	4.60	Tubifast 2436	SS
4673Y	Bandage, large limb size (blue), 10 m	‡1	..	..	32.65	4.60	Tubifast 2438	SS
BANDAGE—TUBULAR (LONG STOCKING)								
4674B	Bandage, small size	2	..	..	*37.05	4.60	Tubigrip 1482	SS
4797L	Bandage, medium size	2	..	..	*37.05	4.60	Tubigrip 1483	SS
4799N	Bandage, large size	2	..	..	*37.05	4.60	Tubigrip 1484	SS
4675C	Bandage, XX/large size	2	..	..	*37.05	4.60	Tubigrip 1486	SS
BANDAGE—TUBULAR (SHORT STOCKING)								
4661H	Bandage, small B/C size	2	..	..	*27.71	4.60	Tubigrip 1479	SS
4815K	Bandage, medium C/D size	2	..	..	*27.71	4.60	Tubigrip 1480	SS
4816L	Bandage, large D/E size	2	..	..	*27.71	4.60	Tubigrip 1481	SS

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
BANDAGE—ZINC PASTE								
NOTE: Used as an adjunct in the management of leg ulceration and associated eczema and skin conditions.								
4668Q	Bandage 7.5 cm x 6 m	2	..	..	*26.53	4.60	Zincaband 3604	SS
4669R	Bandage 7.5 cm x 6 m	2	..	..	*26.97	4.60	Steripaste 3610	SS
4750B	Bandage 7.5 cm x 6 m	2	..	..	*59.23	4.60	Viscopaste 4948	SN
4749Y	Bandage 8 cm x 5 m (compression)	2	..	..	*35.49	4.60	Gelocast Elastic 1080	BV
4670T	Bandage 10 cm x 9.1 m	2	..	..	*26.13	4.60	Flexidress 650941	CC
4760M	Bandages 80 cm (stockings), 4	‡1	..	..	68.37	4.60	ZipZoc 66051550	SN
COTTON WOOL ROLL								
4701K	Roll 100 g	‡1	2	..	8.67	4.60	JJ 02013	JJ
DRESSING—ACTIVATED CHARCOAL (MALODOROUS WOUND)								
4742N	Dressings 10 cm x 10 cm, 10	‡1	..	..	77.31	4.60	CarboFLEX 403202	CC
4681J	Dressing 10.5 cm x 10.5 cm	10	..	..	*83.55	4.60	Actisorb Plus MAC031	JJ
4743P	Dressings 15 cm x 20 cm, 5	‡1	..	..	88.20	4.60	CarboFLEX 403204	CC
DRESSING—ALGINATE (CAVITY WOUND)								
NOTE: This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.								
4832H	Rope 2 g	10	..	..	*113.59	4.60	Kaltostat 168117	CC
				..	*132.55	4.60	Sorbsan 1411	UM
4685N	Ropes 2 g (30 cm), 5	2	..	..	*80.45	4.60	Restore CalciCare 9940	HO
4682K	Ropes 2 g (40 cm), 6	2	..	..	*116.07	4.60	Comfeel SeaSorb Filler 3740	CT
DRESSING—ALGINATE (SUPERFICIAL WOUND)								
NOTE: This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.								
4699H	Dressings 5 cm x 5 cm, 10	‡1	1	..	37.37	4.60	Restore CalciCare 9938	HO
				..	40.19	4.60	Algisite M 66000519	SN
				..	47.08	4.60	Kaltostat 168210	CC

continued ➡

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DRESSING—ALGINATE (SUPERFICIAL WOUND)—cont.								
4684M	Dressing 5 cm x 5 cm	10	1	..	*40.15	4.60	Comfeel SeaSorb Dressing 3705	CT
4683L	Dressings 7.5 cm x 12 cm, 10	‡1	1	..	89.41	4.60	Kaltostat 168212	CC
4831G	Dressing 10 cm x 10 cm	10	1	..	*79.55	4.60	Comfeel SeaSorb Dressing 3710	CT
				..	*81.85	4.60	Sorbsan 1410	UM
4700J	Dressings 10 cm x 10 cm, 10	‡1	1	..	69.97	4.60	Restore CalciCare 9937	HO
				..	78.21	4.60	Algisite M 66000520	SN
4691X	Dressings 15 cm x 20 cm, 10	‡1	1	..	190.65	4.60	Algisite M 66000521	SN
DRESSING with CADEXOMER IODINE								
NOTE: Suitable for yellow sloughy infected and malodorous wounds.								
4931M	Sachets 3 g, 7	‡1	..	..	53.05	4.60	Iodosorb Powder 66051070	SN
4932N	Tubes 10 g, 4	‡1	..	..	85.71	4.60	Iodosorb Ointment 66051240	SN
4933P	Tubes 20 g, 2	‡1	..	..	84.92	4.60	Iodosorb Ointment 66051230	SN
4935R	Sachets 5 g (6 cm x 4 cm), 5	‡1	..	..	76.51	4.60	Iodosorb 66051330	SN
4936T	Sachets 10 g (8 cm x 6 cm), 3	‡1	..	..	110.92	4.60	Iodosorb 66051340	SN
4937W	Sachets 17 g (10 cm x 8 cm), 2	‡1	..	..	116.85	4.60	Iodosorb 66051360	SN
DRESSING—FILM								
4686P	Dressings 6 cm x 7 cm, 8	‡1	..	..	13.57	4.60	Nexcare Tegaderm Transparent H1624	MM
4687Q	Dressings 10 cm x 12 cm, 4	‡1	..	..	17.40	4.60	Nexcare Tegaderm Transparent H1626	MM
4893M	Dressings 10 cm x 12 cm, 10	‡1	..	..	24.56	4.60	Op-Site Flexigrid 4629	SN
4688R	Dressing 15 cm x 20 cm	6	..	..	*27.91	4.60	Tegaderm Transparent 1628	MM

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DRESSING—FILM ISLAND								
4689T	Dressing 5 cm x 7 cm	10	..	..	*14.15	4.60	Tegaderm Transparent Island 3582	MM
4898T	Dressings 5 cm x 7.2 cm, 5	2	..	..	*21.33	4.60	Cutifilm Plus 76309	SN
4899W	Dressings 8 cm x 10 cm, 5	2	..	..	*34.07	4.60	Cutifilm Plus 76308	SN
4690W	Dressing 9 cm x 10 cm	10	..	..	*25.15	4.60	Tegaderm Transparent Island 3586	MM
DRESSING—FOAM with CHARCOAL (MALODOROUS WOUND)								
NOTE: This dressing should remain in place on wounds with odour until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.								
4892L	Dressings 10 cm x 10 cm, 10	2	..	..	*172.39	4.60	Lyof foam C 603025	SS
DRESSING—FOAM—HEAVY EXUDATE								
NOTE: This dressing should remain in place until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.								
4692Y	Dressings (foam alternative) 10 cm x 10 cm, 10	‡1	..	..	53.13	4.60	CombiDERM 651031	CC
4693B	Dressings (foam alternative) 15 cm x 18 cm, 5	‡1	..	..	70.05	4.60	CombiDERM 651027	CC
DRESSING—FOAM—LIGHT EXUDATE								
NOTE: This dressing should remain in place until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.								
4890J	Dressings 7.5 cm x 7.5 cm, 10	‡1	1	..	39.86	4.60	Lyof foam Flat 603092	SS
4891K	Dressings 10 cm x 10 cm, 10	‡1	1	..	47.17	4.60	Lyof foam Flat 603093	SS
4878R	Dressings 20 cm x 15 cm, 10	‡1	..	..	100.19	4.60	Lyof foam Flat 603095	SS

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DRESSING—FOAM—MODERATE EXUDATE								
NOTE: This dressing should remain in place until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.								
4795J	Dressings 10 cm x 10 cm, 10	‡1	..	..	73.39	4.60	Lyof foam Extra 603088	SS
					107.22	4.60	Allevyn 66007637	SN
4590N	Dressings 12.5 cm x 12.5 cm, 10	‡1	..	..	100.99	4.60	Allevyn Adhesive 66000044	SN
NOTE: Care should be taken when changing <i>Allevyn Adhesive</i> dressings to avoid skin tears.								
4880W	Dressings 20 cm x 15 cm, 10	‡1	..	..	187.44	4.60	Lyof foam Extra 603090	SS
4694C	Dressing, cavity, conforming, 20 g	1	..	..	68.59	4.60	Cavicare 4563	SN
4927H	Non-adhesive waterproof semi-permeable absorbent foam pads 10 cm x 10 cm, 10	‡1	1	..	80.43	4.60	Biatain Non-adhesive 3410	CT
4928J	Non-adhesive waterproof semi-permeable absorbent foam pads 15 cm x 15 cm, 5	‡1	2	..	79.00	4.60	Biatain Non-adhesive 3413	CT
4929K	Adhesive waterproof semi-permeable absorbent foam pads 12 cm x 12 cm, 10	‡1	1	..	88.57	4.60	Biatain Adhesive 3420	CT
4930L	Adhesive waterproof semi-permeable absorbent foam pads 18 cm x 18 cm, 5	‡1	2	..	85.82	4.60	Biatain Adhesive 3423	CT
DRESSING—GAUZE (ABSORBENT PAD)								
4707R	Pads 5 cm x 5 cm, 100	‡1	..	..	11.89	4.60	Handy 5672	BV
4708T	Pads 10 cm x 10 cm, 100	‡1	..	..	24.81	4.60	Handy 5674	BV
DRESSING—GAUZE—EYE PAD								
4768Y	Pads, 12	‡1	..	..	10.88	4.60	Curity 4112	KE
DRESSING—GAUZE—PARAFFIN								
4759L	Dressings 10 cm x 10 cm, 10	‡1	..	..	15.10	4.60	Jelonet 7404	SN
DRESSING—GAUZE—PARAFFIN with CHLORHEXIDINE ACETATE								
4845B	Dressings 10 cm x 10 cm, 10	‡1	2	..	19.92	4.60	Bactigras 7457	SN
DRESSING—GAUZE—POVIDONE-IODINE PAD								
4779M	Pads 22.5 cm x 7.5 cm, 12	‡1	2	..	30.69	4.60	Betadine	FH
DRESSING—HYDROACTIVE (CAVITY WOUND)								
4918W	Dressings 5 cm x 6 cm, 10	‡1	1	..	71.73	4.60	Allevyn Plus Cavity 66047571	SN
4919X	Dressings 10 cm x 10 cm, 5	2	1	..	*152.05	4.60	Allevyn Plus Cavity 66047573	SN

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DRESSING—HYDROACTIVE (DEBRIDEMENT)								
4949L	Dressings 4 cm, 8	‡1	..	..	66.23	4.60	TenderWet 24 Active	HR
4948K	Dressings 5.5 cm, 8	‡1	..	..	66.99	4.60	TenderWet Active Cavity	HR
4950M	Dressings 7.5 cm x 7.5 cm, 8	‡1	..	..	90.52	4.60	TenderWet 24 Active	HR

NOTE:

Hartmann products are not available through pharmacy wholesalers. To order please contact the Hartmann Alliance (refer to 'Ordering Hartmann Products' at the foot of the 'Wound Assessment and Dressing Identification' chart at the beginning of this section).

DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—HIGH EXUDATE)								
4695D	Dressings, island, 11 cm x 11 cm, 10	‡1	..	..	84.29	4.60	Tielle MT2440	JJ
4696E	Dressings, island, 18 cm x 18 cm, 5	‡1	..	..	100.26	4.60	Tielle MT2442	JJ
DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—LIGHT EXUDATE)								
4905E	Dressings 5 cm x 6 cm, 10	‡1	1	..	49.40	4.60	Allevyn Thin 66047576	SN
4906F	Dressings 10 cm x 10 cm, 5	2	1	..	*90.45	4.60	Allevyn Thin 66047578	SN
DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—MODERATE EXUDATE)								
4885D	Dressings 5 cm x 6 cm, 10	‡1	1	..	38.25	4.60	Cutinova Hydro 66047441	SN
4886E	Dressings 10 cm x 10 cm, 5	2	1	..	*64.89	4.60	Cutinova Hydro 66047443	SN

DRESSING—HYDROCOLLOID (CAVITY WOUND)

NOTE:

This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

4896Q	Paste 30 g	10	..	..	*143.45	4.60	DuoDERM Paste H7930	CC
4895P	Paste 50 g	6	..	..	*99.79	4.60	Comfeel Paste 4701	CT
4697F	Powder 6 g	2	..	..	*19.07	4.60	Comfeel Powder 4706	CT
4698G	Ropes 2 g (30 cm), 5	‡1	..	..	82.04	4.60	Aquacel 177904	CC

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—LIGHT EXUDATE)								
NOTE: This dressing should be applied to a thickness of 3 mm to 5 mm. It should be covered with a hydrocolloid dressing and may be left in place for up to 7 days.								
4888G	Dressings 5 cm x 7 cm, 10	‡1	1	..	34.87	4.60	Comfeel Plus Transparent 3530	CT
4889H	Dressings 9 cm x 14 cm, 10	‡1	1	..	74.62	4.60	Comfeel Plus Transparent 3536	CT
4908H	Dressings 10 cm x 10 cm, 5	2	1	..	*51.35	4.60	Restore Extra Thin 9921	HO
4907G	Dressings 10 cm x 10 cm, 10	‡1	1	..	70.05	4.60	DuoDERM Extra Thin H7955	CC
4924E	Dressings 10 cm x 10 cm, 10	‡1	1	..	61.20	4.60	Comfeel Plus Transparent 3533	CT
4947J	Dressings 10 cm x 10 cm, 10	‡1	1	..	45.70	4.60	Hydrocoll Thin 900942/1	HR
NOTE: Hartmann products are not available through pharmacy wholesalers. To order please contact the Hartmann Alliance (refer to 'Ordering Hartmann Products' at the foot of the 'Wound Assessment and Dressing Identification' chart at the beginning of this section).								
DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—MODERATE EXUDATE)								
NOTE: This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.								
4897R	Dressings 10 cm x 10 cm, 5	2	1	..	*57.15	4.60	Restore Plus 9956	HO
					*79.73	4.60	DuoDERM CGF H7660	CC
4921B	Dressings 10 cm x 10 cm, 10	‡1	1	..	66.85	4.60	Replicare Ultra 66000434	SN
					99.31	4.60	Aquacel 177902	CC
4945G	Dressings 10 cm x 10 cm, 10	‡1	1	..	45.70	4.60	Hydrocoll 900938/1	HR
NOTE: Hartmann products are not available through pharmacy wholesalers. To order please contact the Hartmann Alliance (refer to 'Ordering Hartmann Products' at the foot of the 'Wound Assessment and Dressing Identification' chart at the beginning of this section).								

continued ➡

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—MODERATE EXUDATE)—cont.								
4922C	Dressings 15 cm x 15 cm, 5	2	1	..	*128.95	4.60	Repicare Ultra 66000435	SN
				..	*207.03	4.60	Aquacel 177903	CC
4946H	Dressings 15 cm x 15 cm, 10	‡1	1	..	88.24	4.60	Hydrocoll 900939/1	HR
NOTE: Hartmann products are not available through pharmacy wholesalers. To order please contact the Hartmann Alliance (refer to 'Ordering Hartmann Products' at the foot of the 'Wound Assessment and Dressing Identification' chart at the beginning of this section).								
4853K	Dressings 15 cm x 20 cm, 3	3	1	..	*131.59	4.60	Restore Plus 9957	HO
4854L	Dressings 20 cm x 20 cm, 3	3	1	..	*156.76	4.60	Restore Plus 9958	HO
4920Y	Dressings 20 cm x 20 cm, 5	2	1	..	*220.65	4.60	DuoDERM CGF H7662	CC
4923D	Dressings with alginate 10 cm x 10 cm, 10	‡1	1	..	71.98	4.60	Comfeel Plus Ulcer Dressing 3110	CT
4842W	Dressings, sacral, 5	2	1	..	*97.93	4.60	Restore Plus Sacral 9959	HO
				..	*153.61	4.60	Repicare Ultra 66000437	SN
DRESSING—HYDROGEL—AMORPHOUS								
NOTE: This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.								
4912M	Tubes 15 g, 10	‡1	1	..	62.81	4.60	DuoDERM Gel H7990	CC
				..	63.40	4.60	Comfeel Purilon Gel 3900	CT
4894N	Tubes 25 g, 10	‡1	1	..	112.84	4.60	Intrasite Gel 7313	SN
4913N	Tubes 30 g, 3	3	1	..	*95.44	4.60	DuoDERM Gel H7987	CC
4914P	Tube 50 g	12	1	..	*91.87	4.60	Solugel 10336	JJ
4599C	Tube 50 g	6	1	..	*38.11	4.60	SoloSite Gel 36361338	SN

REPATRIATION

Code	Name, Restriction, Manner of administration and form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
DRESSING—HYDROGEL—SHEET								
NOTE: This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.								
4911L	Dressings 9.5 cm x 10.2 cm, 5	2	..	..	*81.53	4.60	Nu-Gel 2497	JJ
4806Y	Dressings 11 cm x 10 cm, 5	2	..	..	*51.35	4.60	Aquaclear	HR
NOTE: Hartmann products are not available through pharmacy wholesalers. To order please contact the Hartmann Alliance (refer to 'Ordering Hartmann Products' at the foot of the 'Wound Assessment and Dressing Identification' chart at the beginning of this section).								
DRESSING—NON-ADHERENT								
4860T	Dressings 5 cm x 5 cm, 5	2	..	..	*12.19	4.60	Melolin 36101720	SN
4819P	Dressings 5 cm x 5 cm, 5	2	..	..	*11.69	4.60	Cutilin Non- Stick Wound Pad 76301	SN
4755G	Dressings 5 cm x 7.5 cm, 10	‡1	..	..	9.15	4.60	Telfa 1970C	KE
4909J	Dressing 7.6 cm x 7.6 cm	10	1	..	*13.65	4.60	Adaptic 2012	JJ
4758K	Dressings 7.5 cm x 10 cm, 6	‡1	..	..	9.35	4.60	Telfa 2140C	KE
4944F	Dressings 7.5 cm x 10 cm, 10	‡1	..	..	13.19	4.60	Atrauman 499513	HR
NOTE: Hartmann products are not available through pharmacy wholesalers. To order please contact the Hartmann Alliance (refer to 'Ordering Hartmann Products' at the foot of the 'Wound Assessment and Dressing Identification' chart at the beginning of this section).								
4862X	Dressings 10 cm x 10 cm, 5	2	..	..	*18.75	4.60	Cutilin Non- Stick Wound Pad 76300	SN
4861W	Dressings 10 cm x 10 cm, 10	‡1	..	..	25.83	4.60	Melolin 66974933	SN
4843X	Dressings, self-adhesive, 5 cm x 7.5 cm, 10	‡1	2	..	10.72	4.60	Telfa 6020C	KE
4844Y	Dressings, self-adhesive, 7.5 cm x 10 cm, 6	‡1	2	..	10.11	4.60	Telfa 7650C	KE
GAUZE and COTTON TISSUE (COMBINE ROLL)								
4767X	Wrapped pack 9 cm x 10 m	‡1	..	..	13.33	4.60	BSN 2902165	BV
4761N	Wrapped pack 10 cm x 10 m	‡1	..	..	13.06	4.60	JJ 12010	JJ
GLOVES PLASTIC (DISPOSABLE)								
4772E	Gloves, small, 100	‡1	..	..	10.28	4.60	Handy 4207	BV
4773F	Gloves, medium, 100	‡1	..	..	10.28	4.60	Handy 4208	BV
4774G	Gloves, large, 100	‡1	..	..	10.28	4.60	Handy 4209	BV
LUBRICATING AGENT								
4318G	Jelly 60 g	‡1	..	..	9.04	4.60	Surgical Lubricating Gel	BI

REPATRIATION

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No.of Rpts	Premium	Dispensed Price for Max.Qty \$	Maximum Recordable Value for Safety Net \$	Proprietary Name and Manufacturer	
PRESSURE REDUCING PRODUCTS								
4676D	Sheet 10 cm x 10 cm x 3 mm	2	..	..	*25.87	4.60	Spenco Dermal Pad 10-553	KC
4677E	Sheet 10 cm x 10 cm x 12 mm	2	..	..	*46.99	4.60	Spenco Dermal Pad 10-561	KC
4678F	Butterfly shape 7 cm	5	..	..	*48.15	4.60	Comfeel Plus Pressure Relieving 3350	CT
4679G	Round 10 cm	5	..	..	*52.25	4.60	Comfeel Plus Pressure Relieving 3353	CT
TAPES—NON-WOVEN RETENTION (POLYACRYLATE)								
4915Q	Roll 2.5 cm x 9.1 m	‡1	..	..	10.92	4.60	Medipore 2961	MM
4863Y	Roll 2.5 cm x 10 m	‡1	..	..	14.25	4.60	Hypafix 71443-0	BV
4917T	Roll 2.5 cm x 10 m	‡1	..	..	9.17	4.60	Mefix 310250	MH
4916R	Roll 5 cm x 10 m	‡1	..	..	21.21	4.60	Hypafix 71443-1	BV
TAPES—PLASTER ADHESIVE ELASTIC								
4780N	Roll 2.5 cm x 2.5 m	‡1	..	..	10.81	4.60	Leukoplast 1071	BV
4781P	Roll 5 cm x 2.5 m	‡1	..	..	16.37	4.60	Leukoplast 1072	BV
4782Q	Roll 7.5 cm x 2.5 m	‡1	..	..	19.78	4.60	Leukoplast 1073	BV
4735F	Roll 10 cm x 2.5 m	‡1	..	..	19.99	4.60	Elastoplast 1004	BV
TAPES—PLASTER ADHESIVE HYPOALLERGENIC								
4783R	Roll 1.25 cm x 5 m	‡1	..	..	8.50	4.60	Leukopor 2471	BV
4785W	Roll 1.25 cm x 5 m	‡1	..	..	8.77	4.60	Leukosilk 1021	BV
4794H	Roll 2.5 cm x 5 m	‡1	..	..	10.79	4.60	Leukopor 2472	BV
4787Y	Roll 2.5 cm x 5 m	‡1	..	..	11.27	4.60	Leukosilk 1022	BV
4788B	Stretch roll 5 cm x 5 m	‡1	..	..	15.07	4.60	Leukoflex 1124	BV
4790D	Roll 5 cm x 5 m	‡1	..	..	14.11	4.60	Leukopor 2474	BV
4789C	Roll 5 cm x 5 m	‡1	..	..	14.91	4.60	Leukosilk 1024	BV
4848E	Roll (dispenser) 1.9 cm x 5.4 m	‡1	..	..	9.18	4.60	Nexcare Durable Cloth First Aid Tape 799	MM
4849F	Roll (dispenser) 1.9 cm x 7.3 m	‡1	..	..	9.18	4.60	Nexcare Gentle Paper First Aid Tape 789	MM

Section 2

Standard Packs and Prices

NOTE—

Standard packs and prices (including mark-up, but without dispensing fee and dangerous drug fee) are for items against the price of which an asterisk () is shown in Section 1 of the Schedule.*

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
4579B	Alprostadil	10 mcg	2 @	25.40 PH
4580C		20 mcg	2 @	32.40 PH
4117Q	Aluminium Hydroxide with Magnesium Hydroxide and Simethicone	400 mg-400 mg-30 mg	100 @	9.68 PC
4118R		400 mg-400 mg-30 mg per 5 mL, 500 mL	1 @	7.76 PC
4651T	Bandage—Absorbent Wool	10 cm x 2.7 m	1 @	4.47 BV
4654Y	Bandage—Compression	8 cm x 5 m	1 @	12.86 BV
4736G		7.5 cm x 3 m	1 @	15.54 BV
4656C		7.5 cm x 3.5 m	1 @	12.39 SS
4748X		10 cm x 3 m	1 @	13.30 CC
			1 @	20.57 BV
4657D		10 cm x 3.5 m	1 @	9.85 BV
			1 @	14.43 SS
4658E		Four layer	1 @	38.31 SN
4598B		Four layer	1 @	25.47 SN
4941C		Two layer, 18 cm-22 cm	1 @	37.22 SN
4942D		Two layer, 22 cm-28 cm	1 @	37.22 SN
4943E		Two layer, 28 cm-32 cm	1 @	37.22 SN
4811F	Bandage—Retention—Cohesive— Heavy	5 cm x 1.3 m	1 @	3.63 BK
4812G		7.5 cm x 1.3 m	1 @	5.20 BK
4659F		7.5 cm x 3 m	1 @	6.52 BV
4813H		10 cm x 1.3 m	1 @	7.00 BK
4660G		10 cm x 2 m	1 @	6.19 MM
4814J		15 cm x 1.3 m	1 @	10.41 BK
4719J	Bandage—Retention—Cohesive— Light	6 cm x 4 m	1 @	3.96 BV
4662J		10 cm x 4 m	1 @	5.13 BV
4727T	Bandage—Retention—Cotton Crepe	5 cm x 2.3 m	1 @	5.27 KE
			1 @	5.68 BV
4728W		7.5 cm x 2.3 m	1 @	7.56 KE
			1 @	7.65 BV
4729X		10 cm x 2.3 m	1 @	9.06 KE
			1 @	10.23 BV
4674B	Bandage—Tubular (Long Stocking)	Small	1 @	16.15 SS
4797L		Medium	1 @	16.15 SS
4799N		Large	1 @	16.15 SS
4675C		XX/large	1 @	16.15 SS
4661H	Bandage—Tubular (Short Stocking)	Small B/C	1 @	11.48 SS
4815K		Medium C/D	1 @	11.48 SS
4816L		Large D/E	1 @	11.48 SS
4668Q	Bandage—Zinc Paste	7.5 cm x 6 m	1 @	10.89 SS
4669R		7.5 cm x 6 m	1 @	11.11 SS
4750B		7.5 cm x 6 m	1 @	27.24 SN
4749Y		8 cm x 5 m	1 @	15.37 BV
4670T		10 cm x 9.1 m	1 @	10.69 CC
4150K	Bromazepam	3 mg	30 @	10.02 RO
4151L		6 mg	30 @	12.90 RO
4055K	Calcium Carbonate with Glycine	420 mg-180 mg	100 @	7.47 MM
4081T	Dextropropoxyphene Napsylate	100 mg	10 @	2.33 AS
4681J	Dressing—Activated Charcoal (Malodorous Wound)	10.5 cm x 10.5 cm	1 @	7.88 JJ

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Approved Name	Form/Strength	Pack and Price \$	Manufacturer
4832H	Dressing—Alginate (Cavity Wound)	2 g	1 @ 12.78	UM
			5 @ 54.42	CC
4685N		2 g (30 cm), 5	1 @ 37.85	HO
4682K		2 g (40 cm), 6	1 @ 55.66	CT
4684M	Dressing—Alginate (Superficial Wound)	5 cm x 5 cm	1 @ 3.54	CT
4831G		10 cm x 10 cm	1 @ 7.48	CT
			1 @ 7.71	UM
4688R	Dressing—Film	15 cm x 20 cm	1 @ 3.86	MM
4689T	Dressing—Film Island	5 cm x 7 cm	1 @ 0.94	MM
4898T		5 cm x 7.2 cm, 5	1 @ 8.29	SN
4899W		8 cm x 10 cm, 5	1 @ 14.66	SN
4690W		9 cm x 10 cm	1 @ 2.04	MM
4892L	Dressing—Foam with Charcoal (Malodorous Wound)	10 cm x 10 cm, 10	1 @ 83.82	SS
4919X	Dressing—Hydroactive (Cavity Wound)	10 cm x 10 cm, 5	1 @ 73.65	SN
4906F	Dressing—Hydroactive (Superficial Wound—Light Exudate)	10 cm x 10 cm, 5	1 @ 42.85	SN
4886E	Dressing—Hydroactive (Superficial Wound—Moderate Exudate)	10 cm x 10 cm, 5	1 @ 30.07	SN
4896Q	Dressing—Hydrocolloid (Cavity Wound)	30 g	1 @ 13.87	CC
4895P		50 g	1 @ 15.84	CT
4697F		6 g	1 @ 7.16	CT
4908H	Dressing—Hydrocolloid (Superficial Wound—Light Exudate)	10 cm x 10 cm, 5	1 @ 23.30	HO
4897R	Dressing—Hydrocolloid (Superficial Wound—Moderate Exudate)	10 cm x 10 cm, 5	1 @ 26.20	HO
			1 @ 37.49	CC
4922C		15 cm x 15 cm, 5	1 @ 62.10	SN
			1 @ 101.14	CC
4853K		15 cm x 20 cm, 3	1 @ 42.28	HO
4854L		20 cm x 20 cm, 3	1 @ 50.67	HO
4920Y		20 cm x 20 cm, 5	1 @ 107.95	CC
4842W		5	1 @ 46.59	HO
			1 @ 74.43	SN
4913N	Dressing—Hydrogel—Amorphous	30 g, 3	1 @ 30.23	CC
4914P		50 g	1 @ 7.26	JJ
4599C		50 g	1 @ 5.56	SN
4911L	Dressing—Hydrogel—Sheet	9.5 cm x 10.2 cm, 5	1 @ 38.39	JJ
4806Y		11 cm x 10 cm, 5	1 @ 23.30	HR
4860T	Dressing—Non-Adherent	5 cm x 5 cm, 5	1 @ 3.72	SN
4819P		5 cm x 5 cm, 5	1 @ 3.47	SN
4909J		7.6 cm x 7.6 cm	1 @ 0.89	JJ
4862X		10 cm x 10 cm, 5	1 @ 7.00	SN
4237B	Fexofenadine Hydrochloride	60 mg	20 @ 13.66	AV
4298F	Gatifloxacin	400 mg in 40 mL	1 @ 79.99	BQ
4299G		400 mg in 200 mL	1 @ 79.99	BQ
4246L	Glycerol	2.8 g, 12	1 @ 3.59	PP
4283K	Ipratropium Bromide with Salbutamol Sulfate	20 mcg (anhydrous)- 100 mcg (base) per dose (200 doses)	1 @ 21.20	BY
4330X	Moxifloxacin Hydrochloride	400 mg (base) in 250 mL	1 @ 64.80	BN

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer	
4576W	Nicotine	Approx. 5 mg per 16 hours, 7	1 @	21.95	PH
4571N		Approx. 7 mg per 24 hours, 7	1 @	19.40	AF
4577X		Approx. 10 mg per 16 hours, 7	1 @	33.95	GK
4572P		Approx. 14 mg per 24 hours, 7	1 @	24.12	PH
4578Y		Approx. 15 mg per 16 hours, 7	1 @	21.23	AF
4573Q	Pressure Reducing Products	Approx. 21 mg per 24 hours, 7	1 @	23.43	AF
4676D		10 cm x 10 cm x 3 mm	1 @	33.95	GK
4677E		10 cm x 10 cm x 12 mm	1 @	10.56	KC
4678F		7 cm	1 @	21.12	KC
4679G		10 cm	1 @	8.68	CT
4978B	Ranitidine Hydrochloride	150 mg (base), effervescent	1 @	9.50	CT
4980D		150 mg (base) per 10 mL, 300 mL	30 @	9.71	GK
			1 @	9.71	GK

Generic/Proprietary Index

for

PBS and RPBS Schedules

GENERIC /PROPRIETARY INDEX

A

ABACAVIR SULFATE			
.Section 100	359		
ABACAVIR SULFATE with LAMIVUDINE			
.Section 100	359		
ABACAVIR SULFATE with LAMIVUDINE and ZIDOVUDINE			
.Section 100	359		
Abbacillin-V (SI)			
.Antiinfectives for systemic use	163		
.Dental	339		
Abbacillin-VK Filmtab (SI)			
.Antiinfectives for systemic use	163		
.Dental	338		
ABCIXIMAB	102		
Abilify (BQ)	270, 271		
ACAMPROSATE CALCIUM	287		
ACARBOSE	96		
Accomin Adult Tonic (WT)			
.Repatriation Schedule	475		
Accu-Chek Active (RD)	310		
Accu-Chek Go (RD)	310		
Accu-Chek Integra (RD)	310		
Accupril (PF)	124		
Accuretic 10/12.5mg (PF)	126		
Accuretic 20/12.5mg (PF)	126		
Acenorm 12.5 mg (AF)	121		
Acenorm 25 mg (AF)	122		
Acenorm 50 mg (AF)	122		
ACETAZOLAMIDE	303		
ACETYLCYSTEINE	299		
ACICLOVIR			
.Antiinfectives for systemic use	177		
.Sensory organs	300		
Acihexal (HX)	177, 178		
Aci-Jel (JC)			
.Repatriation Schedule	484		
Acimax Tablets (AL)	82		
ACITRETIN	134		
Aclin (AF)			
.Dental	347		
.Musculo-skeletal system	239		
.Palliative Care	326		
Aclin 200 (AF)			
.Dental	347		
.Musculo-skeletal system	239		
.Palliative Care	326		
Aclor 125 (AW)			
.Antiinfectives for systemic use	168		
.Dental	343		
Aclor 250 (AW)			
.Antiinfectives for systemic use	168		
.Dental	343		
Actilax (AF)	88		
Actilyse (BY)	104		
Actisorb Plus MAC031 (JJ)			
.Repatriation Schedule	504		
Actonel (AV)			
.Musculo-skeletal system	246		
.Repatriation Schedule	491		
Actonel Once-a-Week (AV)			
.Musculo-skeletal system	246		
.Repatriation Schedule	491		
Actos (LY)	97		
Actrapid (NO)	93		
Actrapid Penfill 3 mL (NO)	94		
Acyclo-V 200 (AF)	177		
Acyclo-V 800 (AF)	178		
Adalat 10 (BN)	119		
Adalat 20 (BN)	119		
Adalat Oros 20mg (BN)			
.Cardiovascular system	119, 120		
.Repatriation Schedule	477		
Adalat Oros 30 (BN)	119		
Adalat Oros 60 (BN)	119		
ADALIMUMAB	197		
Adaptic 2012 (JJ)			
.Repatriation Schedule	511		
Addos XR 30 (AW)	119		
Addos XR 60 (AW)	119		
Adefin 10 (AF)	119		
Adefin 20 (AF)	119		
Adefin XL 30 (AF)	119		
Adefin XL 60 (AF)	119		
ADEFOVIR DIPIVOXIL			
.Section 100	359		
ADRENALINE			
.Cardiovascular system	110		
.Dental	334, 356		
.Doctor's Bag Supplies	71		
.Respiratory system	298		
Adriamycin Solution (PH)	185		
ADT (CS)			
.Antiinfectives for systemic use	180		
.Doctor's Bag Supplies	71		
Advantage II (RD)	309		
Advantan (SC)	136		
Aerodiol (SE)	145		
Aeron 250 (AW)	296		
Aeron 500 (AW)	297		
Agenerase (GK)			
.Section 100	360		
Aggrastat (MK)	104		
Airomir (MM)			
.Doctor's Bag Supplies	72		
.Respiratory system	291		
Airomir Autohaler (MM)	291		
Akamin 50 (AF)	161		
Akamin 100 (AF)	160		
Akineton (AB)	266		
Albalon-A (AG)			
.Repatriation Schedule	497		
Albalon Liquifilm (AG)			
.Repatriation Schedule	497		
ALBENDAZOLE	290		
Albey Bee Venom (TH)	308		
Albey Paper Wasp Venom (TH)	308		
Albey Yellow Jacket Venom (TH)	308		
Aldactone (PH)	115		
Aldazine 10 (AF)	269		

GENERIC /PROPRIETARY INDEX

<i>Aldazine 25 (AF)</i>	269	<i>Alphagan (AG)</i>	302
<i>Aldazine 50 (AF)</i>	269	<i>Alphamox 125 (AF)</i>	
<i>Aldazine 100 (AF)</i>	269	Antiinfectives for systemic use.....	162
<i>Aldomet (MK)</i>	112	Dental.....	337
ALEFACEPT		<i>Alphamox 250 (AF)</i>	
Repatriation Schedule.....	488	Antiinfectives for systemic use.....	161, 162
<i>Alendro Once Weekly (AW)</i>	245	Dental.....	337, 338
ALENDRONATE SODIUM.....	245	<i>Alphamox 500 (AF)</i>	
<i>Alepam 15 (AF)</i>		Antiinfectives for systemic use.....	161
Dental.....	355	Dental.....	337
Nervous system.....	273	<i>Alphapress 25 (AF)</i>	113
Palliative Care.....	331	<i>Alphapress 50 (AF)</i>	113
<i>Alepam 30 (AF)</i>		<i>Alphapril (AF)</i>	122, 123
Dental.....	355	<i>Alprax 0.25 (AW)</i>	272
Nervous system.....	273	<i>Alprax 0.5 (AW)</i>	272
Palliative Care.....	331	<i>Alprax 1 (AW)</i>	272
<i>Alfaré (NT)</i>	312	<i>Alprax 2 (AW)</i>	272
<i>Algisite M 66000519 (SN)</i>		ALPRAZOLAM.....	272
Repatriation Schedule.....	504	<i>Alprazolam-DP (DP)</i>	272
<i>Algisite M 66000520 (SN)</i>		<i>Alprim (AF)</i>	169
Repatriation Schedule.....	505	ALPROSTADIL	
<i>Algisite M 66000521 (SN)</i>		Repatriation Schedule.....	485
Repatriation Schedule.....	505	ALTEPLASE (Recombinant tissue-type plasminogen	
<i>Alimta (LY)</i>		activator).....	104
Special Pharmaceutical Benefit.....	75	ALUMINIUM HYDROXIDE with MAGNESIUM	
<i>Alkeran (GK)</i>	181	HYDROXIDE.....	77
ALLANTOIN with GLYCEROL and ICHTHAMMOL		ALUMINIUM HYDROXIDE with MAGNESIUM	
Repatriation Schedule.....	483	HYDROXIDE and SIMETHICONE	
ALLANTOIN with SULFUR, PHENOL, COAL TAR		Repatriation Schedule.....	472
SOLUTION and MENTHOL		ALUMINIUM HYDROXIDE with MAGNESIUM	
Repatriation Schedule.....	480	TRISILICATE and MAGNESIUM HYDROXIDE.....	77
<i>Allegron (AS)</i>	276	<i>Alvesco 80 (AH)</i>	295
<i>Allelyn 66007637 (SN)</i>		<i>Alvesco 160 (AH)</i>	295
Repatriation Schedule.....	507	AMANTADINE HYDROCHLORIDE.....	267
<i>Allelyn Adhesive 66000044 (SN)</i>		<i>Amaryl (AV)</i>	96
Repatriation Schedule.....	507	<i>Amevive (BD)</i>	
<i>Allelyn Plus Cavity 66047571 (SN)</i>		Repatriation Schedule.....	489
Repatriation Schedule.....	507	AMILORIDE HYDROCHLORIDE.....	115
<i>Allelyn Plus Cavity 66047573 (SN)</i>		AMINO ACID FORMULA without METHIONINE,	
Repatriation Schedule.....	507	THREONINE and VALINE and low in	
<i>Allelyn Thin 66047576 (SN)</i>		ISOLEUCINE.....	315
Repatriation Schedule.....	508	AMINO ACID FORMULA without PHENYLALANINE.....	315
<i>Allelyn Thin 66047578 (SN)</i>		AMINO ACID FORMULA without PHENYLALANINE,	
Repatriation Schedule.....	508	TYROSINE and METHIONINE.....	315
<i>Allohexal (HX)</i>	244	AMINO ACID FORMULA with VITAMINS, MINERALS	
ALLOPURINOL.....	244	and LONG CHAIN POLYUNSATURATED FATTY	
<i>Allosig (FM)</i>	244	ACIDS without PHENYLALANINE.....	315
<i>Alodorm (AF)</i>		AMINO ACID FORMULA with VITAMINS and	
Dental.....	356	MINERALS without METHIONINE.....	315
Nervous system.....	262, 274	AMINO ACID FORMULA with VITAMINS and	
Palliative Care.....	331	MINERALS without METHIONINE, THREONINE	
<i>Alpha Keri Bath Oil (MT)</i>		and VALINE and low in ISOLEUCINE.....	316
Repatriation Schedule.....	479	AMINO ACID FORMULA with VITAMINS and	
<i>Alpha Keri Lotion (MT)</i>		MINERALS without PHENYLALANINE.....	316
Repatriation Schedule.....	479	AMINO ACID FORMULA with VITAMINS and	
<i>Alphacin 250 (AF)</i>		MINERALS without PHENYLALANINE and	
Antiinfectives for systemic use.....	162	TYROSINE.....	316
Dental.....	338	AMINO ACID FORMULA with VITAMINS and	
<i>Alphacin 500 (AF)</i>		MINERALS without VALINE, LEUCINE and	
Antiinfectives for systemic use.....	162	ISOLEUCINE.....	317
Dental.....	338	AMINO ACIDS—SYNTHETIC, FORMULA.....	311

GENERIC /PROPRIETARY INDEX

AMINOGLUTETHIMIDE	
.Antineoplastic and immunomodulating agents	192
.Systemic hormonal preparations, excl. sex hormones and insulins	157
AMIODARONE HYDROCHLORIDE	109
AMISULPRIDE	270
AMITRIPTYLINE HYDROCHLORIDE	275
<i>Amizide (AF)</i>	115
AMLODIPINE BESYLATE	
.Cardiovascular system	118
.Repatriation Schedule	476
<i>Amohexal (HX)</i>	
.Antiinfectives for systemic use	161, 162
.Dental	337, 338
AMOROLFINE HYDROCHLORIDE	
.Repatriation Schedule	478
<i>Amoxil (GK)</i>	
.Antiinfectives for systemic use	161, 162
.Dental	337
<i>Amoxil Duo (GK)</i>	162
<i>Amoxil Forte (GK)</i>	
.Antiinfectives for systemic use	162
.Dental	338
AMOXYCILLIN	
.Antiinfectives for systemic use	161, 162
.Dental	337
AMOXYCILLIN with CLAVULANIC ACID	
.Antiinfectives for systemic use	165
.Dental	340
<i>Amoxycillin Sandoz (BG)</i>	162
<i>Amoxycillin-DP (DG)</i>	
.Antiinfectives for systemic use	161
.Dental	337
AMPHOTERICIN	
.Alimentary tract and metabolism	77
.Antiinfectives for systemic use	175
.Dental	333
AMPICILLIN	
.Antiinfectives for systemic use	162
.Dental	338
<i>Amprace 5 (AD)</i>	122
<i>Amprace 10 (AD)</i>	123
<i>Amprace 20 (AD)</i>	123
AMPRENAVIR	
.Section 100	360
<i>Anafranil 25 (NV)</i>	274, 275
ANAKINRA	205
<i>Anamorph (FM)</i>	
.Dental	351
.Nervous system	252
<i>Anandron (AV)</i>	192
<i>Anaprox 550 (RO)</i>	
.Dental	349
.Musculo-skeletal system	241
.Palliative Care	328
ANASTROZOLE	192
<i>Andriol Testocaps (OR)</i>	142
<i>Androcur (SC)</i>	
.Antineoplastic and immunomodulating agents	192
.Genito urinary system and sex hormones	151, 152
<i>Androcur-100 (SC)</i>	
.Antineoplastic and immunomodulating agents	192
.Genito urinary system and sex hormones	152
<i>Androderm (MX)</i>	142
<i>Anginine Stabilised (SI)</i>	
.Cardiovascular system	110
.Dental	335
<i>Angiomax (CS)</i>	105
<i>Anpec 40 (AF)</i>	120
<i>Anpec 80 (AF)</i>	120
<i>Anpec SR (AF)</i>	120
<i>Anselol 50 mg (DP)</i>	116
ANTAZOLINE with NAPHAZOLINE	
.Repatriation Schedule	497
<i>Antenex 2 (AF)</i>	
.Dental	355
.Nervous system	272
.Palliative Care	330
<i>Antenex 5 (AF)</i>	
.Dental	355
.Nervous system	273
.Palliative Care	330
<i>Anthel 125 (AF)</i>	290
<i>Anthel 250 (AF)</i>	290
<i>Antistine-Privine (NV)</i>	
.Repatriation Schedule	497
<i>Antroquoril (EX)</i>	136
<i>Anusol (PC)</i>	
.Repatriation Schedule	476
<i>Anzatax (MX)</i>	185
<i>Anzemet (AV)</i>	84
<i>Apomine (MX)</i>	
.Section 100	360
APOMORPHINE HYDROCHLORIDE	
.Section 100	360
<i>Apoven 250 (DP)</i>	296
<i>Apoven 500 (DP)</i>	297
APRACLOPIDINE HYDROCHLORIDE	302
APREPITANT	86
<i>Aquacare H.P. (AG)</i>	
.Repatriation Schedule	479
<i>Aquacel 177902 (CC)</i>	
.Repatriation Schedule	509
<i>Aquacel 177903 (CC)</i>	
.Repatriation Schedule	510
<i>Aquacel 177904 (CC)</i>	
.Repatriation Schedule	508
<i>Aquaclear (HR)</i>	
.Repatriation Schedule	511
<i>Aquae (HA)</i>	
.Palliative Care	319, 320
.Repatriation Schedule	472
<i>Aquasun Lotion SPF18 (PF)</i>	
.Repatriation Schedule	480
<i>Arabloc (HP)</i>	236
<i>Aranesp (AN)</i>	
.Section 100	371, 372
<i>Aratac 100 (AF)</i>	109
<i>Aratac 200 (AF)</i>	109
<i>Arava (AV)</i>	235, 236

GENERIC /PROPRIETARY INDEX

<i>Aredia 15 mg (NV)</i>		ATENOLOL.....	116
.Musculo-skeletal system	245	ATORVASTATIN CALCIUM	129
.Section 100	373	ATOVAQUONE	289
<i>Aredia 30 mg (NV)</i>		<i>Atrauman 499513 (HR)</i>	
.Musculo-skeletal system	245	.Repatriation Schedule	511
.Section 100	373	ATROPINE SULFATE	
<i>Aredia 90 mg (NV)</i>		.Alimentary tract and metabolism.....	84
.Section 100	373	.Dental.....	333
<i>Aricept (PF)</i>	283	.Doctor's Bag Supplies	71
<i>Arima (AF)</i>	279	.Sensory organs	304
<i>Arima 300 (AF)</i>	279	<i>Atropt (SI)</i>	304
<i>Arimidex (AP)</i>	192	<i>Atrovent (BY)</i>	296, 297
ARIPRAZOLE	270	<i>Atrovent Adult (BY)</i>	297
<i>Aristocort 0.02% (SI)</i>	136	<i>Atrovent Nasal Aqueous (BY)</i>	
<i>Arixtra (GK)</i>	105	.Repatriation Schedule	495
<i>Aromasin (PH)</i>	193	<i>Atrovent Nasal Forte (BY)</i>	
<i>Aropax (GK)</i>	277	.Repatriation Schedule	495
<i>Arsorb 60 (AW)</i>	111	<i>Attenta (AF)</i>	280
<i>Artane (SI)</i>	266	<i>Augmentin (GK)</i>	
<i>Arthrexin (AF)</i>		.Anti-infectives for systemic use	165
.Dental.....	347	.Dental.....	341
.Musculo-skeletal system	238	<i>Augmentin Duo (GK)</i>	
.Palliative Care	326	.Anti-infectives for systemic use	165
<i>Arthrotec 50 (PH)</i>		.Dental.....	340
.Repatriation Schedule.....	491	<i>Augmentin Duo 400 (GK)</i>	
<i>Asasantin SR (BY)</i>	103	.Anti-infectives for systemic use	166
<i>Ascensia Elite (BN)</i>	310	.Dental.....	341
<i>Ascensia Glucodisc (BN)</i>	310	<i>Augmentin Duo forte (GK)</i>	
ASCORBIC ACID		.Anti-infectives for systemic use	165
.Repatriation Schedule.....	475	.Dental.....	340
<i>Asig (SI)</i>	124	AURANOFIN.....	242
<i>Asmol 2.5 uni-dose (AF)</i>		<i>Aurorix (RO)</i>	279
.Doctor's Bag Supplies	72	<i>Aurorix 300 mg (RO)</i>	279
.Respiratory system.....	292	<i>Auscap (SI)</i>	277
<i>Asmol 5 uni-dose (AF)</i>		<i>Ausfam 20 (AW)</i>	78
.Doctor's Bag Supplies	73	<i>Ausfam 40 (AW)</i>	79
.Respiratory system.....	292	<i>Ausgem (SI)</i>	132
<i>Asmol CFC-free (AL)</i>		<i>Auspril (SI)</i>	122, 123
.Doctor's Bag Supplies	72	<i>Ausran (SI)</i>	79, 80
.Respiratory system.....	291	<i>Austrapen (CS)</i>	
<i>Aspalgin (FM)</i>		.Anti-infectives for systemic use	162
.Repatriation Schedule.....	492	.Dental.....	338
<i>Aspen Ampicyn (AS)</i>		<i>Avandia (GK)</i>	98
.Anti-infectives for systemic use	162	<i>Avanza (BP)</i>	280
.Dental.....	338	<i>Avanza SolTab (BP)</i>	280
<i>Aspen Flucil (AS)</i>		<i>Avapro (BQ)</i>	126
.Anti-infectives for systemic use	164	<i>Avapro HCT 150/12.5 (BQ)</i>	127
.Dental.....	339	<i>Avapro HCT 300/12.5 (BQ)</i>	127
ASPIRIN		<i>Avelox (BN)</i>	
.Blood and blood forming organs	102	.Anti-infectives for systemic use	173
.Dental.....	354	.Repatriation Schedule	487
.Nervous system.....	258	<i>Avonex (BD)</i>	195
.Repatriation Schedule.....	475	<i>Axit 30 (AF)</i>	280
<i>Astrix (MX)</i>		<i>Azahexal (HX)</i>	237
.Blood and blood forming organs	102	<i>Azamun (DP)</i>	237
.Repatriation Schedule.....	475	<i>Azapin (AW)</i>	237
<i>Atacand (AP)</i>	126	AZATHIOPRINE	237
<i>Atacand Plus 16/12.5 (AP)</i>	126		
ATAZANAVIR SULFATE			
.Section 100	360		
<i>Atehexal (HX)</i>	116		

GENERIC /PROPRIETARY INDEX

AZITHROMYCIN	
.Antiinfectives for systemic use	170
.Repatriation Schedule	487
.Section 100	360
.Sensory organs	300
<i>Azol 100 (AF)</i>	152
<i>Azol 200 (AF)</i>	152
<i>Azopt (AQ)</i>	303
B	
<i>Baclo (DP)</i>	243
BACLOFEN	
.Musculo-skeletal system	243
.Section 100	360
<i>Baclohexal (HX)</i>	243
<i>Bactigras 7457 (SN)</i>	
.Repatriation Schedule	507
<i>Bactrim (RO)</i>	
.Antiinfectives for systemic use	170
.Dental	344
<i>Bactrim DS (RO)</i>	
.Antiinfectives for systemic use	170
.Dental	344
<i>Bactroban (GK)</i>	
.Repatriation Schedule	480
BALSALAZIDE SODIUM	91
BANDAGE—ABSORBENT WOOL	
.Repatriation Schedule	501
BANDAGE—CALICO	
.Repatriation Schedule	501
BANDAGE—COMPRESSION	
.Repatriation Schedule	501
BANDAGE—RETENTION—COHESIVE—HEAVY	
.Repatriation Schedule	502
BANDAGE—RETENTION—COHESIVE—LIGHT	
.Repatriation Schedule	502
BANDAGE—RETENTION—COTTON CREPE	
.Repatriation Schedule	503
BANDAGE—TUBULAR	
.Repatriation Schedule	503
BANDAGE—TUBULAR (FINGER)	
.Repatriation Schedule	503
BANDAGE—TUBULAR (LIGHTWEIGHT)	
.Repatriation Schedule	503
BANDAGE—TUBULAR (LONG STOCKING)	
.Repatriation Schedule	503
BANDAGE—TUBULAR (SHORT STOCKING)	
.Repatriation Schedule	503
BANDAGE—ZINC PASTE	
.Repatriation Schedule	504
<i>Barbloc 5 (AF)</i>	116
<i>Barbloc 15 (AF)</i>	116
BCG IMMUNOTHERAPEUTIC (Bacillus Calmette-Guérin/ Connaught strain)	
Guérin/ Connaught strain)	196
BCG-TICE (Bacillus Calmette-Guérin/ Tice strain)	
Tice strain)	196
BECLOMETHASONE DIPROPIONATE	
294	
<i>BenPen (CS)</i>	
.Antiinfectives for systemic use	163
.Dental	338
.Doctor's Bag Supplies	71
BENZATHINE PENICILLIN	
.Antiinfectives for systemic use	163
.Dental	338
<i>Benzemul (MG)</i>	290
BENZHEXOL HYDROCHLORIDE	
266	
<i>Benztrop (PL)</i>	266
BENZTROPINE MESYLATE	
.Dental	355
.Doctor's Bag Supplies	71
.Nervous system	266
BENZYDAMINE HYDROCHLORIDE	
.Alimentary tract and metabolism	77
.Dental	333
BENZYL BENZOATE	
290	
BENZYL PENICILLIN	
.Antiinfectives for systemic use	163
.Dental	338
.Doctor's Bag Supplies	71
<i>Betachek (NA)</i>	310
<i>Betachek MERIDIAN (NA)</i>	310
<i>Betadine (FH)</i>	
.Repatriation Schedule	507
<i>Betadine Antiseptic Liquid (FH)</i>	
.Repatriation Schedule	482
<i>Betaferon (SC)</i>	195
<i>Betagan (AG)</i>	303
<i>Betaloc (AP)</i>	117
BETAMETHASONE ACETATE with	
BETAMETHASONE SODIUM PHOSPHATE	
.Dental	335
.Systemic hormonal preparations, excl. sex hormones and insulins	155
BETAMETHASONE DIPROPIONATE	
136	
BETAMETHASONE VALERATE	
.Dermatologicals	136
.Repatriation Schedule	481
<i>Betamin (AV)</i>	
.Repatriation Schedule	475
BETAXOLOL HYDROCHLORIDE	303
BETHANECHOL CHLORIDE	287
<i>Betnovate (SI)</i>	
.Repatriation Schedule	481
<i>Betnovate 1/5 (SI)</i>	136
<i>Betnovate 1/2 (SI)</i>	
.Dermatologicals	136
.Repatriation Schedule	481
<i>Betoptic (AQ)</i>	303
<i>Betoptic S (AQ)</i>	303
<i>BetoQuin (IQ)</i>	303
<i>Bgramin (DP)</i>	
.Antiinfectives for systemic use	161, 162
.Dental	337, 338
<i>Biatain Adhesive 3420 (CT)</i>	
.Repatriation Schedule	507
<i>Biatain Adhesive 3423 (CT)</i>	
.Repatriation Schedule	507
<i>Biatain Non-adhesive 3410 (CT)</i>	
.Repatriation Schedule	507
<i>Biatain Non-adhesive 3413 (CT)</i>	
.Repatriation Schedule	507
<i>Biassig (HP)</i>	171

GENERIC /PROPRIETARY INDEX

BICALUTAMIDE.....	191	BSN 2902165 (BV)	
<i>Bicillin L-A Tubex (AS)</i>		.Repatriation Schedule	511
.Antiinfectives for systemic use.....	163	<i>Budamax Aqueous (PM)</i>	
.Dental.....	338	.Repatriation Schedule	495
<i>Bicor (AL)</i>	116	BUDESONIDE	
BIFONAZOLE		.Repatriation Schedule	495
.Repatriation Schedule.....	477	.Respiratory system	295
BIMATOPROST	304	BUDESONIDE with EFORMOTEROL FUMARATE	
<i>Biodone Forte (MW)</i>		DIHYDRATE	293
.Section 100	428	BUPRENORPHINE	
<i>Bion Tears (AQ)</i>	306	.Nervous system	257
BIPERIDEN HYDROCHLORIDE.....	266	BUPRENORPHINE HYDROCHLORIDE	
BISACODYL		.Section 100	427
.Alimentary tract and metabolism.....	87, 88	BUPROPION HYDROCHLORIDE.....	287
.Palliative Care.....	321, 323	<i>Buscopan (BY)</i>	
<i>Bisalax (AS)</i>		.Palliative Care.....	320
.Alimentary tract and metabolism.....	87, 88	.Repatriation Schedule	473
.Palliative Care.....	321, 323	<i>Buspar (BQ)</i>	
BISOPROLOL FUMARATE	116	.Repatriation Schedule	493
BIVALIRUDIN TRIFLUOROACETATE	105	BUSPIRONE HYDROCHLORIDE	
<i>Blenamax (SI)</i>		.Repatriation Schedule	493
.Special Pharmaceutical Benefit	74	BUSULFAN.....	181
<i>Blenoxane (BQ)</i>		<i>Butamol 2.5 (AW)</i>	
.Special Pharmaceutical Benefit	74	.Doctor's Bag Supplies	72
BLEOMYCIN SULFATE		.Respiratory system	292
.Special Pharmaceutical Benefit	74	<i>Butamol 5 (AW)</i>	
<i>Bleph 10 (AG)</i>	300	.Doctor's Bag Supplies	73
<i>Bonefos (SC)</i>	246	.Respiratory system	292
<i>Bonefos 800 mg (SC)</i>	246	C	
BOSENTAN MONOHYDRATE		<i>Cabaser (PU)</i>	267
.Section 100	361	CABERGOLINE	
<i>Botox (AG)</i>		.Genito urinary system and sex hormones	139
.Section 100	424	.Nervous system	267
BOTULINUM TOXIN TYPE A PURIFIED		<i>Caelyx (SH)</i>	
NEUROTOXIN COMPLEX		.Antineoplastic and immunomodulating agents	186
.Section 100	424	.Section 100	374
<i>Brevinor (PH)</i>	140	CALCIPOTRIOL	134
<i>Brevinor-1 (PH)</i>	140	CALCITRIOL	
<i>Bricanyl (AP)</i>		.Alimentary tract and metabolism.....	99
.Doctor's Bag Supplies	73	.Musculo-skeletal system	247
.Respiratory system.....	298	<i>Calcitriol-DP (DP)</i>	
<i>Bricanyl Respules (AP)</i>	293	.Alimentary tract and metabolism.....	99
<i>Bricanyl Turbuhaler (AP)</i>	293	.Musculo-skeletal system	247
BRIMONIDINE TARTRATE.....	302	CALCIUM.....	99
BRIMONIDINE TARTRATE with TIMOLOL		CALCIUM CARBONATE with GLYCINE	
MALEATE	302	.Repatriation Schedule	472
BRINZOLAMIDE	303	CALCIUM FOLINATE.....	308
<i>BrinzoQuin (IQ)</i>	303	<i>Calmurid (OL)</i>	
BROMAZEPAM		.Repatriation Schedule	479
.Repatriation Schedule.....	493	<i>Cal-Sup (MM)</i>	99
BROMOCRIPTINE MESYLATE		<i>Caltrate (WT)</i>	99
.Genito urinary system and sex hormones	139	<i>Campral (AF)</i>	287
.Nervous system.....	267	<i>Camptosar (PU)</i>	189
<i>Bromohexal (HX)</i>		CANDESARTAN CILEXETIL.....	126
.Genito urinary system and sex hormones	139	CANDESARTAN CILEXETIL with	
.Nervous system.....	267	HYDROCHLOROTHIAZIDE.....	126
<i>Brufen (AB)</i>		<i>Canesten (BN)</i>	
.Dental.....	348	.Repatriation Schedule	477, 483, 484
.Musculo-skeletal system	240	<i>Canesten 1 (BN)</i>	
.Palliative Care.....	327	.Repatriation Schedule	483

GENERIC /PROPRIETARY INDEX

<i>Canesten 3 (BN)</i>		CEFACLOR	
.Repatriation Schedule	484	.Anti-infectives for systemic use	168
CAPECITABINE	183	.Dental	342
<i>Capoten (BQ)</i>	121, 122	<i>Cefazolin Sandoz (SZ)</i>	167
<i>Caprilon (SB)</i>	313	CEFEPIME	169
<i>Captohexal (HX)</i>	121, 122	<i>Cefkor CD (DP)</i>	
CAPTOPRIL	121, 122	.Anti-infectives for systemic use	168
<i>Carafate (AS)</i>	84	.Dental	342
CARBACHOL	302	CEFOTAXIME	
CARBAMAZEPINE		.Anti-infectives for systemic use	168
.Dental	355	.Dental	343
.Nervous system	262	<i>Cefotaxime Sandoz (SZ)</i>	
<i>Carbamazepine-BC (BG)</i>		.Anti-infectives for systemic use	168
.Dental	355	.Dental	343
.Nervous system	262, 263	CEFTRIAXONE	169
<i>Carbamazepine Sandoz (SZ)</i>		<i>Ceftriaxone Sandoz (SZ)</i>	169
.Dental	355	CEFUROXIME AXETIL	
.Nervous system	262, 263	.Anti-infectives for systemic use	168
CARBAMIDE PEROXIDE		.Dental	343
.Repatriation Schedule	498	<i>Celapram (AF)</i>	276
CARBIMAZOLE	158	<i>Celebrex (PH)</i>	241
<i>CarboFLEX 403202 (CC)</i>		CELECOXIB	241
.Repatriation Schedule	504	<i>Celestone Chronodose (SH)</i>	
<i>CarboFLEX 403204 (CC)</i>		.Dental	335
.Repatriation Schedule	504	.Systemic hormonal preparations, excl. sex	
CARBOHYDRATE, FAT, VITAMINS, MINERALS		hormones and insulins	155
and TRACE ELEMENTS	317	<i>Celestone-M (SH)</i>	136
<i>Carbohydrate FreeMixture (SB)</i>	317	<i>CellCept (RO)</i>	
CARBOMER 974	304	.Antineoplastic and immunomodulating agents	236
CARBOMER 980	305	.Section 100	414
CARBOPLATIN	187	<i>Cellufresh (AG)</i>	305
<i>Cardinorm (HX)</i>	109	<i>Celluvisc (AG)</i>	305
<i>Cardiprin 100 (RC)</i>		CEPHALEXIN	
.Repatriation Schedule	475	.Anti-infectives for systemic use	166
<i>Cardizem (AV)</i>	120	.Dental	341
<i>Cardizem CD (AV)</i>	121	CEPHALOTHIN	
<i>Cardol (AF)</i>	109	.Anti-infectives for systemic use	167
<i>CareSens (LB)</i>	310	.Dental	342
CARMELLOSE SODIUM		CEPHAZOLIN	167
.Palliative Care	319	<i>Certican (NV)</i>	
.Sensory organs	305	.Antineoplastic and immunomodulating agents	235
.Repatriation Schedule	472	.Section 100	378
CARMELLOSE SODIUM with PECTIN and GELATIN		<i>Cerumol (AC)</i>	
.Repatriation Schedule	479	.Repatriation Schedule	498
<i>Cartia (GK)</i>		CETIRIZINE HYDROCHLORIDE	
.Repatriation Schedule	475	.Repatriation Schedule	496
CARVEDILOL	118	<i>C-Flox 250 (AL)</i>	172
<i>Catapres (BY)</i>	112	<i>C-Flox 500 (AL)</i>	172
<i>Catapres 100 (BY)</i>	112	<i>C-Flox 750 (AL)</i>	172
CATIONIC CONDITIONER with PANTHENOL		<i>Chem mart Aciclovir (CH)</i>	177
.Repatriation Schedule	483	<i>Chem mart Allopurinol (CH)</i>	244
<i>Caverject Impulse (PH)</i>		<i>Chem mart Alprazolam (CH)</i>	272
.Repatriation Schedule	485	<i>Chem mart Amiodarone (CH)</i>	109
<i>Cavicare 4563 (SN)</i>		<i>Chem mart Amoxycillin (CH)</i>	
.Repatriation Schedule	507	.Anti-infectives for systemic use	161, 162
<i>Ceclor (AS)</i>		.Dental	337, 338
.Anti-infectives for systemic use	168	<i>Chem mart Amoxycillin and Clavulanic Acid (CH)</i>	
.Dental	343	.Anti-infectives for systemic use	165
<i>Ceclor CD (AS)</i>		.Dental	340
.Anti-infectives for systemic use	168	<i>Chem mart Atenolol (CH)</i>	116
.Dental	342	<i>Chem mart Baclofen (CH)</i>	243

GENERIC /PROPRIETARY INDEX

<i>Chem mart Captopril (CH)</i>	121, 122	<i>Chem mart Tramadol (CH)</i>	
<i>Chem mart Cefaclor (CH)</i>		.Dental.....	353
.Antiinfectives for systemic use.....	168	.Nervous system.....	257, 258
.Dental.....	343	<i>Chem mart Trimethoprim with Sulfamethoxazole DS (CH)</i>	
<i>Chem mart Cefaclor CD (CH)</i>		.Antiinfectives for systemic use.....	170
.Antiinfectives for systemic use.....	168	.Dental.....	344
.Dental.....	342	CHLORAMBUCIL.....	181
<i>Chem mart Cephalexin (CH)</i>		CHLORAMPHENICOL	
.Antiinfectives for systemic use.....	166, 167	.Dental.....	356
.Dental.....	341, 342	.Sensory organs.....	300, 307
<i>Chem mart Citalopram (CH)</i>	276	CHLORHEXIDINE GLUCONATE	
<i>Chem mart Clarithromycin (CH)</i>	170	.Repatriation Schedule.....	472
<i>Chem mart Clomipramine (CH)</i>	274, 275	<i>Chloromycetin (PF)</i>	
<i>Chem mart Clotrimazole 3 Day Cream (CH)</i>		.Dental.....	356
.Repatriation Schedule.....	484	.Sensory organs.....	300, 307
<i>Chem mart Clotrimazole 6 Day Cream (CH)</i>		CHLORPROMAZINE HYDROCHLORIDE	
.Repatriation Schedule.....	484	.Doctor's Bag Supplies.....	71
<i>Chem mart Diclofenac (CH)</i>		.Nervous system.....	268
.Dental.....	346	<i>Chlorsig (SI)</i>	
.Musculo-skeletal system.....	238	.Dental.....	356
.Palliative Care.....	324, 325	.Sensory organs.....	300
<i>Chem mart Diltiazem (CH)</i>	120	CHLORTHALIDONE.....	113
<i>Chem mart Diltiazem CD (CH)</i>	121	<i>Chlorvescent (AS)</i>	99
<i>Chem mart Doxycycline (CH)</i>		CHOLESTYRAMINE.....	132
.Antiinfectives for systemic use.....	159, 160	<i>Cialis (LY)</i>	
.Dental.....	336	.Repatriation Schedule.....	485
<i>Chem mart Enalapril (CH)</i>	122, 123	<i>Ciazil (DP)</i>	276
<i>Chem mart Famotidine (CH)</i>	78, 79	CICLESONIDE.....	295
<i>Chem mart Fluoxetine (CH)</i>	277	CICLOPIROX OLAMINE	
<i>Chem mart Frusemide (CH)</i>	114	.Repatriation Schedule.....	478
<i>Chem mart Gemfibrozil (CH)</i>	132	<i>Cicloral (HX)</i>	
<i>Chem mart Gliclazide (CH)</i>	96	.Antineoplastic and immunomodulating agents.....	213
<i>Chem mart Indapamide (CH)</i>	113	.Section 100.....	371
<i>Chem mart Ipratropium (CH)</i>	296, 297	CIDOFOVIR	
<i>Chem mart Isosorbide Mononitrate (CH)</i>	111	.Section 100.....	370
<i>Chem mart Isotretinoin (CH)</i>	137	<i>Cilamox (SI)</i>	
<i>Chem mart Lisinopril (CH)</i>	123, 124	.Antiinfectives for systemic use.....	161, 162
<i>Chem mart Metformin (CH)</i>	95	.Dental.....	337, 338
<i>Chem mart Metoprolol (CH)</i>	117	<i>Cilex (DP)</i>	
<i>Chem mart Moclobemide (CH)</i>	279	.Antiinfectives for systemic use.....	166, 167
<i>Chem mart Nifedipine (CH)</i>	119	.Dental.....	341, 342
<i>Chem mart Norfloxacin (CH)</i>	173	<i>Cilicaine (SI)</i>	
<i>Chem mart Oral Rehydration Salts (CH)</i>	90	.Antiinfectives for systemic use.....	163
<i>Chem mart Paroxetine (CH)</i>	277	.Dental.....	339
<i>Chem mart Piroxicam (CH)</i>		.Doctor's Bag Supplies.....	71
.Dental.....	347, 348	<i>Cilicaine V (FM)</i>	
.Musculo-skeletal system.....	240	.Antiinfectives for systemic use.....	163
<i>Chem mart Piroxicam Dispersible (CH)</i>		.Dental.....	339
.Dental.....	347	<i>Cilicaine VK (FM)</i>	
.Musculo-skeletal system.....	239	.Antiinfectives for systemic use.....	163
<i>Chem mart Prazosin (CH)</i>	112, 113	.Dental.....	338
<i>Chem mart Ranitidine (CH)</i>	79, 80	<i>Cilopen VK (DP)</i>	
<i>Chem mart Salbutamol (CH)</i>		.Antiinfectives for systemic use.....	163
.Doctor's Bag Supplies.....	72, 73	.Dental.....	338
.Respiratory system.....	292	<i>CiloQuin (IQ)</i>	301
<i>Chem mart Sertraline (CH)</i>	278	<i>Ciloxan (AQ)</i>	301
<i>Chem mart Simvastatin (CH)</i>	130, 131	<i>Cimehexal (HX)</i>	77, 78
<i>Chem mart Sotalol (CH)</i>	109	CIMETIDINE	
<i>Chem mart Tamoxifen (CH)</i>	191	.Alimentary tract and metabolism.....	77, 78
		.Repatriation Schedule.....	472

GENERIC /PROPRIETARY INDEX

<i>Cipramil (LU)</i>	276	<i>Clavulin Duo (AW)</i>	
CIPROFLOXACIN		.Antiinfectives for systemic use	165
.Antiinfectives for systemic use	172	.Dental	340
.Sensory organs	301	<i>Clavulin Duo 400 (AW)</i>	
CIPROFLOXACIN HYDROCHLORIDE with		.Antiinfectives for systemic use	166
HYDROCORTISONE		.Dental	341
.Repatriation Schedule	498	<i>Clavulin Duo Forte (AW)</i>	
<i>Ciprofloxacin-BC (BG)</i>	172	.Antiinfectives for systemic use	165
<i>Ciprol 250 (AW)</i>	172	.Dental	340
<i>Ciprol 500 (AW)</i>	172	<i>Cleocin (KR)</i>	
<i>Ciprol 750 (AW)</i>	172	.Antiinfectives for systemic use	171
<i>Ciproxin 250 (BN)</i>	172	.Dental	345
<i>Ciproxin 500 (BN)</i>	172	<i>Clexane (AV)</i>	102
<i>Ciproxin 750 (BN)</i>	172	<i>Climara 25 (SC)</i>	143
<i>Ciproxin HC (AQ)</i>		<i>Climara 50 (SC)</i>	144
.Repatriation Schedule	498	<i>Climara 75 (SC)</i>	144
CISPLATIN	187	<i>Climara 100 (SC)</i>	144
CITALOPRAM HYDROBROMIDE	276	<i>Climen (SC)</i>	148
<i>Citracal (KY)</i>	99	CLINDAMYCIN	
<i>Citralite (MM)</i>		.Antiinfectives for systemic use	171
.Repatriation Schedule	486	.Dental	345
<i>Citravescent Sachets (MM)</i>		<i>Clinistix (BN)</i>	309
.Repatriation Schedule	486	<i>Clinitest (BN)</i>	309
<i>Citrihexal (HX)</i>		<i>Clobemix (DP)</i>	279
.Alimentary tract and metabolism	99	<i>Clofeme (HX)</i>	
.Musculo-skeletal system	247	.Repatriation Schedule	483
CLADRIBINE	182	<i>Clofen 10 (AF)</i>	243
<i>Clamohexal 125mg/31.25mg/5mL (HX)</i>		<i>Clofen 25 (AF)</i>	243
.Antiinfectives for systemic use	165	<i>Clomhexal (HX)</i>	151
.Dental	341	<i>Clomid (AV)</i>	151
<i>Clamohexal Duo 400mg/57mg/5mL (HX)</i>		CLOMIPHENE CITRATE	151
.Antiinfectives for systemic use	166	CLOMIPRAMINE HYDROCHLORIDE	274, 275
.Dental	341	<i>Clonac 25 (AW)</i>	
<i>Clamohexal Duo 500mg/125mg (HX)</i>		.Dental	346
.Antiinfectives for systemic use	165	.Musculo-skeletal system	238
.Dental	340	.Palliative Care	324, 325
<i>Clamohexal Duo Forte 875mg/125mg (HX)</i>		<i>Clonac 50 (AW)</i>	
.Antiinfectives for systemic use	165	.Dental	346
.Dental	340	.Musculo-skeletal system	238
<i>Clamoxyl (AL)</i>		.Palliative Care	325
.Antiinfectives for systemic use	165	CLONAZEPAM	
.Dental	341	.Nervous system	262
<i>Clamoxyl Duo (AL)</i>		.Palliative Care	329
.Antiinfectives for systemic use	165	<i>Clonea (AF)</i>	
.Dental	340	.Repatriation Schedule	477
<i>Clamoxyl Duo 400 (AL)</i>		CLONIDINE	112
.Antiinfectives for systemic use	166	CLOPIDOGREL HYDROGEN SULFATE	
.Dental	341	.Blood and blood forming organs	103
<i>Clamoxyl Duo forte (AL)</i>		.Repatriation Schedule	475
.Antiinfectives for systemic use	165	<i>Clopine 25 (MX)</i>	
.Dental	340	.Section 100	370
<i>Clarac (DP)</i>	170	<i>Clopine 50 (MX)</i>	
<i>Claratyne (SH)</i>		.Section 100	370
.Repatriation Schedule	497	<i>Clopine 100 (MX)</i>	
<i>Clarihexal (HX)</i>	170	.Section 100	370
CLARITHROMYCIN		<i>Clopine 200 (MX)</i>	
.Antiinfectives for systemic use	170	.Section 100	370
.Section 100	370	<i>Clopixol Depot (LU)</i>	269
<i>Clavulin (AW)</i>			
.Antiinfectives for systemic use	165		
.Dental	341		

GENERIC /PROPRIETARY INDEX

CLOSTRIDIUM BOTULINUM TYPE A TOXIN— HAEMAGGLUTININ COMPLEX .Section 100.....	424	Combivent (BY) .Repatriation Schedule.....	496
CloSyn (ZT) .Section 100.....	370	Combivir (GK) .Section 100.....	411
CLOTTRIMAZOLE .Repatriation Schedule.....	477, 483	Comfeel Paste 4701 (CT) .Repatriation Schedule.....	508
CLOZAPINE .Section 100.....	370	Comfeel Plus Pressure Relieving 3350 (CT) .Repatriation Schedule.....	512
Clozaril 25 (NV) .Section 100.....	370	Comfeel Plus Pressure Relieving 3353 (CT) .Repatriation Schedule.....	512
Clozaril 100 (NV) .Section 100.....	370	Comfeel Plus Transparent 3530 (CT) .Repatriation Schedule.....	509
COAL TAR - PREPARED.....	134	Comfeel Plus Transparent 3533 (CT) .Repatriation Schedule.....	509
Coban 1584 (MM) .Repatriation Schedule.....	502	Comfeel Plus Transparent 3536 (CT) .Repatriation Schedule.....	509
Codalgin (FM) .Repatriation Schedule.....	492	Comfeel Plus Ulcer Dressing 3110 (CT) .Repatriation Schedule.....	510
Codalgin Forte (FM) .Dental.....	349	Comfeel Powder 4706 (CT) .Repatriation Schedule.....	508
.Nervous system.....	249	Comfeel Purilon Gel 3900 (CT) .Repatriation Schedule.....	510
Codapane Forte (AL) .Dental.....	349	Comfeel SeaSorb Dressing 3705 (CT) .Repatriation Schedule.....	505
.Nervous system.....	249	Comfeel SeaSorb Dressing 3710 (CT) .Repatriation Schedule.....	505
CODEINE PHOSPHATE .Dental.....	349	Comfeel SeaSorb Filler 3740 (CT) .Repatriation Schedule.....	504
.Nervous system.....	249	Comprilan 1027 (BV) .Repatriation Schedule.....	501
.Respiratory system.....	299	Comtan (NV).....	268
CODEINE PHOSPHATE with ASPIRIN .Repatriation Schedule.....	492	Concorz (HX).....	278
CODEINE PHOSPHATE with PARACETAMOL .Dental.....	349	Condyline Paint (HA) .Repatriation Schedule.....	481
.Nervous system.....	249	Copaxone (AV).....	196
.Repatriation Schedule.....	492	Copius 3629 (BV) .Repatriation Schedule.....	502
Cogentin (MK) .Dental.....	355	COPPER SULFATE.....	309
.Doctor's Bag Supplies.....	71	Coras (AF).....	120
.Nervous system.....	266	Corbeton 20 (AF).....	116
Colazide (PK).....	91	Corbeton 40 (AF).....	116
COLCHICINE.....	244	Cordarone X 100 (SW).....	109
Colese (AF) .Repatriation Schedule.....	473	Cordarone X 200 (SW).....	109
Colestid (PH).....	132	Cordilox 180 SR (KN).....	120
COLESTIPOL HYDROCHLORIDE.....	132	Cordilox SR (KN).....	120
Colgout (AS).....	244	Cortate (AS).....	155
Colifoam (GC).....	90	Cortef (DT) .Dental.....	335
Colofac (SM) .Repatriation Schedule.....	473	.Dermatologicals.....	135
Coloxyl (FM) .Alimentary tract and metabolism.....	87	Cortic-DS 1% (FM) .Dental.....	335
.Palliative Care.....	322	.Dermatologicals.....	135
Coloxyl 50 (FM) .Repatriation Schedule.....	473	CORTISONE ACETATE.....	155
Coloxyl with Senna (FM) .Repatriation Schedule.....	473	Cortival 1/5 (FM).....	136
CombiDERM 651027 (CC) .Repatriation Schedule.....	506	Cortival 1/2 (FM).....	136
CombiDERM 651031 (CC) .Repatriation Schedule.....	506	Cosopt (MK).....	303
Combigan (AG).....	302	Cosudex (AP).....	191
		Cotazym-S Forte (OR).....	93
		COTTON WOOL ROLL .Repatriation Schedule.....	504
		Coumadin (SI).....	101

GENERIC /PROPRIETARY INDEX

<i>Coversyl (SE)</i>	124	<i>Cysporin (MX)</i>	
<i>Coversyl Plus 4/1.25 (SE)</i>	125	.Antineoplastic and immunomodulating agents	213
<i>Creon (SM)</i>	93	.Section 100	371
<i>Creon 5000 (SM)</i>	93	<i>Cytadren 250 (NV)</i>	
<i>Creon Forte (SM)</i>	93	.Antineoplastic and immunomodulating agents	192
<i>Crinone 8% (SG)</i>		.Systemic hormonal preparations, excl. sex	
.Section 100	427	hormones and insulins	157
<i>Crixivan 100 mg (MK)</i>		CYTARABINE	183
.Section 100	389	<i>Cytotec (PH)</i>	80
<i>Crixivan 200 mg (MK)</i>		D	
.Section 100	389	<i>Daivonex (CS)</i>	134
<i>Crixivan 400 mg (MK)</i>		<i>Daktarin (JC)</i>	
.Section 100	389	.Repatriation Schedule	477
<i>Crysanal (MD)</i>		<i>Dalacin C (PH)</i>	
.Dental	349	.Antifectives for systemic use	171
.Musculo-skeletal system	241	.Dental	345
.Palliative Care	328	DALTEPARIN SODIUM (Low Molecular Weight	
<i>Curam 500/125 (SZ)</i>		Heparin Sodium—porcine mucous)	101
.Antifectives for systemic use	165	DANAZOL	152
.Dental	340	<i>Dan Gard (FH)</i>	
<i>Curam 875/125 (SZ)</i>		.Repatriation Schedule	482
.Antifectives for systemic use	165	<i>Dantrium (PU)</i>	243
.Dental	340	DANTROLENE SODIUM	243
<i>Curity 4112 (KE)</i>		<i>Daonil (AV)</i>	95
.Repatriation Schedule	507	<i>Dapa-Tabs (AF)</i>	113
<i>Cutifilm Plus 76308 (SN)</i>		DAPSONE	
.Repatriation Schedule	506	.Antifectives for systemic use	176
<i>Cutifilm Plus 76309 (SN)</i>		.Dermatologicals	137
.Repatriation Schedule	506	<i>Daraprim (GK)</i>	289
<i>Cutilin Non-Stick Wound Pad 76300 (SN)</i>		DARBEPOETIN ALFA	
.Repatriation Schedule	511	.Section 100	371
<i>Cutilin Non-Stick Wound Pad 76301 (SN)</i>		<i>DBL Aspirin 100 mg (FA)</i>	102
.Repatriation Schedule	511	<i>DBL Ceftriaxone (MX)</i>	169
<i>Cutinova Hydro 66047441 (SN)</i>		<i>DBL Doxycycline (FA)</i>	
.Repatriation Schedule	508	.Antifectives for systemic use	159, 160
<i>Cutinova Hydro 66047443 (SN)</i>		.Dental	336
.Repatriation Schedule	508	<i>DBL Erythromycin (FA)</i>	
<i>Cycloblastin (PH)</i>	181	.Antifectives for systemic use	171
CYCLOPHOSPHAMIDE	181	.Dental	344
CYCLOSPORIN		<i>DBL Gabapentin (MX)</i>	
.Antineoplastic and immunomodulating agents	213	.Nervous system	264
.Section 100	370	.Repatriation Schedule	493
<i>Cyklokapron (PH)</i>	106	<i>DBL Ipratropium (MX)</i>	296, 297
<i>Cymevene (RO)</i>		<i>Deca-Durabolin (OR)</i>	100
.Section 100	380	DEFERIPRONE	
CYPROHEPTADINE HYDROCHLORIDE	261	.Section 100	372
<i>Cyprohexal (HX)</i>		DELAVIRDINE MESYLATE	
.Antineoplastic and immunomodulating agents	192	.Section 100	372
.Genito urinary system and sex hormones	151, 152	<i>Depo-Medrol (PH)</i>	
<i>Cyprone (AF)</i>		.Dental	336
.Antineoplastic and immunomodulating agents	192	.Systemic hormonal preparations, excl. sex	
.Genito urinary system and sex hormones	151, 152	hormones and insulins	156
<i>Cyprostat (SY)</i>		<i>Depo-Nisalone (KR)</i>	
.Antineoplastic and immunomodulating agents	192	.Dental	336
.Genito urinary system and sex hormones	151, 152	.Systemic hormonal preparations, excl. sex	
<i>Cyprostat-100 (SY)</i>		hormones and insulins	156
.Antineoplastic and immunomodulating agents	192	<i>Depo-Provera (PH)</i>	141, 146
.Genito urinary system and sex hormones	152	<i>Depo-Ralovera (KR)</i>	141
CYPROTERONE ACETATE		<i>Deptran 10 (AF)</i>	275
.Antineoplastic and immunomodulating agents	192	<i>Deptran 25 (AF)</i>	275
.Genito urinary system and sex hormones	151	<i>Deptran 50 (AF)</i>	275

GENERIC /PROPRIETARY INDEX

<i>Deralin 10 (AF)</i>	116	DICLOFENAC SODIUM	
<i>Deralin 40 (AF)</i>	116	.Dental.....	346
<i>Deralin 160 (AF)</i>	116	.Musculo-skeletal system.....	238
<i>Dermestril 25 (MX)</i>	143	.Palliative Care.....	324
<i>Dermestril 50 (MX)</i>	144	.Sensory organs.....	301
<i>Dermestril 100 (MX)</i>	144	DICLOFENAC SODIUM with MISOPROSTOL	
<i>Deseril (NV)</i>	259	.Repatriation Schedule.....	491
<i>Desferal 500 mg (NV)</i>		<i>Diclofenac-BC (BG)</i>	
.Section 100.....	372	.Dental.....	346
<i>Desferal 2 g (NV)</i>		.Musculo-skeletal system.....	238
.Section 100.....	372	.Palliative Care.....	324, 325
DESFERRIOXAMINE MESYLATE		<i>Diclohexal (HX)</i>	
.Section 100.....	372	.Dental.....	346
DESMOPRESSIN ACETATE.....	154	.Musculo-skeletal system.....	238
DEXAMETHASONE		.Palliative Care.....	324, 325
.Sensory organs.....	301	DICLOXACILLIN	
.Systemic hormonal preparations, excl. sex		.Antiinfectives for systemic use.....	164
hormones and insulins.....	155	.Dental.....	339
DEXAMETHASONE with FRAMYCETIN SULFATE		<i>Dicloxsig (SI)</i>	
and GRAMICIDIN.....	307	.Antiinfectives for systemic use.....	164
DEXAMETHASONE SODIUM PHOSPHATE		.Dental.....	339
.Doctor's Bag Supplies.....	71	DIDANOSINE	
.Systemic hormonal preparations, excl. sex		.Section 100.....	372
hormones and insulins.....	155	<i>Didrocal (PU)</i>	247
DEXAMPHETAMINE SULFATE.....	280	<i>Didronel (PU)</i>	245
<i>Dexmethsone (AS)</i>	155	<i>Difflam (MM)</i>	
DEXTROPROPOXYPHENE NAPSYLATE		.Alimentary tract and metabolism.....	77
.Repatriation Schedule.....	492	.Dental.....	333
<i>Diabex (AL)</i>	95	<i>Diffucan (PF)</i>	175
<i>Diabex 850 (AL)</i>	95	DIFLUNISAL	
<i>Diabex 1000 (AL)</i>	95	.Dental.....	349
<i>Diaformin (AF)</i>	95	.Musculo-skeletal system.....	242
<i>Diaformin 850 (AF)</i>	95	<i>Digestelact (SJ)</i>	314
<i>Diaformin 1000 (AF)</i>	95	DIGOXIN.....	108
<i>Dialamine (SB)</i>	317	<i>Dihydergot (NV)</i>	
<i>Diamicon (SE)</i>	96	.Doctor's Bag Supplies.....	71
<i>Diamicon MR (SE)</i>	95	.Nervous system.....	259
<i>Diamox (SI)</i>	303	DIHYDROERGOTAMINE MESYLATE	
<i>Diastix (BN)</i>	309	.Doctor's Bag Supplies.....	71
DIAZEPAM		.Nervous system.....	259
.Dental.....	355	<i>Dilantin (PF)</i>	261
.Doctor's Bag Supplies.....	71	<i>Dilantin Infatabs (PF)</i>	261
.Nervous system.....	272	<i>Dilantin Sodium (PF)</i>	261
.Palliative Care.....	330	<i>Dilatrend 3.125 (RO)</i>	118
<i>Diazepam-DP (DP)</i>		<i>Dilatrend 6.25 (RO)</i>	118
.Dental.....	355	<i>Dilatrend 12.5 (RO)</i>	118
.Nervous system.....	273	<i>Dilatrend 25 (RO)</i>	118
.Palliative Care.....	330	<i>Dilatrend Titration Pack (RO)</i>	118
<i>Dibenyline (LM)</i>		<i>Dilaudid (AB)</i>	
.Cardiovascular system.....	115	.Dental.....	349, 350
.Genito urinary system and sex hormones.....	153	.Nervous system.....	249, 250
DICHLOROBENZENE with CHLORBUTOL and		<i>Dilaudid-HP (AB)</i>	
TURPENTINE OIL		.Dental.....	349
.Repatriation Schedule.....	498	.Nervous system.....	249
<i>Dicloil (BQ)</i>		<i>Diltahexal (HX)</i>	120
.Antiinfectives for systemic use.....	164	DILTIAZEM HYDROCHLORIDE.....	120
.Dental.....	339	<i>Dilzem 60 mg (DP)</i>	120
		<i>Dilzem CD (DP)</i>	121
		DIMETHICONE with GLYCEROL	
		.Repatriation Schedule.....	479
		<i>Dimetriose (AV)</i>	152

GENERIC /PROPRIETARY INDEX

<i>Dimirel (ML)</i>	96	DORZOLAMIDE HYDROCHLORIDE with TIMOLOL MALEATE.....	303
<i>Dinac (DP)</i>		<i>Dostinex (PH)</i>	139, 140
.Dental.....	346	<i>Dothep 25 (AF)</i>	275
.Musculo-skeletal system.....	238	<i>Dothep 75 (AF)</i>	275
.Palliative Care.....	324, 325	DOTHIEPIN HYDROCHLORIDE.....	275
<i>Dipentum (UC)</i>	92	<i>Douglas Cefaclor-CD (DG)</i> .Anti-infectives for systemic use.....	168
DIPHEMANIL METHYLSULFATE .Repatriation Schedule.....	482	.Dental.....	342
DIPHENOXYLATE HYDROCHLORIDE with ATROPINE SULFATE.....	90	<i>Douglas Gabapentin 300mg (DP)</i> .Nervous system.....	264
DIPHThERIA and TETANUS VACCINE, ADSORBED.....	180	.Repatriation Schedule.....	493
DIPHThERIA and TETANUS VACCINE, ADSORBED, DILUTED FOR ADULT USE .Anti-infectives for systemic use.....	180	<i>Douglas Gabapentin 400mg (DP)</i> .Nervous system.....	264
.Doctor's Bag Supplies.....	71	.Repatriation Schedule.....	493
DIPIVEFRINE HYDROCHLORIDE.....	302	DOXEPIIN HYDROCHLORIDE.....	275
<i>Diprosone (SH)</i>	136	<i>Doxorubicin Ebewe (IT)</i>	185
DIPYRIDAMOLE.....	103	DOXORUBICIN HYDROCHLORIDE.....	185
DIPYRIDAMOLE with ASPIRIN.....	103	DOXORUBICIN HYDROCHLORIDE, PEGYLATED LIPOSOMAL .Antineoplastic and immunomodulating agents.....	186
DISODIUM ETIDRONATE.....	245	.Section 100.....	374
DISODIUM ETIDRONATE and CALCIUM CARBONATE.....	247	<i>Doxsig (SI)</i> .Anti-infectives for systemic use.....	159, 160
DISODIUM PAMIDRONATE .Musculo-skeletal system.....	245	.Dental.....	336
.Section 100.....	373	<i>Doxy-50 (DP)</i>	159
DISOPYRAMIDE.....	108	<i>Doxy-100 (DP)</i> .Anti-infectives for systemic use.....	159, 160
<i>Distaph 250 (AF)</i> .Anti-infectives for systemic use.....	164	.Dental.....	336
.Dental.....	339	DOXYCYCLINE .Anti-infectives for systemic use.....	159
<i>Distaph 500 (AF)</i> .Anti-infectives for systemic use.....	164	.Dental.....	336
.Dental.....	339	<i>Doxyhexal (HX)</i> .Anti-infectives for systemic use.....	159, 160
<i>Dithiazide (PL)</i>	113	.Dental.....	336
<i>Ditropan (AV)</i>	153	<i>Doxylin 50 (AF)</i>	159
DOCETAXEL.....	184	<i>Doxylin 100 (AF)</i> .Anti-infectives for systemic use.....	159, 160
DOCUSATE SODIUM .Repatriation Schedule.....	473, 498	.Dental.....	336
DOCUSATE SODIUM with BISACODYL .Alimentary tract and metabolism.....	87	<i>D-Penamine (AL)</i>	242
.Palliative Care.....	322	DRESSING—ACTIVATED CHARCOAL (MALODOROUS WOUND) .Repatriation Schedule.....	504
DOCUSATE SODIUM with SENNA .Repatriation Schedule.....	473	DRESSING—ALGINATE (CAVITY WOUND) .Repatriation Schedule.....	504
<i>Dolaforte (CO)</i> .Dental.....	349	DRESSING—ALGINATE (SUPERFICIAL WOUND) .Repatriation Schedule.....	504
.Nervous system.....	249	DRESSING with CADEXOMER IODINE .Repatriation Schedule.....	505
DOLASETRON MESYLATE.....	84	DRESSING—FILM .Repatriation Schedule.....	505
<i>Dolobid (MK)</i> .Dental.....	349	DRESSING—FILM ISLAND .Repatriation Schedule.....	506
.Musculo-skeletal system.....	242	DRESSING—FOAM with CHARCOAL (MALODOROUS WOUND) .Repatriation Schedule.....	506
<i>Doloxene (AS)</i> .Repatriation Schedule.....	492	DRESSING—FOAM—HEAVY EXUDATE .Repatriation Schedule.....	506
DOMPERIDONE.....	84	DRESSING—FOAM—LIGHT EXUDATE .Repatriation Schedule.....	506
DONEPEZIL HYDROCHLORIDE.....	282		
DORNASE ALFA .Section 100.....	373		
<i>Doryx (MX)</i> .Anti-infectives for systemic use.....	159, 160		
.Dental.....	336		
DORZOLAMIDE HYDROCHLORIDE.....	303		

GENERIC /PROPRIETARY INDEX

DRESSING—FOAM—MODERATE EXUDATE		<i>DuoDERM Paste H7930 (CC)</i>	
.Repatriation Schedule.....	507	.Repatriation Schedule.....	508
DRESSING—GAUZE (ABSORBENT PAD)		<i>Duphalac (SM)</i>	88
.Repatriation Schedule.....	507	<i>Duphaston (SM)</i>	146
DRESSING—GAUZE—EYE PAD		<i>Duratears (AQ)</i>	306
.Repatriation Schedule.....	507	<i>Duride (AF)</i>	111
DRESSING—GAUZE—PARAFFIN		<i>Durogesic 25 (JC)</i>	255
.Repatriation Schedule.....	507	<i>Durogesic 50 (JC)</i>	255
DRESSING—GAUZE—PARAFFIN with		<i>Durogesic 75 (JC)</i>	255
CHLORHEXIDINE ACETATE		<i>Durogesic 100 (JC)</i>	256
.Repatriation Schedule.....	507	<i>Duro-K (SZ)</i>	99
DRESSING—GAUZE—POVIDONE-IODINE PAD		<i>Durolax (BY)</i>	
.Repatriation Schedule.....	507	.Alimentary tract and metabolism.....	87
DRESSING—HYDROACTIVE (CAVITY WOUND)		.Palliative Care.....	321
.Repatriation Schedule.....	507	<i>Duro-Tuss (MM)</i>	
DRESSING—HYDROACTIVE (DEBRIDEMENT)		.Repatriation Schedule.....	496
.Repatriation Schedule.....	508	DYDROGESTERONE.....	146
DRESSING—HYDROACTIVE (SUPERFICIAL		<i>Dymadon Forte (GK)</i>	
WOUND—HIGH EXUDATE)		.Dental.....	349
.Repatriation Schedule.....	508	.Nervous system.....	249
DRESSING—HYDROACTIVE (SUPERFICIAL		<i>Dymadon P (PC)</i>	
WOUND—LIGHT EXUDATE)		.Dental.....	354
.Repatriation Schedule.....	508	.Nervous system.....	259
DRESSING—HYDROACTIVE (SUPERFICIAL		<i>Dysport (IS)</i>	
WOUND—MODERATE EXUDATE)		.Section 100.....	424
.Repatriation Schedule.....	508	E	
DRESSING—HYDROCOLLOID (CAVITY WOUND)		<i>Ear Clear for Ear Wax Removal (KY)</i>	
.Repatriation Schedule.....	508	.Repatriation Schedule.....	498
DRESSING—HYDROCOLLOID (SUPERFICIAL		<i>Easiphen (SB)</i>	316
WOUND—LIGHT EXUDATE)		<i>Edecrin (MK)</i>	114
.Repatriation Schedule.....	509	<i>Edronax (PH)</i>	280
DRESSING—HYDROCOLLOID (SUPERFICIAL		<i>E.E.S. 200 (AB)</i>	
WOUND—MODERATE EXUDATE)		.Antiinfectives for systemic use.....	171
.Repatriation Schedule.....	509	.Dental.....	344
DRESSING—HYDROGEL—AMORPHOUS		<i>E.E.S. 400 Filmtab (AB)</i>	
.Repatriation Schedule.....	510	.Antiinfectives for systemic use.....	171
DRESSING—HYDROGEL—SHEET		.Dental.....	344
.Repatriation Schedule.....	511	<i>E.E.S. Granules (AB)</i>	
DRESSING—NON-ADHERENT		.Antiinfectives for systemic use.....	171
.Repatriation Schedule.....	511	.Dental.....	344
<i>Drixine (SH)</i>		EFAVIRENZ	
.Repatriation Schedule.....	495	.Section 100.....	374
DROTRECOGIN ALFA (ACTIVATED).....	104	<i>Efexor (WY)</i>	280
<i>Duatrol SR (ME)</i>		<i>Efexor-XR (WY)</i>	280
.Nervous system.....	259	EFORMOTEROL FUMARATE DIHYDRATE.....	291
.Palliative Care.....	329	<i>Efudix (VT)</i>	
<i>Ducene (SU)</i>		.Repatriation Schedule.....	487
.Dental.....	355	<i>Egocort Cream 1% (EO)</i>	135
.Nervous system.....	272, 273	<i>Egoderma Cream (EO)</i>	
.Palliative Care.....	330	.Repatriation Schedule.....	483
<i>Duocal (SB)</i>	318	<i>Egoderma Ointment (EO)</i>	
<i>DuoDERM CGF H7660 (CC)</i>		.Repatriation Schedule.....	483
.Repatriation Schedule.....	509	<i>Egopsoryl-TA (EO)</i>	
<i>DuoDERM CGF H7662 (CC)</i>		.Repatriation Schedule.....	480
.Repatriation Schedule.....	510	<i>Elastocrepe 36102320 (BV)</i>	
<i>DuoDERM Extra Thin H7955 (CC)</i>		.Repatriation Schedule.....	503
.Repatriation Schedule.....	509	<i>Elastocrepe 36102420 (BV)</i>	
<i>DuoDERM Gel H7987 (CC)</i>		.Repatriation Schedule.....	503
.Repatriation Schedule.....	510	<i>Elastocrepe 36102520 (BV)</i>	
<i>DuoDERM Gel H7990 (CC)</i>		.Repatriation Schedule.....	503
.Repatriation Schedule.....	510		

GENERIC /PROPRIETARY INDEX

<i>Elastoplast 1004 (BV)</i>			
.Repatriation Schedule	512		
<i>Elastoplast 2225 (BE)</i>			
.Repatriation Schedule	503		
<i>Elastoplast 2226 (BE)</i>			
.Repatriation Schedule	503		
<i>Elastoplast 2227 (BE)</i>			
.Repatriation Schedule	503		
<i>Eldepryl (DP)</i>	268		
<i>EleCare (AB)</i>	311		
ELECTROLYTE REPLACEMENT (ORAL)	90		
ELECTROLYTE REPLACEMENT SOLUTION	107		
<i>Eleuphrat (EX)</i>	136		
<i>Eleva 50 (AF)</i>	278		
<i>Eleva 100 (AF)</i>	278		
<i>Elidel (NV)</i>	138		
<i>Eligard 1 month (MX)</i>	190		
<i>Eligard 3 month (MX)</i>	190		
<i>Eligard 4 month (MX)</i>	191		
<i>Eligard 6 month (MX)</i>	191		
<i>Elmendos (ME)</i>	265		
<i>Elocon (SH)</i>			
.Dermatologicals	137		
.Repatriation Schedule	481		
<i>Eloflex 2480 (BV)</i>			
.Repatriation Schedule	501		
<i>Eloxatin (SW)</i>	187		
<i>Emend (MK)</i>	86		
EMTRICITABINE			
.Section 100	375		
<i>Emtriva (GI)</i>			
.Section 100	375		
<i>E-Mycin (AF)</i>			
.Anti-infectives for systemic use	171		
.Dental	344		
<i>E-Mycin 200 (AF)</i>			
.Anti-infectives for systemic use	171		
.Dental	344		
<i>E-Mycin 400 (AF)</i>			
.Anti-infectives for systemic use	171		
.Dental	344		
<i>Enahexal (HX)</i>	122, 123		
<i>Enalabell (BF)</i>	122, 123		
ENALAPRIL MALEATE	122		
ENALAPRIL MALEATE with			
HYDROCHLOROTHIAZIDE	125		
<i>Enalapril-DP 5mg (DP)</i>	122		
<i>Enalapril-DP 10mg (DP)</i>	123		
<i>Enalapril-DP 20mg (DP)</i>	123		
<i>Enbrel (WY)</i>			
.Antineoplastic and immunomodulating agents	222, 225, 232, 235		
.Section 100	378		
<i>Endep 10 (AF)</i>	275		
<i>Endep 25 (AF)</i>	275		
<i>Endep 50 (AF)</i>	275		
<i>Endone (SI)</i>			
.Dental	352		
.Nervous system	254		
<i>Endoxan (BX)</i>	181		
<i>Energivit (SB)</i>	317		
ENFUVRTIDE			
.Section 100	375		
<i>Enidin (PE)</i>	302		
ENOXAPARIN SODIUM	102		
ENTACAPONE	268		
<i>Epaq (AW)</i>			
.Doctor's Bag Supplies	72		
.Respiratory system	291		
<i>Epilim (SW)</i>	263		
<i>Epilim EC (SW)</i>	263		
<i>Epilim Liquid (SW)</i>	263		
<i>Epilim Syrup (SW)</i>	263		
<i>EpiPen (CS)</i>			
.Cardiovascular system	110		
.Respiratory system	298		
<i>EpiPen Jr. (CS)</i>			
.Cardiovascular system	110		
.Respiratory system	298		
<i>Epirubicin Ebewe (IT)</i>	186		
EPIRUBICIN HYDROCHLORIDE	186		
EPOETIN ALFA			
.Section 100	375		
<i>Eprex 1000 (JC)</i>			
.Section 100	375		
<i>Eprex 2000 (JC)</i>			
.Section 100	375		
<i>Eprex 3000 (JC)</i>			
.Section 100	375		
<i>Eprex 4000 (JC)</i>			
.Section 100	375		
<i>Eprex 5000 (JC)</i>			
.Section 100	375		
<i>Eprex 6000 (JC)</i>			
.Section 100	375		
<i>Eprex 8000 (JC)</i>			
.Section 100	375		
<i>Eprex 10000 (JC)</i>			
.Section 100	375		
<i>Eprex 20,000 (JC)</i>			
.Section 100	375		
<i>Eprex 40,000 (JC)</i>			
.Section 100	375		
EPROSARTAN MESYLATE	126		
EPROSARTAN MESYLATE with			
HYDROCHLOROTHIAZIDE	126		
EPTIFIBATIDE ACETATE	103		
<i>Eryc (MX)</i>			
.Anti-infectives for systemic use	171		
.Dental	344		
<i>Erythrocin-I.V. (AB)</i>			
.Anti-infectives for systemic use	171		
.Dental	344		
ERYTHROMYCIN			
.Anti-infectives for systemic use	171		
.Dental	344		
ERYTHROMYCIN ETHYL SUCCINATE			
.Anti-infectives for systemic use	171		
.Dental	344		
ERYTHROMYCIN LACTOBIONATE			
.Anti-infectives for systemic use	171		
.Dental	344		

GENERIC /PROPRIETARY INDEX

ESCITALOPRAM OXALATE		
.Special Pharmaceutical Benefit	74	
<i>Eskazole (GK)</i>	290	
ESOMEPRAZOLE MAGNESIUM TRIHYDRATE	81	
ESOMEPRAZOLE MAGNESIUM TRIHYDRATE and		
CLARITHROMYCIN and AMOXICYLLIN	83	
ESSENTIAL AMINO ACIDS FORMULA with		
MINERALS and VITAMIN C	317	
<i>Estalis continuous 50/140 (NV)</i>	146	
<i>Estalis continuous 50/250 (NV)</i>	146	
<i>Estalis sequi 50/140 (NV)</i>	147	
<i>Estalis sequi 50/250 (NV)</i>	147	
<i>Estracombi (NV)</i>	147	
<i>Estraderm 25 (NV)</i>	143	
<i>Estraderm 50 (NV)</i>	144	
<i>Estraderm 100 (NV)</i>	145	
<i>Estraderm MX 25 (NV)</i>	143	
<i>Estraderm MX 50 (NV)</i>	144	
<i>Estraderm MX 100 (NV)</i>	144	
<i>Estradot 25 (NV)</i>	143	
<i>Estradot 37.5 (NV)</i>	143	
<i>Estradot 50 (NV)</i>	143	
<i>Estradot 75 (NV)</i>	144	
<i>Estradot 100 (NV)</i>	144	
ETANERCEPT		
.Antineoplastic and immunomodulating agents	214	
.Section 100	376	
ETHACRYNIC ACID	114	
ETHOSUXIMIDE	262	
ETONOGESTREL	141	
<i>Etopophos (BQ)</i>	184	
ETOPOSIDE	184	
<i>Etoposide Ebewe (IT)</i>	184	
ETOPOSIDE PHOSPHATE	184	
<i>Eucerin (BE)</i>		
.Repatriation Schedule	479	
<i>Eulexin (SH)</i>	192	
<i>Eutroxsig (FM)</i>	157	
EVEROLIMUS		
.Antineoplastic and immunomodulating agents	235	
.Section 100	378	
<i>Evista (LY)</i>	248	
<i>Exelon (NV)</i>	286	
EXEMESTANE	193	
<i>Exorex (EP)</i>	134	
EZETIMIBE	133	
<i>Ezetrol (MK)</i>	133	
F		
FAMCICLOVIR	178	
<i>Famohexal (HX)</i>	78, 79	
FAMOTIDINE	78	
<i>Famvir (NV)</i>	178, 179	
<i>Fareston (SH)</i>	191	
<i>Fasigyn (PF)</i>	174	
<i>Faverin 50 (AW)</i>	277	
<i>Faverin 100 (AW)</i>	277	
<i>Febridol (DG)</i>		
.Dental	354	
.Nervous system	259	
<i>Feldene (PF)</i>		
.Dental	347, 348	
.Musculo-skeletal system	240	
<i>Feldene-D (PF)</i>		
.Dental	347	
.Musculo-skeletal system	239	
FELODIPINE	119	
<i>Felodur ER 2.5 mg (AL)</i>	119	
<i>Felodur ER 5 mg (AL)</i>	119	
<i>Felodur ER 10 mg (AL)</i>	119	
<i>Femara 2.5 mg (NV)</i>	193	
<i>Femoston 2/10 (SM)</i>	147	
<i>Femtran 25 (MM)</i>	143	
<i>Femtran 50 (MM)</i>	144	
<i>Femtran 100 (MM)</i>	144	
<i>Fenac (AF)</i>		
.Dental	346	
.Musculo-skeletal system	238	
.Palliative Care	325	
<i>Fenac 25 (AF)</i>		
.Dental	346	
.Musculo-skeletal system	238	
.Palliative Care	324, 325	
FENOFIBRATE	131	
FENTANYL	255	
<i>Fermil (AW)</i>	151	
<i>Ferriprox (OA)</i>		
.Section 100	372	
<i>Ferro-Liquid (AE)</i>	106	
<i>Ferrosig (SI)</i>	106	
FERROUS SULFATE	106	
<i>Ferrum H (BX)</i>	106	
FEXOFENADINE HYDROCHLORIDE		
.Repatriation Schedule	497	
<i>Fibsol 5 (AW)</i>	123	
<i>Fibsol 10 (AW)</i>	124	
<i>Fibsol 20 (AW)</i>	124	
FILGRASTIM		
.Section 100	378	
FINASTERIDE		
.Repatriation Schedule	486	
<i>Flagyl (AV)</i>		
.Anti-infectives for systemic use	174	
.Dental	345	
<i>Flagyl S (AV)</i>		
.Anti-infectives for systemic use	174	
.Dental	345	
<i>Flarex (AQ)</i>	301	
FLECAINIDE ACETATE	108	
<i>Flecatab (AF)</i>	108	
<i>Fleet Laxative Suppositories (FL)</i>		
.Alimentary tract and metabolism	87	
.Palliative Care	321	
<i>Flexidress 650941 (CC)</i>		
.Repatriation Schedule	504	
<i>Flixotide (GK)</i>	296	
<i>Flixotide Accuhaler (GK)</i>	296	
<i>Flixotide Junior (GK)</i>	295	
<i>Flixotide Junior Accuhaler (GK)</i>	296	
<i>Flomax (CS)</i>		
.Repatriation Schedule	486	

GENERIC /PROPRIETARY INDEX

<i>Flopen (CS)</i>		FOSAMPRENAVIR CALCIUM	
Antiinfectives for systemic use.....	164	Section 100	380
Dental.....	339, 340	FOSCARNET SODIUM	
<i>Florinef (BQ)</i>	155	Section 100	380
<i>Floxapen (GK)</i>		<i>Foscavir (AP)</i>	
Antiinfectives for systemic use.....	164	Section 100	380
Dental.....	339, 340	FOSINOPRIL SODIUM	123
<i>Floxsig (SI)</i>		FOSINOPRIL SODIUM with	
Antiinfectives for systemic use.....	164	HYDROCHLOROTHIAZIDE	125
Dental.....	339, 340	FOTEMUSTINE	181
<i>Fluanxol Concentrated Depot (LU)</i>	269	<i>Fragmin (PH)</i>	101
<i>Fluanxol Depot (LU)</i>	269	<i>Frakas (AW)</i>	159
<i>Fluarix (GK)</i>	180	FRAMYCETIN SULFATE.....	307
<i>Flubiclox (DP)</i>		<i>Frusehexal (HX)</i>	
Antiinfectives for systemic use.....	164	Cardiovascular system	114
Dental.....	339	Doctor's Bag Supplies	71
FLUCLOXACILLIN		<i>Frusehexal 40 mg (HX)</i>	114
Antiinfectives for systemic use.....	164	FRUSEMIDE	
Dental.....	339	Cardiovascular system	114
<i>Flucon (AQ)</i>	301	Doctor's Bag Supplies	71
FLUCONAZOLE.....	175	<i>Frusid (DP)</i>	114
FLUDROCORTISONE ACETATE.....	155	<i>Fucidin (CS)</i>	174
FLUNITRAZEPAM		<i>Fungilin (BQ)</i>	
Repatriation Schedule	494	Alimentary tract and metabolism	77
<i>Fluohexal (HX)</i>	277	Dental	333
FLUOROMETHOLONE	301	<i>Fungizone (BQ)</i>	175
FLUOROMETHOLONE ACETATE	301	FUSIDIC ACID.....	174
FLUOROURACIL		<i>Fuzeon (RO)</i>	
Antineoplastic and immunomodulating agents.....	183	Section 100	375
Repatriation Schedule	487	<i>Fybogel (RC)</i>	
<i>Fluoxebell (BF)</i>	277	Repatriation Schedule	473
FLUOXETINE HYDROCHLORIDE	276	G	
<i>Fluoxetine-DP (DP)</i>	277	<i>Gabahexal 300mg (HX)</i>	264
FLUPENTHIXOL DECANOATE	269	<i>Gabahexal 400mg (HX)</i>	264
FLUPHENAZINE DECANOATE	268	GABAPENTIN	
FLURBIPROFEN SODIUM.....	302	Nervous system	264
FLUTAMIDE	192	Repatriation Schedule	492
<i>Flutamin (AF)</i>	192	<i>Gabitril (MX)</i>	263
FLUTICASONE PROPIONATE	295	GALANTAMINE HYDROBROMIDE	283
FLUTICASONE PROPIONATE with SALMETEROL		GANCICLOVIR	
XINAFOATE.....	294	Section 100	380
FLUVASTATIN SODIUM	129	GANCICLOVIR SODIUM	
<i>Fluvax (CS)</i>	180	Section 100	380
FLUVOXAMINE MALEATE	277	<i>Gantin (AW)</i>	
<i>FML Liquifilm (AG)</i>	301	Nervous system	264
FOLIC ACID	106	Repatriation Schedule	492, 493
FOLLITROPIN ALFA		<i>Gastrogel (FM)</i>	77
Genito urinary system and sex hormones	148	<i>Gastro-Stop Loperamide (AS)</i>	90
Section 100	426	GATIFLOXACIN	
FOLLITROPIN BETA		Repatriation Schedule	487
Genito urinary system and sex hormones	149	GAUZE and COTTON TISSUE (COMBINE ROLL)	
Section 100	426	Repatriation Schedule	511
FONDAPARINUX SODIUM.....	105	<i>Gaviscon P (RC)</i>	83
<i>Foradile (NV)</i>	291	GEFITINIB	188
<i>Formet 500 (AW)</i>	95	GELATIN - SUCCINYLATED	107
<i>Formet 850 (AW)</i>	95	<i>Gelocast Elastic 1080 (BV)</i>	
<i>Formet 1000 (AW)</i>	95	Repatriation Schedule	504
<i>Fortovase (RO)</i>		<i>Gelofusine (BR)</i>	107
Section 100	421	<i>GeITears (BU)</i>	305
<i>Fosamax 40 mg (MK)</i>	245	<i>Gelusil (WW)</i>	77
<i>Fosamax Once Weekly (MK)</i>	245	GEMCITABINE HYDROCHLORIDE	183

GENERIC /PROPRIETARY INDEX

GEMFIBROZIL	132	GenRx Enalapril (GX).....	122, 123
Gemhexal (HX).....	132	GenRx Famotidine (GX).....	78, 79
Gemzar (LY).....	183	GenRx Fluoxetine (GX).....	277
Genlac (AW).....	88	GenRx Frusemide (GX).....	114
Genoptic (AG).....	300	GenRx Gabapentin (GX)	
Genoral 0.625 (KR).....	145	.Nervous system.....	264
Genoral 1.25 (KR).....	145	.Repatriation Schedule.....	493
Genotropin (PH)		GenRx Gemfibrozil (GX).....	132
.Section 100.....	425	GenRx Gliclazide (GX).....	96
Genotropin Mini Quick (PH)		GenRx Indapamide (GX).....	113
.Section 100.....	425	GenRx Ipratropium (GX).....	296, 297
Genox 10 (AF).....	191	GenRx Isosorbide Mononitrate (GX).....	111
Genox 20 (AF).....	191	GenRx Isotretinoin (GX).....	137
GenRx Aciclovir (GX).....	177, 178	GenRx Lactulose (GX).....	88
GenRx Allopurinol (GX).....	244	GenRx Lamotrigine (GX).....	265
GenRx Alprazolam (GX).....	272	GenRx Lisinopril (GX).....	123, 124
GenRx Amiodarone (GX).....	109	GenRx Metformin (GX).....	95
GenRx Amoxicillin (GX)		GenRx Metoprolol (GX).....	117
.Anti-infectives for systemic use.....	161, 162	GenRx Moclobemide (GX).....	279
.Dental.....	337, 338	GenRx Nifedipine (GX).....	119
GenRx Amoxicillin and Clavulanic Acid (GX)		GenRx Norfloxacin (GX).....	173
.Anti-infectives for systemic use.....	165	GenRx Paroxetine (GX).....	277
.Dental.....	340	GenRx Piroxicam (GX)	
GenRx Atenolol (GX).....	116	.Dental.....	347, 348
GenRx Azathioprine (GX).....	237	.Musculo-skeletal system.....	240
GenRx Baclofen (GX).....	243	GenRx Piroxicam Dispersible (GX)	
GenRx Calcitriol (GX)		.Dental.....	347
.Alimentary tract and metabolism.....	99	.Musculo-skeletal system.....	239
.Musculo-skeletal system.....	247	GenRx Prazosin (GX).....	112, 113
GenRx Captopril (GX).....	121, 122	GenRx Ranitidine (GX).....	79, 80
GenRx Cefaclor (GX)		GenRx Salbutamol (GX)	
.Anti-infectives for systemic use.....	168	.Doctor's Bag Supplies.....	72, 73
.Dental.....	343	.Respiratory system.....	292
GenRx Cefaclor CD (GX)		GenRx Sertraline (GX).....	278
.Anti-infectives for systemic use.....	168	GenRx Simvastatin (GX).....	130, 131
.Dental.....	342	GenRx Sotalol (GX).....	109
GenRx Cephalexin (GX)		GenRx Tamoxifen (GX).....	191
.Anti-infectives for systemic use.....	166, 167	GenRx Terbinafine (GX).....	134
.Dental.....	341, 342	GenRx Tramadol (GX)	
GenRx Cimetidine (GX).....	78	.Dental.....	353
GenRx Ciprofloxacin (GX).....	172	.Nervous system.....	257, 258
GenRx Citalopram (GX).....	276	GenRx Trimethoprim with Sulfamethoxazole DS (GX)	
GenRx Clarithromycin (GX).....	170	.Anti-infectives for systemic use.....	170
GenRx Clomiphene (GX).....	151	.Dental.....	344
GenRx Clomipramine (GX).....	274, 275	GENTAMICIN SULFATE.....	171
GenRx Clotrimazole 3 Day Cream (GX)		.Anti-infectives for systemic use.....	171
.Repatriation Schedule.....	484	.Sensory organs.....	300
GenRx Clotrimazole 6 Day Cream (GX)		Genteal (NV).....	306
.Repatriation Schedule.....	484	Genteal gel (NV).....	306
GenRx Cyproterone Acetate (GX)		GESTRINONE.....	152
.Antineoplastic and immunomodulating agents.....	192	GLATIRAMER ACETATE.....	196
.Genito urinary system and sex hormones.....	151, 152	GLIBENCLAMIDE.....	95
GenRx Diclofenac (GX)		GLICLAZIDE.....	95
.Dental.....	346	Glimel (AF).....	95
.Musculo-skeletal system.....	238	GLIMEPIRIDE.....	96
.Palliative Care.....	324, 325	GLIPIZIDE.....	96
GenRx Diltiazem (GX).....	120	Glivec (NV)	
GenRx Diltiazem CD (GX).....	121	.Section 100.....	429, 431, 432
GenRx Doxycycline (GX)		GLOVES PLASTIC (DISPOSABLE)	
.Anti-infectives for systemic use.....	159, 160	.Repatriation Schedule.....	511
.Dental.....	336		

GENERIC /PROPRIETARY INDEX

<i>GlucaGen Hypokit (NO)</i>				H	
.Dental.....	336			<i>Haemacel (AE)</i>	107
.Doctor's Bag Supplies	71			<i>Haldol decanoate (JC)</i>	269
.Systemic hormonal preparations, excl. sex hormones and insulins	158			HALOPERIDOL	
GLUCAGON HYDROCHLORIDE				.Doctor's Bag Supplies	71
.Dental.....	336			.Nervous system	269
.Doctor's Bag Supplies	71			HALOPERIDOL DECANOATE	269
.Systemic hormonal preparations, excl. sex hormones and insulins	158			<i>Hamilton Bath Oil (HA)</i>	
<i>Glucobay 50 (BN)</i>	96			.Repatriation Schedule	479
<i>Glucobay 100 (BN)</i>	96			<i>Hamilton Body Wash (HA)</i>	
<i>Glucocare (DR)</i>	309			.Repatriation Schedule	483
<i>Glucocare Super Sensor (DR)</i>	309			<i>Hamilton Broad Spectrum Milky Lotion 15+ (HA)</i>	
<i>Glucoflex-R (TC)</i>	310			.Repatriation Schedule	480
<i>Glucohexal (HX)</i>	95			<i>Hamilton Pine Tar Solution (HA)</i>	
<i>Glucomen Sensor (GD)</i>	309			.Repatriation Schedule	480
<i>Glucomet 500 mg (DP)</i>	95			<i>Hamilton Solastick 30+ (HA)</i>	
<i>Glucomet 850 mg (DP)</i>	95			.Repatriation Schedule	479
<i>Glucophage (MQ)</i>	95			<i>Hamilton Sunscreen Broad Spectrum Cream 15+ (HA)</i>	
GLUCOSE				.Repatriation Schedule	479
.Blood and blood forming organs	107			<i>Handy 4207 (BV)</i>	
.Dental.....	334			.Repatriation Schedule	511
GLUCOSE INDICATOR—BLOOD	309, 310			<i>Handy 4208 (BV)</i>	
GLUCOSE INDICATOR—URINE	309			.Repatriation Schedule	511
GLUCOSE and KETONE INDICATOR—URINE	309			<i>Handy 4209 (BV)</i>	
<i>Glucostix (BN)</i>	310			.Repatriation Schedule	511
<i>Glucovance 250mg/1.25mg (AL)</i>	96			<i>Handy 5608 (BV)</i>	
<i>Glucovance 500mg/2.5mg (AL)</i>	96			.Repatriation Schedule	501
<i>Glucovance 500mg/5mg (AL)</i>	96			<i>Handy 5672 (BV)</i>	
<i>Glyade (AF)</i>	96			.Repatriation Schedule	507
GLYCEROL				<i>Handy 5674 (BV)</i>	
.Alimentary tract and metabolism.....	89			.Repatriation Schedule	507
.Palliative Care.....	323			<i>Handygauze Cohesive 8631 (BV)</i>	
.Repatriation Schedule	474			.Repatriation Schedule	502
GLYCERYL TRINITRATE				<i>Handygauze Cohesive 8633 (BV)</i>	
.Cardiovascular system	110			.Repatriation Schedule	502
.Dental.....	335			<i>Handygauze Cohesive 8635 (BV)</i>	
.Doctor's Bag Supplies	71			.Repatriation Schedule	502
<i>Gold Cross (BI)</i>				<i>HCU express (VF)</i>	315
.Repatriation Schedule	491, 496			<i>HCU gel (VF)</i>	315
<i>Gonal-f (SG)</i>				<i>healthsense Clotrimazole 3 Day Cream (HS)</i>	
.Genito urinary system and sex hormones	148, 149			.Repatriation Schedule	484
.Section 100	426			<i>healthsense Clotrimazole 6 Day Cream (HS)</i>	
<i>Gonal-f 75 (SG)</i>				.Repatriation Schedule	484
.Genito urinary system and sex hormones	148, 149			HEPARIN SODIUM	102
.Section 100	426			<i>Hepsera (GI)</i>	
<i>Gonal-f Pen (SG)</i>				.Section 100	359
.Genito urinary system and sex hormones	148, 149			<i>Hexal Clotfeme 3 Day Cream (HX)</i>	
.Section 100	426			.Repatriation Schedule	484
<i>Gopten (AB)</i>	125			<i>Hexal Clotfeme 6 Day Cream (HX)</i>	
GOSERELIN ACETATE.....	190			.Repatriation Schedule	484
GRANISETRON HYDROCHLORIDE.....	84			<i>Hexal Konazol 2% Shampoo (HX)</i>	
<i>Granocyte 13 (MX)</i>				.Repatriation Schedule	477
.Section 100	413			HEXAMINE HIPPURATE	175
<i>Granocyte 34 (MX)</i>				<i>Hiprex (MM)</i>	175
.Section 100	413			<i>Hivid (RO)</i>	
GRISOFULVIN	134			.Section 100	423
<i>Grisovin (SI)</i>	134			<i>Holoxan (BX)</i>	181
<i>Grisovin 500 (SI)</i>	134			HOMATROPINE HYDROBROMIDE	304
				<i>HPMC PAA (CV)</i>	306
				<i>Humalog (LY)</i>	93
				<i>Humalog Mix25 (LY)</i>	94

GENERIC /PROPRIETARY INDEX

HUMAN CHORIONIC GONADOTROPHIN	
Genito urinary system and sex hormones	150
Section 100	427
<i>Humatrope</i> (LY)	
Section 100	425
<i>Humira</i> (AB)	202, 204
<i>Humulin 30/70</i> (LY)	94
<i>Humulin NPH</i> (LY)	94
<i>Humulin R</i> (LY)	93, 94
<i>Hycamtin</i> (GK)	189
<i>Hycor</i> (SI)	301
<i>Hydopa</i> (AF)	112
HYDRALAZINE HYDROCHLORIDE	113
<i>Hydrea</i> (BQ)	189
<i>Hydrene 25/50</i> (AF)	115
HYDROCHLOROTHIAZIDE	113
HYDROCHLOROTHIAZIDE with AMILORIDE	
HYDROCHLORIDE	115
HYDROCHLOROTHIAZIDE with TRIAMTERENE	115
<i>Hydrocoll 900938/1</i> (HR)	
Repatriation Schedule	509
<i>Hydrocoll 900939/1</i> (HR)	
Repatriation Schedule	510
<i>Hydrocoll Thin 900942/1</i> (HR)	
Repatriation Schedule	509
HYDROCORTISONE	
Dermatologicals	135
Sensory organs	301
Systemic hormonal preparations, excl. sex hormones and insulins	155
HYDROCORTISONE ACETATE	
Alimentary tract and metabolism	90
Dental	335
Dermatologicals	135
Sensory organs	301
HYDROCORTISONE with CINCHOCAINE	
HYDROCHLORIDE	
Repatriation Schedule	476
HYDROCORTISONE SODIUM SUCCINATE	
Dental	335
Doctor's Bag Supplies	71
Systemic hormonal preparations, excl. sex hormones and insulins	156
HYDROLYZED COLLAGEN PROTEINS	
Repatriation Schedule	483
HYDROMORPHONE HYDROCHLORIDE	
Dental	349
Nervous system	249
HYDROXOCOBALAMIN	106
HYDROXYCHLOROQUINE SULFATE	242
HYDROXYPROPYLCELLULOSE	305
HYDROXYUREA	189
<i>Hygroton 25</i> (NV)	113
HYOSCINE BUTYLBROMIDE	
Palliative Care	320
Repatriation Schedule	473
<i>Hypafix 71443-0</i> (BV)	
Repatriation Schedule	512
<i>Hypafix 71443-1</i> (BV)	
Repatriation Schedule	512
<i>Hypnodorm</i> (AF)	
Repatriation Schedule	494
HYPROMELLOSE	306
HYPROMELLOSE with CARBOMER 980	306
HYPROMELLOSE with DEXTRAN	306
<i>Hypurin Isophane</i> (AS)	94
<i>Hypurin Neutral</i> (AS)	93
<i>Hysone 4</i> (AF)	155
<i>Hysone 20</i> (AF)	155
<i>Hytrin</i> (AB)	
Repatriation Schedule	486
I	
<i>Iallex</i> (LN)	
Antiinfectives for systemic use	166, 167
Dental	341, 342
<i>Ibilex 125</i> (AF)	
Antiinfectives for systemic use	167
Dental	342
<i>Ibilex 250</i> (AF)	
Antiinfectives for systemic use	166, 167
Dental	341, 342
<i>Ibilex 500</i> (AF)	
Antiinfectives for systemic use	166
Dental	341
<i>Ibimicyn</i> (DP)	
Antiinfectives for systemic use	162
Dental	338
IBUPROFEN	
Dental	348
Musculo-skeletal system	240
Palliative Care	327
IDARUBICIN HYDROCHLORIDE	186
IFOSFAMIDE	181
<i>Ikorel</i> (AV)	112
ILOPROST TROMETAMOL	
Section 100	381
IMATINIB MESYLATE	
Section 100	428
<i>Imdur 120 mg</i> (AP)	111
<i>Imdur Durule</i> (AP)	111
<i>Imigran</i> (GK)	260
IMIPRAMINE HYDROCHLORIDE	276
<i>ImmuCyst</i> (AV)	196
<i>Imodium</i> (JC)	90
<i>Imovane</i> (AV)	
Repatriation Schedule	494
<i>Implanon</i> (OR)	141
<i>Improvil 28 Day</i> (KR)	141
<i>Imtrate 60 mg</i> (DP)	111
<i>Imukin</i> (BY)	
Section 100	411
<i>Imuran</i> (GK)	237
<i>In a Wink Moisturising</i> (CV)	306
<i>Indahexal</i> (HX)	113
INDAPAMIDE HEMIHYDRATE	113
<i>Inderal</i> (AP)	116
INDINAVIR SULFATE	
Section 100	389

GENERIC /PROPRIETARY INDEX

<i>Indocid (MK)</i>		<i>Inza 500 (AF)</i>	
.Dental.....	347	.Dental.....	348
.Musculo-skeletal system.....	238	.Musculo-skeletal system.....	240
.Palliative Care.....	326	.Palliative Care.....	327, 328
INDOMETHACIN		<i>Iodosorb 66051330 (SN)</i>	
.Dental.....	347	.Repatriation Schedule.....	505
.Musculo-skeletal system.....	238	<i>Iodosorb 66051340 (SN)</i>	
.Palliative Care.....	326	.Repatriation Schedule.....	505
INFLIXIMAB		<i>Iodosorb 66051360 (SN)</i>	
.Repatriation Schedule.....	490	.Repatriation Schedule.....	505
.Section 100.....	390	<i>Iodosorb Ointment 66051230 (SN)</i>	
INFLUENZA VACCINE.....	180	.Repatriation Schedule.....	505
<i>Influvac (SM)</i>	180	<i>Iodosorb Ointment 66051240 (SN)</i>	
INSECT ALLERGEN EXTRACT—HONEY BEE		.Repatriation Schedule.....	505
VENOM.....	308	<i>Iodosorb Powder 66051070 (SN)</i>	
INSECT ALLERGEN EXTRACT—PAPER WASP		.Repatriation Schedule.....	505
VENOM.....	308	<i>Ionil (GA)</i>	
INSECT ALLERGEN EXTRACT—YELLOW JACKET		.Repatriation Schedule.....	482
VENOM.....	308	<i>Ionil Rinse (GA)</i>	
<i>Insensye (FR)</i>	173	.Repatriation Schedule.....	483
<i>Insig (SI)</i>	113	<i>Ionil-T (GA)</i>	
INSULIN ASPART.....	93	.Repatriation Schedule.....	482
INSULIN ASPART—INSULIN ASPART PROTAMINE		<i>Iopidine 0.5% (AQ)</i>	302
SUSPENSION.....	94	<i>Ipratrin (AF)</i>	296
INSULIN ISOPHANE (N.P.H.).....	94	<i>Ipratrin Adult (AF)</i>	297
INSULIN LISPRO.....	93	IPRATROPIUM BROMIDE	
INSULIN LISPRO—INSULIN LISPRO PROTAMINE		.Repatriation Schedule.....	495
SUSPENSION.....	94	.Respiratory system.....	296
INSULIN NEUTRAL.....	93	IPRATROPIUM BROMIDE with SALBUTAMOL	
INSULIN NEUTRAL—INSULIN ISOPHANE (N.P.H.),		SULFATE	
(MIXED) (Biphasic Isophane).....	94	.Repatriation Schedule.....	496
<i>Intal (AV)</i>	297	<i>Ipravent (PU)</i>	296, 297
<i>Intal CFC-Free (AV)</i>	297	IRBESARTAN.....	126
<i>Intal Forte CFC-Free (AV)</i>	297	IRBESARTAN with HYDROCHLOROTHIAZIDE.....	127
<i>Intal Spincaps (DP)</i>	297	<i>Ircal (PE)</i>	306
<i>Integrilin (SH)</i>	103	<i>Iressa (AP)</i>	189
INTERFERON ALFA-2a		IRINOTECAN HYDROCHLORIDE TRIHYDRATE.....	189
.Antineoplastic and immunomodulating agents.....	193	IRON POLYMALTOSE COMPLEX.....	106
.Section 100.....	409	IRON SUCROSE.....	106
INTERFERON ALFA-2b		<i>Iscover (BQ)</i>	
.Antineoplastic and immunomodulating agents.....	194	.Blood and blood forming organs.....	103
.Section 100.....	410	.Repatriation Schedule.....	475
INTERFERON BETA-1a.....	195	<i>Isohexal (HX)</i>	137
INTERFERON BETA-1b.....	195	<i>Isomonit (HX)</i>	111
INTERFERON GAMMA-1b		ISONIAZID.....	176
.Section 100.....	411	<i>Isoptin (AB)</i>	
<i>Intrasite Gel 7313 (SN)</i>		.Cardiovascular system.....	120
.Repatriation Schedule.....	510	.Doctor's Bag Supplies.....	73
<i>Intron A (SH)</i>		<i>Isoptin 180 SR (AB)</i>	120
.Section 100.....	410, 411	<i>Isoptin SR (AB)</i>	120
<i>Intron A Redipen (SH)</i>		<i>Isopto Carbachol (AQ)</i>	302
.Antineoplastic and immunomodulating agents.....	194	<i>Isopto Carpine (AQ)</i>	302, 303
.Section 100.....	411	<i>Isopto Homatropine (AQ)</i>	304
<i>Invirase (RO)</i>		<i>Isopto Tears (AQ)</i>	306
.Section 100.....	421	<i>Isordil (SI)</i>	111
<i>Inza 250 (AF)</i>		<i>Isordil Sublingual (SI)</i>	111
.Dental.....	348	ISOSORBIDE DINITRATE.....	111
.Musculo-skeletal system.....	240	ISOSORBIDE MONONITRATE.....	111
.Palliative Care.....	327, 328	ISOTRETINOIN.....	137
		ISPAGHULA HUSK	
		.Repatriation Schedule.....	473

GENERIC /PROPRIETARY INDEX

ITRACONAZOLE	176	<i>Keto-Diabur-Test 5000 (RD)</i>	309
IVERMECTIN	290	<i>Keto-Diastix (BN)</i>	309
J		<i>Ketonex-1 (AB)</i>	317
<i>Jelonet 7404 (SN)</i>		<i>Ketonex-2 (AB)</i>	317
.Repatriation Schedule	507	KETOPROFEN	
<i>Jezil (AF)</i>	132	.Dental	348
<i>JJ 02013 (JJ)</i>		.Musculo-skeletal system	240
.Repatriation Schedule	504	<i>Kindergen (SB)</i>	318
<i>JJ 12010 (JJ)</i>		<i>Kineret (AN)</i>	210, 212
.Repatriation Schedule	511	<i>Kinidin Durule (AP)</i>	108
K		<i>Kinson (AF)</i>	266
<i>Kaletra (AB)</i>		<i>Kivexa (GK)</i>	
.Section 100	413	.Section 100	359
<i>Kalixocin (AF)</i>	170	<i>Klacid (AB)</i>	
<i>Kalma 0.25 (AF)</i>	272	.Anti-infectives for systemic use	170
<i>Kalma 0.5 (AF)</i>	272	.Section 100	370
<i>Kalma 1 (AF)</i>	272	<i>Klacid Hp 7 (AB)</i>	83
<i>Kalma 2 (AF)</i>	272	<i>Kliogest (NO)</i>	146
<i>Kaltostat 168117 (CC)</i>		<i>Kliovance (NO)</i>	146
.Repatriation Schedule	504	<i>Kosteo (AW)</i>	
<i>Kaltostat 168210 (CC)</i>		.Alimentary tract and metabolism	99
.Repatriation Schedule	504	.Musculo-skeletal system	247
<i>Kaltostat 168212 (CC)</i>		<i>Kredex (MD)</i>	118
.Repatriation Schedule	505	<i>Krypton 2.5 (AF)</i>	
<i>Kaluril (AF)</i>	115	.Genito urinary system and sex hormones	139
<i>Kapanol (GK)</i>		.Nervous system	267
.Dental	351	<i>Krypton 5 (AF)</i>	
.Nervous system	252, 253	.Genito urinary system and sex hormones	139
<i>Karicare De-Lact (NU)</i>	313, 314	.Nervous system	267
<i>Karlor CD (LN)</i>		<i>Krypton 10 (AF)</i>	
.Anti-infectives for systemic use	168	.Genito urinary system and sex hormones	139
.Dental	342	.Nervous system	267
<i>Karvea (SW)</i>	126	<i>K-Sol (LN)</i>	99
<i>Karvezide 150/12.5 (SW)</i>	127	<i>Kytril (MX)</i>	84
<i>Karvezide 300/12.5 (SW)</i>	127	L	
<i>Keflex (AS)</i>		LABETALOL HYDROCHLORIDE	118
.Anti-infectives for systemic use	166, 167	<i>Lac-Dol (DP)</i>	88
.Dental	341, 342	<i>Lacri-Lube (AG)</i>	306
<i>Keflin Neutral (LY)</i>		<i>Lacrisert (SI)</i>	305
.Anti-infectives for systemic use	167	<i>Lactocur (HX)</i>	88
.Dental	342	LACTULOSE	88
<i>Keflor (AF)</i>		<i>Lamictal (GK)</i>	265
.Anti-infectives for systemic use	168	<i>Lamisil (NC)</i>	
.Dental	343	.Repatriation Schedule	478
<i>Keflor CD (AF)</i>		<i>Lamisil (NV)</i>	
.Anti-infectives for systemic use	168	.Dermatologicals	134
.Dental	342	.Repatriation Schedule	478
<i>Kefzol (LY)</i>	167	<i>Lamisil DermGel (NC)</i>	
<i>Kenacomb (BQ)</i>		.Repatriation Schedule	478
.Repatriation Schedule	481	<i>Lamitrin (HX)</i>	265
<i>Kenacomb Otic (BQ)</i>	307	LAMIVUDINE	
<i>Kenacort-A10 (BQ)</i>		.Section 100	411
.Dental	336	LAMIVUDINE with ZIDOVUDINE	
.Systemic hormonal preparations, excl. sex hormones and insulins	157	.Section 100	411
<i>Keppra (UC)</i>		<i>Lamogine (AF)</i>	265
.Special Pharmaceutical Benefit	75	LAMOTRIGINE	265
KETOCONAZOLE		<i>Lamotrigine-DP (DP)</i>	265
.Anti-infectives for systemic use	175	<i>Lanoxin (SI)</i>	108
.Repatriation Schedule	477	<i>Lanoxin-PG (SI)</i>	108
		LANREOTIDE ACETATE	
		.Section 100	412

GENERIC /PROPRIETARY INDEX

LANSOPRAZOLE	81	LEVONORGESTREL with ETHINYLLOESTRADIOL	140, 141
<i>Lanvis</i> (GK)	183	<i>Lexapro</i> (LU)	74
<i>Largactil</i> (AV)		Special Pharmaceutical Benefit	74
Doctor's Bag Supplies	71	<i>Lexotan</i> (RO)	493
Nervous system	268	Repatriation Schedule	493
<i>Lasix</i> (AV)		LIGNOCAINE HYDROCHLORIDE	
Cardiovascular system	114	Cardiovascular system	108
Doctor's Bag Supplies	71	Dental	334
<i>Lasix-M</i> (AV)	114	Doctor's Bag Supplies	72
LATANOPROST	304	LIGNOCAINE HYDROCHLORIDE with	
LATANOPROST with TIMOLOL MALEATE		CARBOXYMETHYLCELLULOSE	
Repatriation Schedule	497	Repatriation Schedule	480
<i>Lax-Tab</i> (AE)		<i>Lincocin</i> (PH)	
Alimentary tract and metabolism	87	Antifectives for systemic use	171
Palliative Care	321	Dental	345
<i>Ledertrexate</i> (WY)	182, 237	LINCOMYCIN	
LEFLUNOMIDE	235	Antifectives for systemic use	171
<i>Lengout</i> (LN)	244	Dental	345
LENOGRASTIM		<i>Lioresal 10</i> (NV)	243
Section 100	413	<i>Lioresal 25</i> (NV)	243
LERCANIDIPINE HYDROCHLORIDE		<i>Lioresal Intrathecal</i> (NV)	
Cardiovascular system	119	Section 100	360
Repatriation Schedule	476	LIOTHYRONINE SODIUM	157
<i>Lescol</i> (NV)	129	<i>Lipazil 600 mg</i> (DP)	132
LETROZOLE	193	<i>Lipex 5</i> (AD)	129
<i>Leucovorin Calcium</i> (MX)	308	<i>Lipex 10</i> (AD)	130
<i>Leucovorin Calcium</i> (PF)	308	<i>Lipex 20</i> (AD)	130
<i>Leukeran</i> (GK)	181	<i>Lipex 40</i> (AD)	130
<i>Leukoflex 1124</i> (BV)		<i>Lipex 80</i> (AD)	131
Repatriation Schedule	512	<i>Lipidil</i> (LF)	131
<i>Leukoplast 1071</i> (BV)		<i>Lipitor</i> (PF)	129
Repatriation Schedule	512	<i>Lipostat 10</i> (AW)	129
<i>Leukoplast 1072</i> (BV)		<i>Lipostat 20</i> (AW)	129
Repatriation Schedule	512	<i>Lipostat 40</i> (AW)	129
<i>Leukoplast 1073</i> (BV)		<i>Lipostat 80</i> (AW)	129
Repatriation Schedule	512	<i>Liprace</i> (DP)	123, 124
<i>Leukopor 2471</i> (BV)		<i>Liquifilm Forte</i> (AG)	307
Repatriation Schedule	512	<i>Liquifilm Tears</i> (AG)	307
<i>Leukopor 2472</i> (BV)		<i>Lisinobell</i> (BF)	123, 124
Repatriation Schedule	512	LISINAPRIL	123
<i>Leukopor 2474</i> (BV)		<i>Lisinopril Hexal</i> (HX)	123, 124
Repatriation Schedule	512	<i>Lisodur</i> (AF)	123, 124
<i>Leukosilk 1021</i> (BV)		<i>Litak</i> (OA)	182
Repatriation Schedule	512	<i>Lithicarb</i> (AS)	279
<i>Leukosilk 1022</i> (BV)		LITHIUM CARBONATE	279
Repatriation Schedule	512	<i>Livostin</i> (JC)	
<i>Leukosilk 1024</i> (BV)		Repatriation Schedule	495, 497
Repatriation Schedule	512	<i>Locasol</i> (NU)	314
LEUPRORELIN ACETATE	190	<i>Loceryl</i> (GA)	
<i>Leustatin</i> (JC)	182	Repatriation Schedule	478
LEVETIRACETAM		<i>Locilan 28 Day</i> (KR)	141
Special Pharmaceutical Benefit	75	LODOXAMIDE TROMETAMOL	304
<i>Levlen ED</i> (SY)	140	<i>Lofenoxal</i> (KR)	90
LEVOBUNOLOL HYDROCHLORIDE	303	<i>Logicin Rapid Relief</i> (SI)	
LEVOCABASTINE HYDROCHLORIDE		Repatriation Schedule	495
Repatriation Schedule	495, 497	<i>Logicin Sinus</i> (SI)	
LEVODOPA with BENSERAZIDE	266	Repatriation Schedule	496
LEVODOPA with CARBIDOPA	266	<i>Logynon ED</i> (SY)	141
LEVODOPA with CARBIDOPA and ENTACAPONE	267	<i>Lomide</i> (AQ)	304
<i>Levohexal</i> (HX)	266	<i>Lomotil</i> (PH)	90
LEVONORGESTREL	139, 141		

GENERIC /PROPRIETARY INDEX

<i>Loniten (PH)</i>	113	<i>Mapleflex (SB)</i>	317
LOPERAMIDE HYDROCHLORIDE	90	<i>Marevan (FM)</i>	101
<i>Lophlex (SB)</i>	316	<i>Maxamox (SZ)</i>	
<i>Lopid (PF)</i>	132	.Anti-infectives for systemic use	162
LOPINAVIR with RITONAVIR		.Dental	338
.Section 100	413	<i>Maxidex (AQ)</i>	301
<i>Lopresor 50 (NV)</i>	117	<i>Maxipime (BQ)</i>	169
<i>Lopresor 100 (NV)</i>	117	<i>Maxolon (VT)</i>	
LORATADINE		.Alimentary tract and metabolism	84
.Repatriation Schedule	497	.Dental	333
<i>Losec Tablets (AP)</i>	82	.Doctor's Bag Supplies	72
<i>Lovan (AL)</i>	277	<i>MCT Oil (SB)</i>	310
<i>Lovan 20 Tab (AL)</i>	276	MEBENDAZOLE	
<i>Lovan Liquid (AL)</i>	277	.Repatriation Schedule	495
<i>Lovir (DP)</i>	177, 178	MEBEVERINE HYDROCHLORIDE	
LPV (CS)		.Repatriation Schedule	473
.Anti-infectives for systemic use	163	<i>Medipore 2961 (MM)</i>	
.Dental	338	.Repatriation Schedule	512
LUBRICATING AGENT		<i>Medroxyhexal (HX)</i>	146
.Repatriation Schedule	511	MEDROXYPROGESTERONE ACETATE	
<i>Lucrin Depot (AB)</i>	190	.Antineoplastic and immunomodulating agents	189
<i>Lucrin Depot 3 Month Injection (AB)</i>	190	.Genito urinary system and sex hormones	141, 145, 146
<i>Lucrin Depot 4 Month Injection (AB)</i>	190	MEFENAMIC ACID	241
<i>Lumigan (AG)</i>	304	<i>Mefic (PC)</i>	241
<i>Lumin 10 (AF)</i>	279	<i>Mefix 310250 (MH)</i>	
<i>Lumin 20 (AF)</i>	279	.Repatriation Schedule	512
<i>Luvox (SM)</i>	277	<i>Megace (BQ)</i>	190
Lycinate (FM)		<i>Megafol 0.5 (AF)</i>	106
.Cardiovascular system	110	<i>Megafol 5 (AF)</i>	106
.Dental	335	MEGESTROL ACETATE	190
<i>Lyclear (PC)</i>	290	<i>Melipramine (UW)</i>	276
<i>Lyofom C 603025 (SS)</i>		<i>Melizide (AF)</i>	96
.Repatriation Schedule	506	<i>Mellihexal (HX)</i>	96
<i>Lyofom Extra 603088 (SS)</i>		<i>Melolin 36101720 (SN)</i>	
.Repatriation Schedule	507	.Repatriation Schedule	511
<i>Lyofom Extra 603090 (SS)</i>		<i>Melolin 66974933 (SN)</i>	
.Repatriation Schedule	507	.Repatriation Schedule	511
<i>Lyofom Flat 603092 (SS)</i>		MELOXICAM	239
.Repatriation Schedule	506	MELPHALAN	181
<i>Lyofom Flat 603093 (SS)</i>		<i>Menorest 37.5 (NV)</i>	143
.Repatriation Schedule	506	<i>Menorest 50 (NV)</i>	144
<i>Lyofom Flat 603095 (SS)</i>		<i>Menorest 75 (NV)</i>	144
.Repatriation Schedule	506	<i>Menorest 100 (NV)</i>	145
M		<i>Meprazol (HX)</i>	82
<i>Mabthera (RO)</i>	187, 188	MERCAPTOPYRINE	182
<i>Macrodantin (PU)</i>	174	MESALAZINE	91
MACROGOL 3350	88	<i>Mesasal (GK)</i>	91
<i>Madopar (RO)</i>	266	MESNA	309
<i>Madopar 62.5 (RO)</i>	266	<i>Mestinon (VT)</i>	286
<i>Madopar 125 (RO)</i>	266	<i>Mestinon Timespan (VT)</i>	286
<i>Madopar HBS (RO)</i>	266	<i>Metabolic Mineral Mixture (SB)</i>	318
<i>Madopar Rapid 62.5 (RO)</i>	266	<i>Metalyse (BY)</i>	105
<i>Madopar Rapid 125 (RO)</i>	266	<i>Metamucil Regular (PY)</i>	
<i>Magicul 200 (AF)</i>	77	.Repatriation Schedule	473
<i>Magicul 400 (AF)</i>	78	<i>Metamucil Smooth Texture Orange (PY)</i>	
<i>Magicul 800 (AF)</i>	78	.Repatriation Schedule	473
<i>Magmin (BB)</i>		METFORMIN HYDROCHLORIDE	95
.Repatriation Schedule	475	METFORMIN HYDROCHLORIDE with	
MAGNESIUM ASPARTATE		GLIBENCLAMIDE	96
.Repatriation Schedule	475		
<i>Maosig (SI)</i>	279		

GENERIC /PROPRIETARY INDEX

METHADONE HYDROCHLORIDE		
.Nervous system	256	
.Section 100	428	
<i>Methoblastin (PH)</i>	182, 237	
<i>Methopt (SI)</i>	306	
METHOTREXATE	182, 237	
<i>Methotrexate Ebewe (IT)</i>	182	
METHYL SALICYLATE		
.Repatriation Schedule	491	
METHYLDOPA	112	
METHYLPHENIDATE HYDROCHLORIDE	280	
METHYLPREDNISOLONE ACEPONATE	136	
METHYLPREDNISOLONE ACETATE		
.Dental	336	
.Systemic hormonal preparations, excl. sex hormones and insulins	156	
METHYLPREDNISOLONE SODIUM SUCCINATE	156	
METHYSERGIDE	259	
METOCLOPRAMIDE HYDROCHLORIDE		
.Alimentary tract and metabolism	84	
.Dental	333	
.Doctor's Bag Supplies	72	
<i>Metohexal (HX)</i>	117	
<i>Metolol (DP)</i>	117	
METOPROLOL SUCCINATE	117	
METOPROLOL TARTRATE	117	
<i>Metrogyl 200 (AF)</i>		
.Anti-infectives for systemic use	174	
.Dental	345	
<i>Metrogyl 400 (AF)</i>		
.Anti-infectives for systemic use	174	
.Dental	345	
<i>Metrol 100 (AW)</i>	117	
<i>Metrol 50 (AW)</i>	117	
METRONIDAZOLE		
.Anti-infectives for systemic use	174	
.Dental	345	
.Repatriation Schedule	481	
METRONIDAZOLE BENZOATE		
.Anti-infectives for systemic use	174	
.Dental	345	
<i>Metronide 200 (HP)</i>		
.Anti-infectives for systemic use	174	
.Dental	345	
<i>Metronide 400 (HP)</i>		
.Anti-infectives for systemic use	174	
.Dental	345	
MEXILETINE HYDROCHLORIDE	108	
<i>Mexitil (BY)</i>	108	
<i>Miacalcic 50 (NV)</i>	158	
<i>Miacalcic 100 (NV)</i>	158	
MIANSERIN HYDROCHLORIDE	279	
<i>Micardis (BY)</i>	126	
<i>Micardis Plus 40/12.5 mg (BY)</i>	127	
<i>Micardis Plus 80/12.5 mg (BY)</i>	127	
MICONAZOLE		
.Repatriation Schedule	477	
MICONAZOLE NITRATE		
.Repatriation Schedule	478, 484	
<i>Microgynon 30 (SC)</i>	140	
<i>Microgynon 30 ED (SC)</i>	140	
<i>Microgynon 50 ED (SC)</i>	140	
<i>Microlax (PH)</i>		
.Alimentary tract and metabolism	89	
.Palliative Care	323	
.Repatriation Schedule	474	
<i>Microlut 28 (SC)</i>	141	
<i>Micronor (JC)</i>	141	
<i>Microval 28 (WY)</i>	141	
MILK POWDER—LACTOSE FREE FORMULA	313	
MILK POWDER—LACTOSE MODIFIED	314	
MILK POWDER—SYNTHETIC	314	
MILK PROTEIN and FAT FORMULA with VITAMINS and MINERALS—CARBOHYDRATE FREE	317	
<i>Minaphlex (SB)</i>	316	
<i>Minax 50 (AF)</i>	117	
<i>Minax 100 (AF)</i>	117	
MINERAL MIXTURE	318	
<i>Minidiab (PH)</i>	96	
<i>Minipress (PF)</i>	112, 113	
<i>Minirin (FP)</i>	154	
<i>Minirin Nasal Spray (FP)</i>	154	
<i>Minitran 5 (MM)</i>	111	
<i>Minitran 10 (MM)</i>	111	
<i>Minitran 15 (MM)</i>	111	
MINOCYCLINE	160	
<i>Minomycin-50 (SI)</i>	161	
MINOXIDIL	113	
<i>Mirena (SC)</i>	139	
MIRTAZAPINE	280	
<i>Mirtazon (AW)</i>	280	
MISOPROSTOL	80	
MITOZANTRONE HYDROCHLORIDE	186	
<i>Mixtard 20/80 Penfill 3 mL (NO)</i>	94	
<i>Mixtard 30/70 (NO)</i>	94	
<i>Mixtard 30/70 InnoLet (NI)</i>	94	
<i>Mixtard 30/70 Penfill 3 mL (NO)</i>	94	
<i>Mixtard 50/50 (NO)</i>	94	
<i>Mixtard 50/50 Penfill 3 mL (NO)</i>	94	
<i>Mobic (BY)</i>	239	
<i>Mobilis 10 (AF)</i>		
.Dental	347	
.Musculo-skeletal system	240	
<i>Mobilis 20 (AF)</i>		
.Dental	348	
.Musculo-skeletal system	240	
<i>Mobilis D-10 (AF)</i>		
.Dental	347	
.Musculo-skeletal system	239	
<i>Mobilis D-20 (AF)</i>		
.Dental	347	
.Musculo-skeletal system	239	
MOCLOBEMIDE	279	
MODAFINIL	281	
<i>Modavigil (CS)</i>	282	
<i>Moderate (BQ)</i>	268	
<i>Moduretic (MK)</i>	115	
<i>Mogadon (VT)</i>		
.Dental	356	
.Nervous system	262, 274	
.Palliative Care	331	
<i>Moheal (HX)</i>	279	

GENERIC /PROPRIETARY INDEX

MOMETASONE FUROATE			
.Dermatologicals	137		
.Repatriation Schedule	481		
<i>Monistat 7 (JC)</i>			
.Repatriation Schedule	484		
<i>Monistat Derm (JC)</i>			
.Repatriation Schedule	478		
<i>Monodur 60 mg (PM)</i>	111		
<i>Monodur 120 mg (PM)</i>	111		
<i>Monofeme 28 (WX)</i>	140		
<i>Monogen (SB)</i>	313		
<i>Monoplus 10/12.5 (BQ)</i>	125		
<i>Monoplus 20/12.5 (BQ)</i>	125		
<i>Monopril (BQ)</i>	123		
MONTELUKAST SODIUM	299		
MORPHINE HYDROCHLORIDE			
.Dental	350		
.Nervous system	250		
MORPHINE SULFATE			
.Dental	350		
.Doctor's Bag Supplies	72		
.Nervous system	251		
.Repatriation Schedule	492		
MORPHINE TARTRATE	253		
<i>Motilium (JC)</i>	84		
<i>Movalis 7.5 (AW)</i>	239		
<i>Movalis 15 (AW)</i>	239		
<i>Movicol (NE)</i>	88		
<i>Movox 50 (AL)</i>	277		
<i>Movox 100 (AF)</i>	277		
<i>Moxacin (CS)</i>			
.Antiinfectives for systemic use	161, 162		
.Dental	337		
<i>Moxacin 250 (CS)</i>			
.Antiinfectives for systemic use	162		
.Dental	338		
MOXIFLOXACIN HYDROCHLORIDE			
.Antiinfectives for systemic use	173		
.Repatriation Schedule	487		
<i>MS Contin (MF)</i>			
.Dental	351		
.Nervous system	252, 253		
.Repatriation Schedule	492		
<i>MS Contin Suspension 20 mg (MF)</i>			
.Dental	351		
.Nervous system	253		
<i>MS Contin Suspension 30 mg (MF)</i>			
.Dental	352		
.Nervous system	253		
<i>MS Contin Suspension 60 mg (MF)</i>			
.Dental	352		
.Nervous system	253		
<i>MS Contin Suspension 100 mg (MF)</i>			
.Dental	352		
.Nervous system	253		
<i>MS Contin Suspension 200 mg (MF)</i>	253		
<i>MS Mono (MF)</i>			
.Dental	351		
.Nervous system	252, 253		
<i>MSUD AID III (SB)</i>	317		
<i>MSUD Analog (SB)</i>	317		
<i>MSUD Express (VF)</i>	317		
<i>MSUD-gel (VF)</i>	317		
<i>MSUD Maxamald (SB)</i>	317		
<i>MSUD Maxamum (SB)</i>	317		
<i>Mucomyst (BQ)</i>	299		
<i>Muphoran (SE)</i>	181		
MUPIROCIIN			
.Repatriation Schedule	480		
<i>Murelax (FM)</i>			
.Dental	355		
.Nervous system	273		
.Palliative Care	331		
<i>Muric 500/125 (SL)</i>			
.Antiinfectives for systemic use	165		
.Dental	340		
<i>Muric 875/125 (SL)</i>			
.Antiinfectives for systemic use	165		
.Dental	340		
<i>Mycobutin (PH)</i>			
.Section 100	421		
MYCOPHENOLATE MOFETIL			
.Antineoplastic and immunomodulating agents	236		
.Section 100	414		
MYCOPHENOLATE SODIUM			
.Antineoplastic and immunomodulating agents	236		
.Section 100	414		
<i>Mycospor (BN)</i>			
.Repatriation Schedule	477		
<i>Mycostatin (BQ)</i>			
.Alimentary tract and metabolism	77		
.Dental	333		
.Repatriation Schedule	477		
<i>Myfortic (NV)</i>			
.Antineoplastic and immunomodulating agents	236		
.Section 100	414		
<i>Mylanta Double Strength (PC)</i>			
.Repatriation Schedule	472		
<i>Mylanta P (PC)</i>	77		
<i>Myleran (GK)</i>	181		
<i>Myocrisin (AV)</i>	242		
<i>Mysoline (LM)</i>	261		
N			
NAFARELIN ACETATE	154		
NALOXONE HYDROCHLORIDE			
.Dental	356		
.Doctor's Bag Supplies	72		
.Various	308		
<i>Naloxone Min-I-Jet (CS)</i>			
.Dental	356		
.Doctor's Bag Supplies	72		
.Various	308		
NALTREXONE HYDROCHLORIDE	287		
NANDROLONE DECANOATE	100		
<i>Napamide 2.5 mg (DP)</i>	113		
NAPHAZOLINE HYDROCHLORIDE			
.Repatriation Schedule	497		
NAPHAZOLINE HYDROCHLORIDE with PHENIRAMINE MALEATE			
.Repatriation Schedule	497		
<i>Naphcon-A (AQ)</i>			
.Repatriation Schedule	497		

GENERIC /PROPRIETARY INDEX

<i>Naphcon Forte</i> (AQ)		<i>Neulasta</i> (AN)	
.Repatriation Schedule	497	.Section 100	416
<i>Naprosyn</i> (RO)		<i>Neupogen</i> (AN)	
.Dental	348	.Section 100	379
.Musculo-skeletal system	240	<i>Neurontin</i> (PF)	
.Palliative Care	327, 328	.Nervous system	264
<i>Naprosyn SR750</i> (RO)		.Repatriation Schedule	492, 493
.Dental	348	NEVIRAPINE	
.Musculo-skeletal system	241	.Section 100	414
.Palliative Care	327, 328	<i>Nexcare Durable Cloth First Aid Tape 799</i> (MM)	
<i>Naprosyn SR1000</i> (RO)		.Repatriation Schedule	512
.Dental	348	<i>Nexcare Gentle Paper First Aid Tape 789</i> (MM)	
.Musculo-skeletal system	241	.Repatriation Schedule	512
.Palliative Care	327, 328	<i>Nexcare Tegaderm Transparent H1624</i> (MM)	
NAPROXEN		.Repatriation Schedule	505
.Dental	348	<i>Nexcare Tegaderm Transparent H1626</i> (MM)	
.Musculo-skeletal system	240	.Repatriation Schedule	505
.Palliative Care	327	<i>Nexium</i> (AP)	81
NAPROXEN SODIUM		<i>Nexium Hp7</i> (AP)	83
.Dental	349	<i>Nicabate CQ 7</i> (GK)	
.Musculo-skeletal system	241	.Repatriation Schedule	494
.Palliative Care	328	<i>Nicabate CQ 14</i> (GK)	
<i>Naramig</i> (GK)	260	.Repatriation Schedule	494
NARATRIPTAN HYDROCHLORIDE	260	<i>Nicabate CQ 21</i> (GK)	
<i>Nardil</i> (LM)	278	.Repatriation Schedule	494
<i>Natrilix</i> (SE)	113	NICORANDIL	112
<i>Natrilix SR</i> (SE)	113	<i>Nicorette Patch</i> (PH)	
<i>Navelbine</i> (FB)	184	.Repatriation Schedule	494
<i>Navoban</i> (NV)	85	NICOTINE	
<i>Nebcin</i> (AS)	171	.Repatriation Schedule	494
NEDOCROMIL SODIUM	297	<i>Nidem</i> (AW)	96
NELFINAVIR MESYLATE		NIFEDIPINE	
.Section 100	414	.Cardiovascular system	119
<i>Nemdyn</i> (HA)	307	.Repatriation Schedule	477
<i>Neocate</i> (SB)	311	<i>Nifehexal</i> (HX)	119
<i>Neocate Advance</i> (SB)	311	<i>Nilstat</i> (SI)	
<i>Neo-Cytamen</i> (MX)	106	.Alimentary tract and metabolism	77, 89
<i>Neo-Mercazole</i> (RO)	158	.Dental	333, 334
NEOMYCIN SULFATE	89	.Repatriation Schedule	483
NEOMYCIN UNDECENOATE with BACITRACIN		NILUTAMIDE	192
ZINC	307	NITRAZEPAM	
<i>Neoral</i> (NV)		.Dental	356
.Antineoplastic and immunomodulating agents	213	.Nervous system	262, 274
.Section 100	371	.Palliative Care	331
<i>Neoral 10</i> (NV)		<i>Nitro-Dur 5</i> (SH)	111
.Antineoplastic and immunomodulating agents	213	<i>Nitro-Dur 10</i> (SH)	111
.Section 100	371	<i>Nitro-Dur 15</i> (SH)	111
<i>Neoral 25</i> (NV)		NITROFURANTOIN	174
.Antineoplastic and immunomodulating agents	213	<i>Nitrolingual Pumpspray</i> (AV)	
.Section 100	371	.Cardiovascular system	110
<i>Neoral 50</i> (NV)		.Doctor's Bag Supplies	71
.Antineoplastic and immunomodulating agents	213	<i>Nitrostat</i> (LM)	
.Section 100	371	.Cardiovascular system	110
<i>Neoral 100</i> (NV)		.Dental	335
.Antineoplastic and immunomodulating agents	213	<i>Nizac</i> (LN)	79
.Section 100	371	NIZATIDINE	79
<i>Neosporin</i> (PC)	300	<i>Nizoral</i> (JC)	
<i>Neosulf</i> (AF)	89	.Antifungives for systemic use	175
<i>Neotigason</i> (RO)	134, 135	.Repatriation Schedule	477
<i>Neulactil</i> (AV)	269	<i>Nolvadex</i> (AP)	191
		<i>Nolvadex-D</i> (AP)	191

GENERIC /PROPRIETARY INDEX

<i>Nordette 28 (WY)</i>	140	<i>Nyefax 20 mg (DP)</i>	119
<i>Norditropin NordiFlex (NO)</i>		<i>Nyogel (NV)</i>	303
Section 100	425	<i>Nypine 10 (AW)</i>	119
<i>Norditropin SimpleXx (NO)</i>		<i>Nypine 20 (AW)</i>	119
Section 100	424, 425	NYSTATIN	
NORETHISTERONE	141, 146	Alimentary tract and metabolism	77, 89
NORETHISTERONE with ETHINYLOESTRADIOL		Dental	333, 334
.....	140, 141	Repatriation Schedule	477, 483
NORETHISTERONE with MESTRANOL	141	O	
<i>Norflohexal (HX)</i>	173	OCTREOTIDE	
NORFLOXACIN	173	Section 100	415
<i>Noriday 28 Day (PH)</i>	141	OCTREOTIDE ACETATE	
<i>Norimin 28 Day (KR)</i>	140	Section 100	415
<i>Norimin-1 28 Day (KR)</i>	140	<i>Ocufen (AG)</i>	302
<i>Norinyl-1 (PH)</i>	141	<i>Ocuflox (AG)</i>	301
<i>Norinyl-1/28 (PH)</i>	141	<i>Odrik (KN)</i>	125
<i>Normacol Plus (NE)</i>		OESTRADIOL	
Alimentary tract and metabolism	88	Genito urinary system and sex hormones	143
Palliative Care	322	Repatriation Schedule	485
Repatriation Schedule	474	OESTRADIOL HEMIHYDRATE	145
<i>Normison (SI)</i>		OESTRADIOL with NORETHISTERONE ACETATE	146
Dental	356	OESTRADIOL and OESTRADIOL with	
Nervous system	274, 275	DYDROGESTERONE	147
Palliative Care	332	OESTRADIOL and OESTRADIOL with	
<i>Noroxin (MK)</i>	173	NORETHISTERONE ACETATE	147
<i>Norprolac (FP)</i>	140	OESTRADIOL VALERATE	145
<i>Norspan (MF)</i>	257	OESTRADIOL VALERATE and OESTRADIOL	
NORTRIPTYLINE HYDROCHLORIDE	276	VALERATE with CYPROTERONE ACETATE	148
<i>Norvasc (PF)</i>		OESTRIOL	145
Cardiovascular system	118	OESTROGENS—CONJUGATED	145
Repatriation Schedule	476	OESTROGENS—CONJUGATED with	
<i>Norvir (AB)</i>		MEDROXYPROGESTERONE ACETATE	146
Section 100	421	OFLOXACIN	301
<i>Noten (AF)</i>	116	<i>Ogen .625 (PH)</i>	145
<i>Novantrone (SI)</i>	186	<i>Ogen 1.25 (PH)</i>	145
<i>Novasone (EX)</i>	137	OLANZAPINE	270
<i>NovoMix 30 FlexPen (NF)</i>	94	OLSALAZINE SODIUM	92
<i>NovoMix 30 Penfill 3 mL (NO)</i>	94	OMEPRAZOLE	82
<i>NovoRapid (NO)</i>	93	OMEPRAZOLE and CLARITHROMYCIN and	
<i>NovoRapid FlexPen (NF)</i>	93	AMOXYCILLIN	83
<i>NovoRapid Penfill 3 mL (NO)</i>	93	<i>Omnitest EZ (BR)</i>	309
<i>Nucolox (SI)</i>		<i>OncoTICE (OR)</i>	196
Repatriation Schedule	473	ONDANSETRON	85
<i>Nuelin (MM)</i>	298, 299	<i>Onkotrone (BX)</i>	186
<i>Nuelin-SR 200 (MM)</i>	299	<i>Op-Site Flexigrid 4629 (SN)</i>	
<i>Nuelin-SR 250 (MM)</i>	299	Repatriation Schedule	505
<i>Nuelin-SR 300 (MM)</i>	299	<i>Opticrom (AV)</i>	304
<i>Nufloxib (AF)</i>	173	<i>Optium glucose (MS)</i>	310
<i>Nu-Gel 2497 (JJ)</i>		<i>Orabase (BQ)</i>	
Repatriation Schedule	511	Repatriation Schedule	479
<i>Nupentin 100 (AF)</i>		<i>Oratane (DP)</i>	137
Nervous system	264	<i>Ordine 2 (MF)</i>	
Repatriation Schedule	492	Dental	350
<i>Nupentin 300 (AF)</i>		Nervous system	251
Nervous system	264	<i>Ordine 5 (MF)</i>	
Repatriation Schedule	493	Dental	350
<i>Nupentin 400 (AF)</i>		Nervous system	251
Nervous system	264	<i>Ordine 10 (MF)</i>	
Repatriation Schedule	493	Dental	350
<i>Nutraplus (GA)</i>		Nervous system	251
Repatriation Schedule	479		

ORLISTAT		Panadol Osteo (GC)	
.Repatriation Schedule	474	.Nervous system	259
<i>Oroxine (SI)</i>	157	.Palliative Care	329
<i>O.R.S. (AS)</i>	90	<i>Panafcort (AS)</i>	157
<i>Orudis (AV)</i>		<i>Panafcortelone (AS)</i>	156
.Dental	348	<i>Panamax (SW)</i>	
.Musculo-skeletal system	240	.Dental	354
<i>Orudis SR 200 (AV)</i>		.Nervous system	259
.Dental	348	<i>Panamax 240 Elixir (SW)</i>	
.Musculo-skeletal system	240	.Dental	354
<i>Oruvail SR (HP)</i>		.Nervous system	259
.Dental	348	<i>Panamax Co. (SW)</i>	
.Musculo-skeletal system	240	.Repatriation Schedule	492
<i>Ospolot (PL)</i>	265	PANCREATIC EXTRACT	93
<i>Otocomb Otac (BC)</i>	307	PANCRELIPASE	93
<i>Otodex (QM)</i>	307	PANTOPRAZOLE SODIUM SESQUIHYDRATE	82
<i>Ovestin (OR)</i>	145	<i>Panzyltrat 25000 (TM)</i>	93
<i>Ovestin Ovula (OR)</i>	145	PARACETAMOL	
OXALIPLATIN	187	.Dental	354
<i>Oxandrin (CS)</i>	100	.Nervous system	259
OXANDROLONE	100	.Palliative Care	329
OXAZEPAM		PARAFFIN	306
.Dental	355	<i>Parahexal (HX)</i>	
.Nervous system	273	.Dental	354
.Palliative Care	331	.Nervous system	259
OXCARBAZEPINE	263	<i>Paralgin (FM)</i>	
<i>Oxetine (HX)</i>	277	.Dental	354
<i>Oxis Turbuhaler (AP)</i>	291	.Nervous system	259
OPRENOLOL HYDROCHLORIDE	116	<i>Pariet (JC)</i>	83
OXYBUTYNYN HYDROCHLORIDE	153	<i>Parlodel (NV)</i>	
OXYCODONE		.Genito urinary system and sex hormones	139
.Dental	352	.Nervous system	267
.Nervous system	253	<i>Parmol (AW)</i>	
OXYCODONE HYDROCHLORIDE		.Dental	354
.Dental	352	.Nervous system	259
.Nervous system	254	<i>Parnate (LM)</i>	278
<i>OxyContin (MF)</i>		PAROXETINE HYDROCHLORIDE	277
.Dental	352, 353	<i>Paxam 0.5 (AF)</i>	
.Nervous system	255	.Nervous system	262
OXYMETAZOLINE HYDROCHLORIDE		.Palliative Care	329, 330
.Repatriation Schedule	495	<i>Paxam 2 (AF)</i>	
<i>OxyNorm (MF)</i>		.Nervous system	262
.Dental	352	.Palliative Care	329, 330
.Nervous system	254	<i>Paxtine (AF)</i>	277
<i>OxyNorm Liquid 5mg/5mL (MF)</i>		<i>Peg 7420 (BK)</i>	
.Dental	352	.Repatriation Schedule	502
.Nervous system	254	<i>Peg 7422 (BK)</i>	
P		.Repatriation Schedule	502
<i>PAA (CV)</i>	305	<i>Peg 7423 (BK)</i>	
PACLITAXEL	185	.Repatriation Schedule	502
<i>Paclitaxel Ebewe (IT)</i>	185	<i>Peg 7425 (BK)</i>	
<i>Pamacid 20 (AF)</i>	78	.Repatriation Schedule	502
<i>Pamacid 40 (AF)</i>	79	PEG-Intron Redipen (SH)	
<i>Pamisol (MX)</i>		.Section 100	417
.Musculo-skeletal system	245	<i>Pegasys (RO)</i>	
.Section 100	373	.Section 100	417
<i>Panadeine Forte (SW)</i>		<i>Pegasys RBV (RO)</i>	
.Dental	349	.Section 100	418, 419
.Nervous system	249	<i>Pegatron (SH)</i>	
<i>Panadol (GC)</i>		.Section 100	420, 421
.Palliative Care	329		

PEGFILGRASTIM		Phosphate Sandoz (NV)	309
.Section 100	416	Physeptone (GK)	256
PEGINTERFERON ALFA-2a		PILOCARPINE HYDROCHLORIDE	302
.Section 100	416	<i>Pilopt (PE)</i>	302, 303
PEGINTERFERON ALFA-2b		PIMECROLIMUS	138
.Section 100	417	PINDOLOL	116
PEMETREXED DISODIUM		PINE TAR with CADE OIL, COAL TAR SOLUTION, ARACHIS OIL EXTRACT OF CRUDE COAL TAR and OLEYL ALCOHOL	
.Special Pharmaceutical Benefit	75	.Repatriation Schedule	482
<i>Pendine 300 (AL)</i>	264	PINE TAR with TRIETHANOLAMINE LAURYL SULFATE	
<i>Pendine 400 (AL)</i>	264	.Repatriation Schedule	480
<i>Pendine 800 (AF)</i>	264	<i>Pinetarsol (EO)</i>	
<i>Penhexal VK (HX)</i>		.Repatriation Schedule	480
.Antiinfectives for systemic use	163	PIOGLITAZONE HYDROCHLORIDE	97
.Dental	338	PIPERAZINE OESTRONE SULFATE	145
PENICILLAMINE	242	<i>Pirohexal-D (HX)</i>	
<i>Pentasa (FP)</i>	91, 92	.Dental	347
<i>Peppidine (MK)</i>	79	.Musculo-skeletal system	239
<i>Peppidine M (MK)</i>	78	PIROXICAM	
<i>Pepti-Junior (NU)</i>	312	.Dental	347
<i>Pepzan (DP)</i>	78, 79	.Musculo-skeletal system	239
PERGOLIDE MESYLATE	268	PIZOTIFEN MALATE	261
PERHEXILINE MALEATE	112	<i>PK AID II (SB)</i>	315
<i>Periactin (FR)</i>	261	<i>PK Max (SB)</i>	316
PERICYAZINE	269	<i>PKU-Express (VF)</i>	316
PERINDOPRIL ERBUMINE	124	<i>PKU Express Liquid (VF)</i>	316
PERINDOPRIL ERBUMINE with INDAPAMIDE HEMIHYDRATE	125	<i>PKU-gel (VF)</i>	316
<i>Periogard (Chlorohex) Mouth Rinse (OM)</i>		<i>Placil (AF)</i>	274, 275
.Repatriation Schedule	472	<i>Plaqacide (OB)</i>	
<i>Permax (AS)</i>	268	.Repatriation Schedule	472
PERMETHRIN	290	<i>Plaquenil (SW)</i>	242
<i>Persantin SR (BY)</i>	103	<i>Plasma-Lyte 148 (BX)</i>	107
PETHIDINE HYDROCHLORIDE		<i>Plavix (SW)</i>	
.Doctor's Bag Supplies	72	.Blood and blood forming organs	103
.Dental	353	.Repatriation Schedule	475
.Nervous system	256	<i>Plendil ER (AP)</i>	119
<i>Petrus Bisacodyl Suppositories (PP)</i>		PNEUMOCOCCAL VACCINE, POLYVALENT	180
.Alimentary tract and metabolism	87	<i>Pneumovax 23 (CS)</i>	180
.Palliative Care	321	PODOPHYLLOTOXIN	
<i>Pexsig (SI)</i>	112	.Repatriation Schedule	481
<i>Pharmorubicin Solution (PH)</i>	186	<i>Poly Gel (AQ)</i>	304
PHENELZINE SULFATE	278	<i>Poly Visc (IQ)</i>	306
<i>Phenergan (AV)</i>		POLYETHYLENE GLYCOL 400 with PROPYLENE GLYCOL	306
.Palliative Care	320, 321	POLYGLINE	107
.Repatriation Schedule	496	POLYMYXIN B SULFATE with BACITRACIN and NEOMYCIN SULFATE	300
<i>Phenex-1 (AB)</i>	316	<i>Polytar (SX)</i>	
<i>Phenex-2 (AB)</i>	316	.Repatriation Schedule	482
PHENOBARBITONE	261	<i>Poly-Tears (IQ)</i>	306
PHENOBARBITONE SODIUM	261	POLYVINYL ALCOHOL	307
PHENOXYBENZAMINE HYDROCHLORIDE		<i>Ponstan (PD)</i>	241
.Cardiovascular system	115	<i>Posalfilin (NE)</i>	
.Genito urinary system and sex hormones	153	.Repatriation Schedule	482
PHENOXYMETHYLPENICILLIN		POTASSIUM CHLORIDE	99
.Antiinfectives for systemic use	163	POVIDONE-IODINE	
.Dental	338	.Repatriation Schedule	482
PHENYTOIN	261		
PHENYTOIN SODIUM	261		
<i>Phlexy-10 (SB)</i>	315		
<i>Phlexy-10 Drink Mix (SB)</i>	315		
PHOLCODINE			
.Repatriation Schedule	496		

GENERIC /PROPRIETARY INDEX

<i>Pramin (AF)</i>		<i>Procur 100 (DP)</i>	
Alimentary tract and metabolism.....	84	Antineoplastic and immunomodulating agents	192
Dental.....	333	Genito urinary system and sex hormones	152
<i>Prantal (SH)</i>		<i>Prodeine 15 (SW)</i>	
Repatriation Schedule	482	Repatriation Schedule	492
<i>Pravachol (BQ)</i>	129	<i>Prodeine Forte (DK)</i>	
PRAVASTATIN SODIUM.....	129	Dental.....	349
<i>Prazohexal (HX)</i>	112, 113	Nervous system	249
PRAZOSIN HYDROCHLORIDE.....	112	<i>Profloxin (HX)</i>	172
<i>Precision Plus (MS)</i>	310	<i>Profore 66050016 (SN)</i>	
<i>PredMix (LN)</i>	156	Repatriation Schedule	501
<i>Prednefrin Forte (AG)</i>	301	<i>Profore Lite 66050415 (SN)</i>	
PREDNISOLONE.....	156	Repatriation Schedule	501
PREDNISOLONE ACETATE with PHENYLEPHRINE		PROGESTERONE	
HYDROCHLORIDE.....	301	Section 100	427
PREDNISOLONE SODIUM PHOSPHATE		<i>Progout 100 (AF)</i>	244
Alimentary tract and metabolism.....	91	<i>Progout 300 (AF)</i>	244
Systemic hormonal preparations, excl. sex		<i>Prograf (JC)</i>	
hormones and insulins.....	156	Antineoplastic and immunomodulating agents	237
PREDNISON.....	157	Section 100	422
<i>Predsol (SI)</i>	91	<i>ProGuide 66000780 (SN)</i>	
<i>Predsolone (LN)</i>	156	Repatriation Schedule	502
<i>Predsone (LN)</i>	157	<i>ProGuide 66000781 (SN)</i>	
<i>Pregnyl (OR)</i>		Repatriation Schedule	502
Genito urinary system and sex hormones	151	<i>ProGuide 66000782 (SN)</i>	
Section 100.....	427	Repatriation Schedule	502
<i>Premarin (WY)</i>	145	<i>Progynova (SC)</i>	145
<i>Premia 2.5 Continuous (WY)</i>	146	<i>Proladone (PL)</i>	
<i>Premia 5 Continuous (WY)</i>	146	Dental.....	352
<i>Presolol 100 (AF)</i>	118	Nervous system	254
<i>Presolol 200 (AF)</i>	118	PROMETHAZINE HYDROCHLORIDE	
<i>Pressin 1 (AF)</i>	112	Dental.....	334
<i>Pressin 2 (AF)</i>	112	Doctor's Bag Supplies	72
<i>Pressin 5 (AF)</i>	113	Palliative Care.....	320
PRESSURE REDUCING PRODUCTS		Repatriation Schedule	496
Repatriation Schedule	512	Respiratory system	299
PRIMIDONE.....	261	PROPANTHELINE BROMIDE	153
<i>Primogyn Depot (SC)</i>	145	<i>Pro-Phree (AB)</i>	317
<i>Primolut N (SC)</i>	146	<i>Propine (AG)</i>	302
<i>Primoteston Depot (SC)</i>	142	PROPRANOLOL HYDROCHLORIDE.....	116
<i>Prinivil 5 (MK)</i>	123	PROPYLTHIOURACIL	158
<i>Prinivil 10 (MK)</i>	124	<i>Proquin (DP)</i>	172
<i>Prinivil 20 (MK)</i>	124	<i>Proscar (MK)</i>	
<i>Pro-Banthine (SI)</i>	153	Repatriation Schedule	486
PROBENECID.....	244	<i>Protaphane (NO)</i>	94
<i>Probitor (SZ)</i>	82	<i>Protaphane InnoLet (NI)</i>	94
PROCAINE PENICILLIN		<i>Protaphane NovoLet 3 mL (NL)</i>	94
Antiinfectives for systemic use.....	163	<i>Protaphane Penfill 3 mL (NO)</i>	94
Dental.....	339	PROTEIN HYDROLYSATE FORMULA with MEDIUM	
Doctor's Bag Supplies	71	CHAIN TRIGLYCERIDES	312
PROCHLORPERAZINE		<i>Prothiaden (AB)</i>	275
Alimentary tract and metabolism.....	86	<i>Provera (PH)</i>	
Dental.....	333	Antineoplastic and immunomodulating agents	189
Doctor's Bag Supplies	72	Genito urinary system and sex hormones	145, 146
<i>Pro-Cid (PL)</i>	244	<i>Proxen SR 750 (MD)</i>	
<i>Proctosedyl (AV)</i>		Dental.....	348
Repatriation Schedule	476	Musculo-skeletal system	241
<i>Procur (DP)</i>		Palliative Care.....	327, 328
Antineoplastic and immunomodulating agents	192		
Genito urinary system and sex hormones	151, 152		

GENERIC /PROPRIETARY INDEX

<i>Proxen SR 1000 (MD)</i>		<i>Qvar 50 (MM)</i>	294
.Dental	348	<i>Qvar 50 Autohaler (MM)</i>	295
.Musculo-skeletal system	241	<i>Qvar 100 (MM)</i>	294
.Palliative Care	327, 328	<i>Qvar 100 Autohaler (MM)</i>	295
<i>Prozac 20 (LY)</i>	277	R	
<i>Prozac Tab (LY)</i>	276	RABEPRAZOLE SODIUM	83
PSEUDOEPHEDRINE HYDROCHLORIDE		<i>Rafen 200 (AF)</i>	
.Repatriation Schedule	496	.Dental	348
PSYLLIUM HYDROPHILIC MUCILLOID		.Musculo-skeletal system	240
.Repatriation Schedule	473	.Palliative Care	327
PSYLLIUM HYDROPHILIC MUCILLOID with HIGH		<i>Ralovera (KR)</i>	145, 146
AMYLOSE MAIZE STARCH		RALOXIFENE HYDROCHLORIDE	248
.Repatriation Schedule	473	RALTITREXED	182
<i>Pulmicort Respules (AP)</i>	295	<i>Ramace 1.25 mg (ML)</i>	124
<i>Pulmicort Turbuhaler (AP)</i>	295	<i>Ramace 2.5 mg (ML)</i>	124
<i>Pulmozyme (RO)</i>		<i>Ramace 5 mg (ML)</i>	124
.Section 100	374	RAMIPRIL	
<i>Puregon 100 IU/0.5 mL (OR)</i>		.Cardiovascular system	124, 125
.Genito urinary system and sex hormones	149, 150	.Repatriation Schedule	477
.Section 100	426	<i>Rani 2 (AF)</i>	79, 80
<i>Puregon 150 IU/0.5 mL (OR)</i>		<i>Ranihexal (HX)</i>	79, 80
.Genito urinary system and sex hormones	149, 150	RANITIDINE HYDROCHLORIDE	
.Section 100	426	.Alimentary tract and metabolism	79, 80
<i>Puregon 200 IU/0.5 mL (OR)</i>		.Repatriation Schedule	473
.Genito urinary system and sex hormones	150	<i>Ranoxyl (DP)</i>	79, 80
.Section 100	426	<i>Rapamune (WY)</i>	
<i>Puregon 300 IU/0.36 mL (OR)</i>		.Antineoplastic and immunomodulating agents	237
.Genito urinary system and sex hormones	149, 150	.Section 100	422
.Section 100	426	<i>Rapilysin 10 U (RO)</i>	105
<i>Puregon 600 IU/0.72 mL (OR)</i>		<i>RCF (AB)</i>	318
.Genito urinary system and sex hormones	149, 150	<i>Rebif 44 (SG)</i>	195
.Section 100	426	REBOXETINE MESILATE	280
<i>Puregon 900 IU/1.08 mL (OR)</i>		<i>Redipred (AS)</i>	156
.Genito urinary system and sex hormones	149, 150	<i>Refresh Liquigel (AG)</i>	305
.Section 100	426	<i>Refresh Tears Plus (AG)</i>	305
<i>Purinethol (GK)</i>	182	<i>Remeron (OR)</i>	280
<i>P.V. Carpine (AG)</i>	302, 303	<i>Remicade (SH)</i>	
<i>PVA Forte (PE)</i>	307	.Repatriation Schedule	490
<i>PVA Tears (PE)</i>	307	.Section 100	401, 409
<i>Pyralin EN (KR)</i>	92	<i>Reminyl (JC)</i>	285
PYRANTEL EMBONATE	290	<i>Renitec (MK)</i>	123
PYRIDOSTIGMINE BROMIDE	286	<i>Renitec 20 (MK)</i>	123
PYRIMETHAMINE	289	<i>Renitec M (MK)</i>	122
Q		<i>Renitec Plus 20/6 (MK)</i>	125
<i>Questran Lite (BQ)</i>	132	<i>ReoPro (LY)</i>	102
QUETIAPINE FUMARATE	270	<i>Repalyte New Formulation (AV)</i>	90
<i>Quilonum SR (GK)</i>	279	<i>Replicare Ultra 66000434 (SN)</i>	
QUINAGOLIDE HYDROCHLORIDE	140	.Repatriation Schedule	509
QUINAPRIL HYDROCHLORIDE	124	<i>Replicare Ultra 66000435 (SN)</i>	
QUINAPRIL HYDROCHLORIDE with		.Repatriation Schedule	510
HYDROCHLOROTHIAZIDE	126	<i>Replicare Ultra 66000437 (SN)</i>	
<i>Quinate (AS)</i>	289	.Repatriation Schedule	510
<i>Quinbisul (AS)</i>	289	<i>Rescriptor (PF)</i>	
QUINIDINE BISULFATE	108	.Section 100	372
QUININE BISULFATE	289	<i>Resonium-A (SW)</i>	
QUININE SULFATE	289	.Repatriation Schedule	498
<i>Quinsul (LN)</i>	289	<i>Resprim (AF)</i>	
<i>QuitX (AF)</i>		.Antiinfectives for systemic use	169, 170
.Repatriation Schedule	494	.Dental	344
<i>QV Bath Oil (EO)</i>			
.Repatriation Schedule	479		

GENERIC /PROPRIETARY INDEX

<i>Resprim Forte (AF)</i>		
. Antiinfectives for systemic use	170	
. Dental	344	
<i>Restore CalciCare 9937 (HO)</i>		
. Repatriation Schedule	505	
<i>Restore CalciCare 9938 (HO)</i>		
. Repatriation Schedule	504	
<i>Restore CalciCare 9940 (HO)</i>		
. Repatriation Schedule	504	
<i>Restore Extra Thin 9921 (HO)</i>		
. Repatriation Schedule	509	
<i>Restore Plus 9956 (HO)</i>		
. Repatriation Schedule	509	
<i>Restore Plus 9957 (HO)</i>		
. Repatriation Schedule	510	
<i>Restore Plus 9958 (HO)</i>		
. Repatriation Schedule	510	
<i>Restore Plus Sacral 9959 (HO)</i>		
. Repatriation Schedule	510	
RETEPLASE (Recombinant plasminogen activator)	105	
<i>Retrovir (GK)</i>		
. Section 100	423	
<i>ReVia (BQ)</i>	287	
<i>Reyataz (BQ)</i>		
. Section 100	360	
RIBAVIRIN and PEGINTERFERON ALFA-2a		
. Section 100	418	
RIBAVIRIN and PEGINTERFERON ALFA-2b		
. Section 100	419	
RICINOLEIC ACID with ACETIC ACID and HYDROXYQUINOLINE SULFATE		
. Repatriation Schedule	484	
<i>Ridaura (LM)</i>	242	
RIFABUTIN		
. Section 100	421	
<i>Rifadin (AV)</i>	176	
RIFAMPICIN	176	
<i>Rilutek (AV)</i>	288	
RILUZOLE	288	
<i>Rimycin 150 (AF)</i>	176	
<i>Rimycin 300 (AF)</i>	176	
RISEDRONATE SODIUM		
. Musculo-skeletal system	246	
. Repatriation Schedule	491	
<i>Risperdal (JC)</i>	271	
<i>Risperdal Consta (JC)</i>	271, 272	
<i>Risperdal Quicklet (JC)</i>	271	
RISPERIDONE	271	
<i>Ritalin 10 (NV)</i>	280	
<i>Rithmik 100 (AW)</i>	109	
<i>Rithmik 200 (AW)</i>	109	
RITONAVIR		
. Section 100	421	
RITUXIMAB	187	
RIVASTIGMINE HYDROGEN TARTRATE	285	
<i>Rivotril (RO)</i>		
. Nervous system	262	
. Palliative Care	329, 330	
<i>Roaccutane (RO)</i>	137	
<i>Rocaltrol (RO)</i>		
. Alimentary tract and metabolism	99	
. Musculo-skeletal system	247	
<i>Rocephin (RO)</i>	169	
<i>Roferon-A (RO)</i>		
. Antineoplastic and immunomodulating agents	193, 194	
. Section 100	409, 410	
ROSIGLITAZONE MALEATE	98	
<i>Roxin (AW)</i>	173	
ROXITHROMYCIN	171	
<i>Rozex (GA)</i>		
. Repatriation Schedule	481	
<i>Rulide (AV)</i>	171	
<i>Rulide D (AV)</i>	171	
<i>Rynacrom (AV)</i>		
. Repatriation Schedule	495	
<i>Rythmodan (AV)</i>	108	
S		
<i>S-26 LF (WY)</i>	313, 314	
<i>Sabril (AV)</i>	264	
<i>Saizen 8 mg click.easy (SG)</i>		
. Section 100	425	
<i>Salazopyrin (PH)</i>	92	
<i>Salazopyrin-EN (PH)</i>	92	
SALBUTAMOL SULFATE		
. Doctor's Bag Supplies	72, 73	
. Respiratory system	291, 298	
SALCATONIN	158	
SALICYLIC ACID with BENZALKONIUM CHLORIDE, ALCOHOL and POLYOXYETHYLENE ETHERS		
. Repatriation Schedule	482	
SALICYLIC ACID with BENZALKONIUM CHLORIDE, ALCOHOL, COAL TAR and POLYOXYETHYLENE ETHERS		
. Repatriation Schedule	482	
SALICYLIC ACID with COAL TAR SOLUTION, PINE TAR and UNDECYLENAMIDE		
. Repatriation Schedule	482	
SALICYLIC ACID with PODOPHYLLIN RESIN		
. Repatriation Schedule	482	
SALMETEROL XINAFOATE	292	
<i>Salofalk (OA)</i>	91, 92	
<i>Sandimmun (NV)</i>		
. Section 100	370	
<i>Sandomigran 0.5 (NV)</i>	261	
<i>Sandostatin 0.05 (NV)</i>		
. Section 100	415	
<i>Sandostatin 0.1 (NV)</i>		
. Section 100	415	
<i>Sandostatin 0.5 (NV)</i>		
. Section 100	415	
<i>Sandostatin LAR (NV)</i>		
. Section 100	415	
<i>Sandrena (OR)</i>	143	
SAQUINAVIR		
. Section 100	421	
SAQUINAVIR MESYLATE		
. Section 100	421	
<i>Savacol Mouth and Throat Rinse (OM)</i>		
. Repatriation Schedule	472	

GENERIC /PROPRIETARY INDEX

<i>SciTropin</i> (SA)		SILDENAFIL CITRATE	
Section 100	424	Repatriation Schedule	485
<i>Seaze</i> 5 (AW)	265	<i>Silic</i> 15 (EO)	
<i>Seaze</i> 25 (AW)	265	Repatriation Schedule	479
<i>Seaze</i> 50 (AW)	265	<i>Silvazine</i> (SN)	135
<i>Seaze</i> 100 (AW)	265	SILVER SULFADIAZINE with CHLORHEXIDINE	
<i>Seaze</i> 200 (AW)	265	GLUCONATE	135
<i>SebiRinse</i> Conditioner (EO)		<i>Simplotan</i> (GP)	174
Repatriation Schedule	483	<i>Simvabell</i> (BF)	129, 130, 131
<i>Sebitar</i> (EO)		<i>Simvahexal</i> (HX)	129, 130, 131
Repatriation Schedule	482	<i>Simvar</i> 10 (AW)	130
<i>Sebizole</i> (DP)		<i>Simvar</i> 20 (AW)	130
Repatriation Schedule	477	<i>Simvar</i> 40 (AW)	130
SELEGILINE HYDROCHLORIDE	268	<i>Simvar</i> 80 (AW)	131
SELENIUM SULFIDE		SIMVASTATIN	129
Repatriation Schedule	482	<i>Simvastatin-DP</i> (DP)	130, 131
<i>Selgene</i> (AF)	268	<i>Simvastatin-RL</i> (RE)	129, 130, 131
<i>Selsun</i> (AB)		<i>Sinemet</i> (MK)	266
Repatriation Schedule	482	<i>Sinemet</i> 100/25 (MK)	266
SENEGA and AMMONIA		<i>Sinemet</i> CR (MK)	266
Repatriation Schedule	496	<i>Sinequan</i> (PF)	275
SENNA STANDARDISED		<i>Singulair</i> (MK)	299
Repatriation Schedule	473	SIROLIMUS	
<i>Senokot</i> (RC)		Antineoplastic and immunomodulating agents	237
Repatriation Schedule	473	Section 100	422
<i>SensoCard</i> (PX)	310	<i>Sitriol</i> (AF)	
<i>Septin</i> (SI)		Alimentary tract and metabolism	99
Antiinfectives for systemic use	169, 170	Musculo-skeletal system	247
Dental	344	<i>Skelid</i> (MX)	246
<i>Septin Forte</i> (SI)		SKIN CLEANSER	
Antiinfectives for systemic use	170	Repatriation Schedule	483
Dental	344	SKIN EMOLLIENT	
<i>Serenace</i> (SI)		Repatriation Schedule	479
Doctor's Bag Supplies	71	<i>Slow-K</i> (NV)	99
Nervous system	269	<i>Sodibic</i> (AS)	
<i>Serepax</i> (SI)		Repatriation Schedule	486
Dental	355	SODIUM ACID PHOSPHATE	309
Nervous system	273	SODIUM ALGINATE with CALCIUM CARBONATE	
Palliative Care	331	and SODIUM BICARBONATE	83
<i>Seretide Accuhaler</i> 100/50 (GK)	294	SODIUM AUROTHIOMALATE	242
<i>Seretide Accuhaler</i> 250/50 (GK)	294	SODIUM BICARBONATE	
<i>Seretide Accuhaler</i> 500/50 (GK)	294	Repatriation Schedule	486
<i>Seretide MDI</i> 50/25 (GK)	294	SODIUM CHLORIDE	
<i>Seretide MDI</i> 125/25 (GK)	294	Blood and blood forming organs	107
<i>Seretide MDI</i> 250/25 (GK)	294	Dental	334
<i>Serevent</i> (GK)	292	Repatriation Schedule	476
<i>Serevent Accuhaler</i> (GK)	292	SODIUM CHLORIDE COMPOUND	107
<i>Serophene</i> (SG)	151	SODIUM CHLORIDE with GLUCOSE	
<i>Seroquel</i> (AP)	270	Blood and blood forming organs	107
SERTRALINE HYDROCHLORIDE	278	Dental	334
<i>Sertraline-DP</i> (DP)	278	SODIUM CITRO-TARTRATE	
<i>Setopress</i> 3504 (SS)		Repatriation Schedule	486
Repatriation Schedule	501	SODIUM CLODRONATE TETRAHYDRATE	246
<i>Setopress</i> 3505 (SS)		SODIUM CROMOGLYCATE	
Repatriation Schedule	501	Repatriation Schedule	495
<i>Sevredol</i> (MF)	251	Respiratory system	297
<i>Sigmacort</i> (SI)		Sensory organs	304
Dental	335	SODIUM LACTATE COMPOUND	107
Dermatologicals	135	SODIUM POLYSTYRENE SULFONATE	
<i>Sigmaxin</i> (FM)	108	Repatriation Schedule	498
<i>Sigmaxin-PG</i> (FM)	108	SODIUM VALPROATE	263

GENERIC /PROPRIETARY INDEX

<i>Soffban 7224 (BV)</i>		<i>Sporanox (JC)</i>	176
.Repatriation Schedule	501	<i>Stalevo 50/12.5/200mg (NV)</i>	267
<i>Sofradex (AV)</i>	307	<i>Stalevo 100/25/200mg (NV)</i>	267
<i>Soframycin (AV)</i>	307	<i>Stalevo 150/37.5/200mg (NV)</i>	267
<i>SofTact (MS)</i>	310	<i>Staphylex 250 (AF)</i>	
<i>Solavert (AW)</i>	109	.Anti-infectives for systemic use	164
<i>Solian 100 (SW)</i>	270	.Dental	339
<i>Solian 200 (SW)</i>	270	<i>Staphylex 500 (AF)</i>	
<i>Solian 400 (SW)</i>	270	.Anti-infectives for systemic use	164
<i>Solian Solution (SW)</i>	270	.Dental	340
<i>Solone (FM)</i>	156	STAVUDINE	
<i>SoloSite Gel 36361338 (SN)</i>		.Section 100	422
.Repatriation Schedule	510	<i>Stelax 10 (AW)</i>	243
<i>Solprin (RC)</i>		<i>Stelax 25 (AW)</i>	243
.Blood and blood forming organs	102	<i>Stelazine (LM)</i>	268
.Dental	354	<i>Stemetil (AV)</i>	
.Nervous system	258	.Alimentary tract and metabolism	86
<i>Solu-Cortef (PH)</i>		.Dental	333
.Dental	335	.Doctor's Bag Supplies	72
.Doctor's Bag Supplies	71	<i>Stemzine (HP)</i>	
.Systemic hormonal preparations, excl. sex		.Alimentary tract and metabolism	86
hormones and insulins	156	.Dental	333
<i>Solugel 10336 (JJ)</i>		STERCULIA with FRANGULA BARK	
.Repatriation Schedule	510	.Alimentary tract and metabolism	88
<i>Solu-Medrol (PF)</i>	156	.Palliative Care	322
<i>Somac (PH)</i>	82, 83	.Repatriation Schedule	474
SOMATROPIN (Recombinant human growth hormone)		<i>Steripaste 3610 (SS)</i>	
.Section 100	424	.Repatriation Schedule	504
<i>Somatuline Autogel (IS)</i>		<i>Stieprox Liquid (SX)</i>	
.Section 100	412	.Repatriation Schedule	478
<i>Somatuline LA (IS)</i>		<i>Stocrin (MK)</i>	
.Section 100	412	.Section 100	374
<i>Sone (FM)</i>	157	<i>Stromectol (MK)</i>	290
<i>Sorbidin (AF)</i>	111	<i>Subutex (RC)</i>	
SORBITOL with SODIUM CITRATE and SODIUM		.Section 100	427
LAURYL SULFOACETATE		SUCRALFATE	84
.Alimentary tract and metabolism	89	<i>Sudafed Sinus & Nasal Decongestant (PC)</i>	
.Palliative Care	323	.Repatriation Schedule	496
.Repatriation Schedule	474	SULFACETAMIDE SODIUM	300
<i>Sorbsan 1410 (UM)</i>		SULFASALAZINE	92
.Repatriation Schedule	505	SULINDAC	
<i>Sorbsan 1411 (UM)</i>		.Dental	347
.Repatriation Schedule	504	.Musculo-skeletal system	239
<i>Sotab (DP)</i>	109	.Palliative Care	326
<i>Sotacor (BQ)</i>	109	SULTHIAME	265
<i>Sotahexal (HX)</i>	109	SUMATRIPTAN	260
SOTALOL HYDROCHLORIDE	109	SUMATRIPTAN SUCCINATE	260
SOY PROTEIN and FAT FORMULA with VITAMINS		SUNSCREENS	
and MINERALS—CARBOHYDRATE FREE	318	.Repatriation Schedule	479
<i>Span-K (AS)</i>	99	<i>SunSense Cream SPF30+ (EO)</i>	
<i>Spenco Dermal Pad 10-553 (KC)</i>		.Repatriation Schedule	479
.Repatriation Schedule	512	<i>SunSense Ultra SPF30+ (EO)</i>	
<i>Spenco Dermal Pad 10-561 (KC)</i>		.Repatriation Schedule	480
.Repatriation Schedule	512	<i>Surepress 650947 (CC)</i>	
<i>Spiractin 25 (AF)</i>	115	.Repatriation Schedule	501
<i>Spiractin 100 (AF)</i>	115	<i>Surepress 650948 (CC)</i>	
<i>Spiriva (BY)</i>	297	.Repatriation Schedule	501
SPIRONOLACTONE	115	<i>Surgam (AV)</i>	241
<i>Sporahexal (HX)</i>		<i>Surgical Lubricating Gel (BI)</i>	
.Anti-infectives for systemic use	166	.Repatriation Schedule	511
.Dental	341	<i>Sustanon 100 (OR)</i>	142

GENERIC /PROPRIETARY INDEX

<i>Sustanon 250 (OR)</i>	142	<i>Tegretol Liquid (NV)</i>	
<i>Suvalan 50 (AW)</i>	260	.Dental	355
<i>Symbicort Turbuhaler 100/6 (AP)</i>	293	.Nervous system	263
<i>Symbicort Turbuhaler 200/6 (AP)</i>	293	<i>Telfa 1970C (KE)</i>	
<i>Symbicort Turbuhaler 400/12 (AP)</i>	293	.Repatriation Schedule	511
<i>Symmetrel 100 (NV)</i>	267	<i>Telfa 2140C (KE)</i>	
<i>Synacthen Depot 1 mg/1 mL (NV)</i>	154	.Repatriation Schedule	511
<i>Synarel (PH)</i>	154	<i>Telfa 6020C (KE)</i>	
<i>Synphasic (PH)</i>	141	.Repatriation Schedule	511
<i>Systane (AQ)</i>	306	<i>Telfa 7650C (KE)</i>	
T		.Repatriation Schedule	511
TACROLIMUS		<i>Telfa 8252F (KE)</i>	
.Antineoplastic and immunomodulating agents	237	.Repatriation Schedule	503
.Section 100	422	<i>Telfa 8253F (KE)</i>	
TADALAFIL		.Repatriation Schedule	503
.Repatriation Schedule	485	<i>Telfa 8254F (KE)</i>	
<i>Tagamet (GK)</i>	77, 78	.Repatriation Schedule	503
<i>Tagamet 800 Express (GK)</i>		<i>Telfast (AV)</i>	
.Alimentary tract and metabolism	78	.Repatriation Schedule	497
.Repatriation Schedule	472	<i>Telfast 120 (AV)</i>	
<i>Talam (AW)</i>	276	.Repatriation Schedule	497
<i>Talohexal (HX)</i>	276	TELMISARTAN	126
<i>Tambocor (MM)</i>	108	TELMISARTAN with HYDROCHLOROTHIAZIDE	127
<i>Tamosin (SI)</i>	191	<i>Telzir (GK)</i>	
<i>Tamoxen 10 mg (DP)</i>	191	.Section 100	380
<i>Tamoxen 20 mg (DP)</i>	191	Temaze (AF)	
TAMOXIFEN CITRATE	191	.Dental	356
<i>Tamoxifen Hexal (HX)</i>	191	.Nervous system	274, 275
<i>Tamsil (AW)</i>	134	.Palliative Care	332
TAMSULOSIN HYDROCHLORIDE		TEMAZEPAM	
.Repatriation Schedule	486	.Dental	356
TAPES—NON-WOVEN RETENTION (POLYACRYLATE)		.Nervous system	274
.Repatriation Schedule	512	.Palliative Care	332
TAPES—PLASTER ADHESIVE ELASTIC		<i>Temodal (SH)</i>	181, 182
.Repatriation Schedule	512	TEMOZOLOMIDE	181
TAPES—PLASTER ADHESIVE HYPOALLERGENIC		<i>Temtabs (FM)</i>	
.Repatriation Schedule	512	.Dental	356
<i>Taxol (BQ)</i>	185	.Nervous system	274, 275
<i>Taxotere (AV)</i>	184	.Palliative Care	332
<i>Tazac (AS)</i>	79	<i>TenderWet 24 Active (HR)</i>	
<i>Tears Naturale (AQ)</i>	306	.Repatriation Schedule	508
<i>Tegaderm Transparent 1628 (MM)</i>		<i>TenderWet Active Cavity (HR)</i>	
.Repatriation Schedule	505	.Repatriation Schedule	508
<i>Tegaderm Transparent Island 3582 (MM)</i>		TENECTEPLASE	105
.Repatriation Schedule	506	TENOFOVIR DISOPROXIL FUMARATE	
<i>Tegaderm Transparent Island 3586 (MM)</i>		.Section 100	422
.Repatriation Schedule	506	<i>Tenopt (SI)</i>	303
<i>Tegretol 100 (NV)</i>		<i>Tenormin (AP)</i>	116
.Dental	355	<i>Tensig (SI)</i>	116
.Nervous system	262	<i>Tensogrip 36361259 (BV)</i>	
<i>Tegretol 200 (NV)</i>		.Repatriation Schedule	503
.Dental	355	<i>Tensopress 66004347 (BV)</i>	
.Nervous system	263	.Repatriation Schedule	501
<i>Tegretol CR 200 (NV)</i>		<i>Tensopress 66004348 (BV)</i>	
.Dental	355	.Repatriation Schedule	501
.Nervous system	263	<i>Tequin (BQ)</i>	
<i>Tegretol CR 400 (NV)</i>		.Repatriation Schedule	487
.Dental	355	TERAZOSIN HYDROCHLORIDE	
.Nervous system	263	.Repatriation Schedule	486
		<i>Terbihexal (HX)</i>	134

GENERIC /PROPRIETARY INDEX

TERBINAFINE		
.Repatriation Schedule	478	
TERBINAFINE HYDROCHLORIDE		
.Dermatologicals	134	
.Repatriation Schedule	478	
Terbinafine-DP (DP)	134	
TERBUTALINE SULFATE		
.Doctor's Bag Supplies	73	
.Respiratory system	293, 298	
Teril (AF)		
.Dental	355	
.Nervous system	263	
Terry White Chemists Aciclovir (TW)	177	
Terry White Chemists Allopurinol (TW)	244	
Terry White Chemists Alprazolam (TW)	272	
Terry White Chemists Amiodarone (TW)	109	
Terry White Chemists Amoxycillin (TW)		
.Anti-infectives for systemic use	161, 162	
.Dental	337, 338	
Terry White Chemists Amoxycillin and Clavulanic Acid (TW)		
.Anti-infectives for systemic use	165	
.Dental	340	
Terry White Chemists Atenolol (TW)	116	
Terry White Chemists Baclofen (TW)	243	
Terry White Chemists Captopril (TW)	121, 122	
Terry White Chemists Cefaclor (TW)		
.Anti-infectives for systemic use	168	
.Dental	343	
Terry White Chemists Cefaclor CD (TW)		
.Anti-infectives for systemic use	168	
.Dental	342	
Terry White Chemists Cephalexin (TW)		
.Anti-infectives for systemic use	166, 167	
.Dental	341, 342	
Terry White Chemists Citalopram (TW)	276	
Terry White Chemists Clarithromycin (TW)	170	
Terry White Chemists Clomipramine (TW)	274, 275	
Terry White Chemists Clotrimazole 3 Day Cream (TW)		
.Repatriation Schedule	484	
Terry White Chemists Clotrimazole 6 Day Cream (TW)		
.Repatriation Schedule	484	
Terry White Chemists Diclofenac (TW)		
.Dental	346	
.Musculo-skeletal system	238	
.Palliative Care	324, 325	
Terry White Chemists Diltiazem (TW)	120	
Terry White Chemists Diltiazem CD (TW)	121	
Terry White Chemists Doxycycline (TW)		
.Anti-infectives for systemic use	159, 160	
.Dental	336	
Terry White Chemists Enalapril (TW)	122, 123	
Terry White Chemists Famotidine (TW)	78, 79	
Terry White Chemists Fluoxetine (TW)	277	
Terry White Chemists Frusemide (TW)	114	
Terry White Chemists Gemfibrozil (TW)	132	
Terry White Chemists Glimepiride (TW)	96	
Terry White Chemists Indapamide (TW)	113	
Terry White Chemists Ipratropium (TW)	296, 297	
Terry White Chemists Isosorbide Mononitrate (TW)	111	
Terry White Chemists Isotretinoin (TW)	137	
Terry White Chemists Lisinopril (TW)	123, 124	
Terry White Chemists Metformin (TW)	95	
Terry White Chemists Metoprolol (TW)	117	
Terry White Chemists Moclobemide (TW)	279	
Terry White Chemists Nifedipine (TW)	119	
Terry White Chemists Norfloxacin (TW)	173	
Terry White Chemists Oral Rehydration Salts (TW) ...	90	
Terry White Chemists Paracetamol (TW)		
.Dental	354	
.Nervous system	259	
Terry White Chemists Paroxetine (TW)	277	
Terry White Chemists Piroxicam (TW)		
.Dental	347, 348	
.Musculo-skeletal system	240	
Terry White Chemists Piroxicam Dispersible (TW)		
.Dental	347	
.Musculo-skeletal system	239	
Terry White Chemists Prazosin (TW)	112, 113	
Terry White Chemists Ranitidine (TW)	79, 80	
Terry White Chemists Salbutamol (TW)		
.Doctor's Bag Supplies	72, 73	
.Respiratory system	292	
Terry White Chemists Sertraline (TW)	278	
Terry White Chemists Simvastatin (TW)	130, 131	
Terry White Chemists Sotalol (TW)	109	
Terry White Chemists Tamoxifen (TW)	191	
Terry White Chemists Tramadol (TW)		
.Dental	353	
.Nervous system	257, 258	
Terry White Chemists Trimethoprim with Sulfamethoxazole DS (TW)		
.Anti-infectives for systemic use	170	
.Dental	344	
Tertroxin (SI)	157	
Testogel (SC)	142	
TESTOSTERONE	141	
TESTOSTERONE ENANTHATE	142	
TESTOSTERONE ESTERS	142	
TESTOSTERONE UNDECANOATE	142	
TETANUS VACCINE, ADSORBED		
.Anti-infectives for systemic use	180	
.Dental	346	
.Doctor's Bag Supplies	73	
TETRABENAZINE	270	
TETRACOSACTRIN	154	
Tet-Tox (CS)		
.Anti-infectives for systemic use	180	
.Dental	346	
.Doctor's Bag Supplies	73	
Teveten (SM)	126	
Teveten Plus 600/12.5 (SM)	126	
THEOPHYLLINE	298	
TheraTears (CX)	305	
THIAMINE HYDROCHLORIDE		
.Repatriation Schedule	475	
THIOGUANINE	183	
Thioprine (AF)	237	
THIURIDAZINE HYDROCHLORIDE	269	
THIOTEPA	181	
3TC (GK)		
.Section 100	411	

GENERIC /PROPRIETARY INDEX

THYROXINE SODIUM.....	157	<i>Tramal</i> (CS)	
TIAGABINE HYDROCHLORIDE.....	263	.Dental.....	353, 354
TIAPROFENIC ACID.....	241	.Nervous system.....	257, 258
TICARCILLIN with CLAVULANIC ACID		<i>Tramal 100</i> (CS)	
.Antiinfectives for systemic use.....	166	.Dental.....	354
.Dental.....	341	.Doctor's Bag Supplies.....	72
<i>Ticlid</i> (RO).....	104	.Nervous system.....	258
<i>Ticlopidine Hexal</i> (HX).....	104	<i>Tramal SR 100</i> (CS)	
TICLOPIDINE HYDROCHLORIDE.....	103	.Dental.....	353
<i>Tielle MT2440</i> (JJ)		.Nervous system.....	258
.Repatriation Schedule.....	508	<i>Tramal SR 150</i> (CS)	
<i>Tielle MT2442</i> (JJ)		.Dental.....	353
.Repatriation Schedule.....	508	.Nervous system.....	258
<i>Tilade CFC-Free</i> (AV).....	297	<i>Tramal SR 200</i> (CS)	
<i>Tilodene</i> (AF).....	104	.Dental.....	354
TILUDRONATE DISODIUM.....	246	.Nervous system.....	258
<i>Timentin</i> (GK)		<i>Trandate</i> (SI).....	118
.Antiinfectives for systemic use.....	166	TRANOLAPRIL.....	125
.Dental.....	341	TRANEXAMIC ACID.....	106
TIMOLOL MALEATE.....	303	<i>Transiderm-Nitro 25</i> (NV).....	111
<i>Timoptol</i> (FR).....	303	<i>Transiderm-Nitro 50</i> (NV).....	111
<i>Timoptol XE</i> (MK).....	303	TRANLYCYPROMINE SULFATE.....	278
<i>Tinaderm</i> (SH)		<i>Travatan</i> (AQ).....	304
.Repatriation Schedule.....	478	TRAVOPROST.....	304
TINIDAZOLE.....	174	TRIAMCINOLONE ACETONIDE	
TIOTROPIUM BROMIDE MONOHYDRATE.....	297	.Dental.....	336
TIROFIBAN HYDROCHLORIDE.....	104	.Dermatologicals.....	136
<i>Titralac</i> (MM)		.Systemic hormonal preparations, excl. sex	
.Repatriation Schedule.....	472	hormones and insulins.....	157
TOBRAMYCIN.....	300	TRIAMCINOLONE ACETONIDE with NEOMYCIN	
TOBRAMYCIN SULFATE.....	171	SULFATE, GRAMICIDIN and NYSTATIN	
<i>Tobrex</i> (AQ).....	300	.Repatriation Schedule.....	481
<i>Tofranil 10</i> (NV).....	276	.Sensory organs.....	307
<i>Tofranil 25</i> (NV).....	276	<i>Tricortone</i> (FM).....	136
TOLNAFTATE		<i>Trifeme 28</i> (WX).....	141
.Repatriation Schedule.....	478	TRIFLUOPERAZINE HYDROCHLORIDE.....	268
<i>Tolvon</i> (OR).....	279	TRIGLYCERIDES, MEDIUM CHAIN.....	310
<i>Tomudex</i> (AP).....	182	TRIGLYCERIDES—MEDIUM CHAIN, FORMULA.....	313
<i>Topace</i> (FM).....	121, 122	TRIGLYCERIDES, MEDIUM CHAIN and LONG	
<i>Topamax</i> (JC)		CHAIN with GLUCOSE POLYMER.....	318
.Special Pharmaceutical Benefit.....	76	<i>Trileptal</i> (NV).....	263
<i>Topamax Sprinkle</i> (JC)		TRIMETHOPRIM.....	169
.Special Pharmaceutical Benefit.....	76	TRIMETHOPRIM with SULFAMETHOXAZOLE	
TOPIRAMATE		.Antiinfectives for systemic use.....	169
.Special Pharmaceutical Benefit.....	76	.Dental.....	344
TOPOTECAN HYDROCHLORIDE.....	189	<i>Triphasil 28</i> (WY).....	141
<i>Toprol-XL 23.75</i> (AP).....	117	<i>Triprim</i> (SI).....	169
<i>Toprol-XL 47.5</i> (AP).....	117	<i>Triquilar ED</i> (SC).....	141
<i>Toprol-XL 95</i> (AP).....	117	<i>Trisequens</i> (NO).....	147
<i>Toprol-XL 190</i> (AP).....	117	<i>Tritace 1.25 mg</i> (AV).....	124
<i>Toprol-XL Titration Pack</i> (AP).....	117	<i>Tritace 2.5 mg</i> (AV).....	124
TOREMIFENE CITRATE.....	191	<i>Tritace 5 mg</i> (AV).....	124
<i>Tracleer</i> (AT)		<i>Tritace 10 mg</i> (AV)	
.Section 100.....	369	.Cardiovascular system.....	124, 125
TRAMADOL HYDROCHLORIDE		.Repatriation Schedule.....	477
.Dental.....	353	<i>Tritace Titration Pack</i> (AV).....	125
.Doctor's Bag Supplies.....	72	<i>Trizivir</i> (GK)	
.Nervous system.....	257	.Section 100.....	359
<i>Tramahexal SR</i> (HX)		TROPISETRON HYDROCHLORIDE.....	85
.Dental.....	353, 354	<i>TrueSense</i> (MS).....	310
.Nervous system.....	258	<i>TrueTrack</i> (DR).....	309

GENERIC /PROPRIETARY INDEX

<i>Trusopt (MK)</i>	303
<i>Tryptanol (MK)</i>	275
<i>Tubegauz 0501633 (SS)</i>	
.Repatriation Schedule	503
<i>Tubegauz 0501658 (SS)</i>	
.Repatriation Schedule	503
<i>Tubifast 2434 (SS)</i>	
.Repatriation Schedule	503
<i>Tubifast 2436 (SS)</i>	
.Repatriation Schedule	503
<i>Tubifast 2438 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip 1479 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip 1480 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip 1481 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip 1482 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip 1483 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip 1484 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip 1486 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip B 1520 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip C 1545 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip D 1546 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip E 1547 (SS)</i>	
.Repatriation Schedule	503
<i>Tubigrip F 1548 (SS)</i>	
.Repatriation Schedule	503
<i>Tylenol (JT)</i>	
.Dental.....	354
.Nervous system	259
<i>TYR Express (VF)</i>	316
<i>TYR gel (VF)</i>	316
U	
<i>Ulcye (AF)</i>	84
<i>Ural Sachets (SI)</i>	
.Repatriation Schedule	486
UREA	
.Repatriation Schedule	479
<i>Urederm (HA)</i>	
.Repatriation Schedule	479
<i>Uremide (AF)</i>	114
<i>Urex (FM)</i>	114
<i>Urex-Forte (FM)</i>	114
<i>Urex-M (FM)</i>	114
<i>Uro-Carb (HA)</i>	287
<i>Uromitexan (BX)</i>	309
URSODEOXYCHOLIC ACID	86
<i>Ursofalk (OA)</i>	86

V

<i>Vagifem (NO)</i>	143
VALACICLOVIR HYDROCHLORIDE	
.Anti-infectives for systemic use	179
.Section 100	422
<i>Valcyte (RO)</i>	
.Section 100	423
VALGANCICLOVIR HYDROCHLORIDE	
.Section 100	423
<i>Valium (RO)</i>	
.Dental	355
.Nervous system	272, 273
.Palliative Care	330
<i>Valpam 2 (AW)</i>	
.Dental	355
.Nervous system	272
.Palliative Care	330
<i>Valpam 5 (AW)</i>	
.Dental	355
.Nervous system	273
.Palliative Care	330
<i>Valpro 200 (AF)</i>	263
<i>Valpro 500 (AF)</i>	263
<i>Valtrex (GK)</i>	
.Anti-infectives for systemic use	179
.Section 100	422
<i>Vancocin (LY)</i>	
.Alimentary tract and metabolism	90
.Anti-infectives for systemic use	173, 174
.Dental	345
VANCOMYCIN	
.Alimentary tract and metabolism	90
.Anti-infectives for systemic use	173
.Dental	345
<i>Vasocardol (HP)</i>	120
<i>Vasocardol CD (HP)</i>	121
<i>Vastin (AP)</i>	129
<i>Vaxigrip (AX)</i>	180
<i>Velbe (AS)</i>	183
VENLAFAXINE HYDROCHLORIDE	280
<i>Venofer (BX)</i>	106
<i>Ventavis (SC)</i>	
.Section 100	389
<i>Ventolin (GK)</i>	298
<i>Ventolin CFC-free (GK)</i>	
.Doctor's Bag Supplies	72
.Respiratory system	291
<i>Ventolin Nebules (GK)</i>	
.Doctor's Bag Supplies	72, 73
.Respiratory system	292
<i>Ventolin Rotacaps (GK)</i>	291
<i>Vepesid (BQ)</i>	184
<i>Veracaps SR (SI)</i>	120
<i>Verahexal (HX)</i>	120
VERAPAMIL HYDROCHLORIDE	
.Cardiovascular system	120
.Doctor's Bag Supplies	73

GENERIC /PROPRIETARY INDEX

<i>Vermox (JC)</i>		X	
.Repatriation Schedule.....	495	<i>Xalacom (PU)</i>	
<i>Viagra (PF)</i>		.Repatriation Schedule	497
.Repatriation Schedule.....	485	<i>Xalatan (PU)</i>	304
<i>Vibramycin (PF)</i>		<i>Xanax (PH)</i>	272
.Antiinfectives for systemic use.....	159, 160	<i>Xanax Tri-Score (PH)</i>	272
.Dental.....	336	<i>Xeloda (RO)</i>	183
<i>Vibra-Tabs (PF)</i>	159	<i>Xenical (RO)</i>	
<i>Videx EC (BQ)</i>		.Repatriation Schedule	474
.Section 100	372, 373	<i>Xergic (AF)</i>	
VIGABATRIN.....	264	.Repatriation Schedule	497
VINBLASTINE SULFATE	183	<i>Xigris (LY)</i>	105
VINCRISTINE SULFATE	183	<i>XMET Analog (SB)</i>	315
VINORELBINE TARTRATE.....	184	<i>XMET Maxamaid (SB)</i>	315
<i>Viracept (RO)</i>		<i>XMET Maxamum (SB)</i>	315
.Section 100	414	<i>XMTVI Analog (SB)</i>	316
<i>Viramune (BY)</i>		<i>XMTVI Asadon (SB)</i>	315
.Section 100	414	<i>XMTVI Maxamaid (SB)</i>	316
<i>Viread (GI)</i>		<i>XMTVI Maxamum (SB)</i>	316
.Section 100	422	<i>XP Analog (SB)</i>	316
<i>Viscopaste 4948 (SN)</i>		<i>XP Analog LCP (SB)</i>	315
.Repatriation Schedule.....	504	<i>XP Maxamaid (SB)</i>	316
<i>Viscotears (NV)</i>	305	<i>XP Maxamum (SB)</i>	316
<i>Viscotears Liquid Gel (NV)</i>	305	<i>XPhen, Tyr Analog (SB)</i>	316
<i>Visken 5 (NV)</i>	116	<i>XPhen, Tyr Maxamaid (SB)</i>	316
<i>Visken 15 (NV)</i>	116	<i>XPhen, Tyr Maxamum (SB)</i>	316
<i>Vistide (PU)</i>		<i>XPTM Tyrosidon (SB)</i>	315
.Section 100	370	<i>Xydep 50 (AW)</i>	278
<i>Vistil (AE)</i>	307	<i>Xydep 100 (AW)</i>	278
<i>Vistil Forte (AE)</i>	307	<i>Xylocaine Viscous (AP)</i>	
VITAMIN B GROUP COMPLEX		.Repatriation Schedule	480
.Repatriation Schedule.....	475	<i>Xylocard 500 (AP)</i>	108
<i>Vitelle Vitamin C (FH)</i>			
.Repatriation Schedule.....	475	Z	
<i>Vitrasert (BU)</i>		<i>Zabel (AF)</i>	
.Section 100	380	.Dermatologicals	134
<i>Voltaren 25 (NV)</i>		.Repatriation Schedule	478
.Dental.....	346	<i>Zactin (AF)</i>	277
.Musculo-skeletal system	238	ZALCITABINE	
.Palliative Care	324, 325	.Section 100	423
<i>Voltaren 50 (NV)</i>		<i>Zamhexal 0.25mg (HX)</i>	272
.Dental.....	346	<i>Zamhexal 0.5mg (HX)</i>	272
.Musculo-skeletal system	238	<i>Zamhexal 1.0mg (HX)</i>	272
.Palliative Care	325	<i>Zamhexal 2mg (HX)</i>	272
<i>Voltaren 100 (NV)</i>		<i>Zanidip (SM)</i>	
.Dental.....	346	.Cardiovascular system	119
.Musculo-skeletal system	238	.Repatriation Schedule	476
.Palliative Care	325	<i>Zantac (GK)</i>	
<i>Voltaren Ophtha (NV)</i>	301	.Alimentary tract and metabolism.....	79, 80
<i>Voxam (HX)</i>	277	.Repatriation Schedule	473
W		<i>Zantac Syrup (GK)</i>	
WARFARIN SODIUM.....	101	.Alimentary tract and metabolism.....	80
<i>Waxsol (NE)</i>		.Repatriation Schedule	473
.Repatriation Schedule.....	498	<i>Zarontin (PF)</i>	262
<i>Wellvone (GK)</i>	289	<i>Zavedos (PH)</i>	186
WHEY PROTEIN FORMULA supplemented with		<i>Zavedos Solution (PH)</i>	186
AMINO ACIDS, VITAMINS and MINERALS, and		<i>Zeffix (GK)</i>	
low in PROTEIN, PHOSPHATE, POTASSIUM and		.Section 100	411
LACTOSE	318	<i>Zentel (GK)</i>	290
WOOL ALCOHOLS		<i>Zerit (BQ)</i>	
.Repatriation Schedule.....	479	.Section 100	422

GENERIC /PROPRIETARY INDEX

<i>Zestril (AP)</i>	123, 124	ZOLEDRONIC ACID	
<i>Ziagen (GK)</i>		.Section 100	423
.Section 100	359	ZOLMITRIPTAN	261
ZIDOVUDINE		<i>Zoloft (PF)</i>	278
.Section 100	423	<i>Zometa (NV)</i>	
<i>Zimstat (AF)</i>	129, 130, 131	.Section 100	423
ZINC OXIDE		<i>Zomig (AP)</i>	261
.Repatriation Schedule	476	ZOPICLONE	
ZINC OXIDE with STARCH and CHLORPHENESIN		.Repatriation Schedule	494
.Repatriation Schedule	483	<i>Zoton (WY)</i>	81
ZINC PYRITHIONE		<i>Zovirax (GK)</i>	300
.Repatriation Schedule	482	<i>Zovirax 200 mg (GK)</i>	177
ZINC SULFATE with PHENYLEPHRINE		<i>Zovirax 800 mg (GK)</i>	178
HYDROCHLORIDE		<i>Z.S.C. (SI)</i>	
.Repatriation Schedule	497	.Repatriation Schedule	483
<i>Zincaband 3604 (SS)</i>		ZUCLOPENTHIXOL DECANOATE.....	269
.Repatriation Schedule	504	<i>Zumenon (SM)</i>	143
<i>Zincfrin (AQ)</i>		<i>Zyban (GK)</i>	287
.Repatriation Schedule	497	<i>Zyclir 200 (AW)</i>	177
<i>Zinnat (GK)</i>		<i>Zyclir 800 (AW)</i>	178
.Antifectives for systemic use	168	<i>Zydol (AW)</i>	
.Dental.....	343	.Dental	353
<i>ZipZoc 66051550 (SN)</i>		.Nervous system	257, 258
.Repatriation Schedule	504	<i>Zydol SR 100 (AW)</i>	
<i>Zithromax (PF)</i>		.Dental	353
.Antifectives for systemic use	170	.Nervous system	258
.Repatriation Schedule	487	<i>Zydol SR 150 (AW)</i>	
.Section 100	360	.Dental	353
.Sensory organs.....	300	.Nervous system	258
<i>Zocor (MK)</i>	129, 130, 131	<i>Zydol SR 200 (AW)</i>	
<i>Zofran (GK)</i>	85	.Dental	354
<i>Zofran syrup 50 mL (GK)</i>	85	.Nervous system	258
<i>Zofran Zydys (GK)</i>	85	<i>Zyloprim (SI)</i>	244
<i>Zoladex 10.8 Implant (AP)</i>	190	<i>Zyprexa (LY)</i>	270
<i>Zoladex Implant (AP)</i>	190	<i>Zyprexa Zydys (LY)</i>	270
		<i>Zyrtec (PC)</i>	
		.Repatriation Schedule	496

THERAPEUTIC GROUP PREMIUM POLICY

PHARMACEUTICAL BENEFIT ITEMS WHICH HAVE A THERAPEUTIC GROUP PREMIUM WITH EFFECT FROM 1 DECEMBER 2005

The Schedule of Pharmaceutical Benefits shows differences in price in some therapeutic groups where alternative drugs may have a therapeutic group premium.

The Therapeutic Group Premium Policy applies within narrowly defined therapeutic sub-groups where the drugs concerned are of similar safety and health outcomes.

The Australian Government, through the PBS, subsidises up to the price of the lowest priced drug in the group. This means that consumers may have to pay for more expensive drugs (those with a therapeutic group premium). This extra amount does not count towards their PBS safety net threshold.

Therapeutic group premiums apply where a prescriber has prescribed a drug within a therapeutic group that attracts a therapeutic group premium and has not sought an exemption from Medicare Australia on clinical grounds.

The exemption provisions are:

- adverse effects occurring with all of the base-priced drugs; or
- drug interactions occurring with all of the base-priced drugs; or
- drug interactions expected to occur with all of the base-priced drugs; or
- transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.

The premiums are not a Government charge but reflect the fact that the supplier(s) of the drug charge a price higher than the Government is willing to subsidise.

Under the Therapeutic Group Premium Policy drug substitution by pharmacists is not permitted.

For ease of prescribing and dispensing, and in the interests of your patients, the following list shows those PBS drugs that attract a therapeutic group premium.

Premium Priced Brand	Form and Strength	Max. Qty	Therapeutic Group Premium \$

H₂-RECEPTOR ANTAGONISTS

<i>Tagamet 800 Express</i>	Effervescent tablet 800 mg (as hydrochloride)	30	7.01
<i>Zantac</i>	Effervescent tablet 150 mg (base)	60	2.14
<i>Zantac Syrup</i>	Syrup 150 mg (base) per 10 mL, 300 mL	2	2.14

The base-priced drugs in this therapeutic group are cimetidine (except cimetidine effervescent tablet 800 mg (as hydrochloride)), famotidine, nizatidine, and ranitidine hydrochloride (except ranitidine hydrochloride effervescent tablet 150 mg (base) and syrup 150 mg (base) per 10 mL, 300 mL).

DIHYDROPYRIDINE-DERIVATIVE CALCIUM CHANNEL BLOCKERS

<i>Adalat Oros 20mg</i>	Tablet 20 mg (controlled release)	30	2.20
<i>Norvasc</i>	Tablet 5 mg (base)	30	4.62
<i>Norvasc</i>	Tablet 10 mg (base)	30	6.70
<i>Zanidip</i>	Tablet 10 mg	30	2.15
<i>Zanidip</i>	Tablet 20 mg	30	3.60

The base-priced drugs in this therapeutic group are felodipine and nifedipine (except nifedipine controlled release tablet 20 mg).

ACE INHIBITORS, PLAIN

<i>Tritace 10 mg</i>	Capsule 10 mg	30	1.35

The base-priced drugs in this therapeutic group are captopril, enalapril maleate, fosinopril sodium, lisinopril, perindopril erbumine, quinapril hydrochloride, ramipril (except ramipril capsule 10 mg) andtrandolapril.

BRAND PREMIUM POLICY

BRANDS OF PHARMACEUTICAL BENEFIT ITEMS WHICH HAVE A BRAND PREMIUM AND THAT MAY BE SUBSTITUTED WITH EFFECT FROM 1 DECEMBER 2005

The Schedule of Pharmaceutical Benefits shows differences in price between some alternative brands of the same drug product.

Manufacturers can develop generic equivalents and apply to have them listed on the PBS. In doing this, manufacturers need to ensure that they comply with the relevant legislation applicable to patents. These brands are clinically equivalent and must undergo the same strict quality controls. Although these brands are designed to act on the body in exactly the same way, they are usually cheaper than the originator brands.

The Australian Government, through the PBS, subsidises up to the price of the lowest priced brand (except in those instances where the lowest priced brand has, as part of its price, a therapeutic group premium). This means that consumers may have to pay extra for more expensive brands (those with a brand premium). This extra amount does not count towards their PBS safety net threshold.

Brand substitution by pharmacists without reference to the prescriber is permitted for PBS prescriptions where:

- the patient agrees to the substitution;
- the brands are identified in the Schedule of Pharmaceutical Benefits as being interchangeable;
- the prescriber has not indicated on the prescription form that substitution is not to occur; and
- substitution is permitted under the relevant State or Territory legislation.

Prescription forms supplied by Medicare Australia contain a box to be ticked where brand substitution is not to take place.

Prescribers not using these prescription forms should endorse the prescription if brand substitution is not permitted. Where a stamp is used for this purpose, the prescriber will be required to initial the stamped statement.

For ease of prescribing and dispensing, and in the interests of your patients, the following list shows those PBS drugs that attract a brand premium and that can be substituted where permitted. They are listed alphabetically, by brand name, with the brand premium and benchmark brand(s) cited in the last column.

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Abbecillin-V</i>	Paediatric oral suspension 125 mg per 5 mL, 100 mL	2	1.44	<i>Cilicaine V</i>
	Oral suspension 250 mg per 5 mL, 100 mL	2	1.46	<i>Cilicaine V</i>
<i>Adalat Oros 30</i>	Tablet 30 mg (controlled release)	30	3.10	<i>Addos XR 30; Adefin XL 30</i>
<i>Adalat Oros 60</i>	Tablet 60 mg (controlled release)	30	3.70	<i>Addos XR 60; Adefin XL 60</i>
<i>Adalat 10</i>	Tablet 10 mg	60	1.78	<i>Adefin 10; Nypine 10</i>
<i>Adalat 20</i>	Tablet 20 mg	60	3.31	<i>Adefin 20; Chem mart Nifedipine; GenRx Nifedipine; Nifehexal; Nyefax 20 mg; Nypine 20; Terry White Chemists Nifedipine</i>
<i>Airomir</i>	Oral pressurised inhalation 100 micrograms (base) per dose (200 doses), CFC-free formulation	2	0.62	<i>Asmol CFC-free; Epag</i>
<i>Aldactone</i>	Tablet 25 mg	100	1.89	<i>Spiractin 25</i>
	Tablet 100 mg	100	2.62	<i>Spiractin 100</i>
<i>Aldomet</i>	Tablet 250 mg	100	2.95	<i>Hydopa</i>
<i>Alphagan</i>	Eye drops 2 mg per mL (0.2%), 5 mL	1	1.50	<i>Enidin</i>
<i>Amaryl</i>	Tablet 1 mg	30	2.50	<i>Dimirel</i>
	Tablet 2 mg	30	2.50	<i>Dimirel</i>
	Tablet 3 mg	30	2.50	<i>Dimirel</i>
	Tablet 4 mg	30	2.50	<i>Dimirel</i>
<i>Amoxil</i>	Capsule 250 mg	20	1.00	<i>Alphamox 250; Amohexal; Amoxycillin-DP; Bgramin; Chem mart Amoxycillin; Cilamox; GenRx Amoxycillin; Moxacin; Terry White Chemists Amoxycillin</i>
	Capsule 500 mg	20	1.00	<i>Alphamox 500; Amohexal; Amoxycillin-DP; Bgramin; Chem mart Amoxycillin; Cilamox; GenRx Amoxycillin; Moxacin; Terry White Chemists Amoxycillin</i>
	Powder for syrup 125 mg per 5 mL, 100 mL	1	1.00	<i>Alphamox 125; Amohexal; Bgramin; Chem mart Amoxycillin; Cilamox; GenRx Amoxycillin; Moxacin; Terry White Chemists Amoxycillin</i>
<i>Amoxil Duo</i>	Tablet 1 g	14	1.49	<i>Amoxycillin Sandoz</i>
<i>Amoxil Forte</i>	Powder for syrup 250 mg per 5 mL, 100 mL	1	1.01	<i>Alphamox 250; Amohexal; Bgramin; Chem mart Amoxycillin; Cilamox; GenRx Amoxycillin; Moxacin 250; Terry White Chemists Amoxycillin</i>
<i>Amprace 5</i>	Tablet 5 mg	30	2.45	<i>Alphapril; Auspril; Chem mart Enalapril; Enahexal; Enalabell; Enalapril-DP 5mg; GenRx Enalapril; Terry White Chemists Enalapril</i>
<i>Amprace 10</i>	Tablet 10 mg	30	2.45	<i>Alphapril; Auspril; Chem mart Enalapril; Enahexal; Enalabell; Enalapril-DP 10mg; GenRx Enalapril; Terry White Chemists Enalapril</i>
<i>Amprace 20</i>	Tablet 20 mg	30	2.46	<i>Alphapril; Auspril; Chem mart Enalapril; Enahexal; Enalabell; Enalapril-DP 20mg; GenRx Enalapril; Terry White Chemists Enalapril</i>
<i>Anafranil 25</i>	Tablet 25 mg	50	2.94	<i>Chem mart Clomipramine; GenRx Clomipramine; Placil; Terry White Chemists Clomipramine</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Anaprox 550</i>	Tablet 550 mg	50	3.00	<i>Crysanal</i>
<i>Androcur</i>	Tablet 50 mg	20	1.50	<i>Cyprohexal; Cyprone; Cyprostat; GenRx Cyproterone Acetate; Procur</i>
	Tablet 50 mg	100	3.28	<i>Cyprohexal; Cyprone; Cyprostat; GenRx Cyproterone Acetate; Procur</i>
<i>Androcur-100</i>	Tablet 100 mg	50	1.00	<i>Cyprohexal; Cyprostat-100; GenRx Cyproterone Acetate; Procur 100</i>
<i>Anginine Stabilised</i>	Tablets 600 micrograms, 100	1	1.30	<i>Lycinat</i>
<i>Arava</i>	Tablet 10 mg	30	1.22	<i>Arabloc</i>
	Tablet 20 mg	30	1.22	<i>Arabloc</i>
<i>Aristocort 0.02%</i>	Cream 200 micrograms per g (0.02%), 100 g	2	2.60	<i>Tricortone</i>
	Ointment 200 micrograms per g (0.02%), 100 g	2	2.60	<i>Tricortone</i>
<i>Aropax</i>	Tablet 20 mg (base)	30	1.10	<i>Chem mart Paroxetine; GenRx Paroxetine; Oxetine; Paxtine; Terry White Chemists Paroxetine</i>
<i>Astrix</i>	Tablet 100 mg	112	1.36	<i>DBL Aspirin 100 mg</i>
<i>Atrovent</i>	Nebuliser solution single dose units 250 micrograms (anhydrous) in 1 mL, 30	2	0.94	<i>Aeron 250; Apoven 250; Chem mart Ipratropium; DBL Ipratropium; GenRx Ipratropium; Ipratrin; Ipravent; Terry White Chemists Ipratropium</i>
<i>Atrovent Adult</i>	Nebuliser solution single dose units 500 micrograms (anhydrous) in 1 mL, 30	2	0.94	<i>Aeron 500; Apoven 500; Chem mart Ipratropium; DBL Ipratropium; GenRx Ipratropium; Ipratrin Adult; Ipravent; Terry White Chemists Ipratropium</i>
<i>Augmentin</i>	Powder for syrup 125 mg-31.25 mg per 5 mL, 75 mL	1	0.96	<i>Clamohexal 125mg/31.25mg/5mL; Clamoxyl; Clavulin</i>
<i>Augmentin Duo</i>	Tablet 500 mg-125 mg	10	0.99	<i>Clamohexal Duo 500mg/125mg; Clamoxyl Duo; Clavulin Duo; Curam 500/125; Muric 500/125</i>
<i>Augmentin Duo forte</i>	Tablet 875 mg-125 mg	10	1.30	<i>Chem mart Amoxycillin and Clavulanic Acid; Clamohexal Duo Forte 875mg/125mg; Clamoxyl Duo forte; Clavulin Duo Forte; Curam 875/125; GenRx Amoxycillin and Clavulanic Acid; Muric 875/125; Terry White Chemists Amoxycillin and Clavulanic Acid</i>
<i>Augmentin Duo 400</i>	Powder for syrup 400 mg-57 mg per 5 mL, 60 mL	1	0.98	<i>Clamohexal Duo 400mg/57mg/5mL; Clamoxyl Duo 400; Clavulin Duo 400</i>
<i>Aurorix</i>	Tablet 150 mg	60	0.95	<i>Arima; Chem mart Moclobemide; Clobemix; GenRx Moclobemide; Mohexal; Terry White Chemists Moclobemide</i>
<i>Aurorix 300 mg</i>	Tablet 300 mg	60	1.93	<i>Arima 300; Chem mart Moclobemide; Clobemix; GenRx Moclobemide; Maosig; Mohexal; Terry White Chemists Moclobemide</i>
<i>Avanza</i>	Tablet 30 mg	30	1.25	<i>Axit 30; Mirtazon</i>
<i>Azopt</i>	Eye drops 10 mg per mL (1%), 5 mL	1	1.00	<i>BrinzoQuin</i>
<i>Betaloc</i>	Tablet 50 mg	100	3.26	<i>Chem mart Metoprolol; GenRx Metoprolol; Metohexal; Metolol; Metrol 50; Minax 50; Terry White Chemists Metoprolol</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Betaloc</i>	Tablet 100 mg	60	3.25	<i>Chem mart Metoprolol; GenRx Metoprolol; Metohexal; Metolol; Metrol 100; Minax 100; Terry White Chemists Metoprolol</i>
<i>Betnovate 1/5</i>	Cream 200 micrograms (base) per g (0.02%), 100 g	2	2.60	<i>Cortival 1/5</i>
<i>Betnovate 1/2</i>	Cream 500 micrograms (base) per g (0.05%), 15 g	1	1.29	<i>Cortival 1/2</i>
	Ointment 500 micrograms (base) per g (0.05%), 15 g	1	1.29	<i>Cortival 1/2</i>
<i>Betoptic</i>	Eye drops, solution, 5 mg (base) per mL (0.5%), 5 mL	1	2.01	<i>BetoQuin</i>
<i>Blenoxane</i>	Powder for injection 15,000 i.u.	10	79.48	<i>Blenamax</i>
<i>Brevinor</i>	Tablets 500 micrograms-35 micrograms, 21	4	7.96	<i>Norimin 28 Day</i>
	Pack containing 21 tablets 500 micrograms-35 micrograms and 7 inert tablets	4	7.96	<i>Norimin 28 Day</i>
<i>Brevinor-1</i>	Tablets 1 mg-35 micrograms, 21	4	7.96	<i>Norimin-1 28 Day</i>
	Pack containing 21 tablets 1 mg-35 micrograms and 7 inert tablets	4	7.96	<i>Norimin-1 28 Day</i>
<i>Capoten</i>	Tablet 12.5 mg	90	2.46	<i>Acenorm 12.5 mg; Captopril; Chem mart Captopril; GenRx Captopril; Terry White Chemists Captopril; Topace; DP brand</i>
	Tablet 25 mg	90	2.46	<i>Acenorm 25 mg; Captopril; Chem mart Captopril; GenRx Captopril; Terry White Chemists Captopril; Topace; DP brand</i>
	Tablet 50 mg	90	2.45	<i>Acenorm 50 mg; Captopril; Chem mart Captopril; GenRx Captopril; Terry White Chemists Captopril; Topace; DP brand</i>
<i>Carafate</i>	Tablet equivalent to 1 g anhydrous sucralfate	120	2.21	<i>Ulcyte</i>
<i>Cardizem</i>	Tablet 60 mg	90	2.64	<i>Chem mart Diltiazem; Coras; Diltahexal; Dilzem 60 mg; GenRx Diltiazem; Terry White Chemists Diltiazem; Vasocardol</i>
<i>Cardizem CD</i>	Capsule 180 mg (controlled delivery)	30	2.64	<i>Chem mart Diltiazem CD; Dilzem CD; GenRx Diltiazem CD; Terry White Chemists Diltiazem CD; Vasocardol CD</i>
	Capsule 240 mg (controlled delivery)	30	2.64	<i>Chem mart Diltiazem CD; Dilzem CD; GenRx Diltiazem CD; Terry White Chemists Diltiazem CD; Vasocardol CD</i>
	Capsule 360 mg (controlled delivery)	30	3.21	<i>Vasocardol CD</i>
<i>Ceclor</i>	Powder for oral suspension 125 mg per 5 mL, 100 mL	1	1.53	<i>Aclor 125; Chem mart Cefaclor; GenRx Cefaclor; Keflor; Terry White Chemists Cefaclor</i>
	Powder for oral suspension 250 mg per 5 mL, 75 mL	1	1.55	<i>Aclor 250; Chem mart Cefaclor; GenRx Cefaclor; Keflor; Terry White Chemists Cefaclor</i>
<i>Ceclor CD</i>	Tablet 375 mg (sustained release)	10	1.54	<i>Cefkor CD; Chem mart Cefaclor CD; Douglas Cefaclor-CD; GenRx Cefaclor CD; Karlor CD; Keflor CD; Terry White Chemists Cefaclor CD</i>
<i>Chlorvescent</i>	Effervescent tablet 14 mmol K ⁺ and 8 mmol Cl ⁻	60	2.63	<i>K-Sol</i>
<i>Cicloral</i>	Capsule 25 mg	60	2.38	<i>Cysporin</i>
	Capsule 50 mg	60	2.38	<i>Cysporin</i>
	Capsule 100 mg	60	2.40	<i>Cysporin</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Ciloxan</i>	Eye drops 3 mg per mL (0.3%), 5 mL	2	2.48	<i>CiloQuin</i>
<i>Cipramil</i>	Tablet 20 mg (base)	28	3.20	<i>Celapram</i> ; <i>Chem mart Citalopram</i> ; <i>Ciazil</i> ; <i>GenRx Citalopram</i> ; <i>Talam</i> ; <i>Talohexal</i> ; <i>Terry White Chemists Citalopram</i>
<i>Ciproxin 250</i>	Tablet 250 mg	14	1.90	<i>C-Flox 250</i> ; <i>Ciprofloxacin-BC</i> ; <i>Ciprol 250</i> ; <i>GenRx Ciprofloxacin</i> ; <i>Profloxin</i>
<i>Ciproxin 500</i>	Tablet 500 mg	14	1.90	<i>C-Flox 500</i> ; <i>Ciprofloxacin-BC</i> ; <i>Ciprol 500</i> ; <i>GenRx Ciprofloxacin</i> ; <i>Profloxin</i> ; <i>Proquin</i>
<i>Ciproxin 750</i>	Tablet 750 mg	14	1.90	<i>C-Flox 750</i> ; <i>Ciprol 750</i> ; <i>GenRx Ciprofloxacin</i> ; <i>Profloxin</i> ; <i>Proquin</i>
<i>Clomid</i>	Tablet 50 mg	10	3.74	<i>Clomhexal</i> ; <i>Fermil</i> ; <i>GenRx Clomiphene</i>
<i>Colgout</i>	Tablet 500 micrograms	100	0.93	<i>Lengout</i>
<i>Cordarone X 100</i>	Tablet 100 mg	30	1.00	<i>Aratac 100</i> ; <i>Cardinorm</i> ; <i>GenRx Amiodarone</i> ; <i>Rithmik 100</i>
<i>Cordarone X 200</i>	Tablet 200 mg	30	1.00	<i>Aratac 200</i> ; <i>Cardinorm</i> ; <i>Chem mart Amiodarone</i> ; <i>GenRx Amiodarone</i> ; <i>Rithmik 200</i> ; <i>Terry White Chemists Amiodarone</i>
<i>Dalacin C</i>	Capsule 150 mg	25	1.47	<i>Cleocin</i>
<i>Daonil</i>	Tablet 5 mg	100	1.25	<i>Glimel</i>
<i>Depo-Medrol</i>	Injection 40 mg in 1 mL	5	0.75	<i>Depo-Nisolone</i>
<i>Depo-Provera</i>	Injection 150 mg in 1 mL	1	3.36	<i>Depo-Ralovera</i>
<i>Diabex</i>	Tablet 500 mg	100	1.80	<i>Chem mart Metformin</i> ; <i>Diaformin</i> ; <i>Formet 500</i> ; <i>GenRx Metformin</i> ; <i>Glucohexal</i> ; <i>Glucomet 500 mg</i> ; <i>Terry White Chemists Metformin</i>
<i>Diabex 850</i>	Tablet 850 mg	60	1.80	<i>Chem mart Metformin</i> ; <i>Diaformin 850</i> ; <i>Formet 850</i> ; <i>GenRx Metformin</i> ; <i>Glucohexal</i> ; <i>Glucomet 850 mg</i> ; <i>Terry White Chemists Metformin</i>
<i>Diabex 1000</i>	Tablet 1 g	90	1.81	<i>Diaformin 1000</i> ; <i>Formet 1000</i>
<i>Diamicron</i>	Tablet 80 mg	100	1.35	<i>Chem mart Gliclazide</i> ; <i>GenRx Gliclazide</i> ; <i>Glyade</i> ; <i>Mellihexal</i> ; <i>Nidem</i> ; <i>Terry White Chemists Gliclazide</i>
<i>Doryx</i>	Capsule 100 mg	7	1.52	<i>DBL Doxycycline</i>
	Capsule 50 mg	25	1.71	<i>DBL Doxycycline</i>
	Capsule 100 mg	28	6.08	<i>DBL Doxycycline</i>
	Capsule 100 mg	21	2.75	<i>DBL Doxycycline</i>
<i>Duphalac</i>	Mixture 3.34 g per 5 mL, 500 mL	1	2.20	<i>Actilax</i> ; <i>Genlac</i> ; <i>GenRx Lactulose</i> ; <i>Lac-Dol</i> ; <i>Lactocur</i>
<i>Duratears</i>	Compound eye ointment 3.5 g	2	2.00	<i>Poly Visc</i>
<i>E.E.S. Granules</i>	Powder for oral liquid 400 mg (base) per 5 mL, 100 mL	1	1.75	<i>E-Mycin 400</i>
<i>E.E.S. 200</i>	Powder for oral liquid 200 mg (base) per 5 mL, 100 mL	1	2.19	<i>E-Mycin 200</i>
<i>E.E.S. 400 Filmtab</i>	Tablet 400 mg (base)	25	2.89	<i>E-Mycin</i>
<i>Epilim EC</i>	Tablet 200 mg (enteric coated)	200	1.32	<i>Valpro 200</i>
	Tablet 500 mg (enteric coated)	200	1.64	<i>Valpro 500</i>
<i>Eryc</i>	Capsule 250 mg	25	1.77	<i>DBL Erythromycin</i>
<i>Fasigyn</i>	Tablet 500 mg	4	2.50	<i>Simplotan</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Feldene</i>	Capsule 10 mg	50	2.75	<i>Chem mart Piroxicam;</i> <i>GenRx Piroxicam; Mobilis 10;</i> <i>Terry White Chemists Piroxicam</i>
	Capsule 20 mg	25	2.73	<i>Chem mart Piroxicam;</i> <i>GenRx Piroxicam; Mobilis 20;</i> <i>Terry White Chemists Piroxicam</i>
<i>Feldene-D</i>	Dispersible tablet 10 mg	50	2.75	<i>GenRx Piroxicam Dispersible;</i> <i>Mobilis D-10; Pirohexal-D</i>
	Dispersible tablet 20 mg	25	2.73	<i>Chem mart Piroxicam Dispersible;</i> <i>GenRx Piroxicam Dispersible;</i> <i>Mobilis D-20; Pirohexal-D;</i> <i>Terry White</i> <i>Chemists Piroxicam Dispersible</i>
<i>Ferrum H</i>	Injection 100 mg (iron) in 2 mL	5	1.22	<i>Ferrosig</i>
<i>Flagyl</i>	Tablet 200 mg	21	1.98	<i>Metrogyl 200; Metronide 200</i>
	Tablet 400 mg	21	2.07	<i>Metrogyl 400; Metronide 400</i>
<i>Floxapen</i>	Capsule 250 mg	24	0.46	<i>Flopen; Floxsig; Staphylex 250</i>
	Capsule 500 mg	24	0.58	<i>Flopen; Floxsig; Staphylex 500</i>
	Powder for syrup 125 mg per 5 mL, 100 mL	1	0.06	<i>Flopen</i>
	Powder for syrup 250 mg per 5 mL, 100 mL	1	0.10	<i>Flopen</i>
<i>Fosamax Once Weekly</i>	Tablet equivalent to 70 mg alendronic acid	4	0.24	<i>Alendro Once Weekly</i>
<i>Genteal</i>	Eye drops 3 mg per mL (0.3%), 15 mL (contains sodium perborate as preservative)	1	1.86	<i>In a Wink Moisturising</i>
<i>Genteal gel</i>	Ocular lubricating gel 3 mg-2 mg per g (0.3%-0.2%), 10 g	1	1.86	<i>HPMC PAA</i>
<i>Glucophage</i>	Tablet 500 mg	100	1.41	<i>Chem mart Metformin; Diaformin;</i> <i>Formet 500; GenRx Metformin;</i> <i>Glucohexal; Glucomet 500 mg;</i> <i>Terry White Chemists Metformin</i>
	Tablet 850 mg	60	1.41	<i>Chem mart Metformin;</i> <i>Diaformin 850; Formet 850;</i> <i>GenRx Metformin; Glucohexal;</i> <i>Glucomet 850 mg;</i> <i>Terry White Chemists Metformin</i>
<i>Imdur Durule</i>	Tablet 60 mg (sustained release)	30	2.85	<i>Arsorb 60;</i> <i>Chem mart Isosorbide Mononitrate;</i> <i>Duride; GenRx Isosorbide</i> <i>Mononitrate; Imtrate 60 mg;</i> <i>Isomonit; Monodur 60 mg;</i> <i>Terry White Chemists Isosorbide</i> <i>Mononitrate</i>
<i>Imdur 120 mg</i>	Tablet 120 mg (sustained release)	30	3.00	<i>Monodur 120 mg</i>
<i>Imodium</i>	Capsule 2 mg	12	0.92	<i>Gastro-Stop Loperamide</i>
<i>Imuran</i>	Tablet 25 mg	100	1.57	<i>Azahexal</i>
	Tablet 50 mg	100	1.50	<i>Azahexal; Azamun; Azapin;</i> <i>GenRx Azathioprine; Thioprine</i>
<i>Indocid</i>	Capsule 25 mg	100	3.06	<i>Arthrexin</i>
<i>Insensye</i>	Tablet 400 mg	14	1.73	<i>Chem mart Norfloxacin; GenRx</i> <i>Norfloxacin; Norflohexal; Nufloxib;</i> <i>Roxin; Terry White Chemists</i> <i>Norfloxacin; DP brand</i>
<i>Isoptin</i>	Tablet 40 mg	100	1.00	<i>Anpec 40; Verahexal</i>
	Tablet 80 mg	100	1.00	<i>Anpec 80; Verahexal</i>
	Tablet 120 mg	100	1.00	<i>Verahexal</i>
<i>Isoptin SR</i>	Tablet 240 mg (sustained release)	30	3.00	<i>Anpec SR; Cordilox SR</i>
<i>Isoptin 180 SR</i>	Tablet 180 mg (sustained release)	30	3.00	<i>Cordilox 180 SR</i>
<i>Isordil</i>	Tablet 10 mg	200	3.70	<i>Sorbidin</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Keflex</i>	Capsule 250 mg	20	1.76	<i>Chem mart Cephalixin; Cilex; GenRx Cephalixin; Ialex; Ibilex 250; Sporaheal; Terry White Chemists Cephalixin</i>
	Capsule 500 mg	20	1.87	<i>Chem mart Cephalixin; Cilex; GenRx Cephalixin; Ialex; Ibilex 500; Sporaheal; Terry White Chemists Cephalixin</i>
	Granules for syrup 125 mg per 5 mL, 100 mL	1	1.78	<i>Chem mart Cephalixin; Cilex; GenRx Cephalixin; Ialex; Ibilex 125; Terry White Chemists Cephalixin</i>
	Granules for syrup 250 mg per 5 mL, 100 mL	1	1.87	<i>Chem mart Cephalixin; Cilex; GenRx Cephalixin; Ialex; Ibilex 250; Terry White Chemists Cephalixin</i>
<i>Keflin Neutral</i>	Powder for injection 1 g	10	0.30	MX brand
<i>Kefzol</i>	Powder for injection 1 g	10	2.51	<i>Cefazolin Sandoz; MX brand</i>
<i>Kenacomb Otic</i>	Ear drops 1 mg-2.5 mg (base)- 250 micrograms-100,000 units per g (0.1%- 0.25%-0.025%-100,000 units per g), 7.5 mL	1	1.10	<i>Otocomb Otic</i>
	Ear ointment 1 mg-2.5 mg (base)- 250 micrograms-100,000 units per g (0.1%- 0.25%-0.025%-100,000 units per g), 5 g	1	1.10	<i>Otocomb Otic</i>
<i>Klacid</i>	Tablet 250 mg	14	2.00	<i>Chem mart Clarithromycin; Clarac; Clarihexas; GenRx Clarithromycin; Kalixocin; Terry White Chemists Clarithromycin</i>
<i>Lacri-Lube</i>	Pack containing 2 tubes compound eye ointment 3.5 g	1	2.01	<i>Ircal</i>
<i>Lamictal</i>	Tablet 5 mg	56	1.00	<i>Elmendos; Lamitrin; Lamogine; Seaze 5</i>
	Tablet 25 mg	56	1.00	<i>Elmendos; GenRx Lamotrigine; Lamitrin; Lamogine; Lamotrigine- DP; Seaze 25</i>
	Tablet 50 mg	56	0.99	<i>Elmendos; GenRx Lamotrigine; Lamitrin; Lamogine; Lamotrigine- DP; Seaze 50</i>
	Tablet 100 mg	56	1.00	<i>Elmendos; GenRx Lamotrigine; Lamitrin; Lamogine; Lamotrigine- DP; Seaze 100</i>
	Tablet 200 mg	56	1.00	<i>Elmendos; GenRx Lamotrigine; Lamitrin; Lamogine; Lamotrigine- DP; Seaze 200</i>
<i>Lamisil</i>	Tablet 250 mg (base)	42	2.00	<i>GenRx Terbinafine; Tamsil; Terbihexas; Terbinafine-DP; Zabel</i>
<i>Lanoxin</i>	Tablet 250 micrograms	100	1.30	<i>Sigmaxin</i>
<i>Lanoxin-PG</i>	Tablet 62.5 micrograms	200	1.30	<i>Sigmaxin-PG</i>
<i>Lasix</i>	Tablet 40 mg	100	1.11	<i>Chem mart Frusemide; Frusehexal 40 mg; Frusid; GenRx Frusemide; Terry White Chemists Frusemide; Uremide</i>
<i>Lasix-M</i>	Tablet 20 mg	100	1.12	<i>Chem mart Frusemide; Frusid; GenRx Frusemide; Terry White Chemists Frusemide</i>
<i>Ledertrexate</i>	Tablet 2.5 mg	30	0.08	<i>Methoblastin; MX brand</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Lioresal 10</i>	Tablet 10 mg	100	3.00	<i>Baclo</i> ; <i>Baclohexal</i> ; <i>Chem mart Baclofen</i> ; <i>Clofen 10</i> ; <i>GenRx Baclofen</i> ; <i>Stelax 10</i> ; <i>Terry White Chemists Baclofen</i>
<i>Lioresal 25</i>	Tablet 25 mg	100	3.01	<i>Baclo</i> ; <i>Baclohexal</i> ; <i>Chem mart Baclofen</i> ; <i>Clofen 25</i> ; <i>GenRx Baclofen</i> ; <i>Stelax 25</i> ; <i>Terry White Chemists Baclofen</i>
<i>Lipex 10</i>	Tablet 10 mg	30	0.71	<i>Chem mart Simvastatin</i> ; <i>GenRx Simvastatin</i> ; <i>Simvabell</i> ; <i>Simvahexal</i> ; <i>Simvar 10</i> ; <i>Simvastatin-DP</i> ; <i>Simvastatin-RL</i> ; <i>Terry White Chemists Simvastatin</i> ; <i>Zimstat</i>
<i>Lipex 20</i>	Tablet 20 mg	30	0.70	<i>Chem mart Simvastatin</i> ; <i>GenRx Simvastatin</i> ; <i>Simvabell</i> ; <i>Simvahexal</i> ; <i>Simvar 20</i> ; <i>Simvastatin-DP</i> ; <i>Simvastatin-RL</i> ; <i>Terry White Chemists Simvastatin</i> ; <i>Zimstat</i>
<i>Lipex 40</i>	Tablet 40 mg	30	0.70	<i>Chem mart Simvastatin</i> ; <i>GenRx Simvastatin</i> ; <i>Simvabell</i> ; <i>Simvahexal</i> ; <i>Simvar 40</i> ; <i>Simvastatin-DP</i> ; <i>Simvastatin-RL</i> ; <i>Terry White Chemists Simvastatin</i> ; <i>Zimstat</i>
<i>Lipex 80</i>	Tablet 80 mg	30	0.70	<i>Chem mart Simvastatin</i> ; <i>GenRx Simvastatin</i> ; <i>Simvabell</i> ; <i>Simvahexal</i> ; <i>Simvar 80</i> ; <i>Simvastatin-DP</i> ; <i>Simvastatin-RL</i> ; <i>Terry White Chemists Simvastatin</i> ; <i>Zimstat</i>
<i>Liquifilm Forte</i>	Eye drops 30 mg per mL (3%), 15 mL	1	5.98	<i>PVA Forte</i>
<i>Liquifilm Tears</i>	Eye drops 14 mg per mL (1.4%), 15 mL	1	1.70	<i>PVA Tears</i>
<i>Lomotil</i>	Tablet 2.5 mg-25 micrograms	20	1.83	<i>Lofenoxal</i>
<i>Lopid</i>	Tablet 600 mg	60	3.49	<i>Ausgem</i> ; <i>Chem mart Gemfibrozil</i> ; <i>Gemhexal</i> ; <i>GenRx Gemfibrozil</i> ; <i>Jezil</i> ; <i>Lipazil 600 mg</i> ; <i>Terry White Chemists Gemfibrozil</i>
<i>Losec Tablets</i>	Tablet 20 mg	30	2.75	<i>Acimax Tablets</i> ; <i>Meprazol</i>
<i>Luvox</i>	Tablet 50 mg	30	3.90	<i>Faverin 50</i> ; <i>Movox 50</i> ; <i>Voxam</i>
	Tablet 100 mg	30	3.90	<i>Faverin 100</i> ; <i>Movox 100</i> ; <i>Voxam</i>
<i>Maxamox</i>	Tablet 1 g	14	1.49	<i>Amoxycillin Sandoz</i>
<i>Microgynon 30</i>	Tablets 150 micrograms-30 micrograms, 21	4	9.76	<i>Levlen ED</i>
<i>Microgynon 30 ED</i>	Pack containing 21 tablets 150 micrograms-30 micrograms and 7 inert tablets	4	9.76	<i>Levlen ED</i>
<i>Minidiab</i>	Tablet 5 mg	100	4.10	<i>Melizide</i>
<i>Minipress</i>	Tablet 1 mg (base)	100	3.05	<i>Chem mart Prazosin</i> ; <i>GenRx Prazosin</i> ; <i>Prazohexal</i> ; <i>Pressin 1</i> ; <i>Terry White Chemists Prazosin</i>
	Tablet 2 mg (base)	100	3.15	<i>Chem mart Prazosin</i> ; <i>GenRx Prazosin</i> ; <i>Prazohexal</i> ; <i>Pressin 2</i> ; <i>Terry White Chemists Prazosin</i>
	Tablet 5 mg (base)	100	3.40	<i>Chem mart Prazosin</i> ; <i>GenRx Prazosin</i> ; <i>Prazohexal</i> ; <i>Pressin 5</i> ; <i>Terry White Chemists Prazosin</i>
<i>Minomycin-50</i>	Tablet 50 mg	60	1.12	<i>Akamin 50</i>
<i>Mobic</i>	Tablet 7.5 mg	30	1.85	<i>Movalis 7.5</i>
	Tablet 15 mg	30	1.85	<i>Movalis 15</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Moduretic</i>	Tablet 50 mg-5 mg	100	3.44	<i>Amizide</i>
<i>Mogadon</i>	Tablet 5 mg	25	2.02	<i>Alodorm</i>
	Tablet 5 mg	50	4.04	<i>Alodorm</i>
<i>Mylanta P</i>	Tablet 200 mg-200 mg	200	0.40	<i>Gelusil</i>
	Oral suspension 200 mg-200 mg per 5 mL, 500 mL	2	0.32	<i>Gelusil</i>
<i>Naprosyn</i>	Tablet 250 mg	100	3.10	<i>Inza 250</i>
	Tablet 500 mg	50	1.80	<i>Inza 500</i>
<i>Naprosyn SR750</i>	Tablet 750 mg (sustained release)	28	1.68	<i>Proxen SR 750</i>
<i>Naprosyn SR1000</i>	Tablet 1 g (sustained release)	28	1.77	<i>Proxen SR 1000</i>
<i>Natrilix</i>	Tablet 2.5 mg	90	3.37	<i>Chem mart Indapamide; Dapa-Tabs; GenRx Indapamide; Indahexal; Insig; Napamide 2.5 mg; Terry White Chemists Indapamide</i>
<i>Neoral 25</i>	Capsule 25 mg	60	4.80	<i>Cysporin</i>
<i>Neoral 50</i>	Capsule 50 mg	60	4.76	<i>Cysporin</i>
<i>Neoral 100</i>	Capsule 100 mg	60	4.80	<i>Cysporin</i>
<i>Nolvadex</i>	Tablet 10 mg (base)	60	2.16	<i>Genox 10; Tamoxen 10 mg; Tamoxifen Hexal</i>
<i>Nolvadex-D</i>	Tablet 20 mg (base)	60	3.94	<i>Chem mart Tamoxifen; Genox 20; GenRx Tamoxifen; Tamosin; Tamoxen 20 mg; Tamoxifen Hexal; Terry White Chemists Tamoxifen</i>
<i>Nordette 28</i>	Pack containing 21 tablets 150 micrograms- 30 micrograms and 7 inert tablets	4	9.76	<i>Monofeme 28</i>
<i>Noriday 28 Day</i>	Tablets 350 micrograms, 28	4	4.00	<i>Locilan 28 Day</i>
<i>Normison</i>	Tablet 10 mg	25	1.06	<i>Temaze; Temtabs</i>
	Tablet 10 mg	50	2.12	<i>Temaze; Temtabs</i>
<i>Noroxin</i>	Tablet 400 mg	14	3.73	<i>Chem mart Norfloxacin; GenRx Norfloxacin; Norflohexal; Nufloxib; Roxin; Terry White Chemists Norfloxacin; DP brand</i>
<i>Odrik</i>	Capsule 500 micrograms	28	1.00	<i>Gopten</i>
	Capsule 1 mg	28	1.01	<i>Gopten</i>
	Capsule 2 mg	28	0.99	<i>Gopten</i>
<i>Ogen .625</i>	Tablet 730 micrograms (equivalent to 625 micrograms sodium oestrone sulfate)	56	1.48	<i>Genoral 0.625</i>
<i>Ogen 1.25</i>	Tablet 1.46 mg (equivalent to 1.25 mg sodium oestrone sulfate)	56	1.49	<i>Genoral 1.25</i>
<i>Oroxine</i>	Tablet equivalent to 50 micrograms anhydrous thyroxine sodium	200	1.30	<i>Eutroxsig</i>
	Tablet equivalent to 100 micrograms anhydrous thyroxine sodium	200	1.30	<i>Eutroxsig</i>
	Tablet equivalent to 200 micrograms anhydrous thyroxine sodium	200	1.30	<i>Eutroxsig</i>
<i>Orudis SR 200</i>	Capsule 200 mg (sustained release)	28	2.01	<i>Oruvail SR</i>
<i>Panadeine Forte</i>	Tablet 30 mg-500 mg	20	1.89	<i>Codalgin Forte; Codapane Forte; Dolaforte; Dymadon Forte; Prodeine Forte</i>
	Tablet 30 mg-500 mg	60	5.67	<i>Codalgin Forte; Codapane Forte; Dolaforte; Dymadon Forte; Prodeine Forte</i>
<i>Panadol Osteo</i>	Tablet 665 mg (modified release)	192	4.88	<i>Duatrol SR</i>
<i>Panafcort</i>	Tablet 1 mg	100	0.62	<i>Predsone</i>
<i>Panafcortelone</i>	Tablet 1 mg	100	0.62	<i>Predsolone</i>
<i>Parlodel</i>	Tablet 2.5 mg (base)	30	2.94	<i>Bromohexal; Krypton 2.5</i>
	Tablet 2.5 mg (base)	60	3.00	<i>Bromohexal; Krypton 2.5</i>
	Capsule 5 mg (base)	60	3.00	<i>Bromohexal; Krypton 5</i>
	Capsule 10 mg (base)	100	3.00	<i>Krypton 10</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Pepcidine</i>	Tablet 40 mg	30	5.41	<i>Ausfam 40; Chem mart Famotidine; Famohexal; GenRx Famotidine; Pamacid 40; Pepzan; Terry White Chemists Famotidine</i>
<i>Pepcidine M</i>	Tablet 20 mg	60	5.41	<i>Ausfam 20; Chem mart Famotidine; Famohexal; GenRx Famotidine; Pamacid 20; Pepzan; Terry White Chemists Famotidine</i>
<i>Plendil ER</i>	Tablet 2.5 mg (extended release)	30	3.00	<i>Felodur ER 2.5 mg</i>
	Tablet 5 mg (extended release)	30	3.36	<i>Felodur ER 5 mg</i>
	Tablet 10 mg (extended release)	30	4.65	<i>Felodur ER 10 mg</i>
<i>Pravachol</i>	Tablet 10 mg	30	1.75	<i>Lipostat 10</i>
	Tablet 20 mg	30	1.74	<i>Lipostat 20</i>
	Tablet 40 mg	30	1.75	<i>Lipostat 40</i>
	Tablet 80 mg	30	1.75	<i>Lipostat 80</i>
<i>Prinivil 5</i>	Tablet 5 mg	30	1.95	<i>Chem mart Lisinopril; Fibsol 5; GenRx Lisinopril; Liprace; Lisinobell; Lisinopril Hexal; Lisodur; Terry White Chemists Lisinopril</i>
<i>Prinivil 10</i>	Tablet 10 mg	30	1.95	<i>Chem mart Lisinopril; Fibsol 10; GenRx Lisinopril; Liprace; Lisinobell; Lisinopril Hexal; Lisodur; Terry White Chemists Lisinopril</i>
<i>Prinivil 20</i>	Tablet 20 mg	30	1.95	<i>Chem mart Lisinopril; Fibsol 20; GenRx Lisinopril; Liprace; Lisinobell; Lisinopril Hexal; Lisodur; Terry White Chemists Lisinopril</i>
<i>Prothiaden</i>	Capsule 25 mg	50	1.07	<i>Dothep 25</i>
	Tablet 75 mg	30	1.05	<i>Dothep 75</i>
<i>Provera</i>	Tablet 5 mg	56	1.77	<i>Ralovera</i>
	Tablet 10 mg	30	1.78	<i>Medroxyhexal; Ralovera</i>
	Tablet 10 mg	100	1.65	<i>Ralovera</i>
<i>Prozac Tab</i>	Tablet 20 mg (base) (dispersible)	28	4.10	<i>Lovan 20 Tab</i>
<i>Prozac 20</i>	Capsule 20 mg (base)	28	4.10	<i>Auscap; Chem mart Fluoxetine; Fluohexal; Fluoxebell; Fluoxetine-DP; GenRx Fluoxetine; Lovan; Terry White Chemists Fluoxetine; Zactin</i>
<i>P.V. Carpine</i>	Eye drops 5 mg per mL (0.5%), 15 mL	1	2.00	<i>Pilopt</i>
	Eye drops 10 mg per mL (1%), 15 mL	1	2.00	<i>Pilopt</i>
	Eye drops 20 mg per mL (2%), 15 mL	1	2.12	<i>Pilopt</i>
	Eye drops 30 mg per mL (3%), 15 mL	1	2.60	<i>Pilopt</i>
	Eye drops 40 mg per mL (4%), 15 mL	1	2.60	<i>Pilopt</i>
	Eye drops 60 mg per mL (6%), 15 mL	1	1.98	<i>Pilopt</i>
<i>Quinate</i>	Tablet 300 mg	50	2.19	<i>Quinsul</i>
<i>Redipred</i>	Oral solution equivalent to 5 mg prednisolone per mL, 30 mL	1	1.53	<i>PredMix</i>
<i>Remeron</i>	Tablet 30 mg	30	26.69	<i>Axit 30; Mirtazon</i>
<i>Renitec</i>	Tablet 10 mg	30	3.15	<i>Alphapril; Auspril; Chem mart Enalapril; Enahexal; Enalabell; Enalapril-DP 10mg; GenRx Enalapril; Terry White Chemists Enalapril</i>
<i>Renitec M</i>	Tablet 5 mg	30	3.15	<i>Alphapril; Auspril; Chem mart Enalapril; Enahexal; Enalabell; Enalapril-DP 5mg; GenRx Enalapril; Terry White Chemists Enalapril</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Renitec 20</i>	Tablet 20 mg	30	3.15	<i>Alphapril; Auspril; Chem mart Enalapril; Enahexal; Enalabell; Enalapril-DP 20mg; GenRx Enalapril; Terry White Chemists Enalapril</i>
<i>Ritalin 10</i>	Tablet 10 mg	100	1.28	<i>Attenta</i>
<i>Rivotril</i>	Tablet 500 micrograms	200	4.74	<i>Paxam 0.5</i>
	Tablet 2 mg	200	5.36	<i>Paxam 2</i>
<i>Roaccutane</i>	Capsule 20 mg	60	2.50	<i>Chem mart Isotretinoin; GenRx Isotretinoin; Isohexal; Oratane; Terry White Chemists Isotretinoin</i>
<i>Rulide</i>	Tablet 150 mg	10	2.50	<i>Biaxsig</i>
	Tablet 300 mg	5	2.50	<i>Biaxsig</i>
<i>Salazopyrin-EN</i>	Tablet 500 mg (enteric coated)	200	1.64	<i>Pyralin EN</i>
<i>Septtrin</i>	Tablet 80 mg-400 mg	10	1.54	<i>Resprim</i>
	Oral suspension 40 mg-200 mg per 5 mL, 100 mL	1	1.91	<i>Resprim</i>
<i>Septtrin Forte</i>	Tablet 160 mg-800 mg	10	1.46	<i>Bactrim DS; Chem mart Trimethoprim with Sulfamethoxazole DS; GenRx Trimethoprim with Sulfamethoxazole DS; Resprim Forte; Terry White Chemists Trimethoprim with Sulfamethoxazole DS</i>
<i>Serepax</i>	Tablet 15 mg	25	1.44	<i>Alepam 15</i>
	Tablet 30 mg	25	1.45	<i>Alepam 30; Murelax</i>
	Tablet 15 mg	50	2.88	<i>Alepam 15</i>
	Tablet 30 mg	50	2.90	<i>Alepam 30; Murelax</i>
<i>Serophene</i>	Tablet 50 mg	10	0.18	<i>Clomhexal; Fermil; GenRx Clomiphene</i>
<i>Sigmacort</i>	Cream 10 mg per g (1%), 30 g	1	1.30	<i>Cortic-DS 1%</i>
	Cream 10 mg per g (1%), 50 g	1	1.30	<i>Cortic-DS 1%</i>
	Topical ointment 10 mg per g (1%), 30 g	1	1.30	<i>Cortic-DS 1%</i>
	Topical ointment 10 mg per g (1%), 50 g	1	1.30	<i>Cortic-DS 1%</i>
<i>Sinemet</i>	Tablet 250 mg-25 mg	100	3.20	<i>Levohexal</i>
<i>Sinemet 100/25</i>	Tablet 100 mg-25 mg	100	5.41	<i>Kinson</i>
<i>Slow-K</i>	Tablet 600 mg (sustained release)	200	2.66	<i>Duro-K</i>
<i>Sofradex</i>	Ear drops 500 micrograms-5 mg- 50 micrograms per mL, 8 mL	1	1.80	<i>Otodex</i>
<i>Sotacor</i>	Tablet 80 mg	60	1.66	<i>GenRx Sotalol; Solavert; Sotahexal</i>
	Tablet 160 mg	60	1.80	<i>Cardol; Chem mart Sotalol; GenRx Sotalol; Solavert; Sotab; Sotahexal; Terry White Chemists Sotalol</i>
<i>Stemetil</i>	Tablet 5 mg	25	2.15	<i>Stemzine</i>
<i>Synphasic</i>	Pack containing 12 tablets 500 micrograms- 35 micrograms, 9 tablets 1 mg- 35 micrograms and 7 inert tablets	4	7.96	<i>Improvil 28 Day</i>
<i>Tagamet</i>	Tablet 200 mg	120	2.49	<i>Cimehexal; Magicul 200</i>
	Tablet 400 mg	60	2.49	<i>Cimehexal; GenRx Cimetidine; Magicul 400</i>
	Tablet 800 mg	30	4.96	<i>Cimehexal; GenRx Cimetidine; Magicul 800</i>
<i>Tambocor</i>	Tablet 100 mg	60	2.86	<i>Flecatab</i>
<i>Tazac</i>	Capsule 150 mg	60	2.60	<i>Nizac</i>
	Capsule 300 mg	30	2.60	<i>Nizac</i>
<i>Tears Naturale</i>	Eye drops 3 mg-1 mg per mL (0.3%-0.1%), 15 mL	1	1.67	<i>Poly-Tears</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Tegretol 100</i>	Tablet 100 mg	200	2.50	<i>Carbamazepine Sandoz;</i> <i>Carbamazepine-BC</i>
<i>Tegretol 200</i>	Tablet 200 mg	200	3.00	<i>Carbamazepine Sandoz;</i> <i>Carbamazepine-BC; Teril</i>
<i>Tenormin</i>	Tablet 50 mg	30	3.55	<i>Anselol 50 mg; Atehexal;</i> <i>Chem mart Atenolol;</i> <i>GenRx Atenolol; Noten; Tensig;</i> <i>Terry White Chemists Atenolol</i>
<i>Ticlid</i>	Tablet 250 mg	60	2.26	<i>Ticlopidine Hexal; Tilodene</i>
<i>Timoptol</i>	Eye drops 2.5 mg (base) per mL (0.25%), 5 mL	1	1.09	<i>Tenopt</i>
	Eye drops 5 mg (base) per mL (0.5%), 5 mL	1	1.15	<i>Tenopt</i>
<i>Tolvon</i>	Tablet 10 mg	50	1.29	<i>Lumin 10</i>
	Tablet 20 mg	50	1.60	<i>Lumin 20</i>
<i>Tramal</i>	Capsule 50 mg	20	0.60	<i>Chem mart Tramadol; GenRx</i> <i>Tramadol; Terry White Chemists</i> <i>Tramadol; Zydol</i>
<i>Tramal SR 100</i>	Tablet 100 mg (sustained release)	20	1.04	<i>Tramahexal SR; Zydol SR 100</i>
<i>Tramal SR 150</i>	Tablet 150 mg (sustained release)	20	1.04	<i>Tramahexal SR; Zydol SR 150</i>
<i>Tramal SR 200</i>	Tablet 200 mg (sustained release)	20	1.05	<i>Tramahexal SR; Zydol SR 200</i>
<i>Trandate</i>	Tablet 100 mg	100	2.84	<i>Presolol 100</i>
	Tablet 200 mg	100	2.97	<i>Presolol 200</i>
<i>Triphasil 28</i>	Pack containing 6 tablets 50 micrograms- 30 micrograms, 5 tablets 75 micrograms- 40 micrograms, 10 tablets 125 micrograms- 30 micrograms and 7 inert tablets	4	9.76	<i>Trifeme 28</i>
<i>Triprim</i>	Tablet 300 mg	7	1.36	<i>Alprim</i>
<i>Triquilar ED</i>	Pack containing 6 tablets 50 micrograms- 30 micrograms, 5 tablets 75 micrograms- 40 micrograms, 10 tablets 125 micrograms- 30 micrograms and 7 inert tablets	4	9.76	<i>Logynon ED</i>
<i>Tritace 1.25 mg</i>	Tablet 1.25 mg	30	2.00	<i>Ramace 1.25 mg</i>
<i>Tritace 2.5 mg</i>	Tablet 2.5 mg	30	2.01	<i>Ramace 2.5 mg</i>
<i>Tritace 5 mg</i>	Tablet 5 mg	30	2.00	<i>Ramace 5 mg</i>
<i>Tryptanol</i>	Tablet 25 mg	50	1.25	<i>Endep 25</i>
<i>Valium</i>	Tablet 2 mg	50	1.30	<i>Antenex 2; Valpam 2</i>
	Tablet 5 mg	50	1.32	<i>Antenex 5; Diazepam-DP; Valpam 5</i>
<i>Vancocin</i>	Powder for injection 500 mg (500,000 i.u.) vancomycin activity	2	0.36	<i>MX brand</i>
	Powder for injection 500 mg (500,000 i.u.) vancomycin activity	5	0.90	<i>MX brand</i>
<i>Ventolin CFC-free</i>	Oral pressurised inhalation 100 micrograms (base) per dose (200 doses), CFC-free formulation	2	1.00	<i>Asmol CFC-free; Epaq</i>
<i>Ventolin Nebules</i>	Nebuliser solution single dose units 2.5 mg (base) in 2.5 mL, 30	2	2.20	<i>Asmol 2.5 uni-dose; Butamol 2.5;</i> <i>Chem mart Salbutamol; GenRx</i> <i>Salbutamol; Terry White Chemists</i> <i>Salbutamol; PU brand</i>
	Nebuliser solution single dose units 5 mg (base) in 2.5 mL, 30	2	2.20	<i>Asmol 5 uni-dose; Butamol 5;</i> <i>Chem mart Salbutamol;</i> <i>GenRx Salbutamol;</i> <i>Terry White Chemists Salbutamol</i>
<i>Vibramycin</i>	Tablet 100 mg	7	1.60	<i>Chem mart Doxycycline; Doxsig;</i> <i>Doxy-100; Doxyhexal; Doxylin 100;</i> <i>GenRx Doxycycline;</i> <i>Terry White Chemists Doxycycline</i>
	Tablet 100 mg	28	6.40	<i>Chem mart Doxycycline; Doxsig;</i> <i>Doxy-100; Doxyhexal; Doxylin 100;</i> <i>GenRx Doxycycline;</i> <i>Terry White Chemists Doxycycline</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Vibramycin</i>	Tablet 100 mg	21	4.80	<i>Chem mart Doxycycline; Doxsig; Doxy-100; Doxyhexal; Doxylin 100; GenRx Doxycycline; Terry White Chemists Doxycycline</i>
<i>Vibra-Tabs</i>	Tablet 50 mg	25	1.65	<i>Chem mart Doxycycline; Doxy-50; Doxyhexal; Doxylin 50; Frakas; GenRx Doxycycline; Terry White Chemists Doxycycline</i>
<i>Viscotears Liquid Gel</i>	Ocular lubricating gel 2 mg per g (0.2%), 10 g	1	1.00	<i>PAA</i>
<i>Visken 5</i>	Tablet 5 mg	100	2.80	<i>Barbloc 5</i>
<i>Visken 15</i>	Tablet 15 mg	50	2.80	<i>Barbloc 15</i>
<i>Voltaren 25</i>	Tablet 25 mg (enteric coated)	100	3.30	<i>Chem mart Diclofenac; Clonac 25; Diclofenac-BC; Diclohexal; Dinac; Fenac 25; GenRx Diclofenac; Terry White Chemists Diclofenac</i>
<i>Voltaren 50</i>	Tablet 50 mg (enteric coated)	50	3.21	<i>Chem mart Diclofenac; Clonac 50; Diclohexal; Dinac; Fenac; GenRx Diclofenac; Terry White Chemists Diclofenac</i>
<i>Xanax</i>	Tablet 250 micrograms	50	1.08	<i>Alprax 0.25; Kalma 0.25; Zamhexal 0.25mg</i>
	Tablet 500 micrograms	50	1.16	<i>Alprax 0.5; Kalma 0.5; Zamhexal 0.5mg</i>
	Tablet 1 mg	50	1.35	<i>Alprax 1; Alprazolam-DP; Chem mart Alprazolam; GenRx Alprazolam; Kalma 1; Terry White Chemists Alprazolam; Zamhexal 1.0mg</i>
<i>Xanax Tri-Score</i>	Tablet 2 mg	50	1.65	<i>Alprax 2; Alprazolam-DP; Chem mart Alprazolam; GenRx Alprazolam; Kalma 2; Terry White Chemists Alprazolam; Zamhexal 2mg</i>
<i>Zantac</i>	Tablet 150 mg (base)	60	2.17	<i>Ausran; Chem mart Ranitidine; GenRx Ranitidine; Rani 2; Ranihexal; Ranoxyl; Terry White Chemists Ranitidine</i>
	Tablet 300 mg (base)	30	2.17	<i>Ausran; Chem mart Ranitidine; GenRx Ranitidine; Rani 2; Ranihexal; Ranoxyl; Terry White Chemists Ranitidine</i>
<i>Zestril</i>	Tablet 5 mg	30	2.15	<i>Chem mart Lisinopril; Fibsol 5; GenRx Lisinopril; Liprace; Lisinobell; Lisinopril Hexal; Lisodur; Terry White Chemists Lisinopril</i>
	Tablet 10 mg	30	2.15	<i>Chem mart Lisinopril; Fibsol 10; GenRx Lisinopril; Liprace; Lisinobell; Lisinopril Hexal; Lisodur; Terry White Chemists Lisinopril</i>
	Tablet 20 mg	30	2.15	<i>Chem mart Lisinopril; Fibsol 20; GenRx Lisinopril; Liprace; Lisinobell; Lisinopril Hexal; Lisodur; Terry White Chemists Lisinopril</i>
<i>Zocor</i>	Tablet 5 mg	30	0.71	<i>Simvabell; Simvahexal; Simvastatin-RL; Zimstat</i>

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Zocor</i>	Tablet 10 mg	30	0.71	<i>Chem mart Simvastatin;</i> <i>GenRx Simvastatin; Simvabell;</i> <i>Simvahexal; Simvar 10;</i> <i>Simvastatin-DP; Simvastatin-RL;</i> <i>Terry White Chemists Simvastatin;</i> <i>Zimstat</i>
	Tablet 20 mg	30	0.70	<i>Chem mart Simvastatin;</i> <i>GenRx Simvastatin; Simvabell;</i> <i>Simvahexal; Simvar 20;</i> <i>Simvastatin-DP; Simvastatin-RL;</i> <i>Terry White Chemists Simvastatin;</i> <i>Zimstat</i>
	Tablet 40 mg	30	0.70	<i>Chem mart Simvastatin;</i> <i>GenRx Simvastatin; Simvabell;</i> <i>Simvahexal; Simvar 40;</i> <i>Simvastatin-DP; Simvastatin-RL;</i> <i>Terry White Chemists Simvastatin;</i> <i>Zimstat</i>
	Tablet 80 mg	30	0.70	<i>Chem mart Simvastatin;</i> <i>GenRx Simvastatin; Simvabell;</i> <i>Simvahexal; Simvar 80;</i> <i>Simvastatin-DP; Simvastatin-RL;</i> <i>Terry White Chemists Simvastatin;</i> <i>Zimstat</i>
<i>Zoloft</i>	Tablet 50 mg (base)	30	0.55	<i>Chem mart Sertraline; Concorz;</i> <i>Eleva 50; GenRx Sertraline;</i> <i>Sertraline-DP;</i> <i>Terry White Chemists Sertraline;</i> <i>Xydep 50</i>
	Tablet 100 mg (base)	30	0.55	<i>Chem mart Sertraline; Concorz;</i> <i>Eleva 100; GenRx Sertraline;</i> <i>Sertraline-DP;</i> <i>Terry White Chemists Sertraline;</i> <i>Xydep 100</i>
	Tablet 50 mg (base)	30	0.55	<i>Eleva 50; Xydep 50</i>
	Tablet 100 mg (base)	30	0.55	<i>Eleva 100; Xydep 100</i>
<i>Zovirax 200 mg</i>	Tablet 200 mg	50	5.78	<i>Acihexal; Acyclo-V 200;</i> <i>GenRx Aciclovir; Lovir</i>
	Tablet 200 mg	90	4.28	<i>Acihexal; Acyclo-V 200;</i> <i>Chem mart Aciclovir;</i> <i>GenRx Aciclovir; Lovir;</i> <i>Terry White Chemists Aciclovir;</i> <i>Zyclir 200</i>
<i>Zovirax 800 mg</i>	Tablet 800 mg	35	2.09	<i>Acihexal; Acyclo-V 800;</i> <i>GenRx Aciclovir; Lovir; Zyclir 800</i>
<i>Zyloprim</i>	Tablet 800 mg	120	7.19	<i>Acihexal; Acyclo-V 800; Lovir</i>
	Tablet 100 mg	200	2.43	<i>Allohexal; Allosig;</i> <i>Chem mart Allopurinol;</i> <i>GenRx Allopurinol; Progout 100;</i> <i>Terry White Chemists Allopurinol</i>
	Tablet 300 mg	60	2.28	<i>Allohexal; Allosig;</i> <i>Chem mart Allopurinol;</i> <i>GenRx Allopurinol; Progout 300;</i> <i>Terry White Chemists Allopurinol</i>

**Preparations Available to Members of
Colostomy and Ileostomy Associations
Effective 1 DECEMBER 2005**

Code	Name, Manner of Administration and Form	Max. Qty	Unit Price \$	Proprietary Name and Manufacturer	
3206R	CHARCOAL, ACTIVATED Tablets 300 mg, 500	1	15.00	Karbons	EG

**Preparations Available to Members of
Paraplegic and Quadriplegic Associations
Effective 1 DECEMBER 2005**

Code	Name, Manner of Administration and Form	Max. Qty	Unit Price \$	Proprietary Name and Manufacturer	
	BISACODYL				
3261P	Tablets 5 mg, 200	2	7.50	Bisalax Lax-Tab	AS AE
3250C	Suppositories 10 mg, 10	9	5.08	Durolax Petrus Bisacodyl Suppositories	BY PP
3276K	Suppositories 10 mg, 12	8	3.86	Fleet Laxative Suppositories Petrus Bisacodyl Suppositories	FL PP
3263R	Enemas 10 mg in 5 mL, 25	2	28.20	Bisalax	AS
	DOCUSATE SODIUM with BISACODYL				
3253F	Suppositories 100 mg-10 mg, 5	18	2.54	Coloxyl	FM
	GLYCEROL				
3265W	Suppositories 700 mg (for infants), 12	8	3.04	PP	
3266X	Suppositories 1.4 g (for children), 12	8	3.15	PP	
3267Y	Suppositories 2.8 g (for adults), 12	8	3.26	PP	
	SORBITOL with SODIUM CITRATE and SODIUM LAURYL SULFOACETATE				
3274H	Enemas 3.125 g-450 mg-45 mg in 5 mL, 12	4	13.54	Microlax	PH
	STERCULIA with FRANGULA BARK				
3275J	Granules 620 mg-80 mg per g (62%-8%), 500 g	2	16.64	Normacol Plus	NE