



Australian Government

Department of Health and Ageing

SCHEDULE OF PHARMACEUTICAL BENEFITS

This Schedule is also available on the internet at
www.pbs.gov.au

**EFFECTIVE 1 January 2011 -
31 January 2011
(ALL PREVIOUS EDITIONS CANCELLED)**

© Commonwealth of Australia 2011

ISSN 1037-3667

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced by any process without prior written permission from the Commonwealth. Requests and inquiries concerning reproduction and rights should be addressed to the Commonwealth Copyright Administration, Attorney General's Department, Robert Garran Offices, National Circuit, Barton ACT 2600 or posted at <http://www.ag.gov.au/cca>

This Schedule provides information on the arrangements for the prescribing and supply of pharmaceutical benefits. These arrangements operate under the *National Health Act 1953*. However, at the time of printing, the relevant legislation giving authority for the changes included in this issue of the Schedule may still be subject to the usual Parliamentary scrutiny. This book is not a legal document, and, in cases of discrepancy, the legislation will be the source document for payment for the supply of pharmaceutical benefits. The legislation is available from the Federal Register of Legislative Instruments website at <http://www.frl.gov.au>.

The information is not intended to give or replace any legal, medical, dental or optometrical advice. This document is not a legal document and does not constitute legal advice. Neither the information nor this document can be relied upon without first seeking and obtaining independent legal, medical, dental or optometrical advice beforehand. To the extent permitted by law, the Commonwealth of Australia will not be held responsible, nor accept any liability (whether arising out of negligence or otherwise), for any injury, damages, costs, expenses and losses suffered or incurred by a person where such a person has relied on this document or used the information in it as legal, medical, dental or optometrical advice.

Contents

Addresses — Medicare Australia.....	14
Authority Prescription Applications.....	15
Requests for Drugs via the Special Access Scheme (SAS).....	15
Department of Veterans' Affairs.....	15
Telephone Interpreter Service.....	15
Poisons Information Centres.....	16
Drug Information Centres.....	16
List of Contact Officers for Recalls of Therapeutic Goods.....	17
Section 1 — Explanatory Notes.....	27
Introduction.....	27
1. The Schedule — Where to Find What.....	27
Section 1	27
Section 2	27
Section 3	28
Section 4	28
Repatriation Schedule of Pharmaceutical Benefits	28
2. Prescribing Medicines — Information for PBS Prescribers	28
PBS prescribers	28
PBS Prescription forms.....	28
Ordering forms	29
Preparing general PBS prescriptions	29
Do's and Don't's.....	29
Writing the PBS prescription.....	30
Restrictions	30
Authority PBS prescriptions	31
Authority required PBS Prescriptions	31
Authority required (STREAMLINED)	
PBS Prescriptions	31
Writing authority PBS prescriptions.....	32
Maximum quantities and repeats	32
Regulation 24	33
Urgent cases.....	33
Drugs of addiction	33
Emergency drug supplies	33
Availability of Methoxyflurane for emergency	
treatment only	33
Improving the capacity of the PBS to meet particular	
Aboriginal and Torres Strait Islander health needs	34
How to prescribe these items?	34
Aboriginal and Torres Strait Islander	
identification.....	34
Asking about Aboriginal and/or Torres	
Strait Islander identification	34
Aboriginal and Torres Strait Islander health	35
Communication and cultural issues	35
3. Supplying Medicines — What Pharmacists Need to Know	36
Eligible suppliers	36
Approval conditions for pharmacists	36
Before supplying pharmaceutical benefits	36
Supplying pharmaceutical benefits	37
Do's and Don'ts	37
What to do if the Schedule changes	37
Suspected forgery	37
Regulation 24	37
Repeat authorisations	38
Preparing Repeat Authorisation Forms.....	38
Repeat authorisations for injectables	
and solvents.....	38
Repeat authorisations for deferred supply	38
Authority PBS prescriptions	39
Urgent cases.....	39
Receipts.....	39
Emergency drug supplies	39
4. Patient Charges.....	39
Type of patient	39
Establishing entitlement	40
What to charge	40
Patient contribution.....	40
Patient contributions for early supply of	
some PBS medicines	40
Special patient contributions, brand	
premiums and therapeutic	
group premiums	41
Solvents	41
Increased quantities	41
Regulation 24	41
After hours.....	41
Delivery.....	41
5. The Safety Net Scheme.....	41
Safety net thresholds	42
Safety net cross-over arrangements	42
Recording PBS prescriptions.....	43
Hospital prescription record forms	43
Multi-item prescription forms.....	43
Qualifying PBS prescriptions	43
Lost prescription record forms.....	43
Retrospective entitlement and patient refunds	44
Applying for a Safety Net Entitlement/Concession Card	44
Issuing a Safety Net Entitlement/Concession Card.....	44
Issuing supplementary cards.....	44
Notification to Medicare Australia and claim for payment	45
Lost Safety Net Entitlement/Concession Cards	45
Pharmacy record of issued cards.....	45
6. Medicare Australia Entitlement Checks	45

General Patients	45	Special provisions for extemporaneous PBS prescriptions outside the Standard Formulae List	51
Concessional Patients	45	10. Miscellaneous	52
Entitlement checking procedures	45	References	52
General Patients	45	Standards	52
Concessional Patients	46	Legislation	52
Step by step	46	Nurse practitioner PBS prescribing.....	52
7. How Pharmacists Claim Reimbursement:		Midwife PBS prescribing.....	53
Information Required	46	SECTION 2	61
PBS Prescription identification	46	Contents	62
Serial numbers	46	Symbols	62
Repeat authorisations for authority PBS prescriptions	47	Emergency Drug Supplies	64
Repeat authorisations for deferred supply	47	Special Pharmaceutical Benefits	66
Injectable item ordered with a solvent	47	General Pharmaceutical Benefits	68
Dropper containers	47	Pharmaceutical Benefits for Palliative Care	468
Extemporaneously-prepared pharmaceutical benefits not listed in the Standard Formulae List	47	Pharmaceutical Benefits for Dental Use	486
PBS prescriptions paid on an average price basis	47	Pharmaceutical Benefits for Optometrical Use	514
Pricing non-pre-priced extemporaneous preparations	47	Items Available under Special Arrangements (section 100)	522
RPBS prescriptions for items not included in either the PBS or RPBS Schedule	47	SECTION 3 - STANDARD PACKS AND PRICES	801
Payment to Pharmacists for Dispensing Premium-free Substitutable Medicines.....	48	Container Prices and Fees for Ready Prepared Pharmaceutical Benefits	801
8. How Pharmacists Claim Reimbursement: Documents		Standard Packs and Prices for Ready Prepared Pharmaceutical Benefits	801
to be Submitted	48	SECTION 4	
Completing the claim form.....	48	- EXTemporaneously-Prepared	
Lodging claims.....	48	PHARMACEUTICAL BENEFITS	813
Reconciliation statements.....	48	Drug Tariff	814
9. Pricing PBS Prescriptions	49	Container Prices	818
Pricing principles	49	Standard Formula Preparations Explanation	819
Pricing dates.....	49	Table of Codes, Maximum Quantities, and Number of Repeats for Extemporaneously Prepared Pharmaceutical Benefits	822
Pricing ready-prepared items.....	49	REPatriation PHARMACEUTICAL BENEFITS.....	823
For maximum quantities.....	49	Beneficiaries' Entitlement Cards	824
For lesser quantities	49	Explanatory Notes	826
Wastage table percentage	50	Therapeutic Index for RPBS Schedule	833
Pricing extemporaneously-prepared items	50	SECTION 1 - Drugs, Medicines and Dressing	836
General	50	SECTION 2 - Standard Packs and Prices	876
Pricing of ingredients	50	GENERIC/PROPRIETARY INDEX	879
Pricing PBS prescriptions where extra ingredients are added to a formula	51		
Containers	51		

PHARMACEUTICAL BENEFITS

These changes to the Schedule of Pharmaceutical Benefits are effective from 1 January 2011. The Schedule is updated on the first day of each month and is available on the Internet at www.pbs.gov.au.

Fees, Patient Contributions and Safety Net Thresholds

The following fees, patient contributions and safety net thresholds apply as at 1 January 2011 and are included, where applicable, in prices published in the Schedule —

Dispensing Fees:	Ready-prepared	\$6.42
	Dangerous drug fee	\$2.71
	Extemporaneously-prepared	\$8.46
	Allowable additional patient charge*	\$3.92
Additional Fees (for safety net prices):	Ready-prepared	\$1.07
	Extemporaneously-prepared	\$1.41
Patient Co-payments:	General	\$34.20
	Concessional	\$5.60
Safety Net Thresholds:	General	\$1317.20
	Concessional	\$336.00
Safety Net Card Issue Fee:		\$8.58

*The allowable additional patient charge is a discretionary charge to general patients if a pharmaceutical item has a dispensed price for maximum quantity less than the general patient co-payment. The pharmacist may charge general patients the allowable additional fee but the fee cannot take the cost of the prescription above the general patient co-payment for the medicine. This fee does not count towards the Safety Net threshold.

SUMMARY OF CHANGES

ADDITIONS

Additions – Items

5459H	Amlodipine With Valsartan , Tablet 5 mg (as besylate)-320 mg (Exforge 5/320)
5460J	Amlodipine With Valsartan , Tablet 10 mg (as besylate)-320 mg (Exforge 10/320)
5464N	Docetaxel , Solution concentrate for I.V. infusion 80 mg in 4 mL (Taxotere)
5462L	Docetaxel , Solution concentrate for I.V. infusion 20 mg in 1 mL (Taxotere)
5463M	Docetaxel , Solution concentrate for I.V. infusion 20 mg in 1 mL (Taxotere) (Diff.Max. Qty)
5461K	Mesalazine , Suppository (moulded) 1 g (Salofalk)
5458G	Quetiapine , Tablet (modified release) 150 mg (as fumarate) (Seroquel XR)

Additions – Brands

8255L	<i>APO-Carvedilol, TX</i> – Carvedilol , Tablet 3.125 mg
9354H	<i>Clopidogrel Actavis, GQ</i> – Clopidogrel , Tablet 75 mg (as besilate)
1370D	<i>Enalapril-GA, GM</i> – Enalapril , Tablet containing enalapril maleate 5 mg
8049P	<i>Gemcitabine Kabi, PK; Gemcitabine Sun, ZF</i> – Gemcitabine , Powder for I.V. infusion 200 mg (as hydrochloride)
8050Q	<i>Gemcitabine Kabi, PK; Gemcitabine Sun, ZF</i> – Gemcitabine , Powder for I.V. infusion 1 g (as hydrochloride)
9414L	<i>Gemcitabine Kabi, PK</i> – Gemcitabine , Powder for I.V. infusion 2 g (as hydrochloride)
8679T	<i>Lercan, SI; Lercanidipine Sandoz, SZ</i> – Lercanidipine Hydrochloride , Tablet 20 mg
8534E	<i>Lercan, SI; Lercanidipine Sandoz, SZ</i> – Lercanidipine Hydrochloride , Tablet 10 mg
1327W	<i>Omepro-GA, GM; Pemzo, SI; Pharmacor Omeprazole 20, CR</i> – Omeprazole , Capsule 20 mg
1326T	<i>Omepro-GA, GM; Pemzo, SI; Pharmacor Omeprazole 20, CR</i> – Omeprazole , Capsule 20 mg
8847P	<i>Oxaliplatin Kabi, PK</i> – Oxaliplatin , Solution concentrate for I.V. infusion 50 mg in 10 mL
8539K	<i>Xalox, WQ</i> – Oxaliplatin , Powder for I.V. infusion 50 mg
8540L	<i>Xalox, WQ</i> – Oxaliplatin , Powder for I.V. infusion 100 mg
8360B	<i>Paclitaxel Kabi, PK</i> – Paclitaxel , Solution concentrate for I.V. infusion 300 mg in 50 mL
3026G	<i>Paclitaxel Kabi, PK</i> – Paclitaxel , Solution concentrate for I.V. infusion 30 mg in 5 mL

- 8018B *Paclitaxel Kabi, PK – Paclitaxel*, Solution concentrate for I.V. infusion 100 mg in 16.7 mL
 9121C *Ramipril-GA, GM – Ramipril*, Capsule 2.5 mg
 8470T *Ramipril-GA, GM – Ramipril*, Capsule 10 mg
 8144P *APO-Sumatriptan, TX; Chem mart Sumatriptan, CH; Terry White Chemists Sumatriptan, TW – Sumatriptan*, Tablet 50 mg (as succinate)

Additions – Bioequivalence Indicator

- 9414L *DBL Gemcitabine for injection, HH – Gemcitabine*, Powder for I.V. infusion 2 g (as hydrochloride)
 1326T *Probitor, SZ – Omeprazole*, Capsule 20 mg
 1327T *Probitor, SZ – Omeprazole*, Capsule 20 mg

DELETIONS

Deletion – Items

- 2670M **Testosterone Esters**, Injection 100 mg (Sustanon 100)
 5413X **Fentanyl**, Lozenges 200 micrograms (as citrate), 3 (Actiq)
 5414Y **Fentanyl**, Lozenges 400 micrograms (as citrate), 3 (Actiq)
 5415B **Fentanyl**, Lozenges 600 micrograms (as citrate), 3 (Actiq)
 5416C **Fentanyl**, Lozenges 800 micrograms (as citrate), 3 (Actiq)
 5417D **Fentanyl**, Lozenges 1200 micrograms (as citrate), 3 (Actiq)
 5418E **Fentanyl**, Lozenges 1600 micrograms (as citrate), 3 (Actiq)
 5425M **Methylnaltrexone**, Solution for injection containing methylnaltrexone bromide 12 mg in 0.6 mL, (*Relistor*)

Deletion – Brands

- 8559L Gabahexal 600mg, SZ – **Gabapentin**, Tablet 600 mg
 8272J Klacid Hp 7, AB – **Omeprazole and Clarithromycin and Amoxycillin**, Pack containing 14 capsules omeprazole 20 mg, 14 tablets clarithromycin 500 mg and 28 capsules amoxycillin 500 mg
 1977C Ranihexal, HX – **Ranitidine Hydrochloride**, Tablet 300 mg (base)

Deletion – Bioequivalence Indicators

- 8272J Probitor Hp 7, SZ – **Omeprazole and Clarithromycin and Amoxycillin**, Pack containing 14 capsules omeprazole 20 mg, 14 tablets clarithromycin 500 mg and 28 capsules amoxycillin 500 mg

Deletion – Note

- 5385K **Benzydamine Hydrochloride**, Mouth and throat rinse 22.5 mg per 15 mL, 500 mL, (*Difflam*) (**Palliative Care**)
 5301B **Bisacodyl**, Tablet 5 mg, (*Bisalax, Lax-Tab*) (**Palliative Care**)
 5304E **Bisacodyl**, Suppositories 10 mg, 12, (*Petrus Bisacodyl Suppositories*) (**Palliative Care**)
 5303D **Bisacodyl**, Suppositories 10 mg, 10, (*Dulcolax, Petrus Bisacodyl Suppositories*) (**Palliative Care**)
 5302C **Bisacodyl**, Enemas 10 mg in 5 mL, 25, (*Bisalax*) (**Palliative Care**)
 5333Q **Carmellose Sodium**, Mouth spray 10 mg per mL, 25 mL, (*Aquae*) (**Palliative Care**)
 5334R **Carmellose Sodium**, Mouth spray 10 mg per mL, 100 mL, (*Aquae*) (**Palliative Care**)
 5362F **Diclofenac Sodium**, Tablet 50 mg (enteric coated), (*APO-Diclofenac, Chem mart Diclofenac, Clonac 50, Diclofenac-GA, Diclofenac Sandoz, Fenac, Terry White Chemists Diclofenac, Voltaren 50*) (**Palliative Care**)
 5361E **Diclofenac Sodium**, Tablet 25 mg (enteric coated), (*APO-Diclofenac, Chem mart Diclofenac, Clonac 25, Diclofenac-GA, Diclofenac Sandoz, Fenac 25, Terry White Chemists Diclofenac, Voltaren 25*) (**Palliative Care**)
 5363G **Diclofenac Sodium**, Suppository 100 mg, (*Voltaren 100*) (**Palliative Care**)
 5311M **Glycerol**, Suppositories 700 mg (for infants), 12, (*Petrus Pharmaceuticals Pty Ltd*) (**Palliative Care**)
 5313P **Glycerol**, Suppositories 2.8 g (for adults), 12, (*Petrus Pharmaceuticals Pty Ltd*) (**Palliative Care**)
 5312N **Glycerol**, Suppositories 1.4 g (for children), 12, (*Petrus Pharmaceuticals Pty Ltd*) (**Palliative Care**)
 5317W **Hyoscine Butylbromide**, Injection 20 mg in 1 mL, (*Buscopan*) (**Palliative Care**)
 5421H **Hypromellose**, Oral gel 20 mg per g, 100 g, (*Aquae Gel*) (**Palliative Care**)
 5368M **Ibuprofen**, Tablet 400 mg, (*Brufen*) (**Palliative Care**)
 5377B **Indomethacin**, Capsule 25 mg, (*Arthrexin, Indocid*) (**Palliative Care**)
 5378C **Indomethacin**, Suppository 100 mg, (*Indocid*) (**Palliative Care**)
 5387M **Lactulose**, Mixture 3.34 g per 5 mL, 500 mL, (*Actilax, Duphalac, Genlac, GenRx Lactulose, Lac-Dol, Lactocur*) (**Palliative Care**)
 5389P **Macrogol 3350**, Sachets containing powder for solution 13.125 g with electrolytes, 30, (*Movicol*) (**Palliative Care**)
 5426N **Macrogol 3350**, Powder for oral solution 510 g, (*MediHealth ClearLax, OsmoLax*) (**Palliative Care**)

5347K **Naproxen**, Tablet 750 mg (sustained release), (*Naprosyn SR750, Proxen SR 750*) **(Palliative Care)**

5346J **Naproxen**, Tablet 500 mg, (*Inza 500, Naprosyn*) **(Palliative Care)**

5345H **Naproxen**, Tablet 250 mg, (*Inza 250, Naprosyn*) **(Palliative Care)**

5348L **Naproxen**, Tablet 1 g (sustained release), (*Naprosyn SR1000, Proxen SR 1000*) **(Palliative Care)**

5397C **Naproxen**, Oral suspension 125 mg per 5 mL, 474 mL, (*Naprosyn*) **(Palliative Care)**

5353R **Naproxen Sodium**, Tablet 550 mg, (*Anaprox 550, Crysanal*) **(Palliative Care)**

5343F **Paracetamol**, Tablet 665 mg (modified release), (*Panadol Osteo*) **(Palliative Care)**

5319Y **Paracetamol**, Suppositories 500 mg, 24, (*Panadol*) **(Palliative Care)**

5326H **Promethazine Hydrochloride**, Tablet 25 mg, (*Phenergan*) **(Palliative Care)**

5325G **Promethazine Hydrochloride**, Tablet 10 mg, (*Phenergan*) **(Palliative Care)**

5327J **Promethazine Hydrochloride**, Oral liquid 5 mg per 5 mL, 100 mL, (*Phenergan*) **(Palliative Care)**

5331N **Sorbitol With Sodium Citrate And Sodium Lauryl Sulfoacetate**, Enemas 3.125 g-450 mg-45 mg in 5 mL, 12, (*Micolette, Microlax*) **(Palliative Care)**

5322D **Sterculia With Frangula Bark**, Granules 620 mg-80 mg per g (62%-8%), 500 g, (*Normacol Plus*) **(Palliative Care)**

5382G **Sulindac**, Tablet 200 mg, (*Acclin 200*) **(Palliative Care)**

5381F **Sulindac**, Tablet 100 mg, (*Acclin*) **(Palliative Care)**

ALTERATIONS

Alterations — Proprietary Name

From:

2688L **Azathioprine**, Tablet 25 mg, (*Azahexal*)

To:

2688L **Azathioprine**, Tablet 25 mg, (*Azathioprine Sandoz*)

From:

2413B **Frusemide**, Injection 20 mg in 2 mL, (*Frusehexal*)

To:

2413B **Frusemide**, Injection 20 mg in 2 mL, (*Frusemide Sandoz*)

From:

3466K **Frusemide**, Injection 20 mg in 2 mL, (*Frusehexal*) **(Emergency Drug Supply)**

To:

3466K **Frusemide**, Injection 20 mg in 2 mL, (*Frusemide Sandoz*) **(Emergency Drug Supply)**

From:

8524P **Tramadol hydrochloride**, Tablet 150 mg (twice daily sustained release), (*Tramahexal SR*)

To:

8524P **Tramadol hydrochloride**, Tablet 150 mg (twice daily sustained release), (*Tramadol Sandoz SR*)

From:

5235M **Tramadol hydrochloride**, Tablet 150 mg (twice daily sustained release), (*Tramahexal SR*) **(Dental)**

To:

5235M **Tramadol hydrochloride**, Tablet 150 mg (twice daily sustained release), (*Tramadol Sandoz SR*) **(Dental)**

Alterations — Item Description

From:

1370D **Enalapril Maleate**, Tablet 5 mg (base), (*Alphapril, Auspril, Chem mart Enalapril, Enahexal, Enalabell, Enalapril-DP 5mg, Enalapril-GA, Enalapril generichealth, Enalapril Sandoz, Enalapril Winthrop, GenRx Enalapril, Renitec M, Terry White Chemists Enalapril*)

To:

1370D **Enalapril**, Tablet containing enalapril maleate 5 mg, (*Alphapril, Auspril, Chem mart Enalapril, Enahexal, Enalabell, Enalapril-DP 5mg, Enalapril-GA, Enalapril generichealth, Enalapril Sandoz, Enalapril Winthrop, GenRx Enalapril, Renitec M, Terry White Chemists Enalapril*)

From:

1368B **Enalapril Maleate**, Tablet 10 mg (base), (*Alphapril, Auspril, Chem mart Enalapril, Enahexal, Enalabell, Enalapril-DP 10mg, Enalapril-GA, Enalapril generichealth, Enalapril Sandoz, Enalapril Winthrop, GenRx Enalapril, Renitec,*

Terry White Chemists Enalapril)

To:

1368B **Enalapril**, Tablet containing enalapril maleate 10 mg, (*Alphapril, Auspril, Chem mart Enalapril, Enahexal, Enalabell, Enalapril-DP 10mg, Enalapril-GA, Enalapril generichealth, Enalapril Sandoz, Enalapril Winthrop, GenRx Enalapril, Renitec, Terry White Chemists Enalapril*)

From:

1369C **Enalapril Maleate**, Tablet 20 mg (base), (*Alphapril, Auspril, Chem mart Enalapril, Enahexal, Enalabell, Enalapril-DP 20mg, Enalapril-GA, Enalapril generichealth, Enalapril Sandoz, GenRx Enalapril, Renitec 20, Terry White Chemists Enalapril*)

To:

1369C **Enalapril**, Tablet containing enalapril maleate 20 mg, (*Alphapril, Auspril, Chem mart Enalapril, Enahexal, Enalabell, Enalapril-DP 20mg, Enalapril-GA, Enalapril generichealth, Enalapril Sandoz, GenRx Enalapril, Renitec 20, Terry White Chemists Enalapril*)

From:

8050Q **Gemcitabine Hydrochloride**, Powder for I.V. infusion 1 g (base), (*DBL Gemcitabine for Injection, Gemcitabine Actavis, Gemcitabine Ebewe, Gemcitabine Kabi, Gemcitabine Sun, Gemcite, Gemzar*)

To:

8050Q **Gemcitabine**, Powder for I.V. infusion 1 g (as hydrochloride), (*DBL Gemcitabine for Injection, Gemcitabine Actavis, Gemcitabine Ebewe, Gemcitabine Kabi, Gemcitabine Sun, Gemcite, Gemzar*)

From:

9414L **Gemcitabine Hydrochloride**, Powder for I.V. infusion 2 g (base), (*DBL Gemcitabine for Injection, Gemcitabine Kabi*)

To:

9414L **Gemcitabine**, Powder for I.V. infusion 2 g (as hydrochloride), (*DBL Gemcitabine for Injection, Gemcitabine Kabi*)

From:

8049P **Gemcitabine Hydrochloride**, Powder for I.V. infusion 200 mg (base), (*DBL Gemcitabine for Injection, Gemcitabine Actavis, Gemcitabine Ebewe, Gemcitabine Kabi, Gemcitabine Sun, Gemcite, Gemzar*)

To:

8049P **Gemcitabine**, Powder for I.V. infusion 200 mg (as hydrochloride), (*DBL Gemcitabine for Injection, Gemcitabine Actavis, Gemcitabine Ebewe, Gemcitabine Kabi, Gemcitabine Sun, Gemcite, Gemzar*)

From:

9402W **Gemcitabine Hydrochloride**, Solution concentrate for I.V. infusion 1000 mg (base) in 100 mL, (*Gemcitabine Ebewe*)

To:

9402W **Gemcitabine**, Solution concentrate for I.V. infusion 1000 mg (as hydrochloride) in 100 mL, (*Gemcitabine Ebewe*)

From:

9401T **Gemcitabine Hydrochloride**, Solution concentrate for I.V. infusion 200 mg (base) in 20 mL, (*Gemcitabine Ebewe*)

To:

9401T **Gemcitabine**, Solution concentrate for I.V. infusion 200 mg (as hydrochloride) in 20 mL, (*Gemcitabine Ebewe*)

From:

9463C **Gemcitabine Hydrochloride**, Solution concentrate for I.V. infusion 500 mg (base) in 50 mL, (*Gemcitabine Ebewe*)

To:

9463C **Gemcitabine**, Solution concentrate for I.V. infusion 500 mg (as hydrochloride) in 50 mL, (*Gemcitabine Ebewe*)

Alterations — Max Quantity

		From	To
5318X	Hyoscine Butylbromide , Injection 20 mg in 1 mL, (<i>Buscopan</i>) (Palliative Care)	5	30
5317W	Hyoscine Butylbromide , Injection 20 mg in 1 mL, (<i>Buscopan</i>) (Palliative Care)	5	30
5388N	Lactulose , Mixture 3.34 g per 5 mL, 500 mL, (<i>Actilax, Duphalac, Genlac, GenRx Lactulose, Lac-Dol, Lactocur</i>) (Palliative Care)	1	3
5387M	Lactulose , Mixture 3.34 g per 5 mL, 500 mL, (<i>Actilax, Duphalac, Genlac, GenRx Lactulose, Lac-Dol, Lactocur</i>) (Palliative Care)	1	3
5390Q	Macrogol 3350 , Sachets containing powder for solution 13.125 g with electrolytes, 30, (<i>Movicol</i>) (Palliative Care)	1	2
5389P	Macrogol 3350 , Sachets containing powder for solution 13.125 g with electrolytes, 30, (<i>Movicol</i>) (Palliative Care)	1	2
5427P	Macrogol 3350 , Powder for oral solution 510 g, (<i>MediHealth ClearLax, OsmoLax</i>) (Palliative Care)	1	2

5426N	Macrogol 3350 , Powder for oral solution 510 g, (<i>MediHealth ClearLax, OsmoLax</i>) (Palliative Care)	1	2
5320B	Paracetamol , Suppositories 500 mg, 24, (<i>Panadol</i>) (Palliative Care)	1	4
5319Y	Paracetamol , Suppositories 500 mg, 24, (<i>Panadol</i>) (Palliative Care)	1	4

Alterations — Number of Repeats

		From	To
5412W	Fentanyl , Lozenges 1600 micrograms (as citrate), 3, (<i>Actiq</i>)	2	0
5411T	Fentanyl , Lozenges 1200 micrograms (as citrate), 3, (<i>Actiq</i>)	2	0
5410R	Fentanyl , Lozenges 800 micrograms (as citrate), 3, (<i>Actiq</i>)	2	0
5409Q	Fentanyl , Lozenges 600 micrograms (as citrate), 3, (<i>Actiq</i>)	2	0
5408P	Fentanyl , Lozenges 400 micrograms (as citrate), 3, (<i>Actiq</i>)	2	0
5407N	Fentanyl , Lozenges 200 micrograms (as citrate), 3, (<i>Actiq</i>)	2	0
5424L	Methylnaltrexone , Solution for injection containing methylnaltrexone bromide 12 mg in 0.6 mL, (<i>Relistor</i>)	3	0

Alterations — Manufacturer's Codes

		From	To
9297H	Glucose Indicator—Blood , Test strips, 50 (<i>Bionime Rightest</i>)	CQ	QB
9298J	Glucose Indicator—Blood , Test strips, 50 (<i>Bionime Rightest</i>) (Diff. Max. Rpts.)	CQ	QB
3439B	Metformin Hydrochloride , Tablet 1 g (extended release), (<i>Diabex XR 1000</i>)	AF	AL

Alterations — Restrictions and Notes

Amendments have been made to restrictions and notes for all Palliative Care items effective 1 January 2011. Please see the Palliative Care listings in the PBS Schedule for amended restrictions and note text. The Schedule is also available on the internet at www.pbs.gov.au.

The following are now “Authority required (STREAMLINED)” items effective 1 January 2011:

1007B	Aciclovir , Tablet 200 mg, (<i>Aciclovir 200, Acihexal, Acyclo-V 200, Chem mart Aciclovir, GenRx Aciclovir, Lovir, Ozvir, Terry White Chemists Aciclovir, Zovirax 200 mg</i>)
1003T	Aciclovir , Tablet 200 mg, (<i>Acihexal, Acyclo-V 200, GenRx Aciclovir, Lovir, Zovirax 200 mg</i>)
1052J	Aciclovir , Tablet 800 mg, (<i>Aciclovir 800, Acihexal, Acyclo-V 800, GenRx Aciclovir, Zovirax 800 mg</i>)
8234J	Aciclovir , Tablet 800 mg, (<i>Acihexal, Acyclo-V 800</i>)
8808N	Aprepitant , Pack containing 1 capsule 125 mg and 2 capsules 80 mg, (<i>Emend</i>)
8897G	Famciclovir , Tablet 500 mg, (<i>Famvir, Favic 500</i>)
8896F	Famciclovir , Tablet 500 mg, (<i>Ezovir, Famvir, Favic 500</i>)
2274Q	Famciclovir , Tablet 250 mg, (<i>APO-Famciclovir, Ezovir, Famciclovir Sandoz, Famvir, Favic 250</i>)
8217L	Famciclovir , Tablet 250 mg, (<i>APO-Famciclovir, Ezovir, Famciclovir Sandoz, Famvir, Favic 250</i>)
8002E	Famciclovir , Tablet 250 mg, (<i>APO-Famciclovir, Ezovir, Famciclovir Sandoz, Famvir, Favic 250</i>)
8092X	Famciclovir , Tablet 125 mg, (<i>APO-Famciclovir, Ezovir, Famvir, Favic 125</i>)
1757L	Fluconazole , Solution for I.V. infusion 400 mg in 200 mL, (<i>Baxter Healthcare Pty Ltd</i>)
1474N	Fluconazole , Solution for I.V. infusion 200 mg in 100 mL, (<i>Baxter Healthcare Pty Ltd, Diflucan, Fluconazole-Clarix, Fluconazole Hexal, Fluconazole Sandoz</i>)
1473M	Fluconazole , Solution for I.V. infusion 100 mg in 50 mL, (<i>Diflucan, Fluconazole-Clarix, Fluconazole Hexal</i>)
1471K	Fluconazole , Capsule 50 mg, (<i>DBL Fluconazole, Diflucan, Dizole 50, Fluconazole Sandoz, Fluzole 50, Ozole</i>)
1475P	Fluconazole , Capsule 200 mg, (<i>APO-Fluconazole, DBL Fluconazole, Diflucan, Dizole 200, Fluconazole Sandoz, Fluzole 200, Ozole</i>)
1472L	Fluconazole , Capsule 100 mg, (<i>DBL Fluconazole, Diflucan, Dizole 100, Fluconazole Sandoz, Fluconazole Winthrop, Ozole</i>)
8015W	Gestrinone , Capsule 2.5 mg, (<i>Dimetriose</i>)
8873B	Granisetron Hydrochloride , Tablet 2 mg (base), (<i>Kytril</i>)
8730L	Granisetron Hydrochloride , Concentrated injection 3 mg (base) in 3 mL, (<i>Granisetron Kabi, Kytril</i>)
8196J	Itraconazole , Capsule 100 mg, (<i>Sporanox</i>)
1573T	Ketoconazole , Tablet 200 mg, (<i>Nizoral</i>)
1572R	Ketoconazole , Tablet 200 mg, (<i>Nizoral</i>)
8413T	Ondansetron , Wafer 8 mg, (<i>Ondaz Zydix, Zofran Zydix</i>)
8412R	Ondansetron , Wafer 4 mg, (<i>Ondaz Zydix, Zofran Zydix</i>)
1595Y	Ondansetron , Tablet 8 mg, (<i>APO-Ondansetron, Ondaz, Onsetron 8, Zofran</i>)
1594X	Ondansetron , Tablet 4 mg, (<i>APO-Ondansetron, Ondaz, Onsetron 4, Zofran</i>)
8233H	Ondansetron , Syrup 4 mg per 5 mL, 50 mL, (<i>Zofran syrup 50 mL</i>)
1597C	Ondansetron , I.V. injection 8 mg in 4 mL, (<i>Ondansetron-Clarix, Ondaz, Onsetron, Pfizer Australia Pty Ltd, Zofran</i>)

1596B	Ondansetron , I.V. injection 4 mg in 2 mL, (<i>Ondansetron-Claris, Ondaz, Onsetron, Pfizer Australia Pty Ltd, Zofran</i>)
8134D	Valaciclovir Hydrochloride , Tablet 500 mg (base), (<i>Valtrex</i>)
8133C	Valaciclovir Hydrochloride , Tablet 500 mg (base), (<i>Valtrex</i>)
8064K	Valaciclovir Hydrochloride , Tablet 500 mg (base), (<i>Valtrex</i>)

SECTION 100 — HIGHLY SPECIALISED DRUGS PROGRAM ALTERATIONS

Alterations — Proprietary Name and Item Description

From:

5603X **Abacavir sulfate with Lamivudine**, Tablet 600 mg (base)- 300 mg, (*Kivexa*) **(Public)**

To:

5603X **Abacavir with Lamivudine**, Tablet containing abacavir 600 mg (as sulfate) with lamivudine 300 mg, (*Kivexa*) **(Public)**

From:

6458X **Abacavir sulfate with Lamivudine**, Tablet 600 mg (base)- 300 mg, (*Kivexa*) **(Private)**

To:

6458X **Abacavir with Lamivudine**, Tablet containing abacavir 600 mg (as sulfate) with lamivudine 300 mg, (*Kivexa*) **(Private)**

From:

5604Y **Abacavir sulfate with Lamivudine and Zidovudine**, Tablet 300 mg (base)- 150 mg-300 mg, (*Trizivir*) **(Public)**

To:

5604Y **Abacavir with Lamivudine and Zidovudine**, Tablet containing abacavir 300 mg (as sulfate) with lamivudine 150 mg and zidovudine 300 mg, (*Trizivir*) **(Public)**

From:

6327B **Abacavir sulfate with Lamivudine and Zidovudine**, Tablet 300 mg (base)- 150 mg-300 mg, (*Trizivir*) **(Private)**

To:

6327B **Abacavir with Lamivudine and Zidovudine**, Tablet containing abacavir 300 mg (as sulfate) with lamivudine 150 mg and zidovudine 300 mg, (*Trizivir*) **(Private)**

Alterations — Restrictions

5601T **Abacavir**, Tablet 300 mg (as sulfate), (*Ziagen*) **(Public)**

6264Q **Abacavir**, Tablet 300 mg (as sulfate), (*Ziagen*) **(Private)**

5602W **Abacavir**, Oral solution 20 mg (as sulfate) per mL, 240 mL, (*Ziagen*) **(Public)**

6265R **Abacavir**, Oral solution 20 mg (as sulfate) per mL, 240 mL, (*Ziagen*) **(Private)**

5604Y **Abacavir Sulfate With Lamivudine And Zidovudine**, Tablet containing abacavir 300 mg (as sulfate) with lamivudine 150 mg and zidovudine 300 mg, (*Trizivir*) **(Public)**

6327B **Abacavir Sulfate With Lamivudine And Zidovudine**, Tablet containing abacavir 300 mg (as sulfate) with lamivudine 150 mg and zidovudine 300 mg, (*Trizivir*) **(Private)**

5603X **Abacavir With Lamivudine**, Tablet containing abacavir 600 mg (as sulfate) with lamivudine 300 mg, (*Kivexa*) **(Public)**

6458X **Abacavir With Lamivudine**, Tablet containing abacavir 600 mg (as sulfate) with lamivudine 300 mg, (*Kivexa*) **(Private)**

5615M **Atazanavir**, Capsule 100 mg (as sulfate), (*Reyataz*) **(Public)**

9646Q **Atazanavir**, Capsule 100 mg (as sulfate), (*Reyataz*) **(Private)**

5613K **Atazanavir**, Capsule 150 mg (as sulfate), (*Reyataz*) **(Public)**

6451M **Atazanavir**, Capsule 150 mg (as sulfate), (*Reyataz*) **(Private)**

5614L **Atazanavir**, Capsule 200 mg (as sulfate), (*Reyataz*) **(Public)**

6452N **Atazanavir**, Capsule 200 mg (as sulfate), (*Reyataz*) **(Private)**

5612J **Atazanavir**, Capsule 300 mg (as sulfate), (*Reyataz*) **(Public)**

9614B **Atazanavir**, Capsule 300 mg (as sulfate), (*Reyataz*) **(Private)**

5653M **Darunavir**, Tablet 150 mg (as ethanolate), (*Prezista*) **(Public)**

9581G **Darunavir**, Tablet 150 mg (as ethanolate), (*Prezista*) **(Private)**

5652L **Darunavir**, Tablet 300 mg (as ethanolate), (*Prezista*) **(Public)**

9616D **Darunavir**, Tablet 300 mg (as ethanolate), (*Prezista*) **(Private)**

5663C **Didanosine**, Capsule 125 mg (containing enteric coated beadlets), (*Videx EC*) **(Public)**

6298L **Didanosine**, Capsule 125 mg (containing enteric coated beadlets), (*Videx EC*) **(Private)**

5664D **Didanosine**, Capsule 200 mg (containing enteric coated beadlets), (*Videx EC*) **(Public)**

6299M **Didanosine**, Capsule 200 mg (containing enteric coated beadlets), (*Videx EC*) **(Private)**

5665E **Didanosine**, Capsule 250 mg (containing enteric coated beadlets), (*Videx EC*) **(Public)**

6300N **Didanosine**, Capsule 250 mg (containing enteric coated beadlets), (*Videx EC*) **(Private)**

5666F **Didanosine**, Capsule 400 mg (containing enteric coated beadlets), (*Videx EC*) **(Public)**

6301P **Didanosine**, Capsule 400 mg (containing enteric coated beadlets), (*Videx EC*) **(Private)**

5708K **Efavirenz**, Tablet 200 mg, (*Stocrin*) **(Public)**

9618F **Efavirenz**, Tablet 200 mg, (*Stocrin*) **(Private)**

5706H **Efavirenz**, Tablet 600 mg, (*Stocrin*) **(Public)**

6356M **Efavirenz**, Tablet 600 mg, (*Stocrin*) **(Private)**

5707J **Efavirenz**, Oral solution 30 mg per mL, 180 mL, (*Stocrin*) **(Public)**

6372J **Efavirenz**, Oral solution 30 mg per mL, 180 mL, (*Stocrin*) **(Private)**

5709L **Emtricitabine**, Capsule 200 mg, (*Emtriva*) **(Public)**

6137B **Emtricitabine**, Capsule 200 mg, (*Emtriva*) **(Private)**

5710M **Enfuvirtide**, Pack containing 60 vials powder for injection 90 mg with 60 vials water for injections 1.1 mL (with syringes and swabs), (*Fuzeon*) **(Public)**

6455R **Enfuvirtide**, Pack containing 60 vials powder for injection 90 mg with 60 vials water for injections 1.1 mL (with syringes and swabs), (*Fuzeon*) **(Private)**

5736X **Etravirine**, Tablet 100 mg, (*Intelence*) **(Public)**

9639H **Etravirine**, Tablet 100 mg, (*Intelence*) **(Private)**

5746K **Fosamprenavir**, Tablet 700 mg (as calcium), (*Telzir*) **(Public)**

6453P **Fosamprenavir**, Tablet 700 mg (as calcium), (*Telzir*) **(Private)**

5745J **Fosamprenavir**, Oral liquid 50 mg (as calcium) per mL, 225 mL, (*Telzir*) **(Public)**

6454Q **Fosamprenavir**, Oral liquid 50 mg (as calcium) per mL, 225 mL, (*Telzir*) **(Private)**

5752R **Indinavir**, Capsule 400 mg (as sulfate), (*Crixivan 400 mg*) **(Public)**

6202K **Indinavir**, Capsule 400 mg (as sulfate), (*Crixivan 400 mg*) **(Private)**

5755X **Infliximab**, Powder for I.V. infusion 100 mg, (*Remicade*) **(Public)**

9612X **Infliximab**, Powder for I.V. infusion 100 mg, (*Remicade*) **(Private)**

5772T **Lamivudine**, Tablet 150 mg, (*3TC*) **(Public)**

6193Y **Lamivudine**, Tablet 150 mg, (*3TC*) **(Private)**

5774X **Lamivudine**, Tablet 300 mg, (*3TC*) **(Public)**

6435Q **Lamivudine**, Tablet 300 mg, (*3TC*) **(Private)**

5773W **Lamivudine**, Oral solution 10 mg per mL, 240 mL, (*3TC*) **(Public)**

6194B **Lamivudine**, Oral solution 10 mg per mL, 240 mL, (*3TC*) **(Private)**

5775Y **Lamivudine With Zidovudine**, Tablet 150 mg-300 mg, (*Combivir*) **(Public)**

6234D **Lamivudine With Zidovudine**, Tablet 150 mg-300 mg, (*Combivir*) **(Private)**

5790R **Lopinavir With Ritonavir**, Tablet 100 mg-25 mg, (*Kaletra*) **(Public)**

9633B **Lopinavir With Ritonavir**, Tablet 100 mg-25 mg, (*Kaletra*) **(Private)**

5791T **Lopinavir With Ritonavir**, Tablet 200 mg-50 mg, (*Kaletra*) **(Public)**

6495W **Lopinavir With Ritonavir**, Tablet 200 mg-50 mg, (*Kaletra*) **(Private)**

5789Q **Lopinavir With Ritonavir**, Oral liquid 400 mg-100 mg per 5 mL, 60 mL, (*Kaletra*) **(Public)**

6341R **Lopinavir With Ritonavir**, Oral liquid 400 mg-100 mg per 5 mL, 60 mL, (*Kaletra*)

5792W **Maraviroc**, Tablet 150 mg, (*Celsentri*) **(Public)**

9572T **Maraviroc**, Tablet 150 mg, (*Celsentri*) **(Private)**

5793X **Maraviroc**, Tablet 300 mg, (*Celsentri*) **(Public)**

9573W **Maraviroc**, Tablet 300 mg, (*Celsentri*) **(Private)**

9506H **Nevirapine**, Tablet 200 mg, (*Viramune*) **(Public)**

6215D **Nevirapine**, Tablet 200 mg, (*Viramune*) **(Private)**

9507J **Nevirapine**, Oral suspension 50 mg (as hemihydrate) per 5 mL, 240 mL, (*Viramune*) **(Public)**

9571R **Nevirapine**, Oral suspension 50 mg (as hemihydrate) per 5 mL, 240 mL, (*Viramune*) **(Private)**

9523F **Raltegravir**, Tablet 400 mg (as potassium), (*Isentress*) **(Public)**

9629T **Raltegravir**, Tablet 400 mg (as potassium), (*Isentress*) **(Private)**

9660K **Ritonavir**, Tablet 100 mg, (*Norvir*) **(Public)**

9677H **Ritonavir**, Tablet 100 mg, (*Norvir*) **(Private)**

9542F **Ritonavir**, Oral solution 600 mg per 7.5 mL (80 mg per mL), 90 mL, (*Norvir*) **(Public)**

6494T **Ritonavir**, Oral solution 600 mg per 7.5 mL (80 mg per mL), 90 mL, (*Norvir*) **(Private)**

9545J **Saquinavir**, Tablet 500 mg (as mesylate), (*Invirase*) **(Public)**

6498B **Saquinavir**, Tablet 500 mg (as mesylate), (*Invirase*) **(Private)**

9552R **Stavudine**, Powder for oral solution 1 mg per mL, 200 mL, (*Zerit*) **(Public)**

6250Y **Stavudine**, Powder for oral solution 1 mg per mL, 200 mL, (*Zerit*) **(Private)**

9553T **Stavudine**, Capsule 20 mg, (*Zerit*) **(Public)**

6186N **Stavudine**, Capsule 20 mg, (*Zerit*) **(Private)**

9554W **Stavudine**, Capsule 30 mg, (*Zerit*) **(Public)**

6189R **Stavudine**, Capsule 30 mg, (*Zerit*) **(Private)**

9556Y	Stavudine , Capsule 40 mg, (<i>Zerit</i>) (Public)
6190T	Stavudine , Capsule 40 mg, (<i>Zerit</i>) (Private)
9563H	Tenofovir , Tablet containing tenofovir disoproxil fumarate 300 mg, (<i>Viread</i>) (Public)
6358P	Tenofovir , Tablet containing tenofovir disoproxil fumarate 300 mg, (<i>Viread</i>) (Private)
9564J	Tenofovir With Emtricitabine , Tablet containing tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg, (<i>Truvada</i>) (Public)
6468K	Tenofovir With Emtricitabine , Tablet containing tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg, (<i>Truvada</i>) (Private)
9565K	Tenofovir With Emtricitabine And Efavirenz , Tablet containing tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg and efavirenz 600 mg, (<i>Atripla</i>) (Public)
9650X	Tenofovir With Emtricitabine And Efavirenz , Tablet containing tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg and efavirenz 600 mg, (<i>Atripla</i>) (Private)
9656F	Tipranavir , Oral liquid 100 mg per mL, 95 mL, (<i>Aptivus</i>) (Public)
9676G	Tipranavir , Oral liquid 100 mg per mL, 95 mL, (<i>Aptivus</i>) (Private)
9567M	Tipranavir , Capsule 250 mg, (<i>Aptivus</i>) (Public)
9610T	Tipranavir , Capsule 250 mg, (<i>Aptivus</i>) (Private)
9570Q	Zidovudine , Syrup 10 mg per mL, 200 mL, (<i>Retrovir</i>) (Public)
6155Y	Zidovudine , Syrup 10 mg per mL, 200 mL, (<i>Retrovir</i>) (Private)
9651Y	Zidovudine , Capsule 100 mg, (<i>Retrovir</i>) (Public)
6153W	Zidovudine , Capsule 100 mg, (<i>Retrovir</i>) (Private)
9652B	Zidovudine , Capsule 250 mg, (<i>Retrovir</i>) (Public)
6154X	Zidovudine , Capsule 250 mg, (<i>Retrovir</i>) (Private)

ADVANCE NOTICES DELETIONS

Advanced Notice – Deletion of Items

Contrary to previous advice this item will remain as a pharmaceutical benefit after 1 January 2011 and will be deleted on 1 April 2011.

1350C **Dydrogesterone**, Tablet 10 mg, (*Duphaston*)

The following Section 100 items will be deleted from the Schedule of Pharmaceutical Benefits on 1 February 2011:

Items discontinued by the manufacturer—

9552P **Stavudine**, Powder for oral solution 1 mg per mL, 200 mL (*Zerit*) **(Public)**

6250Y **Stavudine**, Powder for oral solution 1 mg per mL, 200 mL (*Zerit*) **(Private)**

The following items will be deleted from the Schedule of Pharmaceutical Benefits on 1 April 2011:

Items discontinued by the manufacturer—

1350C **Dydrogesterone**, Tablet 10 mg, (*Duphaston*)

2772X **Norethisterone with ethinylloestradiol**, Tablets 500 micrograms-35 micrograms, 21, (*Brevinor*)

2773Y **Norethisterone with ethinylloestradiol**, Tablets 1 micrograms-35 micrograms, 21, (*Brevinor*)

8556H **Pancreatic Extract**, Capsule (containing enteric coated minimicrospheres) providing not less than 5,000 BP units of lipase activity, (*Creon 5000*)

Advance Notice — Deletion of Brands

The following brands will be deleted from the Schedule of Pharmaceutical Benefits on 1 February 2011:

Brands discontinued by the manufacturer—

1370D *Enahexal, HX* — **Enalapril maleate**, Tablet 5 mg

1368B *Enahexal, HX* — **Enalapril maleate**, Tablet 10 mg

1369C *Enahexal, HX* — **Enalapril maleate**, Tablet 20 mg

2487X *Famohexal, HX* — **Famotidine**, Tablet 20 mg

The following brands will be deleted from the Schedule of Pharmaceutical Benefits on 1 March 2011:

Brands discontinued by the manufacturer—

2315W *Blenoxane, BQ* — **Bleomycin sulfate**, Powder for injection 15,000 i.u. (solvent required) **(Special Pharmaceutical Benefit)**

1978D *Ranixhexal, SZ* — **Ranitidine hydrochloride**, Tablet 150 mg (base)

Addresses — Medicare Australia

Medicare Australia has responsibility for the operational aspects of the Pharmaceutical Benefits Scheme (PBS). This responsibility covers the processing of pharmaceutical benefit and safety net claims, authority applications and supply of PBS stationery used by medical practitioners, participating dental practitioners and approved pharmacists.

Procedures for ordering prescription forms are set out in Introduction of this Schedule .

New South Wales and Australian Capital Territory

Pharmaceutical Benefits Branch
130 George Street
Parramatta NSW 2150

General and IME enquiries — Tel: 132 290

Orange Service Centre
189 Anson Street
Orange NSW 2800

General and IME enquiries — Tel: 132 290

Victoria

Pharmaceutical Branch
Level 10
595 Collins Street
Melbourne Vic 3000

General and IME enquiries — Tel: 132 290

Queensland

Pharmaceutical Services Branch
143 Turbot Street
Brisbane Qld 4000

General and IME enquiries — Tel: 132 290

South Australia and Northern Territory

Pharmaceutical Services Branch
209 Greenhill Road
Eastwood SA 5063

General and IME enquiries — Tel: 132 290

Western Australia

Pharmaceutical Benefits Branch
Level 5, Work Distribution Centre, (Reception on Level 4)
130 Stirling Street
Northbridge WA 6003

General and IME enquiries — Tel: 132 290

Tasmania

Pharmaceutical Branch
242 Liverpool Street
Hobart Tas 7000

General and IME enquiries — Tel: 132 290

National Program Management

Pharmaceutical Benefits Branch
Medicare Australia
134 Reed Street
Tuggeranong ACT 2900
Telephone — (02) 6124 6333
Website — www.medicareaustralia.gov.au Email — pbs@medicareaustralia.gov.au

Authority Prescription Applications

Authority required benefits fall into two categories – *Authority required* and *Authority required (STREAMLINED)*. The process in which an authority PBS prescription can be prescribed will depend on the type of Authority required benefit.

Prior approval is required for Authority required items as well as all requests for increased quantities and/or repeats for any category of PBS item.

Prior approval is not required for Authority required (STREAMLINED) items except if increased quantities and/or repeats are required (see Explanatory Notes for details).

Mail Applications: REPLY PAID No. 9857
PBS Authorities Section
Medicare Australia
GPO Box 9857
In your Capital City

Telephone Applications: Free call 1800 888 333
Australia-wide 24 hour service PBS Authorities Section

For telephone applications please have the following information available:

Patient: Medicare Number
Surname
First name
Full residential address (including post code)

PBS Authority Prescription Number: Top right hand side of the handwritten PBS Authority Form

Your Prescriber Number: Located below your address block on the personalised forms

Drug Information: PBS item
Quantity required and number of repeats
Daily dose
Disease or purpose information

Requests for Drugs via the Special Access Scheme (SAS)

Requests for individual patient approval to obtain drugs that are available only through the SAS may be directed to a delegate within the Drug Safety and Evaluation Branch, Therapeutic Goods Administration, telephone (02) 6232 8111, facsimile (02) 6232 8112, or by mail to PO Box 100 Woden ACT 2606.

Department of Veterans' Affairs

Details of the approving authority for the Department of Veterans' Affairs are listed at the front of the Repatriation Schedule of Pharmaceutical Benefits.

Telephone Interpreter Service

A 24-hour, seven days a week telephone service is available by contacting 131 450.

The translating service (TIS) can provide immediate assistance over the telephone or arrange for an interpreter to go to a location specified in either city or country areas. The TIS service has access to 2000 professional interpreters, covering over 100 languages and dialects.

Poisons Information Centres

Phone 131 126 from anywhere in Australia — 24 hours — form information and advice on the treatment of poisoning, bites and stings

NSW

The New Children's Hospital
Hawkesbury Road
Westmead NSW 2148
Tel: (02) 9845 3111

VIC

Austin Hospital
Studley Road
Heidelberg VIC 3084
Tel: (03) 9496 4410
www.austin.org.au/poisons

QLD

Pharmacy Department
Royal Children's Hospital
Herston QLD 4029
Tel: 131 126

WA

Sir Charles Gairdner Hospital
Hospital Avenue
Nedlands WA 6009
Tel: 131 126

TAS

Tel: 131 126

NT

Tel: 131 126

ACT

Tel: 131 126

Drug Information Centres

NSW

Drug Information Pharmacist
New South Wales Medicines Information
Centre
PO Box 766
Darlinghurst NSW 2010
Tel: (02) 8382 2136

OR

Drug Information Pharmacist
Hunter Drug Information Service
Newcastle Mater Misericordiae Hospital
Locked Bag 7
Hunter Regional Mail Centre NSW 2310
Tel: (02) 4921 1278
Tel: (02) 4921 1328

VIC

Drug Information Pharmacist
Austin & Repatriation Medical Centre
Studley Road
Heidelberg Vic 3084
Tel: (03) 9496 5668

OR

Drug Information Pharmacist
Drug Information Centre
Southern Health Care Network
Monash Medical Centre
246 Clayton Road
Clayton Vic 3168
Tel: (03) 9594 2361

QLD

Assistant Director of Pharmacy
Queensland Drug Information Ctr
Royal Brisbane Hospital
E Floor, Block 7
Herston Road
Herston Qld 4029
Tel: (07) 3636 7098
(07) 3636 7599

SA

Drug Information Pharmacist
Royal Adelaide Hospital
North Terrace
Adelaide SA 5000
Tel: (08) 8222 5546

OR

Drug Information Pharmacist
Flinders Medical Centre
Bedford Park SA 5042
Tel: (08) 8204 5301

OR

Drug Information Pharmacist
Queen Elizabeth Hospital
Woodville Road
Woodville SA 5011
Tel: (08) 8222 6777

WA

Drug Information Pharmacist
Sir Charles Gairdner Hospital
Hospital Avenue
Nedlands WA 6009
Tel: (08) 9346 2923

TAS

Drug Information Pharmacist
Royal Hobart Hospital
GPO Box 1061L
Hobart Tas 7001
Tel: (03) 6222 8737

NT

Drug Information Pharmacist
Royal Darwin Hospital
PO Box 41326
Casuarina NT 0811
Tel: (08) 8922 8424

ACT

Drug Information Pharmacist
Canberra Hospital
Yamba Drive
Garran ACT 2605
Tel: (02) 6244 3333

National Prescribing Service (NPS)

Therapeutic Advice and Information
Service (TAIS)
Level 7, 418A Elizabeth Street
Surry Hills NSW 2010
Tel: 1300 138 677
Fax: (03) 9459 4546
Email: tais@nps.org.au
Web: www.nps.org.au

List of Contact Officers for Recalls of Therapeutic Goods

For details of consumer level recalls only — telephone 1800 020 512

These officers may be contacted —

- to obtain information about current recalls
- to report suspected problems relating to the quality, safety or efficacy of a therapeutic good

Australian Recall Coordinator

Office of Devices, Blood and Tissues
Therapeutic Goods Administration
Department of Health and Ageing
PO Box 100
Woden ACT 2606
Mr P K Harrison (02) 6232 8636
Mr T Byrne (02) 6232 8637

Australian Capital Territory

ACT Health
GPO Box 825
Canberra ACT 2601
Ms J Strang (02) 6205 0961

New South Wales

Department of Health, NSW
PO Box 103
Gladesville NSW 1675
Mr J E Lumby (02) 9879 3214

Victoria

Department of Human Services
Drugs and Poisons Unit
GPO Box 1670N
Melbourne VIC 3001
Mr K Moyle 1300 364 545
Mr R Bell 1300 364 545

Queensland

Queensland Department of Health
GPO Box 48
Brisbane QLD 4001
Drugs —
Mr A Hawkins (07) 3234 0349
Mr C Healey (07) 3234 0960
Therapeutic Devices—
Mr C Healey (07) 3234 096
Mr D Jones (07) 3406 8068

South Australia

Drug Policies and Programs
Metropolitan Health Division
South Australian Department of Health
PO Box 287
Rundle Mall SA 5000
Mr W Dollman (08) 8226 7110
Ms E Anear (08) 8226 7387

Western Australia

Health Department of WA
PO Box 8172, Perth Business Centre
Perth WA 6849
Mr M Patterson (08) 9388 4980

Tasmania

Department of Health and Human Services
GPO Box 125B
Hobart TAS 7001
Drugs —
Mr J Galloway (03) 6233 2064
Ms M Sharpe (03) 6233 3766
Therapeutic Devices —
Mr A L Wilkins (03) 6233 3913

Northern Territory

Department of Health and Community Services
PO Box 40596
Casuarina NT 0811
Ms H Stone (08) 8922 7035

Index of Manufacturers' codes

<i>Code</i>	<i>Manufacturer</i>
AB	Abbott Australasia Pty Ltd Sir Joseph Banks Corporate Park 32-34 Lord Street Botany NSW 2019 Tel: (02) 9384 9700 Fax: (02) 9384 9800
AC	Alberto Culver Company 14 Loyalty Road North Rocks NSW 2151 Tel: (02) 9630 5099 Fax: (02) 9683 5026
AE	AFT Pharmaceuticals Pty Ltd Level 1, 296 Burns Bay Road Lane Cove NSW 2066 Tel: 1800 097 639 Fax: 1800 097 810
AF	Alphapharm Pty Limited Chase Building 2 Wentworth Park Road Glebe NSW 2037 Tel: (02) 9298 3999 Fax: (02) 9566 4686
AG	Allergan Australia Pty Ltd Level 4, 810 Pacific Highway Gordon NSW 2072 Tel: 1800 252 224 Fax: (02) 9498 0290
AL	Alphapharm Medical A Division of Alphapharm Pty Limited Chase Building 2 Wentworth Park Road Glebe NSW 2037 Tel: (02) 9298 3999 Fax: (02) 9566 4686
AN	Amgen Australia Pty Ltd Level 7, 123 Epping Road North Ryde NSW 2113 Tel: (02) 9870 1333 Fax: (02) 9870 1344
AO	Advanced Medical Optics Australia Pty Ltd Level 3, Building 2 20 Bridge Street Pymble NSW 2073 Tel: 1800 266 111 Fax: 1800 266 222
AP	AstraZeneca Pty Ltd Alma Road North Ryde NSW 2113 Tel: (02) 9978 3500 Fax: (02) 9978 3700
AQ	Alcon Laboratories (Australia) Pty Ltd Allambie Grove Park 25 Frenchs Forest Road East Frenchs Forest NSW 2086 Tel: 1800 025 004 Fax: (02) 9452 5209

<i>Code</i>	<i>Manufacturer</i>
AS	Aspen Pharmacare Australia Pty Ltd First Floor 34-36 Chandos Street St Leonards NSW 2065 Tel: (02) 8436 8300 Fax: (02) 9901 3540
AT	Actelion Pharmaceuticals Australia Pty Ltd Level 2 West, Suites 48-50 7 Narabang Way Belrose NSW 2085 Tel: (02) 9486 4600 Fax: (02) 9986 1344
AV	Aventis Pharma Division of Sanofi-Aventis Australia Pty Limited Building D, Talavera Corporate Centre 12-24 Talavera Road Macquarie Park NSW 2113 Tel: (02) 8666 2000 Fax: (02) 8666 3000
AW	Arrow Pharmaceuticals Pty Ltd A member of Sigma Group of Companies 96 Merrindale Drive Croydon Vic 3136 Tel: (03) 9839 2800 Fax: (03) 9839 2753
BB	Blackmores Ltd 23 Roseberry Street Balgowlah NSW 2093 Tel: (02) 9951 0111 Fax: (02) 9949 1954
BD	Biogen Idec Australia Pty Ltd Suite 2, Level 4 123 Epping Road North Ryde NSW 2113 Tel: (02) 8875 3900 Fax: (02) 9889 1162
BE	Beiersdorf Australia Limited 4 Khartoum Road North Ryde NSW 2113 Tel: (02) 9888 0977 Fax: (02) 9887 3487
BF	Bellwether Pharma Ltd Suite 1, Level 1 1175 Toorak Road Camberwell Vic 3124 Tel: (03) 9809 7900 Fax: (03) 9809 7999
BG	Biochemie Australia A Division of Sandoz Pty Ltd Level 4, Suite 7-19 100 Harris Street Pyrmont NSW 2009 Tel: (02) 9566 1500 Fax: (02) 9566 1458
BI	Biotech Pharmaceuticals Pty Ltd 83 Cherry Lane Laverton North Vic 3026 Tel: (03) 9278 7555 Fax: (03) 9369 6730

<i>Code</i>	<i>Manufacturer</i>
BK	Becton Dickinson Pty Ltd 80 Rushdale Street Knoxfield Vic 3180 Tel: (03) 9764 2444 Fax: (03) 9764 2550
BN	Bayer Australia Limited 875 Pacific Highway Pymble NSW 2073 Tel: (02) 9391 6000 Fax: (02) 9988 3311
BQ	Bristol-Myers Squibb Pharmaceuticals A Division of Bristol-Myers Squibb Australia Pty Ltd 556 Princes Highway Noble Park Vic 3174 Tel: (03) 9213 4000 Fax: (03) 9701 1518
BR	B. Braun Australia Pty Ltd Norwest Business Park 17 Lexington Drive Bella Vista NSW 2153 Tel: (02) 9629 0200 Fax: (02) 9629 0299
BU	Bausch & Lomb Surgical A Division of Bausch & Lomb (Australia) Pty Ltd Level 4, 113 Wicks Road North Ryde NSW 2113 Tel: (02) 9887 1444 Fax: (02) 9888 9642
BV	B.S.N. 315 Ferntree Gully Road Mount Waverley Vic 3149 Tel: (03) 8540 6777 Fax: 1800 671 000
BX	Baxter Healthcare Pty Limited 1 Baxter Drive Old Toongabbie NSW 2146 Tel: (02) 9848 1111 Fax: (02) 9848 1123
BY	Boehringer Ingelheim Pty Limited 78 Waterloo Road North Ryde NSW 2113 Tel: (02) 8875 8800 Fax: (02) 8875 8801
CC	ConvaTec A Division of Bristol-Myers Squibb Australia Pty Ltd 606 Hawthorn Road East Brighton Vic 3187 Tel: 1800 335 276 Fax: (03) 9525 0920
CH	Chem mart Pty Limited Level 7, 5 Queens Road Melbourne Vic 3004 Tel: (03) 9918 2500 Fax: (03) 9918 2006

<i>Code</i>	<i>Manufacturer</i>
CJ	Celgene Pty Ltd Level 7, 607 St Kilda Road Melbourne Vic 3004 Tel: (03) 9539 5500 Fax: (03) 9539 5566
CO	Chemists' Own Pty Ltd A member of Sigma Group of Companies 96 Merrindale Drive Croydon Vic 3136 Tel: (03) 9839 2800 Fax: (03) 9839 2753
CR	Pharmacor Limited 5/36 Campbell Avenue Cromer NSW 2099 Tel: (02) 9981 4470 Fax: (02) 9981 4475
CS	CSL Limited 45 Poplar Road Parkville Vic 3052 Tel: (03) 9389 1911 Fax: (03) 9388 2351
CT	Coloplast Pty Ltd 33 Gilby Road Mount Waverley Vic 3149 Tel: 1800 673 317 Fax: (03) 9541 1199
CU	Care Pharmaceuticals Pty Ltd Suite 303, Level 3, 59-75 Grafton Street Bondi Junction NSW 2022 Tel: 1800 788 870 Fax:
CX	Contact Lens Centre Australia Pty Ltd Unit D6, Hallmark Business Park Cnr Westall and Centre Roads Clayton Vic 3168 Tel: (03) 9543 1811 Fax: (03) 9543 8066
DQ	Church & Dwight (Australia) Pty Ltd Unit 1/108 Old Pittwater Road Brookvale NSW 2100 Tel: 1800 222 099 Fax:
EH	Entra Health Systems Pty Ltd 12/60 Castlereagh Street Sydney NSW 2000 Tel: (02) 8005 4745 Fax: (02) 8088 7105
EO	Ego Pharmaceuticals Pty Ltd 21-31 Malcolm Road Braeside Vic 3195 Tel: (03) 9587 1088 Fax: (03) 9580 7647
EX	Essex Laboratories Level 4, 66 Waterloo Road North Ryde NSW 2113 Tel: (02) 8988 8000 Fax: (02) 9852 7500

<i>Code</i>	<i>Manufacturer</i>
FB	Pierre Fabre Medicament Australia Pty Limited Unit 26B, Parkview Business Centre 1 Maitland Place Baulkham Hills NSW 2153 Tel: (02) 8858 2800 Fax: (02) 8858 2888
FK	Invida Australia Pty Ltd Level 8, 67 Albert Avenue Chatswood NSW 2067 Tel: (02) 9080 7200 Fax: (02) 9080 7201
FM	Fawns and McAllan Pty Ltd A member of Sigma Group of Companies 96 Merrindale Drive Croydon Vic 3136 Tel: (03) 9839 2800 Fax: (03) 9839 2753
FP	Ferring Pharmaceuticals Pty Ltd Suite 2, Level 1, Building 1 Pymble Corporate Centre 20 Bridge Street Pymble NSW 2073 Tel: (02) 9497 2300 Fax: (02) 9497 2399
FR	Charles E. Frosst Division of Merck Sharp & Dohme (Australia) Pty Ltd 54-68 Ferndell Street South Granville NSW 2142 Tel: (02) 9795 9500 Fax: (02) 9795 9595
GA	Galderma Australia Pty Ltd Suite 4, 13B Narabang Way Belrose NSW 2085 Tel: (02) 9479 0600 Fax: (02) 9986 1699
GC	GlaxoSmithKline Consumer Healthcare 82 Hughes Avenue Ermington NSW 2115 Tel: (02) 9684 0888 Fax: (02) 9684 6958
GH	Goldshield Healthcare (Australia) Pty Ltd Suite 3, Level 1 118-124 Willoughby Road Crows Nest NSW 2059 Tel: (02) 9431 6333 Fax: (02) 9906 7147
GI	Gilead Sciences Pty Ltd Level 1, 128 Jolimont Road East Melbourne Vic 3002 Tel: (03) 9272 4400 Fax: (03) 9272 4435
GK	GlaxoSmithKline Australia Pty Ltd Level 4, 436-438 Johnston Street Abbotsford Vic 3067 Tel: (03) 9413 7300 Fax: (03) 8761 2410

<i>Code</i>	<i>Manufacturer</i>
GM	Ascent Pharma Pty Ltd 151-153 Clarendon Street South Melbourne Vic 3205 Tel: 1800 678 302 Fax: (03) 8677 6666
GN	Ascent Pharmaceuticals Limited 151-153 Clarendon Street South Melbourne Vic 3205 Tel: 1800 678 302 Fax: (03) 8677 6666
GP	GP Laboratories A Division of Pfizer Pty Limited 38-42 Wharf Road West Ryde NSW 2114 Tel: (02) 9850 3333 Fax: (02) 9858 1347
GQ	Generic Health Pty Ltd Suite 1, Level 1 1175 Toorak Road Camberwell Vic 3124 Tel: (03) 9809 7900 Fax: (03) 9809 7999
GX	GenRx A Division of Apotex Pty Ltd 66 Waterloo Road North Ryde NSW 2113 Tel: (02) 8877 8333 Fax: (02) 8877 8377
GZ	Genzyme Australasia Pty Ltd Level 1, Building C 12-24 Talavera Road North Ryde NSW 2113 Tel: (02) 9978 3900 Fax: (02) 9889 3900
HA	Hamilton Laboratories Pty Ltd 217 Flinders Street Adelaide SA 5000 Tel: (08) 8223 2957 Fax: (08) 8232 1480
HC	Biotech Healthcare A division of Biotech Pharmaceuticals Pty Ltd 83 Cherry Lane Laverton North Vic 3026 Tel: (03) 9278 7555 Fax: (03) 9369 6730
HE	HealthSense Products Pty Ltd 87 Pitfield Crescent Rowville Vic 3178 Tel: 1300 462 188 Fax:
HH	Hospira Pty Ltd (David Bull Laboratories, Faulding Pharmaceuticals) Level 6, 390 St Kilda Road Melbourne Vic 3004 Tel: (03) 9868 0700 Fax: (03) 9868 0111

<i>Code</i>	<i>Manufacturer</i>
HL	Helex-A Pty Ltd 9/7 Anella Avenue Castle Hill NSW 2154 Tel: (02) 9846 1911 Fax: (02) 9846 1930
HR	Paul Hartmann Pty Ltd 27-28/11-21 Underwood Road Homebush NSW 2140 Tel: 1800 805 839 Fax: (02) 8762 7100
HX	Hexal Australia A division of Sandoz Pty Ltd Level 4, Suite 7-19 100 Harris Street Pyrmont NSW 2009 Tel: (02) 9566 1500 Fax: (02) 9566 1458
IA	iNova Pharmaceuticals (Australia) Pty Limited 9-15 Chilvers Road Thornleigh NSW 2120 Tel: (02) 9875 6333 Fax: (02) 9875 6416
IQ	loquin A Division of Alcon Laboratories (Australia) Pty Ltd Allambie Grove Park 25 Frenchs Forest Road East Frenchs Forest NSW 2086 Tel: 1800 025 004 Fax: (02) 9452 5209
IS	Ipsen Pty Ltd Suite 6, 40 Montclair Avenue Glen Waverley Vic 3150 Tel: (03) 8544 8100 Fax: (03) 9562 5152
IT	InterPharma Pty Ltd Suite 3, 14 Sydney Road Manly NSW 2095 Tel: (02) 9976 6876 Fax: (02) 9976 6859
JC	Janssen-Cilag Pty Ltd 1-5 Khartoum Road North Ryde NSW 2113 Tel: (02) 8875 3333 Fax: (02) 8875 3300
JJ	Johnson & Johnson Medical 1-5 Khartoum Road North Ryde NSW 2113 Tel: (02) 9878 9111 Fax: 1800 808 233
JT	Johnson & Johnson Pacific Pty Limited 45 Jones Street Ultimo NSW 2007 Tel: 13 1565 Fax: (02) 8260 8102
KE	Kendall Australasia Pty Ltd 22 Giffnock Avenue North Ryde NSW 2113 Tel: 1800 252 467 Fax: (02) 9888 7378

<i>Code</i>	<i>Manufacturer</i>
KN	Knoll A Division of Abbott Australasia Pty Ltd Captain Cook Drive Kurnell NSW 2231 Tel: (02) 9668 9711 Fax: (02) 9668 8459
KP	KwikPen Products of Eli Lilly Australia Pty Limited 112 Wharf Road West Ryde NSW 2114 Tel: (02) 9325 4444 Fax: (02) 9325 4410
KR	Kenral Division of Pharmacia Australia Pty Limited 59 Kirby Street Rydalmere NSW 2116 Tel: (02) 9848 3000 Fax: (02) 9848 3333
KY	Key Pharmaceuticals Pty Ltd 12 Lyonpark Road Macquarie Park NSW 2113 Tel: (02) 8113 6200 Fax: (02) 8113 6222
LB	Life Bioscience Pty Ltd 10 Atherton Road Oakleigh Vic 3166 Tel: 1800 114 610 Fax: (03) 8660 2785
LM	Link Medical Products Pty Ltd Level 1, Bridgepoint Centre 3 Brady Street Mosman NSW 2088 Tel: (02) 9960 0150 Fax: (02) 9960 0149
LN	Lennon Healthcare A Division of Aspen Pharmacare Australia Pty Ltd First Floor 34-36 Chandos Street St Leonards NSW 2065 Tel: (02) 8436 8300 Fax: (02) 9901 3540
LU	Lundbeck Australia Pty Ltd Unit 1, 10 Inglewood Place Norwest Business Park Baulkham Hills NSW 2153 Tel: (02) 9836 1655 Fax: (02) 9836 1755
LY	Eli Lilly Australia Pty Limited 112 Wharf Road West Ryde NSW 2114 Tel: (02) 9325 4444 Fax: (02) 9325 4410
MD	Macarthur Research Division of Roche Products Pty Ltd 4-10 Inman Road Dee Why NSW 2099 Tel: (02) 9454 9000 Fax: (02) 9981 3229

<i>Code</i>	<i>Manufacturer</i>
MF	Mundipharma Pty Ltd Level 33, 50 Bridge Street Sydney NSW 2000 Tel: (02) 9231 7200 Fax: (02) 9223 0011
MH	Molnlycke Health Care Pty Ltd Building 1, Ground Floor 14 Aquatic Drive Frenchs Forest NSW 2086 Tel: (02) 9453 1144 Fax: (02) 9453 1155
MI	Meditech Int. Pty Ltd Unit 5, 36 Campbell Avenue Cromer NSW 2099 Tel: (02) 9981 4470 Fax: (02) 9981 4475
MK	Merck Sharp & Dohme (Australia) Pty Ltd 54-68 Ferndell Street South Granville NSW 2142 Tel: (02) 9795 9500 Fax: (02) 9795 9595
MM	3M Pharmaceuticals Australia Pty Ltd 9-15 Chilvers Road Thornleigh NSW 2120 Tel: (02) 9875 6333 Fax: (02) 9875 6416
MQ	Alphapharm Pharmaceuticals Chase Building 2 Wentworth Park Road Glebe NSW 2037 Tel: (02) 9298 3999 Fax: (02) 9566 4686
MS	Abbott Diabetes Care (A Division of Abbott Australasia Pty Ltd) 666 Doncaster Road Doncaster Vic 3108 Tel: (03) 9843 7100 Fax: (03) 9855 8020
MT	Mentholatum Australasia Pty Ltd 12-16 Janine Street Scoresby Vic 3179 Tel: (03) 9763 0322 Fax: (03) 9763 2699
MW	Biomed Aust Pty Ltd c/- Robinson Legal Level 4, 350 Kent Street Sydney NSW 2000 Tel: (02) 9299 2100 Fax: (02) 9299 2201
NA	National Diagnostic Products 22/39 Herbert Street St Leonards NSW 2065 Tel: (02) 9432 8100 Fax: (02) 9432 1151
NC	Novartis Consumer Health Australasia Pty Ltd 327-333 Police Road Mulgrave Vic 3170 Tel: (03) 9701 2711 Fax: (03) 9701 2911

<i>Code</i>	<i>Manufacturer</i>
NE	Norgine Pty Limited 3/14 Rodborough Road Frenchs Forest NSW 2086 Tel: (02) 9972 7500 Fax: (02) 9972 7522
NF	FlexPen Products of Novo Nordisk Pharmaceuticals Pty Ltd Level 3, 21 Solent Circuit Baulkham Hills NSW 2153 Tel: (02) 8858 3600 Fax: (02) 8858 3799
NH	Nycomed Healthcare Pty Limited 2 Lyon Park Road Macquarie Park North Ryde NSW 2113 Tel: (02) 9859 6900 Fax: (02) 9859 6950
NI	InnoLet Products of Novo Nordisk Pharmaceuticals Pty Ltd Level 3, 21 Solent Circuit Baulkham Hills NSW 2153 Tel: (02) 8858 3600 Fax: (02) 8858 3799
NL	NovoLet Products of Novo Nordisk Pharmaceuticals Pty Ltd Level 3, 21 Solent Circuit Baulkham Hills NSW 2153 Tel: (02) 8858 3600 Fax: (02) 8858 3799
NM	Novartis Medicines A Division of Novartis Pharmaceuticals Australia Pty Ltd 54 Waterloo Road North Ryde NSW 2113 Tel: (02) 9805 3555 Fax: (02) 9887 4551
NO	Novo Nordisk Pharmaceuticals Pty Ltd Level 3, 21 Solent Circuit Baulkham Hills NSW 2153 Tel: (02) 8858 3600 Fax: (02) 8858 3799
NQ	Nycomed Pty Ltd 2 Lyon Park Road Macquarie Park North Ryde NSW 2113 Tel: (02) 9859 6900 Fax: (02) 9859 6950
NT	Nestlé Australia Ltd 60 Bathurst Street Sydney NSW 2000 Tel: (02) 9931 2345 Fax: (02) 9931 2610

<i>Code</i>	<i>Manufacturer</i>
NU	Nutricia Australia Pty Limited Talavera Corporate Centre Level 4, Building D 12-24 Talavera Road North Ryde NSW 2113 Tel: (02) 8875 0300 Fax: (02) 8978 4841
NV	Novartis Pharmaceuticals Australia Pty Ltd 54 Waterloo Road North Ryde NSW 2113 Tel: (02) 9805 3555 Fax: (02) 9887 4551
NX	Nipro Australia Pty Ltd Suite 2, 20 Churchill Crescent Cammeray NSW 2062 Tel: 1800 451 737 Fax: (03) 9879 9945
NZ	Nycomed Services Pty Limited 2 Lyon Park Road Macquarie Park North Ryde NSW 2113 Tel: (02) 9859 6900 Fax: (02) 9859 6950
OA	Orphan Australia Pty Ltd 300 Frankston-Dandenong Road Dandenong Vic 3175 Tel: (03) 9709 2200 Fax: (03) 9709 2299
OB	Oral B Laboratories Pty Ltd Level 3, 90 Mount Street North Sydney NSW 2060 Tel: (02) 9957 6499 Fax: (02) 9957 5383
OE	Omegapharm Pty Ltd 21 Queen Street Ormond Vic 3204 Tel: (03) 9483 0070 Fax: (03) 9483 0070
OI	Boian Surgical Pty Ltd 486 King Georges Road Beverly Hills NSW 2209 Tel: (02) 9580 7447 Fax: (02) 9580 7450
OL	Owen Laboratories Division of Galderma Australia Pty Ltd 9 Rodborough Road Frenchs Forest NSW 2086 Tel: 1800 800 765 Fax: (02) 9975 5374
OM	Colgate Oral Care 345 George Street Sydney NSW 2000 Tel: (02) 9229 5600 Fax: (02) 9232 8448

<i>Code</i>	<i>Manufacturer</i>
ON	Orion Laboratories Pty Ltd 25-29 Delawney Street Balcatta WA 6021 Tel: (08) 9441 7800 Fax: (08) 9441 7888
OZ	Medical Specialties Australia Pty Ltd 54 Gibbes Street Chatswood NSW 2067 Tel: (02) 9417 7955 Fax: (02) 9417 5779
PD	Parke Davis Pty Ltd 32 Cawarra Road Caringbah NSW 2229 Tel: (02) 9710 6500 Fax: (02) 9710 6400
PE	Pacific EyeCare A Division of Allergan Australia Pty Ltd Level 4, 810 Pacific Highway Gordon NSW 2072 Tel: 1800 252 224 Fax: (02) 9498 0290
PF	Pfizer Pty Limited 38-42 Wharf Road West Ryde NSW 2114 Tel: (02) 9850 3333 Fax: (02) 9858 1347
PH	Pharmacia Australia Pty Limited 38-42 Wharf Road West Ryde NSW 2114 Tel: (02) 9850 3333 Fax: (02) 9858 1347
PK	Fresenius Kabi Australia Pty Limited 964 Pacific Highway Pymble NSW 2073 Tel: 1300 732 001 Fax: 1300 304 384
PL	Phebra 332 Burns Bay Road Lane Cove NSW 2066 Tel: (02) 9420 9199 Fax: (02) 9420 9177
PM	PMC Pharma A Division of AstraZeneca Pty Ltd Alma Road North Ryde NSW 2113 Tel: (02) 9978 3500 Fax: (02) 9978 3700
PP	Petrus Pharmaceuticals Pty Ltd Level 3, IBM Building 1060 Hay Street West Perth WA 6005 Tel: (08) 9368 5954 Fax: (08) 9368 6692
PQ	PMIP Pty Ltd Unit 18 6a Prosperity Parade Warriewood NSW 2102 Tel: (02) 9997 7176 Fax: (02) 9960 1049

<i>Code</i>	<i>Manufacturer</i>
PX	Point of Care Diagnostics Australia Pty Ltd Unit 14, 76 Reserve Road Artarmon NSW 2064 Tel: (02) 9437 1355 Fax: (02) 9437 1399
PY	Procter & Gamble Pharmaceuticals Australia Pty Ltd 99 Phillip Street Parramatta NSW 2150 Tel: (02) 9685 4500 Fax: (02) 9685 4777
PZ	Prohealth Asia Pacific Pty Ltd Suite 108A, 20 Lexington Drive Bella Vista NSW 2153 Tel: 1300 024 784 Fax: 1300 008 463
QB	Bionime Australia Pty Ltd Level 7, 60 York Street Sydney NSW 2000 Tel: (02) 9262 6900 Fax: (02) 9262 6922
RA	Ranbaxy Australia Pty Limited Suite 4.02, Level 4 Building D 12-24 Talavera Road North Ryde NSW 2113 Tel: (02) 9647 1172 Fax: (02) 9647 1172
RB	BioRevive Pty Ltd Level 1, 263 Mary Street Richmond Vic 3121 Tel: (03) 8416 0399 Fax: (03) 8416 0345
RC	Reckitt Benckiser (Australia) Pty Limited 44 Wharf Road West Ryde NSW 2114 Tel: (02) 9857 2000 Fax: (02) 9857 2004
RD	Roche Diagnostics Australia Pty Ltd 31 Victoria Avenue Castle Hill NSW 2154 Tel: (02) 9899 7999 Fax: (02) 9634 4696
RO	Roche Products Pty Ltd 4-10 Inman Road Dee Why NSW 2099 Tel: (02) 9454 9000 Fax: (02) 9971 7401
RZ	Dr Reddy's Laboratories (Australia) Pty Ltd Level 1, 181 Bay Street Brighton Vic 3186 Tel: (03) 9595 3812 Fax: (03) 9595 3800
SB	Nutricia Australia - Clinical A division of Nutricia Australia Pty Limited Talavera Corporate Centre Level 4, Building D 12-24 Talavera Road North Ryde NSW 2113 Tel: (02) 8875 0300 Fax: (02) 8978 4841

<i>Code</i>	<i>Manufacturer</i>
SC	Schering Pty Ltd Australian Subsidiary of Schering AG, Berlin 875 Pacific Highway Pymble NSW 2073 Tel: (02) 9391 6000 Fax: (02) 9988 3311
SE	Servier Laboratories (Aust.) Pty Ltd 8 Cato Street Hawthorn Vic 3122 Tel: (03) 8823 7333 Fax: (03) 9822 9790
SG	Merck Serono Australia Pty Ltd Unit 3-4, 25 Frenchs Forest Road East Frenchs Forest NSW 2086 Tel: (02) 8977 4100 Fax: (02) 9975 1516
SH	Schering-Plough Pty Ltd Level 4, 66 Waterloo Road North Ryde NSW 2113 Tel: (02) 8988 8000 Fax: (02) 9852 7500
SI	Sigma Pharmaceuticals (Australia) Pty Ltd A member of Sigma Group of Companies 96 Merrindale Drive Croydon Vic 3136 Tel: (03) 9839 2800 Fax: (03) 9839 2753
SJ	Sharpe Laboratories Pty Ltd 12 Hope Street Ermington NSW 2115 Tel: (02) 9858 5622 Fax: (02) 9858 5957
SM	Abbott Products Pty Ltd Level 1, Building 2 Pymble Corporate Centre 20 Bridge Street Pymble NSW 2073 Tel: (02) 9440 0977 Fax: (02) 9440 0910
SN	Smith & Nephew Healthcare 315 Ferntree Gully Road Mount Waverley Vic 3149 Tel: (03) 8540 6777 Fax: 1800 671 000
SS	SSL Australia Pty Ltd 225 Beach Road Mordialloc Vic 3195 Tel: 1800 999 155 Fax: (03) 9587 6870
SW	Sanofi-Aventis Australia Pty Ltd Building D, Talavera Corporate Centre 12-24 Talavera Road Macquarie Park NSW 2113 Tel: (02) 8666 2000 Fax: (02) 8666 3000

<i>Code</i>	<i>Manufacturer</i>
SY	Schering AG 875 Pacific Highway Pymble NSW 2073 Tel: (02) 9391 6000 Fax: (02) 9988 3311
SZ	Sandoz Pty Ltd Level 4, Suite 7-19 100 Harris Street Pyrmont NSW 2009 Tel: (02) 9566 1500 Fax: (02) 9566 1458
TA	Actavis Australia Pty Ltd Upper Ground Floor 183 Melbourne Street North Adelaide SA 5006 Tel: (08) 8267 1545 Fax: (08) 8267 2642
TM	Technipro Marketing Pty Ltd Unit 10, 13 Berry Street Clyde NSW 2142 Tel: (02) 9897 5899 Fax: (02) 9897 5799
TS	Specialised Therapeutics Australia Pty Ltd Suite 3-4, 6 Westbrook Street East Kew Vic 3102 Tel: (03) 9859 1493 Fax: (03) 9859 6950
TW	Terry White Chemists Level 7, 5 Queens Road Melbourne Vic 3004 Tel: (03) 9918 2500 Fax: (03) 9918 2006
TX	Apotex Pty Ltd 66 Waterloo Road North Ryde NSW 2113 Tel: (02) 8877 8333 Fax: (02) 8877 8377
UC	UCB Pharma A Division of UCB Australia Pty Ltd Level 1, 1155 Malvern Road Malvern Vic 3144 Tel: (03) 9828 1800 Fax: (03) 9828 1860
UM	Unomedical Pty Ltd 11-17 Wilmette Place Mona Vale NSW 2103 Tel: (02) 9997 8033 Fax: (02) 9997 3760
VF	Vitaflo Australia Pty Ltd 110 Fyans Street South Geelong Vic 3220 Tel: (03) 5229 8222 Fax: (03) 5229 8225
VI	ViiV Healthcare Pty Ltd Level 4, 436-438 Johnston Street Abbotsford Vic 3067 Tel: (03) 9413 7300 Fax: (03) 8761 2456

<i>Code</i>	<i>Manufacturer</i>
VT	Valeant Pharmaceuticals Australasia Pty Ltd Level 7, Suite 7.02 3 Rider Boulevard Rhodes NSW 2138 Tel: 1800 630 056 Fax: (02) 9743 4053
WA	Winthrop Pharmaceuticals Division of Sanofi- Aventis Australia Pty Limited Building D, Talavera Corporate Centre 12-24 Talavera Road Macquarie Park NSW 2113 Tel: (02) 8666 2000 Fax: (02) 8666 3000
WF	MedWatchDoc 27 Goodwin Street West Ryde NSW 2114 Tel: (02) 9809 0665 Fax: (02) 9989 8469
WQ	Willow Pharmaceuticals Pty Limited Level 31, ABN Amro Tower 88 Phillip Street Sydney NSW 2000 Tel: (02) 9518 1735 Fax: (02) 9518 1835
WT	Wyeth Consumer Healthcare Pty Ltd 17-19 Solent Circuit Norwest Business Park Baulkham Hills NSW 2153 Tel: 1800 555 057 Fax: (02) 9023 0016
WX	Wyeth Australia Pty Limited 38-42 Wharf Road West Ryde NSW 2114 Tel: (02) 9850 3333 Fax: (02) 9813 4011
WY	Wyeth Pharmaceuticals Division of Wyeth Australia Pty Limited 38-42 Wharf Road West Ryde NSW 2114 Tel: (02) 9850 3333 Fax: (02) 9813 4011
XF	Max Pharma Pty Ltd Suite 1, Level 1 1175 Toorak Road Camberwell Vic 3124 Tel: (03) 9809 7900 Fax: (03) 9809 7999
XP	Aaxis Pacific Pty Ltd 24-32 Forge Street Blacktown NSW 2148 Tel: (02) 9881 3333 Fax: (02) 9881 3322
YM	Symbion Pharmacy Services Pty Ltd Level 7, 5 Queens Road Melbourne Vic 3004 Tel: (03) 9918 2000 Fax: (03) 9918 2006

<i>Code</i>	<i>Manufacturer</i>
YN	Mayne Pharma International Pty Ltd 1538 Main North Road Salisbury SA 5106 Tel: (08) 8209 2666 Fax: (08) 8281 6998
YT	Mayne Products Pty Ltd 1538 Main North Road Salisbury SA 5106 Tel: (08) 8209 2666 Fax: (08) 8281 6998
ZF	Sun Pharmaceutical Industries (Australia) Pty Ltd 1053 Burwood Highway Ferntree Gully Vic 3156 Tel: (03) 9568 6102 Fax: (03) 9568 6610

<i>Code</i>	<i>Manufacturer</i>
ZI	Shire Australia Pty Limited Level 9, Avaya House 123 Epping Road North Ryde NSW 2113 Tel: 1800 012 612 Fax: (02) 8875 7977
ZP	Spirit Pharmaceuticals Pty Ltd 117 Harrington Street The Rocks Sydney NSW 2000 Tel: (02) 9251 1088 Fax: (02) 9251 1099

Section 1 — Explanatory Notes

Introduction

These Explanatory Notes are provided to help PBS prescribers and pharmacists work within the Australian Government's Pharmaceutical Benefits Scheme (PBS).

The PBS is a system of subsidising the cost of most prescription medicines. The subsidies are available to all Australian residents and eligible foreign visitors, i.e., people from countries which have Reciprocal Health Care Agreements with Australia. These countries are the United Kingdom, Ireland, New Zealand, Malta, Italy, Sweden, the Netherlands, Finland, Norway and Belgium.

The aim of the PBS, which has been in operation since 1948, is to provide reliable and affordable access to a wide range of necessary medicines.

The Schedule of Pharmaceutical Benefits referred to throughout as the 'Schedule' – lists all the medicinal products available under the PBS, and explains the uses for which they can be subsidised.

The Schedule is produced monthly by the Australian Department of Health and Ageing (effective on the first day of each month).

It is vital therefore that PBS prescribers and pharmacists remain up to date with information on which medicines are included in or excluded from the Schedule, which PBS prescribers may prescribe certain medicines, whether restrictions apply to the medicines, and how much patients should pay.

Queries relating to the PBS can be made to the Pharmaceutical Benefits Branch of Medicare Australia (telephone 132 290 open 24 hours a day, 7 days a week). Queries relating to the Repatriation Pharmaceutical Benefits Scheme (RPBS) can be made to the State offices of the Department of Veterans' Affairs (DVA) (telephone 1800 552 580).

1. The Schedule — Where to Find What

The Schedule of Pharmaceutical Benefits is divided into sections. At the start of the Schedule, immediately after the table of contents, is a summary of any changes to listed items. This is followed by a list of important information sources, contacts and addresses, then an index of manufacturers' codes.

The last pages of the Schedule provide a generic/proprietary index of PBS and RPBS ready-prepared items.

Section 1

Section 1 is what you are reading, the Explanatory Notes. It outlines the correct way to prescribe and supply pharmaceutical benefits; patient charges; who qualifies for concessions; how the Safety Net system works; and, for pharmacists, how to claim reimbursement for PBS items.

Please note that except where indicated, the term '**prescriber**' is used in this section to cover doctors, dentists, optometrists, midwives and nurse practitioners who are approved to prescribe PBS medicines under the National Health Act 1953.

And except where stated otherwise, the term '**pharmacist**' means a pharmacist approved to supply medicines under the PBS.

Section 2

This section lists ready-prepared items, and includes the form, manner of administration, brand and brand equivalents which may be prescribed, and the maximum quantity and number of repeats for each item.

Emergency drug supplies are also listed at the beginning of this section.

Any medicines that have restrictions on how they can be prescribed are printed in ***bold italics***. Items appearing in more than one therapeutic group are cross-referenced.

The second page of Section 2 explains symbols used throughout the Schedule.

The use of 'NOTE' in this section is used to clarify how some pharmaceutical benefits should be prescribed.

The use of 'CAUTION' is to warn of known adverse reactions from, or precautions to be taken with, a particular pharmaceutical benefit. (The absence of a cautionary note does not imply reactions may not happen.)

Separate lists at the end of Section 2 relate to items that can be prescribed by dentists and optometrists who work within the PBS. These are followed by a list of items that are made available under special arrangements for doctors to prescribe.

Section 3

This section lists container prices, fees related to dispensing, standard packs and prices for ready-prepared preparations.

Section 4

This section deals with extemporaneous preparations. It lists the ingredients which can be used, a table of maximum quantities and number of repeats, container prices, and a list of standard formula preparations and prices (based on formularies in common use and referred to in the Schedule as the Standard Formulae List).

Restrictions applying to the use of a pharmaceutical benefit are indicated against the item.

Repatriation Schedule of Pharmaceutical Benefits

After Section 4, the Schedule provides information about pharmaceutical benefits under the RPBS. These may only be prescribed to DVA beneficiaries holding one of the repatriation health cards (see details under '4. Patient Charges').

2. Prescribing Medicines – Information for PBS Prescribers

PBS prescribers

Pharmaceutical benefits can only be prescribed by doctors, dentists, optometrists, midwives and nurse practitioners who are approved to prescribe PBS medicines under the *National Health Act 1953*.

PBS Prescription forms

Standard PBS prescription forms are available from Medicare Australia for prescribing pharmaceutical benefits.

For doctors:

- *Personalised forms* — are printed with the doctor's name, qualifications, practice address/es, telephone number and prescriber number (which relates to pharmaceutical benefits). They are only provided to doctors who have a Medicare provider number.
- *Non-personalised (blank) forms* — are distributed as an emergency supply (usually when a doctor has temporarily run out of personalised forms).
- *Locum forms* — have the doctor's name, prescriber number and telephone number (if available) and a space to record the practice where the doctor is working.
- *PBS/RPBS Authority Prescription Forms* — can be in personalised, non-personalised or locum format.
- *Computer PBS prescription forms* — are either continuous or single sheet. On the reverse side they list the name, address and telephone number of the practice, and in the case of a sole doctor practice, the doctor's name.

For dentists:

- *Personalised forms* — have the dentist's name, qualifications, practice address/es, telephone number and prescriber number.
- *Non-personalised (blank) forms* — are distributed for emergency supply only.

For optometrists:

- *Personalised forms* — have the optometrist's name, qualifications, practice address/es, telephone number and prescriber number. These forms can be also be used to prescribe authority-required PBS/RPBS items.

For midwives:

- *Personalised forms* — have the midwife's name, qualifications, practice address/es, telephone number and prescriber number.
- *Non-personalised (blank) forms* — are distributed for emergency supply only.

For nurse practitioners:

- *Personalised forms* — have the nurse practitioner's name, qualifications, practice address/es, telephone number and prescriber number.
- *Non-personalised (blank) forms* — are distributed for emergency supply only.

PBS prescription forms for PBS prescribers are supplied free of charge.

The inclusion of the prescriber number on a PBS prescription enables the pharmacist to be sure the prescription is from a legitimate prescriber and satisfies State/Territory legislation. A PBS prescription written by a dentist, an optometrist, a midwife or a nurse practitioner must include the person's approval number as a PBS prescriber.

PBS prescriptions should be provided to the patient in duplicate, as both parts make up a valid PBS prescription. The patient should be reminded to present both the original and the duplicate copy to the pharmacist.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner providing medical treatment or a midwife providing midwifery treatment or a nurse practitioner providing nurse practitioner treatment within a participating public hospital may prescribe PBS subsidised medication. The States of Victoria, Queensland, South Australia and Western Australia, and the Northern Territory have agreed to implement these arrangements.

Ordering forms

Prescribers are asked not to over order. Getting the right amount of forms helps to reduce the cost to taxpayers and helps to reduce paper wastage. Also, the pads may deteriorate if stored over time.

Order forms for standard and authority PBS prescription forms are available from Medicare Australia stationery officers. Contact details are listed in the front of the Schedule. Order forms for computer PBS prescription form stationery are obtained from Medicare Australia (at the address below). Orders should be sent to:

Prescription Pad Order Clerk
Pharmaceutical Branch
Medicare Australia
GPO Box 9826
Sydney NSW 2001
Telephone (02) 9895 3295

Orders for PBS prescription stationery will only be accepted by application in writing and through the channels mentioned above.

Preparing general PBS prescriptions

Do's and Don't's

A PBS prescription is only valid when it is written by a doctor, a dentist, an optometrist, a midwife or a nurse practitioner.

The PBS prescription must be for the treatment of the person named on the PBS prescription. A PBS prescription may only be written for the treatment of one person.

A prescriber cannot write more than one PBS prescription for the same pharmaceutical benefit for the same person on the same day.

Up to **three** pharmaceutical benefit items may be included on a single PBS prescription form except for Authority required, Authority required (STREAMLINED) items and optometrist items. These items must be written on individual forms. Pharmaceutical benefits and non-pharmaceutical benefits should not be listed together on the one PBS prescription form.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner providing medical treatment or a midwife providing midwifery treatment or a nurse practitioner providing nurse practitioner treatment within a participating public hospital may prescribe PBS subsidised medication. The States of Victoria, Queensland, South Australia and Western Australia, and the Northern Territory have agreed to implement these arrangements.

If an item has a particular manner of administration it may not, as a pharmaceutical benefit, be administered in any other way, e.g., an ophthalmic preparation may not be prescribed for topical use.

If an item is restricted, and the use for the patient is different from the use specified in the restriction, it cannot be prescribed as a pharmaceutical benefit. The prescriber should write the prescription as a non-PBS private prescription. If a standard PBS prescription form is used for this purpose the 'PBS/RPBS' text must be clearly struck out. It should also be endorsed 'non-PBS'.

Prescribers must heed State/Territory laws when prescribing drugs listed as narcotic, specified or restricted in the poisons legislation of the particular State or Territory. Legislative requirements in some States/Territories are such that prescribers may be required to prescribe a drug of addiction on a separate PBS prescription. Prescribers must ensure that prescriptions written under the PBS fall within the limits of the prescribing approval granted to the person under State or Territory requirements. It is the prescriber's responsibility to ensure that PBS prescriptions comply with all aspects of his/her prescriber approval. Inclusion of a PBS medicine for prescribing does NOT confer approval for a particular prescriber to prescribe that medicine if it is not authorised to be prescribed in a particular State or Territory

A prescriber cannot prescribe a narcotic drug for him/herself.

Prescribers are issued with individual PBS prescription pads by Medicare Australia for their own use — these pads should not be used by other prescribers, as this can cause confusion through incorrect pharmacy records.

Doctors should, and dentists and optometrists, midwives and nurse practitioners are required to, include their prescriber number on non-personalised PBS prescriptions.

The following admixtures are not pharmaceutical benefits:

- the admixture of two or more ready-prepared items listed in the Schedule; or
- the admixture of a ready-prepared item and one or more extemporaneous drugs listed in Section 4 of the Schedule; or
- the admixture of a non-pharmaceutical benefit item with a pharmaceutical benefit item.

Writing the PBS prescription

The following rules apply for writing PBS prescriptions:

- they must be written in indelible form (i.e., ink or ball-point pen) in the prescriber's own handwriting (exceptions must be approved by Medicare Australia's Chief Executive Officer) either on the standard PBS prescription, or on paper approximately 18 cm x 12 cm, or they can be generated by computer on a form approved by Medicare Australia. For patient safety reasons, both the original and the duplicate must be legible;
- they must record the prescriber's name and address (and, in the case of dentists, optometrists, midwives and nurse practitioners, the prescriber number), the patient's name, address and entitlement status, and whether the prescription is under the PBS or RPBS;
- they should completely identify the pharmaceutical benefit by detailing the item, dose, form, strength, quantity and instructions for use;
- they should indicate where brand substitution is not permitted. PBS prescriptions must not be prepared using a computer prescribing program that contains a default which would result in all prescriptions being indicated as Brand Substitution Not Permitted;
- where 'solvent required' is included after the form, the volume and number of ampoules must be specified; and
- they must be signed by the prescriber and dated. Forward or back dating is not permitted.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner providing medical treatment or a midwife providing midwifery treatment or a nurse practitioner providing nurse practitioner treatment within a participating public hospital may prescribe PBS subsidised medication. The States of Victoria, Queensland, South Australia and Western Australia, and the Northern Territory have agreed to implement these arrangements.

Restrictions

Pharmaceutical benefits listed in the Schedule fall into three broad categories:

Unrestricted benefits - have no restrictions on their therapeutic uses;

Restricted benefits - can only be prescribed for specific therapeutic uses (noted as Restricted benefit); and

Authority required benefits - Authority required benefits fall into two categories:

- *Authority required benefits* are restricted benefits that require prior approval from Medicare Australia or the DVA (noted as **Authority required**)
- *Authority required (STREAMLINED) benefits* are restricted benefits that do not require prior approval from Medicare Australia or the DVA but require the recording of a streamlined authority code (noted as **Authority required (STREAMLINED)**).

Authority PBS prescriptions

Authority required benefits fall into two categories - *Authority required* and *Authority required (STREAMLINED)*.

All PBS prescribers (with the exception of dentists) can write authority PBS prescriptions.

Authority PBS prescriptions cannot have retrospective approval.

Authority required PBS Prescriptions

Approval of authority PBS prescriptions by Medicare Australia may be sought by:

- posting an Authority Prescription Form to Medicare Australia - after approval, Medicare Australia will forward both copies of the prescription to the patient or the prescriber (if it is to be sent direct to the patient, the prescriber should mark the box next to the patient's details);
- calling Medicare Australia Authority Freecall service (1800 888 333); or
- using Medicare Australia PBS authorities website at www.medicareaustralia.gov.au/providers.

Approval of authority prescriptions by the DVA may be obtained either by posting an Authority Prescription Form to the DVA, or by using the DVA Authority Freecall service (1800 552 580).

An authority PBS/RPBS prescription is not valid until it has been approved by Medicare Australia or the DVA. Without this approval, a pharmacist must not supply the item as a PBS/RPBS benefit.

Each Authority required PBS/RPBS item must be written on an Authority PBS/RPBS prescription form, one item per form. Authority PBS prescription forms provide for the following:

- the patient/pharmacist copy, which records prescriber, patient, and pharmaceutical benefit item details. Where required a repeat authorisation, which is used for repeat supply, is attached to the pharmacist/patient copy until the last supply is made. The patient/pharmacist copy is then retained by the pharmacist;
- the Medicare Australia/DVA copy which records prescriber, patient, and pharmaceutical benefit item details. After the first dispensing, the Medicare Australia/DVA copy is forwarded to Medicare Australia for processing and payment;
- the prescriber's copy (for computer generated scripts, this is the tear off portion at the base of the script) or Prescriber/Medicare Australia/DVA copy (for handwritten scripts this is the long white copy), is kept by Medicare Australia or the DVA for record purposes when approval is sought in writing. When approval is by telephone or by the authorities website, the prescriber must keep this copy for 12 months. This copy must record the daily dose, details of the disease, clinical justification for using the item, the patient's age (if the patient is a child) and whether the patient has previously received an authority for this pharmaceutical benefit.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner providing medical treatment or a midwife providing midwifery treatment or a nurse practitioner providing nurse practitioner treatment within a participating public hospital may prescribe PBS subsidised medication. The States of Victoria, Queensland, South Australia and Western Australia, and the Northern Territory have agreed to implement these arrangements.

Authority required (STREAMLINED) PBS Prescriptions

Prior approval is not required from Medicare Australia or DVA to prescribe an Authority required (STREAMLINED) item (except where increased quantities and/or repeats are required). Instead the authority prescription form must include a four digit streamlined authority code.

This code is listed with the corresponding restriction for each Authority required (STREAMLINED) item and the prescriber must write the code on the authority PBS/RPBS prescription form. An authority prescription for an Authority required (STREAMLINED) item is not valid unless the code is included on the prescription form. Without the streamlined authority code, a pharmacist must not supply the item as a PBS benefit.

There are no Authority Required (STREAMLINED) items in the Repatriation Schedule of Pharmaceutical Benefits.

Authority required (STREAMLINED) PBS prescriptions must be written on an Authority PBS/RPBS Prescription Form, this includes:

- the pharmacist/patient copy, which records prescriber, patient, and pharmaceutical benefit item details. The prescription is given directly to the patient to be dispensed at their pharmacy;

- the Medicare Australia/DVA copy which records prescriber, patient, and pharmaceutical benefit item details. After the first dispensing, the Medicare Australia/DVA copy is forwarded to Medicare Australia for processing and payment;
- the prescriber's copy is kept by the prescriber for 12 months. This copy must record the daily dose, details of the disease, clinical justification for using the item, the patient's age (if the patient is a child) and whether the patient has previously received an authority for this pharmaceutical benefit.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner providing medical treatment or a midwife providing midwifery treatment or a nurse practitioner providing nurse practitioner treatment within a participating public hospital may prescribe PBS subsidised medication. The States of Victoria, Queensland, South Australia and Western Australia, and the Northern Territory have agreed to implement these arrangements.

Writing authority PBS prescriptions

The following rules apply:

- only one item may be prescribed per PBS prescription;
- PBS prescriptions must be completed by prescribers in writing, unless otherwise approved by Medicare Australia;
- prescribers should include their name, address, telephone number and **prescriber number** (not provider number);
- prescribers must include the patient's name, address and entitlement status (i.e. whether they are a 'concessional' or 'general patient');
- prescribers must indicate when brand substitution is not permitted. PBS prescriptions must not be prepared using a computer prescribing program that contains a default which would result in all PBS prescriptions being indicated as Brand Substitution Not Permitted;
- in certain circumstances, the prescriber must provide additional information to Medicare Australia with the authority application; and
- the PBS prescription must be signed by the prescriber and dated.

Posted applications which lack necessary information, and therefore cannot be approved, will be returned for correction. If the matter can be clarified via telephone, an Authority to Prescribe Form may be prepared by Medicare Australia or the DVA and sent to the prescriber.

In the case of authority PBS prescriptions approved by telephone, the approval number must be included on the PBS prescription to enable the pharmacist to supply the medication. A prescriber who is granted approval but decides not to continue with the therapy should advise Medicare Australia.

In the case of Authority required (STREAMLINED) prescriptions, the streamlined authority code must be written on the PBS/RPBS prescription form. This enables the pharmacist to supply the medication as a PBS benefit.

There are separate arrangements for PBS prescriptions in certain public hospitals. To gain access to pharmaceutical benefits under this arrangement a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner providing medical treatment or a midwife providing midwifery treatment or a nurse practitioner providing nurse practitioner treatment within a participating public hospital may prescribe PBS subsidised medication. The States of Victoria, Queensland, South Australia and Western Australia, and the Northern Territory have agreed to implement these arrangements.

Maximum quantities and repeats

The maximum quantity and number of repeats allowed for PBS items are recommended by the Pharmaceutical Benefits Advisory Committee (PBAC). In the case of RPBs items, the recommendations are made by the Repatriation Pharmaceutical Reference Committee (RPRC).

All PBS prescribers (with the exception of dentists) can prescribe repeats.

PBS prescriptions and repeats can be for any quantity up to the maximum. It is not necessary to prescribe the maximum quantity if a lesser quantity is sufficient for the patient's needs. Please clearly indicate the number of tablets, capsules, etc. required and the number of repeats needed, and **do not use** abbreviations such as 'Max. Qty', 'M.Q.', or 'M.R.'.

If a prescriber feels the maximum quantity or number of repeats should be increased for a particular patient, he or she must complete an Authority PBS Prescription Form (see procedures above under 'Authority PBS Prescriptions'). The provision of increased quantities and repeats on authority PBS prescriptions is intended to provide approximately one

month's therapy which may be repeated (if clinically appropriate) to provide 6 months' therapy in total. This situation usually arises where higher than normal dosages are required.

Approval for increased quantities and repeats of Authority required, Authority required (STREAMLINED) and Restricted benefit PBS items will be granted only where the reason for the PBS prescription is consistent with the indications published in the Schedule.

Approval for increased quantities and repeats extends only to the provision of a pharmaceutical benefit for the patient and does not imply approval of any aspects of the patient's care, which are the responsibility of the treating prescriber.

Regulation 24

Under this regulation, original and repeat supplies of pharmaceutical benefits can be supplied at the one time if a medical practitioner, a midwife or a nurse practitioner is first satisfied that certain conditions apply, then endorses the PBS prescription 'Regulation 24'. RPBS prescriptions may be endorsed 'hardship conditions apply'.

The medical practitioner, midwife or nurse practitioner must first be satisfied all the following conditions apply:

- the maximum PBS quantity is insufficient for the patient's treatment; **AND**
- the patient has a chronic illness or lives in a remote area where access to PBS supplies is limited; **AND**
- the patient would suffer great hardship trying to get the pharmaceutical benefit on separate occasions.

Regulation 24 does not apply for supply of pharmaceutical benefits on optometrist prescriptions.

Urgent cases

In urgent cases and where State/Territory law allows, a prescriber may telephone a pharmacist and ask that a PBS prescription be supplied. He/she must then forward the written PBS prescription and duplicate to the pharmacist within **seven days of the date of supply**.

This also applies to 'Authority required' authority PBS prescriptions provided prior approval has been given by Medicare Australia or DVA. The follow-up written PBS prescription must include the approval number provided over the phone by Medicare Australia or DVA.

Drugs of addiction

Prescribers must heed State/Territory laws when prescribing drugs listed as narcotic, specified or restricted and must notify, or receive approval from, the appropriate health authority.

When a PBS/RPBS authority application is for a drug of addiction (other than dexamphetamine sulfate), the following guidelines apply:

- the maximum quantity authorised is generally for one month's therapy (e.g., one week's therapy with three repeats);
- where supply for a longer period is warranted, quantities are usually for up to three months' therapy;
- telephone approvals are limited to one month's therapy.

Prescribers should also state the interval of repeat where repeats are called for, and ensure State/Territory health authorities are notified about ongoing treatment.

Emergency drug supplies

Certain pharmaceutical benefits are provided without charge to prescribers who in turn can supply them free to patients for emergency use.

The Emergency Drug Supply Order Form must be completed in triplicate, signed, and the original and duplicate given to a pharmacist. Each form is valid for the month indicated on the form.

Prescribers may order the maximum quantity of an item provided they do not already have the maximum quantity on hand. The items can only be obtained once a month. Prescribers may also ask for a particular brand of a pharmaceutical benefit. If it is unavailable, they must specify another listed brand, and initial the alteration.

A receipt must be signed by the prescriber, or by an authorised representative, when supplies are received.

Availability of Methoxyflurane for emergency treatment only

A new Emergency Treatment Program (ETP) for medical practitioners has been established to provide for medicines such as Methoxyflurane to be supplied as items for emergency treatment, other than hospital treatment. Unlike other emergency drug supplies, Methoxyflurane, liquid for inhalation 999.9 mg per g, 3 mL (with inhaler) (*Penthrox*[®]) is not available for prescribing as a general pharmaceutical benefit.

Methoxyflurane is therefore PBS-listed as a 'special pharmaceutical product' under section 100AA, only for emergency treatment, other than hospital treatment. As such, the availability of this drug is provided for under special arrangements under section 100 (1) of the *National Health Act 1953*. The legislative instrument can be viewed on the Federal Register of Legislative instruments at www.frli.gov.au.

For the purposes of administration, Methoxyflurane will be listed with other emergency drug supplies, as outlined above, and be managed by Medicare Australia in the same manner as other emergency drug supply items with the same supply and claiming procedures.

Improving the capacity of the PBS to meet particular Aboriginal and Torres Strait Islander health needs

The PBS includes listings to support the treatment of conditions common in Aboriginal and Torres Strait Islander health settings. These listings are specifically for your patients who identify as Aboriginal and/or Torres Strait Islander persons. Some listings will be medicines recently added to the PBS; others may contain specific restrictions for existing PBS items.

A significant proportion of the higher levels of illness experienced by Aboriginal and Torres Strait Islanders may be addressed through better access to appropriate medicines. The PBS aims to provide greater choice in therapeutic options and to address:

- the greater burden of disease experienced by Aboriginal and Torres Strait Islander peoples; and
- morbidity almost exclusively seen in this population.

How to prescribe these items?

These items are available as "Authority PBS prescriptions". You should obtain approval from Medicare Australia before prescribing these items for patients who identify as Aboriginal and/or Torres Strait Islander persons through the Authority Freecall service [1800 888 333], on line or by mail.

All PBS prescribers except dentists can write Authority PBS prescriptions and your patients will be required to pay their normal PBS co-payment.

Special arrangements apply in remote area Aboriginal Health Services for supplying these PBS items.

Aboriginal and Torres Strait Islander identification

Establishing a client's background may have clinical significance and should be part of routine medical history taking. In the case of Aboriginal and Torres Strait Islander people, this is also relevant to establish eligibility for services such as health checks, specific immunisation programs, and the some PBS items.

Improving the level of identification of Aboriginal and Torres Strait Islander people will also assist in developing initiatives to meet particular needs.

For the purposes of these PBS items a person is Aboriginal and/or Torres Strait Islander if the person identifies himself or herself as being an Aboriginal and/or Torres Strait Islander. Clients should be asked to self-identify either verbally or by completing a form.

- Some people may give this information without being asked.
- It is important not to assume that a person is or is not Aboriginal or Torres Strait Islander.

Asking about Aboriginal and/or Torres Strait Islander identification

Practitioners should ensure that each person attending their practice has the opportunity to identify if they are Aboriginal or Torres Strait Islander. An environment which maintains confidentiality and provides an explanation for this question if requested will assist this process.

- The inquiry may be made verbally and recorded by the general practitioner as part of routine medical history taking at first consultation, or by a receptionist or other staff member. An appropriate question to ask is: *"Are you (is this child) of Aboriginal or Torres Strait Islander origin?"*
- Alternatively, the question may be included on a client self-history or practice record form, using a standard question such as: *"Are you (is this child) of Aboriginal or Torres Strait Islander origin?"*
 - Yes - Aboriginal
 - Yes - Torres Strait Islander
 - Yes - Aboriginal and Torres Strait Islander

O No

Aboriginal and Torres Strait Islander health

Major causes of excess mortality in Aboriginal and Torres Strait Islander peoples are:

- circulatory conditions (including ischaemic heart disease, hypertension, cerebrovascular disease and rheumatic heart disease);
- external causes (including accident and injury);
- endocrine causes (mainly type two diabetes and its complications); and
- respiratory conditions.

Causes of morbidity vary but include the risk factors and precursors of all of these. They also include infections of the respiratory system, the ears (in particular, chronic suppurative otitis media), the eyes (trachoma in some settings), the skin and the gastrointestinal system. End-stage renal disease is a major cause of hospitalisations, and much early renal disease remains undetected. In some settings, sexually transmissible infections are common.

Living environments affect health and may be compromised by overcrowding, limited access to clean water and sanitation, and poverty. Social and family life may be negatively influenced by an excessive burden of care for family members, by substance use and sometimes by family violence.

Communication and cultural issues

Aboriginal cultures are numerous and diverse in language, customs, non-verbal and verbal communication, geographical locations and experiences. Torres Strait Islanders are a separate people with a distinctly different culture and identity. Aboriginal and Torres Strait Islander people often perceive health differently from other Australians.

For Aboriginal and Torres Strait Islander peoples' health does not just entail the freedom of the individual from sickness but requires support for healthy and interdependent relationships between families, communities, land, sea and spirit. The focus must be on spiritual, cultural, emotional and social well-being as well as physical health

Source: National Aboriginal and Torres Strait Islander Health Council. National Strategic Framework for Aboriginal and Torres Strait Islander Health 2003-2013, Context. Canberra: Commonwealth of Australia; 2004.

To provide effective primary health care to Aboriginal and Torres Strait Islander clients, you need to be aware of the issues surrounding this diversity, and which may have an impact on the delivery of services.

- Aboriginal and Torres Strait Islander people may be reluctant to use mainstream medical services. This may be because of a lack of understanding of the mainstream health system and previous negative experiences within the mainstream health care system.
- Access to adequate health care may be hindered by family obligations (often extended family), lack of transport or money, or geographical isolation.
- English may be the person's second, third or even fourth language. Therefore it may be appropriate to consider the use of an interpreter.
- Aboriginal and Torres Strait Islander people may be reluctant to consult a health care provider of the opposite sex, particularly with regard to women's and men's health issues.

The differences between the cultural and language backgrounds of health service providers and patients, whether urban, rural or remote, may range from minor to extreme.

You should:

- Make efforts to ensure waiting rooms are welcoming to Aboriginal and Torres Strait Islander people, including displaying relevant posters and pamphlets;
- Provide a relaxed setting for the consultation (e.g. sit next to your patient rather than across a desk);
- Allow time at the first consultation to build rapport and trust;
- Ensure the person understands clearly what the service entails and the details of any procedures involved, and possible follow-up or referral requirements;
- Obtain health promotion information appropriate for Aboriginal and Torres Strait Islander patients;
- Allow the patient to have family members present if desired. When inviting family or community members to accompany a patient, ensure the patient fully consents to their attendance and that the community/family members are fully aware of the need for confidentiality;

- Provide gender appropriate staff where possible, for both male and female patients, especially in regard to pap smears, mammograms, sexual health checks, pregnancy checks, antenatal care and postnatal care;
- Encourage all staff in the practice to attend Aboriginal and Torres Strait Islander Cultural Awareness programs, which are widely available;
- Ensure practice staff have awareness of appropriate referral and/or support organisations for Aboriginal and Torres Strait Islander patients; and
- Develop partnerships with local Aboriginal and Torres Strait Islander community organisations.

For more information, pbs-indigenous@health.gov.au

3. Supplying Medicines — What Pharmacists Need to Know

Eligible suppliers

Pharmaceutical benefits are mainly supplied by approved pharmacists – pharmacists who comply with certain conditions. These pharmacists are approved to dispense pharmaceutical benefits from a particular pharmacy.

Other suppliers include approved doctors (usually practising in isolated areas), Friendly Society pharmacies, and approved hospitals. All suppliers are issued with approval numbers by Medicare Australia. They should follow the procedures in these Explanatory Notes.

Unapproved pharmacists *cannot* supply pharmaceutical benefits.

Approval conditions for pharmacists

A pharmacist approved to supply medicines under the PBS:

- can only supply benefits from the pharmacy that he/she is operating;
- will not supply to anyone any pharmaceutical benefit that attracts a Commonwealth contribution for free, or for a price that is less than the relevant patient contribution;
- will clearly advertise that any offer for free or cut-price medicines does not include pharmaceutical benefits which have a Commonwealth contribution;
- will not pay rebates or refunds of patient contributions;
- will publicly display a notice setting out the pharmacy's normal trading hours;
- is obliged to supply pharmaceutical benefits at the pharmacy at any hour if a PBS prescription is marked 'urgent' and initialled by the prescriber;
- will keep adequate stocks for the supply of pharmaceutical benefits;
- may be called on by Medicare Australia to provide details of stocks of pharmaceutical benefits or preparations for pharmaceutical benefits; and
- must keep the duplicates of all old format PBS prescriptions, and the patient/pharmacist copies of all new format PBS prescriptions, with a Commonwealth contribution for at least one year from the date of supply. This includes PBS prescriptions ordering repeats when it is the final supply, and order forms for emergency drug supplies. Please note that some State/Territory laws require these copies to be kept for longer periods.

Before supplying pharmaceutical benefits

Several steps must be taken before a pharmaceutical benefit is supplied.

Firstly, a pharmacist must endorse the PBS prescription and duplicate with his/her name and approved supplier number.

Secondly, a PBS prescription identifying number must be given to the PBS prescription item on both the PBS prescription and duplicate. Any recognised series of numbers may be used.

If more than one item is on a PBS prescription, a separate identifying number should be allocated to each item.

In the case of a repeat authorisation, the same PBS prescription identifying number(s) must be carried through for each item. A pharmacist must also allocate his/her own identifying number on the repeat authorisation. It must be written alongside the date and place of supply.

Supplying pharmaceutical benefits

Do's and Don'ts

Except in urgent cases (see details under '2. Prescribing Medicines ... Urgent cases'), pharmacists are authorised to supply pharmaceutical benefits only after they receive:

- the pharmacist/patient and Medicare Australia or DVA copies of a valid PBS prescription which is not more than 12 months old; or
- the pharmacist/patient and Medicare Australia or DVA copies of an approved authority PBS prescription or an authority to prescribe which is not more than 12 months old; or
- a repeat authorisation attached to a patient/pharmacist PBS prescription not more than 12 months after the date of the original PBS prescription.

A pharmacist must not supply an Authority required (STREAMLINED) item unless the prescriber has written the four digit streamlined authority code on an authority PBS/RPBS prescription.

A pharmaceutical benefit cannot be supplied more times than specified in the PBS prescription.

A pharmacist cannot add to, delete from, or alter a PBS prescription in any other way. However, there may be circumstances where after contacting a prescriber, the pharmacist can clarify the prescriber's intentions and endorse the PBS prescription accordingly.

Once a pharmaceutical benefit has been supplied to a patient, it may not be supplied to that patient again:

- on the same day or within the next 20 days, if it is a benefit (other than an eye preparation) that has five or more repeats allowed in the Schedule; or
- on the same day or within the next four days (e.g., if a pharmaceutical benefit is supplied on a Monday, it cannot be supplied again to that patient until the next Saturday) in the case of other benefits.

Exceptions to this are:

- when a PBS prescription is endorsed with the words 'Regulation 24' or 'hardship conditions apply' (see below under 'Regulation 24'); and
- If a pharmacist believes a repeat supply is needed without delay for the treatment of the person, or a previous supply has been destroyed, lost or stolen. In this case, the pharmacist can provide another supply but must write 'immediate supply necessary' and sign the PBS prescription.

A pharmacist can supply an alternative pharmaceutical benefit without reference to the prescriber, provided that:

- the PBS prescription does not indicate that only the pharmaceutical benefit prescribed is to be supplied (ie substitution is not permitted); and
- the Schedule states that the prescribed benefit and the substitute benefits are equivalent; and
- supply of the substitute benefit does not contravene relevant State/Territory law; and
- the substitute benefit is a listed brand in the Schedule.

Pharmacists must heed State/Territory laws when supplying drugs listed as narcotic, specified or restricted in legislation of the particular State or Territory.

What to do if the Schedule changes

If an item or brand is deleted from the Schedule, it *cannot* be supplied as a pharmaceutical benefit from the date the deletion takes effect – regardless of whether the PBS prescription was written before this date. This includes repeat authorisations. (Special conditions applying to RPBS prescriptions are detailed in the RPBS Explanatory Notes.)

However, if restrictions on the prescribing of a pharmaceutical benefit change, or the maximum quantity or number of repeats is altered in the Schedule, valid PBS prescriptions written before the date of effect of the change *may* still be supplied as pharmaceutical benefits, under the conditions applying at the date of prescribing.

Suspected forgery

Pharmacists should take all reasonable steps to satisfy themselves that all items on a PBS prescription were written by a medical practitioner, a dentist, an optometrist, a midwife or a nurse practitioner.

Regulation 24

This regulation allows pharmacists to supply a pharmaceutical benefit and all of its repeats at the one time.

The PBS prescription must be endorsed by the medical practitioner, midwife or nurse practitioner with the words 'Regulation 24' if it is an item under the PBS, or 'hardship conditions apply' if it is being supplied under the RPBS. (For more information see under '2. Prescribing Medicines ... Regulation 24'). Regulation 24 does not apply for supply of pharmaceutical benefits on optometrist prescriptions.

Repeat authorisations

When a PBS prescription calls for repeat supplies, the pharmacist shall prepare a Repeat Authorisation Form, except when the PBS prescription is marked 'Regulation 24'.

The repeat may be requested on a standard PBS prescription, an authority PBS prescription or an Authority to Prescribe Form, or on an earlier repeat authorisation. In the latter case, it must come with the duplicate PBS prescription, or in the new format, the "patient/pharmacist copy".

Preparing Repeat Authorisation Forms

A Repeat Authorisation Form must show:

- the category of benefit (concession or general) – by placing a cross (x) in the relevant box;
- the patient's name and full address;
- in the case of repeats authorised on authority PBS prescriptions, the authority prescription number;
- details of the original PBS prescription stating the item, form, strength, quantity and directions;
- if substitution has occurred, the name of the brand actually supplied;
- for the first supply, the pharmacy name, address and approval number, the date of the original PBS prescription and the allotted PBS prescription identifying number;
- for subsequent supplies, the pharmacy approval number, and the date and PBS prescription number of the original prescription;
- the number of times the item is to be repeated and the number of times it has been supplied;
- the name and pharmacy approval number of the pharmacist issuing the repeat authorisation; and
- the date of supply.

When a repeat authorisation is prepared for any further repeats or deferred supply, a pharmacist must attach the duplicate copy of an old format PBS prescription, or the patient/pharmacist copy of a new format PBS prescription, and give both to the patient at the time of supply.

Repeat authorisations for injectables and solvents

Where an injectable pharmaceutical benefit requires a solvent, both items should be treated as one pharmaceutical benefit. If repeats are needed, only one repeat authorisation is to be prepared. Details of the injectable and the solvent should appear in the space provided for the 'original prescription transcription'.

Repeat authorisations for deferred supply

When a PBS prescription orders a number of pharmaceutical benefit items, but the patient does not need all of the items at the same time, a separate repeat authorisation for each deferred item must be prepared. The words 'original supply deferred' should be indicated across the relevant item on the original PBS prescription, its duplicate, and on the repeat authorisation.

Deferred items must not be claimed on the original PBS prescription.

The Repeat Authorisation Form when it is used for a deferred supply, is issued in the same way as normal repeat authorisations except that:

- '0' is to be inserted in the space for 'no. of times already dispensed'; and
- if no repeats are ordered, '0' is to be inserted in the space for 'no. of repeats authorised'.

Supplying a benefit on a deferred supply repeat authorisation is to be treated as if it is the first time of supply. If repeats are directed, the normal procedure for repeat authorisations applies. Details of the pharmacy at which the deferred supply was authorised are to be written onto subsequent repeat authorisations.

Authority PBS prescriptions

If a pharmacist is presented with an authority PBS prescription and is not sure if it has been approved, he or she should contact Medicare Australia. Please note that Medicare Australia will not provide clinical information.

If the authority PBS/RPBS prescription is for an Authority required (STREAMLINED) item the pharmacist should ensure that the prescriber has written the four digit streamlined authority code on the prescription, this enables the pharmacist to supply the item as a PBS benefit.

Urgent cases

In urgent cases and where State/Territory law allows, pharmacists can supply a pharmaceutical benefit to a person without a PBS prescription, provided details of the prescription are given by the prescriber via telephone or other means. The prescriber must then forward the written PBS prescription and duplicate to the pharmacist within **seven days of the date of supply**.

Where a pharmaceutical benefit needs prior approval from Medicare Australia or the DVA, the prescriber must obtain approval and then advise the pharmacist of the PBS prescription and approval details. Only an original supply can be provided in this manner, not repeats.

Receipts

A person receiving a pharmaceutical benefit item must sign and date a receipt for it. If the person is not the patient, that person must also endorse the PBS prescription or repeat authorisation with his/her address. A receipt cannot be obtained until supply of the benefit has been made.

If a pharmaceutical benefit has to be sent through the post, by rail, or by other means, and a receipt is not practical, the pharmacist must certify on the PBS prescription or repeat authorisation that the benefit has been supplied, and write the date of supply and details of how it was sent. For example, if a pharmaceutical benefit is mailed to a patient on 1 April 2008, the pharmacist should write: "Certified supplied – mailed to patient 1 April 2008 (name of pharmacist) (signature of pharmacist) (date of certification)".

If an item is supplied in an urgent case, or to a person who cannot read or write, the pharmacist should sign and date a statement on the PBS prescription or repeat authorisation, stating the item has been supplied and the date on which it was supplied, and explaining why there is no receipt. For example, if a pharmaceutical benefit is supplied to a patient with a broken arm on 1 May 2008, the pharmacist should write: "Certified supplied 1 May 2008 – patient has a broken arm and is unable to sign (name of pharmacist) (signature of pharmacist) (date of certification)".

Only the pharmacist approved to supply pharmaceutical benefits can certify supply.

Emergency drug supplies

Pharmacists may supply certain pharmaceutical benefit items free of charge to medical practitioners or other authorised prescribers for emergencies if they receive an Emergency Drug Supply Order Form in duplicate, signed by the medical practitioners or other authorised prescriber.

Pharmacists must be satisfied the form was completed by a medical practitioner or other authorised prescribers and includes the medical practitioner's or other authorised prescriber's name and address. If a pharmacist does not know the medical practitioner or other authorised prescriber, he/she should confirm the medical practitioner's or other authorised prescriber's registration and endorse this on the back of the form.

For more information about emergency supplies see under 2. Prescribing Medicines ... Emergency drug supplies'.

4. Patient Charges

Type of patient

There are two types of PBS beneficiaries, general patients, who hold a Medicare card and concessional patients who hold a Medicare card and one of the following:

- Pensioner Concession Card
- Commonwealth Seniors Health Card
- Health Care Card
- Repatriation Health Card for All Conditions (gold) — concessional patients under RPBS
- Repatriation Health Card for Specific Conditions (white) — only regarded as concessional patients for RPBS prescriptions unless they hold a separate entitlement from Centrelink, otherwise they are general patients

- Repatriation Pharmaceutical Benefits Card (orange) — concessional patients under RPBS
- Safety Net Concession Card or Safety Net Entitlement Card — issued by Medicare Australia.

Concessional patients are recognised by public hospitals in all States and Territories apart from South Australia (where DVA beneficiaries are treated as general patients) and New South Wales (where holders of a white DVA card are treated as general patients).

Under the Reciprocal Health Care Agreements, visitors from participating countries (see the introduction of this section for the list of countries) are treated as general patients and do not have concessional entitlements. To receive pharmaceutical benefits these visitors may need to present a temporary Medicare card or their passport. Pharmacists should contact Medicare Australia if they have enquiries about these arrangements.

Establishing entitlement

PBS prescription forms supplied by Medicare Australia have spaces provided for details of a patient's entitlement status. Anyone can enter this information, which must include:

- a cross (x) in the appropriate box to indicate the level of patient contribution;
- the complete Medicare number (including individual reference number) or complete Veteran file number on the card; and
- if applicable, the complete concession number on the card.

The person who signs the receipt for pharmaceutical benefits also accepts responsibility for the validity of the entitlement information on the PBS prescription.

All PBS prescriptions must have a Medicare or Veteran file number. All concessional PBS prescriptions must have a concession number. However, it is not necessary for the Medicare (Veteran file) or the concession number to be endorsed on the PBS prescription if it is included in the electronic prescription details supplied by a pharmacist who is using the Claims Transmission System.

What to charge

Patient contribution

Under the PBS, the maximum cost for a pharmaceutical benefit item at a pharmacy is \$34.20 for general patients and \$5.60 for concessional patients, plus any applicable special patient contribution, brand premium or therapeutic group premium. General patients who have reached the safety net threshold (see details under '5. The Safety Net Scheme') may receive pharmaceutical benefits at the concessional rate, plus any applicable special patient contribution, brand premium or therapeutic group premium.

Patients who have a Safety Net Entitlement Card (see details under '5. The Safety Net Scheme') may receive PBS items free of charge, except for any applicable special patient contribution, brand premium or therapeutic group premium.

The contribution rate for general patients as outpatients at public hospitals in most of Australia is \$27.40. The exceptions are in Queensland and in hospitals participating in the pharmaceutical reforms where they pay the safety net value of an item listed in the Schedule (see details under '5. The Safety Net Scheme'), or up to the general co-payment amount for items not listed in the Schedule. The public hospital pharmaceutical reforms enable participating public hospitals to prescribe and supply pharmaceutical medication from the PBS to outpatients and patients upon discharge. A range of chemotherapy drugs is also available for day-admitted and non-admitted chemotherapy patients.

The contribution rate for concessional patients in all public hospitals is equal to the concessional co-payment amount.

The supply of a pharmaceutical benefit or a Repatriation pharmaceutical benefit to a patient is GST-free. Goods and services tax must not be included in the price charged to a patient for the supply of a PBS or RPBS script.

It is the patient's responsibility to pay any charge lawfully imposed by an approved pharmacist or supply may be refused.

The patient contribution rates are adjusted on 1 January each year in line with inflation.

Patient contributions for early supply of some PBS medicines

Prescriptions for some PBS and RPBS pharmaceutical benefits are not eligible for safety net benefits if re-supplied within 20 days of a supply of the same pharmaceutical benefit for the same person. This is known as the 'Safety Net 20 day rule' and came into effect on 1 January 2006.

Where a prescription is subject to the Safety Net 20 day rules:

- the patient contribution does not count towards the Safety Net, and

- after the Safety Net threshold is reached, the usual patient co-payment amount for the corresponding entitlement level (not the Safety Net amount) applies.

For example: The payment for such a prescription for a patient with a Safety Net Entitlement Card would be the concessional co-payment amount — not free. For a general patient with a Safety Net Concession Card, the usual general co-payment amount would apply — not the concessional amount.

The Safety Net 20 day rule does not apply to PBS/RPBS prescriptions originating from hospitals or day hospital facilities.

Special patient contributions, brand premiums and therapeutic group premiums

A special patient contribution is payable for a pharmaceutical benefit when a supplier will not supply it at the benchmark price. Any extra charge for a higher priced benefit is paid by the patient, together with their usual patient contribution. Other than for bleomycin sulfate, exemptions on medical grounds are available, but must be granted by Medicare Australia. For RPBS special patient contribution arrangements see the RPBS Explanatory Notes.

Under the brand premium arrangements, reimbursement to pharmacists is based on the lowest-priced brand. Any extra charge for a higher priced brand is paid by the patient, together with their usual patient contribution.

Under the therapeutic group premium arrangements, reimbursement to pharmacists is based on the lowest priced benefit items within identified therapeutic groups. Any extra charge for a higher priced benefit is paid by the patient, together with their usual patient contribution. Exemptions on medical grounds are available, but must be granted by Medicare Australia.

Special patient contributions, brand premiums and therapeutic group premiums apply to maximum quantities. When a quantity is less than, or — on an authority or 'Regulation 24' PBS prescription — more than, the maximum, the contributions or premiums will be a factor of the maximum quantity, using standard pricing rules.

There are separate arrangements for PBS prescriptions in certain public hospitals. To obtain pharmaceutical benefits under these arrangements a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner providing medical treatment or a midwife providing midwifery treatment or a nurse practitioner providing nurse practitioner treatment in a participating public hospital may prescribe PBS subsidised medication. Victoria, Queensland, South Australia, Western Australia and the Northern Territory have these arrangements.

Solvents

Where a solvent is prescribed as a part of a pharmaceutical benefit, only one patient contribution is charged.

Increased quantities

Where a prescriber has written an authority PBS prescription for a quantity greater than the maximum, the patient contribution should be made for each supply of the increased maximum quantity.

Regulation 24

For 'Regulation 24' PBS prescriptions, a pharmacist should charge the usual patient contribution for the original and for each repeat quantity needed to make up the total supply (plus any applicable special patient contribution, brand premium or therapeutic group premium, for the original and each repeat quantity in the total supply).

After hours

A pharmacist may charge an extra fee if supplying a PBS item outside normal trading hours. This charge is paid by the patient and does not count towards the safety net.

Delivery

A charge can be added for delivering pharmaceutical benefits from the pharmacy. This charge does not count towards the safety net. For RPBS delivery arrangements refer to the RPBS Explanatory Notes.

5. The Safety Net Scheme

The PBS safety net protects patients and their families requiring a large number of PBS or RPBS items. For the purposes of the scheme, the family includes the person:

- the partner or de facto partner;
- children under the age of 16 who are in the care and control of the person; or
- dependent full-time students under the age of 25.

The scheme requires pharmacists, on request by patients, to record the supply of PBS and RPBS items on prescription record forms. When a patient reaches the Safety Net threshold within a calendar year, they qualify to receive PBS or RPBS

items at a cheaper price or free of charge for the rest of that year. Any applicable special patient contributions, brand premiums or therapeutic group premiums must still be met by the patient.

The safety net threshold is reached by accumulating eligible patient contributions for PBS prescriptions supplied through community pharmacies and private hospitals and for out-patient medication supplied by public hospitals.

Pharmaceutical benefits (including authority items) can only be counted towards the safety net threshold when prescribed and supplied according to PBS conditions. A medicine supplied by a pharmacist not approved to supply pharmaceutical benefits cannot count towards the safety net.

Prescriptions for some pharmaceutical benefits are not eligible for safety net arrangements if re-supplied within 20 days of supply of the same item for the same person and the patient contribution cannot count towards the safety net (see also details under '4. Patient Charges' and '7. How Pharmacists Claim Reimbursement'). This does not apply to out-patient medications in public hospitals or to any prescriptions originating from a hospital or day hospital facility.

There are separate arrangements for PBS prescriptions in certain public hospitals. To obtain pharmaceutical benefits under these arrangements a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner providing medical treatment or a midwife providing midwifery treatment or a nurse practitioner providing nurse practitioner treatment in a participating public hospital may prescribe PBS subsidised medication. Victoria, Queensland, South Australia, Western Australia and the Northern Territory have these arrangements.

Safety net thresholds

There are two safety net thresholds. The general patient safety net threshold is currently \$1317.20. When a person and/or their family's total applicable co-payments reach this amount, they may apply for a safety net concession card and pay the concessional co-payment amount of \$5.60 plus any applicable premium for pharmaceutical benefits for the rest of that calendar year.

The concessional safety net threshold is \$336.00 (this also applies to gold, white or orange card holders under the RPBS). When a patient and/or their family's total applicable co-payments reach this amount, they may apply for a safety net entitlement card and may receive pharmaceutical benefits free of charge (except for any applicable premium) for the rest of that calendar year.

Brand premiums, therapeutic group premiums and special patient contributions do not count towards the safety net thresholds.

The safety net thresholds are adjusted on 1 January each year in line with inflation.

Safety net cross-over arrangements

Some patients and/or members of their families will change between general patient and concessional patient status during a calendar year. Patients should apply for the safety net card appropriate to their status at the time they apply.

Concessional patients who were previously general patients can apply for a safety net entitlement card when they reach the concessional safety net threshold. In this case, any pharmaceutical benefits previously supplied at the general co-payment rate in that calendar year will be counted at the concessional rate per item.

General patients who were previously concessional patients can apply for a safety net concession card when they reach the general safety net threshold. In this case, any pharmaceutical benefits previously supplied at the concessional rate in that calendar year will be counted at the concessional rate per item.

In the case of families where one parent holds a concession card and other family members are general patients, the family can choose to apply for either a safety net entitlement card or a safety net concession card.

To receive a safety net entitlement card, all pharmaceutical benefits (including general pharmaceutical benefits) are counted at the concessional rate per item until the concessional threshold is reached. To receive a safety net concession card, general pharmaceutical benefits are counted at the general co-payment rate per item and concessional pharmaceutical benefits at the concessional rate per item, until the general safety net threshold is reached.

White DVA card holders may either be general or concessional patients (depending on their Centrelink entitlements). If they are receiving treatment for a specific disability accepted by the DVA, they are also supplied with specified items under the RPBS at the concessional rate per item. Therefore, these patients are encouraged to maintain a concessional prescription record form, plus a general prescription record form for items not covered under the RPBS.

White card holders may choose at any time to count contributions made at the general level towards the concessional safety net threshold and receive credits equal to the concessional co-payment amount for each pharmaceutical benefit purchased. Alternatively, white card holders can count contributions at the concessional level towards the general safety net, and receive credits equal to the concessional co-payment amount for each pharmaceutical benefit purchased.

Gold or orange DVA card holders may receive all of their prescription items under the RPBS, and only pay the concessional co-payment amount for each item.

Dependants of white, gold or orange card holders are treated separately and may be either general patients or concessional patients. Their prescriptions may be included in the cross-over arrangements.

Recording PBS prescriptions

There are two types of prescription record forms to record PBS prescription items. A blue form, used for items obtained at community pharmacies and available from community pharmacies, Medicare offices and Medicare Australia; and a grey form, used by out-patients who pay for items at public hospital pharmacies and available from hospital out-patient departments or Medicare Australia.

Patients should record their general or concessional status on the prescription record form, enter their Centrelink, DVA and/or Safety Net Concession/Entitlement Card number, and list family members covered. General patients must also record their Medicare number when applying for a safety net concession card.

Details to be entered on the form by the pharmacist are:

- date of supply;
- PBS/RPBS code number of the item (for community pharmacies only);
- the safety net value of the item (for community pharmacies only);
- pharmacist's approval number (for community pharmacies only);
- item identification — medicine code, name of medicine or abbreviation (for public hospitals only);
- hospital charge (for public hospitals only);
- hospital safety net number (for public hospitals only); and
- signature of the authorised person making the entry.

Community pharmacists should record in the 'safety net value' column:

- the patient contribution when it is less than the PBS dispensed price; or
- the safety net value shown in the Schedule, or any lesser amount charged, if the PBS dispensed price is less than or equal to the patient contribution. The pharmacist may discount the price for these items.

Some computer software suppliers provide a special label to record this information on the prescription record forms. Some suppliers also provide a computer printout as a prescription record form.

The patient is responsible for maintenance and storage of their prescription record form. However, it may be kept in the pharmacy. A person (or family) may have more than one prescription record form.

Hospital prescription record forms

Items to be recorded on hospital prescription record forms must be approved by the hospital's pharmaceutical advisory committee and may be listed on a hospital's formulary (a list of pharmaceutical items approved by the committee for the treatment of particular illnesses), or authorised on a patient-by-patient basis.

Multi-item prescription forms

If a patient submits a multi-item PBS prescription form, which would take the total co-payments past the safety net threshold, any items in excess are treated as entitled items once a safety net entitlement/concession card is issued.

Excess items should be treated as 'deferred supply' items.

For example, if a family has a new PBS prescription for three items and the first takes the family up to the threshold, then this item should be supplied at the general rate. If the second item takes the family over the threshold, the pharmacist should then issue a safety net concession card and supply both this and the third item at the concessional rate. This involves the deferral of two items, recording the safety net concession card number, and the subsequent supply of these items.

Qualifying PBS prescriptions

A PBS prescription should be supplied at the concessional rate or free of charge plus any applicable premium, when the safety net value or hospital charge for that PBS prescription takes the total co-payments over the qualifying amount for a safety net entitlement/concession card.

Lost prescription record forms

If a prescription record form has been lost, stolen or destroyed, a pharmacist may prepare a duplicate copy, but is under no obligation to do so.

Retrospective entitlement and patient refunds

Responsibility for claiming entitlements rests with the patient. If items recorded on a prescription record form have exceeded the safety net threshold, the cost of those items in excess of the limit cannot be refunded by a pharmacist.

However, if the patient failed to apply for a safety net entitlement/concession card on reaching the safety net threshold they should write to Medicare Australia and provide copies of pharmacy accounts or a signed statement from the pharmacist giving the date of supply, description and cost of items supplied and paid for. A copy of the relevant prescription record form should also be provided. If these are not available, the patient should give the name of the pharmacy where the card was issued and the number on the card so that Medicare Australia can locate the prescription record form in its records. Cash refunds are not available. Medicare Australia contact details are provided in the 'Addresses — Medicare Australia' part of the Schedule.

If the patient cannot satisfy a pharmacist that they have a current entitlement and is charged the general patient price, the pharmacist should issue the patient with a receipt and a claim form (provided by Medicare Australia). The patient can then obtain a refund via Medicare offices or PBS processing centres. RPBS prescription refunds are paid at DVA State offices.

Medicare Australia can only pay refunds for PBS items supplied through approved pharmacies. Refunds for hospital supplied items should be referred to the relevant hospital or health department. Refunds cannot be made where the patient was charged the general or concessional amount instead of the safety net concessional or safety net entitlement amount as a result of the safety net 20 day rule. Receipts for prescriptions where the safety net 20 day rule has applied must include 'SN20DR' to indicate the reason for the amount charged.

There are separate arrangements for PBS prescriptions in some public hospitals. To obtain pharmaceutical benefits under these arrangements a patient must attend a participating public hospital and be a discharge patient or non-admitted patient. Only a medical practitioner providing medical treatment or a midwife providing midwifery treatment or a nurse practitioner providing nurse practitioner treatment in a participating public hospital may prescribe PBS subsidised medication. Victoria, Queensland, South Australia, Western Australia and the Northern Territory have these arrangements.

Applying for a Safety Net Entitlement/Concession Card

Once the safety net threshold has been reached, the person covered by a prescription record form may complete the application and declaration to get a safety net entitlement/concession card. Please note that software packages that produce computer generated applications must be approved by Medicare Australia.

If the card is issued to a dependent child or student, it should be in the name of a parent.

When issuing entitlement/concession cards, pharmacists do not have to check all prescription record form details. However, they should ensure each entry has been signed and that the prescription record form total qualifies the patient for the relevant safety net card.

When appropriate the pharmacist should check that the patient's Medicare card number is on the prescription record form.

Issuing a Safety Net Entitlement/Concession Card

When satisfied that the individual or family is entitled, the pharmacist should issue the next blank safety net entitlement/concession card with the following details:

- the names of family members covered. If there are more than eight family members, a second card should be issued listing the card holder and family members not listed on the first card. The prescription record form has space to record that two cards have been issued, and
- the two-character code to indicate the relationship to the card holder. Applicable codes are:
 - SP - partner;
 - DC - child under 16 years; and
 - DS - dependent full-time student under 25 years.

The pharmacist should be satisfied that only family members are listed on the card. The unused space on the card should be ruled through to prevent extra names being added. The sticky label from the safety net entitlement/concession card, pre-printed with the card number, should be attached to the prescription record form. The pharmacist should sign and stamp each prescription record form with the pharmacy stamp and enter the card issue details on a safety net — claim for payment form.

Issuing supplementary cards

A pharmacist may give a card holder a supplementary card for a partner or dependant only at the time the original card is issued. The duplicate card should be recorded in the additional box on the prescription record form.

Later requests for supplementary cards and requests to add a new family member to the original card are to be referred to Medicare Australia.

Notification to Medicare Australia and claim for payment

Payment for issuing a safety net entitlement/concession card is made after the safety net — claim for payment form is sent to Medicare Australia, no later than one month after a card is issued.

Each form must be accompanied by all supporting documentation (prescription record form and cancelled or void safety net entitlement/concession cards).

Payment will not be made for void cards.

Lost Safety Net Entitlement/Concession Cards

When a card has been lost, damaged, stolen or destroyed, a pharmacist cannot re-issue a person with a replacement card. The original card holder (or partner) must apply to Medicare Australia.

Pharmacy record of issued cards

A record of all cards issued must be kept at the pharmacy from which the pharmacist is approved to supply pharmaceutical benefits. The duplicate ('bookfast') copy in the safety net — claim for payment book is provided for this purpose.

6. Medicare Australia Entitlement Checks

General Patients

Medicare Australia validates a patient's entitlement to pharmaceutical benefits by checking Medicare and/or Veteran file numbers in pharmacist's claims. If a number is not recorded correctly, a patient cannot be identified against Medicare Australia's Pharmaceutical Benefits Entitlement File and entitlement cannot be established.

If the Medicare or Veteran file number provided in the pharmacists' claims is incorrect or the number and the name supplied do not match Medicare Australia records to enable patient identification, an appropriate warning or rejection code will be returned to the pharmacy. These notifications of missing or incorrect Medicare or Veteran file numbers are provided to pharmacists in their reconciliation statement produced after the claim period has been paid by Medicare Australia.

Special numbers are available for use in certain circumstances for eligible people who are unable to provide a Medicare number.

Concessional Patients

Medicare Australia routinely validates a patient's entitlement to free or concessional benefits by checking concessional numbers in pharmacist's claims. If a number is not recorded correctly, a patient cannot be identified against Medicare Australia's Pharmaceutical Benefits Entitlement File and entitlement cannot be established.

When a number is found to be from a card which was incorrect, expired at the time of supply or entitlement was withdrawn, warning or rejection codes will be returned to the pharmacy to assist with validation of concessional entitlement in relation to future claims from the same patient.

Entitlement checking procedures

General Patients

Once a pharmacist has been notified by Medicare Australia of an incorrect Medicare or Veteran file number he/she should correct the number for future claims by:

- updating his/her system to reflect the correct number provided by Medicare Australia (if patient consent to do so has been obtained); or
- speaking to the patient; or
- obtaining patient consent and calling Medicare Australia on the Improved Monitoring of Entitlements (IME) (132 290 — select option 1).

If the patient presents a Medicare card that appears correct, but according to Medicare Australia is not a valid number, or not a valid number for that person, a pharmacist may use a special number. A photocopy of the card, or a form must accompany the use of this number. The form is available on Medicare Australia's website or by calling 132 290.

Concessional Patients

Once a pharmacist has been notified by Medicare Australia of an incorrect concessional entitlement number, he/she should view the entitlement card to confirm the entitlement number, and start and end dates, when the patient next presents a PBS prescription.

Step by step

Pharmacists should take the following steps where concession entitlement does not appear to be valid or current:

- Re-confirm entitlement with the cardholder/customer;
- Contact Medicare Australia on 132 290, with consent, to confirm the cardholder/customer concession status;
- If Medicare Australia advises that the cardholder/customer is concessionally entitled to receive the PBS medicines on that day, supply the prescription as a concessional entitlement;
- If Medicare Australia advises that the cardholder/customer is not concessionally entitled to receive the PBS medicines on that day, supply as a general prescription. Provide the customer with the information sheet "Your entitlement card" which explains entitlement checking to the customer and the steps they can follow if they are concessionally entitled.

7. How Pharmacists Claim Reimbursement: Information Required

Medicare Australia uses a computerised system for pricing PBS prescriptions, repeat authorisations and emergency drug supply orders, and for calculating claims.

The payment system is designed to pay pharmacists correctly for the pharmaceutical benefits they supply. It is essential instructions are followed carefully and that each document includes all relevant information. Accurate and complete data ensures claim payment is not delayed.

PBS Prescription identification

Pharmacists must include certain information on each PBS prescription sent in for claim, as specified below. It is important that this information is entered correctly and in the right place on the PBS prescription. This information will be included in a sticker produced by pharmacy software.

The sticker should be placed on the extreme left front of a PBS prescription, opposite each item being claimed. It must not obscure any details written by the prescriber. Most prescribers use PBS prescriptions, which have space for the sticker. If a sticker is not used, a PBS prescription identification stamp can be used or the information can be written in the same place, and in the same order.

Pharmacists should avoid writing over, or placing the sticker over, the prescriber number pre-printed on PBS/RPBS prescriptions, or the prescriber number box on PBS dental and optometrist, midwife and nurse practitioner prescriptions.

The sticker is not necessary for current repeat authorisation, emergency drug supplies, or for old style authority PBS prescription and authority to prescribe forms, as they have printed spaces for the necessary details. However, it is required for the new format authority PBS prescription forms.

The following information should be entered next to the appropriate letter on the sticker or stamp:

- 'S' — the serial number for the claim
- 'A' —
 - a. the price claimed for pricing elected PBS prescriptions, exceptional PBS prescriptions and RPBS non-scheduled prescriptions (see under 'Extemporaneously-prepared pharmaceutical benefits not listed in the Standard Formulae List' for explanations of pricing elected PBS prescriptions and exceptional PBS prescriptions); and/or
 - b. confirmation that the PBS prescription is endorsed 'Regulation 24' or the RPBS prescription is endorsed 'hardship conditions apply'; and/or
 - c. a claim for a glass dropper bottle where applicable; and/or
 - d. any clarification of the prescription which will assist Medicare Australia payment processing.
- 'No.' — the PBS prescription identifying number.

Serial numbers

PBS prescription, repeat authorisation, authority PBS prescription, and emergency drug supplies forms submitted in each claim must bear consecutive serial numbers starting with:

- 1 – for emergency drug supplies;
- 1 – for general benefits;
- C1 – for concessional and Safety Net Concession Card benefits;
- E1 – for Safety Net Entitlement Card benefits; and
- R1 – for RPBS benefits.

Each serial number should also be noted on any document kept by the pharmacist for record purposes.

Each emergency drug supply item should be given a serial number, e.g., if there are five items on the first form in the claim, the first item on the second form in the claim will start with the serial number 6.

For prescriptions subject to the Safety Net 20 day rule, the serial number corresponds to the resulting payment category for the pharmaceutical benefit as supplied, not the patient's entitlement category.

Repeat authorisations for authority PBS prescriptions

When a benefit is supplied on a repeat authorisation which needed an authority PBS prescription, the serial number must be prefixed with the letter 'A' for a general benefit; 'AC' for a concessional benefit or a benefit supplied to a Safety Net Concession Card holder; 'AE' for a Safety Net Entitlement Card holder; or 'AR' for a RPBS benefit.

Repeat authorisations for deferred supply

When a benefit is supplied on a repeat authorisation prepared for deferred supply, the serial number must be prefixed with the letter 'D' for a general benefit; 'DC' for a concessional benefit or a benefit supplied to a Safety Net Concession Card holder; 'DE' for a Safety Net Entitlement Card holder; or 'DR' for a RPBS benefit.

Injectable item ordered with a solvent

When both an injectable item and a solvent are to be supplied, only one serial number is used. This number should be placed on the left hand side of the prescription, opposite the injectable item.

Dropper containers

Dispensed prices for extemporaneously-prepared eye drops, ear drops and nasal instillations include the price of a polythene dropper container. However, if a glass dropper container is supplied, payment should be claimed by writing 'glass bottle' in box 'A' of the stamp.

Extemporaneously-prepared pharmaceutical benefits not listed in the Standard Formulae List

When a formula is not listed on the Standard Formulae List, the PBS prescription is paid at an average of 10 g/mL rate for the type of preparation, unless the pharmacist elects otherwise. A pharmacist may price an exceptional PBS prescription, or elect to price all non-pre-priced extemporaneous PBS prescriptions.

PBS prescriptions paid on an average price basis

If the PBS prescription is to be claimed as an exceptional PBS prescription, the pharmacist should write details of the formula supplied on the PBS prescription or repeat authorisation form; price the PBS prescription in accordance with the pricing principles (as detailed in '9. Pricing PBS Prescriptions'); and enter the calculated price on the sticker.

An exceptional PBS prescription is for an extemporaneously-prepared pharmaceutical benefit that is not included in the Standard Formulae List and for which the price of the ingredients (based on basic pricing rules) is twice or more than the recovery price of the ingredients calculated on an average price basis. Further information on pricing PBS prescriptions can be accessed from the book let titled *Explanation of Current Pricing* on the Medicare Australia's website at www.medicareaustralia.gov.au (PBS publications for Health Care Providers).

Pricing non-pre-priced extemporaneous preparations

Pharmacists should notify Medicare Australia when they elect to price non-pre-priced extemporaneous preparations. Each PBS prescription should be priced in accordance with the pricing principles and that price entered on the sticker.

RPBS prescriptions for items not included in either the PBS or RPBS Schedule

When a prescription for a RPBS patient is for an item not included in either the PBS or the RPBS Schedule, the price claimed should be entered on the sticker. Full details on pricing and availability of such items under the RPBS are set out in the RPBS Explanatory Notes.

Payment to Pharmacists for Dispensing Premium-free Substitutable Medicines

Premium Free Dispensing Incentive payments will commence for eligible PBS listed products dispensed from 1 August 2008. Premium Free Dispensing Incentive payments will be available to approved suppliers to dispense a substitutable, premium-free medicine. The payment will be available only for PBS items which attract a Government subsidy. This includes PBS items supplied to DVA entitled consumers.

A number of conditions and criteria apply to receive this payment. Scripts will be assessed for validity and the Premium Free Dispensing Incentive payment will be paid by Medicare Australia. Further information on this payment can be found on the Medicare Australia website at:

<http://www.medicareaustralia.gov.au/provider/pbs/pharmacists/reforms.shtml#dispensing>

8. How Pharmacists Claim Reimbursement: Documents to be Submitted

A claim for pharmaceutical benefits consists of:

- the original and duplicate of a completed Claim for Payment Form;
- the original orders for emergency drug supplies in a separate bundle;
- the originals of all old format PBS prescriptions and authority PBS prescriptions, the Medicare Australia/DVA copies of new format PBS prescriptions and authority PBS prescriptions, and all repeat authorisations, separated into four bundles for benefits supplied to the general public; concessional beneficiaries/Safety Net Concession Card holders; Safety Net Entitlement Card holders and RPBS patients.

PBS prescriptions in each bundle should be in serial number order, with serial number 1 at the top of the bundle.

PBS prescriptions subject to the Safety Net 20 day rule are bundled according to the resulting payment category. For prescription forms with multiple PBS items, where the Safety Net 20 day rule would result in different payment categories for different items, dispensing via 'deferred supply' should be used where necessary to allow all items to be included in the correct bundles.

PBS prescriptions in the wrong bundle may be returned to the pharmacist for clarification. If appropriate, they can be resubmitted in the correct bundle in the next claim period.

Completing the claim form

The claimant's name, address of the pharmacy from which the pharmacist is approved to supply pharmaceutical benefits, approval number, and claim period number should be entered on the Claim for Payment Form. These details should match the latest written information held by Medicare Australia, or payments can be delayed while clarification is sought.

The claim period number should state how many claims have been submitted so far in a calendar year, e.g., the sixth claim submitted by an approved pharmacist in 2005 should have a claim period number of 0506.

The first and last serial numbers given to items in each bundle are to be entered on the Claim for Payment Form.

A total claim amount is not required – this will be calculated by Medicare Australia after the PBS prescriptions have been individually priced.

The declaration must be signed by the pharmacist approved to supply pharmaceutical benefits, unless he/she has made arrangements through Medicare Australia for another pharmacist to sign it.

Lodging claims

A claim may be lodged at any time during the month at the relevant Medicare Australia State office. Unless other arrangements have been made with Medicare Australia, the following conditions apply:

- only one claim period can exist and only one claim can be lodged per month;
- the claim period shall cover pharmaceutical benefits supplied during one month; and
- the claim shall be sent within 30 days from when the benefits were supplied.

Claims for pharmaceutical benefits supplied over 18 months earlier may not be accepted for computer processing. Pharmacists with such claims should contact Medicare Australia.

Reconciliation statements

As mentioned earlier, a pharmacist will receive a PBS reconciliation statement after a claim period has been processed. It provides details of each prescription for each brand of each pharmaceutical benefit item supplied in that claim period.

Reasons for non-payment of any item are coded, with the code numbers explained in the statement.

PBS prescriptions and repeat authorisations not accepted for payment will be returned, with the exception of PBS prescriptions with a dispensed price equal to or less than the patient contribution. Any other items on those PBS prescriptions that have been paid will have been cancelled.

If a PBS prescription was not accepted and can be re-submitted, it must be given a new serial number and included in a subsequent claim period.

If a PBS prescription is finally rejected for payment and a pharmacist is not satisfied with the decision, he/she may apply to the Administrative Appeals Tribunal for a review of that decision.

9. Pricing PBS Prescriptions

Pricing principles

The same pricing principles apply to all PBS prescriptions.

For ready-prepared pharmaceutical benefits, payment is made on the basis of the lowest-priced brand.

For a pharmaceutical benefit not listed as a ready-prepared item, and where a formulation title is stated but no formulary specified, payment is made on the basis of precedence given to formularies by State/Territory legislation.

Prices published in the Schedule do not include any component for goods and services tax (GST).

Further information on pricing PBS prescriptions can be accessed from the booklet titled *Explanation of Current Pricing* on the Medicare Australia's website at www.medicareaustralia.gov.au (PBS publications for Health Care Providers).

Pricing dates

Ready-prepared pharmaceutical benefits are priced on the first day of April, August and December for items supplied as from each of those days respectively.

Extemporaneously-prepared pharmaceutical benefits and containers are priced on the first day of May each year for items supplied as from the first day of August that year.

Pricing ready-prepared items

For maximum quantities

The price payable for a pharmaceutical benefit is shown in the Schedule against the item. The price is for the maximum quantity available.

If the prescription is for an injectable item and solvent, the price of each is added together, but only one dispensing fee is payable.

The maximum quantity of some pharmaceutical benefits, such as eye drops and oral suspensions, has been determined as a single pack corresponding to the manufacturer's pack. These packs cannot be broken, so if a PBS prescription calls for less, the maximum quantity should be supplied and claimed from Medicare Australia. Packs not to be broken are indicated by a double dagger (‡) in the Schedule.

For lesser quantities

For items where the standard pack is the same as the maximum quantity, and the pack can be broken, the price payable for a lesser quantity is established as follows:

- an amount equal to the dispensing fee, and if applicable the dangerous drug fee, is deducted from the benefit price as shown in the Schedule;
- to this new amount, a wastage percentage is applied, determined from the Wastage Factor Table;
- then the amount equal to the dispensing fee, dangerous drug fee (if applicable), and appropriate container fee, is added.

In no case shall the price for a broken quantity be more than the dispensed price of the Schedule's maximum quantity.

When a standard pack is not the same as the maximum quantity, the price of the pharmaceutical benefit concerned has an asterisk next to it and the standard pack rate is set out in Section 3 of the Schedule. The price payable for the quantity supplied is established by:

- applying the appropriate wastage table percentage to the standard pack rate;
- then adding an amount equivalent to the dispensing fee, the dangerous drug fee where applicable, and the appropriate container fee.

In no case shall the supply of a broken quantity, which is less than the item's maximum quantity, cost more than the dispensed price for the maximum quantity.

No container fee is payable when the quantity of pharmaceutical benefit supplied is more than the quantity contained in the standard pack.

Wastage table percentage

The following Wastage Factor Table is used to calculate the price payable for quantities supplied from the standard pack.

Wastage Factor Table

Column A - 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100

Column B - 10, 18, 26, 32, 38, 44, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98, 100

The appropriate wastage table percentage is as follows:

- the percentage of the amount supplied from the amount in the standard pack is determined; and
- where this percentage is the same as a percentage listed in Column A of the table, the percentage used is the figure shown in Column B; or
- where the percentage is not the same as a percentage in Column A, then the nearest upward percentage in Column A applies, and the percentage used is the figure in Column B.

For example, 24 tablets are supplied from a standard pack of 100. Thus 24 per cent of the number contained in the standard pack is supplied. As this percentage does not appear in Column A, the next higher (i.e., 25 per cent) is used. Reading down from 25 per cent to Column B, the wastage table percentage is found to be 38 per cent.

Pricing extemporaneously-prepared items

General

The price payable for supplying the maximum quantity of standard formula preparations is shown in the Standard Formulae List.

The following principles apply in determining prices of all pre-priced extemporaneous formulae on the list.

They also apply when a pharmacist elects to price extemporaneous PBS prescriptions outside the list, including exceptional PBS prescriptions.

The amount payable is the sum of:

- the recovery price of each ingredient as shown in the Drug Tariff;
- the price of the appropriate container as shown in the price section; and
- a dispensing fee as shown in the price section.

Pricing of ingredients

When the quantity dispensed is not specified in the Drug Tariff, the recovery price is as follows:

1. determine the basic pricing unit relative to the quantity dispensed by referring to the following table:

Quantity	Basic Pricing Unit
Up to and including 700 mg	100 mg price rate
Over 700 mg and up to and including 1 g	price as if 1 g
Over 1 g and up to and including 7 g	1 g price rate
Over 7 g and up to and including 10 g	price as if 10 g
Over 10 g and up to and including 80 g	10 g price rate
Over 80 g and up to and including 90 g	price as if 80 g
Over 90 g	100 g price rate

2. find the recovery price of the basic pricing unit by applying the following quantity divisors to the recovery price shown for the ingredient in the Drug Tariff:
 - 100 g price is 500 g price divided by 5, or 1 kg price divided by 10

- 10 g price is 100 g price plus 12.5 per cent divided by 10
 - 1 g price is 10 g price plus 25 per cent divided by 10
 - 100 mg price is 1 g price plus 25 per cent divided by 10
3. find the recovery price by multiplying the price of the basic pricing unit – as established in 2 – by the fraction that the quantity dispensed bears to the basic pricing unit.

For pricing purposes the quantity is to be taken to the next upward 50 milligrams or 0.05 millilitres.

The minimum recovery price for any ingredient is one cent. In other cases where a fraction of a cent occurs, the price is to be taken to the nearest cent (a half cent being taken up to the next cent).

In no case shall the recovery price for a quantity of an ingredient exceed the recovery price for a greater quantity of that ingredient.

Where liquids are purchased by weight, the recovery price includes the 'Specific Gravity Factor'.

Special pricing provisions apply to drugs marked '(a)' or '(b)' in the Drug Tariff.

For drugs marked '(a)', the pricing rules shown above apply to quantities up to the quantity listed in the Drug Tariff. Greater quantities are priced on a linear basis: the recovery price is ascertained by multiplying the fraction that the quantity dispensed bears to the quantity listed in the Drug Tariff by the price shown for the quantity listed.

Drugs marked '(b)' are packed sterile or are unstable, and all quantities are priced as if whole pack(s) were required. The recovery price is ascertained by multiplying the fraction that the quantity dispensed bears to the quantity listed in the Drug Tariff, taken to the next whole number, by the price shown for the quantity listed.

Pricing PBS prescriptions where extra ingredients are added to a formula

Where the vehicle is liquid and one or more solid ingredients are added, displacement of the liquid by the solid ingredients is disregarded for pricing purposes.

Containers

When a quantity is for more than the container sizes listed in this Schedule, payment will be made as if that quantity had been supplied in the minimum number of containers necessary to supply that quantity.

A double size container is allowed for bulk powders.

Special provisions for extemporaneous PBS prescriptions outside the Standard Formulae List

If a pharmacist elects to price extemporaneous PBS prescriptions outside the Standard Formulae List, there can be no variation for three months. This applies to all extemporaneously-prepared formulae not on the list, and includes both PBS and RPBS prescriptions.

If a pharmacist does not elect to price out these PBS prescriptions, he/she will be paid at an average reimbursement rate.

Under this system, payment is made on the basis of an average 10 g/mL rate applied to the category of preparation concerned, i.e., the price will be determined by multiplying the appropriate 10 g/mL rate by the number of 10 g/mL units supplied and adding container and dispensing fees. For example, an 80 mL mixture would be priced at eight times the average 10 mL rate for mixtures, with container and dispensing fee added.

The average 10 g/mL rate for each type of preparation is calculated monthly. It applies to PBS prescriptions supplied in the following month.

PBS prescriptions ordering a combination of standard formula preparations fall outside the scope of the Standard Formulae List and therefore are subject to this section.

Any variant to a formula included in the list (adding or deleting an ingredient or varying the dose) takes the formula dispensed outside the list.

When an ingredient is added to a standard formula and the recovery price for the standard formula plus additive under the average price system is less than for the standard formula alone, the pharmacist may have the PBS prescription priced as a basic standard formula item.

10. Miscellaneous

References

This Schedule identifies monographs of the British Pharmacopoeia, the British Pharmaceutical Codex, and the Australian Pharmaceutical Formulary and Handbook by the letters BP, BPC and APF respectively. References to all editions of the BPC and to earlier editions of the BP and APF also include the year of publication or the number of the edition.

Standards

Pharmacists can only supply under the PBS medicines which, or whose ingredients, conform to the standards of composition or purity prescribed. These standards are those specified in the *Therapeutic Goods Act 1989*.

Legislation

Copies of the *National Health Act 1953* and the *National Health (Pharmaceutical Benefits) Regulations 1960* are available from Government AusInfo shops in each capital city. The Act and the Regulations may also be accessed through the Attorney-General's Department website at www.comlaw.gov.au.

Nurse practitioner PBS prescribing

MEDICINES WHICH MAY BE PRESCRIBED BY AUTHORISED NURSE PRACTITIONERS

From 1 September 2010, nurse practitioners endorsed to prescribe under state or territory legislation can apply for approval as PBS prescribers (*authorised nurse practitioners*). Information for nurse practitioners to become authorised PBS prescribers is available from Medicare Australia.

The medicines listed for prescribing by authorised nurse practitioners are identified by 'NP' in the PBS Schedule. Nurse practitioners must not write PBS prescriptions for other medicines.

PBS prescribing is limited by a nurse practitioner's scope of practice, and state and territory prescribing rights. Prescribing of PBS medicines is also contingent on a prescriber being an *authorised nurse practitioner* and having collaborative arrangements in place, as required by amendments to the *National Health Act 1953*.

The Pharmaceutical Benefits Advisory Committee (PBAC) is responsible for making recommendations to the Minister for Health and Ageing regarding medicines for prescribing by authorised nurse practitioners.

Further to prescribing within collaborative arrangements, certain medicines also have additional conditions for prescribing by nurse practitioners, as recommended by the PBAC. These medicines are identified by the codes 'CTO' for continuation therapy only or 'SCM' for prescribing within a shared care model, as outlined below:

- *Continuing therapy only model*

Where the patient's treatment and prescribing of a medicine has been initiated by a medical practitioner, but prescribing is continued by a nurse practitioner. (This is similar to existing arrangements between specialists and medical practitioners for prescribing certain medicines.)

- *Shared care model*

Where care is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed plan to manage the patient, in a patientcentred model of care. The details surrounding shared care arrangements will depend on the practitioners involved, patient needs and the healthcare context.

Some medicines are included in more than one section of the Schedule, and for more than one prescriber type. For a prescription to be eligible for subsidy, prescribers must ensure that they prescribe under the PBS only those medicines, and in accordance with the restrictions, listed for their prescriber type. Listing details for the same product may differ between sections and different PBS item codes apply for each prescriber type.

Nurse practitioner PBS prescriptions are identifiable by colour, and include the indicator 'NP' on personalised forms and a tick box on non-personalised (blank) forms.

Prescriptions must include the nurse practitioner's PBS prescriber number. For unrestricted and restricted PBS medicines, midwives/nurse practitioners can use the personalised or non-personalised PBS prescriber forms. For authority required and authority required (streamlined) PBS medicines, midwives/nurse practitioners can use the authority personalised or non-personalised PBS prescriber forms. Nurse practitioner PBS prescriptions may include repeats.

Regulation 24 applies for nurse practitioner prescribing. A nurse practitioner can direct that original and repeat supplies of pharmaceutical benefits be supplied at the one time, if certain conditions are satisfied.

Authority prescriptions: Authority prescriptions for authority required items, or for increased quantities or repeats, require prior approval from Medicare Australia for each prescription. (Refer to Prescribing Medicines — Information for PBS prescribers and Supplying medicines — What Pharmacists Need to Know, for more information on authority prescriptions.)

State and territory requirements: Nurse practitioners may prescribe medicines as private prescriptions according to their state/territory prescribing accreditation. The medicines which can be prescribed differ between states and territories. It is the nurse practitioner's responsibility to ensure adherence to State/Territory law for all prescriptions (PBS and private) and additionally to all PBS requirements for PBS prescriptions.

Midwife PBS prescribing

MEDICINES WHICH MAY BE PRESCRIBED BY AUTHORISED MIDWIVES

From 1 September 2010, midwives endorsed to prescribe under state or territory legislation can apply for approval as PBS prescribers (*authorised midwives*). Information for midwives to become authorised PBS prescribers is available from Medicare Australia.

The medicines listed for prescribing by authorised midwives are identified by 'MW' in the PBS Schedule. Midwives must not write PBS prescriptions for other medicines.

PBS prescribing by midwives is limited by state and territory prescribing rights. It is also contingent on a prescriber being an *authorised midwife* and having collaborative arrangements in place, as required by amendments to the *National Health Act 1953*.

The Pharmaceutical Benefits Advisory Committee (PBAC) is responsible for making recommendations to the Minister for Health and Ageing regarding medicines for prescribing by authorised midwives.

Some medicines are included in more than one section of the Schedule, and for more than one prescriber type. For a prescription to be eligible for subsidy, prescribers must ensure that they prescribe under the PBS only those medicines, and in accordance with the restrictions, listed for their prescriber type. Listing details for the same product may differ between sections and different PBS item codes apply for each prescriber type.

Midwife PBS prescriptions are identifiable by colour, and include the indicator 'MW' on personalised forms and a tick box on non-personalised (blank) forms. Prescriptions must include the midwife's PBS prescriber number. For unrestricted and restricted PBS medicines, midwives/nurse practitioners can use the personalised or non-personalised PBS prescriber forms. For authority required and authority required (streamlined) PBS medicines, midwives/nurse practitioners can use the authority personalised or non-personalised PBS prescriber forms. Midwife PBS prescriptions may include repeats.

Regulation 24 applies for midwife prescribing. A midwife can direct that original and repeat supplies of pharmaceutical benefits be supplied at the one time, if certain conditions are satisfied.

Authority prescriptions: Authority prescriptions for authority required items, or for increased quantities or repeats, require prior approval from Medicare Australia for each prescription. (Refer to Prescribing Medicines—Information for PBS prescribers and Supplying Medicines — What Pharmacists Need to Know, for more information on authority prescriptions.)

State and Territory requirements: Midwives may prescribe medicines as private prescriptions according to their state/territory prescribing accreditation. The medicines which can be prescribed differ between states and territories. It is the midwife's responsibility to ensure adherence to state/territory law for all prescriptions (PBS and private) and additionally to all PBS requirements for PBS prescriptions.

THERAPEUTIC INDEX

Alimentary tract and metabolism 68*Stomatological preparations 68*

Stomatological preparations 68

Drugs for acid related disorders 68

Antacids 68

Drugs for peptic ulcer and gastro-oesophageal reflux disease (GORD) 68

Drugs for functional gastrointestinal disorders 76

Propulsives 76

Antiemetics and antinauseants 76

Antiemetics and antinauseants 76

Bile and liver Therapy 79

Bile therapy 79

Laxatives 80

Laxatives 80

Antidiarrheals, intestinal antiinflammatory/ antiinfective agents 82

Intestinal antiinfectives 82

Electrolytes with carbohydrates 82

Antipropulsives 82

Intestinal antiinflammatory agents 82

Digestives, incl. enzymes 85

Digestives, incl. enzymes 85

Drugs used in diabetes 87

Insulins and analogues 87

Blood glucose lowering drugs, excl. insulins 88

Vitamins 95

Vitamin A and D, incl. combinations of the two 95

Vitamin B 1 , plain and in combination with vitamin B 6 and vitamin B 12 95

Mineral supplements 95

Calcium 95

Potassium 95

Anabolic agents for systemic use 96

Anabolic steroids 96

Blood and blood forming organs 97*Antithrombotic agents 97*

Antithrombotic agents 97

Antihemorrhagics 104

Antifibrinolytics 104

Antianemic preparations 104

Iron preparations 104

Vitamin B 12 and folic acid 104

Blood substitutes and perfusion solutions 105

Blood and related products 105

I.V. solutions 105

Cardiovascular system 107*Cardiac therapy 107*

Cardiac glycosides 107

Antiarrhythmics, class I and III 107

Cardiac stimulants excl. cardiac glycosides 108

Vasodilators used in cardiac diseases 109

Antihypertensives 110

Antiadrenergic agents, centrally acting 110

Antiadrenergic agents, peripherally acting 111

Arteriolar smooth muscle, agents acting on 111

Diuretics 111

Low-ceiling diuretics, thiazides 111

Low-ceiling diuretics, excl. thiazides 112

High-ceiling diuretics 112

Potassium-sparing agents	113
Diuretics and potassium-sparing agents in combination	113
<i>Peripheral vasodilators</i>	114
Peripheral vasodilators	114
<i>Beta blocking agents</i>	114
Beta blocking agents	114
<i>Calcium channel blockers</i>	118
Selective calcium channel blockers with mainly vascular effects	118
Selective calcium channel blockers with direct cardiac effects	120
<i>Agents acting on the renin-angiotensin system</i>	121
ACE inhibitors, plain	121
ACE inhibitors, combinations	128
Angiotensin II antagonists, plain	130
Angiotensin II antagonists, combinations	131
<i>Lipid modifying agents</i>	137
Lipid modifying agents, plain	137
Lipid modifying agents, combinations	147
Dermatologicals	149
<i>Antifungals for dermatological use</i>	149
Antifungals for topical use	149
Antifungals for systemic use	150
<i>Antipsoriatics</i>	150
Antipsoriatics for topical use	150
Antipsoriatics for systemic use	151
<i>Antibiotics and chemotherapeutics for dermatological use</i>	151
Chemotherapeutics for topical use	151
<i>Corticosteroids, dermatological preparations</i>	152
Corticosteroids, plain	152
<i>Anti-acne preparations</i>	154
Anti-acne preparations for systemic use	154
<i>Other dermatological preparations</i>	154
Other dermatological preparations	154
Genito urinary system and sex hormones	156
<i>Other gynecologicals</i>	156
Contraceptives for topical use	156
Other gynecologicals	156
<i>Sex hormones and modulators of the genital system</i>	157
Hormonal contraceptives for systemic use	157
Androgens	158
Estrogens	159
Progestogens	161
Progestogens and estrogens in combination	161
Gonadotropins and other ovulation stimulants	162
Antiandrogens	163
Other sex hormones and modulators of the genital system	164
<i>Urologicals</i>	165
Other urologicals, incl. antispasmodics	165
Systemic hormonal preparations, excl. sex hormones and insulins	167
<i>Pituitary and hypothalamic hormones and analogues</i>	167
Anterior pituitary lobe hormones and analogues	167
Posterior pituitary lobe hormones	167
Hypothalamic hormones	168
<i>Corticosteroids for systemic use</i>	168
Corticosteroids for systemic use, plain	168
<i>Thyroid therapy</i>	171
Thyroid preparations	171
Antithyroid preparations	171

<i>Pancreatic hormones</i>	172
Glycogenolytic hormones	172
<i>Calcium homeostasis</i>	172
Parathyroid hormones and analogues	172
Anti-parathyroid agents	173
Antiinfectives for systemic use	175
<i>Antibacterials for systemic use</i>	175
Tetracyclines	175
Beta-lactam antibacterials, penicillins	177
Other beta-lactam antibacterials	181
Sulfonamides and trimethoprim	184
Macrolides, lincosamides and streptogramins	185
Aminoglycoside antibacterials	187
Quinolone antibacterials	187
Other antibacterials	188
<i>Antimycotics for systemic use</i>	190
Antimycotics for systemic use	190
<i>Antimycobacterials</i>	193
Drugs for treatment of tuberculosis	193
Drugs for treatment of lepra	193
<i>Antivirals for systemic use</i>	194
Direct acting antivirals	194
<i>Vaccines</i>	198
Bacterial vaccines	198
Antineoplastic and immunomodulating agents	199
<i>Antineoplastic agents</i>	199
Alkylating agents	199
Antimetabolites	200
Plant alkaloids and other natural products	203
Cytotoxic antibiotics and related substances	206
Other antineoplastic agents	207
<i>Endocrine therapy</i>	230
Hormones and related agents	230
Hormone antagonists and related agents	231
<i>Immunostimulants</i>	234
Immunostimulants	234
<i>Immunosuppressants</i>	236
Immunosuppressants	236
Musculo-skeletal system	358
<i>Antiinflammatory and antirheumatic products</i>	358
Antiinflammatory and antirheumatic products, non-steroids	358
Specific antirheumatic agents	362
<i>Muscle relaxants</i>	363
Muscle relaxants, centrally acting agents	363
Muscle relaxants, directly acting agents	364
<i>Antigout preparations</i>	364
Antigout preparations	364
<i>Drugs for treatment of bone diseases</i>	365
Drugs affecting bone structure and mineralization	365
Nervous system	375
<i>Analgesics</i>	375
Opioids	375
Other analgesics and antipyretics	384
Antimigraine preparations	385
<i>Antiepileptics</i>	386
Antiepileptics	386
<i>Anti-Parkinson drugs</i>	395

Anticholinergic agents	395
Dopaminergic agents	396
<i>Psycholeptics</i>	400
Antipsychotics	400
Anxiolytics	407
Hypnotics and sedatives	409
<i>Psychoanaleptics</i>	411
Antidepressants	411
Psychostimulants, agents used for ADHD and nootropics	419
Anti-dementia drugs	421
<i>Other nervous system drugs</i>	427
Parasympathomimetics	427
Drugs used in addictive disorders	427
Other nervous system drugs	428
Antiparasitic products, insecticides and repellents	430
<i>Antiprotozoals</i>	430
Agents against amoebiasis and other protozoal diseases	430
Antimalarials	430
<i>Anthelmintics</i>	431
Antitrepatodals	431
Antinematodal agents	431
<i>Ectoparasitocides, incl. scabicides, insecticides and repellents</i>	432
Ectoparasitocides, incl. scabicides	432
Respiratory system	433
<i>Nasal preparations</i>	433
Decongestants and other nasal preparations for topical use	433
<i>Drugs for obstructive airway diseases</i>	433
Adrenergics, inhalants	433
Other drugs for obstructive airway diseases, inhalants	435
Adrenergics for systemic use	437
Other systemic drugs for obstructive airway diseases	438
<i>Cough and cold preparations</i>	439
Cough suppressants, excl. combinations with expectorants	439
<i>Antihistamines for systemic use</i>	439
Antihistamines for systemic use	439
Sensory organs	440
<i>Ophthalmologicals</i>	440
Antiinfectives	440
Antiinflammatory agents	441
Antiglaucoma preparations and miotics	441
Mydriatics and cycloplegics	443
Decongestants and antiallergics	444
Ocular vascular disorder agents	444
Other ophthalmologicals	445
<i>Otologicals</i>	450
Antiinfectives	450
Corticosteroids and antiinfectives in combination	451
<i>Ophthalmological and otological preparations</i>	451
Antiinfectives	451
Various	452
<i>Allergens</i>	452
Allergens	452
<i>All other therapeutic products</i>	452
All other therapeutic products	452
<i>Diagnostic agents</i>	454
Urine tests	454
Other diagnostic agents	455

- General nutrients* 457
- Other nutrients 457
- All other non-therapeutic products* 467
- All other non-therapeutic products 467

PHARMACEUTICAL BENEFITS FOR PALLIATIVE CARE

Alimentary tract and metabolism 469

- Stomatological preparations* 469
- Stomatological preparations 469
- Drugs for functional gastrointestinal disorders* 470
- Belladonna and derivatives, plain 470
- Antiemetics and antinauseants* 470
- Antiemetics and antinauseants 470
- Laxatives* 471
- Laxatives 471

Musculo-skeletal system 475

- Antiinflammatory and antirheumatic products* 475
- Antiinflammatory and antirheumatic products, non-steroids 475

Nervous system 480

- Analgesics* 480
- Opioids 480
- Other analgesics and antipyretics 482
- Antiepileptics* 483
- Antiepileptics 483
- Psycholeptics* 483
- Anxiolytics 483
- Hypnotics and sedatives 484

PHARMACEUTICAL BENEFITS FOR DENTAL USE

Alimentary tract and metabolism 486

- Stomatological preparations* 486
- Stomatological preparations 486
- Drugs for functional gastrointestinal disorders* 486
- Propulsives 486
- Antiemetics and antinauseants* 486
- Antiemetics and antinauseants 486
- Antidiarrheals, intestinal antiinflammatory/ antiinfective agents* 487
- Intestinal antiinfectives 487

Blood and blood forming organs 488

- Blood substitutes and perfusion solutions* 488
- I.V. solutions 488

Cardiovascular system 489

- Cardiac therapy* 489
- Antiarrhythmics, class I and III 489
- Cardiac stimulants excl. cardiac glycosides 489
- Vasodilators used in cardiac diseases 489

Dermatologicals 490

- Corticosteroids, dermatological preparations* 490
- Corticosteroids, plain 490

Systemic hormonal preparations, excl. sex hormones and insulins 491

- Corticosteroids for systemic use* 491
- Corticosteroids for systemic use, plain 491
- Pancreatic hormones* 491

Glycogenolytic hormones 491

Antiinfectives for systemic use 492

Antibacterials for systemic use 492

Tetracyclines 492

Beta-lactam antibacterials, penicillins 492

Other beta-lactam antibacterials 495

Sulfonamides and trimethoprim 497

Macrolides, lincosamides and streptogramins 498

Other antibacterials 499

Musculo-skeletal system 501

Antiinflammatory and antirheumatic products 501

Antiinflammatory and antirheumatic products, non-steroids 501

Nervous system 504

Analgesics 504

Opioids 504

Other analgesics and antipyretics 508

Antiepileptics 509

Antiepileptics 509

Anti-Parkinson drugs 509

Anticholinergic agents 509

Psycholeptics 509

Anxiolytics 509

Hypnotics and sedatives 510

Respiratory system 511

Drugs for obstructive airway diseases 511

Adrenergics for systemic use 511

Sensory organs 512

Ophthalmologicals 512

Antiinfectives 512

Various 513

All other therapeutic products 513

All other therapeutic products 513

All other non-therapeutic products 513

All other non-therapeutic products 513

PHARMACEUTICAL BENEFITS FOR OPTOMETRICAL USE

Sensory organs 516

Ophthalmologicals 516

Antiinfectives 516

Antiinflammatory agents 516

Antiglaucoma preparations and miotics 516

Decongestants and antiallergics 518

Other ophthalmologicals 518

Ophthalmological and otological preparations 521

Antiinfectives 521

Section 2

Emergency Drug Supplies

Special Pharmaceutical Benefits

General Pharmaceutical Benefits

Pharmaceutical Benefits for Palliative Care

Pharmaceutical Benefits for Dental Use

Pharmaceutical Benefits for Optometrical Use

Items Available Under Special Arrangements (s.100)

SYMBOLS USED IN THE SCHEDULE

An asterisk (*) against the dispensed price of a benefit indicates that the manufacturer's pack does not coincide with the maximum quantity.

A double dagger (‡) in the maximum quantity column indicates an item for which the maximum quantity has been specially determined to correspond to the manufacturer's pack and the manufacturer's standard pack should be prescribed and supplied. For any item where a maximum quantity greater than 1 is marked with a double dagger (‡), that maximum quantity should be prescribed and supplied.

A gauge sign (#) against the dispensed price of a benefit indicates that the product is not preconstituted and that an extemporaneously-prepared dispensing fee is included in the dispensed price and, where appropriate, an amount for purified water.

Where a STATE is indicated after a manufacturer's code, that brand may be available only in the State indicated. NSW–(N); Vic–(V); Qld–(Q); SA–(S); WA–(W); Tas–(T).

RESTRICTED BENEFITS

All restricted items have separate headings for authority and non-authority items. In each case these items may be prescribed as pharmaceutical benefits only for use for one of the specified indications. Where more than one indication is specified for an Authority required or Restricted pharmaceutical benefit, each indication is separated from the preceding indication by a semi-colon and commences on the next line. In the case of Authority required (STREAMLINED) items, each indication will also include a four digit streamlined authority code. The drug may be prescribed as a pharmaceutical benefit for a patient who qualifies under any of the specified indications.

A straight line is drawn between entries for different forms and strengths of an item to indicate clearly the different restrictions which apply to these various forms and strengths.

The maximum quantity and/or number of repeats in respect of an item shown in the Schedule may be varied by the Chief Executive Officer of Medicare Australia when approving an Authority Prescription or an Authority to Prescribe. The quantity and number of repeats shown on the authority shall be supplied. (See Explanatory Notes). Payment will be made on the basis of the price shown for that item in the Schedule.

CODES FOR INJECTABLE ITEMS WITH ALLOWABLE SOLVENTS

The entry in this schedule of those pharmaceutical benefit injectable items which require a solvent includes the codes of the items with the relevant solvents. For each such item the code is for the injectable with 10mL sodium chloride injection 9 mg per mL (0.9%).

BRAND EQUIVALENCE

'a' located immediately before brand names of a particular strength of an item indicates that the sponsors of these brands have submitted evidence that they have been demonstrated to be bioequivalent or therapeutically equivalent, or that justification for not needing bioequivalence or therapeutic equivalence data has been provided to and accepted by the Therapeutic Goods Administration. It would thus be expected that these brands may be interchanged without differences in clinical effect.

For other brands of an item, i.e., those not indicated as above, it is unknown whether or not they are equivalent. There may be several reasons for this, such as bioequivalence data not being considered necessary when the products were approved for marketing, or that advice or data have not been forthcoming from sponsors. This does not necessarily suggest a lack of safety or efficacy, but in these circumstances caution should be taken if brands are interchanged.

'b' attached to brand names indicates that these brands are also equivalent, but that it is not known if there is equivalence between brands marked 'a' and brands marked 'b'.

BRAND PREMIUM POLICY

The Brand Premium Policy was introduced on 1 December 1990 to increase price competition by allowing pharmaceutical manufacturers to set their own price on multi-branded items listed on the Pharmaceutical Benefits Scheme and to encourage the development of the generic pharmaceutical industry in Australia. The policy does this by increasing prescribers' and patients' consciousness about the price of drugs. In effect, it makes both groups question whether it is necessary for the patient to pay more for the drugs when a cheaper brand is available. The policy also allows companies to establish prices taking into account competition and consumer acceptance.

The policy operates where there is more than one brand of a particular drug available through the Pharmaceutical Benefits Scheme and where the brands are therapeutically interchangeable. Due to this, the policy mainly applies to out of patent drugs.

Basically the policy operates by:

- the Australian Government subsidising a drug to the level of the lowest priced brand (except in those instances where the lowest priced brand has, as part of its price, a therapeutic group premium);
- suppliers of other brands of that drug being able to set a price above the price charged by the supplier(s) of the lowest priced brand(s); and
- the patient paying the brand premium which is the price difference between the lowest price brand and the brand prescribed.

If a prescription is written generically or for the lowest priced brand, and the lowest priced brand is supplied, there is no brand premium payable.

'B' located immediately before an amount in the premium column indicates a brand premium which applies to that particular brand of the item.

If a brand of a drug which is subject to a therapeutic group premium also has a brand premium, there will be two amounts shown on separate lines in the premium column, prefixed by 'T' and 'B' respectively.

If a brand of a drug which is subject to a special patient contribution also has a brand premium, there will be two amounts shown on separate lines in the premium column, prefixed by 'S' and 'B' respectively.

THERAPEUTIC GROUP PREMIUM POLICY

The Therapeutic Group Premium Policy was introduced on 1 February 1998 as an extension of the Brand Premium Policy to encourage greater competition between manufacturers of drugs and to make doctors and patients more aware of the costs of medicines.

The Therapeutic Group Premium policy applies within narrowly defined therapeutic sub-groups where the drugs concerned are of similar safety, efficacy and health outcomes.

Basically the policy operates by:

- the Australian Government subsidising drugs within a defined therapeutic sub-group to the level of the lowest priced drug in the sub-group;
- suppliers of other drugs within that sub-group being able to set prices above the price charged by the supplier(s) of the lowest priced drug; and
- the patient paying the therapeutic group premium which is the price difference between the lowest price drug and the drug prescribed.

'T' located immediately before an amount in the premium column indicates a therapeutic group premium which applies to that particular item.

If a brand of a drug which is subject to a therapeutic group premium also has a brand premium, there will be two amounts shown on separate lines in the premium column, prefixed by 'T' and 'B' respectively.

The success of the Government in controlling prices of products supplied through the Pharmaceutical Benefits Scheme has often been criticised by the pharmaceutical industry. Under both the Brand Premium Policy and the Therapeutic Group Premium Policy, suppliers of multi-branded items and therapeutically similar drugs are able to set their own prices at a level that they think the market will bear. At the same time, the prescriber and the patient can decide whether it is necessary to pay more for a particular brand or drug when a cheaper one is available and is therapeutically interchangeable.

The brand premium or therapeutic group premium does not count toward the patient's safety net.

It should be noted that the brand premium or therapeutic group premium is not a Government charge or revenue. The premium arises from the manufacturer's price and the majority goes to the manufacturer with wholesalers and pharmacists receiving a small percentage.

EMERGENCY DRUG SUPPLIES

Code	Name, Manner of Administration and Form	Max. Qty	Dispensed Price for Max. Qty \$	Proprietary Name and Manufacturer	
3451P NP	ADRENALINE Injection 1 mg in 1 mL (1 in 1,000)	5	20.34	AstraZeneca Pty Ltd	AP
3457Y NP	BENZTROPINE MESYLATE Injection 2 mg in 2 mL	5	103.59	Cogentin	FK
3486L NP or 3485K NP	BENZYLPENICILLIN Powder for injection 600 mg or PROCAINE PENICILLIN Injection 1.5 g	10 5	*42.92 92.22	BenPen Cilicaine	CS SI
3487M NP	BENZYLPENICILLIN Powder for injection 3 g	1	12.75	BenPen	CS
3455W NP or 3456X NP	CHLORPROMAZINE HYDROCHLORIDE Injection 50 mg in 2 mL or HALOPERIDOL Injection 5 mg in 1 mL	10 10	20.48 22.28	Largactil Serenace	SW SI
3478C NP	CLONAZEPAM Oral liquid 2.5 mg per mL, 10 mL	‡1	10.73	Rivotril	RO
3472R NP or 3470P NP or 3471Q NP	DEXAMETHASONE SODIUM PHOSPHATE Injection equivalent to 4 mg dexamethasone phosphate in 1 mL or HYDROCORTISONE SODIUM SUCCINATE Injection equivalent to 100 mg hydrocortisone with 2 mL solvent or HYDROCORTISONE SODIUM SUCCINATE Injection equivalent to 250 mg hydrocortisone with 2 mL solvent	5 2 1	18.08 *16.72 15.72	Hospira Pty Limited Solu-Cortef Solu-Cortef	HH PF PF
3458B NP	DIAZEPAM Injection 10 mg in 2 mL	5	12.29	Hospira Pty Limited	HH
3460D NP	DIHYDROERGOTAMINE MESYLATE Injection 1 mg in 1 mL	5	17.06	Dihydergot	NV
3463G NP	DIPHtheria and TETANUS VACCINE, ADSORBED, DILUTED FOR ADULT USE Injection 0.5 mL in pre-filled syringe	20	*275.02	ADT Booster	CS
3466K NP	FRUSEMIDE Injection 20 mg in 2 mL	5	10.48	a Frusemide Sandoz a Lasix a Frusemide-Clarix	SZ SW AE
3467L NP	GLUCAGON HYDROCHLORIDE Injection set containing 1 mg (1 i.u.) and 1 mL solvent in disposable syringe	1	45.63	GlucaGen Hypokit	NO
3475X NP	GLYCERYL TRINITRATE Sublingual spray (pump pack) 400 micrograms per dose (200 doses)	‡1	20.13	Nitrolingual Pumpspray	SW
3473T NP	HYOSCINE BUTYLBROMIDE Injection 20 mg in 1 mL	5	24.21	Buscopan	BY
3474W NP	LIGNOCAINE HYDROCHLORIDE Injection 100 mg in 5 mL	5	37.33	Pfizer Australia Pty Ltd	PF
3489P	METHOXYFLURANE Liquid for inhalation 999.9 mg per g, 3 mL (with inhaler)	1	44.78	Penthrox	NQ
3476Y NP or 3477B NP	METOCLOPRAMIDE HYDROCHLORIDE Injection 10 mg in 2 mL or PROCHLORPERAZINE Injection containing prochlorperazine mesylate 12.5 mg in 1 mL	10 10	12.99 16.82	Maxolon Stemetil	VT SW

EMERGENCY DRUG SUPPLIES

Code	Name, Manner of Administration and Form	Max. Qty	Dispensed Price for Max. Qty \$	Proprietary Name and Manufacturer	
3479D NP	MORPHINE SULFATE Injection 15 mg in 1 mL	5	14.45	Hospira Pty Limited	HH
or	or				
3480E NP	MORPHINE SULFATE Injection 30 mg in 1 mL	5	15.90	Hospira Pty Limited	HH
3482G NP	NALOXONE HYDROCHLORIDE Injection 2 mg in 5 mL	2	*78.08	Naloxone Min-I- Jet	CS
3488N NP	PROMETHAZINE HYDROCHLORIDE Injection 50 mg in 2 mL	10	*22.32	Hospira Pty Limited	HH
3495Y NP	SALBUTAMOL SULFATE Oral pressurised inhalation 100 micrograms (base) per dose (200 doses), CFC-free formulation	‡1	10.82	a Asmol CFC-free	AL
				a Airomir	IA
3495Y NP	SALBUTAMOL SULFATE Oral pressurised inhalation 100 micrograms (base) per dose (200 doses), CFC-free formulation	‡1	11.41	a Ventolin CFC-free	GK
or	or				
3496B NP	SALBUTAMOL SULFATE Nebuliser solution single dose units 2.5 mg (base) in 2.5 mL, 30	‡1	12.68	a Asmol 2.5 uni- dose	AF
				a GenRx Salbutamol	GX
				a Butamol 2.5	SI
				a Pfizer Australia Pty Ltd	PF
				a Pharmacor Salbutamol 2.5	CR
				a Salbutamol Sandoz	SZ
				a Salbutamol-GA	GM
3496B NP	SALBUTAMOL SULFATE Nebuliser solution single dose units 2.5 mg (base) in 2.5 mL, 30	‡1	13.41	a Ventolin Nebules	GK
3497C NP	SALBUTAMOL SULFATE Nebuliser solution single dose units 5 mg (base) in 2.5 mL, 30	‡1	13.03	a Asmol 5 uni-dose	AF
				a GenRx Salbutamol	GX
				a Butamol 5	SI
				a Pharmacor Salbutamol 5	CR
				a Salbutamol Sandoz	SZ
				a Salbutamol-GA	GM
3497C NP	SALBUTAMOL SULFATE Nebuliser solution single dose units 5 mg (base) in 2.5 mL, 30	‡1	13.76	a Ventolin Nebules	GK
3491R NP	TERBUTALINE SULFATE Injection 500 micrograms in 1 mL	5	30.59	Bricanyl	AP
3484J NP	TRAMADOL HYDROCHLORIDE Injection 100 mg in 2 mL	5	14.30	a Tramal 100	CS
				a Tramahexal	SZ
3494X NP	VERAPAMIL HYDROCHLORIDE Injection 5 mg in 2 mL	5	12.38	Isoptin	AB

SPECIAL PHARMACEUTICAL BENEFITS

The special patient contribution is payable by all patients in addition to the relevant patient contribution for concessional and general patients. Other than for bleomycin sulfate, exemptions on medical grounds are available. For eligible veterans under RPBS provisions, see RPBS EXPLANATORY NOTES, paragraph 32.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Reimbursement Price for Max. Qty \$	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	

GENERAL PHARMACEUTICAL BENEFITS

AMOXYCILLIN

1888J NP	Powder for paediatric oral drops 100 mg per mL, 20 mL	\$1	1	\$0.61	#13.18	#13.79	14.59	Amoxil	GK

AMOXYCILLIN

Authority required

Treatment of infections suspected or proven to be due to a susceptible organism in patients who require a liquid formulation and in whom the syrup formulations are unsuitable.

9714G NP	Powder for paediatric oral drops 100 mg per mL, 20 mL	\$1	1	..	#13.79	#13.79	15.20	Amoxil	GK

BLEOMYCIN SULFATE

Restricted benefit

Germ cell neoplasms;

Lymphoma.

2315W	Powder for injection 15,000 i.u. (solvent required) (code 6896Y applies to above item with approved solvent)	10	..	\$419.65	473.02	892.67	34.20	^a Blenamax	SI
				\$419.70	*473.02	*892.72	34.20	^a Hospira Pty Limited	HH
				^B 72.32	473.02	964.99	34.20	^a Blenoxane	BQ
				\$419.65					

NARATRIPTAN

Caution

Naratriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.

Authority required

Migraine attack in a patient where attacks in the past have usually failed to respond to analgesics.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8298R NP	Tablet 2.5 mg (as hydrochloride)	4	5	\$2.78	*25.90	*28.68	26.97	Naramig	GK

NARATRIPTAN

Caution

Naratriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.

Authority required

Migraine attack in a patient where attacks in the past have usually failed to respond to analgesics, and where:

- adverse events have occurred with other suitable PBS-listed drugs; or
- drug interactions have occurred with other suitable PBS-listed drugs; or
- drug interactions are expected to occur with other suitable PBS-listed drugs; or
- transfer to another suitable PBS-listed drug would cause patient confusion resulting in problems with compliance; or
- transfer to another suitable PBS-listed drug is likely to result in adverse clinical consequences.

SPECIAL PHARMACEUTICAL BENEFITS

The special patient contribution is payable by all patients in addition to the relevant patient contribution for concessional and general patients. Other than for bleomycin sulfate, exemptions on medical grounds are available. For eligible veterans under RPBS provisions, see RPBS EXPLANATORY NOTES, paragraph 32.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Reimbursement Price for Max. Qty \$	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9734H NP	Tablet 2.5 mg (as hydrochloride)	4	5	..	*28.68	*28.68	29.75	Naramig	GK

ZOLMITRIPTAN

Caution

Zolmitriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.

Authority required

Migraine attack in a patient where attacks in the past have usually failed to respond to analgesics.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8266C NP	Tablet 2.5 mg	4	5	§2.76	*25.84	*28.60	26.91	Zomig	AP

ZOLMITRIPTAN

Caution

Zolmitriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.

Authority required

Migraine attack in a patient where attacks in the past have usually failed to respond to analgesics, and where:

- (a) adverse events have occurred with other suitable PBS-listed drugs; or
- (b) drug interactions have occurred with other suitable PBS-listed drugs; or
- (c) drug interactions are expected to occur with other suitable PBS-listed drugs; or
- (d) transfer to another suitable PBS-listed drug would cause patient confusion resulting in problems with compliance; or
- (e) transfer to another suitable PBS-listed drug is likely to result in adverse clinical consequences.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9736K NP	Tablet 2.5 mg	4	5	..	*28.60	*28.60	29.67	Zomig	AP

PHARMACEUTICAL BENEFITS FOR DENTAL USE

AMOXYCILLIN

3310F	Powder for paediatric oral drops 100 mg per mL, 20 mL	±1	..	§0.61	#13.18	#13.79	14.59	Amoxil	GK

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	

Alimentary tract and metabolism

Stomatological preparations

Stomatological preparations

Antiinfectives and antiseptics for local oral treatment

AMPHOTERICIN								
2931G NP	Lozenge 10 mg	20	1	..	12.03	13.10	Fungilin	SI
NYSTATIN								
3033P NP	Oral suspension 100,000 units per mL, 24 mL	1	1	..	10.85	11.92	Mycostatin	FM
							Nilstat	SI

Other agents for local oral treatment

BENZYDAMINE HYDROCHLORIDE								
<u>Restricted benefit</u>								
Radiation induced mucositis.								
1121B NP	Mouth and throat rinse 22.5 mg per 15 mL, 500 mL	1	1	..	22.26	23.33	Diffiam	IA

Drugs for acid related disorders

Antacids

Combinations and complexes of aluminium, calcium and magnesium compounds

ALUMINIUM HYDROXIDE with MAGNESIUM HYDROXIDE								
2157M NP	Oral suspension 200 mg-200 mg per 5 mL, 500 mL	2	5	..	*15.72	16.79	Mylanta P	JT
ALUMINIUM HYDROXIDE with MAGNESIUM TRISILICATE and MAGNESIUM HYDROXIDE								
2159P NP	Oral suspension 250 mg-120 mg-120 mg per 5 mL, 500 mL	2	5	..	*17.70	18.77	Gastrogel	FM

Drugs for peptic ulcer and gastro-oesophageal reflux disease (GORD)

H₂-receptor antagonists

Note

The base-priced drugs in this therapeutic group are cimetidine, famotidine, nizatidine and ranitidine hydrochloride (except ranitidine hydrochloride effervescent tablet 150 mg (base) and syrup 150 mg (base) per 10 mL, 300 mL).

CIMETIDINE

Note

Helicobacter pylori eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.

1157X NP	Tablet 200 mg	120	5	..	19.12	20.19	Magicul 200	AF
1158Y NP	Tablet 400 mg	60	5	..	19.12	20.19 ^a	GenRx Cimetidine	GX
							^a Magicul 400	AF
1159B NP	Tablet 800 mg	30	5	..	19.12	20.19 ^a	GenRx Cimetidine	GX
							^a Magicul 800	AF

FAMOTIDINE

Note

Helicobacter pylori eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.

2487X NP	Tablet 20 mg	60	5	..	17.41	18.48 ^a	Ausfam 20	SI

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer			
2488Y NP	Tablet 40 mg	30	5	B4.95	22.36	18.48	a Chem mart Famotidine CH			
							a Famohexal HX			
							a Famotidine Sandoz SZ			
							a GenRx Famotidine GX			
							a Pamacid 20 AF			
							a Pepzan GM			
							a Terry White Chemists Famotidine TW			
				a Pepcidine M MK						
				a Ausfam 40 SI						
				a Chem mart Famotidine CH						
				a Famotidine Sandoz SZ						
				a GenRx Famotidine GX						
				a Pamacid 40 AF						
				a Pepzan GM						
				a Terry White Chemists Famotidine TW						
				B5.41	22.82	18.48	a Pepcidine MK			
				NIZATIDINE						
Note										
Helicobacter pylori eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.										
1504E NP	Capsule 300 mg	30	5	..	19.06	20.13	a Nizac LN			
1505F NP	Capsule 150 mg	60	5	..	19.06	20.13	a Tacidine AF			
							B5.60	24.66	20.13	a Tazac AS
							a Nizac LN			
							a Tacidine AF			
							B5.60	24.66	20.13	a Tazac AS
RANITIDINE HYDROCHLORIDE										
Note										
Helicobacter pylori eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.										
1937Y NP	Effervescent tablet 150 mg (base)	60	5	T3.26	*21.16	18.97	Zantac GK			
1977C NP	Tablet 300 mg (base)	30	5	..	17.87	18.94	a Ausran SI			
1978D NP,MW	Tablet 150 mg (base)	60	5	B1.71	19.58	18.94	a Chem mart Ranitidine CH			
							a GenRx Ranitidine GX			
							a Rani 2 AF			
							a Ranitidine Sandoz SZ			
				a Terry White Chemists Ranitidine TW						
				a Ulcaid RA						
				a Zantac GK						
				a Ausran SI						
a Chem mart Ranitidine CH										
a GenRx Ranitidine GX										
a Rani 2 AF										

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
							^a Ranihexal	HX
							^a Ranitidine Sandoz	SZ
							^a Ranoxyl	GM
							^a Terry White Chemists Ranitidine	TW
							^a Ulcaid	RA
				^B 1.71	19.58	18.94	^a Zantac	GK
8162N NP	Syrup 150 mg (base) per 10 mL, 300 mL	2	5	^T 2.20	*23.92	22.79	Zantac Syrup	GK

RANITIDINE HYDROCHLORIDE

Note

Helicobacter pylori eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.

Authority required

Adverse effects occurring with all of the base-priced drugs;

Drug interactions occurring with all of the base-priced drugs;

Drug interactions expected to occur with all of the base-priced drugs;

Transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.

8903N NP	Effervescent tablet 150 mg (base)	60	5	..	*21.16	22.23	Zantac	GK
8905Q NP	Syrup 150 mg (base) per 10 mL, 300 mL	2	5	..	*23.92	24.99	Zantac Syrup	GK

Prostaglandins

MISOPROSTOL

Caution

Misoprostol is a prostaglandin analogue. It should not be used in pregnant women.

Authority required (STREAMLINED)

2630

Reduction in the incidence of gastrointestinal complications in patients who have a history of peptic ulcer disease and where NSAID therapy is essential;

2631

Duodenal ulcer (including pyloric and stomal ulcers), proven by current or prior x-ray, endoscopy or surgery. The date and the method by which the ulcer was proven must be documented in the patient's medical records when treatment is initiated;

2632

Gastric ulcer, proven by x-ray, endoscopy or surgery within the previous 2 years. The date and the method by which the ulcer was proven must be documented in the patient's medical records when treatment is initiated.

1648R	Tablet 200 micrograms	120	2	..	52.12	34.20	Cytotec	PF

Proton pump inhibitors

ESOMEPRAZOLE MAGNESIUM TRIHYDRATE

Restricted benefit

Initial treatment of gastric ulcer.

Note

Helicobacter pylori eradication therapy should be considered.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8886Q NP	Tablet (enteric coated), equivalent to 20 mg esomeprazole	30	1	..	33.36	34.20	Nexium	AP

ESOMEPRAZOLE MAGNESIUM TRIHYDRATE

Restricted benefit

Healing of gastro-oesophageal reflux disease.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
	Note No applications for increased maximum quantities and/or repeats will be authorised.							
8601Q NP	Tablet (enteric coated), equivalent to 40 mg esomeprazole	30	1	..	50.81	34.20	Nexium	AP
	ESOMEPRAZOLE MAGNESIUM TRIHYDRATE Restricted benefit Maintenance of healed gastro-oesophageal reflux disease; Scleroderma oesophagus; Pathological hypersecretory conditions including Zollinger-Ellison syndrome and idiopathic hypersecretion.							
	Note No applications for increased maximum quantities will be authorised.							
8600P NP	Tablet (enteric coated), equivalent to 20 mg esomeprazole	30	5	..	33.36	34.20	Nexium	AP
	ESOMEPRAZOLE MAGNESIUM TRIHYDRATE Authority required Pathological hypersecretory conditions including Zollinger-Ellison syndrome and idiopathic hypersecretion.							
	Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
3401B NP	Tablet (enteric coated), equivalent to 40 mg esomeprazole	30	5	..	50.81	34.20	Nexium	AP
	LANSOPRAZOLE Restricted benefit Initial treatment of peptic ulcer.							
	Note Helicobacter pylori eradication therapy should be considered.							
	No applications for increased repeats will be authorised.							
	Note Bioequivalence has been demonstrated between lansoprazole capsule 30 mg and lansoprazole tablet 30 mg (orally disintegrating).							
2240X NP	Capsule 30 mg	28	1	..	29.54	30.61 ^a	AP0-Lansoprazole	TX
						^a	Lanzopran	RA
						^a	Zopral	AF
9477T NP	Tablet 30 mg (orally disintegrating)	28	1	..	29.54	30.61 ^a	Zoton FasTabs	WX
	LANSOPRAZOLE Restricted benefit Gastro-oesophageal reflux disease; Scleroderma oesophagus.							
8198L NP	Capsule 15 mg	30	5	..	20.27	21.34	Zopral	AF
9331D NP	Tablet 15 mg (orally disintegrating)	28	5	..	19.33	20.40	Zoton FasTabs	WX
	LANSOPRAZOLE Restricted benefit Gastro-oesophageal reflux disease;							

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
	Scleroderma oesophagus.							
	Note							
	Bioequivalence has been demonstrated between lansoprazole capsule 30 mg and lansoprazole tablet 30 mg (orally disintegrating).							
2241Y NP	Capsule 30 mg	28	5	..	29.54	30.61	^a	APO-Lansoprazole TX
							^a	Lanzopran RA
							^a	Zopral AF
9478W NP	Tablet 30 mg (orally disintegrating)	28	5	..	29.54	30.61	^a	Zoton FasTabs WX
	OMEPRAZOLE							
	Restricted benefit							
	Initial treatment of peptic ulcer.							
	Note							
	Helicobacter pylori eradication therapy should be considered.							
	No applications for increased repeats will be authorised.							
	Note							
	Bioequivalence has been demonstrated between omeprazole tablet 20 mg and omeprazole tablet 20 mg (as magnesium).							
8331L NP	Tablet 20 mg	30	1	..	29.34	30.41	^a	APO-Omeprazole TX
							^a	Chem mart CH
							^a	Omeprazole GX
							^a	GenRx Omeprazole
							^a	Meprazol SZ
							^a	Omeprazole-GA GM
							^a	Omeprazole GQ
							^a	generichealth
							^a	Omeprazole RA
							^a	Ranbaxy
							^a	Omeprazole WA
							^a	Winthrop
							^a	Ozmep ZP
							^a	Terry White TW
							^a	Chemists
9109K NP	Tablet 20 mg (as magnesium)	30	1	..	29.34	30.41	^a	Omeprazole AL
							^a	Acimax Tablets
							^a	Omepral PM
				^B 3.75	33.09	30.41	^a	Losec Tablets AP
	OMEPRAZOLE							
	Restricted benefit							
	Initial treatment of peptic ulcer.							
	Note							
	Helicobacter pylori eradication therapy should be considered.							
	No applications for increased repeats will be authorised.							
1326T NP	Capsule 20 mg	30	1	..	29.34	30.41	^a	Omepro-GA GM
							^a	Pemzo SI
							^a	Pharmacor CR
							^a	Omeprazole 20
							^a	Probitor SZ

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
OMEPRAZOLE								
<u>Restricted benefit</u>								
Gastro-oesophageal reflux disease;								
Scleroderma oesophagus;								
Zollinger-Ellison syndrome.								
<u>Note</u>								
Bioequivalence has been demonstrated between omeprazole tablet 20 mg and omeprazole tablet 20 mg (as magnesium).								
8333N NP	Tablet 20 mg	30	5	..	29.34	30.41	^a	APO-Omeprazole TX
							^a	Chem mart CH
								Omeprazole
							^a	GenRx Omeprazole GX
							^a	Meprazol SZ
							^a	Omeprazole-GA GM
							^a	Omeprazole GQ
								generichealth
							^a	Omeprazole RA
							^a	Ranbaxy
9110L NP	Tablet 20 mg (as magnesium)	30	5	..	29.34	30.41	^a	Omeprazole Winthrop WA
							^a	Ozmep ZP
							^a	Terry White Chemists TW
							^a	Omeprazole
							^a	Acimax Tablets AL
							^a	Omepral PM
							^a	Losec Tablets AP
					^B 3.75	33.09		
					OMEPRAZOLE			
<u>Restricted benefit</u>								
Gastro-oesophageal reflux disease;								
Scleroderma oesophagus;								
Zollinger-Ellison syndrome.								
1327W NP	Capsule 20 mg	30	5	..	29.34	30.41	^a	Omepro-GA GM
							^a	Pemzo SI
							^a	Pharmacor CR
							^a	Omeprazole 20 SZ
8332M NP	Tablet 10 mg (as magnesium)	30	5	..	22.58	23.65		Probitor
								Losec Tablets AP
PANTOPRAZOLE SODIUM SESQUIHYDRATE								
<u>Restricted benefit</u>								
Initial treatment of peptic ulcer.								
<u>Note</u>								
Helicobacter pylori eradication therapy should be considered.								
<u>Note</u>								
No applications for increased repeats will be authorised.								
8007K NP	Tablet (enteric coated), equivalent to 40 mg pantoprazole	30	2	..	31.98	33.05	^a	APO-Pantoprazole TX
							^a	Chem mart CH
								Pantoprazole
							^a	Ozpan RA
							^a	Panto NZ
							^a	Pantofast 40 RZ
						^a	Pantoloc NH	

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer	
							a	Pantoprazole-GA	GM
							a	Pantoprazole generichealth	GQ
							a	Pantoprazole Sandoz	SZ
							a	Salpraz	AF
							a	Somac	NQ
							a	Sozol	SI
							a	Terry White Chemists Pantoprazole	TW
9423Y NP	Sachet containing granules 40 mg	30	2	..	31.98	33.05		Somac	NQ
PANTOPRAZOLE SODIUM SESQUIHYDRATE									
<u>Restricted benefit</u>									
Gastro-oesophageal reflux disease.									
8399C NP	Tablet (enteric coated), equivalent to 20 mg pantoprazole	30	5	..	18.89	19.96	a	APO-Pantoprazole	TX
							a	Chem mart Pantoprazole	CH
							a	Ozpan	RA
							a	Panto	NZ
							a	Pantofast 20	RZ
							a	Pantoloc	NH
							a	Pantoprazole-GA	GM
							a	Pantoprazole generichealth	GQ
							a	Pantoprazole Sandoz	SZ
							a	Salpraz	AF
							a	Somac	NQ
							a	Terry White Chemists Pantoprazole	TW
PANTOPRAZOLE SODIUM SESQUIHYDRATE									
<u>Restricted benefit</u>									
Gastro-oesophageal reflux disease.									
<u>Restricted benefit</u>									
Scleroderma oesophagus;									
Zollinger-Ellison syndrome.									
8008L NP	Tablet (enteric coated), equivalent to 40 mg pantoprazole	30	5	..	31.98	33.05	a	APO-Pantoprazole	TX
							a	Chem mart Pantoprazole	CH
							a	Ozpan	RA
							a	Panto	NZ
							a	Pantofast 40	RZ
							a	Pantoloc	NH
							a	Pantoprazole-GA	GM
							a	Pantoprazole generichealth	GQ
							a	Pantoprazole Sandoz	SZ
							a	Salpraz	AF
							a	Somac	NQ

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
							^a Sozol	SI
							^a Terry White Chemists Pantoprazole Somac	TW NQ
9424B NP	Sachet containing granules 40 mg	30	5	..	31.98	33.05		

RABEPRAZOLE SODIUM

Restricted benefit

Initial treatment of peptic ulcer.

Note

Helicobacter pylori eradication therapy should be considered.

Note

No applications for increased repeats will be authorised.

8509W NP	Tablet 20 mg (enteric coated)	30	2	..	37.78	34.20	Pariet	JC

RABEPRAZOLE SODIUM

Restricted benefit

Gastro-oesophageal reflux disease;

Scleroderma oesophagus.

8507R NP	Tablet 10 mg (enteric coated)	28	5	..	37.78	34.20	Pariet	JC
8508T NP	Tablet 20 mg (enteric coated)	30	5	..	37.78	34.20	Pariet	JC

Combinations for eradication of Helicobacter pylori

ESOMEPRAZOLE MAGNESIUM TRIHYDRATE and CLARITHROMYCIN and AMOXYCILLIN

Restricted benefit

Eradication of Helicobacter pylori associated with peptic ulcer disease.

8738X NP	Pack containing 14 tablets (enteric coated) equivalent to 20 mg esomeprazole, 14 tablets clarithromycin 500 mg and 28 capsules amoxicillin 500 mg	£1	..	..	81.36	34.20	Nexium Hp7	AP

OMEPRAZOLE and CLARITHROMYCIN and AMOXYCILLIN

Restricted benefit

Eradication of Helicobacter pylori associated with peptic ulcer disease.

8272J NP	Pack containing 14 capsules omeprazole 20 mg, 14 tablets clarithromycin 500 mg and 28 capsules amoxicillin 500 mg	£1	..	..	65.71	34.20	Probitor Hp7	SZ

Other drugs for peptic ulcer and gastro-oesophageal reflux disease (GORD)

SODIUM ALGINATE with CALCIUM CARBONATE and SODIUM BICARBONATE

2014B NP	Oral liquid 1 g-320 mg-534 mg in 20 mL, 500 mL	2	5	..	*14.68	15.75	Gaviscon P	RC

SUCRALFATE

2055E NP	Tablet equivalent to 1 g anhydrous sucralfate	120	2	..	23.69	24.76	^a Ulcyte	AF
				^B 2.11	25.80	24.76	^a Carafate	AS

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Drugs for functional gastrointestinal disorders							

Propulsives

Propulsives

DOMPERIDONE								
1347X NP	Tablet 10 mg	25	..	..	8.89	9.96	Motilium	JC
METOCLOPRAMIDE HYDROCHLORIDE								
1206L NP,MW	Injection 10 mg in 2 mL	10	..	..	12.99	14.06	Maxolon	VT
1207M NP,MW	Tablet 10 mg	25	..	..	8.20	9.27	Pramin	AF
				^B 3.02	11.22	9.27	Maxolon	VT

Antiemetics and antinauseants

Antiemetics and antinauseants

Serotonin (5HT₃) antagonists

DOLASETRON MESYLATE

Restricted benefit

Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

8191D NP	Tablet 200 mg	2	..	..	50.72	34.20	Anzemet	SW
8192E NP	I.V. injection 100 mg in 5 mL	1	..	..	29.29	30.36	Anzemet	SW

GRANISETRON HYDROCHLORIDE

Restricted benefit

Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

8728J NP	Tablet 2 mg (base)	2	..	..	*58.98	34.20	Kytril	HH
8729K NP	Concentrated injection 3 mg (base) in 3 mL	1	..	..	*37.85	34.20	^a Granisetron Kabi	PK
							^a Kytril	HH

GRANISETRON HYDROCHLORIDE

Authority required (STREAMLINED)

3611

Management of nausea and vomiting associated with radiotherapy being used to treat malignancy.

8730L NP	Concentrated injection 3 mg (base) in 3 mL	1	..	..	*37.85	34.20	^a Granisetron Kabi	PK
							^a Kytril	HH
8873B NP	Tablet 2 mg (base)	5	1	..	137.84	34.20	Kytril	HH

ONDANSETRON

Restricted benefit

Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
8224W NP	Tablet 4 mg	4	..	..	39.44	34.20	^a	APO-Ondansetron TX
							^a	Ondaz SZ
							^a	Onsetron 4 ZP
							^a	Zofran GK
8225X NP	Tablet 8 mg	4	..	..	55.89	34.20	^a	APO-Ondansetron TX
							^a	Ondaz SZ
							^a	Onsetron 8 ZP
							^a	Zofran GK
8226Y NP	I.V. injection 4 mg in 2 mL	1	..	..	19.93	21.00	^a	Ondansetron-Claris AE
							^a	Ondaz SZ
							^a	Onsetron ZP
							^a	Pfizer Australia Pty Ltd PF
							^a	Zofran GK
8227B NP	I.V. injection 8 mg in 4 mL	1	..	..	27.90	28.97	^a	Ondansetron-Claris AE
							^a	Ondaz SZ
							^a	Onsetron ZP
							^a	Pfizer Australia Pty Ltd PF
							^a	Zofran GK
8410P NP	Wafer 4 mg	4	..	..	39.44	34.20	^a	Ondaz Zydis SZ
							^a	Zofran Zydis GK
8411Q NP	Wafer 8 mg	4	..	..	55.89	34.20	^a	Ondaz Zydis SZ
							^a	Zofran Zydis GK
9441X NP	Syrup 4 mg per 5 mL, 50 mL	±1	..	..	85.38	34.20		Zofran syrup 50 mL GK

ONDANSETRON

Authority required (STREAMLINED)

3611

Management of nausea and vomiting associated with radiotherapy being used to treat malignancy.

1594X NP	Tablet 4 mg	10	1	..	85.38	34.20	^a	APO-Ondansetron TX
							^a	Ondaz SZ
							^a	Onsetron 4 ZP
							^a	Zofran GK
1595Y NP	Tablet 8 mg	10	1	..	130.12	34.20	^a	APO-Ondansetron TX
							^a	Ondaz SZ
							^a	Onsetron 8 ZP
							^a	Zofran GK
1596B NP	I.V. injection 4 mg in 2 mL	1	..	..	19.93	21.00	^a	Ondansetron-Claris AE
							^a	Ondaz SZ
							^a	Onsetron ZP
							^a	Pfizer Australia Pty Ltd PF
							^a	Zofran GK
1597C NP	I.V. injection 8 mg in 4 mL	1	..	..	27.90	28.97	^a	Ondansetron-Claris AE
							^a	Ondaz SZ
							^a	Onsetron ZP

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
							^a Pfizer Australia Pty Ltd	PF
							^a Zofran	GK
8233H NP	Syrup 4 mg per 5 mL, 50 mL	1	1	..	85.38	34.20	Zofran syrup 50 mL	GK
8412R NP	Wafer 4 mg	10	1	..	85.38	34.20	^a Ondaz Zydis	SZ
							^a Zofran Zydis	GK
8413T NP	Wafer 8 mg	10	1	..	130.12	34.20	^a Ondaz Zydis	SZ
							^a Zofran Zydis	GK

PALONOSETRON

Restricted benefit

Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Note

No applications for increased maximum quantities will be authorised. Palonosetron is not PBS-subsidised for administration with oral 5-HT3 antagonists.

5295Q	Injection 250 micrograms (as hydrochloride) in 5 mL	1	..	..	47.86	34.20	Aloxi	TS

TROPISETRON HYDROCHLORIDE

Restricted benefit

Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

2745L NP	Capsule 5 mg (base)	2	..	..	50.72	34.20	Navoban	NV
2746M NP	I.V. injection 5 mg (base) in 5 mL	1	..	..	29.29	30.36	Navoban	NV

Other antiemetics

APREPITANT

Note

Aprepitant is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.

Authority required (STREAMLINED)

3619

Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat malignancy, in combination with a 5HT3 antagonist and dexamethasone, where any 1 of the following chemotherapy agents are to be administered:

- (a) altretamine;
- (b) carmustine;
- (c) cisplatin when a single dose constitutes a cycle of chemotherapy;
- (d) cyclophosphamide at a dose of 1500 mg per square metre per day or greater;
- (e) dacarbazine;
- (f) procarbazine when a single dose constitutes a cycle of chemotherapy;
- (g) streptozocin.

No more than 1 pack containing 1 x 125 mg capsule and 2 x 80 mg capsules will be authorised per cycle of cytotoxic chemotherapy;

3620

Management of nausea and vomiting associated with cytotoxic chemotherapy being used to treat breast cancer, in combination with a 5HT3 antagonist and dexamethasone, where cyclophosphamide and an anthracycline are to be co-administered.

No more than 1 pack containing 1 x 125 mg capsule and 2 x 80 mg capsules will be authorised per cycle of cytotoxic chemotherapy;

3621

Management of nausea and vomiting associated with moderately emetogenic cytotoxic chemotherapy being used to treat malignancy, in combination with a 5HT3 antagonist and dexamethasone on day 1, where the patient has had a prior episode of chemotherapy induced nausea or vomiting where any 1 of the following intravenous chemotherapy agents is to be administered:

- (a) arsenic trioxide;
- (b) azacitidine;
- (c) carboplatin;
- (d) cyclophosphamide at a dose of less than 1500 mg per square metre per day;
- (e) cytarabine at a dose of greater than 1 g per square metre per day;
- (f) dactinomycin;

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	(g) daunorubicin; (h) doxorubicin; (i) epirubicin; (j) fotemustine; (k) idarubicin; (l) ifosfamide; (m) irinotecan; (n) melphalan; (o) methotrexate at a dose of 250 mg to 1 g per square metre; (p) oxaliplatin; (q) raltitrexed. No more than one pack containing 1 x 125 mg capsule and 2 x 80 mg capsules will be authorised per cycle of cytotoxic chemotherapy. Concomitant use of a 5HT3 antagonist should not occur with aprepitant on days 2 and 3 of any chemotherapy cycle.						
	Note No applications for increased maximum quantities will be authorised. Prescribers should advise Medicare Australia of the number of cycles planned when requesting approval for repeats.						
8808N NP	Pack containing 1 capsule 125 mg and 2 capsules 80 mg	£1	..	..	138.89	34.20	Emend MK
	PROCHLORPERAZINE						
	Caution Prochlorperazine may be associated with parkinsonism and tardive dyskinesia and should be used for short-term treatment only.						
	Note As prochlorperazine may be associated with parkinsonism and tardive dyskinesia it should be used for short-term treatment only. However, authorities for increased maximum quantities and/or repeats of prochlorperazine tablets will be granted for the treatment of emesis associated with malignant disease.						
2369Q NP	Injection containing prochlorperazine mesylate 12.5 mg in 1 mL	10	..	..	16.82	17.89	Stemetil SW
2893G NP	Tablet containing prochlorperazine maleate 5 mg	25	..	..	9.51	10.58	^a APO-Prochlorperazine TX ^a ProCalm SI ^a Prochlorperazine-GA GM ^a Stemetil AV
				^B 3.45	12.96	10.58	^a Stemetil SW
2895J NP	Suppositories containing prochlorperazine equivalent to 25 mg prochlorperazine maleate, 5	£1	2	..	19.93	21.00	Stemetil SW

Bile and liver Therapy

Bile therapy

Bile acid preparations

URSODEOXYCHOLIC ACID

Authority required (STREAMLINED)

1700

Primary biliary cirrhosis.

Note

Not for use in the treatment of sclerosing cholangitis or cholelithiasis.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8448P NP	Capsule 250 mg	200	2	..	*372.60	34.20	Ursosalk OA

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	
Laxatives							

Laxatives

Contact laxatives

BISACODYL

Restricted benefit

Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;

Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;

Patients receiving palliative care;

Terminal malignant neoplasia;

Anorectal congenital abnormalities;

Megacolon.

1258F NP	Suppositories 10 mg, 12	3	4	..	*18.57	19.64	Petrus Bisacodyl Suppositories	PP
1259G NP	Tablet 5 mg	200	2	..	14.27	15.34	Bisalax	AS
							Lax-Tab	AE
1260H NP	Suppositories 10 mg, 10	3	5	..	*21.27	22.34	^a Petrus Bisacodyl Suppositories	PP
				^B 1.08	*22.35	22.34	^a Dulcolax	BY

Bulk producers

STERCULIA with FRANGULA BARK

Restricted benefit

Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;

Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;

Patients receiving palliative care;

Terminal malignant neoplasia;

Anorectal congenital abnormalities;

Megacolon.

1104D NP	Granules 620 mg-80 mg per g (62%-8%), 500 g	£1	1	..	24.95	26.02	Normacol Plus	NE

Osmotically acting laxatives

LACTULOSE

Restricted benefit

Hepatic coma or precoma (chronic porto-systemic encephalopathy);

Constipation in patients with malignant neoplasia.

3064G NP	Mixture 3.34 g per 5 mL, 500 mL	£1	5	..	14.23	15.30	^a Actilax	AF
							^a Genlac	SI
							^a GenRx Lactulose	GX
							^a Lac-Dol	GM
							^a Lactocur	SZ
				^B 1.67	15.90	15.30	^a Duphalac	SM

MACROGOL 3350

Restricted benefit

Constipation in patients with malignant neoplasia;

Chronic constipation or faecal impaction not adequately controlled with first line interventions such as bulk-forming agents;

Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function not responding to other oral therapies;

Patients receiving palliative care.

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
3416T NP	Powder for oral solution 510 g	£1	5	..	20.55	21.62 ^a	MediHealth ClearLax	ON
						^a	OsmoLax	KY
8612G NP	Sachets containing powder for solution 13.125 g with electrolytes, 30	£1	5	..	20.55	21.62	Movicol	NE

Enemas

BISACODYL

Restricted benefit

Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;

Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;

Patients receiving palliative care;

Terminal malignant neoplasia;

Anorectal congenital abnormalities;

Megacolon.

1263L NP	Enemas 10 mg in 5 mL, 25	£1	2	..	38.59	34.20	Bisalax	AS

SORBITOL with SODIUM CITRATE and SODIUM LAURYL SULFOACETATE

Restricted benefit

Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;

Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;

Patients receiving palliative care;

Terminal malignant neoplasia;

Anorectal congenital abnormalities;

Megacolon.

2091C NP	Enemas 3.125 g-450 mg-45 mg in 5 mL, 12	2	2	..	*32.80	33.87 ^a	Micolette	AE
						^a	Microlax	JT

Other laxatives

GLYCEROL

Restricted benefit

Paraplegic and quadriplegic patients and others with severe neurogenic impairment of bowel function;

Patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult;

Patients receiving palliative care;

Terminal malignant neoplasia;

Anorectal congenital abnormalities;

Megacolon.

2555L NP	Suppositories 700 mg (for infants), 12	3	5	..	*18.84	19.91	Petrus Pharmaceuticals Pty Ltd	PP
2556M NP	Suppositories 1.4 g (for children), 12	3	5	..	*19.26	20.33	Petrus Pharmaceuticals Pty Ltd	PP
2557N NP	Suppositories 2.8 g (for adults), 12	3	5	..	*19.74	20.81	Petrus Pharmaceuticals Pty Ltd	PP

Alimentary tract and metabolism

					Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium			Brand Name and Manufacturer	
Antidiarrheals, intestinal antiinflammatory/ antiinfective agents								

Intestinal antiinfectives

Antibiotics

NEOMYCIN SULFATE								
2325J NP	Tablet 500 mg	25	1	..	15.05	16.12	Neosulf	AF

NYSTATIN								
1696G NP	Tablet 500,000 units	50	..	..	17.98	19.05	Nilstat	SI
1699K NP	Capsule 500,000 units	50	..	..	17.98	19.05	Nilstat	SI

VANCOMYCIN

Authority required

Antibiotic associated pseudomembranous colitis due to Clostridium difficile which is unresponsive to metronidazole;

Antibiotic associated pseudomembranous colitis due to Clostridium difficile where there is intolerance to metronidazole.

Note

Metronidazole has similar efficacy to vancomycin but may have less selective pressure to vancomycin resistant enterococci and is therefore the preferred treatment.

3113W	Capsule 125 mg (125,000 i.u.) vancomycin activity	40	..	..	*236.50	34.20	Vancocin	AS
3114X	Capsule 250 mg (250,000 i.u.) vancomycin activity	40	..	..	*448.54	34.20	Vancocin	AS

Electrolytes with carbohydrates

Oral rehydration salt formulations

ELECTROLYTE REPLACEMENT (ORAL)

Note

Each sachet contains sodium chloride 470 mg, potassium chloride 300 mg, sodium acid citrate 530 mg and glucose 3.56 g.

3196F NP	Sachets containing powder for oral solution 4.9 g, 10	±1	..	..	13.06	14.13	^a O.R.S.	AS
							^a Repalyte New Formulation	SW
							^a restore O.R.S.	GM

Antipropulsives

Antipropulsives

DIPHENOXYLATE HYDROCHLORIDE with ATROPINE SULFATE								
2501P NP	Tablet 2.5 mg-25 micrograms	20	..	..	8.52	9.59	^a Lofenoxal	HC
				^B 1.76	10.28	9.59	^a Lomotil	BI
LOPERAMIDE HYDROCHLORIDE								
1571Q NP	Capsule 2 mg	12	..	..	8.50	9.57	^a Gastro-Stop Loperamide	AS
				^B 0.91	9.41	9.57	^a Imodium	JT

Intestinal antiinflammatory agents

Corticosteroids acting locally

HYDROCORTISONE ACETATE

Restricted benefit

Proctitis;

Ulcerative colitis.

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
1502C NP	Rectal foam 90 mg per applicatorful, 14 applications, aerosol 21.1 g	2	3	..	*37.70	34.20	Colifoam AS
PREDNISOLONE SODIUM PHOSPHATE Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
1920C NP	Retention enema equivalent to 20 mg prednisolone in 100 mL	28	3	..	*221.18	34.20	Predsol SI
PREDNISOLONE SODIUM PHOSPHATE Restricted benefit Proctitis; Ulcerative colitis. Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
2554K NP	Suppositories equivalent to 5 mg prednisolone, 10	3	3	..	*43.32	34.20	Predsol SI
Aminosalicylic acid and similar agents BALSALAZIDE SODIUM Authority required (STREAMLINED) 1708 Ulcerative colitis where hypersensitivity to sulfonamides exists; 1709 Ulcerative colitis where intolerance to sulfasalazine exists. Note Not for the treatment of Crohn disease. Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
8845M NP	Capsule 750 mg	180	5	..	124.85	34.20	Colazide PK
MESALAZINE Authority required (STREAMLINED) 1708 Ulcerative colitis where hypersensitivity to sulfonamides exists; 1709 Ulcerative colitis where intolerance to sulfasalazine exists; 2268 Crohn disease where hypersensitivity to sulfonamides exists; 2269 Crohn disease where intolerance to sulfasalazine exists. Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
1611T NP	Tablet 250 mg (enteric coated)	100	5	..	93.43	34.20	Mesasal GK

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
2214M NP	Tablet 500 mg (prolonged release)	200	5	..	*297.44	34.20	Pentasa	FP
2234N NP	Sachet containing prolonged release granules, 1 g per sachet	100	5	..	279.63	34.20	Pentasa	FP
2287J NP	Sachet containing prolonged release granules, 2 g per sachet	60	5	..	312.30	34.20	Pentasa	FP
3413P NP	Tablet 1 g (prolonged release)	120	5	..	*330.68	34.20	Pentasa	FP
8731M NP	Tablet 500 mg (enteric coated)	200	5	..	*297.44	34.20	Salofalk	OA

MESALAZINE

Authority required (STREAMLINED)

1708

Ulcerative colitis where hypersensitivity to sulfonamides exists;

1709

Ulcerative colitis where intolerance to sulfasalazine exists.

Note

Not for the treatment of Crohn disease.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8598M NP	Sachet containing granules, 500 mg per sachet	200	5	..	*297.44	34.20	Salofalk	OA
8599N NP	Sachet containing granules, 1 g per sachet	100	5	..	279.63	34.20	Salofalk	OA
9206M NP	Sachet containing granules, 1.5 g per sachet	60	5	..	244.92	34.20	Salofalk	OA
9353G NP	Tablet 1.2 g (prolonged release)	60	5	..	220.99	34.20	Mezavant	ZI

MESALAZINE

Restricted benefit

Acute episode of mild to moderate ulcerative proctitis.

Note

Not for the treatment of Crohn disease.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

5461K NP	Suppository (moulded) 1 g	30	1	..	136.39	34.20	Salofalk	OA
8752P NP	Suppository 1 g	28	1	..	127.72	34.20	Pentasa	FP

MESALAZINE

Authority required (STREAMLINED)

1707

Acute episode of mild to moderate ulcerative colitis.

Note

Not for the treatment of Crohn disease.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Alimentary tract and metabolism

					Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	\$	\$	Brand Name and Manufacturer	
Note								
No applications for increased maximum quantities and/or repeats will be authorised.								
8616L NP	Enemas 2 g in 60 mL, 7	4	1	..	*336.22	34.20	Salofalk	OA
8617M NP	Enemas 4 g in 60 mL, 7	4	1	..	*445.90	34.20	Salofalk	OA
8753Q NP	Enemas 1 g in 100 mL, 7	4	1	..	*336.22	34.20	Pentasa	FP
8768L NP	Rectal foam 1 g per applicatorful, 14 applications, aerosol 80 g	4	1	..	*336.22	34.20	Salofalk	OA

OLSALAZINE SODIUM

Authority required (STREAMLINED)

1708

Ulcerative colitis where hypersensitivity to sulfonamides exists;

1709

Ulcerative colitis where intolerance to sulfasalazine exists.

Note

Not for the treatment of Crohn disease.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1728Y NP	Capsule 250 mg	100	5	..	61.41	34.20	Dipentum	UC
8086N NP	Tablet 500 mg	100	5	..	103.29	34.20	Dipentum	UC

SULFASALAZINE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2093E NP	Tablet 500 mg	200	5	..	*51.08	34.20	Salazopyrin	PF
2096H NP	Tablet 500 mg (enteric coated)	200	5	..	*55.12	34.20	^a Pyralin EN	KR
				^B 1.96	*57.08	34.20	^a Salazopyrin-EN	PF

SULFASALAZINE

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9208P	Tablet 500 mg	200	11	..	*51.08	34.20	Salazopyrin	PF
9209Q	Tablet 500 mg (enteric coated)	200	11	..	*55.12	34.20	^a Pyralin EN	KR
				^B 1.96	*57.08	34.20	^a Salazopyrin-EN	PF

Digestives, incl. enzymes

Digestives, incl. enzymes

Enzyme preparations

PANCREATIC EXTRACT

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8020D NP	Capsule (containing enteric coated minimicrospheres) providing not less than 10,000 BP units of lipase activity	500	10	..	*170.77	34.20	Creon 10,000	SM
8021E NP	Capsule (containing enteric coated minimicrospheres) providing not less than 25,000 BP units of lipase activity	200	10	..	*137.90	34.20	Creon 25,000	SM
8556H NP	Capsule (containing enteric coated minimicrospheres) providing not less than 5,000 BP units of lipase activity	500	10	..	*119.22	34.20	Creon 5000	SM
9412J NP	Capsule (containing enteric coated minimicrospheres) providing not less than 40,000 BP units of lipase activity	200	10	..	*215.62	34.20	Creon 40,000	SM

PANCREATIC EXTRACT

5453B	Granules (enteric coated) providing not less than 5,000 BP units of lipase activity per 100 mg, 20 g	3	10	..	*141.78	34.20	Creon Micro	SM

PANCREATIC EXTRACT

Restricted benefit

For use in patients with cystic fibrosis, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

5454C	Granules (enteric coated) providing not less than 5,000 BP units of lipase activity per 100 mg, 20 g	3	21	..	*141.78	34.20	Creon Micro	SM
9225M	Capsule (containing enteric coated minimicrospheres) providing not less than 5,000 BP units of lipase activity	500	21	..	*119.22	34.20	Creon 5000	SM
9226N	Capsule (containing enteric coated minimicrospheres) providing not less than 10,000 BP units of lipase activity	500	21	..	*170.77	34.20	Creon 10,000	SM
9227P	Capsule (containing enteric coated minimicrospheres) providing not less than 25,000 BP units of lipase activity	200	21	..	*137.90	34.20	Creon 25,000	SM
9413K	Capsule (containing enteric coated minimicrospheres) providing not less than 40,000 BP units of lipase activity	200	21	..	*215.62	34.20	Creon 40,000	SM

PANCRELIPASE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8366H NP	Capsule (containing enteric coated microtablets) providing not less than 25,000 BP units of lipase activity	200	10	..	*137.90	34.20	Panzytrat 25000	TM

PANCRELIPASE

Restricted benefit

For use in patients with cystic fibrosis, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9229R	Capsule (containing enteric coated microtablets) providing not less than 25,000 BP units of lipase activity	200	21	..	*137.90	34.20	Panzytrat 25000	TM

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Drugs used in diabetes								
Insulins and analogues								
Insulins and analogues for injection, fast-acting								
INSULIN ASPART								
8435Y NP	Injections (human analogue) 100 units per mL, 3 mL, 5	5	1	..	*264.22	34.20	NovoRapid FlexPen	NF
							NovoRapid Penfill 3 mL	NO
8571D NP	Injection (human analogue) 100 units per mL, 10 mL	5	2	..	*159.27	34.20	NovoRapid	NO
INSULIN GLULISINE								
1921D NP	Injections (human analogue) 100 units per mL, 3 mL, 5	5	1	..	*264.22	34.20	Apidra	AV
							Apidra SoloStar	SW
9224L NP	Injection (human analogue) 100 units per mL, 10 mL	5	2	..	*159.27	34.20	Apidra	SW
INSULIN LISPRO								
8084L NP	Injection (human analogue) 100 units per mL, 10 mL	5	2	..	*159.27	34.20	Humalog	LY
8212F NP	Injections (human analogue) 100 units per mL, 3 mL, 5	5	1	..	*264.22	34.20	Humalog	LY
							Humalog KwikPen	KP
INSULIN NEUTRAL								
1531N NP	Injection (human) 100 units per mL, 10 mL	5	2	..	*133.82	34.20	Actrapid	NO
							Humulin R	LY
1713E NP	Injection (bovine) 100 units per mL, 10 mL	5	2	..	*172.02	34.20	Hypurin Neutral	AS
1762R NP	Injections (human) 100 units per mL, 3 mL, 5	5	1	..	*224.32	34.20	Actrapid Penfill 3 mL	NO
							Humulin R	LY
Insulins and analogues for injection, intermediate-acting								
INSULIN ISOPHANE (N.P.H.)								
1533Q NP	Injection (human) 100 units per mL, 10 mL	5	2	..	*133.82	34.20	Humulin NPH	LY
							Protaphane	NO
1711C NP	Injection (bovine) 100 units per mL, 10 mL	5	2	..	*172.02	34.20	Hypurin Isophane	AS
1761Q NP	Injections (human) 100 units per mL, 3 mL, 5	5	1	..	*224.32	34.20	Humulin NPH	LY
							Protaphane InnoLet	NI
							Protaphane NovoLet 3 mL	NL
							Protaphane Penfill 3 mL	NO
Insulins and analogues for injection, intermediate-acting combined with fast-acting								
INSULIN ASPART—INSULIN ASPART PROTAMINE SUSPENSION								
8609D NP	Injections (human analogue) 100 units (30 units- 70 units) per mL, 3 mL, 5	5	1	..	*264.22	34.20	NovoMix 30 FlexPen	NF
							NovoMix 30 Penfill 3 mL	NO
INSULIN LISPRO—INSULIN LISPRO PROTAMINE SUSPENSION								
8390N	Injections (human analogue) 100 units (25 units-	5	1	..	*264.22	34.20	Humalog Mix25	LY

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
<i>NP</i>	75 units) per mL, 3 mL, 5						Humalog Mix25 KwikPen	KP
8874C <i>NP</i>	Injections (human analogue) 100 units (50 units- 50 units) per mL, 3 mL, 5	5	1	..	*264.22	34.20	Humalog Mix50	LY
							Humalog Mix50 KwikPen	KP
INSULIN NEUTRAL—INSULIN ISOPHANE (N.P.H.), (MIXED) (Biphasic Isophane)								
1426C <i>NP</i>	Injection (human) 100 units (30 units-70 units) per mL, 10 mL	5	2	..	*133.82	34.20	Humulin 30/70	LY
1763T <i>NP</i>	Injections (human) 100 units (30 units-70 units) per mL, 3 mL, 5	5	1	..	*224.32	34.20	Humulin 30/70	LY
							Mixtard 30/70 InnoLet	NI
							Mixtard 30/70 Penfill 3 mL	NO
2062M <i>NP</i>	Injections (human) 100 units (50 units-50 units) per mL, 3 mL, 5	5	1	..	*224.32	34.20	Mixtard 50/50 Penfill 3 mL	NO

Insulins and analogues for injection, long-acting

INSULIN DETEMIR

Restricted benefit

Type 1 diabetes.

9040T <i>NP</i>	Injections (human analogue) 100 units per mL, 3 mL, 5	5	1	..	*432.72	34.20	Levemir FlexPen	NF
							Levemir Penfill	NO

INSULIN GLARGINE

9039R <i>NP</i>	Injections (human analogue) 100 units per mL, 3 mL, 5	5	1	..	*432.72	34.20	Lantus	SW
							Lantus SoloStar	AV

Blood glucose lowering drugs, excl. insulins

Biguanides

METFORMIN HYDROCHLORIDE

1801T <i>NP</i>	Tablet 850 mg	60	5	..	13.09	14.16	^a Ascent Pharmaceuticals Limited	GN
							^a Chem mart Metformin	CH
							^a Diaformin 850	AF
							^a Formet 850	SI
							^a GenRx Metformin	GX
							^a Glucohexal	HX
							^a Metformin 850	CR
							^a Metformin-GA	GM
							^a Metformin generichealth	GQ
							^a Metformin Ranbaxy	RA
							^a Metformin Sandoz	SZ
							^a Terry White Chemists Metformin	TW
				^B 1.10	14.19	14.16	^a Glucophage	MQ
				^B 1.79	14.88	14.16	^a Diabex 850	AL
2430X <i>NP</i>	Tablet 500 mg	100	5	..	13.09	14.16	^a Ascent Pharmaceuticals Limited	GN
							^a Chem mart	CH

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
							Metformin
							^a Diaformin AF
							^a Formet 500 SI
							^a GenRx Metformin GX
							^a Glucohexal HX
							^a Metformin 500 CR
							^a Metformin-GA GM
							^a Metformin GQ
							^a generichealth
							^a Metformin RA
							^a Ranbaxy
							^a Metformin Sandoz SZ
							^a Terry White Chemists TW
				^B 1.10	14.19	14.16	^a Metformin
				^B 1.79	14.88	14.16	^a Glucophage MQ
				..	17.09	18.16	^a Diabex AL
3439B NP	Tablet 1 g (extended release)	60	5	..	17.09	18.16	^a Diabex XR 1000 AL
8607B NP	Tablet 1 g	90	5	..	18.04	19.11	^a Diaformin 1000 AF
							^a Formet 1000 SI
							^a Glucohexal HX
							^a Metformin-GA GM
							^a Metformin GQ
							^a generichealth
							^a 1000
							^a Metformin RA
							^a Ranbaxy 1000
							^a Metformin Sandoz SZ
							^a Pharmacor CR
				^B 1.80	19.84	19.11	^a Metformin 1000
9435N NP	Tablet 500 mg (extended release)	120	5	..	17.09	18.16	^a Diabex 1000 AL
							^a Diabex XR AL
							^a Diaformin XR AF
							^a Metex XR SI

Sulfonamides, urea derivatives

GLIBENCLAMIDE

Caution

Sulfonylureas may cause hypoglycaemia, particularly in the elderly.

2939Q NP	Tablet 5 mg	100	5	..	11.49	12.56	^a Glimel AF
				^B 1.44	12.93	12.56	^a Daonil SW

GLICLAZIDE

Caution

Sulfonylureas may cause hypoglycaemia, particularly in the elderly.

2449X NP	Tablet 80 mg	100	5	..	13.52	14.59	^a Chem mart CH
							^a Gliclazide
							^a GenRx Gliclazide GX
							^a Glyade AF
							^a Mellihexal SZ
							^a Nidem SI
							^a Terry White Chemists TW
							^a Gliclazide
8535F	Tablet 30 mg (modified release)	100	5	..	13.72	14.79	^a APO-Gliclazide MR TX

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
NP							
							^a Chem mart CH
							^a Gliclazide MR AF
							^a Glyade MR RA
							^a Oziclide MR RA
							^a Terry White TW
							^a Chemists
							^a Gliclazide MR
9302N	Tablet 60 mg (modified release)	60	5	..	15.18	16.25	Diamcron 60mg SE
NP							MR
GLIMEPIRIDE							
Caution							
Sulfonylureas may cause hypoglycaemia, particularly in the elderly.							
8450R	Tablet 1 mg	30	5	..	9.10	10.17	^a APO-Glimepiride TX
NP							
							^a Aylide 1 AF
							^a Diapride 1 SI
							^a Dimirel AV
							^a Glimepiride Sandoz SZ
				^B 2.81	11.91	10.17	^a Amaryl SW
8451T	Tablet 2 mg	30	5	..	11.55	12.62	^a APO-Glimepiride TX
NP							
							^a Aylide 2 AF
							^a Diapride 2 SI
							^a Dimirel AV
							^a Glimepiride Sandoz SZ
				^B 2.81	14.36	12.62	^a Amaryl SW
8452W	Tablet 4 mg	30	5	..	14.46	15.53	^a APO-Glimepiride TX
NP							
							^a Aylide 4 AF
							^a Diapride 4 SI
							^a Dimirel AV
							^a Glimepiride Sandoz SZ
				^B 2.80	17.26	15.53	^a Amaryl SW
8533D	Tablet 3 mg	30	5	..	13.00	14.07	^a APO-Glimepiride TX
NP							
							^a Aylide 3 AF
							^a Diapride 3 SI
							^a Dimirel AV
							^a Glimepiride Sandoz SZ
				^B 2.80	15.80	14.07	^a Amaryl SW
GLIPIZIDE							
Caution							
Sulfonylureas may cause hypoglycaemia, particularly in the elderly.							
2440K	Tablet 5 mg	100	5	..	11.58	12.65	^a Melizide AF
NP							
				^B 3.91	15.49	12.65	^a Minidiab PF

Combinations of oral blood glucose lowering drugs

METFORMIN HYDROCHLORIDE with GLIBENCLAMIDE

Caution

Sulfonylureas may cause hypoglycaemia, particularly in the elderly.

8810Q	Tablet 500 mg-2.5 mg	90	5	..	15.51	16.58	Glucovance	AL
NP							500mg/2.5mg	
8811R	Tablet 500 mg-5 mg	90	5	..	16.53	17.60	Glucovance	AL

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
NP							500mg/5mg
8838E	Tablet 250 mg-1.25 mg	90	5	..	13.24	14.31	Glucovance
NP							250mg/1.25mg

ROSIGLITAZONE with METFORMIN

Note

Rosiglitazone with metformin fixed dose combination tablet is not PBS-subsidised when used in combination with insulin.

Authority required (STREAMLINED)

3544

Type 2 diabetes in a patient whose HbA1c is greater than 7% prior to initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or a glucagon-like peptide-1 despite treatment with metformin and where a sulfonylurea is contraindicated or not tolerated.

The date and level of the qualifying HbA1c must be documented in the patient's medical records at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) clinical conditions with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) red cell transfusion within the previous 3 months.

A patient in these circumstances will be eligible for treatment where blood glucose monitoring over a 2 week period shows blood glucose levels greater than 10 mmol per L in more than 20% of tests. The results of this blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone or a glucagon-like peptide-1, must be documented in the patient's medical records.

9059T	Tablet containing 2 mg rosiglitazone (as maleate) with 500 mg metformin hydrochloride	56	5	..	65.09	34.20	Avandamet	GK
NP								
9060W	Tablet containing 2 mg rosiglitazone (as maleate) with 1 g metformin hydrochloride	56	5	..	68.43	34.20	Avandamet	GK
NP								
9061X	Tablet containing 4 mg rosiglitazone (as maleate) with 500 mg metformin hydrochloride	56	5	..	94.76	34.20	Avandamet	GK
NP								
9062Y	Tablet containing 4 mg rosiglitazone (as maleate) with 1 g metformin hydrochloride	56	5	..	98.09	34.20	Avandamet	GK
NP								

SITAGLIPTIN with METFORMIN

Note

Sitagliptin with metformin fixed dose combination tablet is not PBS-subsidised for use in combination with a sulfonylurea (triple oral therapy), as initial therapy or in combination with a thiazolidinedione (glitazone).

Authority required (STREAMLINED)

3543

Type 2 diabetes in a patient whose HbA1c is greater than 7% prior to initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or a glucagon-like peptide-1 despite treatment with metformin and where a combination of metformin and a sulfonylurea is contraindicated or not tolerated.

The date and level of the qualifying HbA1c must be documented in the patient's medical records at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) clinical conditions with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) red cell transfusion within the previous 3 months.

A patient in these circumstances will be eligible for treatment where blood glucose monitoring over a 2 week period shows blood glucose levels greater than 10 mmol per L in more than 20% of tests. The results of this blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone or a glucagon-like peptide-1, must be documented in the patient's medical records.

Authority required (STREAMLINED)

3149

Continuation of therapy in type 2 diabetes mellitus in a patient who has previously received and been stabilised on a PBS-subsidised regimen of oral diabetic medicines which includes metformin and sitagliptin.

9449H	Tablet containing 50 mg sitagliptin (as phosphate monohydrate) with 500 mg metformin hydrochloride	56	5	..	94.82	34.20	Janumet	MK
NP								
9450J	Tablet containing 50 mg sitagliptin (as phosphate monohydrate) with 850 mg metformin hydrochloride	56	5	..	97.28	34.20	Janumet	MK
NP								
9451K	Tablet containing 50 mg sitagliptin (as phosphate monohydrate) with 1000 mg metformin hydrochloride	56	5	..	97.90	34.20	Janumet	MK
NP								

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Alpha glucosidase inhibitors

ACARBOSE

8188Y NP	Tablet 50 mg	90	5	..	30.90	31.97	Glucobay 50	BN
8189B NP	Tablet 100 mg	90	5	..	40.92	34.20	Glucobay 100	BN

Thiazolidinediones

PIOGLITAZONE

Note

Pioglitazone hydrochloride is not PBS-subsidised as monotherapy.

Authority required (STREAMLINED)

3540

Dual oral combination therapy with metformin or a sulfonylurea

Type 2 diabetes, in combination with either metformin or a sulfonylurea, in a patient whose HbA1c is greater than 7% prior to initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or a glucagon-like peptide-1 despite treatment with either metformin or a sulfonylurea and where a combination of metformin and a sulfonylurea is contraindicated or not tolerated.

The date and level of the qualifying HbA1c must be documented in the patient's medical records at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) clinical conditions with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) red cell transfusion within the previous 3 months.

A patient in these circumstances will be eligible for treatment where blood glucose monitoring over a 2 week period shows blood glucose levels greater than 10 mmol per L in more than 20% of tests. The results of this blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone or a glucagon-like peptide-1, must be documented in the patient's medical records.

Authority required (STREAMLINED)

3541

Combination therapy with insulin

Type 2 diabetes, in combination with insulin, in a patient whose HbA1c is greater than 7% prior to initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or a glucagon-like peptide-1 despite treatment with insulin and oral anti-diabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c must be documented in the patient's medical records at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) clinical conditions with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) red cell transfusion within the previous 3 months.

A patient in these circumstances will be eligible for treatment where blood glucose monitoring over a 2 week period shows blood glucose levels greater than 10 mmol per L in more than 20% of tests. The results of this blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone or a glucagon-like peptide-1, must be documented in the patient's medical records.

Authority required (STREAMLINED)

3542

Triple oral combination therapy with metformin and a sulfonylurea

Type 2 diabetes, in combination with metformin and a sulfonylurea, in a patient whose HbA1c is greater than 7% prior to initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or a glucagon-like peptide-1 despite treatment with maximally tolerated doses of metformin and a sulfonylurea.

The date and level of the qualifying HbA1c must be documented in the patient's medical records at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) clinical conditions with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) red cell transfusion within the previous 3 months.

A patient in these circumstances will be eligible for treatment where blood glucose monitoring over a 2 week period shows blood glucose levels greater than 10 mmol per L in more than 20% of tests. The results of this blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone or a glucagon-like peptide-1, must be documented in the patient's medical records.

8694N NP	Tablet 15 mg (as hydrochloride)	28	5	..	61.52	34.20	Actos	LY
8695P NP	Tablet 30 mg (as hydrochloride)	28	5	..	91.19	34.20	Actos	LY

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
8696Q NP	Tablet 45 mg (as hydrochloride)	28	5	..	116.64	34.20	Actos LY

ROSIGLITAZONE

Note

Rosiglitazone maleate is not PBS-subsidised as monotherapy.

Authority required (STREAMLINED)

3540

Dual oral combination therapy with metformin or a sulfonylurea

Type 2 diabetes, in combination with either metformin or a sulfonylurea, in a patient whose HbA1c is greater than 7% prior to initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or a glucagon-like peptide-1 despite treatment with either metformin or a sulfonylurea and where a combination of metformin and a sulfonylurea is contraindicated or not tolerated.

The date and level of the qualifying HbA1c must be documented in the patient's medical records at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) clinical conditions with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) red cell transfusion within the previous 3 months.

A patient in these circumstances will be eligible for treatment where blood glucose monitoring over a 2 week period shows blood glucose levels greater than 10 mmol per L in more than 20% of tests. The results of this blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone or a glucagon-like peptide-1, must be documented in the patient's medical records.

8689H NP	Tablet 4 mg (as maleate)	28	5	..	61.52	34.20	Avandia GK
8690J NP	Tablet 8 mg (as maleate)	28	5	..	91.19	34.20	Avandia GK

Dipeptidyl peptidase 4 (DPP-4) inhibitors

SITAGLIPTIN

Note

Sitagliptin is not PBS-subsidised for use in combination with metformin and a sulfonylurea (triple oral therapy), as monotherapy or in combination with a thiazolidinedione (glitazone).

Authority required (STREAMLINED)

3540

Dual oral combination therapy with metformin or a sulfonylurea

Type 2 diabetes, in combination with either metformin or a sulfonylurea, in a patient whose HbA1c is greater than 7% prior to initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or a glucagon-like peptide-1 despite treatment with either metformin or a sulfonylurea and where a combination of metformin and a sulfonylurea is contraindicated or not tolerated.

The date and level of the qualifying HbA1c must be documented in the patient's medical records at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) clinical conditions with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) red cell transfusion within the previous 3 months.

A patient in these circumstances will be eligible for treatment where blood glucose monitoring over a 2 week period shows blood glucose levels greater than 10 mmol per L in more than 20% of tests. The results of this blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone or a glucagon-like peptide-1, must be documented in the patient's medical records.

9180E NP	Tablet 25 mg (as phosphate monohydrate)	28	5	..	91.19	34.20	Januvia MK
9181F NP	Tablet 50 mg (as phosphate monohydrate)	28	5	..	91.19	34.20	Januvia MK
9182G NP	Tablet 100 mg (as phosphate monohydrate)	28	5	..	91.19	34.20	Januvia MK

VILDAGLIPTIN

Note

Vildagliptin is not PBS-subsidised for use in combination with metformin and a sulfonylurea (triple oral therapy), as monotherapy or in combination with a thiazolidinedione (glitazone).

Authority required (STREAMLINED)

3540

Dual oral combination therapy with metformin or a sulfonylurea

Type 2 diabetes, in combination with either metformin or a sulfonylurea, in a patient whose HbA1c is greater than 7% prior to initiation of a

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or a glucagon-like peptide-1 despite treatment with either metformin or a sulfonylurea and where a combination of metformin and a sulfonylurea is contraindicated or not tolerated.						
	The date and level of the qualifying HbA1c must be documented in the patient's medical records at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated.						
	Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances: (a) clinical conditions with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or (b) red cell transfusion within the previous 3 months. A patient in these circumstances will be eligible for treatment where blood glucose monitoring over a 2 week period shows blood glucose levels greater than 10 mmol per L in more than 20% of tests. The results of this blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone or a glucagon-like peptide-1, must be documented in the patient's medical records.						
3415R	Tablet 50 mg	60	5	..	97.24	34.20	Galvus NV

Other blood glucose lowering drugs, excl. insulins

EXENATIDE

Authority required

Dual combination therapy with metformin or a sulfonylurea

Initiation of therapy, in combination with either metformin or a sulfonylurea, in a patient with type 2 diabetes who has an HbA1c greater than 7% prior to initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or a glucagon-like peptide-1 and in whom a combination of metformin and a sulfonylurea is contraindicated or not tolerated.

The date and level of the HbA1c must be documented in the patient's medical records at the time therapy with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated;

Continuation of therapy, in combination with either metformin or a sulfonylurea, in a patient with type 2 diabetes where the patient has previously been issued with an authority prescription for exenatide.

Note

Exenatide is not PBS-subsidised as monotherapy or in combination with an insulin, a thiazolidinedione (glitazone) or a dipeptidyl peptidase 4 inhibitor (gliptin).

Blood glucose monitoring as an alternative assessment to HbA1c levels will be accepted in the following circumstances:

- (a) clinical conditions with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) red cell transfusion within the previous 3 months.

Patients in these circumstances will be eligible for treatment where blood glucose monitoring over a 2 week period shows blood glucose levels greater than 10 mmol per L in more than 20% of tests. The results of this blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone or a glucagon-like peptide-1, must be documented in the patient's medical record.

Authority required

Triple combination therapy with metformin and a sulfonylurea

Initiation of therapy, in combination with metformin and a sulfonylurea, in a patient with type 2 diabetes who has an HbA1c greater than 7% prior to initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or a glucagon-like peptide-1 despite maximally tolerated doses of metformin and a sulfonylurea.

The date and level of the HbA1c must be documented in the patient's medical records at the time therapy with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone or a glucagon-like peptide-1 is initiated;

Continuation of therapy, in combination with metformin and a sulfonylurea, in a patient with type 2 diabetes where the patient has previously been issued with an authority prescription for exenatide.

Note

Exenatide is not PBS-subsidised as monotherapy or in combination with an insulin, a thiazolidinedione (glitazone) or a dipeptidyl peptidase 4 inhibitor (gliptin).

Blood glucose monitoring as an alternative assessment to HbA1c levels will be accepted in the following circumstances:

- (a) clinical conditions with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) red cell transfusion within the previous 3 months.

Patients in these circumstances will be eligible for treatment where blood glucose monitoring over a 2 week period shows blood glucose levels greater than 10 mmol per L in more than 20% of tests. The results of this blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone or a glucagon-like peptide-1, must be documented in the patient's medical record.

Note

Special Pricing Arrangements apply.

3423E	Injection solution 5 micrograms per dose in pre-filled pen, 60 doses	1	5	..	176.39	34.20	Byetta 5 microgram	LY
NP								
3424F	Injection solution 10 micrograms per dose in pre-filled pen, 60 doses	1	5	..	176.39	34.20	Byetta 10 microgram	LY
NP								

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$ ^a	
				^B 2.72	*15.74	14.09	Slow-K NV
POTASSIUM CHLORIDE with POTASSIUM BICARBONATE							
3012M NP	Effervescent tablet 14 mmol potassium and 8 mmol chloride	60	1	..	15.31	16.38	Chlorvescent AS

Anabolic agents for systemic use

Anabolic steroids

Estren derivatives

NANDROLONE DECANOATE

Authority required

Monotherapy for osteoporosis, where other treatment has failed and where specialist advice confirms that this is the only suitable treatment option for the patient. Specialist advice need only be obtained for the first authority approval;

Monotherapy for osteoporosis, where other treatment is not tolerated and where specialist advice confirms that this is the only suitable treatment option for the patient. Specialist advice need only be obtained for the first authority approval;

Monotherapy for osteoporosis, where other treatment is contraindicated and where specialist advice confirms that this is the only suitable treatment option for the patient. Specialist advice need only be obtained for the first authority approval;

Patients receiving PBS-subsidised therapy with this drug for osteoporosis prior to 1 February 2004;

Patients on long-term treatment with corticosteroids.

Note

Monotherapy for the treatment of osteoporosis does not exclude calcium supplementation.

1671Y	Injection 50 mg in 1 mL, disposable syringe	1	7	..	21.20	22.27	Deca-Durabolin SH

Blood and blood forming organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	

Blood and blood forming organs

Antithrombotic agents

Antithrombotic agents

Vitamin K antagonists

WARFARIN SODIUM

Caution

The listed brands have NOT been shown to be bioequivalent and should not be interchanged.

2209G NP	Tablet 2 mg	50	2	..	12.77	13.84	Coumadin	SI
2211J NP	Tablet 5 mg	50	2	..	14.03	15.10	Coumadin	SI
2843P NP	Tablet 1 mg	50	2	..	12.42	13.49	Marevan Coumadin	FM SI
2844Q NP	Tablet 3 mg	50	2	..	12.69	13.76	Marevan Marevan	FM FM

Heparin group

DALTEPARIN SODIUM (Low Molecular Weight Heparin Sodium—porcine mucous)

2816F NP	Injection 5,000 units (anti-Xa) in 0.2 mL single dose pre-filled syringe	10	1	..	57.65	34.20	Fragmin	PF
8269F NP	Injection 10,000 units (anti-Xa) in 1 mL single dose pre-filled syringe	10	1	..	109.38	34.20	Fragmin	PF
8271H NP	Injection 7,500 units (anti-Xa) in 0.75 mL single dose pre-filled syringe	10	1	..	83.65	34.20	Fragmin	PF
8603T NP	Injection 2,500 units (anti-Xa) in 0.2 mL single dose pre-filled syringe	10	1	..	55.61	34.20	Fragmin	PF

DALTEPARIN SODIUM (Low Molecular Weight Heparin Sodium—porcine mucous)

Restricted benefit

Haemodialysis.

8641T NP	Injection 2,500 units (anti-Xa) in 0.2 mL single dose pre-filled syringe	20	3	..	*104.74	34.20	Fragmin	PF
8642W NP	Injection 5,000 units (anti-Xa) in 0.2 mL single dose pre-filled syringe	20	3	..	*108.88	34.20	Fragmin	PF
8643X NP	Injection 7,500 units (anti-Xa) in 0.75 mL single dose pre-filled syringe	20	3	..	*160.88	34.20	Fragmin	PF

ENOXAPARIN SODIUM

8262W NP	Injection 60 mg (6,000 i.u. anti-Xa) in 0.6 mL pre-filled syringe	10	1	..	79.68	34.20	Clexane	SW
8263X NP	Injection 80 mg (8,000 i.u. anti-Xa) in 0.8 mL pre-filled syringe	10	1	..	90.70	34.20	Clexane	SW
8264Y NP	Injection 100 mg (10,000 i.u. anti-Xa) in 1 mL pre-filled syringe	10	1	..	109.08	34.20	Clexane	SW
8510X NP	Injection 40 mg (4,000 i.u. anti-Xa) in 0.4 mL pre-filled syringe	20	..	..	*108.88	34.20	Clexane	SW
8558K NP	Injection 20 mg (2,000 i.u. anti-Xa) in 0.2 mL pre-filled syringe	20	..	..	*104.74	34.20	Clexane	SW
9195Y NP	Solution for injection 40 mg (4,000 i.u. anti-Xa) in 0.4 mL	20	..	..	*108.88	34.20	Clexane	SW

ENOXAPARIN SODIUM

Restricted benefit

Haemodialysis.

Blood and blood forming organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8639Q NP	Injection 40 mg (4,000 i.u. anti-Xa) in 0.4 mL pre-filled syringe	20	3	..	*108.88	34.20	Clexane	SW
8640R NP	Injection 60 mg (6,000 i.u. anti-Xa) in 0.6 mL pre-filled syringe	20	3	..	*152.94	34.20	Clexane	SW
8716R NP	Injection 20 mg (2,000 i.u. anti-Xa) in 0.2 mL pre-filled syringe	20	3	..	*104.74	34.20	Clexane	SW
9196B NP	Solution for injection 40 mg (4,000 i.u. anti-Xa) in 0.4 mL	20	3	..	*108.88	34.20	Clexane	SW
HEPARIN SODIUM								
1076P NP	Injection 35,000 units in 35 mL	12	5	..	*278.58	34.20	Hospira Pty Limited	HH
1463B NP	Injection (preservative-free) 5,000 units in 5 mL	50	5	..	66.93	34.20	Pfizer Australia Pty Ltd	PF
1466E NP	Injection 5,000 units in 0.2 mL	5	5	..	15.48	16.55	Hospira Pty Limited	HH

Platelet aggregation inhibitors excl. heparin

ABCIXIMAB

Authority required (STREAMLINED)

1716

Patients undergoing percutaneous coronary balloon angioplasty;

1717

Patients undergoing percutaneous coronary atherectomy;

1718

Patients undergoing percutaneous coronary stent placement.

8048N	I.V. injection 10 mg in 5 mL	3	..	..	*1453.11	34.20	ReoPro	LY

ASPIRIN

1010E NP	Tablet 300 mg (dispersible)	96	1	..	8.55	9.62	Solprin	RC
8202Q NP	Tablet 100 mg	112	1	..	8.06	9.13 ^a	DBL Aspirin 100 mg	YT
				^B 1.32	9.38	9.13 ^a	Astrix	YN

CLOPIDOGREL

Authority required (STREAMLINED)

1719

Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events in patients with a history of symptomatic cerebrovascular ischaemic episodes while on therapy with low-dose aspirin;

1720

Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events in patients where low-dose aspirin poses an unacceptable risk of gastrointestinal bleeding;

1721

Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events in patients where there is a history of anaphylaxis, urticaria or asthma within 4 hours of ingestion of aspirin, other salicylates, or NSAIDs;

1722

Prevention of recurrence of myocardial infarction or unstable angina in patients with a history of symptomatic cardiac ischaemic events while on therapy with low-dose aspirin;

1723

Prevention of recurrence of myocardial infarction or unstable angina in patients where low-dose aspirin poses an unacceptable risk of gastrointestinal bleeding;

1724

Prevention of recurrence of myocardial infarction or unstable angina in patients where there is a history of anaphylaxis, urticaria or asthma within 4 hours of ingestion of aspirin, other salicylates, or NSAIDs.

Note

Not for prophylaxis of DVT or peripheral arterial disease.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Blood and blood forming organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net		Brand Name and Manufacturer
					\$	\$		
	Note							
	Clopidogrel (as besilate) 75 mg tablets have been established to be bioequivalent to clopidogrel (as hydrogen sulfate) 75 mg tablets.							
8358X NP	Tablet 75 mg (as hydrogen sulfate)	28	5	..	71.61	34.20	^a	APO-Clopidogrel TX
							^a	Chem mart CH
							^a	Clopidogrel Sandoz SZ
							^a	Clopidogrel Winthrop WA
							^a	Iscover BQ
							^a	Piax AF
							^a	Plavix SW
							^a	Terry White Chemists Clopidogrel TW
9354H NP	Tablet 75 mg (as besilate)	28	5	..	71.61	34.20	^a	Clopidogrel Actavis GQ
							^a	Clopidogrel-GA GM
							^a	Clovix 75 SI

CLOPIDOGREL

Authority required (STREAMLINED)

3245

Treatment of acute coronary syndromes (myocardial infarction or unstable angina) in combination with aspirin;

3146

Treatment in combination with aspirin following cardiac stent insertion.

Note

Not for prophylaxis of DVT or peripheral arterial disease.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9317J NP	Tablet 75 mg (as hydrogen sulfate)	28	5	..	71.61	34.20	^a	Clopidogrel Winthrop WA
							^a	Iscover BQ
							^a	Plavix SW

CLOPIDOGREL with ASPIRIN

Authority required (STREAMLINED)

3246

Treatment of acute coronary syndromes (myocardial infarction or unstable angina);

3219

Treatment following cardiac stent insertion;

1722

Prevention of recurrence of myocardial infarction or unstable angina in patients with a history of symptomatic cardiac ischaemic events while on therapy with low-dose aspirin.

Note

Not for prophylaxis of DVT or peripheral arterial disease.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9296G NP	Tablet 75 mg (as hydrogen sulfate)-100 mg	30	5	..	76.27	34.20	^a	CoPlavix SW
							^a	DuoCover BQ

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
DIPYRIDAMOLE								
<u>Restricted benefit</u>								
Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events:								
(1) as adjunctive therapy with low-dose aspirin; or								
(2) where low-dose aspirin poses an unacceptable risk of gastrointestinal bleeding; or								
(3) where there is a history of anaphylaxis, urticaria or asthma within 4 hours of ingestion of aspirin, other salicylates, or NSAIDs.								
<u>Note</u>								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8335Q NP	Capsule 200 mg (sustained release)	60	5	..	36.96	34.20	Persantin SR	BY
DIPYRIDAMOLE with ASPIRIN								
<u>Restricted benefit</u>								
Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events.								
<u>Note</u>								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8382E NP	Capsule 200 mg (sustained release)-25 mg	60	5	..	37.19	34.20	Asasantin SR	BY
EPTIFIBATIDE ACETATE								
<u>Authority required (STREAMLINED)</u>								
1884								
Patients undergoing non-urgent percutaneous intervention with intracoronary stenting.								
8683B	Solution for I.V. injection 20 mg (base) in 10 mL	2	..	..	*262.54	34.20	Integrilin	SH
8684C	Solution for I.V. infusion 75 mg (base) in 100 mL	3	..	..	*1020.36	34.20	Integrilin	SH
PRASUGREL								
<u>Authority required (STREAMLINED)</u>								
3208								
Treatment of acute coronary syndrome (myocardial infarction or unstable angina) managed by percutaneous coronary intervention in combination with aspirin.								
<u>Note</u>								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
9495R NP	Tablet 5 mg (as hydrochloride)	28	5	..	96.43	34.20	Effient	LY
9496T NP	Tablet 10 mg (as hydrochloride)	28	5	..	106.43	34.20	Effient	LY
TICLOPIDINE HYDROCHLORIDE								
<u>Caution</u>								
Severe neutropenia is common in the early months of therapy. Haematological monitoring should be undertaken at commencement and every two weeks in the first four months of therapy.								
<u>Authority required (STREAMLINED)</u>								
1719								
Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events in patients with a history of symptomatic cerebrovascular ischaemic episodes while on therapy with low-dose aspirin;								
1720								
Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events in patients where low-dose aspirin poses an unacceptable risk of gastrointestinal bleeding;								
1721								
Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events in patients where there is a history of anaphylaxis, urticaria or asthma within 4 hours of ingestion of aspirin, other salicylates, or NSAIDs;								

Blood and blood forming organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
1260							
Patients established on this drug as a pharmaceutical benefit prior to 1 November 1999.							
<u>Note</u>							
Shared Care Model:							
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
2095G NP	Tablet 250 mg	60	5	..	140.93	34.20	Tilodene AF

TIROFIBAN HYDROCHLORIDE

Authority required (STREAMLINED)

1729

Patients with high risk unstable angina who have new transient or persistent ST-T ischaemic changes and anginal pain lasting longer than 20 minutes;

1730

Patients with high risk unstable angina who have new transient or persistent ST-T ischaemic changes and repetitive episodes of angina at rest or during minimal exercise in the previous 12 hours;

1275

Patients with non-Q-wave myocardial infarction.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8350L NP	Solution concentrate for I.V. infusion 12.5 mg (base) in 50 mL	1	2	..	363.11	34.20	Aggrastat AS

Enzymes

DROTRECOGIN ALFA (ACTIVATED)

Authority required

Adult patients with severe sepsis who have a high risk of death as determined by acute dysfunction in at least 2 organs or modified Apache II score of at least 25.

Acute organ dysfunction is defined as follows:

- (1) For cardiovascular-system dysfunction, an arterial systolic blood pressure of less than or equal to 90 mmHg or mean arterial pressure of less than or equal to 70 mmHg for at least 1 hour despite adequate fluid resuscitation, adequate intravascular volume status or the use of vasopressors in an attempt to maintain a systolic blood pressure of greater than or equal to 90 mmHg or a mean arterial pressure of greater than or equal to 70 mmHg;
- (2) For kidney dysfunction, urine output of less than 0.5 mL per kg of body weight per hour for 1 hour despite adequate fluid resuscitation;
- (3) For respiratory-system dysfunction, a ratio of PaO₂ to FiO₂ of less than or equal to 250;
- (4) For haematologic dysfunction, a platelet count of less than 80,000 per cubic millimetre or which has decreased by 50 percent in the previous 3 days;
- (5) In the case of unexplained metabolic acidosis, a pH of less than or equal to 7.30 or a base deficit of greater than or equal to 5.0 mmol per L in association with a plasma lactate level of greater than 1.5 times the upper limit of the normal value for the reporting laboratory.

Note

Medical practitioners should request the appropriate quantity of vials at the time of the authority application, according to the weight of the patient, to achieve a dose of 24 micrograms per kg per hour over a maximum of 96 hours.

8614J	Powder for I.V. infusion 5 mg	1	..	..	467.14	34.20	Xigris LY

RETEPLASE (Recombinant plasminogen activator)

Restricted benefit

Treatment of acute myocardial infarction within 6 hours of onset of attack.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8253J NP	Pack containing 2 vials powder for injection 10 units, 2 single use pre-filled syringes with solvent, 2 reconstitution spikes and 2 needles	±1	..	..	2066.96	34.20	Rapilysin 10 U TA

TENECTEPLASE

Restricted benefit

Treatment of acute myocardial infarction within 12 hours of onset of attack.

Blood and blood forming organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
<u>Note</u>							
Shared Care Model:							
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
8526R <i>NP</i>	Powder for injection 40 mg with solvent	1	..	..	1960.76	34.20	Metalyse BY
8527T <i>NP</i>	Powder for injection 50 mg with solvent	1	..	..	2057.06	34.20	Metalyse BY

Direct thrombin inhibitors

BIVALIRUDIN TRIFLUOROACETATE

Authority required (STREAMLINED)

3075

A patient undergoing percutaneous coronary intervention.

8844L	Powder for I.V. injection 250 mg (base)	1	..	..	671.75	34.20	Angiomax CS

DABIGATRAN ETEXILATE

Authority required

Prevention of venous thromboembolism in a patient undergoing total hip replacement.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9318K NP	Capsule 75 mg (as mesilate)	20	1	..	*81.16	34.20	Pradaxa BY
9319L NP	Capsule 110 mg (as mesilate)	20	1	..	*81.16	34.20	Pradaxa BY

DABIGATRAN ETEXILATE

Authority required

Prevention of venous thromboembolism in a patient undergoing total hip replacement.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

No applications for increased maximum quantities and/or repeats will be authorised for the pack of 60 capsules.

9320M NP	Capsules 75 mg (as mesilate), 60	£1	..	..	228.21	34.20	Pradaxa BY
9321N NP	Capsules 110 mg (as mesilate), 60	£1	..	..	228.21	34.20	Pradaxa BY

DABIGATRAN ETEXILATE

Authority required

Prevention of venous thromboembolism in a patient undergoing total knee replacement.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9322P NP	Capsule 75 mg (as mesilate)	20	..	..	*81.16	34.20	Pradaxa BY
9323Q NP	Capsule 110 mg (as mesilate)	20	..	..	*81.16	34.20	Pradaxa BY

Blood and blood forming organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Other antithrombotic agents

FONDAPARINUX SODIUM

Authority required (STREAMLINED)

2005

Prevention of venous thromboembolic events in patients undergoing major hip surgery;

2006

Prevention of venous thromboembolic events in patients undergoing total knee replacement.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8775W NP	Injection 2.5 mg in 0.5 mL single dose pre-filled syringe	7	..	..	140.54	34.20	Arixtra	GK

RIVAROXABAN

Authority required

Prevention of venous thromboembolism in a patient undergoing total hip replacement.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9465E NP	Tablets 10 mg, 10	£1	1	..	101.14	34.20	Xarelto	BN
9466F NP	Tablet 10 mg	15	1	..	148.66	34.20	Xarelto	BN

RIVAROXABAN

Authority required

Prevention of venous thromboembolism in a patient undergoing total hip replacement.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

No applications for increased maximum quantities and/or repeats will be authorised for the 30 tablet pack.

9467G NP	Tablets 10 mg, 30	£1	..	..	279.89	34.20	Xarelto	BN

RIVAROXABAN

Authority required

Prevention of venous thromboembolism in a patient undergoing total knee replacement.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9468H NP	Tablets 10 mg, 10	£1	..	..	101.14	34.20	Xarelto	BN
9469J NP	Tablet 10 mg	15	..	..	148.66	34.20	Xarelto	BN

Blood and blood forming organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Antihemorrhagics								
Antifibrinolytics								
Amino acids								
TRANEXAMIC ACID								
Note								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
2180R NP	Tablet 500 mg	100	2	..	51.68	34.20	Cyklokapron	PF
Antianemic preparations								
Iron preparations								
Iron bivalent, oral preparations								
FERROUS SULFATE								
8815Y NP	Oral liquid 30 mg per mL, 250 mL	‡1	2	..	19.35	20.42	Ferro-Liquid	AE
Iron trivalent, parenteral preparations								
IRON POLYMALTOSE COMPLEX								
2593L NP	Injection 100 mg (iron) in 2 mL	5	..	..	50.36	34.20 ^a	Ferrosig	SI
						34.20 ^a	Ferrum H	AS
IRON SUCROSE								
Authority required (STREAMLINED)								
2070								
Iron deficiency anaemia, in combination with either epoetin alfa or darbepoetin alfa, in patients undergoing chronic haemodialysis who have had a documented hypersensitivity reaction to iron polymaltose and in whom continued intravenous iron therapy is appropriate.								
8807M NP	Concentrate for solution for infusion 2.7 g (equivalent to 100 mg iron (III)) in 5 mL	5	..	..	139.48	34.20	Venofer	AS
Iron in combination with folic acid								
FERROUS FUMARATE with FOLIC ACID								
9011G NP	Tablet 310 mg (equivalent to 100 mg iron)-350 micrograms	60	1	..	12.79	13.86	Ferro-f-tab	AE
Vitamin B ₁₂ and folic acid								
Vitamin B ₁₂ (cyanocobalamin and derivatives)								
HYDROXOCOBALAMIN								
Restricted benefit								
Pernicious anaemia;								
Other proven vitamin B ₁₂ deficiencies;								
Prophylaxis after gastrectomy.								
Note								
One injection of hydroxocobalamin 1 mg every three months provides appropriate maintenance therapy in vitamin B ₁₂ deficiencies.								
9048F NP	Injection 1 mg (as chloride) in 1 mL	3	..	..	15.87	16.94 ^a	Hydroxo-B12	AS
						16.94 ^a	Neo-B12	HH
Folic acid and derivatives								
FOLIC ACID								
2958Q NP	Tablet 500 micrograms	200	..	..	*13.78	14.85	Megafof 0.5	AF

Blood and blood forming organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	
FOLIC ACID							
<u>Note</u>							
The 5 mg strength tablet should be used in malabsorption states only.							
1437P NP	Tablet 5 mg	200	1	..	*14.02	15.09	Megafol 5 AF

Blood substitutes and perfusion solutions

Blood and related products

Blood substitutes and plasma protein fractions

GELATIN - SUCCINYLATED								
8444K NP	I.V. infusion 20 g per 500 mL, 500 mL	3	..	..	*45.75	34.20	Gelofusine	BR
HYDROXYETHYL STARCH 130/0.4								
9487H NP	I.V. infusion 30 g per 500 mL, 500 mL	3	..	..	*45.75	34.20	Voluven 6%	PK
POLYGELINE								
2334W NP	I.V. infusion 17.5 g per 500 mL (3.5%) with electrolytes, 500 mL	3	..	..	*45.75	34.20	Haemacel	AE

I.V. solutions

Solutions for parenteral nutrition

GLUCOSE								
2245E NP	I.V. infusion 278 mmol (anhydrous) per L (5%), 1 L	5	1	..	*23.17	24.24	^a B. Braun Australia Pty Ltd	BR
							^a Baxter Healthcare Pty Ltd	BX
							^a Fresenius Kabi Australia Pty Limited	PK
9444C NP	I.V. infusion 139 mmol (anhydrous) per 500 mL (5%), 500 mL	5	1	..	*18.07	19.14	^a B. Braun Australia Pty Ltd	BR
							^a Fresenius Kabi Australia Pty Limited	PK
9445D NP	I.V. infusion 278 mmol (anhydrous) per 500 mL (10%), 500 mL	5	1	..	*18.07	19.14	Fresenius Kabi Australia Pty Limited	PK
9474P NP	I.V. infusion 69.5 mmol (anhydrous) per 250 mL (5%), 250 mL	5	1	..	*24.02	25.09	^a B. Braun Australia Pty Ltd	BR
							^a Glucose 5% Freeflex	PK

Solutions affecting the electrolyte balance

ELECTROLYTE REPLACEMENT SOLUTION								
3199J NP	I.V. infusion 1 L	2	1	..	*21.96	23.03	Plasma-Lyte 148	BX
SODIUM CHLORIDE								
2260Y NP	I.V. infusion 513 mmol per L (3%), 1 L	2	1	..	*16.54	17.61	Baxter Healthcare Pty Ltd	BX
2264E NP	I.V. infusion 154 mmol per L (0.9%), 1 L	5	1	..	*23.17	24.24	^a B. Braun Australia Pty Ltd	BR
							^a Baxter Healthcare Pty Ltd	BX
							^a Fresenius Kabi Australia Pty Limited	PK

Blood and blood forming organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
9392H NP	I.V. infusion 77 mmol per 500 mL (0.9%), 500 mL	5	1	..	*18.07	19.14	^a B. Braun Australia Pty Ltd	BR
							^a Fresenius Kabi Australia Pty Limited	PK
9473N NP	I.V. infusion 38.5 mmol per 250 mL (0.9%), 250 mL	5	1	..	*24.02	25.09	^a B. Braun Australia Pty Ltd	BR
							^a Sodium Chloride 0.9% Freeflex	PK
SODIUM CHLORIDE COMPOUND								
2266G NP	I.V. infusion 1 L	4	1	..	*30.02	31.09	Baxter Healthcare Pty Ltd	BX
SODIUM CHLORIDE with GLUCOSE								
2278X NP	I.V. infusion 39 mmol-69 mmol (anhydrous) per 500 mL (0.45%-2.5%), 500 mL	5	1	..	*28.77	29.84	Baxter Healthcare Pty Ltd	BX
2279Y NP	I.V. infusion 19 mmol-104 mmol (anhydrous) per 500 mL (0.225%-3.75%), 500 mL	5	1	..	*28.77	29.84	Baxter Healthcare Pty Ltd	BX
2281C NP	I.V. infusion 31 mmol-222 mmol (anhydrous) per L (0.18%-4%), 1 L	5	1	..	*23.52	24.59	Baxter Healthcare Pty Ltd	BX
SODIUM LACTATE COMPOUND								
2286H NP	I.V. infusion 1 L	5	1	..	*23.17	24.24	^a B. Braun Australia Pty Ltd	BR
							^a Baxter Healthcare Pty Ltd	BX
							^a Fresenius Kabi Australia Pty Limited	PK
9416N NP	I.V. infusion 500 mL	5	1	..	*18.07	19.14	^a B. Braun Australia Pty Ltd	BR
							^a Fresenius Kabi Australia Pty Limited	PK

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer

Cardiovascular system

Cardiac therapy

Cardiac glycosides

Digitalis glycosides

DIGOXIN

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1322N NP	Tablet 250 micrograms	100	1	..	10.80	11.87 ^a	Sigmaxin	FM
				^B 3.00	13.80	11.87 ^a	Lanoxin	SI
2605D NP	Tablet 62.5 micrograms	200	1	..	10.50	11.57 ^a	Sigmaxin-PG	FM
				^B 3.00	13.50	11.57 ^a	Lanoxin-PG	SI
3164M NP	Oral solution for children 50 micrograms per mL, 60 mL	2	3	..	*28.68	29.75	Lanoxin	SI

Antiarrhythmics, class I and III

Antiarrhythmics, class IA

DISOPYRAMIDE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2923W NP	Capsule 100 mg	100	5	..	29.13	30.20	Rythmodan	SW
2924X NP	Capsule 150 mg	100	5	..	46.51	34.20	Rythmodan	SW

Antiarrhythmics, class IB

LIGNOCAINE HYDROCHLORIDE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2875H NP	Injection 100 mg in 5 mL	5	..	..	37.33	34.20	Pfizer Australia Pty Ltd	PF
2876J NP	Infusion 500 mg in 5 mL	10	..	..	29.59	30.66	Xylocard 500	AP

Antiarrhythmics, class IC

FLECAINIDE ACETATE

Caution

Flecainide acetate should be avoided in patients with poor cardiac function.

Restricted benefit

Serious supra-ventricular cardiac arrhythmias;

Serious ventricular cardiac arrhythmias where treatment is initiated in a hospital (in-patient or out-patient).

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1088G NP	Tablet 50 mg	60	5	..	38.39	34.20	Tambocor	IA
1090J NP	Tablet 100 mg	60	5	..	45.38	34.20 ^a	Flecatab	AF
						^a	Tambocor	IA

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Antiarrhythmics, class III							
AMIODARONE HYDROCHLORIDE							
Caution							
Amiodarone hydrochloride has been reported to cause frequent and potentially serious toxicity. Regular monitoring of hepatic and thyroid function is recommended.							
Restricted benefit							
Severe cardiac arrhythmias.							
Note							
Shared Care Model:							
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
2343H NP	Tablet 200 mg	30	5	..	21.72	22.79	^a Aratac 200 AF
							^a Cardinorm SZ
							^a Chem mart CH
							^a Amiodarone Cordarone X 200 SW
							^a GenRx Amiodarone GX
							^a Rithmik 200 SI
							^a Terry White Chemists TW
							^a Amiodarone Aratac 100 AF
2344J NP	Tablet 100 mg	30	5	..	15.02	16.09	^a Cardinorm SZ
							^a Cordarone X 100 SW
							^a Rithmik 100 SI
SOTALOL HYDROCHLORIDE							
Restricted benefit							
Severe cardiac arrhythmias.							
Note							
Shared Care Model:							
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
2043M NP	Tablet 160 mg	60	5	..	26.50	27.57	^a Cardol AF
							^a Chem mart Sotalol CH
							^a GenRx Sotalol GX
							^a Solavert SI
							^a Sotalol Sandoz SZ
							^a Terry White Chemists Sotalol TW
				^B 5.00	31.50	27.57	^a Sotacor FM
8398B NP	Tablet 80 mg	60	5	..	15.78	16.85	^a GenRx Sotalol GX
							^a Solavert SI
							^a Sotalol Sandoz SZ
				^B 5.00	20.78	16.85	^a Sotacor FM

Cardiac stimulants excl. cardiac glycosides Adrenergic and dopaminergic agents

1016L NP	ADRENALINE Injection 1 mg in 1 mL (1 in 1,000)	5	1	..	20.34	21.41	AstraZeneca Pty Ltd AP

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
ADRENALINE							
<u>Authority required</u>							
Initial sole PBS-subsidised supply for anticipated emergency treatment of acute allergic reactions with anaphylaxis in a patient who:							
(a) has been assessed to be at significant risk of anaphylaxis by, or in consultation with, a clinical immunologist, allergist, paediatrician or respiratory physician. The name of the specialist consulted must be provided at the time of application for initial supply; or							
(b) has been discharged from hospital or an emergency department after treatment with adrenaline for acute allergic reaction with anaphylaxis;							
Continuing sole PBS-subsidised supply for anticipated emergency treatment of acute allergic reactions with anaphylaxis, where the patient has previously been issued with an authority prescription for this drug.							
<u>Note</u>							
The auto-injector should be provided in the framework of a comprehensive anaphylaxis prevention program and an emergency action plan including training in recognition of the symptoms of anaphylaxis and the use of the auto-injector device. (For further information see the Australasian Society of Clinical Immunology and Allergy website at www.allergy.org.au .)							
<u>Note</u>							
Authority approvals will be limited to a maximum quantity of 2 auto-injectors (Anapen or EpiPen) at any one time.							
No repeats will be issued.							
<u>Caution</u>							
EpiPen and Anapen products have different administration techniques and should not be prescribed to the same patient without training in their use.							
3408J NP	I.M. injection 150 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	106.00	34.20	Anapen Junior LM
3409K NP	I.M. injection 300 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	106.00	34.20	Anapen LM
8697R NP	I.M. injection 150 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	106.00	34.20	EpiPen Jr. AL
8698T NP	I.M. injection 300 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	106.00	34.20	EpiPen AL

Vasodilators used in cardiac diseases

Organic nitrates

GLYCERYL TRINITRATE							
1459T NP	Tablets 600 micrograms, 100	1	5	..	15.00	16.07 ^a	Lycinate FM
				^B 3.00	18.00	16.07 ^a	Anginine Stabilised SI
1515R NP	Transdermal patch releasing approximately 5 mg per 24 hours	30	5	..	27.74	28.81	Transiderm-Nitro 25 NV
1516T NP	Transdermal patch releasing approximately 10 mg per 24 hours	30	5	..	34.38	34.20	Transiderm-Nitro 50 NV
8010N NP	Transdermal patch releasing approximately 5 mg per 24 hours	30	5	..	27.74	28.81	Nitro-Dur 5 SH
8011P NP	Transdermal patch releasing approximately 10 mg per 24 hours	30	5	..	34.38	34.20	Nitro-Dur 10 SH
8026K NP	Transdermal patch releasing approximately 15 mg per 24 hours	30	5	..	34.38	34.20	Nitro-Dur 15 SH
8027L NP	Transdermal patch releasing approximately 5 mg per 24 hours	30	5	..	27.74	28.81	Minitran 5 IA
8028M NP	Transdermal patch releasing approximately 10 mg per 24 hours	30	5	..	34.38	34.20	Minitran 10 IA
8119H NP	Transdermal patch releasing approximately 15 mg per 24 hours	30	5	..	34.38	34.20	Minitran 15 IA

GLYCERYL TRINITRATE

Note

The spray should not be inhaled.

8171C NP	Sublingual spray (pump pack) 400 micrograms per dose (200 doses)	1	5	..	20.13	21.20	Nitrolingual Pumpspray SW
ISOSORBIDE DINITRATE							
2587E NP	Tablet 10 mg	200	2	..	*13.82	14.89	Sorbidin AF
2588F NP	Sublingual tablet 5 mg	200	2	..	*14.72	15.79	Isordil Sublingual SI

Cardiovascular system

					Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$			
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium			Brand Name and Manufacturer		
ISOSORBIDE MONONITRATE									
1558B NP	Tablet 60 mg (sustained release)	30	5	..	12.88	13.95	^a	Chem mart	CH
								Isosorbide	
								Mononitrate	
							^a	Duride	AF
							^a	GenRx Isosorbide	GX
								Mononitrate	
							^a	Imtrate 60 mg	GM
							^a	Isomonit	SZ
^a	Monodur 60 mg	PM							
^a	Terry White	TW							
								Chemists	
								Isosorbide	
								Mononitrate	
				^B 2.84	15.72	13.95	^a	Imdur Durule	AP
8273K NP	Tablet 120 mg (sustained release)	30	5	..	21.74	22.81	^a	Monodur 120 mg	PM
							^B 3.00	24.74	22.81

Other vasodilators used in cardiac diseases

NICORANDIL

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8228C NP	Tablets 10 mg, 60	£1	5	..	24.14	25.21		Ikorel	SW
8229D NP	Tablets 20 mg, 60	£1	5	..	31.26	32.33		Ikorel	SW

PERHEXILINE MALEATE

Caution

Regular monitoring of drug serum levels is recommended.

Authority required (STREAMLINED)

1023

Angina not responding to other therapy.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1822X NP	Tablet 100 mg	100	5	..	62.62	34.20		Pexsig	SI

Antihypertensives

Antiadrenergic agents, centrally acting

Methyldopa

METHYLDOPA

1629R NP	Tablet 250 mg	100	5	..	13.65	14.72	^a	Hydopa	AF
				^B 2.65	16.30	14.72	^a	Aldomet	AS

Imidazoline receptor agonists

CLONIDINE

3141H NP	Tablet 150 micrograms	100	5	..	37.44	34.20		Catapres	BY
3145M NP	Tablet 100 micrograms	100	5	..	28.88	29.95		Catapres 100	BY

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
MOXONIDINE							
<u>Restricted benefit</u>							
Hypertension in patients receiving concurrent antihypertensive therapy.							
9019Q NP	Tablet 200 micrograms	30	5	..	19.53	20.60	Physiotens SM
9020R NP	Tablet 400 micrograms	30	5	..	28.78	29.85	Physiotens SM
Antiadrenergic agents, peripherally acting							
<i>Alpha-adrenoceptor antagonists</i>							
PRAZOSIN HYDROCHLORIDE							
1478T NP	Tablet 5 mg (base)	100	5	..	20.85	21.92	^a Chem mart CH
							^a Prazosin GX
							^a GenRx Prazosin TW
							^a Terry White Chemists Prazosin
				^B 3.29	24.14	21.92	^a Minipress PF
1479W NP	Tablet 1 mg (base)	100	5	..	12.62	13.69	^a Chem mart CH
							^a Prazosin GX
							^a GenRx Prazosin TW
							^a Terry White Chemists Prazosin
				^B 2.95	15.57	13.69	^a Minipress PF
1480X NP	Tablet 2 mg (base)	100	5	..	14.93	16.00	^a Chem mart CH
							^a Prazosin GX
							^a GenRx Prazosin TW
							^a Terry White Chemists Prazosin
				^B 3.04	17.97	16.00	^a Minipress PF

Arteriolar smooth muscle, agents acting on *Hydrazinophthalazine derivatives*

HYDRALAZINE HYDROCHLORIDE							
1639G NP	Tablet 50 mg	200	2	..	*17.42	18.49	Alphapress 50 AF
1640H NP	Tablet 25 mg	200	2	..	*15.50	16.57	Alphapress 25 AF

Pyrimidine derivatives

MINOXIDIL

Authority required (STREAMLINED)

2759

Severe refractory hypertension. Treatment must be initiated by a consultant physician.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2313R NP	Tablet 10 mg	100	5	..	52.83	34.20	Loniten PF

Diuretics

Low-ceiling diuretics, thiazides *Thiazides, plain*

HYDROCHLOROTHIAZIDE							
1484D NP	Tablet 25 mg	100	1	..	21.24	22.31	Dithiazide PL

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Low-ceiling diuretics, excl. thiazides							
<i>Sulfonamides, plain</i>							
CHLORTHALIDONE							
1585K NP	Tablet 25 mg	100	1	..	*13.64	14.71	Hygroton 25 LM
INDAPAMIDE HEMIHYDRATE							
2436F NP	Tablet 2.5 mg	90	1	..	17.48	18.55	^a Chem mart CH
							^a Indapamide AF
							^a Dapa-Tabs GX
							^a GenRx Indapamide GM
							^a Indapamide-GA SZ
							^a Indapamide Sandoz SI
							^a Insig TW
							^a Terry White Chemists Indapamide
				^B 2.56	20.04	18.55	^a Natrilix SE
8532C NP	Tablet 1.5 mg (sustained release)	90	1	..	19.31	20.38	Natrilix SR SE
High-ceiling diuretics							
<i>Sulfonamides, plain</i>							
FRUSEMIDE							
2411X NP	Oral solution 10 mg per mL, 30 mL	‡1	3	..	17.06	18.13	Lasix SW
2412Y NP	Tablet 40 mg	100	1	..	8.79	9.86	^a Chem mart CH
							^a Frusemide SZ
							^a Frusemide Sandoz GM
							^a Frusid GX
							^a GenRx Frusemide TW
							^a Terry White Chemists Frusemide
							^a Uremide AF
							Urex FM
				^B 2.40	11.19	9.86	^a Lasix SW
2413B NP	Injection 20 mg in 2 mL	5	..	..	10.48	11.55	^a Frusemide-Clarix AE
							^a Frusemide Sandoz SZ
							^a Lasix SW
2414C NP	Tablet 20 mg	100	1	..	*9.02	10.09	Urex-M FM
				^B 1.90	*10.92	10.09	^a Lasix-M SW
					9.03	10.10	^a Chem mart CH
							^a Frusemide GM
							^a Frusid GX
							^a GenRx Frusemide TW
							^a Terry White Chemists Frusemide
2415D NP	Tablet 500 mg	50	3	..	18.74	19.81	Urex-Forte FM

Aryloxyacetic acid derivatives

ETHACRYNIC ACID

Restricted benefit

Patients hypersensitive to other oral diuretics.

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8748K NP	Tablet 25 mg	200	1	..	*197.30	34.20	Edecrin	FK

Potassium-sparing agents *Aldosterone antagonists*

EPLERENONE

Caution

Serum electrolytes should be checked regularly.

Authority required (STREAMLINED)

2637

Heart failure with a left ventricular ejection fraction of 40% or less occurring within 3 to 14 days following an acute myocardial infarction. Treatment with eplerenone must be commenced within 14 days of an acute myocardial infarction.

The date of the acute myocardial infarction and the date of initiation of eplerenone treatment must be documented in the patient's medical records when PBS-subsidised treatment is initiated.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8879H NP	Tablet 25 mg	30	5	..	112.77	34.20	Inspira	PF
8880J NP	Tablet 50 mg	30	5	..	112.77	34.20	Inspira	PF

SPIRONOLACTONE

Caution

Appropriate contraceptive measures should be taken by women of child-bearing age in whom spironolactone therapy has been initiated.

Caution

Serum electrolytes should be checked regularly.

2339D NP	Tablet 25 mg	100	5	..	12.49	13.56 ^a	Spiractin 25	AF
				^B 1.84	14.33	13.56 ^a	Aldactone	PF
2340E NP	Tablet 100 mg	100	5	..	30.32	31.39 ^a	Spiractin 100	AF
				^B 2.53	32.85	31.39 ^a	Aldactone	PF

Other potassium-sparing agents

AMILORIDE HYDROCHLORIDE

Caution

Serum electrolytes should be checked regularly.

3109P NP	Tablet 5 mg	100	1	..	*10.98	12.05	Kaluril	AF

Diuretics and potassium-sparing agents in combination *Low-ceiling diuretics and potassium-sparing agents*

HYDROCHLOROTHIAZIDE with AMILORIDE HYDROCHLORIDE

Caution

Serum electrolytes should be checked regularly.

1486F NP	Tablet 50 mg-5 mg	100	1	..	*13.88	14.95	Moduretic	AS

HYDROCHLOROTHIAZIDE with TRIAMTERENE

Caution

Serum electrolytes should be checked regularly.

1280J NP	Tablet 25 mg-50 mg	100	1	..	12.89	13.96	Hydrene 25/50	AF

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Peripheral vasodilators							

Peripheral vasodilators

Other peripheral vasodilators

PHENOXYBENZAMINE HYDROCHLORIDE

Restricted benefit

Phaeochromocytoma;

Neurogenic urinary retention.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1166J NP	Capsules 10 mg, 30	3	5	..	*204.90	34.20	Dibenyline	GH
1862B NP	Capsule 10 mg	100	5	..	67.36	34.20	Dibenyline	GH
9286R NP	Capsules 10 mg, 100	±1	5	..	1164.47	34.20	Dibenzyliline	GH

Beta blocking agents

Beta blocking agents

Beta blocking agents, non-selective

OXPRENOLOL HYDROCHLORIDE

2942W NP	Tablet 20 mg	100	5	..	10.00	11.07	Corbeton 20	AF
2961W NP	Tablet 40 mg	100	5	..	11.78	12.85	Corbeton 40	AF

PINDOLOL

3062E NP	Tablet 5 mg	100	5	..	11.48	12.55	^a Barbloc 5	AF
				^B 2.71	14.19	12.55	^a Visken 5	NV
3065H NP	Tablet 15 mg	50	5	..	13.71	14.78	^a Barbloc 15	AF
				^B 2.69	16.40	14.78	^a Visken 15	NV

PROPRANOLOL HYDROCHLORIDE

2565B NP	Tablet 10 mg	100	5	..	10.19	11.26	Deralin 10	AF
				^B 3.14	13.33	11.26	Inderal	AP
2566C NP	Tablet 40 mg	100	5	..	10.56	11.63	Deralin 40	AF
				^B 3.14	13.70	11.63	Inderal	AP
2899N NP	Tablet 160 mg	50	5	..	11.01	12.08	Deralin 160	AF

SOTALOL HYDROCHLORIDE

Restricted benefit

Severe cardiac arrhythmias.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2043M NP	Tablet 160 mg	60	5	..	26.50	27.57	^a Cardol	AF
							^a Chem mart Sotalol	CH
							^a GenRx Sotalol	GX
							^a Solavert	SI

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
8398B NP	Tablet 80 mg	60	5	..	15.78	16.85	^a Sotalol Sandoz SZ
							^a Terry White Chemists Sotalol TW
							^a Sotacor FM
							^a GenRx Sotalol GX
				^B 5.00	20.78	16.85	^a Solavert SI
							^a Sotalol Sandoz SZ
							^a Sotacor FM

Beta blocking agents, selective

1081X NP	ATENOLOL Tablet 50 mg	30	5	..	10.27	11.34	^a APO-Atenolol TX
							^a Atenolol-GA GN
							^a Atenolol Sandoz SZ
							^a Chem mart Atenolol CH
				^B 3.54	13.81	11.34	^a Noten AF
							^a Tensig SI
							^a Terry White Chemists Atenolol TW
							^a Tenormin AP

BISOPROLOL FUMARATE

Authority required (STREAMLINED)

3234

Moderate to severe heart failure in a patient stabilised on conventional therapy which must include an ACE inhibitor or Angiotensin II antagonist, if tolerated.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8604W NP	Tablet 2.5 mg	28	5	..	49.16	34.20	^a Bicar AL
							^a Bisoprolol Sandoz SZ
							^a Bispro 2.5 AF
8605X NP	Tablet 5 mg	28	5	..	59.00	34.20	^a Bicar AL
							^a Bisoprolol Sandoz SZ
							^a Bispro 5 AF
8606Y NP	Tablet 10 mg	28	5	..	72.13	34.20	^a Bicar AL
							^a Bisoprolol Sandoz SZ
							^a Bispro 10 AF

METOPROLOL SUCCINATE

Authority required (STREAMLINED)

3234

Moderate to severe heart failure in a patient stabilised on conventional therapy which must include an ACE inhibitor or Angiotensin II antagonist, if tolerated.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8732N NP	Tablet 23.75 mg (controlled release)	15	..	..	21.04	22.11	Toprol-XL 23.75	AP
8733P	Tablet 47.5 mg (controlled release)	30	5	..	72.46	34.20	Toprol-XL 47.5	AP

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
NP							
8734Q	Tablet 95 mg (controlled release)	30	5	..	88.96	34.20	Toprol-XL 95 AP
NP							
8735R	Tablet 190 mg (controlled release)	30	5	..	109.60	34.20	Toprol-XL 190 AP
NP							
METOPROLOL TARTRATE							
1324Q	Tablet 50 mg	100	5	..	10.84	11.91	^a Chem mart CH
NP							Metoprolol
							^a GenRx Metoprolol GX
							^a Metohexal SZ
							^a Metrol 50 SI
							^a Minax 50 AF
							^a Terry White Chemists TW
							Metoprolol
				^B 2.35	13.19	11.91	Lopresor 50 NV
				^B 3.25	14.09	11.91	^a Betaloc AP
1325R	Tablet 100 mg	60	5	..	12.03	13.10	^a Chem mart CH
NP							Metoprolol
							^a GenRx Metoprolol GX
							^a Metohexal SZ
							^a Metrol 100 SI
							^a Minax 100 AF
							^a Terry White Chemists TW
							Metoprolol
				^B 2.35	14.38	13.10	Lopresor 100 NV
				^B 3.25	15.28	13.10	^a Betaloc AP

NEBIVOLOL

Authority required (STREAMLINED)

3234

Moderate to severe heart failure in a patient stabilised on conventional therapy which must include an ACE inhibitor or Angiotensin II antagonist, if tolerated.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9310B	Tablet 1.25 mg (as hydrochloride), 28	1	5	..	29.25	30.32	Nebilet	CS
NP								
9311C	Tablet 5 mg (as hydrochloride)	28	5	..	60.94	34.20	Nebilet	CS
NP								
9312D	Tablet 10 mg (as hydrochloride)	28	5	..	68.02	34.20	Nebilet	CS
NP								
9316H	Tablet 1.25 mg (as hydrochloride)	56	5	..	*50.62	34.20	Nebilet	CS
NP								

Alpha and beta blocking agents

CARVEDILOL

Authority required (STREAMLINED)

3234

Moderate to severe heart failure in a patient stabilised on conventional therapy which must include an ACE inhibitor or Angiotensin II antagonist, if tolerated;

1735

Patients receiving this drug as a pharmaceutical benefit prior to 1 August 2002.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
8255L <i>NP</i>	Tablet 3.125 mg	30	..	..	15.53	16.60	a	APO-Carvedilol TX
							a	Chem mart CH
							a	Carvedilol 3.125 mg Dilasig 3.125 FM
							a	Dilatrend 3.125 RO
							a	GenRx Carvedilol GX
							a	GN-Carvedilol GM
							a	Kredex MD
							a	Terry White Chemists Carvedilol 3.125 mg Veditol 3.125 SI
8256M <i>NP</i>	Tablet 6.25 mg	60	5	..	48.40	34.20	a	APO-Carvedilol TX
							a	Carvedilol generichealth GQ
							a	Carvedilol Sandoz SZ
							a	Chem mart Carvedilol 6.25 mg Dicarz AF
							a	Dilasig 6.25 FM
							a	Dilatrend 6.25 RO
							a	GenRx Carvedilol GX
							a	GN-Carvedilol GM
							a	Kredex MD
							a	Terry White Chemists Carvedilol 6.25 mg Veditol 6.25 SI
8257N <i>NP</i>	Tablet 12.5 mg	60	5	..	57.97	34.20	a	APO-Carvedilol TX
							a	Carvedilol generichealth GQ
							a	Carvedilol Sandoz SZ
							a	Chem mart Carvedilol 12.5 mg Dicarz AF
							a	Dilasig 12.5 FM
							a	Dilatrend 12.5 RO
							a	GenRx Carvedilol GX
							a	GN-Carvedilol GM
							a	Kredex MD
							a	Terry White Chemists Carvedilol 12.5 mg Veditol 12.5 SI
8258P <i>NP</i>	Tablet 25 mg	60	5	..	70.85	34.20	a	APO-Carvedilol TX
							a	Carvedilol generichealth GQ
							a	Carvedilol Sandoz SZ
							a	Chem mart Carvedilol 25 mg Dicarz AF

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
							^a Dilasig 25 FM
							^a Dilatrend 25 RO
							^a GenRx Carvedilol GX
							^a GN-Carvedilol GM
							^a Kredex MD
							^a Terry White Chemists TW
							^a Carvedilol 25 mg Vedilol 25 SI
LABETALOL HYDROCHLORIDE							
1566K NP	Tablet 100 mg	100	5	..	15.46	16.53	^a Presolol 100 AF
				^B 3.20	18.66	16.53	^a Trandate SI
1567L NP	Tablet 200 mg	100	5	..	21.30	22.37	^a Presolol 200 AF
				^B 3.20	24.50	22.37	^a Trandate SI

Calcium channel blockers

Selective calcium channel blockers with mainly vascular effects

Dihydropyridine derivatives

Note

The base-priced drugs in this therapeutic group are amlodipine, felodipine, lercanidipine hydrochloride and nifedipine (except nifedipine controlled release tablet 20 mg).

AMLODIPINE

Note

Bioequivalence has been demonstrated between the tablet containing 5 mg amlodipine (as besylate) and the tablet containing 5 mg amlodipine (as maleate).

1343Q NP	Tablet 5 mg (as maleate)	30	5	..	16.68	17.75	^a Amlo 5 ZP
2751T NP	Tablet 5 mg (as besylate)	30	5	..	16.68	17.75	^a Amlodipine-DRLA RZ
							^a Amlodipine-GA GM
							^a Amlodipine generichealth GQ
							^a Amlodipine Sandoz SZ
							^a APO-Amlodipine TX
							^a Chem mart Amlodipine CH
							^a Nordip AF
							^a Norvapine GN
							^a Ozlodip RA
							^a Perivasc AL
							^a Pharmacor Amlodipine 5 CR
							^a Terry White Chemists TW
				^B 3.91	20.59	17.75	^a Amlodipine Norvasc PF

AMLODIPINE

Note

Bioequivalence has been demonstrated between the tablet containing 10 mg amlodipine (as besylate) and the tablet containing 10 mg amlodipine (as maleate).

1345T NP	Tablet 10 mg (as maleate)	30	5	..	24.39	25.46	^a Amlo 10 ZP
2752W	Tablet 10 mg (as besylate)	30	5	..	24.39	25.46	^a Amlodipine-DRLA RZ

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
<i>NP</i>							
							^a Amlodipine-GA GM
							^a Amlodipine generichealth GQ
							^a Amlodipine Sandoz SZ
							^a APO-Amlodipine TX
							^a Chem mart Amlodipine CH
							^a Nordip AF
							^a Norvapine GN
							^a Ozlodip RA
							^a Perivasc AL
							^a Pharmacor Amlodipine 10 CR
							^a Terry White Chemists Amlodipine TW
				^B 5.69	30.08	25.46	^a Norvasc PF
	FELODIPINE						
2361G <i>NP</i>	Tablet 2.5 mg (extended release)	30	5	..	13.11	14.18	^a Felodur ER 2.5 mg AL
				^B 4.99	18.10	14.18	^a Plendil ER AP
2366M <i>NP</i>	Tablet 5 mg (extended release)	30	5	..	15.98	17.05	^a Felodil XR 5 SI
							^a Felodur ER 5 mg AL
				^B 5.00	20.98	17.05	^a Plendil ER AP
2367N <i>NP</i>	Tablet 10 mg (extended release)	30	5	..	23.57	24.64	^a Felodil XR 10 SI
							^a Felodur ER 10 mg AL
				^B 5.00	28.57	24.64	^a Plendil ER AP
	LERCANIDIPINE HYDROCHLORIDE						
8534E <i>NP</i>	Tablet 10 mg	28	5	..	16.18	17.25	^a APO-Lercanidipine TX
							^a Chem mart Lercanidipine CH
							^a Lercan SI
							^a Lercanidipine Sandoz SZ
							^a Terry White Chemists Lercanidipine TW
				^B 1.95	18.13	17.25	^a Zanidip SM
8679T <i>NP</i>	Tablet 20 mg	28	5	..	22.74	23.81	^a APO-Lercanidipine TX
							^a Chem mart Lercanidipine CH
							^a Lercan SI
							^a Lercanidipine Sandoz SZ
							^a Terry White Chemists Lercanidipine TW
				^B 3.44	26.18	23.81	^a Zanidip SM
	NIFEDIPINE						
1694E <i>NP</i>	Tablet 10 mg	60	5	..	15.86	16.93	^a Adefin 10 AF
				^B 1.17	17.03	16.93	^a Adalat 10 BN
1695F	Tablet 20 mg	60	5	..	18.17	19.24	^a Adefin 20 AF

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
NP								
							^a GenRx Nifedipine	GX
							^a Nifehexal	SZ
				^B 2.21	20.38	19.24	^a Adalat 20	BN
1906H	Tablet 30 mg (controlled release)	30	5	..	19.07	20.14	^a Addos XR 30	SI
NP							^a Adefin XL 30	AF
							^a APO-Nifedipine XR	TX
				^B 2.54	21.61	20.14	^a Adalat Oros 30	BN
1907J	Tablet 60 mg (controlled release)	30	5	..	22.31	23.38	^a Addos XR 60	SI
NP							^a Adefin XL 60	AF
							^a APO-Nifedipine XR	TX
				^B 2.81	25.12	23.38	^a Adalat Oros 60	BN
8610E	Tablet 20 mg (controlled release)	30	5	^T 2.26	20.40	19.21	Adalat Oros 20mg	BN
NP								

NIFEDIPINE

Authority required

Adverse effects occurring with all of the base-priced drugs;

Drug interactions occurring with all of the base-priced drugs;

Drug interactions expected to occur with all of the base-priced drugs;

Transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.

8938K	Tablet 20 mg (controlled release)	30	5	..	20.40	21.47	Adalat Oros 20mg	BN
NP								

Selective calcium channel blockers with direct cardiac effects

Phenylalkylamine derivatives

VERAPAMIL HYDROCHLORIDE

Caution

The myocardial depressant effects of this drug and of beta-blocking drugs are additive.

1060T	Injection 5 mg in 2 mL	5	..	..	12.38	13.45	Isoptin	AB
NP								
1241H	Tablet 240 mg (sustained release)	30	5	..	16.15	17.22	^a Cordilox SR	KN
NP								
				^B 2.26	18.41	17.22	^a Isoptin SR	AB
1248Q	Tablet 40 mg	100	5	..	11.51	12.58	^a Anpec 40	AF
NP								
				^B 0.76	12.27	12.58	^a Isoptin	AB
1250T	Tablet 80 mg	100	5	..	15.46	16.53	^a Anpec 80	AF
NP								
				^B 0.76	16.22	16.53	^a Isoptin	AB
1253Y	Tablet 160 mg	60	5	..	18.44	19.51	Isoptin	AB
NP								
1254B	Tablet 120 mg	100	5	..	19.52	20.59	Isoptin	AB
NP								
2206D	Capsule 160 mg (sustained release)	30	5	..	12.45	13.52	Veracaps SR	SI
NP								
2207E	Capsule 240 mg (sustained release)	30	5	..	16.24	17.31	Veracaps SR	SI
NP								
2208F	Tablet 180 mg (sustained release)	30	5	..	13.72	14.79	^a Cordilox 180 SR	KN
NP								
				^B 2.27	15.99	14.79	^a Isoptin 180 SR	AB

Benzothiazepine derivatives

DILTIAZEM HYDROCHLORIDE

Caution

The myocardial depressant effects of this drug and of beta-blocking drugs are additive.

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
1312C NP	Capsule 180 mg (controlled delivery)	30	5	..	17.26	18.33	^a	Cardizem CD SW
							^a	Chem mart CH
							^a	Diltiazem CD
							^a	Diltahexal CD SZ
							^a	Dilzem CD GM
							^a	GenRx Diltiazem CD GX
							^a	Terry White Chemists TW
							^a	Diltiazem CD
							^a	Vasocardol CD AV
1313D NP	Capsule 240 mg (controlled delivery)	30	5	..	21.07	22.14	^a	Cardizem CD SW
							^a	Chem mart CH
							^a	Diltiazem CD
							^a	Diltahexal CD SZ
							^a	Dilzem CD GM
							^a	GenRx Diltiazem CD GX
							^a	Terry White Chemists TW
							^a	Diltiazem CD
							^a	Vasocardol CD AV
1335G NP	Tablet 60 mg	90	5	..	16.20	17.27	^a	Cardizem SW
							^a	Chem mart CH
							^a	Diltiazem
							^a	Coras AF
							^a	Diltiazem Sandoz SZ
							^a	Dilzem 60 mg GM
							^a	GenRx Diltiazem GX
							^a	Terry White Chemists TW
							^a	Diltiazem
							^a	Vasocardol AV
8480H NP	Capsule 360 mg (controlled delivery)	30	5	..	25.64	26.71	^a	Cardizem CD SW
							^a	Diltahexal CD SZ
							^a	Vasocardol CD AV

Agents acting on the renin-angiotensin system

ACE inhibitors, plain

ACE inhibitors, plain

Caution

Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

CAPTOPRIL

1147J NP	Tablet 12.5 mg	90	5	..	16.68	17.75	^a	Captopril Sandoz SZ
							^a	GenRx Captopril GX
							^a	Zedace AF
1148K NP	Tablet 25 mg	90	5	..	21.27	22.34	^a	Ascent Pharma Pty Ltd GM
							^a	Captopril Sandoz SZ
							^a	GenRx Captopril GX
							^a	Zedace AF
				^B 5.00	26.27	22.34	^a	Capoten SI
1149L NP	Tablet 50 mg	90	5	..	35.70	34.20	^a	Ascent Pharma Pty Ltd GM
							^a	Captopril Sandoz SZ

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
							^a GenRx Captopril GX
							^a Zedace AF
				^B 5.00	40.70	34.20	^a Capoten SI
CAPTOPRIL							
<u>Restricted benefit</u>							
For patients unable to take a solid dose form of an ACE inhibitor.							
8760C NP	Oral solution 5 mg per mL, 95 mL	#1	5	..	111.82	34.20	Capoten SI
ENALAPRIL							
1368B NP	Tablet containing enalapril maleate 10 mg	30	5	..	17.45	18.52	^a Alphapril AF
							^a Auspril SI
							^a Chem mart Enalapril CH
							^a Enahexal HX
							^a Enalabell BF
							^a Enalapril-DP 10mg GN
							^a Enalapril-GA GM
							^a Enalapril generichealth GQ
							^a Enalapril Sandoz SZ
							^a Enalapril Winthrop WA
							^a GenRx Enalapril GX
							^a Terry White Chemists Enalapril TW
				^B 4.90	22.35	18.52	^a Renitec MK
1369C NP	Tablet containing enalapril maleate 20 mg	30	5	..	20.46	21.53	^a Alphapril AF
							^a Auspril SI
							^a Chem mart Enalapril CH
							^a Enahexal HX
							^a Enalabell BF
							^a Enalapril-DP 20mg GN
							^a Enalapril-GA GM
							^a Enalapril generichealth GQ
							^a Enalapril Sandoz SZ
							^a GenRx Enalapril GX
							^a Terry White Chemists Enalapril TW
				^B 4.90	25.36	21.53	^a Renitec 20 MK
1370D NP	Tablet containing enalapril maleate 5 mg	30	5	..	13.14	14.21	^a Alphapril AF
							^a Auspril SI
							^a Chem mart Enalapril CH
							^a Enahexal HX
							^a Enalabell BF
							^a Enalapril-DP 5mg GN
							^a Enalapril-GA GM
							^a Enalapril generichealth GQ

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
							^a Enalapril Sandoz	SZ
							^a Enalapril Winthrop	WA
							^a GenRx Enalapril	GX
							^a Terry White Chemists Enalapril	TW
							^B 4.90	18.04
FOSINOPRIL SODIUM								
1182F NP	Tablet 10 mg	30	5	..	15.65	16.72	^a Fosinopril Sandoz	SZ
							^a Fosipril 10	SI
							^a GenRx Fosinopril	GX
							^a Monace 10	AF
							^a Monopril	BQ
1183G NP	Tablet 20 mg	30	5	..	20.25	21.32	^a Fosinopril Sandoz	SZ
							^a Fosipril 20	SI
							^a GenRx Fosinopril	GX
							^a Monace 20	AF
							^a Monopril	BQ
LISINOPRIL								
2456G NP	Tablet 5 mg	30	5	..	14.13	15.20	^a APO-Lisinopril	TX
							^a Chem mart Lisinopril	CH
							^a Fibsol 5	SI
							^a GenRx Lisinopril	GX
							^a Liprace	GM
							^a Lisinopril 5	CR
							^a Lisinopril-DRLA	RZ
							^a Lisinopril-GA	GN
							^a Lisinopril generichealth	GQ
							^a Lisinopril Ranbaxy	RA
							^a Lisinopril Sandoz	SZ
							^a Lisinopril Winthrop	WA
							^a Lisodur	AF
							^a Terry White Chemists Lisinopril	TW
							^B 2.08	16.21
^B 3.95	18.08	15.20	^a Prinivil 5	MK				
2457H NP	Tablet 10 mg	30	5	..	17.99	19.06	^a APO-Lisinopril	TX
							^a Chem mart Lisinopril	CH
							^a Fibsol 10	SI
							^a GenRx Lisinopril	GX
							^a Liprace	GM
							^a Lisinopril 10	CR
							^a Lisinopril-DRLA	RZ
							^a Lisinopril-GA	GN
							^a Lisinopril generichealth	GQ
							^a Lisinopril Hexal	HX

PERINDOPRIL

Note

3050M NP	Tablet containing 2 mg perindopril erbumine	30	5	..	12.58	13.65	^a	APO-Perindopril	TX
							^a	Chem mart Perindopril	CH
							^a	GenRx Perindopril	GX
							^a	Indopril 2	SI
							^a	Ozapace	RA
							^a	Perindo	AF
							^a	Perindopril 2	CR
							^a	Perindopril-DP	GN
							^a	Perindopril-GA	GM
							^a	Terry White Chemists Perindopril	TW
9006B NP	Tablet containing 2.5 mg perindopril arginine	30	5	..	12.58	13.65	^a	Coversyl 2.5mg	SE

PERINDOPRIL

Note

3051N NP	Tablet containing 4 mg perindopril erbumine	30	5	..	17.94	19.01	^a	APO-Perindopril	TX
							^a	Chem mart Perindopril	CH

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
9007C NP	Tablet containing 5 mg perindopril arginine	30	5	..	17.94	19.01	^a GenRx Perindopril	GX
							^a Indopril 4	SI
							^a Ozapace	RA
							^a Perindo	AF
							^a Perindopril 4	CR
							^a Perindopril-DP	GN
							^a Perindopril-GA	GM
							^a Terry White Chemists Perindopril	TW
^a Coversyl 5mg	SE							
PERINDOPRIL								
Note								
Bioequivalence has been demonstrated between perindopril erbumine 8 mg and perindopril arginine 10 mg.								
8704D NP	Tablet containing 8 mg perindopril erbumine	30	5	..	24.07	25.14	^a APO-Perindopril	TX
							^a Chem mart Perindopril	CH
							^a GenRx Perindopril	GX
							^a Indopril 8	SI
							^a Ozapace	RA
							^a Perindo	AF
							^a Perindopril 8	CR
							^a Perindopril-DP	GN
9008D NP	Tablet containing 10 mg perindopril arginine	30	5	..	24.07	25.14	^a Perindopril-GA	GM
							^a Terry White Chemists Perindopril	TW
							^a Coversyl 10mg	SE
QUINAPRIL								
1968N NP	Tablet 5 mg (as hydrochloride)	30	5	..	13.96	15.03	^a Acquin 5	SI
							^a APO-Quinapril	TX
							^a Filpril	AL
							^a Pharmacor Quinapril 5	CR
							^a Qpril 5	AF
							^a Quinapril-DP	GN
							^a Quinapril generichealth	GQ
							^a Quinapril Sandoz	SZ
1969P NP	Tablet 10 mg (as hydrochloride)	30	5	..	17.16	18.23	^a Accupril	PF
							^a Acquin 10	SI
							^a APO-Quinapril	TX
							^a Filpril	AL
							^a Pharmacor Quinapril 10	CR
							^a Qpril 10	AF
							^a Quinapril-DP	GN
							^a Quinapril generichealth	GQ
				^B 0.66	17.82	18.23	^a Accupril	PF

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
1970Q NP	Tablet 20 mg (as hydrochloride)	30	5	..	19.73	20.80	^a	Acquin 20 SI
							^a	APO-Quinapril TX
							^a	Filpril AL
							^a	Pharmacor CR
							^a	Quinapril 20 Qpril 20 AF
							^a	Quinapril-GA GM
							^a	Quinapril generichealth GQ
							^a	Quinapril Sandoz SZ
				^B 1.00	20.73	20.80	^a	Accupril PF
RAMIPRIL								
Note								
Ramipril 1.25 mg tablets and capsules are bioequivalent.								
1944H NP	Tablet 1.25 mg	30	5	..	11.60	12.67	^a	APO-Ramipril TX
							^a	Chem mart CH
							^a	Ramipril SI
							^a	Prilace 1.25 SI
							^a	Ramace 1.25 mg AV
							^a	Ramipril Sandoz SZ
							^a	Ramipril Winthrop WA
							^a	Terry White Chemists TW
							^a	Ramipril SW
							^a	Tritace 1.25 mg SW
							^a	Tryzan Tabs 1.25 AF
9120B NP	Capsule 1.25 mg	30	5	..	11.60	12.67	^a	Pharmacor CR
							^a	Ramipril 1.25 GN
							^a	Ramipril-DP GN
							^a	Ramipril-GA GM
							^a	Ramipril generichealth GQ
							^a	Tryzan Caps 1.25 AF
RAMIPRIL								
Note								
Ramipril 2.5 mg tablets and capsules are bioequivalent.								
1945J NP	Tablet 2.5 mg	30	5	..	14.00	15.07	^a	APO-Ramipril TX
							^a	Chem mart CH
							^a	Ramipril SI
							^a	Prilace 2.5 SI
							^a	Ramace 2.5 mg AV
							^a	Ramipril Sandoz SZ
							^a	Ramipril Winthrop WA
							^a	Terry White Chemists TW
							^a	Ramipril SW
							^a	Tritace 2.5 mg SW
							^a	Tryzan Tabs 2.5 AF
9121C NP	Capsule 2.5 mg	30	5	..	14.00	15.07	^a	Pharmacor CR
							^a	Ramipril 2.5 GN
							^a	Ramipril-DP GN
							^a	Ramipril-GA GM

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
							^a	
								RAMIPRIL
							^a	generichealth GQ
							^a	Tryzan Caps 2.5 AF
RAMIPRIL								
Note								
Ramipril 5 mg tablets and capsules are bioequivalent.								
1946K NP	Tablet 5 mg	30	5	..	15.93	17.00	^a	APO-Ramipril TX
							^a	Chem mart CH
							^a	Ramipril
							^a	Prilace 5 SI
							^a	Ramace 5 mg AV
							^a	Ramipril Sandoz SZ
							^a	Ramipril Winthrop WA
							^a	Terry White TW
							^a	Chemists
							^a	Ramipril
							^a	Tritace 5 mg SW
							^a	Tryzan Tabs 5 AF
9122D NP	Capsule 5 mg	30	5	..	15.93	17.00	^a	Pharmacor CR
							^a	Ramipril 5
							^a	Ramipril-DP GN
							^a	Ramipril GQ
							^a	generichealth
							^a	Tryzan Caps 5 AF
RAMIPRIL								
Note								
Ramipril 10 mg tablets and capsules are bioequivalent.								
1316G NP	Tablet 10 mg	30	5	..	23.23	24.30	^a	APO-Ramipril TX
							^a	Chem mart CH
							^a	Ramipril
							^a	Ramipril Sandoz SZ
							^a	Terry White TW
							^a	Chemists
							^a	Ramipril
							^a	Tritace SW
							^a	Tryzan Tabs 10 AF
8470T NP	Capsule 10 mg	30	5	..	23.23	24.30	^a	GenRx Ramipril GX
							^a	Pharmacor CR
							^a	Ramipril 10
							^a	Prilace 10 SI
							^a	Ramace 10 mg AV
							^a	Ramipril-DP GN
							^a	Ramipril-GA GM
							^a	Ramipril GQ
							^a	generichealth
							^a	Ramipril Sandoz SZ
							^a	Ramipril Winthrop WA
							^a	Tritace 10 mg SW
							^a	Tryzan Caps 10 AF

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8668F NP	RAMIPRIL Pack containing 7 tablets 2.5 mg, 21 tablets 5 mg and 10 capsules 10 mg	±1	..	..	22.03	23.10	Tritace Titration Pack	SW
2791X NP	TRANDOLAPRIL Capsule 500 micrograms	28	5	..	9.28	10.35	^a APO-Trandolapril	TX
							^a Dolapril 0.5	SI
							^a Tranalpha	AF
							^a Trandolapril-DP	GN
							^a Trandolapril generichealth	GQ
				^B 1.87	11.15	10.35	^a Gopten	AB
2792Y NP	Capsule 1 mg	28	5	..	14.00	15.07	^a APO-Trandolapril	TX
							^a Dolapril 1	SI
							^a Tranalpha	AF
							^a Trandolapril-DP	GN
							^a Trandolapril generichealth	GQ
				^B 1.87	15.87	15.07	^a Gopten	AB
2793B NP	Capsule 2 mg	28	5	..	15.57	16.64	^a APO-Trandolapril	TX
							^a Dolapril 2	SI
							^a Tranalpha	AF
							^a Trandolapril-DP	GN
							^a Trandolapril generichealth	GQ
				^B 1.87	17.44	16.64	^a Gopten	AB
8758Y NP	Capsule 4 mg	28	5	..	23.60	24.67	^a APO-Trandolapril	TX
							^a Dolapril 4	SI
							^a Tranalpha	AF
							^a Trandolapril-DP	GN
							^a Trandolapril generichealth	GQ
				^B 1.88	25.48	24.67	^a Gopten	AB

ACE inhibitors, combinations

ACE inhibitors and diuretics

Caution

Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

ENALAPRIL MALEATE with HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

8477E NP	Tablet 20 mg-6 mg	30	5	..	28.04	29.11	^a Enalapril/HCT Sandoz	SZ
							^a Renitec Plus 20/6	MK

FOSINOPRIL SODIUM with HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

8400D NP	Tablet 10 mg-12.5 mg	30	5	..	21.62	22.69	^a APO-Fosinopril HCTZ 10/12.5	TX
							^a Fosinopril/HCT Sandoz 10mg/12.5mg	SZ
							^a Fosinopril/HCTZ-GA 10/12.5	GM

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8401E NP	Tablet 20 mg-12.5 mg	30	5	..	28.75	29.82	^a Hyforil	RA
							^a Monoplus 10/12.5	BQ
							^a APO-Fosinopril HCTZ 20/12.5	TX
							^a Fosetic 20/12.5	ZP
							^a Fosinopril/HCT Sandoz 20mg/12.5mg	SZ
							^a Fosinopril/HCTZ-GA 20/12.5	GM
							^a Hyforil	RA
							^a Monoplus 20/12.5	BQ
PERINDOPRIL with INDAPAMIDE HEMIHYDRATE								
2190G NP	Tablet containing 2.5 mg perindopril arginine- 0.625 mg indapamide hemihydrate	30	5	..	16.32	17.39	Coversyl Plus LD 2.5mg/0.625mg	SE
PERINDOPRIL with INDAPAMIDE HEMIHYDRATE								
<u>Restricted benefit</u>								
Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.								
<u>Note</u>								
Bioequivalence has been demonstrated between perindopril erbumine/indapamide hemihydrate tablet 4 mg-1.25 mg and perindopril arginine/indapamide hemihydrate tablet 5 mg-1.25 mg.								
2845R NP	Tablet containing 5 mg perindopril arginine- 1.25 mg indapamide hemihydrate	30	5	..	28.67	29.74	^a Coversyl Plus 5mg/1.25mg	SE
8449Q NP	Tablet containing 4 mg perindopril erbumine- 1.25 mg indapamide hemihydrate	30	5	..	28.67	29.74	^a Chem mart Perindopril/ Indapamide 4/1.25	CH
							^a GenRx Perindopril/ Indapamide 4/1.25	GX
							^a Perindo Combi 4/1.25	AF
							^a Terry White Chemists Perindopril/ Indapamide 4/1.25	TW
QUINAPRIL HYDROCHLORIDE with HYDROCHLOROTHIAZIDE								
<u>Restricted benefit</u>								
Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.								
8589C NP	Tablet 10 mg (base)-12.5 mg	30	5	..	19.39	20.46	Accuretic 10/12.5mg	PF
8590D NP	Tablet 20 mg (base)-12.5 mg	30	5	..	21.96	23.03	Accuretic 20/12.5mg	PF

ACE inhibitors and calcium channel blockers

Caution

Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

LERCANIDIPINE HYDROCHLORIDE with ENALAPRIL MALEATE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

9144G NP	Tablet 10 mg-10 mg	28	5	..	26.49	27.56	Zan-Extra 10/10	SM
9145H NP	Tablet 10 mg-20 mg	28	5	..	29.29	30.36	Zan-Extra 10/20	SM

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
PERINDOPRIL with AMLODIPINE								
Note								
Treatment should not be initiated with this combination.								
Restricted benefit								
Hypertension in a patient who is not adequately controlled with either of the drugs in the combination;								
Stable coronary heart disease in a patient who is stabilised on treatment with perindopril and amlodipine at the same doses.								
9346X NP	Tablet containing 5 mg perindopril arginine with 5 mg amlodipine (as besylate)	30	5	..	28.20	29.27	Coveram	SE
9347Y NP	Tablet containing 5 mg perindopril arginine with 10 mg amlodipine (as besylate)	30	5	..	35.92	34.20	Coveram	SE
9348B NP	Tablet containing 10 mg perindopril arginine with 5 mg amlodipine (as besylate)	30	5	..	34.33	34.20	Coveram	SE
9349C NP	Tablet containing 10 mg perindopril arginine with 10 mg amlodipine (as besylate)	30	5	..	41.90	34.20	Coveram	SE
RAMIPRIL with FELODIPINE								
Restricted benefit								
Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.								
2626F NP	Tablet 2.5 mg-2.5 mg (modified release)	30	5	..	20.69	21.76	Triasyn 2.5/2.5	SW
2629J NP	Tablet 5 mg-5 mg (modified release)	30	5	..	25.49	26.56	Triasyn 5.0/5.0	SW
TRANDOLAPRIL with VERAPAMIL HYDROCHLORIDE								
Caution								
The myocardial depressant effects of verapamil hydrochloride and of beta-blocking drugs are additive.								
Restricted benefit								
Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.								
2857J NP	Tablet 4 mg-240 mg (sustained release)	28	5	..	32.69	33.76	Tarka 4/240	AB
9387C NP	Tablet 2 mg-180 mg (sustained release)	28	5	..	22.41	23.48	Tarka 2/180	AB
Angiotensin II antagonists, plain								
Angiotensin II antagonists, plain								
CANDESARTAN CILEXETIL								
8295N NP	Tablet 4 mg	30	5	..	19.25	20.32	Atacand	AP
8296P NP	Tablet 8 mg	30	5	..	22.75	23.82	Atacand	AP
8297Q NP	Tablet 16 mg	30	5	..	28.91	29.98	Atacand	AP
8889W NP	Tablet 32 mg	30	5	..	46.62	34.20	Atacand	AP
EPROSARTAN MESYLATE								
8397Y NP	Tablet 400 mg (base)	56	5	T2.00	*31.08	30.15	Teveten	SM
8447N NP	Tablet 600 mg (base)	28	5	..	30.03	31.10	Teveten	SM
EPROSARTAN MESYLATE								
Authority required								
Adverse effects occurring with all of the base-priced drugs;								
Drug interactions occurring with all of the base-priced drugs;								
Drug interactions expected to occur with all of the base-priced drugs;								
Transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.								
8951D NP	Tablet 400 mg (base)	56	5	..	*31.08	32.15	Teveten	SM

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
IRBESARTAN								
8246B NP	Tablet 75 mg	30	5	..	21.11	22.18	^a Avapro	BQ
							^a Karvea	SW
8247C NP	Tablet 150 mg	30	5	..	25.15	26.22	^a Avapro	BQ
							^a Karvea	SW
8248D NP	Tablet 300 mg	30	5	..	30.24	31.31	^a Avapro	BQ
							^a Karvea	SW
OLMESARTAN MEDOXOMIL								
2147B NP	Tablet 20 mg	30	5	..	25.15	26.22	Olmotec	SH
2148C NP	Tablet 40 mg	30	5	..	30.24	31.31	Olmotec	SH
TELMISARTAN								
8355R NP	Tablet 40 mg	28	5	..	21.66	22.73	Micardis	BY
8356T NP	Tablet 80 mg	28	5	..	28.91	29.98	Micardis	BY
VALSARTAN								
9368C NP	Tablet 40 mg	28	..	..	16.01	17.08	Diovan	NV
9369D NP	Tablet 80 mg	28	5	..	20.13	21.20	Diovan	NV
9370E NP	Tablet 160 mg	28	5	..	23.90	24.97	Diovan	NV
VALSARTAN								
<u>Note</u>								
No applications for increased maximum quantities and/or repeats will be authorised for the 320 mg tablet.								
9371F NP	Tablet 320 mg	28	5	..	28.65	29.72	Diovan	NV

Angiotensin II antagonists, combinations

Angiotensin II antagonists and diuretics

CANDESARTAN CILEXETIL with HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

8504N NP	Tablet 16 mg-12.5 mg	30	5	..	31.13	32.20	Atacand Plus 16/12.5	AP
9314F NP	Tablet 32 mg-12.5 mg	30	5	..	48.55	34.20	Atacand Plus 32/12.5	AP
9315G NP	Tablet 32 mg-25 mg	30	5	..	50.49	34.20	Atacand Plus 32/25	AP

EPROSARTAN MESYLATE with HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

8624X NP	Tablet 600 mg (base)-12.5 mg	28	5	..	32.11	33.18	Teveten Plus 600/12.5	SM

IRBESARTAN with HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
2136K NP	Tablet 300 mg-25 mg	30	5	..	34.69	34.20	^a Avapro HCT 300/25	BQ
							^a Karvezide 300/25	SW
8404H NP	Tablet 150 mg-12.5 mg	30	5	..	27.37	28.44	^a Avapro HCT 150/12.5	BQ
							^a Karvezide 150/12.5	SW
8405J NP	Tablet 300 mg-12.5 mg	30	5	..	32.46	33.53	^a Avapro HCT 300/12.5	BQ
							^a Karvezide 300/12.5	SW

OLMESARTAN MEDOXOMIL with HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

2161R NP	Tablet 20 mg-12.5 mg	30	5	..	27.36	28.43	Olmotec Plus	SH
2166B NP	Tablet 40 mg-12.5 mg	30	5	..	32.44	33.51	Olmotec Plus	SH
2170F NP	Tablet 40 mg-25 mg	30	5	..	34.69	34.20	Olmotec Plus	SH

TELMISARTAN with HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

8622T NP	Tablet 40 mg-12.5 mg	28	5	..	23.73	24.80	Micardis Plus 40/12.5 mg	BY
8623W NP	Tablet 80 mg-12.5 mg	28	5	..	30.98	32.05	Micardis Plus 80/12.5 mg	BY
9381R NP	Tablet 80 mg-25 mg	28	5	..	33.07	34.14	Micardis Plus 80/25 mg	BY

VALSARTAN with HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

9372G NP	Tablet 80 mg-12.5 mg	28	5	..	22.21	23.28	Co-Diovan 80/12.5	NV
9373H NP	Tablet 160 mg-12.5 mg	28	5	..	25.98	27.05	Co-Diovan 160/12.5	NV
9374J NP	Tablet 160 mg-25 mg	28	5	..	28.06	29.13	Co-Diovan 160/25	NV

VALSARTAN with HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

Note

No applications for increased maximum quantities and/or repeats will be authorised for the tablets containing 320 mg valsartan.

9481B NP	Tablet 320 mg-12.5 mg	28	5	..	30.73	31.80	Co-Diovan 320/12.5	NV
9482C NP	Tablet 320 mg-25 mg	28	5	..	32.80	33.87	Co-Diovan 320/25	NV

Angiotensin II antagonists and calcium channel blockers

AMLODIPINE with VALSARTAN

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

5459H NP	Tablet 5 mg (as besylate)-320 mg	28	5	..	38.22	34.20	Exforge 5/320	NV
5460J NP	Tablet 10 mg (as besylate)-320 mg	28	5	..	44.84	34.20	Exforge 10/320	NV
9375K NP	Tablet 5 mg (as besylate)-80 mg	28	5	..	29.71	30.78	Exforge 5/80	NV

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$		
9376L NP	Tablet 5 mg (as besylate)-160 mg	28	5	..	33.48	34.20	Exforge 5/160	NV
9377M NP	Tablet 10 mg (as besylate)-160 mg	28	5	..	40.69	34.20	Exforge 10/160	NV

OLMESARTAN with AMLODIPINE

Restricted benefit

Hypertension in a patient who is not adequately controlled with either of the drugs in the combination.

5292M	Tablet containing olmesartan medoxomil 20 mg with amlodipine 5 mg (as besylate)	30	5	..	35.41	34.20	Sevikar 20/5	SH
5293N	Tablet containing olmesartan medoxomil 40 mg with amlodipine 5 mg (as besylate)	30	5	..	40.49	34.20	Sevikar 40/5	SH
5294P	Tablet containing olmesartan medoxomil 40 mg with amlodipine 10 mg (as besylate)	30	5	..	47.26	34.20	Sevikar 40/10	SH

Angiotensin II antagonists, other combinations

AMLODIPINE with VALSARTAN and HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension in a patient who is not adequately controlled with any two of the drugs in the combination.

5285E	Tablet 5 mg (as besylate)-160 mg-12.5 mg	28	5	..	35.55	34.20	Exforge HCT 5/160/12.5	NV
5286F	Tablet 5 mg (as besylate)-160 mg-25 mg	28	5	..	37.62	34.20	Exforge HCT 5/160/25	NV
5287G	Tablet 10 mg (as besylate)-160 mg-12.5 mg	28	5	..	42.52	34.20	Exforge HCT 10/160/12.5	NV
5288H	Tablet 10 mg (as besylate)-160 mg-25 mg	28	5	..	44.32	34.20	Exforge HCT 10/160/25	NV
5289J	Tablet 10 mg (as besylate)-320 mg-25 mg	28	5	..	48.45	34.20	Exforge HCT 10/320/25	NV

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

GENERAL STATEMENT FOR LIPID-LOWERING DRUGS PRESCRIBED AS PHARMACEUTICAL BENEFITS

Use the following criteria to determine patient eligibility for subsidisation under the PBS for the following drugs:

- atorvastatin calcium
- fluvastatin sodium
- pravastatin sodium
- rosuvastatin calcium
- simvastatin
- fenofibrate
- gemfibrozil

By writing a PBS prescription, the prescriber is certifying the patient satisfies the qualifying criteria set out below and the use is in accordance with the registered indications which differ between agents in this class - refer to the current Product Information for details. Note also that patients already established on a particular lipid-lowering drug, where use satisfies the PBS qualifying criteria, but is outside the registered indications for that drug, are not required to switch to another drug in the class to retain PBS eligibility.

Patients in very high risk categories (see below) may commence drug therapy with statins or fibrates immediately (ie simultaneously with an appropriate diet). For all other patients, dietary therapy should be trialled prior to initiation of drug therapy.


Dietary therapy should be continued concurrently with pharmacological therapy and should be reviewed on at least an annual basis.

Patients identified as being in one of the following very high risk categories may commence drug therapy with statins or fibrates at any cholesterol level:

- coronary heart disease which has become symptomatic
- cerebrovascular disease which has become symptomatic
- peripheral vascular disease which has become symptomatic
- diabetes mellitus with microalbuminuria (defined as urinary albumin excretion rate of >20mcg/min or urinary albumin to creatinine ratio of > 2.5 for males, > 3.5 for females)
- diabetes mellitus in Aboriginal or Torres Strait Islander patients
- diabetes mellitus in patients aged 60 years or more
- family history of coronary heart disease which has become symptomatic before the age of 55 years in two or more first degree relatives
- family history of coronary heart disease which has become symptomatic before the age of 45 years in one or more first degree relatives

If your patient is not identified as being in any of the above very high risk categories, then use the flow-chart and table below to determine whether your patient satisfies the following criteria for subsidisation under the PBS. Document how the patient meets each of these steps in the patient record. Lipid levels must be measured at an accredited laboratory.

Cardiovascular system



POST-DIETARY QUALIFYING CRITERIA

PATIENT CATEGORY

LIPID LEVELS FOR PBS SUBSIDY

Aboriginal or Torres Strait Islander patients	total cholesterol > 6.5 mmol/L
Patients with hypertension	or total cholesterol > 5.5 mmol/L and HDL cholesterol < 1 mmol/L

Patients with HDL cholesterol < 1 mmol/L total cholesterol > 6.5 mmol/L

Patients with familial hypercholesterolaemia identified by:

- DNA mutation; or
- tendon xanthomas in the patient or their first or second degree relative

If aged 18 years or less at treatment initiation:
LDL cholesterol > 4 mmol/L

If aged more than 18 years at treatment initiation:

Patients with:

- family history of coronary heart disease which has become symptomatic before the age of 60 years in one or more first degree relatives; or
- family history of coronary heart disease which has become symptomatic before the age of 50 years in one or more second degree relatives

LDL cholesterol > 5 mmol/L
or
total cholesterol > 6.5 mmol/L
or
total cholesterol > 5.5 mmol/L and
HDL cholesterol < 1 mmol/L

Patients not eligible under the above:

- men aged 35 to 75 years
- post-menopausal women aged up to 75 years

total cholesterol > 7.5 mmol/L
or
triglyceride > 4 mmol/L

Patients not otherwise included	total cholesterol > 9 mmol/L or triglyceride > 8 mmol/L

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Lipid modifying agents

Lipid modifying agents, plain HMG CoA reductase inhibitors

ATORVASTATIN

Restricted benefit

For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.

8213G NP	Tablet 10 mg (as calcium)	30	5	..	42.70	34.20	Lipitor	PF
8214H NP	Tablet 20 mg (as calcium)	30	5	..	58.00	34.20	Lipitor	PF
8215J NP	Tablet 40 mg (as calcium)	30	5	..	79.05	34.20	Lipitor	PF
8521L NP	Tablet 80 mg (as calcium)	30	5	..	110.25	34.20	Lipitor	PF

ATORVASTATIN

Restricted benefit

For use in patients who meet the criteria set out in the General Statement for Lipid-Lowering Drugs, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9230T	Tablet 10 mg (as calcium)	30	11	..	42.70	34.20	Lipitor	PF
9231W	Tablet 20 mg (as calcium)	30	11	..	58.00	34.20	Lipitor	PF
9232X	Tablet 40 mg (as calcium)	30	11	..	79.05	34.20	Lipitor	PF
9233Y	Tablet 80 mg (as calcium)	30	11	..	110.25	34.20	Lipitor	PF

FLUVASTATIN

Restricted benefit

For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.

2863Q NP	Tablet (prolonged release) 80 mg (as sodium)	28	5	..	46.12	34.20	Lescol XL	NV
8023G NP	Capsule 20 mg (as sodium)	28	5	..	25.86	26.93	^a Lescol	NV
				^B 3.13	28.99	26.93	^a Vastin	NM
8024H NP	Capsule 40 mg (as sodium)	28	5	..	30.24	31.31	^a Lescol	NV
				^B 3.44	33.68	31.31	^a Vastin	NM

FLUVASTATIN

Restricted benefit

For use in patients who meet the criteria set out in the General Statement for Lipid-Lowering Drugs, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9234B	Capsule 20 mg (as sodium)	28	11	..	25.86	26.93	^a Lescol	NV
				^B 3.13	28.99	26.93	^a Vastin	NM
9235C	Capsule 40 mg (as sodium)	28	11	..	30.24	31.31	^a Lescol	NV
				^B 3.44	33.68	31.31	^a Vastin	NM
9236D	Tablet (prolonged release) 80 mg (as sodium)	28	11	..	46.12	34.20	Lescol XL	NV

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
PRAVASTATIN								
<u>Restricted benefit</u>								
For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.								
2833D NP	Tablet containing pravastatin sodium 10 mg	30	5	..	21.55	22.62	^a	APO-Pravastatin TX
							^a	Chem mart CH
							^a	Pravastatin AF
							^a	Cholstat 10 AF
							^a	GenRx Pravastatin GX
							^a	Lipostat 10 SI
							^a	Pravastatin 10 CR
							^a	Pravastatin-GA 10 GM
							^a	Pravastatin generichealth GQ
							^a	Pravastatin Sandoz SZ
							^a	Pravastatin Winthrop WA
							^a	Terry White Chemists Pravastatin TW
2834E NP	Tablet containing pravastatin sodium 20 mg	30	5	..	30.50	31.57	^B 3.99	25.54
							^a	22.62
							^a	Pravachol FM
							^a	APO-Pravastatin TX
							^a	Chem mart CH
							^a	Pravastatin AF
							^a	Cholstat 20 AF
							^a	GenRx Pravastatin GX
							^a	Lipostat 20 SI
							^a	Pravastatin 20 CR
							^a	Pravastatin-GA 20 GM
							^a	Pravastatin generichealth GQ
8197K NP	Tablet containing pravastatin sodium 40 mg	30	5	..	43.50	34.20	^B 4.00	34.50
							^a	31.57
							^a	Pravachol FM
							^a	APO-Pravastatin TX
							^a	Chem mart CH
							^a	Pravastatin AF
							^a	Cholstat 40 AF
							^a	GenRx Pravastatin GX
							^a	Lipostat 40 SI
							^a	Pravastatin 40 CR
							^a	Pravastatin-GA 40 GM
							^a	Pravastatin generichealth GQ
8829Q NP	Tablet containing pravastatin sodium 80 mg	30	5	..	61.51	34.20	^B 4.00	47.50
							^a	34.20
							^a	Pravachol FM
							^a	APO-Pravastatin TX
							^a	Chem mart CH
							^a	Pravastatin AF
							^a	Cholstat 40 AF
							^a	GenRx Pravastatin GX
							^a	Lipostat 40 SI
							^a	Pravastatin 40 CR
							^a	Pravastatin-GA 40 GM
							^a	Pravastatin generichealth GQ
							^a	Pravastatin Sandoz SZ
							^a	Pravastatin Winthrop WA
							^a	Terry White Chemists Pravastatin TW
							^a	Vastoran RA

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
							^a	Chem mart CH
								Pravastatin
							^a	Lipostat 80 SI
							^a	Pravastatin-GA 80 GM
							^a	Pravastatin GQ
								generichealth
							^a	Pravastatin Sandoz SZ
							^a	Terry White TW
								Chemists
								Pravastatin
				^B 4.00	65.51	34.20	^a	Pravachol FM
PRAVASTATIN								
<u>Restricted benefit</u>								
For use in patients who meet the criteria set out in the General Statement for Lipid-Lowering Drugs, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.								
<u>Note</u>								
No applications for increased maximum quantities and/or repeats will be authorised.								
9237E	Tablet containing pravastatin sodium 10 mg	30	11	..	21.55	22.62	^a	APO-Pravastatin TX
							^a	Chem mart CH
								Pravastatin
							^a	Cholstat 10 AF
							^a	GenRx Pravastatin GX
							^a	Lipostat 10 SI
							^a	Pravastatin 10 CR
							^a	Pravastatin-GA 10 GM
							^a	Pravastatin GQ
								generichealth
							^a	Pravastatin Sandoz SZ
							^a	Pravastatin WA
								Winthrop
							^a	Terry White TW
								Chemists
								Pravastatin
				^B 3.99	25.54	22.62	^a	Pravachol FM
9238F	Tablet containing pravastatin sodium 20 mg	30	11	..	30.50	31.57	^a	APO-Pravastatin TX
							^a	Chem mart CH
								Pravastatin
							^a	Cholstat 20 AF
							^a	GenRx Pravastatin GX
							^a	Lipostat 20 SI
							^a	Pravastatin 20 CR
							^a	Pravastatin-GA 20 GM
							^a	Pravastatin GQ
								generichealth
							^a	Pravastatin Sandoz SZ
							^a	Pravastatin WA
								Winthrop
							^a	Terry White TW
								Chemists
								Pravastatin
				^B 4.00	34.50	31.57	^a	Vastoran RA
							^a	Pravachol FM
9239G	Tablet containing pravastatin sodium 40 mg	30	11	..	43.50	34.20	^a	APO-Pravastatin TX
							^a	Chem mart CH
								Pravastatin
							^a	Cholstat 40 AF
							^a	GenRx Pravastatin GX

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	
9240H	Tablet containing pravastatin sodium 80 mg	30	11	..	61.51	34.20	^a Lipostat 40 SI
							^a Pravastatin 40 CR
							^a Pravastatin-GA 40 GM
							^a Pravastatin generichealth GQ
							^a Pravastatin Sandoz SZ
							^a Pravastatin Winthrop WA
							^a Terry White Chemists TW
							Pravastatin Vastoran RA
							^a Pravachol FM
							^a APO-Pravastatin TX
							^a Chem mart CH
							Pravastatin Lipostat 80 SI
							^a Pravastatin-GA 80 GM
							^a Pravastatin generichealth GQ
							^a Pravastatin Sandoz SZ
							^a Terry White Chemists TW
							Pravastatin Pravachol FM
				^B 4.00	47.50	34.20	
				^B 4.00	65.51	34.20	

ROSUVASTATIN

Restricted benefit

For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.

9042X NP	Tablet 5 mg (as calcium)	30	5	..	45.75	34.20	Crestor	AP
9043Y NP	Tablet 10 mg (as calcium)	30	5	..	62.50	34.20	Crestor	AP
9044B NP	Tablet 20 mg (as calcium)	30	5	..	86.21	34.20	Crestor	AP
9045C NP	Tablet 40 mg (as calcium)	30	5	..	120.19	34.20	Crestor	AP

ROSUVASTATIN

Restricted benefit

For use in patients who meet the criteria set out in the General Statement for Lipid-Lowering Drugs, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

3402C	Tablet 5 mg (as calcium)	30	11	..	45.75	34.20	Crestor	AP
3403D	Tablet 10 mg (as calcium)	30	11	..	62.50	34.20	Crestor	AP
3404E	Tablet 20 mg (as calcium)	30	11	..	86.21	34.20	Crestor	AP
3405F	Tablet 40 mg (as calcium)	30	11	..	120.19	34.20	Crestor	AP

SIMVASTATIN

Restricted benefit

For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.

2011W NP	Tablet 10 mg	30	5	..	25.04	26.11	^a APO-Simvastatin TX
							^a Chem mart Simvastatin CH
							^a GenRx Simvastatin GX

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer				
2012X NP	Tablet 20 mg	30	5	..	33.20	34.20	^a Pharmacor Simvastatin 10 MI				
							^a Ransim RA				
							^a Simvahexal SZ				
							^a Simvar 10 SI				
							^a Simvastatin-DP GM				
							^a Simvastatin-GA 10 GN				
							^a Simvastatin generichealth GQ				
							^a Simvastatin-Spirit 10 ZP				
							^a Simvastatin Winthrop WA				
							^a Simvasyn CR				
							^a Terry White Chemists Simvastatin TW				
							^a Zimstat AF				
				^B 3.51	28.55	26.11	^a Lipex 10 FR				
				^a Zocor MK							
				2013Y NP	Tablet 5 mg	30	5	..	19.89	20.96	^a APO-Simvastatin TX
											^a Chem mart Simvastatin CH
											^a GenRx Simvastatin GX
											^a Pharmacor Simvastatin 20 MI
											^a Ransim RA
											^a Simvahexal SZ
											^a Simvar 20 SI
											^a Simvastatin-DP GM
											^a Simvastatin-GA 20 GN
											^a Simvastatin generichealth GQ
											^a Simvastatin-Spirit 20 ZP
											^a Simvastatin Winthrop WA
											^a Simvasyn CR
											^a Terry White Chemists Simvastatin TW
											^a Zimstat AF
^B 3.50	36.70	34.20	^a Lipex 20 FR								
^a Zocor MK											
8173E NP	Tablet 40 mg	30	5	..	44.45	34.20	^a Simvahexal SZ				
							^a Simvasyn CR				
							^a Zimstat AF				
							^a Zocor MK				
							^a APO-Simvastatin TX				
							^a Chem mart Simvastatin CH				
							^a GenRx Simvastatin GX				
^a Pharmacor Simvastatin 40 MI											
^a Ransim RA											
^a Simvahexal SZ											
^a Simvar 40 SI											

SIMVASTATIN

For use in patients who meet the criteria set out in the General Statement for Lipid-Lowering Drugs, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

No applications for increased maximum quantities and/or repeats will be authorised.

9241J	Tablet 5 mg	30	11	..	19.89	20.96	^a	Simvahexal	SZ
							^a	Simvasyn	CR
							^a	Zimstat	AF
							^a	Zocor	MK
9242K	Tablet 10 mg	30	11	..	25.04	26.11	^b 3.50		
							^a	APO-Simvastatin	TX
							^a	Chem mart	CH
							^a	Simvastatin	
							^a	GenRx Simvastatin	GX
							^a	Pharmacor	MI
							^a	Simvastatin 10	
							^a	Ransim	RA

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
9243L	Tablet 20 mg	30	11	..	33.20	34.20	^a Simvahexal SZ
							^a Simvar 10 SI
							^a Simvastatin-DP GM
							^a Simvastatin-GA 10 GN
							^a Simvastatin generichealth GQ
							^a Simvastatin-Spirit 10 ZP
							^a Simvastatin Winthrop WA
							^a Simvasyn CR
							^a Terry White Chemists TW
							^a Simvastatin Zimstat AF
							^a Lipex 10 FR
							^a Zocor MK
							^a APO-Simvastatin TX
							^a Chem mart CH
							^a Simvastatin GenRx Simvastatin GX
							^a Pharmacor MI
							^a Simvastatin 20 Ransim RA
							^a Simvahexal SZ
							^a Simvar 20 SI
							^a Simvastatin-DP GM
							^a Simvastatin-GA 20 GN
							^a Simvastatin generichealth GQ
							^a Simvastatin-Spirit 20 ZP
							^a Simvastatin Winthrop WA
							^a Simvasyn CR
							^a Terry White Chemists TW
							^a Simvastatin Zimstat AF
9244M	Tablet 40 mg	30	11	..	44.45	34.20	^B 3.51 28.55 26.11 ^a Lipex 20 FR
							^a Zocor MK
							^a APO-Simvastatin TX
							^a Chem mart CH
							^a Simvastatin GenRx Simvastatin GX
							^a Pharmacor MI
							^a Simvastatin 40 Ransim RA
							^a Simvahexal SZ
							^a Simvar 40 SI
							^a Simvastatin-DP GM
							^a Simvastatin-GA 40 GN
							^a Simvastatin generichealth GQ
							^a Simvastatin-Spirit 40 ZP
							^a Simvastatin Winthrop WA
							^a Simvasyn CR
							^a Terry White Chemists TW
							^a Simvastatin Zimstat AF

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
9245N	Tablet 80 mg	30	11	..	59.42	34.20	Simvastatin	
							^a Zimstat	AF
							^a Lipex 40	FR
							^a Zocor	MK
							^a APO-Simvastatin	TX
							^a Chem mart	CH
							Simvastatin	
							^a GenRx Simvastatin	GX
							^a Pharmacor	MI
							Simvastatin 80	
							^a Ransim	RA
							^a Simvahexal	SZ
							^a Simvar 80	SI
							^a Simvastatin-DP	GM
							^a Simvastatin-GA 80	GN
							^a Simvastatin generichealth	GQ
							^a Simvastatin-Spirit 80	ZP
							^a Simvastatin Winthrop	WA
							^a Simvasyn	CR
							^a Terry White Chemists	TW
							Simvastatin	
							^a Zimstat	AF
				^B 3.50	62.92	34.20	^a Lipex 80	FR
							^a Zocor	MK

Fibrates

FENOFIBRATE

Note

The risk of serious muscle toxicity is increased if fenofibrate is used concomitantly with HMG CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution in patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and patients monitored closely for chronic signs of muscle toxicity.

Restricted benefit

For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.

9022W <i>NP</i>	Tablet 48 mg	60	5	..	30.05	31.12	Lipidil	SM
9023X <i>NP</i>	Tablet 145 mg	30	5	..	41.75	34.20	Lipidil	SM

FENOFIBRATE

Note

The risk of serious muscle toxicity is increased if fenofibrate is used concomitantly with HMG CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution in patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and patients monitored closely for chronic signs of muscle toxicity.

Restricted benefit

For use in patients who meet the criteria set out in the General Statement for Lipid-Lowering Drugs, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9246P	Tablet 48 mg	60	11	..	30.05	31.12	Lipidil	SM
9247Q	Tablet 145 mg	30	11	..	41.75	34.20	Lipidil	SM

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
GEMFIBROZIL								
<u>Note</u>								
The risk of serious muscle toxicity is increased if gemfibrozil is used concomitantly with HMG CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution in patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and patients monitored closely for chronic signs of muscle toxicity.								
<u>Restricted benefit</u>								
For use in patients that meet the criteria set out in the General Statement for Lipid-Lowering Drugs.								
1453L NP	Tablet 600 mg	60	5	..	29.70	30.77	^a	Ausgem SI
							^a	Chem mart CH
							^a	Gemfibrozil Gemhexal SZ
							^a	GenRx Gemfibrozil GX
							^a	Jezil GN
							^a	Lipazil 600 mg GM
							^a	Lipigem AF
							^a	Pharmacor CR
							^a	Gemfibrozil 600 Terry White Chemists TW
							^a	Gemfibrozil Lipid PF
				^B 2.95	32.65	30.77	^a	

GEMFIBROZIL

Note

The risk of serious muscle toxicity is increased if gemfibrozil is used concomitantly with HMG CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution in patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and patients monitored closely for chronic signs of muscle toxicity.

Restricted benefit

For use in patients who meet the criteria set out in the General Statement for Lipid-Lowering Drugs, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9248R	Tablet 600 mg	60	11	..	29.70	30.77	^a	Ausgem SI
							^a	Chem mart CH
							^a	Gemfibrozil Gemhexal SZ
							^a	GenRx Gemfibrozil GX
							^a	Jezil GN
							^a	Lipazil 600 mg GM
							^a	Lipigem AF
							^a	Pharmacor CR
							^a	Gemfibrozil 600 Terry White Chemists TW
							^a	Gemfibrozil Lipid PF
				^B 2.95	32.65	30.77		

Bile acid sequestrants

CHOLESTYRAMINE

2967E NP	Sachets 4.7 g (equivalent to 4 g cholestyramine), 50	2	5	..	*71.94	34.20		Questran Lite SI

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
CHOLESTYRAMINE							
<u>Restricted benefit</u>							
For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised.							
9249T	Sachets 4.7 g (equivalent to 4 g cholestyramine), 50	2	11	..	*71.94	34.20	Questran Lite SI
COLESTIPOL HYDROCHLORIDE							
1224K NP	Sachets 5 g, 120	1	5	..	85.04	34.20	Colestid PF
COLESTIPOL HYDROCHLORIDE							
<u>Restricted benefit</u>							
For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised.							
9250W	Sachets 5 g, 120	1	11	..	85.04	34.20	Colestid PF

Other lipid modifying agents

EZETIMIBE

Authority required (STREAMLINED)

Treatment, in conjunction with dietary therapy and exercise, for co-administration with an HMG CoA reductase inhibitor (statin) in patients whose cholesterol levels are inadequately controlled with a statin and who have:

2649

(a) coronary heart disease; or

2650

(b) diabetes mellitus; or

2651

(c) peripheral vascular disease; or

2652

(d) heterozygous familial hypercholesterolaemia; or

2653

(e) symptomatic cerebrovascular disease; or

2667

(f) family history of coronary heart disease; or

2668

(g) hypertension.

Inadequate control with a statin is defined as follows:

(1) where the patient falls into a category for which the General Statement for Lipid-Lowering Drugs includes an initial cholesterol threshold for PBS-subsidy, a cholesterol level in excess of that threshold after at least 3 months of treatment at a daily dose of 40 mg or greater of a statin, in conjunction with dietary therapy and exercise. The dose and duration of statin treatment and the cholesterol level which shows inadequate control must be documented in the patient's medical records when ezetimibe is initiated. The cholesterol level which shows inadequate control must be no more than 2 months old when ezetimibe is initiated; or

(2) where the patient falls into a category for which the General Statement for Lipid-Lowering Drugs allows PBS-subsidised treatment with a statin at any cholesterol level, a cholesterol level in excess of 4 mmol per L after at least 3 months of treatment at a daily dose of 40 mg or greater of a statin, in conjunction with dietary therapy and exercise. The dose and duration of statin treatment and the cholesterol level which shows inadequate control must be documented in the patient's medical records when ezetimibe is initiated. The cholesterol level which shows inadequate control must be no more than 2 months old when ezetimibe is initiated.

Authority required (STREAMLINED)

1989

Patients eligible for PBS-subsidised lipid-lowering medication (according to the criteria set out in the General Statement for Lipid-Lowering Drugs) where treatment with an HMG CoA reductase inhibitor (statin) is contraindicated;

2669

Patients eligible for PBS-subsidised lipid-lowering medication (according to the criteria set out in the General Statement for Lipid-Lowering Drugs) where treatment with an HMG CoA reductase inhibitor (statin) must be discontinued or reduced to a dose of 20 mg or less per day, because the patient developed a clinically important product-related adverse event during treatment with a statin.

A clinically important product-related adverse event is defined as follows:

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	
	(i) Severe myalgia (muscle symptoms without CK elevation) which is proven to be temporally associated with statin treatment; or (ii) Myositis (clinically important CK elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or (iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.						
	<u>Authority required (STREAMLINED)</u>						
	1991						
	Homozygous sitosterolaemia;						
	2438						
	Patients with homozygous familial hypercholesterolaemia who are eligible for PBS-subsidised lipid-lowering medication (according to the criteria set out in the General Statement for Lipid-Lowering Drugs), in combination with an HMG CoA reductase inhibitor (statin).						
	<u>Note</u>						
	Continuing Therapy Only:						
	For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.						
8757X NP	Tablet 10 mg	30	5	..	70.97	34.20	Ezetrol MK

Lipid modifying agents, combinations

HMG CoA reductase inhibitors in combination with other lipid modifying agents

EZETIMIBE with SIMVASTATIN

Authority required (STREAMLINED)

2431

Patients with homozygous familial hypercholesterolaemia who are eligible for PBS-subsidised lipid-lowering medication (according to the criteria set out in the General Statement for Lipid-Lowering Drugs);

3194

Patients eligible for PBS-subsidised lipid-lowering medication (according to the criteria set out in the General Statement for Lipid-Lowering Drugs) where treatment with an HMG CoA reductase inhibitor (statin) must be reduced to a dose of 20 mg or less per day, because the patient developed a clinically important product-related adverse event during treatment with a statin.

A clinically important product-related adverse event is defined as follows:

- (i) Severe myalgia (muscle symptoms without CK elevation) which is proven to be temporally associated with statin treatment; or
- (ii) Myositis (clinically important CK elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- (iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9483D NP	Tablet 10 mg-10 mg	30	5	..	88.79	34.20	Vytorin MK
9484E NP	Tablet 10 mg-20 mg	30	5	..	96.59	34.20	Vytorin MK

EZETIMIBE with SIMVASTATIN

Authority required (STREAMLINED)

Treatment, in conjunction with dietary therapy and exercise, in patients whose cholesterol levels are inadequately controlled with an HMG CoA reductase inhibitor (statin) and who have:

2654

(a) coronary heart disease; or

2655

(b) diabetes mellitus; or

2656

(c) peripheral vascular disease; or

2657

(d) heterozygous familial hypercholesterolaemia; or

2658

(e) cerebrovascular disease which has become symptomatic; or

2678

(f) family history of coronary heart disease; or

2679

(g) hypertension;

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Inadequate control with a statin is defined as follows: (1) where the patient falls into a category for which the General Statement for Lipid-Lowering Drugs includes an initial cholesterol threshold for PBS-subsidy, a cholesterol level in excess of that threshold after at least 3 months of treatment at a daily dose of 40 mg or greater of a statin, in conjunction with dietary therapy and exercise. The dose and duration of statin treatment and the cholesterol level which shows inadequate control must be documented in the patient's medical records when the ezetimibe component is initiated. The cholesterol level which shows inadequate control must be no more than 2 months old when the ezetimibe component is initiated; or (2) where the patient falls into a category for which the General Statement for Lipid-Lowering Drugs allows PBS-subsidised treatment with a statin at any cholesterol level, a cholesterol level in excess of 4 mmol per L after at least 3 months of treatment at a daily dose of 40 mg or greater of a statin, in conjunction with dietary therapy and exercise. The dose and duration of statin treatment and the cholesterol level which shows inadequate control must be documented in the patient's medical records when the ezetimibe component is initiated. The cholesterol level which shows inadequate control must be no more than 2 months old when the ezetimibe component is initiated. 2431 Patients with homozygous familial hypercholesterolaemia who are eligible for PBS-subsidised lipid-lowering medication (according to the criteria set out in the General Statement for Lipid-Lowering Drugs). Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
8881K NP	Tablet 10 mg-40 mg	30	5	..	107.85	34.20	Vytorin MK
8882L NP	Tablet 10 mg-80 mg	30	5	..	123.97	34.20	Vytorin MK

HMG CoA reductase inhibitors, other combinations

AMLODIPINE BESYLATE with ATORVASTATIN CALCIUM

Restricted benefit

For use in patients who have hypertension and/or angina and who meet the criteria set out in the General Statement for Lipid-Lowering Drugs, and:

(a) who are currently receiving treatment with a dihydropyridine calcium channel blocker; OR

(b) whose blood pressure and/or angina is inadequately controlled with other classes of antihypertensive and/or anti-anginal agent, and in whom adjunctive therapy with a dihydropyridine calcium channel blocker would be appropriate; OR

(c) who are intolerant of the side effects of other classes of antihypertensive and/or anti-anginal agent, and in whom replacement therapy with a dihydropyridine calcium channel blocker would be appropriate.

9049G NP	Tablet 5 mg (base)-10 mg (base)	30	5	..	51.62	34.20	Caduet 5/10 PF
9050H NP	Tablet 5 mg (base)-20 mg (base)	30	5	..	67.81	34.20	Caduet 5/20 PF
9051J NP	Tablet 5 mg (base)-40 mg (base)	30	5	..	88.87	34.20	Caduet 5/40 PF
9052K NP	Tablet 5 mg (base)-80 mg (base)	30	5	..	120.06	34.20	Caduet 5/80 PF
9053L NP	Tablet 10 mg (base)-10 mg (base)	30	5	..	58.57	34.20	Caduet 10/10 PF
9054M NP	Tablet 10 mg (base)-20 mg (base)	30	5	..	75.19	34.20	Caduet 10/20 PF
9055N NP	Tablet 10 mg (base)-40 mg (base)	30	5	..	96.25	34.20	Caduet 10/40 PF
9056P NP	Tablet 10 mg (base)-80 mg (base)	30	5	..	127.44	34.20	Caduet 10/80 PF

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Dermatologicals

Antifungals for dermatological use

Antifungals for topical use

Antibiotics

NYSTATIN

Authority required (STREAMLINED)

2354

Treatment of a fungal or a yeast infection in an Aboriginal or a Torres Strait Islander person.

1698J NP	Cream 100,000 units per g, 15 g	2	3	..	*18.56	19.63	Mycostatin	FM

Imidazole and triazole derivatives

CLOTRIMAZOLE

Authority required (STREAMLINED)

2354

Treatment of a fungal or a yeast infection in an Aboriginal or a Torres Strait Islander person.

1017M NP	Cream 10 mg per g (1%), 20 g	2	3	..	*11.26	12.33	Clonea	AF

KETOCONAZOLE

Authority required (STREAMLINED)

2354

Treatment of a fungal or a yeast infection in an Aboriginal or a Torres Strait Islander person.

1574W NP	Shampoo 20 mg per g (2%), 60 mL	£1	1	..	18.31	19.38	Nizoral 2%	JT
9024Y NP	Cream 20 mg per g (2%), 30 g	£1	2	..	23.12	24.19	Nizoral 2% Cream	JT
9025B NP	Shampoo 10 mg per g (1%), 100 mL	£1	1	..	17.60	18.67	Nizoral 1%	JT

MICONAZOLE

Authority required (STREAMLINED)

2354

Treatment of a fungal or a yeast infection in an Aboriginal or a Torres Strait Islander person.

9031H NP	Tincture 20 mg per mL (2%), 30 mL	£1	2	..	19.47	20.54	Daktarin	JT

MICONAZOLE NITRATE

Authority required (STREAMLINED)

2354

Treatment of a fungal or a yeast infection in an Aboriginal or a Torres Strait Islander person.

9026C NP	Cream 20 mg per g (2%), 15 g	2	3	..	*15.90	16.97	Daktarin	JT
9027D NP	Cream 20 mg per g (2%), 30 g	£1	2	..	14.79	15.86	Daktarin	JT
9028E NP	Cream 20 mg per g (2%), 70 g	£1	1	..	16.79	17.86	Daktarin	JT
9029F NP	Powder 20 mg per g (2%), 30 g	£1	2	..	15.56	16.63	Daktarin	JT
9030G NP	Lotion 20 mg per mL (2%), 30 g	£1	2	..	16.72	17.79	Daktarin	JT

Other antifungals for topical use

TERBINAFINE

Authority required (STREAMLINED)

2354

Treatment of a fungal or a yeast infection in an Aboriginal or a Torres Strait Islander person;

Dermatologicals

					Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	\$	\$	Brand Name and Manufacturer	
3243								
Treatment of a fungal or a yeast infection in a patient aged up to 18 years inclusive.								
9160D NP	Cream containing terbinafine hydrochloride 10 mg per g (1%), 15 g	2	3	..	*37.36	34.20	Lamisil	NC

Antifungals for systemic use

Antifungals for systemic use

GRISEOFULVIN								
1460W NP	Tablet 125 mg	100	2	..	25.87	26.94	Grisovin	SI
2982Y NP	Tablet 500 mg	28	2	..	26.99	28.06	Grisovin 500	SI

TERBINAFINE

Authority required

Treatment of a dermatophyte infection in an Aboriginal or a Torres Strait Islander person where topical treatment has failed;

Treatment of a dermatophyte infection in a patient aged up to 18 years inclusive where topical treatment and griseofulvin have failed.

2285G NP	Tablet 250 mg (as hydrochloride)	42	..	..	102.82	34.20	^a GenRx Terbinafine	GX
							^a Sebifin 250	RA
							^a Tamsil	SI
							^a Terbihexal	SZ
							^a Terbinafine 250	CR
							^a Terbinafine-DRLA	RZ
							^a Terbinafine-GA	GM
							^a Terbix 250	MI
							^a Zabel	AF
				^B 1.46	104.28	34.20	^a Lamisil	NV

TERBINAFINE

Authority required

Proximal or extensive (greater than 80% nail involvement) onychomycosis due to dermatophyte infection where topical treatment has failed. This infection must be proven by microscopy or culture and confirmed by an Approved Pathology Authority. The date of the pathology report must be provided at the time of application and must not be more than 12 months old.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

2804N NP	Tablet 250 mg (as hydrochloride)	42	1	..	102.82	34.20	^a GenRx Terbinafine	GX
							^a Sebifin 250	RA
							^a Tamsil	SI
							^a Terbihexal	SZ
							^a Terbinafine 250	CR
							^a Terbinafine-DRLA	RZ
							^a Terbinafine-GA	GM
							^a Terbix 250	MI
							^a Zabel	AF
				^B 1.46	104.28	34.20	^a Lamisil	NV

Antipsoriatics

Antipsoriatics for topical use

Tars

COAL TAR - PREPARED								
8864M	Gel 10 mg per g (1%), 100 mL	1	2	..	33.08	34.15	Exorex	GM

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	

Other antipsoriatics for topical use

CALCIPOTRIOL

Restricted benefit

Chronic stable plaque type psoriasis vulgaris.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2080L NP	Cream 50 micrograms per g (0.005%), 30 g	\$1	1	..	28.06	29.13	Daivonex	CS

CALCIPOTRIOL

Restricted benefit

Chronic stable plaque type psoriasis vulgaris of the scalp.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9163G NP	Scalp solution 50 micrograms per mL (0.005%), 30 mL	\$1	1	..	28.06	29.13	Daivonex	CS

CALCIPOTRIOL with BETAMETHASONE DIPROPIONATE

Restricted benefit

Chronic stable plaque type psoriasis vulgaris in a patient who is not adequately controlled with either calcipotriol or potent topical corticosteroid monotherapy.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9494Q NP	Ointment 50 micrograms-500 micrograms (base) per g (0.005%-0.05%), 30 g	\$1	1	..	42.14	34.20	Daivobet	CS

Antipsoriatics for systemic use

Retinoids for treatment of psoriasis

ACITRETIN

Caution

This drug is a potent teratogen—pregnancy should be avoided for at least two years after cessation of therapy.

Note

Care must be taken to comply with the provisions of State/Territory law when prescribing acitretin.

Authority required (STREAMLINED)

1366

Severe intractable psoriasis;

1363

Severe forms of disorders of keratinisation.

2019G	Capsule 10 mg	100	2	..	205.77	34.20	Neotigason	TA
2020H	Capsule 25 mg	100	2	..	393.21	34.20	Neotigason	TA

Antibiotics and chemotherapeutics for dermatological use

Chemotherapeutics for topical use

Sulfonamides

SILVER SULFADIAZINE

Restricted benefit

Prevention and treatment of infection in partial or full skin thickness loss due to burns;

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$		
Prevention and treatment of infection in partial or full skin thickness loss due to epidermolysis bullosa; Stasis ulcers.								
9479X NP	Cream 10 mg per g (1%), 50 g	‡1	..	..	19.15	20.22	Flamazine	SN

Corticosteroids, dermatological preparations

Corticosteroids, plain

Corticosteroids, weak (group I)

HYDROCORTISONE ACETATE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

2881P NP	Cream 10 mg per g (1%), 50 g	‡1	1	..	8.61	9.68	^a	Cortic-DS 1%	FM
				^B 0.08	8.69	9.68		Cortef	VT
				^B 2.76	11.37	9.68	^a	Sigmacort	SI
2882Q NP	Topical ointment 10 mg per g (1%), 50 g	‡1	1	..	8.61	9.68	^a	Cortic-DS 1%	FM
				^B 2.76	11.37	9.68	^a	Sigmacort	SI
2887Y NP	Cream 10 mg per g (1%), 30 g	‡1	1	..	8.94	10.01	^a	Cortic-DS 1%	FM
				^B 2.75	11.69	10.01	^a	Sigmacort	SI
2888B NP	Topical ointment 10 mg per g (1%), 30 g	‡1	1	..	8.94	10.01	^a	Cortic-DS 1%	FM
				^B 2.75	11.69	10.01	^a	Sigmacort	SI

Corticosteroids, moderately potent (group II)

TRIAMCINOLONE ACETONIDE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

2117K NP	Cream 200 micrograms per g (0.02%), 100 g	2	..	..	*14.56	15.63	^a	Tricortone	FM
				^B 3.84	*18.40	15.63	^a	Aristocort 0.02%	SI
2118L NP	Ointment 200 micrograms per g (0.02%), 100 g	2	..	..	*14.56	15.63	^a	Tricortone	FM
				^B 3.84	*18.40	15.63	^a	Aristocort 0.02%	SI

Corticosteroids, potent (group III)

BETAMETHASONE DIPROPIONATE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1115Q NP	Cream 500 micrograms (base) per g (0.05%), 15 g	‡1	1	..	13.27	14.34	^a	Eleuphrat	EX
				^B 2.50	15.77	14.34	^a	Diprosone	SH
1119X NP	Ointment 500 micrograms (base) per g (0.05%), 15 g	‡1	1	..	13.27	14.34	^a	Eleuphrat	EX
				^B 2.50	15.77	14.34	^a	Diprosone	SH

BETAMETHASONE VALERATE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

2812B NP	Cream 200 micrograms (base) per g (0.02%), 100 g	2	..	..	*24.60	25.67	^a	Antroquoril	EX
							^b	Cortival 1/5	FM

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer		
					\$	\$			
2820K NP	Ointment 200 micrograms (base) per g (0.02%), 100 g	2	..	^B 2.50	*27.10	25.67	^a	Celestone-M	SH
				^B 7.00	*31.60	25.67	^b	Betnovate 1/5	SI
				..	*24.60	25.67	^a	Antroquoril	EX
				^B 2.50	*27.10	25.67	^a	Celestone-M	SH

BETAMETHASONE VALERATE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2813C NP	Cream 500 micrograms (base) per g (0.05%), 15 g	‡1	1	..	8.46	9.53	^b	Cortival 1/2	FM
				^B 3.00	11.46	9.53	^b	Betnovate 1/2	SI
2815E NP	Ointment 500 micrograms (base) per g (0.05%), 15 g	‡1	1	..	8.46	9.53	^b	Cortival 1/2	FM
				^B 3.00	11.46	9.53	^b	Betnovate 1/2	SI

METHYLPREDNISOLONE ACEPONATE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8054X NP	Cream 1 mg per g (0.1%), 15 g	‡1	..	..	14.13	15.20		Advantan	CS
8055Y NP	Ointment 1 mg per g (0.1%), 15 g	‡1	..	..	14.13	15.20		Advantan	CS
8128T NP	Fatty ointment 1 mg per g (0.1%), 15 g	‡1	..	..	14.13	15.20		Advantan	CS

METHYLPREDNISOLONE ACEPONATE

Restricted benefit

Eczema.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8618N NP	Lotion 1 mg per g (0.1%), 20 g	‡1	..	..	14.83	15.90		Advantan	CS

MOMETASONE FUROATE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1913Q NP	Cream 1 mg per g (0.1%), 15 g	‡1	..	..	14.07	15.14	^a	Novasone	EX
				^B 2.50	16.57	15.14	^a	Elocon	SH
1915T NP	Ointment 1 mg per g (0.1%), 15 g	‡1	..	..	14.07	15.14	^a	Novasone	EX
				^B 2.50	16.57	15.14	^a	Elocon	SH

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	
8043H	Lotion 1 mg per g (0.1% w/w), 30 mL	‡1	..	..	18.47	19.54 ^a	Novasone EX
NP				^B 2.50	20.97	19.54 ^a	Elocon SH

Anti-acne preparations

Anti-acne preparations for systemic use

Retinoids for treatment of acne

ISOTRETINOIN

Caution

This drug causes birth defects. Isotretinoin has been reported to cause other frequent and potentially serious toxicity.

Note

Care must be taken to comply with the provisions of State/Territory law when prescribing isotretinoin.

Authority required (STREAMLINED)

1354

Severe cystic acne not responsive to other therapy.

2549E	Capsule 40 mg	30	3	..	125.48	34.20	Oratane	GM
2591J	Capsule 10 mg	60	3	..	91.07	34.20 ^a	Oratane	GM
						^a	Roaccutane	RO
2592K	Capsule 20 mg	60	3	..	138.34	34.20 ^a	GenRx Isotretinoin	GX
						^a	Oratane	GM
				^B 2.29	140.63	34.20 ^a	Roaccutane	RO

Other dermatological preparations

Other dermatological preparations

Agents for atopic dermatitis, excluding corticosteroids

PIMECROLIMUS

Authority required

Treatment of facial or eyelid atopic dermatitis in patients aged at least 3 months with 1 or more of the following contraindications to topical corticosteroids:

- (i) perioral dermatitis;
- (ii) periorbital dermatitis;
- (iii) rosacea;
- (iv) epidermal atrophy;
- (v) dermal atrophy;
- (vi) allergy to topical corticosteroids;
- (vii) cataracts;
- (viii) glaucoma;
- (ix) raised intraocular pressure.

Authority required

Short-term (up to 3 weeks) intermittent treatment of atopic dermatitis of the face or eyelids in patients aged at least 3 months who fail to achieve satisfactory disease control with intermittent topical corticosteroid therapy, and where more than 3 months have passed since the initial diagnosis of atopic dermatitis.

Failure to achieve satisfactory disease control with intermittent topical corticosteroid therapy is manifest by:

- (i) failure of the facial skin to clear despite at least 2 weeks of topical hydrocortisone 1% applied every day; or
- (ii) failure of the facial skin to clear despite at least 1 week of a moderate or potent topical corticosteroid applied every day; or
- (iii) clearing of the facial skin with at least 2 weeks of topical hydrocortisone 1% applied every day, but almost immediate and significant flare in facial disease (within 48 hours) upon stopping topical corticosteroids, occurring on at least 2 consecutive occasions; or
- (iv) clearing of the facial skin with at least 1 week of a moderate or potent topical corticosteroid applied every day, but almost immediate and significant flare in facial disease (within 48 hours) upon stopping topical corticosteroids, occurring on at least 2 consecutive occasions.

Note

No applications for increased maximum quantities and/or repeats will be authorised. Only 1 authority application per 6 months, per patient, will be authorised.

8802G	Cream 10 mg per g (1%), 15 g	‡1	1	..	33.79	34.20	Elidel	NV

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Other dermatologicals

DAPSONE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1272Y NP	Tablet 100 mg	100	1	..	113.84	34.20	Link Medical Products Pty Ltd	LM
8801F NP	Tablet 25 mg	100	1	..	100.58	34.20	Link Medical Products Pty Ltd	LM

IMIQUIMOD

Authority required

Treatment of biopsy confirmed primary (previously untreated) superficial basal cell carcinoma (sBCC) in patients with normal immune function for whom surgical excision, cryotherapy, or curettage with diathermy are inappropriate and topical drug therapy is required.

The date of the pathology report and name of the Approved Pathology Authority must be provided at the time of application.

Note

The patient or carer must be able to understand and administer the imiquimod dosing regimen.

No applications for increased maximum quantities and/or repeats will be authorised.

Treatment of recurrent (previously treated) lesions will not be authorised.

2546B	Cream 50 mg per g (5%), 250 mg single use sachets, 12	1	1	..	159.95	34.20	Aldara	IA

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Genito urinary system and sex hormones

Other gynecologicals

Contraceptives for topical use

Intrauterine contraceptives

LEVONORGESTREL

Restricted benefit

Contraception;

Idiopathic menorrhagia where oral treatments are ineffective;

Idiopathic menorrhagia where oral treatments are contraindicated.

8633J NP	Intrauterine drug delivery system 52 mg (releasing approximately 20 micrograms per 24 hours)	1	..	..	246.41	34.20	Mirena	SC

Other gynecologicals

Prolactine inhibitors

BROMOCRIPTINE MESYLATE

Restricted benefit

Prevention of the onset of lactation in the puerperium for medical reasons.

1444B NP	Tablet 2.5 mg (base)	30	..	..	19.58	20.65	^a Kripton 2.5	AF
				^B 2.84	22.42	20.65	^a Parlodel	NV

BROMOCRIPTINE MESYLATE

Restricted benefit

Acromegaly;

Parkinson's disease;

Pathological hyperprolactinaemia where surgery is not indicated;

Pathological hyperprolactinaemia where surgery has already been used with incomplete resolution;

Pathological hyperprolactinaemia where radiotherapy is not indicated;

Pathological hyperprolactinaemia where radiotherapy has already been used with incomplete resolution.

Note

Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.

1443Y	Tablet 2.5 mg (base)	60	5	..	32.73	33.80	^a Kripton 2.5	AF
				^B 2.92	35.65	33.80	^a Parlodel	NV
1445C	Capsule 10 mg (base)	100	5	..	155.94	34.20	^a Kripton 10	AF
				^B 3.08	159.02	34.20	^a Parlodel	NV
1446D	Capsule 5 mg (base)	60	5	..	50.25	34.20	^a Kripton 5	AF
				^B 2.91	53.16	34.20	^a Parlodel	NV

CABERGOLINE

Restricted benefit

Prevention of the onset of lactation in the puerperium for medical reasons.

8115D NP	Tablet 500 micrograms	2	..	..	24.07	25.14	Dostinex	PF

CABERGOLINE

Authority required (STREAMLINED)

2659

Pathological hyperprolactinaemia where surgery is not indicated;

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer		
	2660 Pathological hyperprolactinaemia where surgery has already been used with incomplete resolution;								
	2661 Pathological hyperprolactinaemia where radiotherapy is not indicated;								
	2662 Pathological hyperprolactinaemia where radiotherapy has already been used with incomplete resolution.								
8114C	Tablet 500 micrograms	8	5	..	66.27	34.20	^a ^a	Dostinex Tinexa	PF SI
QUINAGOLIDE HYDROCHLORIDE									
<u>Authority required (STREAMLINED)</u>									
	2659 Pathological hyperprolactinaemia where surgery is not indicated;								
	2660 Pathological hyperprolactinaemia where surgery has already been used with incomplete resolution;								
	2661 Pathological hyperprolactinaemia where radiotherapy is not indicated;								
	2662 Pathological hyperprolactinaemia where radiotherapy has already been used with incomplete resolution.								
8822H	Tablet 75 micrograms (base)	30	5	..	54.79	34.20		Norprolac	FP
8860H	Pack containing 3 tablets 25 micrograms (base) and 3 tablets 50 micrograms (base)	±1	..	..	11.47	12.54		Norprolac	FP

Sex hormones and modulators of the genital system

Hormonal contraceptives for systemic use

Progestogens and estrogens, fixed combinations

LEVONORGESTREL with ETHINYLOESTRADIOL								
1394J NP	Pack containing 21 tablets 150 micrograms- 30 micrograms and 7 inert tablets	4	2	..	17.21	18.28	^b Monofeme 28	WX
							^a Levlen ED	SY
				^B 13.82	31.03	18.28	^b Nordette 28	WY
				^B 13.87	31.08	18.28	^a Microgynon 30 ED	SC
1456P NP	Pack containing 21 tablets 125 micrograms- 50 micrograms and 7 inert tablets	4	2	..	17.21	18.28	Microgynon 50 ED	SC
NORETHISTERONE with ETHINYLOESTRADIOL								
<u>Note</u> This product may be interchanged with the products in item 2774B.								
2772X NP	Tablets 500 micrograms-35 micrograms, 21	4	2	^B 7.64	*24.26	17.69	^a Brevinor	PF
NORETHISTERONE with ETHINYLOESTRADIOL								
<u>Note</u> These products may also be interchanged with item 2772X.								
2774B NP	Pack containing 21 tablets 500 micrograms- 35 micrograms and 7 inert tablets	4	2	..	*16.62	17.69	^a Norimin 28 Day	KR
				^B 7.88	*24.50	17.69	^a Brevinor	PF
NORETHISTERONE with ETHINYLOESTRADIOL								
<u>Note</u> This product may be interchanged with the products in item 2775C.								
2773Y NP	Tablets 1 mg-35 micrograms, 21	4	2	^B 7.64	*24.26	17.69	^a Brevinor-1	PF

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
NORETHISTERONE with ETHINYLOESTRADIOL								
Note								
These products may also be interchanged with item 2773Y.								
2775C NP	Pack containing 21 tablets 1 mg-35 micrograms and 7 inert tablets	4	2	..	*16.62	17.69	^a	Norimin-1 28 Day KR
				^B 7.88	*24.50	17.69	^a	Brevinor-1 PF
NORETHISTERONE with MESTRANOL								
3176E NP	Tablets 1 mg-50 micrograms, 21	4	2	..	*16.62	17.69		Norinyl-1 PF
3179H NP	Pack containing 21 tablets 1 mg-50 micrograms and 7 inert tablets	4	2	..	*16.62	17.69		Norinyl-1/28 PF
Progestogens and estrogens, sequential preparations								
LEVONORGESTREL with ETHINYLOESTRADIOL								
1392G NP	Pack containing 6 tablets 50 micrograms- 30 micrograms, 5 tablets 75 micrograms- 40 micrograms, 10 tablets 125 micrograms- 30 micrograms and 7 inert tablets	4	2	..	17.21	18.28	^b	Trifeme 28 WX
							^a	Logynon ED SY
				^B 13.82	31.03	18.28	^b	Triphasil 28 WY
				^B 13.87	31.08	18.28	^a	Triquilar ED SC
NORETHISTERONE with ETHINYLOESTRADIOL								
2776D NP	Pack containing 12 tablets 500 micrograms- 35 micrograms, 9 tablets 1 mg-35 micrograms and 7 inert tablets	4	2	..	*16.62	17.69	^a	Improvil 28 Day KR
				^B 7.88	*24.50	17.69	^a	Synphasic PF
Progestogens								
ETONOGESTREL								
8487Q NP	Subcutaneous implant 68 mg	1	..	..	215.92	34.20		Implanon SH
LEVONORGESTREL								
2913H NP,MW	Tablets 30 micrograms, 28	4	2	..	17.32	18.39		Microlut 28 SC
MEDROXYPROGESTERONE ACETATE								
3118D NP	Injection 150 mg in 1 mL	1	1	..	21.59	22.66	^a	Depo-Ralovera KR
				^B 3.26	24.85	22.66	^a	Depo-Provera PF
NORETHISTERONE								
1967M NP	Tablets 350 micrograms, 28	4	2	..	*16.62	17.69	^a	Locilan 28 Day KR
				^B 4.00	*20.62	17.69	^a	Micronor JC
							^a	Noriday 28 Day PF

Androgens

3-oxoandrostens (4) derivatives

TESTOSTERONE

Authority required

Androgen deficiency in males with established pituitary or testicular disorders;

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$		
	Androgen deficiency in males 40 years and older who do not have established pituitary or testicular disorders other than aging, confirmed by at least 2 morning blood samples taken on different mornings. Androgen deficiency is confirmed by testosterone less than 8 nmol per L, or 8-15 nmol per L with high LH (greater than 1.5 times the upper limit of the eugonadal reference range for young men);							
	Micropenis, pubertal induction, or constitutional delay of growth or puberty, in males under 18 years of age.							
8098F	Subcutaneous implant 100 mg	6	..	..	*209.58	34.20	Schering-Plough Pty Limited	SH
8099G	Subcutaneous implant 200 mg	3	..	..	*209.55	34.20	Schering-Plough Pty Limited	SH
8460G	Transdermal patches 12.2 mg (releasing approximately 2.5 mg per 24 hours), 60	†1	5	..	95.84	34.20	Androderm	HH
8619P	Transdermal patches 24.3 mg (releasing approximately 5 mg per 24 hours), 30	†1	5	..	95.84	34.20	Androderm	HH
8830R	Transdermal gel 50 mg in 5 g sachet, 30	†1	5	..	95.12	34.20	Testogel	SC

TESTOSTERONE ENANTHATE

Authority required

Androgen deficiency in males with established pituitary or testicular disorders;

Androgen deficiency in males 40 years and older who do not have established pituitary or testicular disorders other than aging, confirmed by at least 2 morning blood samples taken on different mornings. Androgen deficiency is confirmed by testosterone less than 8 nmol per L, or 8-15 nmol per L with high LH (greater than 1.5 times the upper limit of the eugonadal reference range for young men);

Micropenis, pubertal induction, or constitutional delay of growth or puberty, in males under 18 years of age.

2114G	Injection 250 mg in 1 mL	3	3	..	33.48	34.20	Primoteston Depot SC

TESTOSTERONE ESTERS

Authority required

Androgen deficiency in males with established pituitary or testicular disorders;

Androgen deficiency in males 40 years and older who do not have established pituitary or testicular disorders other than aging, confirmed by at least 2 morning blood samples taken on different mornings. Androgen deficiency is confirmed by testosterone less than 8 nmol per L, or 8-15 nmol per L with high LH (greater than 1.5 times the upper limit of the eugonadal reference range for young men);

Micropenis, pubertal induction, or constitutional delay of growth or puberty, in males under 18 years of age.

2101N	Injection 250 mg	3	3	..	*33.48	34.20	Sustanon 250 SH

TESTOSTERONE UNDECANOATE

Authority required

Androgen deficiency in males with established pituitary or testicular disorders;

Androgen deficiency in males 40 years and older who do not have established pituitary or testicular disorders other than aging, confirmed by at least 2 morning blood samples taken on different mornings. Androgen deficiency is confirmed by testosterone less than 8 nmol per L, or 8-15 nmol per L with high LH (greater than 1.5 times the upper limit of the eugonadal reference range for young men);

Micropenis, pubertal induction, or constitutional delay of growth or puberty, in males under 18 years of age.

2115H	Capsule 40 mg	60	5	..	37.53	34.20	Andriol Testocaps SH
9004X	I.M. injection 1,000 mg in 4 mL	1	1	..	147.41	34.20	Reandron 1000 SC

Estrogens

Natural and semisynthetic estrogens, plain

OESTRADIOL

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

Oestradiol should be used in conjunction with an oral progestogen in women with an intact uterus.

1743R NP	Transdermal patches 2 mg (releasing approximately 25 micrograms per 24 hours), 8	†1	5	..	17.31	18.38	Estraderm 25 NV
1745W NP	Transdermal patches 8 mg (releasing approximately 100 micrograms per 24 hours), 8	†1	5	..	19.39	20.46	Estraderm 100 NV
8125P NP	Transdermal patches 3.8 mg (releasing approximately 50 micrograms per 24 hours), 4	†1	5	..	17.31	18.38	Climara 50 SC

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$		
8126Q NP	Transdermal patches 7.6 mg (releasing approximately 100 micrograms per 24 hours), 4	¥1	5	..	19.39	20.46	Climara 100	SC
8140K NP	Transdermal patches 1.5 mg (releasing approximately 50 micrograms per 24 hours), 8	¥1	5	..	17.31	18.38	Estraderm MX 50	NV
8286D NP	Transdermal gel 1 mg in 1 g sachet, 28	¥1	5	..	17.31	18.38	Sandrena	SH
8311K NP	Transdermal patches 750 micrograms (releasing approximately 25 micrograms per 24 hours), 8	¥1	5	..	17.31	18.38	Estraderm MX 25	NV
8312L NP	Transdermal patches 3 mg (releasing approximately 100 micrograms per 24 hours), 8	¥1	5	..	19.39	20.46	Estraderm MX 100	NV
8485N NP	Transdermal patches 2 mg (releasing approximately 25 micrograms per 24 hours), 4	¥1	5	..	17.31	18.38	Climara 25	SC
8486P NP	Transdermal patches 5.7 mg (releasing approximately 75 micrograms per 24 hours), 4	¥1	5	..	19.39	20.46	Climara 75	SC
8761D NP	Transdermal patches 390 micrograms (releasing approximately 25 micrograms per 24 hours), 8	¥1	5	..	17.31	18.38	Estradot 25	NV
8762E NP	Transdermal patches 585 micrograms (releasing approximately 37.5 micrograms per 24 hours), 8	¥1	5	..	17.31	18.38	Estradot 37.5	NV
8763F NP	Transdermal patches 780 micrograms (releasing approximately 50 micrograms per 24 hours), 8	¥1	5	..	17.31	18.38	Estradot 50	NV
8764G NP	Transdermal patches 1.17 mg (releasing approximately 75 micrograms per 24 hours), 8	¥1	5	..	19.39	20.46	Estradot 75	NV
8765H NP	Transdermal patches 1.56 mg (releasing approximately 100 micrograms per 24 hours), 8	¥1	5	..	19.39	20.46	Estradot 100	NV

OESTRADIOL

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1742Q NP	Vaginal tablets 25 micrograms, 15	¥1	2	..	23.24	24.31	Vagifem	NO
8274L NP	Tablet 2 mg	56	2	..	13.70	14.77	Zumenon	SM

OESTRADIOL VALERATE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1663M NP	Tablet 1 mg	56	2	..	11.78	12.85	Progynova	SC
1664N NP	Tablet 2 mg	56	2	..	14.04	15.11	Progynova	SC

OESTRIOL

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1771F NP	Pessaries 500 micrograms, 15	¥1	2	..	21.26	22.33	Ovestin Ovula	SH
1781R NP	Vaginal cream 1 mg per g (0.1%), 15 g	¥1	1	..	19.09	20.16	Ovestin	SH

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Progestogens								
<i>Pregnen (4) derivatives</i>								
MEDROXYPROGESTERONE ACETATE								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
2321E NP	Tablet 10 mg	30	2	..	15.48	16.55	^a Medroxyhexal	SZ
							^a Ralovera	KR
				^B 1.68	17.16	16.55	^a Provera	PF
2323G NP	Tablet 5 mg	56	2	..	14.86	15.93	^a Ralovera	KR
				^B 1.68	16.54	15.93	^a Provera	PF
MEDROXYPROGESTERONE ACETATE								
<u>Restricted benefit</u>								
Endometriosis.								
2722G	Tablet 10 mg	100	2	..	31.19	32.26	^a Ralovera	KR
				^B 1.57	32.76	32.26	^a Provera	PF
<i>Pregnadien derivatives</i>								
DYDROGESTERONE								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
1350C NP	Tablet 10 mg	28	2	..	16.62	17.69	Duphaston	SM
<i>Estren derivatives</i>								
NORETHISTERONE								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
2993M NP	Tablet 5 mg	30	2	..	32.48	33.55	Primolut N	SC
Progestogens and estrogens in combination								
<i>Progestogens and estrogens, combinations</i>								
OESTRADIOL with NORETHISTERONE ACETATE								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8427M NP	Transdermal patches 620 micrograms-2.7 mg (releasing approximately 50 micrograms-140 micrograms per 24 hours), 8	¥1	5	..	19.39	20.46	Estalis continuous 50/140	NV
8428N NP	Transdermal patches 510 micrograms-4.8 mg (releasing approximately 50 micrograms-250 micrograms per 24 hours), 8	¥1	5	..	19.39	20.46	Estalis continuous 50/250	NV

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	

Progestogens and estrogens, sequential preparations

OESTRADIOL and OESTRADIOL with DYDROGESTERONE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8244X NP	Pack containing 14 tablets oestradiol 2 mg and 14 tablets oestradiol 2 mg with dydrogesterone 10 mg	£1	5	..	18.76	19.83	Femoston 2/10	SM

OESTRADIOL and OESTRADIOL with NORETHISTERONE ACETATE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8425K NP	Pack containing 4 transdermal patches oestradiol 780 micrograms (releasing approximately 50 micrograms per 24 hours) and 4 transdermal patches oestradiol with norethisterone acetate 620 micrograms- 2.7 mg (releasing approximately 50 micrograms-140 micrograms per 24 hours)	£1	5	..	19.39	20.46	Estalis sequi 50/140	NV
8426L NP	Pack containing 4 transdermal patches oestradiol 780 micrograms (releasing approximately 50 micrograms per 24 hours) and 4 transdermal patches oestradiol with norethisterone acetate 510 micrograms- 4.8 mg (releasing approximately 50 micrograms-250 micrograms per 24 hours)	£1	5	..	19.39	20.46	Estalis sequi 50/250	NV

Gonadotropins and other ovulation stimulants

Gonadotropins

FOLLITROPIN ALFA

Restricted benefit

Anovulatory infertility.

Note

Except in cases of hypopituitarism or primary amenorrhoea, the patient should have been adequately treated with clomiphene citrate and/or gonadorelin and failed to have conceived.

Women who have had apparent ovulation induced by other agents and have failed to conceive should have laparoscopic evidence that there is no other impediment to conception.

Oligomenorrhoea should have been present for at least twelve months or amenorrhoea for at least six months prior to treatment.

Patients with hyperprolactinaemia should have had appropriate surgical or medical treatment prior to treatment.

Restricted benefit

For the treatment of infertility in males due to hypogonadotrophic hypogonadism, following failure of 6 months' treatment with human chorionic gonadotrophin to achieve adequate spermatogenesis. Combined treatment with HCG must be given.

8713N	Injection 300 i.u. in 0.5 mL multi-dose cartridge	3	5	..	*563.43	34.20	Gonal-f Pen	SG
8714P	Injection 450 i.u. in 0.75 mL multi-dose cartridge	3	5	..	*841.92	34.20	Gonal-f Pen	SG
8715Q	Injection 900 i.u. in 1.5 mL multi-dose cartridge	2	5	..	*1115.24	34.20	Gonal-f Pen	SG

FOLLITROPIN BETA

Restricted benefit

Anovulatory infertility.

Note

Except in cases of hypopituitarism or primary amenorrhoea, the patient should have been adequately treated with clomiphene citrate and/or gonadorelin and failed to have conceived.

Women who have had apparent ovulation induced by other agents and have failed to conceive should have laparoscopic evidence that there is no other impediment to conception.

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
Oligomenorrhoea should have been present for at least twelve months or amenorrhoea for at least six months prior to treatment.							
Patients with hyperprolactinaemia should have had appropriate surgical or medical treatment prior to treatment.							
<u>Restricted benefit</u>							
For the treatment of infertility in males due to hypogonadotrophic hypogonadism, following failure of 6 months' treatment with human chorionic gonadotrophin to achieve adequate spermatogenesis. Combined treatment with HCG must be given.							
8565T	Solution for injection 300 i.u. in 0.36 mL multi-dose cartridge	3	5	..	*563.43	34.20	Puregon 300 IU/0.36 mL SH
8566W	Solution for injection 600 i.u. in 0.72 mL multi-dose cartridge	2	5	..	*749.08	34.20	Puregon 600 IU/0.72 mL SH
8871X	Solution for injection 900 i.u. in 1.08 mL multi-dose cartridge	2	5	..	*1115.22	34.20	Puregon 900 IU/1.08 mL SH

HUMAN CHORIONIC GONADOTROPHIN

Restricted benefit

Anovulatory infertility.

Note

Except in cases of hypopituitarism or primary amenorrhoea, the patient should have been adequately treated with clomiphene citrate and/or gonadorelin and failed to have conceived.

Women who have had apparent ovulation induced by other agents and have failed to conceive should have laparoscopic evidence that there is no other impediment to conception.

Oligomenorrhoea should have been present for at least twelve months or amenorrhoea for at least six months prior to treatment.

Patients with hyperprolactinaemia should have had appropriate surgical or medical treatment prior to treatment.

Restricted benefit

For the treatment of infertility in males due to hypogonadotrophic hypogonadism;

For the treatment of infertility in males associated with isolated luteinising hormone deficiency;

For the treatment of males who have combined deficiency of human growth hormone and gonadotrophins and in whom the absence of secondary sexual characteristics indicates a lag in maturation.

Restricted benefit

For the treatment of boys over the age of 16 years who show clinical evidence of hypogonadism or delayed puberty. Treatment must not extend beyond 6 months.

1581F	Injection set containing 3 ampoules powder for injection 1,500 units and 3 ampoules solvent 1 mL	1	5	..	53.47	34.20	Pregnyl	SH

Ovulation stimulants, synthetic

CLOMIPHENE CITRATE

Note

Care must be taken to comply with the provisions of State/Territory law when prescribing clomiphene citrate.

Restricted benefit

Anovulatory infertility;

Patients undergoing in-vitro fertilisation.

1211R	Tablet 50 mg	10	5	..	36.00	34.20	^a Clomid ^a Serophene	SW SG

Antiandrogens

Antiandrogens, plain preparations

CYPROTERONE ACETATE

Authority required (STREAMLINED)

1230

Moderate to severe androgenisation in non-pregnant women (acne alone is not a sufficient indication of androgenisation).

Caution

This drug should not be used during pregnancy as it may result in feminisation of the male foetus.

1269T	Tablet 50 mg	20	5	..	52.74	34.20	^a Cyprohexal ^a Cyprone	SZ AF

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
							^a Cyprostat SY
							^a GenRx Cyproterone Acetate GX
							^a Procur GM
				^B 3.13	55.87	34.20	^a Androcur SC
CYPROTERONE ACETATE							
<u>Authority required (STREAMLINED)</u>							
1014							
Advanced carcinoma of the prostate;							
1404							
To reduce drive in sexual deviations in males.							
1270W	Tablet 50 mg	100	5	..	*207.72	34.20	^a Cyprohexal SZ
							^a Cyprone AF
							^a Cyprostat SY
							^a GenRx Cyproterone Acetate GX
							^a Procur GM
				^B 3.00	*210.72	34.20	^a Androcur SC
8019C	Tablet 100 mg	50	5	..	169.76	34.20	^a Cyprohexal SZ
							^a Cyprostat-100 SY
							^a GenRx Cyproterone Acetate GX
							^a Procur 100 GM
				^B 1.65	171.41	34.20	^a Androcur-100 SC

Other sex hormones and modulators of the genital system

Antigonadotropins and similar agents

DANAZOL

Caution

Pregnancy must be excluded prior to administration of this drug.

Authority required (STREAMLINED)

1090

Endometriosis, visually proven;

1151

Hereditary angio-oedema;

2639

Short-term treatment (up to 6 months) of intractable primary menorrhagia (Treatment of this indication is limited to 6 months. See Australian Product Information);

2640

Short-term treatment (up to 6 months) of severe benign (fibrocystic) breast disease or mastalgia associated with severe symptomatic benign breast disease in patients refractory to other treatments (Treatment of this indication is limited to 6 months. See Australian Product Information).

1285P	Capsule 100 mg	100	5	..	58.58	34.20	Azol 100	AF
1287R	Capsule 200 mg	100	5	..	86.97	34.20	Azol 200	AF

GESTRINONE

Authority required (STREAMLINED)

3652

Short-term treatment (up to 6 months) of visually proven endometriosis (only 1 course of not more than 6 months' therapy may be prescribed).

8015W	Capsule 2.5 mg	8	5	..	81.81	34.20	Dimetriose	SW

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Selective estrogen receptor modulators							
RALOXIFENE HYDROCHLORIDE							
<u>Authority required (STREAMLINED)</u>							
2647							
Treatment as the sole PBS-subsidised anti-resorptive agent for established post-menopausal osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.							
A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.							
<u>Note</u>							
Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.							
8363E NP	Tablet 60 mg	28	5	..	57.87	34.20	Evista LY

Urologicals

Other urologicals, incl. antispasmodics

Urinary antispasmodics

OXYBUTYNIN

Restricted benefit

Detrusor overactivity in a patient who cannot tolerate oral oxybutynin, or who cannot swallow oral oxybutynin.

9454N NP	Transdermal patches 36 mg (releasing approximately 3.9 mg per 24 hours), 8	£1	5	..	35.81	34.20	Oxytrol HH

OXYBUTYNIN HYDROCHLORIDE

Restricted benefit

Detrusor overactivity.

8039D NP	Tablet 5 mg	100	5	..	15.59	16.66	^a Ditropan SW
						^a Oxybutynin Sandoz	SZ
						^a Oxybutynin Winthrop	WA

PROPANTHELINE BROMIDE

Restricted benefit

Detrusor overactivity.

1953T NP	Tablet 15 mg	200	5	..	*26.46	27.53	Pro-Banthine SI

Other urologicals

PHENOXYBENZAMINE HYDROCHLORIDE

Restricted benefit

Phaeochromocytoma;

Neurogenic urinary retention.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1166J NP	Capsules 10 mg, 30	3	5	..	*204.90	34.20	Dibenylene GH
1862B NP	Capsule 10 mg	100	5	..	67.36	34.20	Dibenylene GH
9286R NP	Capsules 10 mg, 100	£1	5	..	1164.47	34.20	Dibenzylene GH

Genito urinary system and sex hormones

					Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	\$	\$	Brand Name and Manufacturer	
SODIUM BICARBONATE								
9470K NP	Capsule 840 mg	100	2	..	14.00	15.07	Sodibic	AS

Systemic hormonal preparations, excl. sex hormones and insulins

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	

Systemic hormonal preparations, excl. sex hormones and insulins

Pituitary and hypothalamic hormones and analogues

Anterior pituitary lobe hormones and analogues

ACTH

TETRACOSACTRIN							
2832C	Injection 1 mg in 1 mL	5	5	..	*71.27	34.20	Synacthen Depot 1 mg/1 mL NV

Thyrotropin

THYROTROPIN ALFA

Authority required (STREAMLINED)

3193

Ablation of thyroid remnant tissue, in combination with radioactive iodine, in a post thyroidectomy patient without known metastatic disease.

2700D	Powder for injection 0.9 mg, 2	1	..	..	1901.42	34.20	Thyrogen GZ

Posterior pituitary lobe hormones

Vasopressin and analogues

DESMOPRESSIN ACETATE

Authority required (STREAMLINED)

1678

Cranial diabetes insipidus.

2129C	Intranasal solution 100 micrograms per mL, 2.5 mL	5	5	..	*161.17	34.20	Minirin FP
8662X	Tablet 200 micrograms	90	5	..	*179.91	34.20	Minirin FP
8711L	Nasal spray (pump pack) 10 micrograms per actuation, 60 actuations, 6 mL	2	5	..	*161.04	34.20	Minirin Nasal Spray FP

DESMOPRESSIN ACETATE

Authority required (STREAMLINED)

2641

Primary nocturnal enuresis in patients aged 6 years or older who are refractory to an enuresis alarm;

2642

Primary nocturnal enuresis in patients aged 6 years or older for whom an enuresis alarm is contraindicated. The reason that an alarm is contraindicated must be documented in the patient's medical records when treatment is initiated.

Note

Not to be used in preference to enuresis alarms.

Desmopressin nasal spray may be associated with an increased risk of hyponatraemia compared to the oral formulations.

Note

Only one application per six months with no more than twice the maximum quantity will be authorised for the tablets.

8663Y NP	Tablet 200 micrograms	30	5	..	64.25	34.20	Minirin FP

DESMOPRESSIN ACETATE

Authority required (STREAMLINED)

2641

Primary nocturnal enuresis in patients aged 6 years or older who are refractory to an enuresis alarm;

2642

Primary nocturnal enuresis in patients aged 6 years or older for whom an enuresis alarm is contraindicated. The reason that an alarm is contraindicated must be documented in the patient's medical records when treatment is initiated.

Systemic hormonal preparations, excl. sex hormones and insulins

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
<u>Note</u> Not to be used in preference to enuresis alarms. Desmopressin nasal spray may be associated with an increased risk of hyponatraemia compared to the oral formulations.							
<u>Note</u> Only one application per 6 months with no more than twice the maximum quantity will be authorised for the wafers.							
9398P NP	Wafer 120 micrograms (base)	30	5	..	70.85	34.20	Minirin Melt FP

DESMOPRESSIN ACETATE

Authority required (STREAMLINED)

2641

Primary nocturnal enuresis in patients aged 6 years or older who are refractory to an enuresis alarm;

2642

Primary nocturnal enuresis in patients aged 6 years or older for whom an enuresis alarm is contraindicated. The reason that an alarm is contraindicated must be documented in the patient's medical records when treatment is initiated.

Note

Not to be used in preference to enuresis alarms.

Desmopressin nasal spray may be associated with an increased risk of hyponatraemia compared to the oral formulations.

8712M NP	Nasal spray (pump pack) 10 micrograms per actuation, 60 actuations, 6 mL	£1	5	..	83.73	34.20	Minirin Nasal Spray FP

Hypothalamic hormones

Gonadotropin-releasing hormones

NAFARELIN ACETATE

Authority required

Initial treatment (up to 6 months) of visually proven endometriosis;

Subsequent treatment (up to 6 months) of visually proven endometriosis, where 2 years or more have elapsed since the end of the previous course and where a recent bone density assessment has been made. The date of the assessment must be provided.

2962X	Nasal spray (pump pack) 200 micrograms (base) per dose (60 doses)	£1	5	..	95.51	34.20	Synarel PF

Corticosteroids for systemic use

Corticosteroids for systemic use, plain

Mineralocorticoids

FLUDROCORTISONE ACETATE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1433K NP	Tablet 100 micrograms	200	1	..	*46.50	34.20	Florinef SI

Glucocorticoids

BETAMETHASONE ACETATE with BETAMETHASONE SODIUM PHOSPHATE

Restricted benefit

Alopecia areata;

For local intra-articular or peri-articular infiltration;

Granulomata, dermal;

Keloid;

Lichen planus hypertrophic;

Lichen simplex chronicus;

Lupus erythematosus, chronic discoid;

Necrobiosis lipoidica;

Uveitis.

Systemic hormonal preparations, excl. sex hormones and insulins

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Note								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
2694T NP	Injection 3 mg-3.9 mg (equivalent to 5.7 mg betamethasone) in 1 mL	5	..	..	25.38	26.45	Celestone Chronodose	SH
CORTISONE ACETATE								
Note								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
1246N NP	Tablet 5 mg	50	4	..	15.30	16.37	Cortate	AS
1247P NP	Tablet 25 mg	60	4	..	17.74	18.81	Cortate	AS
DEXAMETHASONE								
Note								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
1292B NP	Tablet 500 micrograms	30	4	..	8.84	9.91	Dexamethsone	AS
2507Y NP	Tablet 4 mg	30	4	..	12.40	13.47	Dexamethsone	AS
DEXAMETHASONE SODIUM PHOSPHATE								
Note								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
1291Y NP	Injection equivalent to 8 mg dexamethasone phosphate in 2 mL	5	1	..	27.58	28.65	Hospira Pty Limited	HH
2509C NP	Injection equivalent to 4 mg dexamethasone phosphate in 1 mL	5	..	..	18.08	19.15	Hospira Pty Limited	HH
HYDROCORTISONE								
Note								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
1499X NP	Tablet 4 mg	50	4	..	11.01	12.08	Hysone 4	AF
1500Y NP	Tablet 20 mg	60	4	..	14.49	15.56	Hysone 20	AF
HYDROCORTISONE SODIUM SUCCINATE								
1501B NP	Injection equivalent to 100 mg hydrocortisone with 2 mL solvent	2	..	..	*16.72	17.79	Solu-Cortef	PF
3096Y NP	Injection equivalent to 250 mg hydrocortisone with 2 mL solvent	1	..	..	15.72	16.79	Solu-Cortef	PF
HYDROCORTISONE SODIUM SUCCINATE								
Restricted benefit								
For use in a hospital.								
1510L NP	Injection equivalent to 100 mg hydrocortisone with 2 mL solvent	6	..	..	*37.32	34.20	Solu-Cortef	PF
1511M NP	Injection equivalent to 250 mg hydrocortisone with 2 mL solvent	6	..	..	*59.82	34.20	Solu-Cortef	PF

Systemic hormonal preparations, excl. sex hormones and insulins

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
METHYLPREDNISOLONE ACETATE								
<u>Restricted benefit</u>								
For local intra-articular or peri-articular infiltration.								
1928L NP	Injection 40 mg in 1 mL	5	..	..	24.60	25.67 ^a	Depo-Nisolone	KR
				^B 0.73	25.33	25.67 ^a	Depo-Medrol	PF
METHYLPREDNISOLONE SODIUM SUCCINATE								
2981X NP	Powder for injection equivalent to 40 mg methylprednisolone with diluent	5	..	..	35.63	34.20	Solu-Medrol	PF
8834Y NP	Powder for injection equivalent to 1 g methylprednisolone with diluent	1	..	..	95.43	34.20	Solu-Medrol	PF
PREDNISOLONE								
1916W NP	Tablet 25 mg	30	4	..	10.33	11.40	Panafcortelone	AS
							Solone	FM
1917X NP	Tablet 5 mg	60	4	..	8.58	9.65	Panafcortelone	AS
							Solone	FM
3152X NP	Tablet 1 mg	100	4	..	8.43	9.50 ^a	Predsolone	LN
				^B 0.46	8.89	9.50 ^a	Panafcortelone	AS
PREDNISOLONE SODIUM PHOSPHATE								
8285C NP	Oral solution equivalent to 5 mg prednisolone per mL, 30 mL	‡1	5	..	15.14	16.21 ^a	PredMix	LN
				^B 1.86	17.00	16.21 ^a	Redipred	AS
PREDNISONE								
1934T NP	Tablet 1 mg	100	4	..	8.90	9.97 ^a	Predsone	LN
				^B 0.63	9.53	9.97 ^a	Panafcort	AS
1935W NP	Tablet 5 mg	60	4	..	9.24	10.31	Panafcort	AS
							Sone	FM
1936X NP	Tablet 25 mg	30	4	..	11.51	12.58	Panafcort	AS
							Sone	FM
TRIAMCINOLONE ACETONIDE								
<u>Restricted benefit</u>								
Alopecia areata;								
For local intra-articular or peri-articular infiltration;								
Granulomata, dermal;								
Keloid;								
Lichen planus hypertrophic;								
Lichen simplex chronicus;								
Lupus erythematosus, chronic discoid;								
Necrobiosis lipoidica;								
Psoriasis.								
<u>Note</u>								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
2990J NP	Injection 10 mg in 1 mL	5	..	..	25.38	26.45	Kenacort-A10	SI

Systemic hormonal preparations, excl. sex hormones and insulins

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
Thyroid therapy							

Thyroid preparations

Thyroid hormones

LIOTHYRONINE SODIUM

Authority required (STREAMLINED)

1219

Management of patients with thyroid cancer;

1858

Replacement therapy for hypothyroid patients who have documented intolerance to thyroxine sodium;

1859

Replacement therapy for hypothyroid patients who have documented resistance to thyroxine sodium;

1182

Initiation of thyroid therapy in severely hypothyroid patients.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2318B NP	Tablet 20 micrograms	100	2	..	83.53	34.20	Tertroxin	SI

THYROXINE SODIUM

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2173J NP	Tablet equivalent to 200 micrograms anhydrous thyroxine sodium	200	1	..	27.43	28.50 ^a	Eutroxsig	FM
				^B 2.25	29.68	28.50 ^a	Oroxine	SI
2174K NP	Tablet equivalent to 50 micrograms anhydrous thyroxine sodium	200	1	..	23.72	24.79 ^a	Eutroxsig	FM
				^B 2.25	25.97	24.79 ^a	Oroxine	SI
2175L NP	Tablet equivalent to 100 micrograms anhydrous thyroxine sodium	200	1	..	24.34	25.41 ^a	Eutroxsig	FM
				^B 2.25	26.59	25.41 ^a	Oroxine	SI
9287T NP	Tablet equivalent to 75 micrograms anhydrous thyroxine sodium	200	1	..	24.37	25.44 ^a	Eutroxsig	FM
				^B 2.32	26.69	25.44 ^a	Oroxine	SI

Antithyroid preparations

Thiouracils

PROPYLTHIOURACIL

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1955X NP	Tablet 50 mg	200	2	..	*49.64	34.20	PTU	PL

Sulfur-containing imidazole derivatives

CARBIMAZOLE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1153Q NP	Tablet 5 mg	200	2	..	*31.04	32.11	Neo-Mercazole	LM

Systemic hormonal preparations, excl. sex hormones and insulins

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
Pancreatic hormones							
Glycogenolytic hormones							
Glycogenolytic hormones							
GLUCAGON HYDROCHLORIDE							
1449G NP	Injection set containing 1 mg (1 i.u.) and 1 mL solvent in disposable syringe	1	1	..	45.63	34.20	GlucaGen Hypokit NO

Calcium homeostasis

Parathyroid hormones and analogues *Parathyroid hormones and analogues*

TERIPARATIDE

Note

Any queries concerning the arrangements to prescribe teriparatide may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe teriparatide should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial treatment, as the sole PBS-subsidised agent, by a specialist or consultant physician, for severe, established osteoporosis in a patient with a very high risk of fracture who:

- (a) has a bone mineral density (BMD) T-score of -3.0 or less; and
- (b) has had 2 or more fractures due to minimal trauma; and
- (c) has experienced at least 1 symptomatic new fracture after at least 12 months continuous therapy with an anti-resorptive agent at adequate doses.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

If treatment with anti-resorptive therapy is contraindicated according to the relevant TGA-approved Product Information, details of the contraindication must be provided at the time of application.

If an intolerance of a severity necessitating permanent treatment withdrawal develops during the relevant period of use of one anti-resorptive agent, alternate anti-resorptive agents must be trialled so that the patient achieves the minimum requirement of 12 months continuous therapy. Details of accepted toxicities including severity can be found on the Medicare Australia website at www.medicareaustralia.gov.au and must be provided at the time of application.

Anti-resorptive therapies for osteoporosis and their adequate doses which will be accepted for the purposes of administering this restriction are alendronate sodium 10 mg per day or 70 mg once weekly, risedronate sodium 5 mg per day or 35 mg once weekly or 150 mg once monthly, raloxifene hydrochloride 60 mg per day (women only), denosumab 60 mg once every 6 months, disodium etidronate 200 mg with calcium carbonate 1.25 g per day, strontium ranelate 2 g per day and zoledronic acid 5 mg per annum.

Authority applications must be made in writing and must include:

Details of prior anti-resorptive therapy, fracture history including the date(s), site(s), the symptoms associated with the fracture(s) which developed during the course of anti-resorptive therapy and the score of the qualifying BMD measurement.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Authority required

Initial treatment, as the sole PBS-subsidised agent, by a specialist or consultant physician, for severe, established osteoporosis in a patient with a very high risk of fracture who was receiving treatment with teriparatide prior to 1 May 2009.

The authority application must be made in writing and the commencement date of treatment and the number of doses the patient has received of teriparatide must be provided with the application. The patient is eligible to receive a maximum of 18 months therapy of combined PBS-subsidised and non-PBS-subsidised therapy.

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Systemic hormonal preparations, excl. sex hormones and insulins

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
<u>Note</u> No applications for increased maximum quantities and/or repeats will be authorised. <u>Authority required</u> Continuing treatment for severe established osteoporosis where the patient has previously been issued with an authority prescription for this drug. Teriparatide must only be used for a lifetime maximum of 18 months therapy (18 pens). Up to a maximum of 18 pens will be reimbursed through the PBS. Authority applications for continuing treatment may be made by telephone to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). <u>Note</u> No applications for increased maximum quantities and/or repeats will be authorised. <u>Note</u> Special Pricing Arrangements apply.							
9411H	Injection 250 micrograms per mL, 2.4 mL in multi-dose pre-filled pen	1	5	..	438.37	34.20	Forteo LY

Anti-parathyroid agents Calcitonin preparations

SALCATONIN

Note

The maximum quantities for salcatonin shown represent the number of individual ampoules and NOT multiples of the manufacturer's packs. The pack size for both strengths is five ampoules.

Authority required (STREAMLINED)

3256

Symptomatic Paget disease of bone;

1412

Treatment initiated in a hospital (in-patient or out-patient) of hypercalcaemia.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2995P NP	Injection 50 i.u. in 1 mL	30	5	..	*207.66	34.20	Miacalcic 50	NV
2997R NP	Injection 100 i.u. in 1 mL	15	5	..	*161.13	34.20	Miacalcic 100	NV

Other anti-parathyroid agents

CINACALCET HYDROCHLORIDE

Authority required

Maintenance therapy, following initiation and stabilisation of treatment with cinacalcet, of a patient with chronic kidney disease on dialysis who has after 6 months treatment:

(i) a decrease of at least 30% in iPTH concentrations; or

(ii) iPTH greater than 15 pmol per L and an (adjusted) serum calcium concentration of less than 2.6 mmol per L.

Note

During the titration phase, intact PTH should be monitored 4 weekly (measured at least 12 hours post dose) and dose titrated until an appropriate iPTH concentration is achieved. During the titration phase, approval will be limited to sufficient supply for 4 weeks treatment at a time, with doses between 30 and 180 mg per day according to the patient's response and tolerability.

During the maintenance phase, approval will be limited to provide sufficient quantity for 4 weeks treatment up to a maximum of 6 months supply for doses between 30 and 180 mg per day according to the patient's response and tolerability. Intact PTH should be monitored quarterly (measured at least 12 hours post dose) and dose adjusted as necessary to maintain an appropriate iPTH concentration.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

Special Pricing Arrangements apply.

9157Y NP	Tablet 30 mg (base)	28	5	..	343.60	34.20	Sensipar	AN
9158B	Tablet 60 mg (base)	28	5	..	670.32	34.20	Sensipar	AN

Systemic hormonal preparations, excl. sex hormones and insulins

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
NP							
9159C	Tablet 90 mg (base)	28	5	..	1002.27	34.20	Sensipar
NP							AN

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	

Antiinfectives for systemic use

Antibacterials for systemic use

Tetracyclines

Tetracyclines

DOXYCYCLINE

Note

Bioequivalence has been demonstrated between doxycycline tablet 100 mg (as hydrochloride) and doxycycline tablet 100 mg (as monohydrate).

2709N NP	Tablet 100 mg (as hydrochloride)	7	1	..	8.47	9.54	^a	Doxsig	SI
							^a	Doxy-100	GM
							^a	Doxylin 100	AF
9105F NP	Tablet 100 mg (as monohydrate)	7	1	..	8.47	9.54	^B 1.19	Vibramycin	PF
							^a	Chem mart	CH
							^a	Doxycycline	
							^a	Doxyhexal	SZ
							^a	GenRx Doxycycline	GX
							^a	Terry White Chemists Doxycycline	TW

DOXYCYCLINE

2708M NP	Capsule 100 mg (as hydrochloride)	7	1	..	8.47	9.54	^a	Mayne Pharma	YT
							^a	Doryx	YN

DOXYCYCLINE

Restricted benefit

Bronchiectasis in patients aged 8 years or older;

Chronic bronchitis in patients aged 8 years or older;

Severe acne.

Note

Bioequivalence has been demonstrated between doxycycline tablet 50 mg (as hydrochloride) and doxycycline tablet 50 mg (as monohydrate).

2711Q NP	Tablet 50 mg (as hydrochloride)	25	5	..	10.07	11.14	^a	Doxy-50	GM
							^a	Doxylin 50	AF
							^B 1.25	Vibra-Tabs	PF
9106G NP	Tablet 50 mg (as monohydrate)	25	5	..	10.07	11.14	^a	Chem mart	CH
							^a	Doxycycline	
							^a	Doxyhexal	SZ
							^a	Frakas	SI
							^a	GenRx Doxycycline	GX
							^a	Terry White Chemists Doxycycline	TW

DOXYCYCLINE

Restricted benefit

Bronchiectasis in patients aged 8 years or older;

Chronic bronchitis in patients aged 8 years or older;

Severe acne.

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer	
2707L NP	Capsule 50 mg (as hydrochloride)	25	5	..	10.07	11.14	^a	Mayne Pharma Doxycycline	YT
				^B 1.31	11.38	11.14	^a	Doryx	YN

DOXYCYCLINE

Restricted benefit

Pelvic inflammatory disease.

Note

Bioequivalence has been demonstrated between doxycycline tablet 100 mg (as hydrochloride) and doxycycline tablet 100 mg (as monohydrate).

2702F NP	Tablet 100 mg (as hydrochloride)	28	..	..	*14.62	15.69	^a	Doxsig	SI
							^a	Doxy-100	GM
							^a	Doxylin 100	AF
				^B 4.76	*19.38	15.69	^a	Vibramycin	PF
9107H NP	Tablet 100 mg (as monohydrate)	28	..	..	*14.62	15.69	^a	Chem mart Doxycycline	CH
							^a	Doxyhexal	SZ
							^a	GenRx Doxycycline	GX
							^a	Terry White Chemists Doxycycline	TW

DOXYCYCLINE

Restricted benefit

Pelvic inflammatory disease.

2703G NP	Capsule 100 mg (as hydrochloride)	28	..	..	*14.62	15.69	^a	Mayne Pharma Doxycycline	YT
				^B 4.60	*19.22	15.69	^a	Doryx	YN

DOXYCYCLINE

Restricted benefit

Urethritis.

Note

Bioequivalence has been demonstrated between doxycycline tablet 100 mg (as hydrochloride) and doxycycline tablet 100 mg (as monohydrate).

2714W NP	Tablet 100 mg (as hydrochloride)	21	..	..	*12.57	13.64	^a	Doxsig	SI
							^a	Doxy-100	GM
							^a	Doxylin 100	AF
				^B 3.57	*16.14	13.64	^a	Vibramycin	PF
9108J NP	Tablet 100 mg (as monohydrate)	21	..	..	12.56	13.63	^a	GenRx Doxycycline	GX
					*12.57	13.64	^a	Chem mart Doxycycline	CH
							^a	Doxyhexal	SZ
							^a	Terry White Chemists Doxycycline	TW

DOXYCYCLINE

Restricted benefit

Urethritis.

2715X NP	Capsule 100 mg (as hydrochloride)	21	..	..	12.52	13.59	^a	Mayne Pharma Doxycycline	YT
				^B 2.09	14.61	13.59	^a	Doryx	YN

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
MINOCYCLINE								
Caution								
There are concerns about the incidence of benign intracranial hypertension associated with this drug.								
Note								
No applications for increased maximum quantities and/or repeats will be authorised.								
3037W NP	Capsule 100 mg	11	..	..	9.65	10.72	Akamin 100	AF
MINOCYCLINE								
Caution								
There are concerns about the incidence of benign intracranial hypertension associated with this drug.								
Restricted benefit								
Severe acne not responding to other tetracyclines.								
Note								
No applications for increased maximum quantities and/or repeats will be authorised.								
1616C NP	Tablet 50 mg	60	5	..	15.49	16.56	^a Akamin 50	AF
				^B 2.00	17.49	16.56	^a Minomycin-50	SI
Beta-lactam antibacterials, penicillins								
Penicillins with extended spectrum								
AMOXYCILLIN								
1878W NP	Sachet containing oral powder 3 g	1	..	..	9.11	10.18	Amoxil	GK
1884E NP,MW	Capsule 250 mg	20	1	..	8.55	9.62	^a Alphamox 250	AF
							^a Amoxycillin-GA	GM
							^a Amoxycillin Ranbaxy	RA
							^a Amoxycillin Sandoz	SZ
							^a APO-Amoxycillin	TX
							^a Chem mart Amoxycillin	CH
							^a Cilamox	SI
							^a GenRx Amoxycillin	GX
							^a Terry White Chemists Amoxycillin	TW
				^B 0.79	9.34	9.62	^a Amoxil	GK
1886G NP	Powder for syrup 125 mg per 5 mL, 100 mL	‡1	1	..	#10.85	12.26	^a Alphamox 125	AF
							^a Amoxycillin Sandoz	SZ
							^a Bgramin	GM
							^a Chem mart Amoxycillin	CH
							^a GenRx Amoxycillin	GX
							^a Ranmoxy	RA
							^a Terry White Chemists Amoxycillin	TW
				^B 0.95	#11.80	12.26	^a Amoxil	GK
1887H NP	Powder for syrup 250 mg per 5 mL, 100 mL	‡1	1	..	#11.69	13.10	^a Alphamox 250	AF
							^a Amoxycillin Sandoz	SZ
							^a Bgramin	GM
							^a Chem mart Amoxycillin	CH
							^a Cilamox	SI

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
1889K NP,MW	Capsule 500 mg	20	1	B0.79 ..	#12.48 10.65	13.10 11.72	a GenRx Amoxycillin GX
							a Ranmoxy RA
							a Terry White Chemists Amoxycillin TW
							a Amoxil Forte GK
							a Alphamox 500 AF
							a Amoxycillin-GA GM
							a Amoxycillin Ranbaxy RA
							a Amoxycillin Sandoz SZ
							a APO-Amoxycillin TX
							a Chem mart CH
a Amoxycillin Cilamox SI							
a GenRx Amoxycillin GX							
a Terry White Chemists Amoxycillin TW							
8705E NP	Powder for oral suspension 500 mg per 5 mL, 100 mL	#1	1	B0.80 ..	11.45 #14.70	11.72 16.11	a Amoxil GK
							Maxamox SZ
AMOXYCILLIN							
<u>Restricted benefit</u>							
Acute exacerbations of chronic bronchitis.							
8581P NP	Tablet 1 g	14	1	.. B1.17	10.79 11.96	11.86 11.86	a Amoxycillin Sandoz BG
							a Maxamox SZ
AMPICILLIN							
2390T NP	Powder for injection 500 mg	5	1	..	11.08	12.15	a Austrapen LN
2977Q NP	Powder for injection 1 g	5	1	..	14.07	15.14	a Ibimicyn TS
							a Aspen Ampicyn AS
							a Austrapen LN
							a Ibimicyn TS
Beta-lactamase sensitive penicillins							
BENZATHINE BENZYLPENICILLIN							
2267H NP	Injection 900 mg in 2.3 mL single use pre-filled syringe	10	..	..	293.11	34.20	Bicillin L-A AS
BENZYLPENICILLIN							
1775K NP,MW	Powder for injection 600 mg	10	1	..	*42.92	34.20	BenPen CS
2647H NP	Powder for injection 3 g	10	..	..	*66.92	34.20	BenPen CS
PHENOXYMETHYLPENICILLIN							
1787C NP	Tablet 250 mg	50	..	..	*11.58	12.65	Abbecillin-VK Filmtab SI
1789E NP	Capsule 250 mg	50	..	..	11.41	12.48	a Cilicaïne VK FM
							a Cilopen VK GM
							LPV AS

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
2965C NP	Capsule 500 mg	50	..	..	13.84	14.91	^a Cilicaine VK	FM
							^a Cilopen VK	GM
							LPV	AS
3028J NP	Tablet 500 mg	50	..	..	*14.06	15.13	Abbocillin-VK	SI
							Filmstab	
9143F NP	Oral suspension 150 mg (as benzathine) per 5 mL, 100 mL	2	1	..	*22.40	23.47	^a Cilicaine V	FM
				^B 2.00	*24.40	23.47	^a Abbocillin-V	SI

PHENOXYMETHYLPENICILLIN

Restricted benefit

Prophylaxis of recurrent streptococcal infections (including rheumatic fever).

1703P NP	Tablet 250 mg	50	5	..	*11.58	12.65	Abbocillin-VK	SI
							Filmstab	
1705R NP	Capsule 250 mg	50	5	..	11.41	12.48	^a Cilicaine VK	FM
							^a Cilopen VK	GM
							LPV	AS

PROCAINE PENICILLIN

1794K NP	Injection 1.5 g	5	..	..	92.22	34.20	Cilicaine	SI

Beta-lactamase resistant penicillins

DICLOXACILLIN

Restricted benefit

Serious staphylococcal infections.

8121K NP,MW	Capsule 250 mg	24	..	..	11.45	12.52	^a Dicloxsig	SI
							^a Distaph 250	AF
8122L NP,MW	Capsule 500 mg	24	..	..	16.94	18.01	^a Diclocil	BQ
							^a Dicloxsig	SI
							^a Distaph 500	AF

FLUCLOXACILLIN

Caution

Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

1524F NP	Powder for injection 500 mg	5	..	..	15.49	16.56	^a Flubiclox	TS
							^a Flucil	AS
1525G NP	Powder for injection 1 g	5	1	..	20.66	21.73	^a Flubiclox	TS
							^a Flucil	AS
							^a Hospira Pty Limited	HH

FLUCLOXACILLIN

Caution

Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

Restricted benefit

Serious staphylococcal infections.

1526H NP,MW	Capsule 250 mg (as sodium)	24	..	..	11.45	12.52	^a Flopen	AS

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
1527J NP,MW	Capsule 500 mg (as sodium)	24	..	..	16.94	18.01	^a Staphylex 250	AF
							^a Flopen	AS
							^a Staphylex 500	AF
9149M NP	Powder for oral liquid 125 mg (as sodium) per 5 mL, 100 mL	‡1	..	..	#16.37	17.78	Aspen Pharmacare Australia Pty Limited	LN
9150N NP	Powder for oral liquid 250 mg (as sodium) per 5 mL, 100 mL	‡1	..	..	#20.08	21.49	Aspen Pharmacare Australia Pty Limited	LN

Combinations of penicillins, incl. beta-lactamase inhibitors

AMOXYCILLIN with CLAVULANIC ACID

Caution

Hepatotoxicity has been reported with this drug.

Restricted benefit

Infections where resistance to amoxicillin is suspected;

Infections where resistance to amoxicillin is proven.

1891M NP,MW	Tablet 500 mg-125 mg	10	1	..	12.16	13.23	^a Amoxicillin/ Clavulanic Acid 500/125 generichealth	GQ
							^a APO-Amoxicillin/ Clavulanic Acid 500/125	TX
							^a Clamoxyl Duo	AL
							^a Curam Duo 500/125	SZ
							^a GA-Amclav 500/125	GM
							^a Moxiclav Duo 500/125	SI
1892N NP	Powder for syrup 125 mg-31.25 mg per 5 mL, 75 mL	#1	1	^B 1.55	13.71	13.23	^a Augmentin Duo	GK
				..	#12.48	13.89	^a Clamoxyl	AL
							^a Curam	SZ
8254K NP	Tablet 875 mg-125 mg	10	1	^B 1.51	#13.99	13.89	^a Augmentin	GK
				..	14.60	15.67	^a Amoxicillin/ Clavulanic Acid 875/125 generichealth	GQ
							^a Chem mart Amoxicillin and Clavulanic Acid	CH
							^a Clamoxyl Duo forte	AL
							^a Clavycillin 875/125	CR
							^a Curam Duo Forte 875/125	SZ
							^a GA-Amclav Forte 875/125	GM
							^a GenRx Amoxicillin and Clavulanic Acid	GX
							^a Moxiclav Duo Forte 875/125	SI
							^a Terry White Chemists Amoxicillin and Clavulanic Acid	TW
8319W NP	Powder for syrup 400 mg-57 mg per 5 mL, 60 mL	#1	1	^B 2.04	16.64	15.67	^a Augmentin Duo forte	GK
				..	#13.99	15.40	^a Clamoxyl Duo 400	AL
							^a Curam Duo	SZ

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
				^B 1.54	#15.53	15.40 ^a	Augmentin Duo 400 GK

TICARCILLIN with CLAVULANIC ACID

Restricted benefit

Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent;

Septicaemia, suspected;

Septicaemia, proven.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2179Q NP	Powder for injection 3 g-100 mg (solvent required) (code 6884H applies to above item with approved solvent)	10	..	..	163.32	34.20	Timentin GK

Other beta-lactam antibacterials

First-generation cephalosporins

CEFALOTIN

2964B NP	Powder for injection 1 g	10	1	..	26.25	27.32 ^a	Cefalotin Sandoz SZ
						^a	Hospira Pty Limited HH
						^a	Keflin Neutral AS

CEPHALEXIN

3058Y NP,MW	Capsule 250 mg	20	1	..	8.85	9.92 ^a	Cefalexin Sandoz SZ
						^a	Cephabeil BF
						^a	Cephalexin GQ
						^a	generichealth MI
						^a	Cephitrust 250 CH
						^a	Chem mart CH
						^a	Cephalexin GM
						^a	Cilex GM
						^a	GenRx Cephalexin GX
						^a	Ialex LN
						^a	Ibilex 250 AF
						^a	Rancef RA
						^a	Terry White Chemists TW
						^a	Cephalexin AS
				^B 3.30	12.15	9.92 ^a	Keflex AS
3094W NP	Granules for syrup 125 mg per 5 mL, 100 mL	£1	1	..	#11.84	13.25 ^a	APO-Cephalexin TX
						^a	Cefalexin Sandoz SZ
						^a	Chem mart CH
						^a	Cephalexin GM
						^a	Cilex GM
						^a	GenRx Cephalexin GX
						^a	Ialex LN
						^a	Ibilex 125 AF
						^a	Terry White Chemists TW
						^a	Cephalexin AS
				^B 3.55	#15.39	13.25 ^a	Keflex AS
3095X NP	Granules for syrup 250 mg per 5 mL, 100 mL	£1	1	..	#13.24	14.65 ^a	APO-Cephalexin TX
						^a	Cefalexin Sandoz SZ

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
3119E NP,MW	Capsule 500 mg	20	1	..	10.77	11.84	^a Chem mart Cephalexin CH
							^a Cilex GM
							^a GenRx Cephalexin GX
							^a Ialex LN
							^a Ibilex 250 AF
							^a Terry White Chemists Cephalexin TW
							^a Keflex AS
							^a Cefalexin Sandoz SZ
							^a Cephabell BF
							^a Cephalexin generichealth GQ
							^a Cephatrust 500 MI
							^a Chem mart Cephalexin CH
							^a Cilex GM
							^a GenRx Cephalexin GX
							^a Ialex LN
							^a Ibilex 500 AF
							^a Rancef RA
							^a Terry White Chemists Cephalexin TW
							^a Keflex AS

CEPHAZOLIN

Restricted benefit

Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent;

Septicaemia, suspected;

Septicaemia, proven;

Cellulitis.

1256D NP	Powder for injection 500 mg	10	..	..	*39.88	34.20	Hospira Pty Limited	HH
1257E NP	Powder for injection 1 g	10	..	..	*56.92	34.20	^a Hospira Pty Limited	HH
					56.93	34.20	^a Cefazolin Sandoz	SZ
							^a Cephalozin Alphapharm	AF
							^a Kefzol	AS
9326W NP	Powder for injection 2 g	10	..	..	*104.22	34.20	^a Cefazolin Sandoz	SZ
							^a Cephalozin Alphapharm	AF

Second-generation cephalosporins

CEFACLOR

Caution

Serum sickness-like reactions have been reported with this drug, especially in children.

1169M	Tablet 375 mg (sustained release)	10	1	..	12.89	13.96	^a Cefaclor-GA	GN
							^a Chem mart Cefaclor CD	CH
							^a Douglas Cefaclor- CD	GM
							^a GenRx Cefaclor CD	GX
							^a Karlor CD	LN
							^a Keflor CD	AF

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer				
2460L	Powder for oral suspension 125 mg per 5 mL, 100 mL	#1	1	..	#13.50	14.91	^a Ozcef	RA			
							^a Terry White Chemists Cefaclor CD	TW			
							^a Ceclor CD	AS			
							^a Aclor 125	SI			
							^a Cefaclor Sandoz	SZ			
							^a Chem mart Cefaclor	CH			
							^a GenRx Cefaclor	GX			
							^a Keflor	AF			
							^a Ozcef	RA			
							^a Terry White Chemists Cefaclor	TW			
2461M	Powder for oral suspension 250 mg per 5 mL, 75 mL	#1	1	..	#13.83	15.24	^a Ceclor	AS			
							^a Aclor 250	SI			
							^a Cefaclor Sandoz	SZ			
							^a Chem mart Cefaclor	CH			
							^a GenRx Cefaclor	GX			
							^a Keflor	AF			
							^a Ozcef	RA			
							^a Terry White Chemists Cefaclor	TW			
							^B 4.37	#18.20	15.24	^a Ceclor	AS
							CEFUROXIME AXETIL				
8292K	Tablet 250 mg (base)	14	1	..	18.62	19.69	Zinnat	GK			

Third-generation cephalosporins

CEFOTAXIME

Restricted benefit

Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent;

Septicaemia, suspected;

Septicaemia, proven.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1085D NP	Powder for injection 1 g	10	..	..	*26.82	27.89	^a Cefotaxime Sandoz	SZ
					26.86	27.93	^a Hospira Pty Limited	HH
1086E NP	Powder for injection 2 g	10	..	..	*43.62	34.20	^a Cefotaxime Sandoz	SZ
					43.68	34.20	^a Hospira Pty Limited	HH

CEFTRIAZONE

Restricted benefit

Gonorrhoea.

9058R NP	Powder for injection 500 mg	1	..	..	10.33	11.40	Ceftriazone ICP	PP

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
CEFTRIAZONE							
<u>Restricted benefit</u>							
Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent;							
Septicaemia, suspected;							
Septicaemia, proven.							
<u>Note</u>							
Shared Care Model:							
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
1783W NP	Powder for injection 500 mg	5	..	..	*25.97	27.04	Ceftriaxone ICP PP
1784X NP	Powder for injection 1 g	5	..	..	36.96	34.20	^a Max Pharma Pty Ltd XF
					*36.97	34.20	^a Ceftriaxone ICP PP
						^a Ceftriaxone Sandoz SZ	
						^a DBL Ceftriaxone HH	
						^a Rocephin RO	
1785Y NP	Powder for injection 2 g	5	..	..	*60.62	34.20	^a Ceftriaxone ICP PP
						^a Ceftriaxone Sandoz SZ	
						^a DBL Ceftriaxone HH	
						^a Rocephin RO	

Fourth-generation cephalosporins

CEFEPIME

Authority required

Treatment of febrile neutropenia.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8315P NP	Powder for injection 1 g (as hydrochloride) (solvent required) (code 7079N applies to above item with approved solvent)	10	..	..	*161.62	34.20	^a Maxipime	BQ
							^a Omegapharm Pty Ltd	OE
8316Q NP	Powder for injection 2 g (as hydrochloride) (solvent required) (code 7085X applies to above item with approved solvent)	10	..	..	*293.22	34.20	^a Maxipime	BQ
							^a Omegapharm Pty Ltd	OE

Sulfonamides and trimethoprim

Trimethoprim and derivatives

TRIMETHOPRIM

2922T NP	Tablet 300 mg	7	1	..	8.48	9.55	^a Alprim	AF
				^B 2.00	10.48	9.55	^a Triprim	SI

Combinations of sulfonamides and trimethoprim, incl. derivatives

TRIMETHOPRIM with SULFAMETHOXAZOLE

Caution

There is an increased risk of severe adverse reactions with this combination in the elderly.

2949F NP	Tablet 80 mg-400 mg	10	1	..	8.67	9.74	Resprim	AF
2951H NP	Tablet 160 mg-800 mg	10	1	..	9.39	10.46	^a Bactrim DS	RO
							^a Resprim Forte	AF

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
3103H NP	Oral suspension 40 mg-200 mg per 5 mL, 100 mL	1	1	^B 1.54	10.93	10.46	^a Septrin Forte	SI
				..	9.05	10.12	Bactrim	RO
				^B 1.90	10.95	10.12	Septin	SI

Macrolides, lincosamides and streptogramins

Macrolides

AZITHROMYCIN

Restricted benefit

Uncomplicated urethritis due to Chlamydia trachomatis;

Uncomplicated cervicitis due to Chlamydia trachomatis.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8200N NP	Tablet 500 mg	2	..	..	21.39	22.46	^a Azithromycin	SZ
						^a	Sandoz Zithromax	PF

AZITHROMYCIN

Restricted benefit

Trachoma.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8201P NP	Powder for oral suspension 200 mg per 5 mL, 15 mL	1	..	..	#21.35	22.76		Zithromax	PF
8336R NP	Tablet 500 mg	2	2	..	21.39	22.46	^a Azithromycin	SZ	
						^a	Sandoz Zithromax	PF	
8318T NP	Tablet 250 mg	14	1	..	12.68	13.75	^a APO- Clarithromycin	TX	
						^a	Chem mart	CH	
						^a	Clarithromycin		
						^a	Clarac	GM	
						^a	Clarihexal	SZ	
						^a	Clarithro 250	SI	
						^a	GenRx	GX	
						^a	Clarithromycin		
						^a	Kalixocin	AF	
						^a	Terry White Chemists	TW	
				^B 1.96	14.64	13.75	^a Klarid	AB	

CLARITHROMYCIN

Restricted benefit

Bordetella pertussis;

Atypical mycobacterial infections.

9192T NP	Powder for oral liquid 250 mg per 5 mL, 50 mL	1	..	..	#37.23	34.20		Klarid	AB
1404X NP	Capsule 250 mg	25	1	..	9.43	10.50	^a	Mayne Pharma Erythromycin	YT
				^B 1.35	10.78	10.50	^a	Eryc	YN

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
ERYTHROMYCIN ETHYL SUCCINATE								
2424N NP	Powder for oral liquid 200 mg (base) per 5 mL, 100 mL	‡1	1	..	#12.32	13.73	^a	E-Mycin 200 AF
				^B 2.86	#15.18	13.73	^a	E.E.S. 200 LM
2428T NP	Powder for oral liquid 400 mg (base) per 5 mL, 100 mL	‡1	1	..	#13.40	14.81	^a	E-Mycin 400 AF
				^B 2.89	#16.29	14.81	^a	E.E.S. Granules LM
2750R NP	Tablet 400 mg (base)	25	1	..	10.92	11.99	^a	E-Mycin AF
				^B 2.80	13.72	11.99	^a	E.E.S. 400 Filmtab LM
ERYTHROMYCIN LACTOBIONATE								
1397M NP	Powder for I.V. infusion 1 g (base)	5	..	..	*88.92	34.20		Erythrocin-I.V. LM
ROXITHROMYCIN								
1760P NP	Tablet 150 mg	10	1	..	11.76	12.83	^a	APO-Roxithromycin TX
							^a	Biaxsig AV
							^a	Chem mart CH
							^a	Roxithromycin
							^a	Roxar 150 SI
							^a	Roxide SZ
							^a	Roximycin AF
							^a	Roxithromycin-GA GM
							^a	Terry White Chemists TW
				^B 2.60	14.36	12.83	^a	Rulide SW
8016X NP	Tablet 300 mg	5	1	..	11.76	12.83	^a	APO-Roxithromycin TX
							^a	Biaxsig AV
							^a	Chem mart CH
							^a	Roxithromycin
							^a	Roxar 300 SI
							^a	Roxide SZ
							^a	Roximycin AF
							^a	Roxithromycin-GA GM
							^a	Terry White Chemists TW
				^B 2.60	14.36	12.83	^a	Rulide SW
8129W NP	Tablet for oral suspension 50 mg	10	1	..	12.89	13.96		Rulide D SW

Lincosamides

CLINDAMYCIN

Restricted benefit

Gram-positive coccal infections where these cannot be safely and effectively treated with a penicillin.

3138E NP,MW	Capsule 150 mg	24	..	..	20.02	21.09	^a	Cleocin KR
				^B 1.40	21.42	21.09	^a	Dalacin C PF
LINCOMYCIN								
2530E NP,MW	Injection 600 mg in 2 mL	5	..	..	33.74	34.20		Lincocin PF

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Aminoglycoside antibacterials							
<i>Other aminoglycosides</i>							
GENTAMICIN SULFATE							
2824P NP	Injection 80 mg (base) in 2 mL	10	1	..	*19.94	21.01 ^a	Hospira Pty Limited HH
						^a	Pfizer Australia Pty Ltd PF
TOBRAMYCIN SULFATE							
<u>Restricted benefit</u>							
Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent;							
Septicaemia, suspected;							
Septicaemia, proven.							
1356J NP	Injection 80 mg (base) in 2 mL	10	1	..	*65.02	34.20	Hospira Pty Limited HH
8872Y NP	Injection 80 mg (base) in 2 mL (without preservative)	10	1	..	*65.02	34.20	Pfizer Australia Pty Ltd PF
TOBRAMYCIN SULFATE							
<u>Restricted benefit</u>							
Systemic treatment of Pseudomonas aeruginosa infection in a patient with cystic fibrosis.							
9480Y NP	Injection 500 mg (base) in 5 mL (without preservative)	10	1	..	357.37	34.20	Tobra-Day PL
Quinolone antibacterials							
<i>Fluoroquinolones</i>							
CIPROFLOXACIN							
<u>Authority required</u>							
Respiratory tract infection proven or suspected to be caused by Pseudomonas aeruginosa in severely immunocompromised patients;							
Bacterial gastroenteritis in severely immunocompromised patients;							
Treatment of infections proven to be due to Pseudomonas aeruginosa or other gram-negative bacteria resistant to all other oral antimicrobials;							
Treatment of joint and bone infections, epididymo-orchitis, prostatitis or perichondritis of the pinna, suspected or proven to be caused by gram-negative bacteria or gram-positive bacteria resistant to all other appropriate antimicrobials;							
Gonorrhoea.							
1208N NP	Tablet 250 mg	14	..	..	26.33	27.40 ^a	C-Flox 250 AL
						^a	Cifran RA
						^a	Ciprofloxacin-DRLA RZ
						^a	Ciprofloxacin Sandoz SZ
						^a	Ciprol 250 SI
						^a	GenRx GX
						^a	Ciprofloxacin Profloxin HX
				^B 1.45	27.78	27.40 ^a	Ciproxin 250 BN
CIPROFLOXACIN							
<u>Authority required</u>							
Respiratory tract infection proven or suspected to be caused by Pseudomonas aeruginosa in severely immunocompromised patients;							
Bacterial gastroenteritis in severely immunocompromised patients;							
Treatment of infections proven to be due to Pseudomonas aeruginosa or other gram-negative bacteria resistant to all other oral antimicrobials;							
Treatment of joint and bone infections, epididymo-orchitis, prostatitis or perichondritis of the pinna, suspected or proven to be caused by gram-negative bacteria or gram-positive bacteria resistant to all other appropriate antimicrobials.							
1209P	Tablet 500 mg	14	..	..	44.75	34.20 ^a	Ascent GN

Other antibacterials

Glycopeptide antibacterials

Restricted benefit

Prophylaxis of endocarditis in patients hypersensitive to penicillin.

2269K	Powder for injection 1 g (1,000,000 i.u.) vancomycin activity	1	..	..	17.95	19.02	^a	Hospira Pty Limited	HH
							^a	Vancomycin Sandoz	SZ
3130R	Powder for injection 500 mg (500,000 i.u.) vancomycin activity	2	..	..	*17.96	19.03	^a	Hospira Pty Limited	HH

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
							^a Vancocin CP	AS
							^a Vancomycin Sandoz	SZ
VANCOMYCIN								
<u>Restricted benefit</u>								
Endophthalmitis;								
Use initiated in a hospital for infections where vancomycin is an appropriate antibiotic.								
2270L	Powder for injection 1 g (1,000,000 i.u.) vancomycin activity	3	..	..	*41.01	34.20	^a Hospira Pty Limited	HH
							^a Vancomycin Sandoz	SZ
3131T	Powder for injection 500 mg (500,000 i.u.) vancomycin activity	5	..	..	*35.27	34.20	^a Hospira Pty Limited	HH
							^a Vancocin CP	AS
							^a Vancomycin Sandoz	SZ

Steroid antibacterials

FUSIDIC ACID

Restricted benefit

For use in combination with another antibiotic in the treatment of proven serious staphylococcal infections.

2312Q	Tablet (sodium salt) 250 mg	36	1	..	90.89	34.20	Fucidin	CS

Imidazole derivatives

METRONIDAZOLE

1626N NP	Tablet 400 mg	5	2	..	7.88	8.95	Metrogyl 400	AF
1636D NP	Tablet 200 mg	21	1	..	7.96	9.03	^a Metrogyl 200	AF
							^a Metronide 200	AV
				^B 2.30	10.26	9.03	^a Flagyl	SW
1642K NP	Suppositories 500 mg, 10	‡1	..	..	23.16	24.23	Flagyl	SW

METRONIDAZOLE

Restricted benefit

Treatment of anaerobic infections.

1621H NP	Tablet 400 mg	21	1	..	10.03	11.10	^a Metrogyl 400	AF
							^a Metronide 400	AV
				^B 2.30	12.33	11.10	^a Flagyl	SW

METRONIDAZOLE

Restricted benefit

Prophylaxis in large bowel surgery;

Treatment, in a hospital, of acute anaerobic sepsis.

1638F NP	I.V. infusion 500 mg in 100 mL	5	1	..	*31.97	33.04	^a Baxter Healthcare Pty Ltd	BX
					*32.00	33.07	^a DBL Metronidazole Intravenous Infusion	HH
							^a Metronidazole Sandoz	SZ

Antiinfectives for systemic use

					Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	\$	\$	Brand Name and Manufacturer	
	METRONIDAZOLE BENZOATE							
1630T NP	Oral suspension 320 mg per 5 mL (equivalent to 200 mg metronidazole in 5 mL), 100 mL	1	..	..	18.82	19.89	Flagyl S	SW
	TINIDAZOLE							
1465D NP	Tablet 500 mg	4	..	..	10.88	11.95 ^a	Simplotan	GP
				^B 2.46	13.34	11.95 ^a	Fasigyn	PF

Nitrofurantoin derivatives

NITROFURANTOIN

Caution

Nitrofurantoin may cause peripheral neuritis and severe pulmonary reactions.

1692C NP, MW	Capsule 50 mg	30	1	..	20.38	21.45	Macrochantin	PF
1693D NP, MW	Capsule 100 mg	30	1	..	26.26	27.33	Macrochantin	PF

Other antibacterials

HEXAMINE HIPPURATE

3124K NP	Tablet 1 g	100	5	..	42.38	34.20	Hiprex	IA

Antimycotics for systemic use

Antimycotics for systemic use

Imidazole derivatives

KETOCONAZOLE

Authority required (STREAMLINED)

3606

Symptomatic genital candidiasis recurring after treatment of at least 2 episodes with topical therapy.

Caution

Hepatotoxicity has been reported with ketoconazole.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1573T NP	Tablet 200 mg	10	..	..	19.79	20.86	Nizoral	JC

KETOCONAZOLE

Authority required (STREAMLINED)

3604

Oral candidiasis in severely immunocompromised persons where topical therapy has failed;

3605

Systemic or deep mycoses where other forms of therapy have failed.

Caution

Hepatotoxicity has been reported with ketoconazole.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1572R NP	Tablet 200 mg	30	5	..	42.17	34.20	Nizoral	JC

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Triazole derivatives							
FLUCONAZOLE							
<u>Authority required (STREAMLINED)</u>							
3615 Treatment of cryptococcal meningitis;							
3616 Maintenance therapy in patients with cryptococcal meningitis and immunosuppression;							
3613 Treatment of oropharyngeal candidiasis in immunosuppressed patients;							
3614 Treatment of oesophageal candidiasis in immunosuppressed patients;							
3617 Prophylaxis of oropharyngeal candidiasis in immunosuppressed patients;							
3618 Treatment of serious and life-threatening candida infections.							
<u>Note</u>							
Shared Care Model:							
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
1471K NP	Capsule 50 mg	28	5	..	70.56	34.20	^a DBL Fluconazole HH
							^a Diflucan PF
							^a Dizole 50 AF
							^a Fluconazole Sandoz SZ
							^a Fluzole 50 SI
							^a Ozole RA
1472L NP	Capsule 100 mg	28	5	..	131.36	34.20	^a DBL Fluconazole HH
							^a Diflucan PF
							^a Dizole 100 AF
							^a Fluconazole Sandoz SZ
							^a Fluconazole Winthrop WA
							^a Ozole RA
1473M NP	Solution for I.V. infusion 100 mg in 50 mL	7	..	..	*159.37	34.20	^a Diflucan PF
							^a Fluconazole-Claris AE
							^a Fluconazole Hexal SZ
1474N NP	Solution for I.V. infusion 200 mg in 100 mL	7	..	..	*285.93	34.20	^a Baxter Healthcare Pty Ltd BX
							^a Diflucan PF
							^a Fluconazole-Claris AE
							^a Fluconazole Hexal HX
							^a Fluconazole Sandoz SZ
1475P NP	Capsule 200 mg	28	5	..	245.69	34.20	^a APO-Fluconazole TX
							^a DBL Fluconazole HH
							^a Diflucan PF
							^a Dizole 200 AF
							^a Fluconazole Sandoz SZ
							^a Fluzole 200 SI
							^a Ozole RA
1757L NP	Solution for I.V. infusion 400 mg in 200 mL	1	..	..	71.09	34.20	Baxter Healthcare Pty Ltd BX

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
For the treatment and maintenance therapy of other serious invasive mycosis.							
<u>Note</u>							
Shared Care Model:							
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
9363T NP	Tablet 50 mg	56	2	..	700.87	34.20	Vfend PF
9364W NP	Tablet 200 mg	56	2	..	2631.08	34.20	Vfend PF

VORICONAZOLE

Authority required

For the treatment and maintenance therapy of definite or probable invasive aspergillosis in immunocompromised patients;

For the treatment and maintenance therapy of serious fungal infections caused by *Scedosporium* species or *Fusarium* species;

For the treatment and maintenance therapy of serious *Candida* infections where:

(a) the causative species is not susceptible to fluconazole; or

(b) treatment with fluconazole has failed; or

(c) treatment with fluconazole is not tolerated;

For the treatment and maintenance therapy of other serious invasive mycosis.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

Application for an increased maximum quantity to allow for up to 1 month's treatment and repeats sufficient for up to 6 months' treatment may be authorised.

9452L NP	Powder for oral suspension 40 mg per mL, 70 mL	1	..	..	#703.40	34.20	Vfend PF

Antimycobacterials

Drugs for treatment of tuberculosis

Hydrazides

ISONIAZID

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1554T NP	Tablet 100 mg	100	2	..	11.86	12.93	Fawns and McAllan Proprietary Limited	FM

Drugs for treatment of lepra

Drugs for treatment of lepra

DAPSONE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1272Y NP	Tablet 100 mg	100	1	..	113.84	34.20	Link Medical Products Pty Ltd	LM
8801F NP	Tablet 25 mg	100	1	..	100.58	34.20	Link Medical Products Pty Ltd	LM

RIFAMPICIN

Authority required

Leprosy in adults.

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	
1982H NP	Note Shared Care Model: For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.	100	..	..	37.01	34.20	Rimycin 150 AF
	Capsule 150 mg						
	1983J NP	Capsule 300 mg	100	..	..	64.94	34.20

RIFAMPICIN

Restricted benefit

Prophylaxis of meningococcal disease in close contacts and carriers;

Prophylactic treatment of contacts of patients with Haemophilus influenzae type B.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1981G NP	Capsule 150 mg	10	..	..	11.96	13.03	Rimycin 150 AF
1984K NP	Capsule 300 mg	10	..	..	13.55	14.62	Rimycin 300 AF
8025J NP	Syrup 100 mg per 5 mL, 60 mL	1	..	..	28.57	29.64	Rifadin SW

Antivirals for systemic use

Direct acting antivirals

Nucleosides and nucleotides excl. reverse transcriptase inhibitors

ACICLOVIR

Authority required (STREAMLINED)

3632

Moderate to severe initial genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is desirable but need not delay treatment.

Note

Aciclovir 200 mg is not PBS-subsidised for chickenpox, herpes zoster or herpes simplex infections other than genital herpes.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

1003T NP	Tablet 200 mg	50	..	..	*69.54	34.20	^a Acihexal	SZ
							^a Acyclo-V 200	AF
							^a GenRx Aciclovir	GX
							^a Lovir	GM
					^B 4.34	*73.88	^a Zovirax 200 mg	GK

ACICLOVIR

Authority required (STREAMLINED)

3633

Episodic treatment or suppressive therapy of moderate to severe recurrent genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.

Note

Aciclovir 200 mg is not PBS-subsidised for chickenpox, herpes zoster or herpes simplex infections other than genital herpes.

1007B NP	Tablet 200 mg	90	5	..	121.90	34.20	^a Aciclovir 200	CR
							^a Acihexal	SZ
							^a Acyclo-V 200	AF
							^a Chem mart Aciclovir	CH

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer	
							^a	GenRx Aciclovir	GX
							^a	Lovir	GM
							^a	Ozvir	RA
							^a	Terry White Chemists Aciclovir	TW
				^B 3.21	125.11	34.20	^a	Zovirax 200 mg	GK
ACICLOVIR									
<u>Authority required (STREAMLINED)</u>									
3622									
Treatment of patients with herpes zoster within 72 hours of the onset of the rash;									
3631									
Herpes zoster ophthalmicus.									
<u>Note</u>									
Aciclovir is effective only if commenced within 72 hours of onset of rash.									
Aciclovir 800 mg is not PBS-subsidised for herpes simplex or chickenpox.									
<u>Note</u>									
No applications for repeats will be authorised.									
1052J NP	Tablet 800 mg	35	..	..	146.32	34.20	^a	Aciclovir 800	CR
							^a	Acihexal	SZ
							^a	Acyclo-V 800	AF
							^a	GenRx Aciclovir	GX
				^B 1.56	147.88	34.20	^a	Zovirax 800 mg	GK
ACICLOVIR									
<u>Authority required (STREAMLINED)</u>									
3630									
Patients with advanced HIV disease (CD4 cell counts of less than 150 million per litre).									
8234J NP	Tablet 800 mg	120	5	..	446.33	34.20	^a	Acihexal	SZ
							^a	Acyclo-V 800	AF
FAMCICLOVIR									
<u>Authority required (STREAMLINED)</u>									
3624									
Episodic treatment of moderate to severe recurrent genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.									
<u>Note</u>									
Famciclovir 250 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.									
2274Q NP	Tablet 250 mg	20	1	..	134.44	34.20	^a	APO-Famciclovir	TX
							^a	Ezovir	AF
							^a	Famciclovir Sandoz	SZ
							^a	Famvir	NV
							^a	Favic 250	SI
8092X NP	Tablet 125 mg	40	1	..	134.44	34.20	^a	APO-Famciclovir	TX
							^a	Ezovir	AF
							^a	Famvir	NV
							^a	Favic 125	SI

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
FAMCICLOVIR							
<u>Authority required (STREAMLINED)</u>							
3622							
Treatment of patients with herpes zoster within 72 hours of the onset of the rash.							
<u>Note</u>							
Famciclovir is effective only if commenced within 72 hours of onset of rash.							
Famciclovir 250 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.							
<u>Note</u>							
No applications for repeats will be authorised.							
8002E NP	Tablet 250 mg	21	..	..	140.84	34.20	^a APO-Famciclovir TX
							^a Ezovir AF
							^a Famciclovir Sandoz SZ
							^a Famvir NV
							^a Favic 250 SI
FAMCICLOVIR							
<u>Authority required (STREAMLINED)</u>							
3623							
Suppressive therapy of moderate to severe recurrent genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.							
<u>Note</u>							
Famciclovir 250 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.							
8217L NP	Tablet 250 mg	56	5	..	350.28	34.20	^a APO-Famciclovir TX
							^a Ezovir AF
							^a Famciclovir Sandoz SZ
							^a Famvir NV
							^a Favic 250 SI
FAMCICLOVIR							
<u>Authority required (STREAMLINED)</u>							
3625							
Treatment of immunocompromised patients with herpes zoster within 72 hours of the onset of the rash.							
<u>Note</u>							
Famciclovir is effective only if commenced within 72 hours of onset of rash.							
Famciclovir 500 mg is not PBS-subsidised for chickenpox.							
Famciclovir 500 mg is not PBS-subsidised for herpes zoster, genital herpes or other herpes simplex infections in immunocompetent patients.							
<u>Note</u>							
No applications for repeats will be authorised.							
8897G NP	Tablet 500 mg	30	..	..	198.44	34.20	^a Famvir NV
							^a Favic 500 SI
FAMCICLOVIR							
<u>Authority required (STREAMLINED)</u>							
3626							
Episodic treatment or suppressive therapy of moderate to severe recurrent genital herpes in immunocompromised patients. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.							

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
<u>Authority required (STREAMLINED)</u>							
3627							
Episodic treatment of moderate to severe recurrent oral or labial herpes in a patient with HIV infection and a CD4 cell count of less than 500 million per litre. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.							
<u>Authority required (STREAMLINED)</u>							
3628							
Suppressive therapy of moderate to severe recurrent oral or labial herpes in a patient with HIV infection and a CD4 cell count of less than 150 million per litre. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment;							
3629							
Suppressive therapy of moderate to severe recurrent oral or labial herpes in a patient with HIV infection and other opportunistic infections or AIDS defining tumours. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.							
<u>Note</u>							
Famciclovir 500 mg is not PBS-subsidised for chickenpox.							
Famciclovir 500 mg is not PBS-subsidised for herpes zoster, genital herpes or other herpes simplex infections in immunocompetent patients.							
8896F NP	Tablet 500 mg	56	5	..	350.28	34.20 ^a	Ezovir AF
						^a	Famvir NV
						^a	Favic 500 SI
VALACICLOVIR HYDROCHLORIDE							
<u>Authority required (STREAMLINED)</u>							
3632							
Moderate to severe initial genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is desirable but need not delay treatment.							
<u>Note</u>							
Valaciclovir 500 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised.							
8133C NP	Tablet 500 mg (base)	20	..	..	*105.78	34.20	Valtrex GK
VALACICLOVIR HYDROCHLORIDE							
<u>Authority required (STREAMLINED)</u>							
3633							
Episodic treatment or suppressive therapy of moderate to severe recurrent genital herpes. Microbiological confirmation of diagnosis (viral culture, antigen detection or nucleic acid amplification by PCR) is required but need not delay treatment.							
<u>Note</u>							
Valaciclovir 500 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.							
8134D NP	Tablet 500 mg (base)	30	5	..	155.43	34.20	Valtrex GK
VALACICLOVIR HYDROCHLORIDE							
<u>Authority required (STREAMLINED)</u>							
3622							
Treatment of patients with herpes zoster within 72 hours of the onset of the rash;							
3631							
Herpes zoster ophthalmicus.							
<u>Note</u>							
Valaciclovir is effective only if commenced within 72 hours of onset of rash.							
Valaciclovir 500 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.							
<u>Note</u>							
No applications for repeats will be authorised.							
8064K	Tablet 500 mg (base)	42	..	..	214.06	34.20	Valtrex GK

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
NP							

Vaccines

Bacterial vaccines

Pneumococcal vaccines

PNEUMOCOCCAL VACCINE, POLYVALENT

Restricted benefit

Splenectomised persons over 2 years of age;

Persons with Hodgkin's disease;

Persons at high risk of pneumococcal infections.

1903E	Injection 0.5 mL (23 valent)	1	..	..	46.13	34.20	Pneumovax 23	CS
NP								

Tetanus vaccines

DIPHTHERIA and TETANUS VACCINE, ADSORBED, DILUTED FOR ADULT USE

Note

For immunisation of adults and children aged greater than or equal to 8 years.

8783G	Injection 0.5 mL in pre-filled syringe	5	..	..	75.34	34.20	ADT Booster	CS
NP								

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Antineoplastic and immunomodulating agents

Antineoplastic agents

Alkylating agents

Nitrogen mustard analogues

CHLORAMBUCIL							
1163F	Tablet 2 mg	100	2	..	*137.98	34.20	Leukeran AS
CYCLOPHOSPHAMIDE							
1031G	Powder for injection 2 g (solvent required) (code 7055H applies to above item with approved solvent)	1	..	..	56.60	34.20	Endoxan BX
1079T	Powder for injection 500 mg (solvent required) (code 6704W applies to above item with approved solvent)	2	..	..	*40.70	34.20	Endoxan BX
1080W	Powder for injection 1 g (solvent required) (code 6710E applies to above item with approved solvent)	1	..	..	32.65	33.72	Endoxan BX
1266P	Tablet 50 mg	50	2	..	31.29	32.36	Cycloblastin PF

IFOSFAMIDE

Restricted benefit

Relapsed or refractory germ cell tumours following first-line chemotherapy;

Relapsed or refractory sarcomas following first-line chemotherapy.

8076C	Powder for I.V. injection 1 g	5	5	..	*342.42	34.20	Holoxan BX
8077D	Powder for I.V. injection 2 g	5	5	..	*668.37	34.20	Holoxan BX

MELPHALAN

2547C	Tablet 2 mg	25	1	..	50.88	34.20	Alkeran AS

Alkyl sulphonates

BUSULFAN

1128J	Tablet 2 mg	100	..	..	86.26	34.20	Myleran AS

Ethylene imines

THIOTEPA

2345K	Powder for injection 15 mg	2	1	..	*155.66	34.20	Sigma Pharmaceuticals (Australia) Pty Ltd SI

Nitrosoureas

CARMUSTINE

Restricted benefit

Glioblastoma multiforme, suspected or confirmed, at the time of initial surgery.

Note

Carmustine is not PBS-subsidised for use in conjunction with PBS-subsidised temozolomide.

8898H	Implants 7.7 mg, 8	±1	..	..	17539.32	34.20	Gliadel OA

FOTEMUSTINE

Authority required (STREAMLINED)

3181

Metastatic malignant melanoma.

8786K	Powder for injection 208 mg with solvent	1	4	..	1206.86	34.20	Muphoran SE

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Other alkylating agents

TEMOZOLOMIDE

Authority required

Glioblastoma multiforme concomitantly with radiotherapy.

Note

Temozolomide is not PBS-subsidised for use in conjunction with PBS-subsidised carmustine.

Note

Applications for doses above 150 mg per day will not be authorised. No applications for increased repeats will be authorised.

8819E	Capsule 5 mg	15	2	..	*208.05	34.20	Temodal	SH
8820F	Capsule 20 mg	15	2	..	*567.93	34.20	Temodal	SH
8821G	Capsule 100 mg	15	2	..	*2389.29	34.20	Temodal	SH
9361Q	Capsule 140 mg	15	2	..	*3266.10	34.20	Temodal	SH

TEMOZOLOMIDE

Authority required

Recurrence of anaplastic astrocytoma following standard therapy;

Recurrence of glioblastoma multiforme following standard therapy;

Glioblastoma multiforme following radiotherapy.

8378Y	Capsule 5 mg	5	5	..	73.75	34.20	Temodal	SH
8379B	Capsule 20 mg	5	5	..	204.39	34.20	Temodal	SH
8380C	Capsule 100 mg	5	5	..	808.22	34.20	Temodal	SH
8381D	Capsule 250 mg	5	5	..	1863.31	34.20	Temodal	SH
9362R	Capsule 140 mg	5	5	..	1112.18	34.20	Temodal	SH

Antimetabolites

Folic acid analogues

METHOTREXATE

1622J	Tablet 2.5 mg	30	5	..	13.26	14.33	^a Hospira Pty Limited	HH
						^a Methoblastin	PF	
2272N	Tablet 10 mg	15	1	..	22.15	23.22	Methoblastin	PF
2395C	Injection 50 mg in 2 mL	5	..	..	36.12	34.20	^a Hospira Pty Limited	HH
						^a Pfizer Australia Pty Ltd	PF	
2396D	Injection 5 mg in 2 mL	5	..	..	36.81	34.20	Hospira Pty Limited	HH
8851W	Solution concentrate for I.V. infusion 1000 mg in 10 mL	1	..	..	120.12	34.20	^a Hospira Pty Limited	HH
						^a Methotrexate	IT	
						Ebewe		
8852X	Solution concentrate for I.V. infusion 5000 mg in 50 mL	1	..	..	543.90	34.20	Methotrexate	IT
						Ebewe		
8863L	Solution concentrate for I.V. infusion 500 mg in 20 mL	1	..	..	63.27	34.20	Hospira Pty Limited	HH

METHOTREXATE

Restricted benefit

For patients requiring doses greater than 20 mg per week.

1623K	Tablet 10 mg	50	2	..	45.98	34.20	Methoblastin	PF

PEMETREXED DISODIUM

Authority required

Locally advanced or metastatic non-small cell lung cancer, after prior platinum-based chemotherapy.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	Doses greater than 500 mg per metre squared body surface area (BSA) will not be approved for PBS subsidy. The patient's BSA must be provided at the time of the authority approval.						
	<u>Authority required</u>						
	Mesothelioma in combination with cisplatin.						
	Doses greater than 500 mg per metre squared body surface area (BSA) will not be approved for PBS subsidy. The patient's BSA must be provided at the time of the authority approval.						

Authority required

Mesothelioma in combination with cisplatin.

Doses greater than 500 mg per metre squared body surface area (BSA) will not be approved for PBS subsidy. The patient's BSA must be provided at the time of the authority approval.

Note

No applications for increased maximum quantities for the 500 mg vial will be authorised.

9130M	Powder for I.V. infusion 500 mg (base)	1	3	..	1701.41	34.20	Alimta	LY
9131N	Powder for I.V. infusion 100 mg (base)	1	3	..	359.85	34.20	Alimta	LY

RALTITREXED

Authority required (STREAMLINED)

3185

For use as a single agent in the treatment of advanced colorectal cancer.

8284B	Powder for I.V. infusion 2 mg	3	2	..	*856.29	34.20	Tomudex	HH

Purine analogues

CLADRIBINE

Authority required (STREAMLINED)

3180

Hairy cell leukaemia.

1811H	Solution for I.V. infusion 10 mg in 10 mL	7	..	..	*4629.57	34.20	Leustatin	JC
8800E	Injection 10 mg in 5 mL	7	..	..	*4629.57	34.20	Litak	OA

FLUDARABINE PHOSPHATE

Authority required

B-cell chronic lymphocytic leukaemia in combination with cyclophosphamide where the patient has advanced disease (Binet Stage B or C) or evidence of progressive Stage A disease.

Stage A progressive disease is defined by at least one of the following: persistent rise in lymphocyte count with doubling time less than 12 months; a downward trend in haemoglobin or platelets, or both; more than 50% increase in the size of liver, spleen, or lymph nodes, or appearance of these signs if not previously present; constitutional symptoms attributable to disease.

The diagnosis of chronic lymphocytic leukaemia (CLL) must have been established based on:
(a) a lymphocytosis, with more than 5,000 million lymphocytes per L in the peripheral blood; and
(b) a clonal population of B-cells (CD5/CD19) documented by flow cytometry.

9184J	Tablet 10 mg	20	5	..	955.69	34.20	Fludara	GZ

FLUDARABINE PHOSPHATE

Authority required

B-cell chronic lymphocytic leukaemia in combination with cyclophosphamide where the patient has advanced disease (Binet Stage B or C) or evidence of progressive Stage A disease.

Stage A progressive disease is defined by at least one of the following: persistent rise in lymphocyte count with doubling time less than 12 months; a downward trend in haemoglobin or platelets, or both; more than 50% increase in the size of liver, spleen, or lymph nodes, or appearance of these signs if not previously present; constitutional symptoms attributable to disease.

The diagnosis of chronic lymphocytic leukaemia (CLL) must have been established based on:
(a) a lymphocytosis, with more than 5,000 million lymphocytes per L in the peripheral blood; and
(b) a clonal population of B-cells (CD5/CD19) documented by flow cytometry.

Note

The solution for I.V. injection and powder for I.V. injection (after reconstitution) are bioequivalent.

9185K	Powder for I.V. injection 50 mg	5	3	..	1535.81	34.20	^a Fludara	GZ
					*1535.82	34.20	^a Farine	WQ
							^a Fludarabine Actavis	GQ
9207N	Solution for I.V. injection 50 mg in 2 mL	5	3	..	1535.81	34.20	^a Fludarabine Ebewe	IT

Antineoplastic and immunomodulating agents

					Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium			Brand Name and Manufacturer	
	MERCAPTOPURINE							
1598D	Tablet 50 mg	100	2	..	*251.94	34.20	Purinethol	AS
	THIOGUANINE							
1233X	Tablet 40 mg	25	1	..	198.66	34.20	Lanvis	AS

Pyrimidine analogues

CAPECITABINE

Authority required

Advanced breast cancer after failure of prior therapy which includes a taxane and an anthracycline;

Advanced breast cancer where therapy with a taxane and/or an anthracycline is contraindicated;

Advanced breast cancer in combination with docetaxel after failure of prior anthracycline-containing chemotherapy;

Treatment of advanced or metastatic colorectal cancer;

Adjuvant treatment of stage III (Dukes C) colon cancer, following complete resection of the primary tumour;

Advanced (Stage III or IV) oesophago-gastric cancer, previously untreated, in combination with a cisplatin-based regimen, in a patient with a WHO performance status of 2 or less.

Note

In the adjuvant setting, the recommended treatment duration is 24 weeks.

Capecitabine is not PBS-subsidised for the treatment of patients with stage II (Dukes B) colon cancer.

Capecitabine is not PBS-subsidised for the adjuvant treatment of patients with rectal cancer.

8361C	Tablet 150 mg	60	2	..	123.93	34.20	Xeloda	RO
8362D	Tablet 500 mg	120	2	..	695.17	34.20	Xeloda	RO

CYTARABINE

2884T	Injection 100 mg in 5 mL	10	1	..	*125.78	34.20	Pfizer Australia Pty Ltd	PF

FLUOROURACIL

2528C	Injection 500 mg in 10 mL	10	..	..	*55.70	34.20	^a Fluorouracil Ebewe	IT
					*55.72	34.20	^a Hospira Pty Limited	HH
9005Y	Injection 1000 mg in 20 mL	5	..	..	*48.97	34.20	Fluorouracil Ebewe	IT

GEMCITABINE

Authority required

Advanced breast cancer in combination with paclitaxel after failure of prior therapy which includes an anthracycline;

Advanced epithelial ovarian cancer, in combination with carboplatin, in patients who relapse more than 6 months after platinum-based therapy;

Locally advanced or metastatic non-small cell lung cancer;

Locally advanced or metastatic adenocarcinoma of the pancreas;

Locally advanced or metastatic bladder cancer, in combination with cisplatin.

Note

The powder for I.V. infusion 200 mg (as hydrochloride) (after reconstitution) and the solution concentrate for I.V. infusion 200 mg (as hydrochloride) are bioequivalent.

8049P	Powder for I.V. infusion 200 mg (as hydrochloride)	4	2	..	*205.78	34.20	^a DBL Gemcitabine for Injection	HH
							^a Gemcitabine Actavis	GQ
							^a Gemcitabine Ebewe	IT
							^a Gemcitabine Kabi	PK
							^a Gemcitabine Sun	ZF
							^a Gemcitate	ZP
							^a Gemzar	LY
9401T	Solution concentrate for I.V. infusion 200 mg (as hydrochloride) in 20 mL	4	2	..	*205.78	34.20	^a Gemcitabine Ebewe	IT

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer	
GEMCITABINE									
<u>Authority required</u>									
Advanced breast cancer in combination with paclitaxel after failure of prior therapy which includes an anthracycline;									
Advanced epithelial ovarian cancer, in combination with carboplatin, in patients who relapse more than 6 months after platinum-based therapy;									
Locally advanced or metastatic non-small cell lung cancer;									
Locally advanced or metastatic adenocarcinoma of the pancreas;									
Locally advanced or metastatic bladder cancer, in combination with cisplatin.									
<u>Note</u>									
The powder for I.V. infusion 1 g (as hydrochloride) (after reconstitution) and the solution concentrate for I.V. infusion 1000 mg (as hydrochloride) are bioequivalent.									
8050Q	Powder for I.V. infusion 1 g (as hydrochloride)	2	2	..	*472.24	34.20	^a	DBL Gemcitabine for Injection Gemcitabine Actavis Gemcitabine Ebewe Gemcitabine Kabi Gemcitabine Sun Gemcite Gemzar	HH GQ IT PK ZF ZP LY
9402W	Solution concentrate for I.V. infusion 1000 mg (as hydrochloride) in 100 mL	2	2	..	*472.24	34.20	^a	Gemcitabine Ebewe	IT
GEMCITABINE									
<u>Authority required</u>									
Advanced breast cancer in combination with paclitaxel after failure of prior therapy which includes an anthracycline;									
Advanced epithelial ovarian cancer, in combination with carboplatin, in patients who relapse more than 6 months after platinum-based therapy;									
Locally advanced or metastatic non-small cell lung cancer;									
Locally advanced or metastatic adenocarcinoma of the pancreas;									
Locally advanced or metastatic bladder cancer, in combination with cisplatin.									
9414L	Powder for I.V. infusion 2 g (as hydrochloride)	1	2	..	473.26	34.20	^a	DBL Gemcitabine for Injection Gemcitabine Kabi	HH PK
9463C	Solution concentrate for I.V. infusion 500 mg (as hydrochloride) in 50 mL	4	2	..	*472.26	34.20		Gemcitabine Ebewe	IT
Plant alkaloids and other natural products									
<i>Vinca alkaloids and analogues</i>									
VINBLASTINE SULFATE									
2199R	Solution for I.V. injection 10 mg in 10 mL	5	..	..	169.78	34.20		Hospira Pty Limited	HH
VINCRISTINE SULFATE									
2374Y	I.V. injection 1 mg in 1 mL	10	..	..	*155.20	34.20	^a	Hospira Pty Limited	HH
							^a	Pfizer Australia Pty Ltd	PF
VINORELBINE TARTRATE									
<u>Authority required</u>									
Locally advanced or metastatic non-small cell lung cancer.									
9009E	Capsule 20 mg (base)	20	2	..	*2011.82	34.20		Navelbine	FB
9010F	Capsule 30 mg (base)	16	2	..	*2386.26	34.20		Navelbine	FB

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer		
					\$	\$			
VINORELBINE TARTRATE									
<u>Authority required</u>									
Advanced breast cancer after failure of prior therapy which includes an anthracycline;									
Locally advanced or metastatic non-small cell lung cancer.									
8280T	Solution for I.V. infusion 10 mg (base) in 1 mL	16	2	..	*1137.14	34.20	^a	Hospira Pty Limited	HH
							^a	Navelbine	FB
							^a	Vinorelbine Ebewe	IT
							^a	Vinorelbine Link	PK
8281W	Solution for I.V. infusion 50 mg (base) in 5 mL	4	2	..	*1186.02	34.20	^a	Hospira Pty Limited	HH
							^a	Navelbine	FB
							^a	Vinorelbine Ebewe	IT
							^a	Vinorelbine Link	PK

Podophyllotoxin derivatives

ETOPOSIDE								
1389D	Capsule 100 mg	10	..	..	398.21	34.20	Vepesid	BQ
1390E	Solution for I.V. infusion 100 mg in 5 mL	5	..	..	*166.67	34.20	^a Hospira Pty Limited	HH
					166.69	34.20	^a Etoposide Ebewe	IT
1396L	Capsule 50 mg	20	..	..	453.52	34.20	Vepesid	BQ
8120J	Powder for I.V. infusion 100 mg (as phosphate)	5	..	..	*166.67	34.20	Etopophos	BQ
8515E	Powder for I.V. infusion 1 g (as phosphate)	1	..	..	315.76	34.20	Etopophos	BQ

Taxanes

DOCETAXEL

Authority required

Neoadjuvant treatment of a patient with a WHO performance status of 1 or less, with inoperable Stage III, IVa or IVb squamous cell carcinoma of the oral cavity, larynx, oropharynx or hypopharynx, in combination with cisplatin and fluorouracil.

Note

The carcinoma can be considered inoperable for technical or organ preservation reasons.

5462L	Solution concentrate for I.V. infusion 20 mg in 1 mL	1	..	..	334.39	34.20	Taxotere	SW
9291B	Injection set containing 1 single use vial concentrate for I.V. infusion 20 mg (anhydrous) in 0.5 mL with solvent	1	..	..	334.39	34.20	Taxotere	SW

DOCETAXEL

Authority required

Adjuvant treatment of node-positive breast cancer in combination with an anthracycline and cyclophosphamide;

Advanced breast cancer after failure of prior therapy;

Advanced metastatic ovarian cancer after failure of prior therapy which includes a platinum compound;

Locally advanced or metastatic non-small cell lung cancer;

Treatment of HER2 positive early breast cancer in combination with trastuzumab.

Authority required

Treatment of androgen independent (hormone refractory) metastatic carcinoma of the prostate in a patient with a Karnofsky performance-status score of at least 60%. Docetaxel must be used as first-line chemotherapy and administered in three weekly cycles.

Note

A maximum of 10 cycles of treatment with docetaxel will be authorised under this restriction.

Authority required

Adjuvant treatment of operable breast cancer in combination with cyclophosphamide.

Note

A maximum of four cycles of treatment will be authorised under this restriction.

5463M	Solution concentrate for I.V. infusion 20 mg in 1 mL	2	..	..	*651.16	34.20	Taxotere	SW
8071T	Injection set containing 1 single use vial concentrate for I.V. infusion 20 mg	2	..	..	*651.16	34.20	Taxotere	SW

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
	(anhydrous) in 0.5 mL with solvent							
DOCETAXEL								
<u>Authority required</u>								
Neoadjuvant treatment of a patient with a WHO performance status of 1 or less, with inoperable Stage III, IVa or IVb squamous cell carcinoma of the oral cavity, larynx, oropharynx or hypopharynx, in combination with cisplatin and fluorouracil.								
<u>Note</u>								
The carcinoma can be considered inoperable for technical or organ preservation reasons.								
<u>Authority required</u>								
Adjuvant treatment of node-positive breast cancer in combination with an anthracycline and cyclophosphamide;								
Advanced breast cancer after failure of prior therapy;								
Advanced metastatic ovarian cancer after failure of prior therapy which includes a platinum compound;								
Locally advanced or metastatic non-small cell lung cancer;								
Treatment of HER2 positive early breast cancer in combination with trastuzumab.								
<u>Authority required</u>								
Treatment of androgen independent (hormone refractory) metastatic carcinoma of the prostate in a patient with a Karnofsky performance-status score of at least 60%. Docetaxel must be used as first-line chemotherapy and administered in three weekly cycles.								
<u>Note</u>								
A maximum of 10 cycles of treatment with docetaxel will be authorised under this restriction.								
<u>Authority required</u>								
Adjuvant treatment of operable breast cancer in combination with cyclophosphamide.								
<u>Note</u>								
A maximum of four cycles of treatment will be authorised under this restriction.								
5464N	Solution concentrate for I.V. infusion 80 mg in 4 mL	1	..	..	1280.83	34.20	Taxotere	SW
8074Y	Injection set containing 1 single use vial concentrate for I.V. infusion 80 mg (anhydrous) in 2 mL with solvent	1	..	..	1280.83	34.20	Taxotere	SW
 NAB PACLITAXEL								
<u>Authority required</u>								
Metastatic breast cancer after failure of prior therapy.								
9415M	Powder for I.V. injection 100 mg (base)	1	..	..	456.09	34.20	Abraxane	TS
 PACLITAXEL								
<u>Authority required</u>								
Adjuvant treatment of node-positive breast cancer administered sequentially to an anthracycline and cyclophosphamide;								
Advanced breast cancer after failure of prior therapy;								
Advanced metastatic ovarian cancer after failure of prior therapy which includes a platinum compound;								
Primary treatment of ovarian cancer in combination with a platinum compound;								
Locally advanced or metastatic non-small cell lung cancer;								
Treatment of HER2 positive early breast cancer in combination with trastuzumab.								
3017T	Solution concentrate for I.V. infusion 150 mg in 25 mL	2	..	..	*1796.32	34.20	^a Anzatax	HH
							^a Paclitaxel Actavis	GQ
							^a Paclitaxel Ebewe	IT
							^a Plaxel	WQ
3026G	Solution concentrate for I.V. infusion 30 mg in 5 mL	5	..	..	*931.92	34.20	^a Anzatax	HH
							^a Paclitaxel Actavis	GQ
							^a Paclitaxel Kabi	PK
							^a Plaxel	WQ
							^a Taxol	BQ
					931.93	34.20	^a Paclitaxel Ebewe	IT

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
8018B	Solution concentrate for I.V. infusion 100 mg in 16.7 mL	2	..	..	*1234.36	34.20	^a Anzatax	HH
							^a Paclitaxel Actavis	GQ
							^a Paclitaxel Ebewe	IT
							^a Paclitaxel Kabi	PK
							^a Plaxel	WQ
8360B	Solution concentrate for I.V. infusion 300 mg in 50 mL	1	..	..	1813.70	34.20	^a Taxol	BQ
							^a Anzatax	HH
							^a Paclitaxel Actavis	GQ
							^a Paclitaxel Ebewe	IT
							^a Paclitaxel Kabi	PK

Cytotoxic antibiotics and related substances

Anthracyclines and related substances

DOXORUBICIN HYDROCHLORIDE

1336H	Solution for I.V. injection or intravesical administration 10 mg in 5 mL	4	..	..	*40.46	34.20	^a Adriamycin Solution	PF
							^a Doxorubicin Ebewe	IT
							^a Hospira Pty Limited	HH
1340M	Solution for I.V. injection or intravesical administration 20 mg in 10 mL	4	..	..	*64.58	34.20	Adriamycin Solution	PF
1342P	Solution for I.V. injection or intravesical administration 50 mg in 25 mL	3	..	..	*109.59	34.20	^a Adriamycin Solution	PF
							^a Doxorubicin Ebewe	IT
							^a Hospira Pty Limited	HH
8827N	Solution for I.V. injection or intravesical administration 100 mg in 50 mL	1	..	..	75.18	34.20	Doxorubicin Ebewe	IT
8828P	Solution for I.V. injection or intravesical administration 200 mg in 100 mL	1	..	..	143.93	34.20	^a Adriamycin	PF
							^a Doxorubicin Ebewe	IT

DOXORUBICIN HYDROCHLORIDE, PEGYLATED LIPOSOMAL

Authority required

Advanced epithelial ovarian cancer in women who have failed a first-line platinum-based chemotherapy regimen;

Metastatic breast cancer, as monotherapy, after failure of prior therapy which includes capecitabine and a taxane;

Metastatic breast cancer, as monotherapy, where therapy with capecitabine and/or a taxane is contraindicated.

8569B	Suspension for I.V. infusion 20 mg in 10 mL	1	..	..	703.05	34.20	Caelyx	SH
8570C	Suspension for I.V. infusion 50 mg in 25 mL	1	..	..	1621.79	34.20	Caelyx	SH

EPIRUBICIN HYDROCHLORIDE

1375J	Solution for injection 10 mg in 5 mL	4	..	..	*212.02	34.20	^a Epirubicin Ebewe	IT
							^a Pharmorubicin Solution	PF
1376K	Solution for injection 20 mg in 10 mL	4	..	..	*386.54	34.20	Pharmorubicin Solution	PF
1377L	Solution for injection 50 mg in 25 mL	4	..	..	*937.70	34.20	^a Epirubicin Ebewe	IT
							^a Hospira Pty Limited	HH
							^a Pharmorubicin Solution	PF
8817C	Solution for injection 100 mg in 50 mL	2	..	..	*925.20	34.20	^a Epirubicin Ebewe	IT
							^a Hospira Pty Limited	HH
8858F	Solution for injection 200 mg in 100 mL	1	..	..	911.44	34.20	Epirubicin Ebewe	IT

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
IDARUBICIN HYDROCHLORIDE								
<u>Restricted benefit</u>								
Acute myelogenous leukaemia.								
2446R	Capsule 5 mg	3	..	..	*247.11	34.20	Zavedos	PF
2448W	Capsule 10 mg	3	..	..	*451.62	34.20	Zavedos	PF
8530Y	Solution for I.V. injection 5 mg in 5 mL	3	..	..	568.88	34.20	Zavedos Solution	PF
8531B	Solution for I.V. injection 10 mg in 10 mL	6	..	..	2104.80	34.20	Zavedos Solution	PF

MITOZANTRONE HYDROCHLORIDE								
1929M	Injection 20 mg (base) in 10 mL	1	..	..	179.10	34.20	^a Hospira Pty Limited ^a Mitozantrone ^a Ebewe ^a Onkotrone	HH IT BX PF
1930N	Injection 25 mg (base) in 12.5 mL	1	..	..	220.54	34.20	^a Onkotrone ^a Pfizer Australia Pty Ltd	BX PF
1932Q	Injection 10 mg (base) in 5 mL	1	..	..	92.76	34.20	Pfizer Australia Pty Ltd	PF

Other antineoplastic agents *Platinum compounds*

CARBOPLATIN								
1160C	Solution for I.V. injection 50 mg in 5 mL	2	..	..	*65.86	34.20	^a Carboplatin Ebewe ^a Hospira Pty Limited ^a Pfizer Australia Pty Ltd	IT HH PF
1161D	Solution for I.V. injection 150 mg in 15 mL	6	..	..	*415.80	34.20	^a Carboplatin Ebewe ^a Hospira Pty Limited ^a Pfizer Australia Pty Ltd	IT HH PF
1162E	Solution for I.V. injection 450 mg in 45 mL	2	..	..	*270.24	34.20	^a Carboplatin Ebewe ^a Hospira Pty Limited ^a Pfizer Australia Pty Ltd	IT HH PF

CISPLATIN								
2578Q	I.V. injection 10 mg in 10 mL	1	..	..	14.52	15.59	Pfizer Australia Pty Ltd	PF
2579R	I.V. injection 50 mg in 50 mL	1	..	..	28.13	29.20	^a Hospira Pty Limited ^a Pfizer Australia Pty Ltd	HH PF
2580T	I.V. injection 100 mg in 100 mL	1	..	..	58.76	34.20	^a Cisplatin Ebewe ^a Hospira Pty Limited ^a Pfizer Australia Pty Ltd	IT HH PF

OXALIPLATIN

Authority required

Metastatic colorectal cancer in a patient with a WHO performance status of 2 or less, to be used in combination with:

(a) capecitabine; or

(b) 5-fluorouracil and folinic acid;

Adjuvant treatment of stage III (Dukes C) colon cancer, in combination with 5-fluorouracil and folinic acid, following complete resection of the primary tumour.

Note

Oxaliplatin is not PBS-subsidised for the treatment of patients with stage II (Dukes B) colon cancer.

Oxaliplatin is not PBS-subsidised for the adjuvant treatment of patients with rectal cancer.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Note								
The solution concentrate for I.V. infusion 50 mg and powder for I.V. infusion 50 mg (after reconstitution) are bioequivalent.								
8539K	Powder for I.V. infusion 50 mg	1	2	..	356.96	34.20	^a Hospira Pty Limited	HH
							^a Oxalatin	ZP
							^a Oxaliplatin Actavis	GQ
							^a Oxaliplatin	AF
							^a Alphapharm	
							^a Oxaliplatin Ebewe	IT
							^a Oxaliplatin Link	PK
							^a Xalox	WQ
8847P	Solution concentrate for I.V. infusion 50 mg in 10 mL	1	2	..	356.96	34.20	^a DBL Oxaliplatin Concentrate	HH
							^a Eloxatin	SW
							^a Oxaliplatin Kabi	PK
OXALIPLATIN								
Authority required								
Metastatic colorectal cancer in a patient with a WHO performance status of 2 or less, to be used in combination with:								
(a) capecitabine; or								
(b) 5-fluorouracil and folinic acid;								
Adjuvant treatment of stage III (Dukes C) colon cancer, in combination with 5-fluorouracil and folinic acid, following complete resection of the primary tumour.								
Note								
Oxaliplatin is not PBS-subsidised for the treatment of patients with stage II (Dukes B) colon cancer.								
Oxaliplatin is not PBS-subsidised for the adjuvant treatment of patients with rectal cancer.								
Note								
The solution concentrate for I.V. infusion 100 mg and powder for I.V. infusion 100 mg (after reconstitution) are bioequivalent.								
8540L	Powder for I.V. infusion 100 mg	1	2	..	674.47	34.20	^a Hospira Pty Limited	HH
							^a Oxalatin	ZP
							^a Oxaliplatin Actavis	GQ
							^a Oxaliplatin	AF
							^a Alphapharm	
							^a Oxaliplatin Ebewe	IT
							^a Oxaliplatin Link	PK
							^a Winthrop	WA
							^a Oxaliplatin	
							^a Xalox	WQ
8848Q	Solution concentrate for I.V. infusion 100 mg in 20 mL	1	2	..	674.47	34.20	^a DBL Oxaliplatin Concentrate	HH
							^a Eloxatin	SW
							^a Oxaliplatin Kabi	PK
OXALIPLATIN								
Authority required								
Metastatic colorectal cancer in a patient with a WHO performance status of 2 or less, to be used in combination with:								
(a) capecitabine; or								
(b) 5-fluorouracil and folinic acid;								
Adjuvant treatment of stage III (Dukes C) colon cancer, in combination with 5-fluorouracil and folinic acid, following complete resection of the primary tumour.								
Note								
Oxaliplatin is not PBS-subsidised for the treatment of patients with stage II (Dukes B) colon cancer.								
Oxaliplatin is not PBS-subsidised for the adjuvant treatment of patients with rectal cancer.								
2310N	Solution concentrate for I.V. infusion 200 mg in 40 mL	1	2	..	1317.56	34.20	Eloxatin	SW

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Monoclonal antibodies

BEVACIZUMAB

Authority required

Initial PBS-subsidised treatment, in combination with first-line chemotherapy, of a patient with previously untreated metastatic colorectal cancer with a WHO performance status of 0 or 1.

The maximum dose that will be approved is 5 mg per kg every 2 weeks or 7.5 mg per kg every 3 weeks.

Note

Not for use as monotherapy.

Authority required

Continuing PBS-subsidised treatment, in combination with first-line chemotherapy, of a patient with metastatic colorectal cancer who has previously been issued with an authority prescription for bevacizumab and who does not have progressive disease and who remains on first-line chemotherapy. The maximum dose that will be approved is 5 mg per kg every 2 weeks or 7.5 mg per kg every 3 weeks.

Note

Not for use as monotherapy.

Note

Special Pricing Arrangements apply.

9442Y	Solution for I.V. infusion 100 mg in 4 mL	1	..	..	534.77	34.20	Avastin	RO
9443B	Solution for I.V. infusion 400 mg in 16 mL	1	..	..	1866.36	34.20	Avastin	RO

CETUXIMAB

Authority required

Initial treatment of stage III, IVa or IVb squamous cell cancer of the larynx, oropharynx or hypopharynx for the week prior to radiotherapy, where cisplatin is contraindicated according to the TGA-approved Product Information;

Initial treatment of stage III, IVa or IVb squamous cell cancer of the larynx, oropharynx or hypopharynx, in combination with radiotherapy, where cisplatin is not tolerated.

Note

No applications for repeats will be authorised.

9136W	Solution for I.V. infusion 100 mg in 20 mL	1	..	..	391.06	34.20	Erbix	SG
9137X	Solution for I.V. infusion 500 mg in 100 mL	1	..	..	1851.36	34.20	Erbix	SG

CETUXIMAB

Authority required

Continuing treatment of stage III, IVa or IVb squamous cell cancer of the larynx, oropharynx or hypopharynx, in combination with radiotherapy, where cisplatin is either contraindicated or not tolerated.

Note

A maximum lifetime supply for this indication is limited to a maximum of 8 treatments per site and to 10 treatments per site for patients in whom radiotherapy is interrupted.

9138Y	Solution for I.V. infusion 100 mg in 20 mL	1	6	..	391.06	34.20	Erbix	SG
9139B	Solution for I.V. infusion 500 mg in 100 mL	1	6	..	1851.36	34.20	Erbix	SG

RITUXIMAB

Authority required

Relapsed or refractory low-grade B-cell non-Hodgkin's lymphoma;

Relapsed or refractory follicular B-cell non-Hodgkin's lymphoma.

8293L	Solution for I.V. infusion 100 mg in 10 mL	2	3	..	948.07	34.20	Mabthera	RO
8294M	Solution for I.V. infusion 500 mg in 50 mL	1	3	..	2339.99	34.20	Mabthera	RO

RITUXIMAB

Authority required

Treatment of previously untreated, CD20 positive, diffuse large B-cell non-Hodgkin's lymphoma, in combination with chemotherapy;

Treatment of symptomatic patients with previously untreated, CD20 positive, Stage III or IV, follicular, B-cell non-Hodgkin's lymphoma, in combination with chemotherapy.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	
8665C	Solution for I.V. infusion 100 mg in 10 mL	2	7	..	948.07	34.20	Mabthera RO
8666D	Solution for I.V. infusion 500 mg in 50 mL	1	7	..	2339.99	34.20	Mabthera RO

Protein kinase inhibitors

DASATINIB

Note

Any queries concerning the arrangements to prescribe dasatinib may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Any queries concerning patients who are enrolled on the Dasatinib Compassionate Program may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe dasatinib should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001.

Authority required

Initial treatment, as the sole PBS-subsidised therapy, of a patient with chronic myeloid leukaemia in any disease phase bearing the Philadelphia chromosome or expressing the transcript, BCR-ABL, who has active leukaemia (as defined by presence on current pathology assessments of either the Philadelphia chromosome on cytogenetic or FISH analysis, or the presence of the transcript BCR-ABL greater than 1% on the international scale) and who has failed an adequate trial of imatinib.

Failure of an adequate trial of imatinib is defined as:

- (i) Lack of response to initial imatinib therapy, defined as either:
 - failure to achieve a haematological response after a minimum of 3 months therapy with imatinib for patients initially treated in chronic phase; or
 - failure to achieve any cytogenetic response after a minimum of 6 months therapy with imatinib for patients initially treated in chronic phase as demonstrated on bone marrow biopsy by presence of greater than 95% Philadelphia chromosome positive cells; or
 - failure to achieve a major cytogenetic response or a peripheral blood BCR-ABL level of less than 1% after a minimum of 12 months therapy with imatinib; OR
- (ii) Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing imatinib therapy; OR
- (iii) Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing in value by at least 5 fold to a level of greater than 1% confirmed on a subsequent test), during ongoing imatinib therapy; OR
- (iv) Development of accelerated phase or blast crisis in a patient previously prescribed imatinib for any phase of chronic myeloid leukaemia.

Accelerated phase is defined by the presence of 1 or more of the following:

- (1) Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 15% but less than 30%; or
- (2) Percentage of blasts plus promyelocytes in the peripheral blood or bone marrow greater than or equal to 30%; or
- (3) Peripheral basophils greater than or equal to 20%; or
- (4) Progressive splenomegaly to a size greater than or equal to 10 cm below the left costal margin to be confirmed on 2 occasions at least 4 weeks apart, or a greater than or equal to 50% increase in size below the left costal margin over 4 weeks; or
- (5) Karyotypic evolution (chromosomal abnormalities in addition to a single Philadelphia chromosome).

Blast crisis is defined as either:

- (1) Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 30%; or
- (2) Extramedullary involvement other than spleen and liver; OR
- (v) Disease progression (defined as a greater than or equal to 50% increase in peripheral white blood cell count, blast count, basophils or platelets) during first-line imatinib therapy in patients with accelerated phase or blast crisis chronic myeloid leukaemia; OR
- (vi) Grade 3 or 4 non-haematological toxicity that is imatinib related and necessitates permanent cessation of imatinib. For patients with imatinib related toxicities, leukaemia activity does not need to be demonstrated.

Applications for authorisation must be in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Chronic Myeloid Leukaemia Dasatinib/Nilotinib PBS Authority Application - Supporting Information Form; and
- (c) a signed patient acknowledgement; and
- (d) a bone marrow biopsy pathology report demonstrating the patient has active chronic myeloid leukaemia, either manifest as cytogenetic evidence of the Philadelphia chromosome, or RT-PCR level of BCR-ABL transcript greater than 1% on the international scale. (The date of the

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
	relevant pathology report needs to be provided); and (e) where there has been a loss of response to imatinib, a copy of the current confirming pathology report(s) from an Approved Pathology Authority or details of the dates of assessment in the case of progressive splenomegaly or extramedullary involvement; or (f) details of Grade 3 or 4 non-haematological toxicity.						
	Note Dasatinib will only be subsidised for patients with chronic myeloid leukaemia who are not receiving concomitant PBS-subsidised imatinib mesylate, nilotinib or interferon alfa therapy.						
	Patients should be commenced on a dose of dasatinib of at least 100 mg (base) daily. Continuing therapy is dependent on patients demonstrating a major cytogenetic response to dasatinib therapy or a peripheral blood BCR-ABL level of less than 1% at 18 months and thereafter at 12 monthly intervals, irrespective of the daily dasatinib dose received.						
	From 1 November 2008, under the PBS, a patient will be able to trial either dasatinib and/or nilotinib within the initial 18 month treatment period, providing the patient's CML is not resistant to the first second-line agent.						
	Dasatinib is not PBS-subsidised for patients with CML that is resistant to nilotinib.						
9282M	Tablet 20 mg	60	2	..	3246.36	34.20	Sprycel BQ
9283N	Tablet 50 mg	60	2	..	5250.68	34.20	Sprycel BQ
9284P	Tablet 70 mg	60	2	..	6465.01	34.20	Sprycel BQ
9341P	Tablet 100 mg	30	2	..	5250.68	34.20	Sprycel BQ

DASATINIB

Note

Any queries concerning the arrangements to prescribe dasatinib may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Any queries concerning patients who are enrolled on the Dasatinib Compassionate Program may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe dasatinib should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001.

Authority required

Continuing treatment, as the sole PBS-subsidised therapy, of a patient who has received initial treatment with dasatinib as a pharmaceutical benefit for chronic myeloid leukaemia, and who has demonstrated either a major cytogenetic response, or less than 1% BCR-ABL level in the blood, to dasatinib in the preceding 18 months and thereafter at 12 monthly intervals.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Chronic Myeloid Leukaemia Dasatinib/Nilotinib Authority Application Form for continuing treatment; and
- (3) demonstration of continued response to treatment as evidenced by either:
 - (a) major cytogenetic response [see Note explaining definitions of response]. Where this has been supplied within the previous 12 months (or 18 months for the initial supply), only the date of the relevant pathology report needs to be provided; or
 - (b) a peripheral blood level of BCR-ABL of less than 1% on the international scale [see Note explaining definitions of response]. Where this has been supplied within the previous 12 months (or 18 months for the initial supply), only the date of the relevant pathology report needs to be provided.

Note

Definitions of response.

A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells.

A bone marrow or peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the biological equivalent of a major cytogenetic response.

Authority approval requirements.

For the purposes of assessing response to PBS-subsidised treatment with dasatinib, either cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted as follows:

- (i) between 10 and 18 months of the commencement of treatment with dasatinib, at which time patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% has been demonstrated may receive authorisation for a further 12 months of treatment; and

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
(ii) at no greater than 12 month intervals thereafter, to demonstrate that the major cytogenetic response or peripheral blood BCR-ABL level of less than 1% has been sustained. For each authority application where eligibility for continuing PBS-subsidised treatment is to be demonstrated, a copy of the cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or a copy of the quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted as described in (i) and (ii) above. For bone marrow analyses, where the standard karyotyping conducted at the time of application is not informative, a copy of a cytogenetic analysis conducted on the bone marrow using FISH with BCR-ABL specific probe must be submitted with the authority application. A copy of the non-informative standard karyotype analysis must be included with the authority application. Where a patient has previously received PBS-subsidised treatment with dasatinib, no approval will be granted for PBS-subsidised re-treatment where that patient has at any time failed to meet the criteria for continuing treatment.							
2478K	Tablet 20 mg	60	5	..	3246.36	34.20	Sprycel BQ
2482P	Tablet 50 mg	60	5	..	5250.68	34.20	Sprycel BQ
2485T	Tablet 70 mg	60	5	..	6465.01	34.20	Sprycel BQ
9342Q	Tablet 100 mg	30	5	..	5250.68	34.20	Sprycel BQ

DASATINIB

Note

Any queries concerning the arrangements to prescribe dasatinib may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Any queries concerning patients who are enrolled on the Dasatinib Compassionate Program may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe dasatinib should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001.

Authority required

Initial treatment, as monotherapy, of a patient with acute lymphoblastic leukaemia (ALL) bearing the Philadelphia chromosome or expressing the transcript, BCR-ABL, who has failed treatment with chemotherapy AND imatinib and where appropriate, allogeneic haemopoietic stem cell transplantation.

Failure of treatment is defined as either:

- (i) Failure to achieve a complete morphological and cytogenetic remission after a minimum of 2 months treatment with intensive chemotherapy and imatinib;
- (ii) Morphological or cytogenetic relapse of leukaemia after achieving a complete remission induced by chemotherapy and imatinib;
- (iii) Morphological or cytogenetic relapse or persistence of leukaemia after allogeneic haemopoietic stem cell transplantation.

Patients must have active leukaemia, as defined by presence on current pathology assessments of either morphological infiltration of the bone marrow (greater than 5% lymphoblasts) or cerebrospinal fluid or other sites; OR the presence of cells bearing the Philadelphia chromosome on cytogenetic or FISH analysis in the bone marrow of patients in morphological remission.

The first authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Acute Lymphoblastic Leukaemia Dasatinib PBS Authority Application - Supporting Information Form; and
- (c) a signed patient acknowledgement; and
- (d) a pathology report demonstrating that the patient has active acute lymphoblastic leukaemia, either manifest as cytogenetic evidence of the Philadelphia chromosome, or morphological evidence of acute lymphoblastic leukaemia plus qualitative RT-PCR evidence of BCR-ABL transcript. The date of the relevant pathology report(s) need(s) to be provided.

Authority required

Initial treatment, as monotherapy, of a patient with acute lymphoblastic leukaemia bearing the Philadelphia chromosome or expressing the transcript, BCR-ABL, who has been treated prior to 1 December 2007 and has failed treatment with chemotherapy and where appropriate, allogeneic haemopoietic stem cell transplantation.

Patients must have active leukaemia, as defined by presence on current pathology assessments of either morphological infiltration of the bone marrow (greater than 5% lymphoblasts) or cerebrospinal fluid or other sites; OR the presence of cells bearing the Philadelphia chromosome on cytogenetic or FISH analysis in the bone marrow of patients in morphological remission.

The first authority application must be made in writing and must include:

- (a) a completed authority prescription form; and

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
(b) a completed Acute Lymphoblastic Leukaemia Dasatinib PBS Authority Application - Supporting Information Form; and (c) a signed patient acknowledgement; and (d) a pathology report demonstrating that the patient has active acute lymphoblastic leukaemia, either manifest as cytogenetic evidence of the Philadelphia chromosome, or morphological evidence of acute lymphoblastic leukaemia plus qualitative RT-PCR evidence of BCR-ABL transcript. The date of the relevant pathology report(s) need(s) to be provided.							
<u>Authority required</u>							
Continuing treatment, as monotherapy, of a patient with acute lymphoblastic leukaemia bearing the Philadelphia chromosome or expressing the transcript, BCR-ABL, where the patient has previously been issued with an authority prescription for dasatinib and does not have progressive disease.							
Authority applications for continuing treatment may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).							
<u>Note</u>							
Dasatinib will only be subsidised for patients with acute lymphoblastic leukaemia who are not receiving concomitant PBS-subsidised imatinib mesylate and who are not appropriate for an allogeneic haemopoietic stem cell transplant.							
<u>Note</u>							
No applications for increased repeats will be authorised.							
9125G	Tablet 20 mg	60	2	..	3246.36	34.20	Sprycel BQ
9126H	Tablet 50 mg	60	2	..	5250.68	34.20	Sprycel BQ
9127J	Tablet 70 mg	60	2	..	6465.01	34.20	Sprycel BQ
9343R	Tablet 100 mg	30	2	..	5250.68	34.20	Sprycel BQ

ERLOTINIB

Authority required

Initial PBS-subsidised treatment, as monotherapy, in a patient with locally advanced or metastatic (stage IIIB or IV) non-small cell lung cancer with a WHO performance status of 3 or less, after prior treatment with platinum-based chemotherapy, where:

- (1) (a) disease progression has occurred following treatment with docetaxel or pemetrexed; or
- (b) treatment with docetaxel and pemetrexed is either contraindicated or cannot be tolerated; and
- (2) further cytotoxic chemotherapy is not appropriate.

Authority required

Continuing PBS-subsidised treatment, as monotherapy, in a patient with locally advanced or metastatic (stage IIIB or IV) non-small cell lung cancer who has previously been issued with an authority prescription for this drug and who does not have progressive disease.

Authority required

Initial PBS-subsidised treatment, as monotherapy, in a patient with locally advanced or metastatic (stage IIIB or IV) non-small cell lung cancer, after prior treatment with platinum-based chemotherapy:

- (1) where:
 - (a) disease progression occurred following treatment with docetaxel or pemetrexed; or
 - (b) treatment with docetaxel and pemetrexed was either contraindicated or could not be tolerated; and
- (2) who has received treatment with erlotinib under the Erlotinib Access Programme prior to 1 August 2008; and
- (3) who does not have progressive disease.

Note

Special Pricing Arrangements apply.

9166K	Tablet 25 mg (as hydrochloride)	30	3	..	794.19	34.20	Tarceva RO
9167L	Tablet 100 mg (as hydrochloride)	30	3	..	2703.34	34.20	Tarceva RO
9168M	Tablet 150 mg (as hydrochloride)	30	3	..	3309.66	34.20	Tarceva RO

GEFITINIB

Note

Any queries concerning the arrangements to prescribe gefitinib may be directed to Medicare Australia on 1800 700 270.

Written applications for authority to prescribe gefitinib should be forwarded to:

Medicare Australia
 Prior Written Approval of Specialised Drugs
 Reply Paid 9826
 GPO Box 9826
 HOBART TAS 7001.

Authority required

Initial PBS-subsidised treatment, as monotherapy, of locally advanced or metastatic non-small cell lung cancer in patients with a WHO performance status of 2 or less, where:

- (1) disease progression has occurred following treatment with at least 1 chemotherapy agent; and
- (2) there is evidence that the patient has an activating mutation(s) of the epidermal growth factor receptor (EGFR) gene in tumour material.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Gefitinib (Iressa) PBS Authority Application for Use in the Treatment of Locally Advanced or Metastatic Non-Small Cell Lung Cancer - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] and
- (3) details of the prior chemotherapy including the name(s) of drug(s) and date of the most recent treatment cycle; and
- (4) details of the patient's WHO performance status; and
- (5) a copy of the pathology report providing evidence of the presence of activating mutation(s) of the EGFR gene from an Approved Pathology Authority.

Authority required

Continuing PBS-subsidised treatment, as monotherapy, of locally advanced or metastatic non-small cell lung cancer in patients with a WHO performance status of 2 or less, where the patient has previously been issued with an authority prescription for gefitinib.

Applications for continuing treatment may be made in writing or on the telephone by contacting Medicare Australia on 1800 700 270.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8769M	Tablet 250 mg	30	1	..	3851.36	34.20	Iressa	AP

IMATINIB

Note

Imatinib mesylate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Authority required

Initial PBS-subsidised treatment, for up to 3 months, of adult patients with a metastatic or unresectable malignant gastrointestinal stromal tumour which has been histologically confirmed by the detection of CD117 on immunohistochemical staining.

Patients must commence treatment at a dose not exceeding 400 mg per day for at least 3 months. Authority prescriptions for a higher dose will not be approved during this initial 3 month treatment period.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Imatinib Mesylate (Glivec) PBS Authority Application for Use in the Treatment of Gastrointestinal Stromal Tumour - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) a copy of a pathology report from an Approved Pathology Authority supporting the diagnosis of a gastrointestinal stromal tumour and confirming the presence of CD117 on immunohistochemical staining; and
 - (ii) a copy of the most recent (within 2 months of the application) computed tomography (CT) scan, magnetic resonance imaging (MRI) or ultrasound assessment of the tumour(s), including whether or not there is evidence of metastatic disease; and
 - (iii) where the application for authority to prescribe is being sought on the basis of an unresectable tumour, written evidence in support of that claim must be provided.

Authority required

Continuing PBS-subsidised treatment, at a dose of up to 600 mg per day, of adult patients with a metastatic or unresectable malignant gastrointestinal stromal tumour who have previously been issued with an authority prescription for this drug.

Applications for continuing treatment may be made by telephone (1800 700 270, hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients who have failed to respond or are intolerant to imatinib are no longer eligible to receive PBS-subsidised imatinib.

Note

Patients who achieve a response to treatment at an imatinib dose of 400 mg per day should be continued at this dose and assessed for response at regular intervals. Patients who fail to achieve a response to 400 mg per day may have their dose increased to 600 mg per day. Authority applications for doses higher than 600 mg per day will not be approved.

A response to treatment is defined as a decrease from baseline in the sum of the products of the perpendicular diameters of all measurable lesions of 50% or greater. (Response definition based on the Southwest Oncology Group standard criteria, see Demetri et al. N Engl J Med 2002; 347: 472-80.)

Note

No applications for increased repeats will be authorised.

9111M	Tablet 100 mg (as mesylate)	60	2	..	2032.53	34.20	Glivec	NV
9112N	Tablet 400 mg (as mesylate)	30	2	..	3918.69	34.20	Glivec	NV

IMATINIB

Note

Imatinib mesylate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Note

Any queries concerning the arrangements to prescribe imatinib mesylate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe imatinib mesylate should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

For the following diseases, written authority is required at initiation and for continuation:

Chronic myeloid leukaemia (chronic phase);
Dermatofibrosarcoma protuberans;
Hypereosinophilic syndrome;
Chronic eosinophilic leukaemia;
Myelodysplastic or myeloproliferative disorder;
Aggressive systemic mastocytosis with eosinophilia.

Authority required

Initial treatment of patients in the chronic phase of chronic myeloid leukaemia expressing the Philadelphia chromosome or the transcript, bcr-abl tyrosine kinase, and who have a primary diagnosis of chronic myeloid leukaemia.

Applications under this restriction will be limited to provide patients with a maximum of 18 months of therapy from the date the first application for initial treatment was approved.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Imatinib Mesylate (Glivec) PBS Authority Application for Use in the Treatment of Chronic Myeloid Leukaemia - Supporting Information form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of chronic myeloid leukaemia to confirm eligibility for treatment, or a qualitative PCR report documenting the presence of the bcr-abl transcript in either peripheral blood or bone marrow; and
- (4) a copy of a signed patient acknowledgement form indicating that the patient understands and acknowledges that PBS-subsidised treatment with imatinib mesylate for the chronic phase of chronic myeloid leukaemia will cease if subsequent testing demonstrates that:
 - (i) the patient has failed to achieve a major cytogenetic response within the initial 18 months of treatment [see Note defining major cytogenetic response]; or
 - (ii) the patient has failed to sustain a major cytogenetic response for 12 months from the date of the last pathology report that indicated that a major cytogenetic response had been achieved [see Note defining major cytogenetic response].

Note

Imatinib mesylate in the chronic phase of chronic myeloid leukaemia will only be subsidised for patients who are not receiving concomitant PBS-subsidised interferon alfa therapy.

Patients should be commenced on a dose of imatinib mesylate of 400 mg (base) daily and maintained on a minimum dose of imatinib mesylate of 400 mg (base) daily. Prescribing of lower doses should be carefully considered. Continuing therapy is dependent on patients demonstrating a response to imatinib mesylate therapy following the initial 18 months of treatment and at 12 monthly intervals thereafter, irrespective of the daily imatinib mesylate dose received.

Authority required

Continuing treatment of patients who have received initial treatment with imatinib mesylate as a pharmaceutical benefit for the chronic phase of chronic myeloid leukaemia and who have demonstrated either a major cytogenetic response or less than 1% bcr-abl level in the blood in the preceding 12 months.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) demonstration of continued response to treatment as evidenced by either:
 - (a) major cytogenetic response [see Note explaining requirements]. Where this has been supplied within the previous 12 months, only the date of the relevant pathology report need be provided; or
 - (b) a peripheral blood level of bcr-abl of less than 1% on the international scale [see Note explaining requirements]. Where this has been supplied within the previous 12 months, only the date of the relevant pathology report need be provided.

Note

Definitions of response.

A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells.

A peripheral blood bcr-abl level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the biological equivalent of a major cytogenetic response.

Authority approval requirements.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	For the purposes of assessing response to PBS-subsidised treatment with imatinib mesylate, either cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of bcr-abl transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with bcr-abl specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted as follows: (i) between 10 and 12 months of the commencement of treatment with imatinib mesylate, at which time patients in whom a major cytogenetic response or peripheral blood bcr-abl level of less than 1% has been demonstrated may receive authorisation for a further 12 months of treatment; and (ii) within 18 months of the commencement of treatment with imatinib mesylate, in patients who have failed to demonstrate a major cytogenetic response or peripheral blood bcr-abl level of less than 1% at between 10 and 12 months (patients in whom a major cytogenetic response or peripheral blood bcr-abl level of less than 1% is demonstrable by 18 months may also receive authorisation for a further 12 months of treatment); and (iii) at no greater than 12 month intervals thereafter, to demonstrate that the major cytogenetic response or peripheral blood bcr-abl level of less than 1% has been sustained. For each authority application where eligibility for continuing PBS-subsidised treatment is to be demonstrated, a copy of the cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or a copy of the quantitative PCR indicating the relative level of bcr-abl transcript in the peripheral blood using the international scale, must be submitted as described in (i) to (iii) above. For bone marrow analyses, where the standard karyotyping conducted at the time of application is not informative, a copy of a cytogenetic analysis conducted on the bone marrow using FISH with bcr-abl specific probe must be submitted with the authority application. A copy of the non-informative standard karyotype analysis must be included with the authority application. Where a patient has previously received PBS-subsidised treatment with imatinib mesylate, no approval will be granted for PBS-subsidised re-treatment in the chronic phase of chronic myeloid leukaemia, where that patient has at any time failed to meet the criteria for continuing treatment.						
	Note No applications for increased repeats will be authorised.						
9113P	Tablet 100 mg (as mesylate)	60	5	..	2032.53	34.20	Glivec NV
9114Q	Tablet 400 mg (as mesylate)	30	5	..	3918.69	34.20	Glivec NV

IMATINIB

Note

Imatinib mesylate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Note

Any queries concerning the arrangements to prescribe imatinib mesylate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe imatinib mesylate should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

For the following diseases, written authority is required at initiation and for continuation:

Chronic myeloid leukaemia (chronic phase);
Dermatofibrosarcoma protuberans;
Hypereosinophilic syndrome;
Chronic eosinophilic leukaemia;
Myelodysplastic or myeloproliferative disorder;
Aggressive systemic mastocytosis with eosinophilia.

Authority required

Treatment of patients in the accelerated phase of chronic myeloid leukaemia expressing the Philadelphia chromosome or the transcript, bcr-abl tyrosine kinase, and who have a primary diagnosis of chronic myeloid leukaemia. Progress to the accelerated phase is defined by the presence of 1 or more of the following:

- (1) Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 15% but less than 30%; or
- (2) Percentage of blasts plus promyelocytes in the peripheral blood or bone marrow greater than or equal to 30%; or
- (3) Peripheral basophils greater than or equal to 20%; or
- (4) Progressive splenomegaly to a size greater than or equal to 10 cm below the left costal margin to be confirmed on 2 occasions at least 4 weeks apart, or a greater than or equal to 50% increase in size below the left costal margin over 4 weeks; or
- (5) Karyotypic evolution (chromosomal abnormalities in addition to a single Philadelphia chromosome).

Applications for authorisation must be in writing and must include:

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	(a) a completed authority prescription form; and (b) a completed Imatinib Mesylate (Glivec) PBS Authority Application for Use in the Treatment of Chronic Myeloid Leukaemia - Supporting Information form, stating which of the above criteria are satisfied by the patient; and (c) a copy of the confirming pathology report from an Approved Pathology Authority in the case of criteria (1), (2), (3) and (5) above, or details of the dates of assessments in the case of progressive splenomegaly.						
	<u>Authority required</u> Treatment of patients in the blast phase of chronic myeloid leukaemia expressing the Philadelphia chromosome or the transcript, bcr-abl tyrosine kinase, and who have a primary diagnosis of chronic myeloid leukaemia. Progress to myeloid blast crisis is defined as either: (1) Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 30%; or (2) Extramedullary involvement other than spleen and liver.						
	Applications for authorisation must be in writing and must include: (a) a completed authority prescription form; and (b) a completed Imatinib Mesylate (Glivec) PBS Authority Application for Use in the Treatment of Chronic Myeloid Leukaemia - Supporting Information form, stating which of the above criteria are satisfied by the patient; and (c) a copy of the confirming pathology report from an Approved Pathology Authority in the case of criterion (1) above, or details of the date of assessment in the case of extramedullary involvement.						
	<u>Authority required</u> Continuing treatment of patients with chronic myeloid leukaemia expressing the Philadelphia chromosome or the transcript, bcr-abl tyrosine kinase, where the patient has previously received PBS-subsidised treatment with imatinib mesylate of: (i) the accelerated phase of chronic myeloid leukaemia; or (ii) the blast phase of chronic myeloid leukaemia.						
	<u>Note</u> No applications for increased repeats will be authorised.						
9115R	Tablet 100 mg (as mesylate)	60	2	..	2032.53	34.20	Glivec NV
9116T	Tablet 400 mg (as mesylate)	30	2	..	3918.69	34.20	Glivec NV

IMATINIB

Note

Imatinib mesylate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Note

Any queries concerning the arrangements to prescribe imatinib mesylate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe imatinib mesylate should be forwarded to:

Medicare Australia
 Prior Written Approval of Specialised Drugs
 Reply Paid 9826
 GPO Box 9826
 HOBART TAS 7001

For the following diseases, written authority is required at initiation and for continuation:

Chronic myeloid leukaemia (chronic phase);
 Dermatofibrosarcoma protuberans;
 Hypereosinophilic syndrome;
 Chronic eosinophilic leukaemia;
 Myelodysplastic or myeloproliferative disorder;
 Aggressive systemic mastocytosis with eosinophilia.

Authority required

Initial treatment in combination with chemotherapy as induction or consolidation of a newly diagnosed patient with acute lymphoblastic leukaemia (ALL) bearing the Philadelphia chromosome or expressing the transcript, BCR-ABL.

The first authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Acute Lymphoblastic Leukaemia Imatinib PBS Authority Application - Supporting Information Form; and
- (c) a pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of acute lymphoblastic leukaemia to confirm eligibility for treatment, with either cytogenetic evidence of the Philadelphia chromosome, or a qualitative PCR report documenting the presence of the BCR-ABL transcript in either peripheral blood or bone marrow. (The date of the relevant pathology report needs to be provided); and
- (d) a signed patient acknowledgement.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Authority required

Initial treatment of a patient with acute lymphoblastic leukaemia bearing the Philadelphia chromosome or expressing the transcript BCR-ABL who was previously treated with imatinib mesylate under the Imatinib Compassionate Program and who meets all the PBS criteria.

The first authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Acute Lymphoblastic Leukaemia Imatinib PBS Authority Application - Supporting Information Form; and
- (c) a pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of acute lymphoblastic leukaemia to confirm eligibility for treatment, with either cytogenetic evidence of the Philadelphia chromosome, or a qualitative PCR report documenting the presence of the BCR-ABL transcript in either peripheral blood or bone marrow. (The date of the relevant pathology report needs to be provided); and
- (d) a signed patient acknowledgement.

Authority required

Continuing treatment in combination with chemotherapy as maintenance of first complete remission of patients with acute lymphoblastic leukaemia bearing the Philadelphia chromosome or expressing the transcript, BCR-ABL.

Authority applications for continuing treatment may be made by telephone to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Imatinib mesylate is available with a lifetime maximum of 24 months for continuing treatment with imatinib mesylate therapy for patients with acute lymphoblastic leukaemia reimbursed through the PBS.

Any queries concerning the arrangements to prescribe imatinib mesylate beyond 24 months may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note

Allogeneic stem cell transplantation is the preferred therapy for eligible patients achieving a complete remission of Philadelphia positive acute lymphoblastic leukaemia.

Note

No applications for increased repeats will be authorised.

9123E	Tablet 100 mg (as mesylate)	60	2	..	2032.53	34.20	Glivec	NV
9124F	Tablet 400 mg (as mesylate)	30	2	..	3918.69	34.20	Glivec	NV

IMATINIB

Note

Imatinib mesylate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Note

Any queries concerning the arrangements to prescribe imatinib mesylate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe imatinib mesylate should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

For the following diseases, written authority is required at initiation and for continuation:

- Chronic myeloid leukaemia (chronic phase);
- Dermatofibrosarcoma protuberans;
- Hypereosinophilic syndrome;
- Chronic eosinophilic leukaemia;
- Myelodysplastic or myeloproliferative disorder;
- Aggressive systemic mastocytosis with eosinophilia.

Authority required

Initial PBS-subsidised treatment of a patient with unresectable, locally recurrent or metastatic dermatofibrosarcoma protuberans.

Maximum dose: 800 mg per day.

- (1) Where the application for authority to prescribe is being sought on the basis of unresectable tumour, written evidence in support of that claim must be provided; and
- (2) Where the application for authority to prescribe is being sought on the basis of locally recurrent disease, the site of the local recurrence must be

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	specified; and (3) Where the application for authority to prescribe is being sought on the basis of metastatic disease, the site(s) of metastatic disease must be provided.						
	Applications for authorisation for initial treatment must be made in writing and must include: (a) a completed authority prescription form; and (b) a completed Rare Diseases Imatinib PBS Authority Application - Supporting Information Form; and (c) a signed patient acknowledgement.						
	Authority required Continuing PBS-subsidised treatment of a patient with unresectable, locally recurrent or metastatic dermatofibrosarcoma protuberans who has previously been issued with an authority prescription for imatinib and who has demonstrated a response, but whose disease remains unresectable.						
	Maximum dose: 800 mg per day.						
	Applications for authorisation must be made in writing and must include: (a) a completed authority prescription form; and (b) a completed Rare Diseases Imatinib PBS Authority Application - Supporting Information Form; and (c) a statement that the disease has not progressed on imatinib therapy.						
	Note No applications for increased repeats will be authorised.						
9172R	Tablet 100 mg (as mesylate)	60	2	..	2032.53	34.20	Glivec NV
9173T	Tablet 400 mg (as mesylate)	30	2	..	3918.69	34.20	Glivec NV

IMATINIB

Note

Imatinib mesylate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Note

Any queries concerning the arrangements to prescribe imatinib mesylate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe imatinib mesylate should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

For the following diseases, written authority is required at initiation and for continuation:

Chronic myeloid leukaemia (chronic phase);
Dermatofibrosarcoma protuberans;
Hypereosinophilic syndrome;
Chronic eosinophilic leukaemia;
Myelodysplastic or myeloproliferative disorder;
Aggressive systemic mastocytosis with eosinophilia.

Authority required

Initial PBS-subsidised treatment of a patient with hypereosinophilic syndrome or chronic eosinophilic leukaemia requiring treatment and confirmed to carry the FIP1L1-PDGFR fusion gene.

Maximum dose: 400 mg per day.

Applications for authorisation for initial treatment must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Rare Diseases Imatinib PBS Authority Application - Supporting Information Form; and
- (c) a copy of the pathology report confirming the presence of the FIP1L1-PDGFR fusion gene; and
- (d) a copy of the full blood examination report confirming the presence of hypereosinophilic syndrome or chronic eosinophilic leukaemia; and
- (e) details of organ involvement requiring treatment, including a copy of the radiology, nuclear medicine, respiratory function or anatomical pathology reports as appropriate; and
- (f) a signed patient acknowledgement.

Authority required

Continuing PBS-subsidised treatment of a patient with hypereosinophilic syndrome or chronic eosinophilic leukaemia who has previously been issued with an authority prescription for imatinib and who has achieved and maintained a complete haematological response.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Maximum dose: 400 mg per day.							
Applications for authorisation must be made in writing and must include:							
(a) a completed authority prescription form; and							
(b) a completed Rare Diseases Imatinib PBS Authority Application - Supporting Information Form; and							
(c) a copy of the full blood examination report which demonstrates a complete haematological response, with a normal eosinophil count; and							
(d) a statement that the disease has not progressed on imatinib therapy.							
<u>Note</u>							
No applications for increased repeats will be authorised.							
9174W	Tablet 100 mg (as mesylate)	60	2	..	2032.53	34.20	Glivec NV
9175X	Tablet 400 mg (as mesylate)	30	2	..	3918.69	34.20	Glivec NV

IMATINIB

Note

Imatinib mesylate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Note

Any queries concerning the arrangements to prescribe imatinib mesylate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe imatinib mesylate should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

For the following diseases, written authority is required at initiation and for continuation:

Chronic myeloid leukaemia (chronic phase);
Dermatofibrosarcoma protuberans;
Hypereosinophilic syndrome;
Chronic eosinophilic leukaemia;
Myelodysplastic or myeloproliferative disorder;
Aggressive systemic mastocytosis with eosinophilia.

Authority required

Initial PBS-subsidised treatment of a patient with a myelodysplastic or myeloproliferative disorder where:

- (1) there is confirmed evidence of a platelet-derived growth factor receptor (PDGFR) gene re-arrangement either by standard karyotyping, or FISH or PDGFRB fusion gene transcript; and
- (2) the patient has previously failed an adequate trial of one or more of the following conventional therapies:
 - cytarabine;
 - etoposide;
 - hydroxyurea.

Maximum dose: 400 mg per day.

Applications for authorisation for initial treatment must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Rare Diseases Imatinib PBS Authority Application - Supporting Information Form; and
- (c) a copy of the pathology report confirming the platelet-derived growth factor receptor (PDGFR) gene re-arrangement; and
- (d) a copy of the bone marrow biopsy report which demonstrates the presence of a myelodysplastic or myeloproliferative disorder; and
- (e) details of the prior therapy trialled and the response; and
- (f) a signed patient acknowledgement.

Authority required

Continuing PBS-subsidised treatment of a patient with a PDGFRB fusion gene-positive myelodysplastic or myeloproliferative disorder who has previously been issued with an authority prescription for imatinib and who has demonstrated a complete haematological response.

Maximum dose: 400 mg per day.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Rare Diseases Imatinib PBS Authority Application - Supporting Information Form; and

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
(c) a copy of the full blood examination report which demonstrates a complete haematological response; and (d) a statement that the disease has not progressed on imatinib therapy.							
<u>Note</u> No applications for increased repeats will be authorised.							
9176Y	Tablet 100 mg (as mesylate)	60	2	..	2032.53	34.20	Glivec NV
9177B	Tablet 400 mg (as mesylate)	30	2	..	3918.69	34.20	Glivec NV

IMATINIB

Note

Imatinib mesylate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Note

Any queries concerning the arrangements to prescribe imatinib mesylate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe imatinib mesylate should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

For the following diseases, written authority is required at initiation and for continuation:

Chronic myeloid leukaemia (chronic phase);
Dermatofibrosarcoma protuberans;
Hypereosinophilic syndrome;
Chronic eosinophilic leukaemia;
Myelodysplastic or myeloproliferative disorder;
Aggressive systemic mastocytosis with eosinophilia.

Authority required

Initial PBS-subsidised treatment of a patient with aggressive systemic mastocytosis with eosinophilia where:

- (1) there is confirmed evidence of the FIP1L1-PDGFR fusion gene; and
- (2) the patient has previously failed an adequate trial of one or more of the following conventional therapies:
 - corticosteroids;
 - hydroxyurea.

Maximum dose: 400 mg per day.

Applications for authorisation for initial treatment must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Rare Diseases Imatinib PBS Authority Application - Supporting Information Form; and
- (c) a copy of the pathology report confirming the presence of the FIP1L1-PDGFR fusion gene; and
- (d) a copy of the bone marrow biopsy report and/or other tissue biopsy report confirming the diagnosis of aggressive systemic mastocytosis and a copy of the full blood examination report demonstrating eosinophilia; and
- (e) details of symptomatic organ involvement requiring treatment, including a copy of the radiology, nuclear medicine, respiratory function or anatomical pathology reports as appropriate; and
- (f) details of prior treatment trialled and the response; and
- (g) a signed patient acknowledgement.

Authority required

Continuing PBS-subsidised treatment of a patient with aggressive systemic mastocytosis confirmed to carry the FIP1L1-PDGFR fusion gene, who has previously been issued with an authority prescription for imatinib and who has demonstrated a complete haematological response.

Maximum dose: 400 mg per day.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Rare Diseases Imatinib PBS Authority Application - Supporting Information Form; and
- (c) a copy of the full blood examination report which demonstrates a complete haematological response; and
- (d) a statement that the disease has not progressed on imatinib therapy.

Note

No applications for increased repeats will be authorised.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
9178C	Tablet 100 mg (as mesylate)	60	2	..	2032.53	34.20	Glivec	NV
9179D	Tablet 400 mg (as mesylate)	30	2	..	3918.69	34.20	Glivec	NV

LAPATINIB

Note

Any queries concerning the arrangements to prescribe lapatinib may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Lapatinib should not be used in patients with a left ventricular ejection fraction (LVEF) of less than 45% or with symptomatic heart failure. Cardiac function must be tested by a suitable method including, for example, ECHO or MUGA, prior to seeking the initial authority approval and then at 3 monthly intervals during treatment.

Lapatinib is not PBS-subsidised when used in combination with Commonwealth-subsidised trastuzumab.

If disease progression occurs, the prescribing doctor must contact Medicare Australia within one week on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) and lapatinib treatment must be ceased immediately.

Authority required

Initial treatment, in combination with capecitabine, of a patient with HER2 positive metastatic breast cancer (equivalent to Stage IIIC or Stage IV) who has received prior therapy with a taxane, for at least 3 cycles, and whose disease has progressed despite treatment with trastuzumab for metastatic disease.

Authority applications for initial treatment must be made in writing and must include:

- (a) a completed authority prescription form;
- (b) a pathology report demonstrating HER2 positivity has been demonstrated by in situ hybridisation (ISH);
- (c) date of last treatment with a taxane and total number of cycles;
- (d) a signed patient acknowledgment;
- (e) dates of treatment with trastuzumab; and
- (f) date of demonstration of progression whilst on treatment with trastuzumab.

Note

Treatment with trastuzumab for metastatic disease is defined as trastuzumab administered alone or in combination with chemotherapy for at least 6 weeks at standard doses.

If treatment with a taxane is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities, including severity, can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

Authority required

Continuing treatment, in combination with capecitabine, of a patient with HER2 positive metastatic breast cancer who has previously received treatment with PBS-subsidised lapatinib and who does not have progressive disease.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a statement from the prescribing doctor that the disease has not progressed.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9148L	Tablet 250 mg (as ditosylate monohydrate)	140	2	..	*3387.46	34.20	Tykerb	GK

NILOTINIB

Note

Any queries concerning the arrangements to prescribe nilotinib may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Any queries concerning patients who are enrolled on the Nilotinib Compassionate Program may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Applications for authority to prescribe nilotinib should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	GPO Box 9826 HOBART TAS 7001.						
	Authority required						
	Initial treatment, as the sole PBS-subsidised therapy, of a patient with chronic myeloid leukaemia in chronic or accelerated phase bearing the Philadelphia chromosome or expressing the transcript, BCR-ABL, who has active leukaemia (as defined by presence on current pathology assessments of either the Philadelphia chromosome on cytogenetic or FISH analysis, or the presence of the transcript BCR-ABL greater than 1% on the international scale) and who has failed an adequate trial of imatinib.						
	Failure of an adequate trial of imatinib is defined as:						
	(i) Lack of response to initial imatinib therapy, defined as either:						
	— failure to achieve a haematological response after a minimum of 3 months therapy with imatinib for patients initially treated in chronic phase; or						
	— failure to achieve any cytogenetic response after a minimum of 6 months therapy with imatinib for patients initially treated in chronic phase as demonstrated on bone marrow biopsy by presence of greater than 95% Philadelphia chromosome positive cells; or						
	— failure to achieve a major cytogenetic response or a peripheral blood BCR-ABL level of less than 1% after a minimum of 12 months therapy with imatinib; OR						
	(ii) Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing imatinib therapy; OR						
	(iii) Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing in value by at least 5 fold to a level of greater than 1% confirmed on a subsequent test), during ongoing imatinib therapy; OR						
	(iv) Development of accelerated phase in a patient previously prescribed imatinib for the chronic phase of chronic myeloid leukaemia.						
	Accelerated phase is defined by the presence of 1 or more of the following:						
	(1) Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 15% but less than 30%; or						
	(2) Percentage of blasts plus promyelocytes in the peripheral blood or bone marrow greater than or equal to 30%, provided that blast count is less than 30%; or						
	(3) Peripheral basophils greater than or equal to 20%; or						
	(4) Progressive splenomegaly to a size greater than or equal to 10 cm below the left costal margin to be confirmed on 2 occasions at least 4 weeks apart, or a greater than or equal to 50% increase in size below the left costal margin over 4 weeks; or						
	(5) Karyotypic evolution (chromosomal abnormalities in addition to a single Philadelphia chromosome); OR						
	(v) Disease progression (defined as a greater than or equal to 50% increase in peripheral white blood cell count, blast count, basophils or platelets) during first-line imatinib therapy in patients with accelerated phase chronic myeloid leukaemia, provided that blast crisis has been excluded on bone marrow biopsy; OR						
	(vi) Grade 3 or 4 non-haematological toxicity that is imatinib related and necessitates permanent cessation of imatinib. For patients with imatinib related toxicities, leukaemia activity does not need to be demonstrated.						
	Applications for authorisation must be in writing and must include:						
	(a) a completed authority prescription form; and						
	(b) a completed Chronic Myeloid Leukaemia Dasatinib/Nilotinib PBS Authority Application - Supporting Information Form; and						
	(c) a signed patient acknowledgement; and						
	(d) a bone marrow biopsy pathology report demonstrating the patient has active chronic myeloid leukaemia, either manifest as cytogenetic evidence of the Philadelphia chromosome, or RT-PCR level of BCR-ABL transcript greater than 1% on the international scale. (The date of the relevant pathology report needs to be provided); and						
	(e) where there has been a loss of response to imatinib, a copy of the current confirming pathology report(s) from an Approved Pathology Authority; or						
	(f) details of Grade 3 or 4 non-haematological imatinib related toxicity.						
	Note						
	Nilotinib will only be subsidised for patients with chronic myeloid leukaemia who are not receiving concomitant PBS-subsidised imatinib mesylate, dasatinib or interferon alfa therapy.						
	Patients should be commenced on a dose of nilotinib of 400 mg twice daily. Continuing therapy is dependent on patients demonstrating a major cytogenetic response to nilotinib therapy or a peripheral blood BCR-ABL level of less than 1% at 18 months and thereafter at 12 monthly intervals, irrespective of the daily nilotinib dose received.						
	Nilotinib is not PBS-subsidised for patients with CML that is resistant to dasatinib.						
	Nilotinib is not TGA-registered and not PBS-subsidised for patients with CML in blast crisis.						
	Requests for doses of greater than nilotinib 400 mg twice daily will not be approved.						
	From 1 November 2008, under the PBS, a patient will be able to trial either dasatinib and/or nilotinib within the initial 18 month treatment period, providing the patient's CML is not resistant to the first second-line agent.						
9285Q	Capsule 200 mg (as hydrochloride monohydrate)	112	2	..	*5490.42	34.20	Tasigna NV

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
NILOTINIB							
<u>Note</u>							
Any queries concerning the arrangements to prescribe nilotinib may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).							
Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au .							
Any queries concerning patients who are enrolled on the Nilotinib Compassionate Program may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).							
Applications for authority to prescribe nilotinib should be forwarded to:							
Medicare Australia Prior Written Approval of Specialised Drugs Reply Paid 9826 GPO Box 9826 HOBART TAS 7001.							
<u>Authority required</u>							
Continuing treatment, as the sole PBS-subsidised therapy, of a patient who has received initial treatment with nilotinib as a pharmaceutical benefit for chronic myeloid leukaemia, and who has demonstrated either a major cytogenetic response, or less than 1% BCR-ABL level in the blood, to nilotinib in the preceding 18 months and thereafter at 12 monthly intervals.							
Applications for authorisation must be in writing and must include:							
(1) a completed authority prescription form; and							
(2) a completed Chronic Myeloid Leukaemia Dasatinib/Nilotinib Authority Application Form for continuing treatment; and							
(3) demonstration of continued response to treatment as evidenced by either:							
(a) major cytogenetic response [see Note explaining definitions of response]. Where this has been supplied within the previous 12 months (or 18 months for the initial supply), only the date of the relevant pathology report needs to be provided; or							
(b) a peripheral blood level of BCR-ABL of less than 1% on the international scale [see Note explaining definitions of response]. Where this has been supplied within the previous 12 months (or 18 months for the initial supply), only the date of the relevant pathology report needs to be provided.							
<u>Note</u>							
Definitions of response.							
A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells.							
A bone marrow or peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the biological equivalent of a major cytogenetic response.							
Authority approval requirements.							
For the purposes of assessing response to PBS-subsidised treatment with nilotinib, either cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted as follows:							
(i) between 10 and 18 months of the commencement of treatment with nilotinib, at which time patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% has been demonstrated may receive authorisation for a further 12 months of treatment; and							
(ii) at no greater than 12 month intervals thereafter, to demonstrate that the major cytogenetic response or peripheral blood BCR-ABL level of less than 1% has been sustained.							
For each authority application where eligibility for continuing PBS-subsidised treatment is to be demonstrated, a copy of the cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or a copy of the quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted as described in (i) and (ii) above. For bone marrow analyses, where the standard karyotyping conducted at the time of application is not informative, a copy of a cytogenetic analysis conducted on the bone marrow using FISH with BCR-ABL specific probe must be submitted with the authority application. A copy of the non-informative standard karyotype analysis must be included with the authority application.							
Where a patient has previously received PBS-subsidised treatment with nilotinib, no approval will be granted for PBS-subsidised re-treatment where that patient has at any time failed to meet the criteria for continuing treatment.							
9171Q	Capsule 200 mg (as hydrochloride monohydrate)	112	5	..	*5490.42	34.20	Tasigna NV

SORAFENIB

Authority required

Initial treatment, as the sole PBS-subsidised agent, of advanced (BCLC Stage C) hepatocellular carcinoma in a patient with a WHO performance status of 2 or less and Child Pugh class A;

Continuing treatment, as the sole PBS-subsidised agent, of advanced hepatocellular carcinoma in a patient who has previously been treated with PBS-subsidised sorafenib and who does not have progressive disease.

Note

Sorafenib is not PBS-subsidised for adjunctive treatment after resection, ablation or chemoembolization.
Sorafenib is not PBS-subsidised for maintenance therapy after disease progression.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Special Pricing Arrangements apply.

9380Q	Tablet 200 mg (as tosylate)	120	2	..	*6457.08	34.20	Nexavar	BN

SUNITINIB

Authority required

Initial treatment, as the sole PBS-subsidised therapy, of Stage IV clear cell variant renal cell carcinoma (RCC) in a patient who meets the Memorial Sloan Kettering Cancer Centre (MSKCC) low to intermediate risk group and has a WHO performance status of 2 or less.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Special Pricing Arrangements apply.

9417P	Capsule 12.5 mg (as malate)	28	1	..	1834.20	34.20	Sutent	PF
9418Q	Capsule 25 mg (as malate)	28	1	..	3521.76	34.20	Sutent	PF
9419R	Capsule 50 mg (as malate)	28	1	..	6897.44	34.20	Sutent	PF

SUNITINIB

Authority required

Continuing treatment beyond 3 months, as the sole PBS-subsidised therapy, of Stage IV clear cell variant renal cell carcinoma (RCC) in a patient who has previously been issued with an authority prescription for sunitinib and who has stable or responding disease according to RECIST criteria.

Note

RECIST Criteria is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Authority required

Initial treatment, as the sole PBS-subsidised therapy, of Stage IV clear cell variant renal cell carcinoma (RCC) in a patient who was receiving treatment with sunitinib prior to 1 May 2009.

Note

Special Pricing Arrangements apply.

9420T	Capsule 12.5 mg (as malate)	28	3	..	1834.20	34.20	Sutent	PF
9421W	Capsule 25 mg (as malate)	28	3	..	3521.76	34.20	Sutent	PF
9422X	Capsule 50 mg (as malate)	28	3	..	6897.44	34.20	Sutent	PF

SUNITINIB

Authority required

Initial PBS-subsidised treatment as monotherapy of a patient with WHO performance status of 2 or less with a metastatic or unresectable malignant gastrointestinal stromal tumour after failure of imatinib mesylate treatment due to resistance or intolerance.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Sunitinib Malate (Sutent) PBS Authority Application for Use in the Treatment of Gastrointestinal Stromal Tumour - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

Patients who have failed to respond or are intolerant to imatinib are no longer eligible to receive PBS-subsidised imatinib.

Note

Any queries concerning the arrangements to prescribe sunitinib malate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Any queries concerning patients who are enrolled on the Sunitinib Compassionate Program may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Written applications for authority to prescribe sunitinib malate should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Sunitinib malate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Authority required

Continuing PBS-subsidised treatment as monotherapy of a patient with WHO performance status of 2 or less with a metastatic or unresectable malignant gastrointestinal stromal tumour who has previously been issued with an authority prescription for sunitinib and who does not have progressive disease.

Applications for continuing treatment may be made by telephone (1800 700 270, hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients who have failed to respond or who are intolerant to imatinib are no longer eligible to receive PBS-subsidised imatinib.

Patients who have progressive disease on sunitinib are no longer eligible for PBS-subsidised sunitinib.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Special Pricing Arrangements apply.

9488J	Capsule 12.5 mg (as malate)	28	1	..	1834.20	34.20	Sutent	PF
9489K	Capsule 25 mg (as malate)	28	1	..	3521.76	34.20	Sutent	PF
9490L	Capsule 50 mg (as malate)	28	1	..	6897.44	34.20	Sutent	PF

Other antineoplastic agents

ARSENIC TRIOXIDE

Authority required

Induction and consolidation treatment of relapsed acute promyelocytic leukaemia (characterised by the presence of the t(15:17) translocation or PML/RAR-alpha fusion gene transcript) in a patient who is arsenic naive at induction.

9453M	Injection concentrate 10 mg in 10 mL	60	2	..	*24196.0 8	34.20	Phenasen	PL

BORTEZOMIB

Note

Any queries concerning the arrangements to prescribe bortezomib may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Applications for authority to prescribe bortezomib should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001.

Authority required

Initial PBS-subsidised treatment, as monotherapy or in combination with a corticosteroid and/or cyclophosphamide, of a patient with a histological diagnosis of multiple myeloma who has progressive disease after at least 1 prior therapy and who has undergone or is ineligible for a primary stem cell transplant. The patient must have experienced treatment failure after a trial of at least four (4) weeks of thalidomide at a dose of at least 100 mg daily or have failed to achieve at least a minimal response after eight (8) or more weeks of thalidomide-based therapy for progressive disease.

If the dosing requirement for thalidomide cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
 (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
 (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
 (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein and less than 200 mg per 24 hour Bence-Jones proteinuria.

Thalidomide treatment failure is defined as:

- (1) confirmed disease progression during thalidomide treatment or within 6 months of discontinuing thalidomide treatment; or
- (2) severe intolerance or toxicity unresponsive to clinically appropriate dose adjustment.

Severe intolerance due to thalidomide is defined as unacceptable somnolence or sedation interfering with activities of daily living.

Toxicity from thalidomide is defined as peripheral neuropathy (Grade 2 or greater, interfering with function), drug-related seizures, serious Grade 3 or Grade 4 drug-related dermatological reactions, such as Stevens-Johnson Syndrome, or other Grade 3 or 4 toxicity.

Any queries concerning additional details about treatment failure may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Failure to achieve at least a minimal response after 8 or more weeks of thalidomide-based therapy for progressive disease is defined as:

- (1) less than a 25% reduction in serum or urine M protein; or
- (2) in oligo-secretory and non-secretory myeloma patients only, less than a 25% reduction in the difference between involved and uninvolved serum free light chain levels.

Bortezomib will only be subsidised for patients with multiple myeloma who are not receiving concomitant PBS-subsidised lenalidomide.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Multiple Myeloma Authority Application - Supporting Information Form, which includes details of the histological diagnosis of multiple myeloma, prior treatments including name(s) of drug(s) and date of most recent treatment cycle and record of prior stem cell transplant or ineligibility for prior stem cell transplant; details of thalidomide treatment failure; details of the basis of the diagnosis of progressive disease or failure to respond; and nomination of which disease activity parameters will be used to assess response.

To enable confirmation by Medicare Australia, current diagnostic reports of at least one of the following are required:

- (a) the level of serum monoclonal protein; or
- (b) Bence-Jones proteinuria — the results of 24-hour urinary light chain M protein excretion; or
- (c) the serum level of free kappa and lambda light chains; or
- (d) bone marrow aspirate or trephine; or
- (e) if present, the size and location of lytic bone lesions (not including compression fractures); or
- (f) if present, the size and location of all soft tissue plasmacytomas by clinical or radiographic examination i.e. MRI or CT-scan; or
- (g) if present, the level of hypercalcaemia, corrected for albumin concentration.

As these parameters will be used to determine response, results for either (a) or (b) or (c) should be provided for all patients. Where the patient has oligo-secretory or non-secretory multiple myeloma, either (c) or (d) or if relevant (e), (f) or (g) should be provided. Where the prescriber plans to assess response in patients with oligo-secretory or non-secretory multiple myeloma with free light chain assays, evidence of the oligo-secretory or non-secretory nature of the multiple myeloma (either previous or current serum M protein less than 10 g per L and urinary Bence-Jones protein undetectable or less than 200 mg per 24 hours) must be provided; and

- (3) duration of thalidomide and daily dose prescribed; and
- (4) a signed patient acknowledgment.

Authority required

Continuing PBS-subsidised treatment, as monotherapy or in combination with a corticosteroid and/or cyclophosphamide, of multiple myeloma in a patient who has previously received 4 treatment cycles of bortezomib and who, at the time of application, has demonstrated at least a partial response to bortezomib.

If serum M protein and urine Bence-Jones protein levels are measurable, partial response (PR) compared with baseline (prior to treatment with bortezomib) is defined as:

- (a) at least a 50% reduction in the level of serum M protein (monoclonal protein); or
- (b) at least a 90% reduction in 24-hour urinary light chain M protein excretion or to less than 200 mg per 24 hours.

If serum M protein and urine Bence-Jones protein levels are unmeasurable as in non-secretory/oligo-secretory multiple myeloma, partial response compared with baseline is defined as:

- (c) at least a 50% reduction in the difference between involved and uninvolved serum free light chain (FLC) levels.

If serum M protein and urine Bence-Jones protein and serum FLC are unmeasurable/unavailable, partial response compared with baseline is defined as:

- (d) at least a 50% reduction in bone marrow plasma cells; or
- (e) no increase in size or number of lytic bone lesions (development of compression fracture does not exclude response); or
- (f) at least a 50% reduction in the size of soft tissue plasmacytoma (by clinical or applicable radiographic examination, i.e. MRI or CT-Scan); or

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	(g) normalisation of corrected serum calcium to less than or equal to 2.65 mmol per L.						

For the purpose of assessing eligibility for continuing PBS-subsidised bortezomib treatment beyond 4 cycles, the patient must have achieved at least a partial response at the completion of cycle 4. The results of the response assessment must be included in a written application to Medicare Australia for further treatment. Where a response assessment is not submitted to Medicare Australia prior to cycle 5, patients will be deemed to have failed to respond to treatment with bortezomib. Continuing PBS-subsidised supply will not be approved if there is a gap of more than 6 months between the initial application and subsequent applications.

The same parameters provided for the diagnosis of progressive disease are to be used to demonstrate at least a partial response to treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Multiple Myeloma Authority Application - Supporting Information Form; and
- (3) diagnostic reports demonstrating the patient has achieved at least a partial response.

Diagnostic reports must be no more than 1 month old at the time of application.

Patients who fail to demonstrate at least a partial response after 8 cycles will not be eligible to receive further PBS-subsidised treatment with bortezomib.

No more than 2 cycles of treatment beyond the cycle at which a confirmed complete response was first achieved will be authorised. Confirmation requires 2 determinations a minimum of 6 weeks apart.

Note

Special Pricing Arrangements apply.

9117W	Powder for injection 3.5 mg (solvent required) (code 7086Y applies to above item with approved solvent)	4	3	..	*7002.38	34.20	Velcade	JC

BORTEZOMIB

Note

Any queries concerning the arrangements to prescribe bortezomib may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Applications for authority to prescribe bortezomib should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001.

Authority required

Continuing PBS-subsidised treatment, as monotherapy or in combination with a corticosteroid and/or cyclophosphamide, of multiple myeloma in a patient who has previously received 8 treatment cycles with bortezomib and who, at the time of application, has demonstrated at least a partial response to bortezomib but who has not received 2 treatment cycles after first achieving a confirmed complete response.

If serum M protein and urine Bence-Jones protein levels are measurable, partial response (PR) compared with baseline (prior to treatment with bortezomib) is defined as:

- (a) at least a 50% reduction in the level of serum M protein (monoclonal protein); or
- (b) at least a 90% reduction in 24-hour urinary light chain M protein excretion or to less than 200 mg per 24 hours.

If serum M protein and urine Bence-Jones protein levels are unmeasurable as in non-secretory/oligo-secretory multiple myeloma, partial response compared with baseline is defined as:

- (c) the difference between involved and uninvolved serum free light chain (FLC) levels, with at least a 50% reduction in this value.

If serum M protein and urine Bence-Jones protein levels and serum FLC are unmeasurable/unavailable, partial response compared with baseline is defined as:

- (d) at least a 50% reduction in bone marrow plasma cells; or
- (e) no increase in size or number of lytic bone lesions (development of compression fracture does not exclude response); or
- (f) at least a 50% reduction in the size of soft tissue plasmacytoma (by clinical or applicable radiographic examination, i.e. MRI or CT-Scan); or
- (g) normalisation of corrected serum calcium to less than or equal to 2.65 mmol per L.

The same parameters provided for the diagnosis of progressive disease are to be used to demonstrate at least a partial response to treatment.

Diagnostic reports must be within 1 month of the date of application.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty	Safety Net	
	\$	\$					
	For the purpose of assessing eligibility for continuing PBS-subsidised bortezomib treatment beyond 8 cycles, the patient must have achieved at least a partial response at the completion of cycle 8. The results of the response assessment must be included in a written application to Medicare Australia for further treatment. Where a response assessment is not submitted to Medicare Australia prior to cycle 9, patients will be deemed to have failed to respond to treatment with bortezomib. Continuing PBS-subsidised supply will not be approved if there is a gap of more than 10 months between the initial application and an application following completion of 8 treatment cycles.						
	The authority application must be made in writing and must include:						
	(1) a completed authority prescription form; and						
	(2) a completed Multiple Myeloma Authority Application - Supporting Information Form; and						
	(3) diagnostic reports demonstrating the patient has achieved at least a partial response.						
	No more than 2 cycles of treatment beyond the cycle at which the complete response was first achieved will be authorised. Confirmation requires 2 determinations a minimum of 6 weeks apart.						
	Applications for PBS-subsidised treatment with bortezomib that extends beyond 11 cycles will not be approved.						
	Note						
	Special Pricing Arrangements apply.						
9118X	Powder for injection 3.5 mg (solvent required) (code 7087B applies to above item with approved solvent)	4	2	..	*7002.38	34.20	Velcade JC
HYDROXYUREA							
3093T	Capsule 500 mg	100	..	..	76.46	34.20	Hydrea BQ
IRINOTECAN HYDROCHLORIDE TRIHYDRATE							
Authority required (STREAMLINED)							
3184							
Metastatic colorectal cancer in patients with a WHO performance status of 2 or less.							
Note							
In first-line usage, effectiveness and tolerance may be improved when irinotecan is combined with an infusional 5-fluorouracil regimen.							
8414W	I.V. injection 40 mg in 2 mL	1	3	..	137.91	34.20	^a Camptosar PF ^a Hospira Pty Limited HH ^a Irinotecan Actavis GQ ^a Irinotecan Alphapharm AF ^a Irinotecan Ebewe IT ^a Irinotecan Sandoz SZ ^a Omegapharm OE ^a Irinotecan Tecan WQ
8415X	I.V. injection 100 mg in 5 mL	2	3	..	*628.02	34.20	^a Camptosar PF ^a Hospira Pty Limited HH ^a Irinotecan Actavis GQ ^a Irinotecan Alphapharm AF ^a Irinotecan Ebewe IT ^a Irinotecan Sandoz SZ ^a Omegapharm OE ^a Irinotecan Tecan WQ
9119Y	I.V. injection 500 mg in 25 mL	1	3	..	1560.34	34.20	^a Hospira Pty Limited HH ^a Irinotecan Ebewe IT
9410G	I.V. injection 300 mg in 15 mL	1	3	..	938.02	34.20	^a Camptosar PF ^a Irinotecan Ebewe IT

TOPOTECAN HYDROCHLORIDE

Authority required (STREAMLINED)

3186

Advanced metastatic ovarian cancer after failure of prior therapy which includes a platinum compound.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
8199M	Powder for I.V. infusion 4 mg (base)	5	1	..	2126.36	34.20	Hycamtin	GK

Endocrine therapy

Hormones and related agents

Progestogens

MEDROXYPROGESTERONE ACETATE

Restricted benefit

Hormone-dependent advanced breast cancer.

2728N	Tablet 500 mg	30	2	..	128.31	34.20	Provera	PF

MEDROXYPROGESTERONE ACETATE

Restricted benefit

Hormone-dependent breast cancer;

Endometrial cancer.

2316X	Tablet 200 mg	60	2	..	103.74	34.20	Provera	PF
2725K	Tablet 100 mg	100	2	..	92.10	34.20	Provera	PF
2727M	Tablet 250 mg	60	2	..	128.08	34.20	Provera	PF

MEGESTROL ACETATE

Restricted benefit

Hormone-dependent advanced breast cancer.

2734X	Tablet 160 mg	30	2	..	83.39	34.20	Megace	SI

Gonadotropin releasing hormone analogues

GOSERELIN ACETATE

Authority required

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate;

Hormone-dependent locally advanced (equivalent to stage III) or metastatic (equivalent to stage IV) breast cancer in pre-menopausal women;

Short-term treatment (up to 6 months) of visually proven endometriosis (only 1 course of not more than 6 months' therapy will be authorised);

Hormone-dependent breast cancer as an alternative to adjuvant chemotherapy in peri- or pre-menopausal women.

1454M	Subcutaneous implant 3.6 mg (base) in pre-filled injection syringe	1	5	..	333.00	34.20	Zoladex Implant	AP

GOSERELIN ACETATE

Authority required (STREAMLINED)

3229

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate.

8093Y	Subcutaneous implant (long acting) 10.8 mg (base) in pre-filled injection syringe	1	1	..	1108.76	34.20	Zoladex 10.8 Implant	AP

GOSERELIN ACETATE and BICALUTAMIDE

Authority required (STREAMLINED)

3239

Metastatic (equivalent to stage D) prostatic carcinoma in patients for whom a combination of an antiandrogen and a GnRH (LH-RH) agonist is required.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9064C	Pack containing 1 subcutaneous implant goserelin 3.6 mg in pre-filled injection syringe and 28 tablets bicalutamide 50 mg	1	5	..	480.43	34.20	ZolaCos CP 3.6/50	AP
9065D	Pack containing 1 subcutaneous implant goserelin 10.8 mg in pre-filled injection syringe	1	..	..	1251.13	34.20	ZolaCos CP 10.8/50(28)	AP

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
9066E	and 28 tablets bicalutamide 50 mg Pack containing 1 subcutaneous implant goserelin 10.8 mg in pre-filled injection syringe and 84 tablets bicalutamide 50 mg	±1	1	..	1535.89	34.20	ZolaCos CP 10.8/50(84)	AP

LEUPRORELIN ACETATE

Authority required (STREAMLINED)

3229

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate.

8707G	Suspension for subcutaneous injection (modified release), 7.5 mg injection set	1	5	..	420.20	34.20	Eligard 1 month	HH
8708H	Suspension for subcutaneous injection (modified release), 22.5 mg injection set	1	1	..	1108.76	34.20	Eligard 3 month	HH
8709J	Suspension for subcutaneous injection (modified release), 30 mg injection set	1	1	..	1451.33	34.20	Eligard 4 month	HH
8859G	Suspension for subcutaneous injection (modified release), 45 mg injection set	1	..	..	2123.98	34.20	Eligard 6 month	HH
8875D	I.M. injection (modified release), powder for injection 7.5 mg with diluent in pre-filled dual-chamber syringe	1	5	..	420.20	34.20	Lucrin Depot 7.5mg PDS	AB
8876E	I.M. injection (modified release), powder for injection 22.5 mg with diluent in pre-filled dual-chamber syringe	1	1	..	1108.76	34.20	Lucrin Depot 3 Month PDS	AB
8877F	I.M. injection (modified release), powder for injection 30 mg with diluent in pre-filled dual-chamber syringe	1	1	..	1451.33	34.20	Lucrin Depot 4 Month PDS	AB

TRIPTORELIN

Authority required (STREAMLINED)

3229

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate.

5297T	Powder for I.M. injection (prolonged release) 22.5 mg (as embonate) with solvent, syringe and needles	1	..	..	2123.98	34.20	Diphereline	IS
9378N	Powder for I.M. injection (prolonged release) 3.75 mg (as embonate) with solvent, syringe and needles	1	5	..	420.20	34.20	Diphereline	IS
9379P	Powder for I.M. injection (prolonged release) 11.25 mg (as embonate) with solvent, syringe and needles	1	1	..	1108.76	34.20	Diphereline	IS

Hormone antagonists and related agents

Anti-estrogens

TAMOXIFEN CITRATE

Restricted benefit

Treatment of hormone-dependent breast cancer.

Note

This drug is not PBS-subsidised for primary prevention of breast cancer.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2109B NP	Tablet 10 mg (base)	60	5	..	38.88	34.20	Genox 10	AF
2110C NP	Tablet 20 mg (base)	60	5	..	60.24	34.20	^a Genox 20	AF
							^a GenRx Tamoxifen	GX
							^a Tamosin	SI
							^a Tamoxen 20 mg	GM
							^a Tamoxifen Sandoz	SZ
					*64.08	34.20	^a Nolvadex-D	AP

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

TOREMIFENE CITRATE

Restricted benefit

Treatment of hormone-dependent metastatic breast cancer in post-menopausal patients.

Note

This drug is not PBS-subsidised for primary prevention of breast cancer.

8216K	Tablet 60 mg (base)	30	5	..	73.74	34.20	Fareston	SH

Anti-androgens

BICALUTAMIDE

Authority required (STREAMLINED)

3247

Metastatic (equivalent to stage D) prostatic carcinoma in combination with GnRH (LH-RH) agonist therapy.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8094B NP	Tablet 50 mg	28	5	..	168.34	34.20	^a	APO-Bicalutamide	TX
							^a	Bicalutamide-GA	GM
							^a	Bicalutamide	RA
							^a	Ranbaxy	
							^a	Calutex	SI
							^a	Cosamide	AF
							^a	Cosudex	AP

CYPROTERONE ACETATE

Authority required (STREAMLINED)

1014

Advanced carcinoma of the prostate;

1404

To reduce drive in sexual deviations in males.

1270W	Tablet 50 mg	100	5	..	*207.72	34.20	^a	Cyprohexal	SZ
							^a	Cyprone	AF
							^a	Cyprostat	SY
							^a	GenRx Cyproterone	GX
							^a	Acetate	
8019C	Tablet 100 mg	50	5	..	*210.72	34.20	^a	Procur	GM
							^a	Androcur	SC
							^a	Cyprohexal	SZ
							^a	Cyprostat-100	SY
							^a	GenRx Cyproterone	GX
							^a	Acetate	
					^B 3.00		^a	Procur 100	GM
					^B 1.65		^a	Androcur-100	SC

FLUTAMIDE

Authority required (STREAMLINED)

3247

Metastatic (equivalent to stage D) prostatic carcinoma in combination with GnRH (LH-RH) agonist therapy.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
1417N NP	Tablet 250 mg	100	5	..	205.56	34.20 ^a	Eulexin	SH
						^a	Flutamin	AF

NILUTAMIDE

Authority required (STREAMLINED)

3299

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) prostatic carcinoma, in combination with GnRH (LH-RH) agonist therapy;

3300

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) prostatic carcinoma, in conjunction with surgical orchidectomy.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8131Y NP	Tablet 150 mg	30	5	..	236.56	34.20	Anandron	SW

Enzyme inhibitors

ANASTROZOLE

Restricted benefit

Treatment of hormone-dependent breast cancer in post-menopausal women.

Note

This drug is not PBS-subsidised for primary prevention of breast cancer.

This drug is not PBS-subsidised for adjuvant hormonal treatment of early breast cancer extended beyond 5 years.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8179L NP	Tablet 1 mg	30	5	..	180.18	34.20	Arimidex	AP

EXEMESTANE

Restricted benefit

Treatment of hormone-dependent advanced breast cancer in post-menopausal women with disease progression following treatment with tamoxifen citrate;

Treatment of hormone-dependent early breast cancer in post-menopausal women following a minimum of 2 years' treatment with tamoxifen citrate.

Note

This drug is not PBS-subsidised for primary prevention of breast cancer.

This drug is not PBS-subsidised for adjuvant hormonal treatment of early breast cancer extended beyond 5 years, i.e. a patient who has received 2 years of tamoxifen therapy may only receive 3 years of PBS-subsidised treatment with exemestane.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8506Q NP	Tablet 25 mg	30	5	..	180.18	34.20	Aromasin	PF

LETROZOLE

Restricted benefit

Treatment of hormone-dependent advanced breast cancer in post-menopausal women;

Treatment of hormone-dependent early breast cancer in post-menopausal women;

Extended adjuvant treatment of hormone-dependent early breast cancer in post-menopausal women commencing within 6 months of ceasing treatment with tamoxifen citrate.

Note

This drug is not PBS-subsidised for primary prevention of breast cancer.

This drug is not PBS-subsidised for adjuvant hormonal treatment of early breast cancer extended beyond 5 years.

This drug is not PBS-subsidised for extended adjuvant early breast cancer treatment where the total duration of letrozole (or any other aromatase inhibitor) treatment extends beyond 5 years.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	
Note Shared Care Model: For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
8245Y NP	Tablet 2.5 mg	30	5	..	180.18	34.20	Femara 2.5 mg NV

Other hormone antagonists and related agents

DEGARELIX

Authority required (STREAMLINED)

3229

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate.

5455D	Powder for injection 80 mg (as acetate) with solvent, syringe and needles	1	5	..	420.20	34.20	Firmagon 80mg FP

DEGARELIX

Authority required (STREAMLINED)

3229

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate.

Note

No applications for increased maximum quantities and/or repeats will be authorised for the 120 mg powder for injection.

5456E	Powder for injection 120 mg (as acetate) with solvent, syringe and needles, 2	1	..	..	438.72	34.20	Firmagon 120mg FP

Immunostimulants

Immunostimulants

Interferons

INTERFERON ALFA-2a

Caution

Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required

Hairy cell leukaemia;

Myeloproliferative disease with excessive thrombocytosis.

8180M	Injection 3,000,000 i.u. in 0.5 mL single dose pre-filled syringe	15	4	..	*506.22	34.20	Roferon-A RO

INTERFERON ALFA-2a

Caution

Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required

Myeloproliferative disease with excessive thrombocytosis.

8551C	Injection 4,500,000 i.u. in 0.5 mL single dose pre-filled syringe	5	4	..	*264.72	34.20	Roferon-A RO
8552D	Injection 6,000,000 i.u. in 0.5 mL single dose pre-filled syringe	5	4	..	*344.72	34.20	Roferon-A RO
8553E	Injection 9,000,000 i.u. in 0.5 mL single dose pre-filled syringe	5	4	..	*506.12	34.20	Roferon-A RO

INTERFERON ALFA-2a

Caution

Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
<u>Authority required</u>							
Low grade non-Hodgkin's lymphoma with clinical features suggestive of a poor prognosis, in combination with anthracycline-based chemotherapy.							
8181N	Injection 3,000,000 i.u. in 0.5 mL single dose pre-filled syringe	15	5	..	*506.22	34.20	Roferon-A RO
8182P	Injection 4,500,000 i.u. in 0.5 mL single dose pre-filled syringe	5	5	..	*264.72	34.20	Roferon-A RO
8183Q	Injection 6,000,000 i.u. in 0.5 mL single dose pre-filled syringe	5	5	..	*344.72	34.20	Roferon-A RO
8184R	Injection 9,000,000 i.u. in 0.5 mL single dose pre-filled syringe	5	5	..	*506.12	34.20	Roferon-A RO

INTERFERON ALFA-2b

Caution

Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required

Hairy cell leukaemia.

8572E	Solution for injection 18,000,000 i.u. in 1.2 mL multi-dose injection pen	3	4	..	*606.03	34.20	Intron A Redipen SH

INTERFERON ALFA-2b

Caution

Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required

Maintenance treatment of multiple myeloma once remission has been achieved with chemotherapy;

Low grade non-Hodgkin's lymphoma with clinical features suggestive of a poor prognosis, in combination with anthracycline-based chemotherapy.

8348J	Solution for injection 18,000,000 i.u. in 1.2 mL multi-dose injection pen	3	5	..	*606.03	34.20	Intron A Redipen SH
8476D	Solution for injection 30,000,000 i.u. in 1.2 mL multi-dose injection pen	3	5	..	*1005.75	34.20	Intron A Redipen SH

INTERFERON BETA-1a

Authority required

Initial treatment of clinically definite relapsing-remitting multiple sclerosis in ambulatory (without assistance or support) patients who have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to the multiple sclerosis, in the preceding 2 years. The diagnosis must be confirmed by magnetic resonance imaging of the brain and/or spinal cord and the date of the scan included in the authority application, unless the authority application is accompanied by written certification provided by a radiologist that an MRI scan is contraindicated because of the risk of physical (not psychological) injury to the patient. The authority will be limited to the maximum quantity and number of repeats indicated in the schedule;

Continuing treatment of clinically definite relapsing-remitting multiple sclerosis in patients previously issued with an authority prescription for this drug who do not show continuing progression of disability while on treatment with this drug and who have demonstrated compliance with, and an ability to tolerate, this therapy. Authorities will be limited to the maximum quantity and number of repeats indicated in the schedule.

8289G	Injection set comprising 1 vial powder for injection 30 micrograms (6,000,000 i.u.) with diluent	4	5	..	1056.77	34.20	Avonex BD
8403G	Injection 44 micrograms (12,000,000 i.u.) in 0.5 mL single dose pre-filled syringe	12	5	..	1056.77	34.20	Rebif 44 SG
8805K	Injection 30 micrograms (6,000,000 i.u.) in 0.5 mL single dose pre-filled syringe	4	5	..	1056.77	34.20	Avonex BD
9332E	Solution for injection 132 micrograms in 1.5 mL multidose cartridge	4	5	..	1056.77	34.20	Rebif 44 SG

INTERFERON BETA-1b

Authority required

Initial treatment of clinically definite relapsing-remitting multiple sclerosis in ambulatory (without assistance or support) patients who have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to the multiple sclerosis, in the preceding 2 years. The diagnosis must be confirmed by magnetic resonance imaging of the brain and/or spinal cord and the date of the scan included in the authority application, unless the authority application is accompanied by written certification provided by a radiologist that an MRI scan is contraindicated because of the risk of physical (not psychological) injury to the patient. The authority will be limited to the maximum quantity and number of repeats indicated in the schedule;

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
8101J	Continuing treatment of clinically definite relapsing-remitting multiple sclerosis in patients previously issued with an authority prescription for this drug who do not show continuing progression of disability while on treatment with this drug and who have demonstrated compliance with, and an ability to tolerate, this therapy. Authorities will be limited to the maximum quantity and number of repeats indicated in the schedule.						
	Injection set including 1 vial powder for injection 8,000,000 i.u. (250 micrograms) and solvent	15	5	..	1180.16	34.20	Betaferon SC

Other immunostimulants

BCG IMMUNOTHERAPEUTIC (Bacillus Calmette-Guérin/ Connaught strain)

Restricted benefit

Treatment of carcinoma in situ of the urinary bladder.

1140B	Single dose set comprising 1 vial powder for intravesical administration containing 6.6 to 19.2 x 10 ⁸ CFU and 1 vial diluent 3 mL	3	1	..	*459.87	34.20	ImmuCyst SW

BCG-TICE (Bacillus Calmette-Guérin/ Tice strain)

Restricted benefit

Primary and relapsing superficial urothelial carcinoma of the bladder.

1131M	Vial containing powder for intravesical administration approximately 5 x 10 ⁸ CFU	3	1	..	556.39	34.20	OncoTICE SH

GLATIRAMER ACETATE

Authority required

Initial treatment of clinically definite relapsing-remitting multiple sclerosis in ambulatory (without assistance or support) patients who have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to the multiple sclerosis, in the preceding 2 years. The diagnosis must be confirmed by magnetic resonance imaging of the brain and/or spinal cord and the date of the scan included in the authority application, unless the authority application is accompanied by written certification provided by a radiologist that an MRI scan is contraindicated because of the risk of physical (not psychological) injury to the patient. The authority will be limited to the maximum quantity and number of repeats indicated in the schedule;

Continuing treatment of clinically definite relapsing-remitting multiple sclerosis in patients previously issued with an authority prescription for this drug who do not show continuing progression of disability while on treatment with this drug and who have demonstrated compliance with, and an ability to tolerate, this therapy. Authorities will be limited to the maximum quantity and number of repeats indicated in the schedule.

8726G	Injection 20 mg in 1 mL single dose pre-filled syringe	28	5	..	1092.65	34.20	Copaxone SW

Immunosuppressants

Immunosuppressants

Selective immunosuppressants

EVEROLIMUS

Caution

Careful monitoring of patients is mandatory.

Authority required

Maintenance therapy, following initiation and stabilisation of treatment with everolimus and where therapy remains under the supervision and direction of the transplant unit reviewing that patient, of patients with:

(a) renal transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application;

(b) cardiac transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application.

8840G	Tablet 0.25 mg	60	3	..	282.79	34.20	Certican NV
8841H	Tablet 0.5 mg	60	3	..	543.83	34.20	Certican NV
8842J	Tablet 0.75 mg	120	3	..	*1578.62	34.20	Certican NV
9352F	Tablet 1 mg	120	3	..	*2068.76	34.20	Certican NV

LEFLUNOMIDE

Caution

Leflunomide is a category X drug and must not be given to pregnant women. Pregnancy should be avoided for two years after cessation of therapy, unless special wash-out procedures are carried out.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
<u>Authority required (STREAMLINED)</u>							
2643							
Initial treatment of severe active rheumatoid arthritis where other disease modifying anti-rheumatic drugs (including methotrexate) are ineffective and/or inappropriate. Treatment must be initiated by a physician;							
2681							
Initial treatment of severe active psoriatic arthritis where other disease modifying anti-rheumatic drugs (including methotrexate) are ineffective and/or inappropriate. Treatment must be initiated by a physician.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised.							
8373Q	Pack containing 3 tablets leflunomide 100 mg and 30 tablets leflunomide 20 mg	£1	..	..	211.31	34.20	Arava SW

LEFLUNOMIDE

Caution

Leflunomide is a category X drug and must not be given to pregnant women. Pregnancy should be avoided for two years after cessation of therapy, unless special wash-out procedures are carried out.

Authority required (STREAMLINED)

2644

Treatment of severe active rheumatoid arthritis where other disease modifying anti-rheumatic drugs (including methotrexate) are ineffective and/or inappropriate. Treatment must be initiated by a physician;

2682

Treatment of severe active psoriatic arthritis where other disease modifying anti-rheumatic drugs (including methotrexate) are ineffective and/or inappropriate. Treatment must be initiated by a physician.

8374R	Tablet 10 mg	30	5	..	91.91	34.20	^a Arabloc ^a Arava	AV SW
8375T	Tablet 20 mg	30	5	..	136.59	34.20	^a Arabloc ^a Arava	AV SW

MYCOPHENOLATE MOFETIL

Caution

Careful monitoring of patients is mandatory.

Authority required

Maintenance therapy, following initiation and stabilisation of treatment with mycophenolate mofetil and where therapy remains under the supervision and direction of the transplant unit reviewing that patient, of patients with:

(a) renal transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application;

(b) cardiac transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application.

8649F	Capsule 250 mg	300	3	..	*627.81	34.20	CellCept	RO
8650G	Tablet 500 mg	150	3	..	*627.81	34.20	CellCept	RO
8651H	Powder for oral suspension 1 g per 5 mL, 165 mL	£1	3	..	#289.85	34.20	CellCept	RO

MYCOPHENOLATE SODIUM

Caution

Careful monitoring of patients is mandatory.

Authority required

Maintenance therapy, following initiation and stabilisation of treatment with mycophenolate sodium and where therapy remains under the supervision and direction of the transplant unit reviewing that patient, of patients with renal transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application.

8652J	Tablet (enteric coated) 180 mg (mycophenolic acid)	120	3	..	263.42	34.20	Myfortic	NV
8653K	Tablet (enteric coated) 360 mg (mycophenolic acid)	120	3	..	503.53	34.20	Myfortic	NV

SIROLIMUS

Caution

Careful monitoring of patients is mandatory.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty	Safety Net	
\$ \$							
<u>Authority required</u>							
Maintenance therapy, following initiation and stabilisation of treatment with sirolimus and where therapy remains under the supervision and direction of the transplant unit reviewing that patient, of patients with renal transplants. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application.							
8724E	Tablet 1 mg	100	3	..	815.25	34.20	Rapamune WX
8725F	Oral solution 1 mg per mL, 60 mL	±1	3	..	529.74	34.20	Rapamune WX
8833X	Tablet 2 mg	100	3	..	1583.69	34.20	Rapamune WX

Tumor necrosis factor alpha (TNF-alpha) inhibitors

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

(iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (new patient or patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].							

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 16 weeks of treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab.

Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Applications for treatment with adalimumab where the dosing frequency exceeds 40 mg per fortnight will not be approved.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have a documented history of severe active rheumatoid arthritis; and
- (b) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with adalimumab and who wish to re-commence therapy with this drug, must

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised adalimumab treatment, within the timeframes specified below.						
	A maximum of 16 weeks of treatment will be authorised under this restriction.						
	Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).						
	Where the most recent course of PBS-subsidised adalimumab treatment was approved under either of the initial 1 or 2 treatment restrictions, patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.						
	Where the most recent course of PBS-subsidised adalimumab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.						
	Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.						
	<u>Note</u> No applications for increased maximum quantities and/or repeats will be authorised. Applications for treatment with adalimumab where the dosing frequency exceeds 40 mg per fortnight will not be approved.						
	<u>Note</u> Special Pricing Arrangements apply.						
8737W	Injection 40 mg in 0.8 mL pre-filled syringe	2	3	..	1774.36	34.20	Humira AB
9099X	Injection 40 mg in 0.8 mL pre-filled pen	2	3	..	1774.36	34.20	Humira AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.							

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count)

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have a documented history of severe active rheumatoid arthritis; and
- (b) who have demonstrated an adequate response to treatment with adalimumab; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with adalimumab.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).						

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with adalimumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with adalimumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Applications for treatment with adalimumab where the dosing frequency exceeds 40 mg per fortnight will not be approved.

Note

Special Pricing Arrangements apply.

8741C	Injection 40 mg in 0.8 mL pre-filled syringe	2	5	..	1774.36	34.20	Humira	AB
9100Y	Injection 40 mg in 0.8 mL pre-filled pen	2	5	..	1774.36	34.20	Humira	AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents (adalimumab, etanercept, golimumab and infliximab) for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time. Where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, golimumab and infliximab.

From 1 August 2006, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis after 1 August 2010.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients — golimumab only.

Applications for patients who commenced treatment with golimumab prior to 1 March 2010 may apply for initial PBS-subsidised treatment as continuing therapy under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; or
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
prescription or remaining repeats for the biological agent the patient is ceasing.							

(4) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment Cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Authority required

Initial 1

Initial PBS-subsidised treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults who:

- (1) have severe active psoriatic arthritis; and
- (2) have received no prior PBS-subsidised biological treatment for this condition in this Treatment Cycle; and
- (3) have failed to achieve an adequate response to:
 - (a) methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months; and
 - (b) sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; or
 - (c) leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities, including severity, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) an active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 16 weeks treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

The assessment of the patient's response to the initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.							

Authority required

Initial 2

Initial PBS-subsidised treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults who:

- (1) have a documented history of severe active psoriatic arthritis; and
- (2) have received prior PBS-subsidised biological treatment for this condition in this Treatment Cycle and are eligible to receive further biological therapy; and
- (3) have not failed treatment with adalimumab during the current Treatment Cycle.

Applications for patients who have received PBS-subsidised treatment with adalimumab within this Treatment Cycle and who wish to re-commence therapy with this drug within this same Cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised adalimumab treatment, within the timeframes specified below.

A maximum of 16 weeks treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised adalimumab treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised biological therapy or, under this restriction, for patients who have received previous PBS-subsidised biological therapy), patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised adalimumab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Applications for treatment with adalimumab where the dosing frequency exceeds 40 mg per fortnight will not be approved.

9033K	Injection 40 mg in 0.8 mL pre-filled syringe	2	3	..	1774.36	34.20	Humira	AB
9101B	Injection 40 mg in 0.8 mL pre-filled pen	2	3	..	1774.36	34.20	Humira	AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents (adalimumab, etanercept, golimumab and infliximab) for adult patients with severe active psoriatic arthritis.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time. Where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, golimumab and infliximab.

From 1 August 2006, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis after 1 August 2010.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients — golimumab only.

Applications for patients who commenced treatment with golimumab prior to 1 March 2010 may apply for initial PBS-subsidised treatment as continuing therapy under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; or
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment Cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with adalimumab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults:

- (1) who have a documented history of severe active psoriatic arthritis; and
- (2) whose most recent course of PBS-subsidised biological agent for this condition in the current Treatment Cycle was with adalimumab; and
- (3) who, at the time of application, demonstrate an adequate response to treatment with adalimumab.

An adequate response to treatment with adalimumab is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with adalimumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with adalimumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised.							
Applications for treatment with adalimumab where the dosing frequency exceeds 40 mg per fortnight will not be approved.							
9034L	Injection 40 mg in 0.8 mL pre-filled syringe	2	5	..	1774.36	34.20	Humira AB
9102C	Injection 40 mg in 0.8 mL pre-filled pen	2	5	..	1774.36	34.20	Humira AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH ACTIVE ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept, golimumab and infliximab for adult patients with active ankylosing spondylitis. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab, etanercept, golimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 4 TNF-alfa antagonists at any 1 time.

From 1 March 2007, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised TNF-alfa antagonists without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 March 2007 is considered to be in their first cycle as of 1 March 2007.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than once. A patient who, prior to 1 March 2007, was authorised to receive PBS-subsidised initial treatment for ankylosing spondylitis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2007.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty	Safety Net	
					\$	\$	
	(iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).						

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and 18 weeks of treatment for infliximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap to an alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with at least 1 NSAID, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the BASDAI, ESR and/or CRP levels are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for active ankylosing spondylitis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with golimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with golimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial 1 (new patients)

Initial PBS-subsidised treatment with adalimumab, by a rheumatologist, of an adult with active ankylosing spondylitis who has radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis and who has not received any PBS-subsidised treatment with either adalimumab, etanercept, golimumab or infliximab in this treatment cycle; AND

(a) who has at least 2 of the following:

- (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or
- (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI) [for further information on the BASMI please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; or
- (iii) limitation of chest expansion relative to normal values for age and gender [for chest expansion normal values please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; AND

(b) who has failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months.

The application must include details of the NSAIDs trialled, their doses and duration of treatment. If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance. Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Medicare Australia website at www.medicareaustralia.gov.au.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au] which must include the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form [www.medicareaustralia.gov.au]; and
 - (iii) a completed Exercise Program Self Certification Form included in the supporting information form; and
 - (iv) a signed patient acknowledgment form.

The assessment of the patient's response to the initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A maximum of 16 weeks of treatment with adalimumab will be approved under this criterion.

Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
<u>Authority required</u>							
Initial 2 (change or re-commencement for all patients)							
Initial PBS-subsidised treatment with adalimumab, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who, in this treatment cycle, has received prior PBS-subsidised TNF-alfa antagonist treatment for this condition and is eligible to receive further TNF-alfa antagonist therapy, and has not failed PBS-subsidised therapy with adalimumab in the current treatment cycle.							
Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised TNF-alfa antagonist therapy or, under this restriction, for patients who have received previous PBS-subsidised TNF-alfa antagonist therapy) the patient must have been assessed for response to that course following a minimum of 12 weeks of treatment. These assessments must be provided to Medicare Australia no later than 4 weeks from the date the course was ceased. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.							
Where the most recent course of PBS-subsidised adalimumab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.							
Authority applications must be made in writing and must include:							
(a) a completed authority prescription form; and							
(b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au].							
A maximum of 16 weeks of treatment with adalimumab will be approved under this criterion.							
Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.							
Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised.							
Applications for treatment with adalimumab where the dosing frequency exceeds 40 mg per fortnight will not be approved.							
9077R	Injection 40 mg in 0.8 mL pre-filled syringe	2	3	..	1774.36	34.20	Humira AB
9103D	Injection 40 mg in 0.8 mL pre-filled pen	2	3	..	1774.36	34.20	Humira AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH ACTIVE ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept, golimumab and infliximab for adult patients with active ankylosing spondylitis. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab, etanercept, golimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 4 TNF-alfa antagonists at any 1 time.

From 1 March 2007, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised TNF-alfa antagonists without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 March 2007 is considered to be in their first cycle as of 1 March 2007.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

once. A patient who, prior to 1 March 2007, was authorised to receive PBS-subsidised initial treatment for ankylosing spondylitis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2007.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF- α antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF- α antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF- α antagonist under the new treatment cycle.

A patient who has failed fewer than 3 TNF- α antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 TNF- α antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF- α antagonist therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF- α antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF- α antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF- α antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and 18 weeks of treatment for infliximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF- α antagonist.

For second and subsequent courses of PBS-subsidised TNF- α antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific TNF- α antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF- α antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF- α antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF- α antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF- α antagonist is approved, a patient may swap to an alternate TNF- α antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate TNF- α antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF- α antagonist at the time of the application. However, they cannot swap to a particular TNF- α antagonist if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate TNF- α antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF- α antagonist the patient is ceasing.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a TNF- α antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF- α antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with at least 1 NSAID, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the BASDAI, ESR and/or CRP levels are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for active ankylosing spondylitis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with golimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with golimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Continuing treatment for all patients

Continuing PBS-subsidised treatment, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who:

- (a) has demonstrated an adequate response to treatment with adalimumab; and
- (b) whose most recent course of PBS-subsidised therapy in this treatment cycle was with adalimumab.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au].

All measurements provided must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.

All applications for continuing treatment with adalimumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment following an initial treatment course it must be made following a minimum of 12 weeks of treatment with adalimumab. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Patients who fail to demonstrate a response to treatment with adalimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial adalimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF- α antagonist was approved in this cycle and the date of the first application under a new cycle.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Note							
No applications for increased maximum quantities and/or repeats will be authorised.							
Applications for treatment with adalimumab where the dosing frequency exceeds 40 mg per fortnight will not be approved.							
9078T	Injection 40 mg in 0.8 mL pre-filled syringe	2	5	..	1774.36	34.20	Humira AB
9104E	Injection 40 mg in 0.8 mL pre-filled pen	2	5	..	1774.36	34.20	Humira AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and infliximab for adult patients with severe refractory Crohn disease. Where the term 'tumour necrosis factor (TNF) alpha antagonist' appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alfa antagonists at any 1 time.

From 1 August 2008, under the PBS, all patients will be able to commence a treatment cycle where they may trial each PBS-subsidised TNF-alfa antagonist without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 August 2008 is considered to be in their first cycle as of 1 August 2008.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than twice.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2008.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

From 1 August 2008, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap if eligible to the alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, evidence of intestinal inflammation), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab or infliximab.

A patient who commenced treatment with adalimumab for severe refractory Crohn disease prior to 9 November 2007 or infliximab prior to 7 March 2007 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab or

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

infliximab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab or infliximab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial 1 (new patients)

Initial treatment of Crohn disease in a patient assessed by CDAI.

Initial PBS-subsidised treatment with adalimumab by a gastroenterologist or a consultant physician as specified in the NOTE below, of a patient with severe refractory Crohn disease who satisfies the following criteria:

- (a) has confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician as specified in the NOTE below; and
- (b) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (c) has failed to achieve an adequate response to prior systemic therapy including:
 - (i) a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; and
 - (ii) immunosuppressive therapy including:
 - azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
 - 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
 - methotrexate at a dose of at least 15 mg weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) have a severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as assessed.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

The most recent CDAI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the signed patient acknowledgement.

A maximum of 16 weeks treatment will be authorised under this criterion.

Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats. Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

A CDAI assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Authority required

Initial 2

Change or re-commencement of treatment of Crohn disease in a patient assessed by CDAI.

Initial PBS-subsidised treatment with adalimumab by a gastroenterologist or a consultant physician as specified in the NOTE below of a patient who:

- (a) has a documented history of severe refractory Crohn disease; and
- (b) in this treatment cycle, has received prior PBS-subsidised treatment with infliximab or adalimumab for this condition; and
- (c) has not failed PBS-subsidised therapy with adalimumab for this condition more than once in the current treatment cycle.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of TNF-alfa antagonist therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of TNF-alfa antagonist treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of TNF-alfa antagonist.

Authority applications must be made in writing and must include:

- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CAI) Score calculation sheet including the date of the assessment of the patient's condition; and
 - (ii) details of prior TNF alfa antagonist treatment including details of date and duration of treatment.

Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats. Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

A CDAI assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

Authority required

Initial 1

Initial treatment of Crohn disease in a patient with short gut syndrome or an ostomy patient.

Initial PBS-subsidised treatment with adalimumab by a gastroenterologist, or consultant physician as specified in the NOTE below of a patient who satisfies the following criteria:

- (a) has confirmed Crohn disease defined by standard clinical, endoscopic and/or imaging features, including histological evidence with the diagnosis confirmed by a gastroenterologist or consultant physician as specified in the NOTE below; and
- (b) has diagnostic imaging or surgical evidence of short gut syndrome or has an ileostomy or colostomy; and
- (c) has evidence of intestinal inflammation; and
- (d) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (e) has failed to achieve an adequate response to prior systemic drug therapy including:
 - (i) a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; and
 - (ii) immunosuppressive therapy including:
 - azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
 - 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
 - methotrexate at a dose of at least 15 mg weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) have evidence of intestinal inflammation, including:
 - (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; AND/OR
 - (ii) faeces: higher than normal lactoferrin or calprotectin level; AND/OR
 - (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery;
- AND/OR
- (b) be assessed clinically as being in a high faecal output state;
- AND/OR
- (c) be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of adalimumab.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

Applications for authorisation must be made in writing and must include:

- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (ii) reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iii) date of the most recent clinical assessment; and
 - (iv) the signed patient acknowledgement.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application.

A maximum of 16 weeks treatment will be authorised under this criterion.

Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats. Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

Authority required

Initial 2

Change or re-commencement of treatment of Crohn disease in a patient with short gut syndrome, an ostomy patient or a patient with extensive small intestine disease.

Initial PBS-subsidised treatment with adalimumab by a gastroenterologist or a consultant physician as specified in the NOTE below of a patient who:

- (a) has a documented history of severe refractory Crohn disease; and
- (b) in this treatment cycle, has received prior PBS-subsidised treatment with infliximab or adalimumab for this condition; and
- (c) has not failed PBS-subsidised therapy with adalimumab for this condition more than once in the current treatment cycle.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of TNF-alfa antagonist therapy within the timeframes specified in the relevant restriction.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of TNF-alfa antagonist treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of TNF-alfa antagonist.

Authority applications must be made in writing and must include:

- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criteria, if relevant; and
 - (ii) details of prior TNF alfa antagonist treatment including details of date and duration of treatment.

A maximum of 16 weeks of treatment will be approved under this criterion.

Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats. Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

Authority required

Initial 1

Initial treatment of Crohn disease in a patient with extensive small intestine disease.

Initial PBS-subsidised treatment with adalimumab by a gastroenterologist or a consultant physician as specified in the NOTE below, of a patient with severe refractory Crohn disease who satisfies the following criteria:

- (a) has confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or consultant physician as specified in the NOTE below; and
- (b) has extensive small intestinal disease with radiological evidence of intestinal inflammation affecting more than 50 cm of the small intestine; and
- (c) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) has failed to achieve an adequate response to prior systemic therapy including:
 - (i) a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; and
 - (ii) immunosuppressive therapy including:
 - azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
 - 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
 - methotrexate at a dose of at least 15 mg weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) have severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; AND/OR
- (b) have evidence of active intestinal inflammation, including:
 - (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; AND/OR
 - (ii) faeces: higher than normal lactoferrin or calprotectin level; AND/OR
 - (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	lymphadenopathy or fat streaking in the mesentery; AND/OR (c) be assessed clinically as being in a high faecal output state; AND/OR (d) be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of adalimumab.						

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

Applications for authorisation must be made in writing and must include:

- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (ii) (1) reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; or
 - (2) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the dates of assessment of the patient's condition, if relevant; and
 - (iii) date of the most recent clinical assessment; and
 - (iv) the signed patient acknowledgement.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application.

A maximum of 16 weeks treatment of adalimumab will be authorised under this criterion.

Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats. Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy after the first dose so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9186L	Injection 40 mg in 0.8 mL pre-filled syringe, 6	1	..	..	5036.36	34.20	Humira	AB
9187M	Injection 40 mg in 0.8 mL pre-filled pen, 6	1	..	..	5036.36	34.20	Humira	AB
9188N	Injection 40 mg in 0.8 mL pre-filled syringe	2	2	..	1774.36	34.20	Humira	AB
9190Q	Injection 40 mg in 0.8 mL pre-filled pen	2	2	..	1774.36	34.20	Humira	AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty	Recordable Value for Safety Net	
					\$	\$	

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and infliximab for adult patients with severe refractory Crohn disease. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alfa antagonists at any 1 time.

From 1 August 2008, under the PBS, all patients will be able to commence a treatment cycle where they may trial each PBS-subsidised TNF-alfa antagonist without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 August 2008 is considered to be in their first cycle as of 1 August 2008.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than twice.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2008.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2008, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.							

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap if eligible to the alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, evidence of intestinal inflammation), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab or infliximab.

A patient who commenced treatment with adalimumab for severe refractory Crohn disease prior to 9 November 2007 or infliximab prior to 7 March 2007 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab or infliximab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab or infliximab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial 3 (grandfather)

Initial PBS-subsidised treatment of Crohn disease in a patient assessed by CDAI who has previously received non-PBS-subsidised therapy with adalimumab.

Initial PBS-subsidised supply for continuing treatment with adalimumab by a gastroenterologist, a consultant physician as specified in the NOTE below, or other consultant physician in consultation with a gastroenterologist of a patient who:

- has a documented history of severe refractory Crohn disease and was receiving treatment with adalimumab prior to 9 November 2007; and
- had a Crohn Disease Activity Index (CDAI) Score of greater than or equal to 300 prior to commencing treatment with adalimumab. Where a baseline CDAI assessment is not available, please call Medicare Australia on 1800 700 270 to discuss; and
- has signed a patient acknowledgement indicating that they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- has demonstrated or sustained an adequate response to treatment with adalimumab. For advice please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or

Antineoplastic and immunomodulating agents

					Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	\$	\$	Brand Name and Manufacturer
	consultant physicians [general medicine specialising in gastroenterology (code 82)].						

An adequate response to adalimumab treatment is defined as a reduction in Crohn Disease Activity Index (CDAI) Score to no greater than 150.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and baseline Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition; and
 - (ii) the signed patient acknowledgement.

The current CDAI assessment must be no more than 1 month old at the time of application. The baseline CDAI assessment must be from immediately prior to commencing treatment with adalimumab.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with adalimumab.

A maximum of 24 weeks treatment will be approved under this criterion.

Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Authority required

Continuing treatment of Crohn disease in a patient assessed by CDAI.

Continuing PBS-subsidised treatment with adalimumab by a gastroenterologist, a consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease; and
- (b) has demonstrated or sustained an adequate response to treatment with adalimumab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to adalimumab treatment is defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition.

The CDAI assessment must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with adalimumab, a CDAI assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with adalimumab.

Patients are eligible to receive continuing adalimumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

A maximum of 24 weeks treatment will be authorised under this criterion.

Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment of Crohn disease in a patient with short gut syndrome or an ostomy patient.

Continuing PBS-subsidised treatment with adalimumab by a gastroenterologist, a consultant physician as specified in the NOTE below or other

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease with intestinal inflammation and with short gut syndrome or with an ileostomy or colostomy; and
- (b) has demonstrated or sustained an adequate response to treatment with adalimumab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to adalimumab treatment is defined as:

- (a) improvement of intestinal inflammation as demonstrated by:
 - (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; AND/OR
 - (ii) faeces: normalisation of lactoferrin or calprotectin level; AND/OR
 - (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or
- (b) reversal of high faecal output state; or
- (c) avoidance of the need for surgery or total parenteral nutrition (TPN).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy or the date of clinical assessment.

The patient's assessment must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with adalimumab, an assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy after the first dose so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with adalimumab.

Patients are eligible to receive continuing adalimumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

A maximum of 24 weeks of treatment will be authorised under this criterion.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment of Crohn disease in a patient with extensive small intestine disease.

Continuing PBS-subsidised treatment with adalimumab by a gastroenterologist, or consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease with extensive intestinal inflammation affecting more than 50 cm of the small intestine; and
- (b) has demonstrated or sustained an adequate response to treatment with adalimumab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to adalimumab treatment is defined as:

- (a) a reduction in Crohn Disease Activity Index (CDAI) Score to no greater than 150; or
- (b) improvement of intestinal inflammation as demonstrated by:
 - (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; AND/OR
 - (ii) faeces: normalisation of lactoferrin or calprotectin level; AND/OR
 - (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or
- (c) reversal of high faecal output state; or
- (d) avoidance of the need for surgery or total parenteral nutrition (TPN).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition; or
 - (ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy; or

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

(iii) the date of clinical assessment.

All assessments must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with adalimumab, an assessment of the patient's response must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with adalimumab.

Patients are eligible to receive continuing adalimumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

A maximum of 24 weeks treatment will be authorised under this criterion.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial 3

Initial PBS-subsidised treatment of Crohn disease in a patient with short gut syndrome, an ostomy patient, or a patient with extensive small intestine disease, who has previously received non-PBS-subsidised therapy with adalimumab.

Initial PBS-subsidised supply for continuing treatment with adalimumab by a gastroenterologist, a consultant physician as specified in the NOTE below, or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease and was receiving treatment with adalimumab prior to 9 November 2007; and
- (b) (1) has a history of extensive small intestinal disease with radiological evidence of intestinal inflammation affecting more than 50 cm of the small intestine; or
- (2) has diagnostic imaging or surgical evidence of short gut syndrome or has an ileostomy or colostomy with a documented history of intestinal inflammation; and
- (c) has signed a patient acknowledgement indicating that they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) has demonstrated or sustained an adequate response to treatment with adalimumab according to the criteria included in the relevant continuation restriction. For advice please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

The same criteria used to determine an inadequate response to prior treatment at baseline must be used to determine response to treatment and eligibility for continuing therapy, according to the criteria included in the continuing treatment restriction.

An adequate response to adalimumab treatment is defined as:

- (a) a reduction in Crohn Disease Activity Index (CDAI) Score to no greater than 150; or
- (b) improvement of intestinal inflammation as demonstrated by:
 - (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; AND/OR
 - (ii) faeces: normalisation of lactoferrin or calprotectin level; AND/OR
 - (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or
 - (c) reversal of high faecal output state; or
 - (d) avoidance of the need for surgery or total parenteral nutrition (TPN).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) (1) the completed current and baseline Crohn Disease Activity Index (CDAI) Score calculation sheet, where relevant, including the date of the assessment of the patient's condition; or
 - (2) the reports and dates of the current and baseline pathology or diagnostic imaging test(s) in order to assess response to therapy; or
 - (3) the date of clinical assessment(s); and
 - (ii) the signed patient acknowledgement.

The patient's assessment must be no more than 1 month old at the time of application. The baseline CDAI assessments must be from immediately prior to commencing treatment with adalimumab. Where a baseline assessment is not available, please call Medicare Australia on 1800 700 270 to discuss.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with adalimumab.							
Patients are eligible to receive continuing adalimumab treatment in courses of up to 24 weeks providing they continue to sustain the response.							
Patients who fail to demonstrate or sustain a response to treatment with adalimumab for Crohn disease as specified in the criteria for continuing treatment with adalimumab, will not be eligible to recommence PBS-subsidised treatment with this drug within 12 months of the date on which treatment was ceased.							
A maximum of 24 weeks treatment will be authorised under this criterion.							
Where fewer than 5 repeats are requested at the time of this application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).							
Patients may qualify for PBS-subsidised treatment under this restriction once only.							
Note							
No applications for increased maximum quantities and/or repeats will be authorised.							
9189P	Injection 40 mg in 0.8 mL pre-filled syringe	2	5	..	1774.36	34.20	Humira AB
9191R	Injection 40 mg in 0.8 mL pre-filled pen	2	5	..	1774.36	34.20	Humira AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, etanercept, infliximab and ustekinumab, for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, infliximab and ustekinumab.

From 1 March 2010, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial adalimumab, etanercept, infliximab or ustekinumab without having to meet the initial treatment criteria, that is they will not need to experience a disease flare when swapping to an alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

A patient who received PBS-subsidised biological agent treatment for chronic plaque psoriasis prior to 1 March 2010 is considered to be in their first Cycle as of 1 March 2010.

Patients are eligible for PBS-subsidised treatment with only 1 biological agent at any 1 time.

Within the same Treatment Cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for a PBS-subsidised biological agent, they must change to an alternate agent if they wish to continue PBS-subsidised biological treatment. A patient who, prior to 1 March 2010, was authorised to receive PBS-subsidised initial treatment for chronic plaque psoriasis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2010.

Patients must be assessed for response to each course of continuing treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a Treatment Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological agent therapy before they are eligible to commence the next Cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological agent treatment in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Treatment Cycle.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Patients for whom a break in PBS-subsidised therapy of less than 5 years duration has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe chronic plaque psoriasis after 1 March 2010.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made in the following situations:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); or
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under '(4) Swapping therapy' below]; or
- (iii) patients who wish to re-commence treatment following a break in PBS-subsidised therapy with that agent (Initial 2).

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of treatment in the case of adalimumab and etanercept, 22 weeks of treatment in the case of infliximab and 28 weeks of treatment in the case of ustekinumab.

Grandfather patients (adalimumab and ustekinumab only).

Applications for patients who commenced treatment with adalimumab prior to 1 March 2009 or ustekinumab prior to 26 November 2009, may be made for initial PBS-subsidised treatment as continuing therapy under the initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with a biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment. Approval will be based on the criteria included in the relevant restriction.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological agent, a PASI assessment must be conducted after at least 12 weeks of treatment. This assessment must be submitted to Medicare Australia within 1 month of the completion of this initial treatment course. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Application for continuing treatment.

Following the completion of an initial treatment course of a biological agent to which an adequate response has been demonstrated, patients may qualify to receive up to 24 weeks of continuing treatment with that biological agent. Patients are eligible to continue to receive continuous treatment with 24 week courses providing they continue to sustain a response.

For second and subsequent courses of PBS-subsidised treatment with adalimumab, etanercept, infliximab or ustekinumab it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to sustain a response to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

(4) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate agent within the same Treatment Cycle without having to requalify with respect to disease severity (i.e. a PASI score of greater than 15), or prior treatment requirements.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

Patients may trial an alternate biological agent at any time, regardless of whether they are receiving therapy with a biological agent at the time of the application or not. However, they cannot swap to a particular agent if they have failed to respond to treatment with that particular agent within the same Cycle.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Patients who commenced treatment with adalimumab prior to 1 June 2009 or ustekinumab prior to 1 March 2010 access these interchangeability arrangements in the same way as patients who have not.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the agent being ceased.

(5) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a Treatment Cycle and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatment applications.

(6) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent Biological Treatment Cycle, following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment according to the criteria of the relevant restriction and index of disease severity. Patients must have had at least 1 prior treatment, as listed in the criteria, for a minimum of 6 weeks, and must have a PASI assessment conducted preferably whilst still on treatment, but no later than 1 month following cessation of treatment. The PASI assessment must be no older than 1 month at the time of application.

Authority required

Initial treatment [Initial 1, Whole body (New patients — No prior biological agent)]

Initial treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis; and
- (b) have not received any prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (whole body); and
- (d) have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 3 of the following 4 treatments:
 - (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or
 - (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or
 - (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or
 - (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities including severity, associated with phototherapy, methotrexate, cyclosporin and acitretin, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 16 weeks of treatment with adalimumab will be authorised under this restriction.

Where fewer than 4 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

would otherwise extend the initial treatment period beyond 16 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

Authority required

Initial or re-Treatment [Initial 2, Whole body (Received prior biological agent under PBS)]

Treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have a documented history of severe chronic plaque psoriasis; and
- (b) have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have not failed PBS-subsidised therapy with adalimumab for the treatment of this condition in the current Treatment Cycle.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

Applications for patients who have demonstrated a response to PBS-subsidised adalimumab treatment within this Treatment Cycle and who wish to re-commence adalimumab treatment within the same Cycle following a break in therapy, will only be approved where evidence of the patient's response to their most recent course of PBS-subsidised adalimumab treatment has been submitted to Medicare Australia within 1 month of cessation of treatment.

A maximum of 16 weeks of treatment with adalimumab will be authorised under this restriction.

Where fewer than 4 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 16 weeks.

A PASI assessment of the patient's response to this course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Initial treatment [Initial 1, Face, hand, foot (New patients — No prior biological agent)]

Initial treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- (b) have not received any prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (face, hand, foot); and
- (d) have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 3 of the following 4 treatments:
 - (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or
 - (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or
 - (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or
 - (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or where

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

phototherapy is contraindicated, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities including severity, associated with phototherapy, methotrexate, cyclosporin and acitretin, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]]; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 16 weeks of treatment with adalimumab will be authorised under this restriction.

Where fewer than 4 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 16 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Authority required

Initial or re-Treatment [Initial 2, Face, hand, foot (Received prior biological agent under PBS)]

Treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot; and
- (b) have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have not failed PBS-subsidised therapy with adalimumab for the treatment of this condition in the current Treatment Cycle.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]]; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

Applications for patients who have demonstrated a response to PBS-subsidised adalimumab treatment within this Treatment Cycle and who wish to re-commence adalimumab treatment within the same Cycle following a break in therapy, will only be approved where evidence of the patient's response to their most recent course of PBS-subsidised adalimumab treatment has been submitted to Medicare Australia within 1 month of cessation of treatment.

A maximum of 16 weeks of treatment with adalimumab will be authorised under this restriction.

Where fewer than 4 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 16 weeks.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

A PASI assessment of the patient's response to this course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

Special Pricing Arrangements apply.

9425C	Injection 40 mg in 0.8 mL pre-filled syringe	2	4	..	1774.36	34.20	Humira	AB
9426D	Injection 40 mg in 0.8 mL pre-filled pen	2	4	..	1774.36	34.20	Humira	AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, etanercept, infliximab and ustekinumab, for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, infliximab and ustekinumab.

From 1 March 2010, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial adalimumab, etanercept, infliximab or ustekinumab without having to meet the initial treatment criteria, that is they will not need to experience a disease flare when swapping to an alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

A patient who received PBS-subsidised biological agent treatment for chronic plaque psoriasis prior to 1 March 2010 is considered to be in their first Cycle as of 1 March 2010.

Patients are eligible for PBS-subsidised treatment with only 1 biological agent at any 1 time.

Within the same Treatment Cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for a PBS-subsidised biological agent, they must change to an alternate agent if they wish to continue PBS-subsidised biological treatment. A patient who, prior to 1 March 2010, was authorised to receive PBS-subsidised initial treatment for chronic plaque psoriasis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2010.

Patients must be assessed for response to each course of continuing treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a Treatment Cycle and they must

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

have, at a minimum, a 5-year break in PBS-subsidised biological agent therapy before they are eligible to commence the next Cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological agent treatment in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Treatment Cycle.

Patients for whom a break in PBS-subsidised therapy of less than 5 years duration has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe chronic plaque psoriasis after 1 March 2010.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made in the following situations:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); or
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under '(4) Swapping therapy' below]; or
- (iii) patients who wish to re-commence treatment following a break in PBS-subsidised therapy with that agent (Initial 2).

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of treatment in the case of adalimumab and etanercept, 22 weeks of treatment in the case of infliximab and 28 weeks of treatment in the case of ustekinumab.

Grandfather patients (adalimumab and ustekinumab only).

Applications for patients who commenced treatment with adalimumab prior to 1 March 2009 or ustekinumab prior to 26 November 2009, may be made for initial PBS-subsidised treatment as continuing therapy under the initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with a biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment. Approval will be based on the criteria included in the relevant restriction.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological agent, a PASI assessment must be conducted after at least 12 weeks of treatment. This assessment must be submitted to Medicare Australia within 1 month of the completion of this initial treatment course. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Application for continuing treatment.

Following the completion of an initial treatment course of a biological agent to which an adequate response has been demonstrated, patients may qualify to receive up to 24 weeks of continuing treatment with that biological agent. Patients are eligible to continue to receive continuous treatment with 24 week courses providing they continue to sustain a response.

For second and subsequent courses of PBS-subsidised treatment with adalimumab, etanercept, infliximab or ustekinumab it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to sustain a response to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

(4) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate agent within the same Treatment Cycle without having to requalify with respect to disease severity (i.e. a PASI score of greater than 15), or prior treatment requirements.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Patients may trial an alternate biological agent at any time, regardless of whether they are receiving therapy with a biological agent at the time of the application or not. However, they cannot swap to a particular agent if they have failed to respond to treatment with that particular agent within the same Cycle.

Patients who commenced treatment with adalimumab prior to 1 June 2009 or ustekinumab prior to 1 March 2010 access these interchangeability arrangements in the same way as patients who have not.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the agent being ceased.

(5) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a Treatment Cycle and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatment applications.

(6) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent Biological Treatment Cycle, following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment according to the criteria of the relevant restriction and index of disease severity. Patients must have had at least 1 prior treatment, as listed in the criteria, for a minimum of 6 weeks, and must have a PASI assessment conducted preferably whilst still on treatment, but no later than 1 month following cessation of treatment. The PASI assessment must be no older than 1 month at the time of application.

Authority required

Initial treatment [Initial 3, Whole body (Grandfather patients)]

Initial PBS-subsidised supply for continuing treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have a documented history of severe chronic plaque psoriasis and were receiving treatment with adalimumab prior to 1 March 2009; and
- (b) had a Psoriasis Area and Severity Index (PASI) score of greater than 15 prior to commencing treatment with adalimumab; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (whole body); and
- (d) have demonstrated a response as specified in the criterion included in the restriction for continuing PBS-subsidised treatment with adalimumab (whole body).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition at baseline (prior to initiation of adalimumab therapy) and the most recent PASI assessment [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 24 weeks of treatment with adalimumab will be authorised under this restriction.

Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks.

A PASI assessment of the patient's response to this initial PBS-subsidised course of therapy must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The most recent PASI assessment must be no more than 1 month old at the time of application.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Authority required

Continuing treatment (Whole body)

Continuing PBS-subsidised treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over:

- (a) who have a documented history of severe chronic plaque psoriasis; and
- (b) whose most recent course of PBS-subsidised biological treatment for this condition in this Treatment Cycle was with adalimumab; and
- (c) who have demonstrated an adequate response to their most recent course of treatment with adalimumab.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-biological treatment baseline value for this Treatment Cycle.

This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with adalimumab, the assessment of response must be after a minimum of 12 weeks of treatment with an initial course.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet along with the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with adalimumab.

A maximum of 24 weeks of treatment with adalimumab will be authorised under this restriction.

Where fewer than 5 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for continuing authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks.

A PASI assessment of the patient's response must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Initial treatment [Initial 3, Face, hand, foot (Grandfather patients)]

Initial PBS-subsidised supply for continuing treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over:

- (a) who have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot and were receiving treatment with adalimumab prior to 1 March 2009; and
- (b) whose disease, prior to treatment with adalimumab, was of a severity as defined in the initiation criterion included in the initial treatment restriction (Initial 1, New patients — face, hand, foot); and
- (c) who have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (face, hand, foot); and
- (d) who have demonstrated a response as specified in the criterion included in the restriction for continuing PBS-subsidised treatment with adalimumab (face, hand, foot).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams along with the date of the assessment of the patient's condition at baseline (prior to initiation of adalimumab therapy) and the most recent PASI assessment [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

(iii) the signed patient and prescriber acknowledgements.

A maximum of 24 weeks of treatment with adalimumab will be authorised under this restriction.

Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 24 weeks.

A PASI assessment of the patient's response to this initial PBS-subsidised course of therapy must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The most recent PASI assessment must be no more than 1 month old at the time of application.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

The PASI assessment must be performed on the same affected area as assessed at baseline or prior to initiation of adalimumab treatment.

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Authority required

Continuing treatment (Face, hand, foot)

Continuing PBS-subsidised treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over:

- (a) who have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot; and
- (b) whose most recent course of PBS-subsidised biological treatment for this condition in this Treatment Cycle was with adalimumab; and
- (c) who have demonstrated an adequate response to treatment with adalimumab.

An adequate response to adalimumab treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the pre-biological treatment baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the pre-biological treatment baseline value.

This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with adalimumab, the assessment of response must be after a minimum of 12 weeks of treatment with an initial course.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams along with the date of the assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

The most recent PASI assessment must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with adalimumab will be authorised under this restriction.

Where fewer than 5 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for continuing authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks.

A PASI assessment of the patient's response must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised adalimumab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	
the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised.							
<u>Note</u>							
Special Pricing Arrangements apply.							
9427E	Injection 40 mg in 0.8 mL pre-filled syringe	2	5	..	1774.36	34.20	Humira AB
9428F	Injection 40 mg in 0.8 mL pre-filled pen	2	5	..	1774.36	34.20	Humira AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and etanercept for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term bDMARD appears in the following NOTES and restrictions, it refers to adalimumab and etanercept only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 bDMARDs at any one time.

From 1 November 2010, a patient receiving PBS-subsidised bDMARD therapy is considered to be in a treatment cycle where they may swap to the alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy, and
- fail to respond, or to sustain a response to one PBS-subsidised bDMARD twice and the other PBS-subsidised bDMARD once only.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised bDMARD therapy before they are eligible to receive further PBS-subsidised bDMARD therapy. The length of a treatment break is measured from the date of the last approval for PBS-subsidised bDMARD therapy in the last treatment cycle and the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

A patient who was receiving PBS-subsidised bDMARD treatment immediately prior to 1 November 2010 is considered to be in their first cycle as of 1 November 2010. A patient who has had a break in bDMARD treatment of at least 12 months immediately prior to making a new application, on or after 1 November 2010, will commence a new treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle. The length of the break in therapy is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment was stopped to the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 November 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

A patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to the alternate bDMARD without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months.

A patient may trial the alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug twice within the current treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a bDMARD. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to the revised baseline measurement.

(4) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 12 months, must requalify for treatment under the Initial 1 treatment restriction.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for severe active juvenile idiopathic arthritis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must qualify

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 12 month break in PBS-subsidised therapy' above for further details.							

Authority required

Initial 1 (new patient or patient recommencing after a break of more than 12 months).

Initial treatment, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of a patient aged 18 years or older who:

- (a) has a documented history of juvenile idiopathic arthritis with onset prior to the age of 18 years; AND
- (b) has received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) has failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg per day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) an active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
	(1) a completed authority prescription form; and (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and (3) a signed patient acknowledgement.						
	A maximum of 16 weeks of treatment will be authorised under this restriction.						
	Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).						
	Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment.						
	Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab.						
	If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial adalimumab after a minimum of 5 years have elapsed between the date of the last approval for PBS-subsidised bDMARD therapy in the last treatment cycle and the date of the first application under a new treatment cycle.						
	<u>Authority required</u>						
	Initial 2 (change or re-commencement after break of less than 12 months).						
	Initial PBS-subsidised treatment with adalimumab by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of a patient aged 18 years or older who:						
	(a) has a documented history of juvenile idiopathic arthritis with onset prior to the age of 18 years; AND						
	(b) in this treatment cycle, has received prior PBS-subsidised treatment with adalimumab or etanercept for this condition; and						
	(c) has not failed PBS-subsidised therapy with adalimumab for this condition more than once in the current treatment cycle.						
	The authority application must be made in writing and must include:						
	(a) a completed authority prescription form; and						
	(b) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].						
	Applications for a patient who has received PBS-subsidised treatment with adalimumab in this treatment cycle and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised adalimumab treatment, within the timeframes specified below.						
	A maximum of 16 weeks of treatment will be authorised under this restriction.						
	Where fewer than 3 repeats are requested at the time of the initial authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).						
	Where the most recent course of PBS-subsidised adalimumab treatment was approved under either of the initial 1 or 2 treatment restrictions, the patient must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.						
	Where the most recent course of PBS-subsidised adalimumab treatment was approved under the continuing treatment criteria, the patient must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.						
	Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to that particular course of bDMARD.						
	If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial adalimumab after a minimum of 5 years have elapsed between the date of the last approval for PBS-subsidised bDMARD therapy in the last treatment cycle and the date of the first application under a new treatment cycle.						
5281Y	Injection 40 mg in 0.8 mL pre-filled syringe	2	3	..	1774.36	34.20	Humira AB
5282B	Injection 40 mg in 0.8 mL pre-filled pen	2	3	..	1774.36	34.20	Humira AB

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and etanercept for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term bDMARD appears in the following NOTES and restrictions, it refers to adalimumab and etanercept only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 bDMARDs at any one time.

From 1 November 2010, a patient receiving PBS-subsidised bDMARD therapy is considered to be in a treatment cycle where they may swap to the alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy, and
- fail to respond, or to sustain a response to one PBS-subsidised bDMARD twice and the other PBS-subsidised bDMARD once only.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised bDMARD therapy before they are eligible to receive further PBS-subsidised bDMARD therapy. The length of a treatment break is measured from the date of the last approval for PBS-subsidised bDMARD therapy in the last treatment cycle and the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

A patient who was receiving PBS-subsidised bDMARD treatment immediately prior to 1 November 2010 is considered to be in their first cycle as of 1 November 2010. A patient who has had a break in bDMARD treatment of at least 12 months immediately prior to making a new application, on or after 1 November 2010, will commence a new treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle. The length of the break in therapy is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment was stopped to the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 November 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

A patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to the alternate bDMARD without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months.

A patient may trial the alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug twice within the current treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a bDMARD. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to the revised baseline measurement.

(4) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 12 months, must requalify for treatment under the Initial 1 treatment restriction.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for severe active juvenile idiopathic arthritis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must qualify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 12 month break in PBS-subsidised therapy' above for further details.

Authority required

Initial 3 ('grandfather' patients).

Initial PBS-subsidised supply for continuing treatment, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of a patient aged 18 years or older who:

- (a) has a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years; and
- (b) was receiving treatment with adalimumab prior to 1 March 2010; and
- (c) has demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with adalimumab; and
- (d) is receiving treatment with adalimumab at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 24 weeks of treatment with adalimumab will be approved under this criterion.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

The same indices of disease severity used to establish baseline at the commencement of treatment with a bDMARD must be used for assessment of all continuing applications.

The assessment of the patient's response to a continuing course of therapy must be made within 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled in order to ensure continuity of treatment for those patients who meet the continuation criterion.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

Authority required

Continuing treatment.

Continuing PBS-subsidised treatment, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of a patient aged 18 years or older:

- (a) who has a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years; and
- (b) who has demonstrated an adequate response to treatment with adalimumab; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with adalimumab.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; AND either of the following:

- (i) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (ii) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (iii) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with adalimumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with adalimumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial adalimumab after a minimum of 5 years have elapsed between the date of the last approval for PBS-subsidised bDMARD therapy in the last treatment cycle and the date of the first application under a new treatment cycle.

5283C	Injection 40 mg in 0.8 mL pre-filled syringe	2	5	..	1774.36	34.20	Humira	AB
5284D	Injection 40 mg in 0.8 mL pre-filled pen	2	5	..	1774.36	34.20	Humira	AB

CERTOLIZUMAB PEGOL

Note

Any queries concerning the arrangements to prescribe certolizumab pegol may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Further prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe certolizumab pegol should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	GPO Box 9826 HOBART TAS 7001;						

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (new patient or patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with certolizumab pegol, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 18 to 20 weeks of treatment depending on the dosage regimen will be authorised under this restriction.

Where fewer than 5 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 18 or 20 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with certolizumab pegol.

Patients who fail to demonstrate a response to treatment with certolizumab pegol under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with certolizumab pegol, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have a documented history of severe active rheumatoid arthritis; and
- (b) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with certolizumab pegol and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised certolizumab pegol treatment, within the timeframes specified below.

A maximum of 18 to 20 weeks of treatment depending on the dosage regimen will be authorised under this restriction.

Where fewer than 5 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 18 or 20 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised certolizumab pegol treatment was approved under either of the initial 1 or 2 treatment restrictions, patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised certolizumab pegol treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Patients who fail to demonstrate a response to treatment with certolizumab pegol under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 3 ('grandfather' patients)

Initial PBS-subsidised supply for continuing treatment with certolizumab pegol, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of an adult who:

- (a) has a documented history of severe active rheumatoid arthritis; and
- (b) was receiving treatment with certolizumab pegol prior to 1 March 2010; and
- (c) has demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with certolizumab pegol; and
- (d) is receiving treatment with certolizumab pegol at the time of application.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au]; and
- (3) a signed patient acknowledgement.

The same indices of disease severity used to establish baseline at the commencement of treatment with a bDMARD must be used for assessment of all continuing applications.

The assessment of the patient's response to a continuing course of therapy must be made within 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled in order to ensure continuity of treatment for those patients who meet the continuation criterion.

A maximum of 24 weeks of treatment with certolizumab pegol will be approved under this criterion.

Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Patients who fail to demonstrate a response to treatment with certolizumab pegol under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with certolizumab pegol, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have a documented history of severe active rheumatoid arthritis; and
- (b) who have demonstrated an adequate response to treatment with certolizumab pegol; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with certolizumab pegol.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with certolizumab pegol must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with certolizumab pegol, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Patients who fail to demonstrate a response to treatment with certolizumab pegol under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

3425G	Injection 200 mg in 1 mL single use pre-filled syringe	2	5	..	1708.64	34.20	Cimzia	UC

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (new patient or patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with etanercept, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
(ii) at least 4 active joints from the following list of major joints:
— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 16 weeks of treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept.

Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with etanercept, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have a documented history of severe active rheumatoid arthritis; and
- (b) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with etanercept and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised etanercept treatment, within the timeframes specified below.

A maximum of 16 weeks of treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised etanercept treatment was approved under either of the initial 1 or 2 treatment restrictions, patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised etanercept treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
<u>Note</u> No applications for increased maximum quantities and/or repeats will be authorised.							
<u>Note</u> Special Pricing Arrangements apply.							
8637N	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	3	..	*1829.00	34.20	Enbrel WX
9089J	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	3	..	1774.37	34.20	Enbrel WX
9459W	Injection 50 mg in 1 mL single use auto-injector, 4	1	3	..	1774.37	34.20	Enbrel WX

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

[illegible]

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with etanercept, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have a documented history of severe active rheumatoid arthritis; and
- (b) who have demonstrated an adequate response to treatment with etanercept; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with etanercept.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;
AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
All applications for continuing treatment with etanercept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with etanercept, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course. Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. <u>Note</u> No applications for increased maximum quantities and/or repeats will be authorised. <u>Note</u> Special Pricing Arrangements apply.							
8638P	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	5	..	*1829.00	34.20	Enbrel WX
9090K	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	5	..	1774.37	34.20	Enbrel WX
9460X	Injection 50 mg in 1 mL single use auto-injector, 4	1	5	..	1774.37	34.20	Enbrel WX

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents (adalimumab, etanercept, golimumab and infliximab) for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time. Where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, golimumab and infliximab.

From 1 August 2006, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis after 1 August 2010.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients — golimumab only.

Applications for patients who commenced treatment with golimumab prior to 1 March 2010 may apply for initial PBS-subsidised treatment as continuing therapy under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialed it on the PBS; or
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment Cycle and Medicare Australia will assess response according to these revised baseline measurements.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Authority required

Initial 1

Initial PBS-subsidised treatment with etanercept, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults who:

- (1) have severe active psoriatic arthritis; and
- (2) have received no prior PBS-subsidised biological treatment for this condition in this Treatment Cycle; and
- (3) have failed to achieve an adequate response to:
 - (a) methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months; and
 - (b) sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; or
 - (c) leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities, including severity, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) an active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 16 weeks treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

The assessment of the patient's response to the initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial etanercept after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Initial 2

Initial PBS-subsidised treatment with etanercept, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults who:

- (1) have a documented history of severe active psoriatic arthritis; and
- (2) have received prior PBS-subsidised biological treatment for this condition in this Treatment Cycle and are eligible to receive further biological therapy; and

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

(3) have not failed treatment with etanercept during the current Treatment Cycle.

Applications for patients who have received PBS-subsidised treatment with etanercept within this Treatment Cycle and who wish to re-commence therapy with this drug within this same Cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised etanercept treatment, within the timeframes specified below.

A maximum of 16 weeks treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised etanercept treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised biological therapy or, under this restriction, for patients who have received previous PBS-subsidised biological therapy), patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised etanercept treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial etanercept after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9035M	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	3	..	*1829.00	34.20	Enbrel	WX
9087G	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	3	..	1774.37	34.20	Enbrel	WX
9457R	Injection 50 mg in 1 mL single use auto-injector, 4	1	3	..	1774.37	34.20	Enbrel	WX

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents (adalimumab, etanercept, golimumab and infliximab) for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time. Where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, golimumab and infliximab.

From 1 August 2006, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis after 1 August 2010.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients — golimumab only.

Applications for patients who commenced treatment with golimumab prior to 1 March 2010 may apply for initial PBS-subsidised treatment as continuing therapy under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; or
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment Cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with etanercept, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults:

- (1) who have a documented history of severe active psoriatic arthritis; and
- (2) whose most recent course of PBS-subsidised biological agent for this condition in the current Treatment Cycle was with etanercept; and
- (3) who, at the time of application, demonstrate an adequate response to treatment with etanercept.

An adequate response to treatment with etanercept is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with etanercept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with etanercept, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial etanercept after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
9036N	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	5	..	*1829.00	34.20	Enbrel	WX
9088H	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	5	..	1774.37	34.20	Enbrel	WX
9458T	Injection 50 mg in 1 mL single use auto-injector, 4	1	5	..	1774.37	34.20	Enbrel	WX

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH ACTIVE ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept, golimumab and infliximab for adult patients with active ankylosing spondylitis. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab, etanercept, golimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 4 TNF-alfa antagonists at any 1 time.

From 1 March 2007, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised TNF-alfa antagonists without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 March 2007 is considered to be in their first cycle as of 1 March 2007.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than once. A patient who, prior to 1 March 2007, was authorised to receive PBS-subsidised initial treatment for ankylosing spondylitis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2007.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

(Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and 18 weeks of treatment for infliximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF- α antagonist.

For second and subsequent courses of PBS-subsidised TNF- α antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific TNF- α antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF- α antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF- α antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF- α antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF- α antagonist is approved, a patient may swap to an alternate TNF- α antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate TNF- α antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF- α antagonist at the time of the application. However, they cannot swap to a particular TNF- α antagonist if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate TNF- α antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF- α antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a TNF- α antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF- α antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with at least 1 NSAID, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the BASDAI, ESR and/or CRP levels are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for active ankylosing spondylitis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with golimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with golimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial 1 (new patients)

Initial PBS-subsidised treatment with etanercept, by a rheumatologist, of an adult with active ankylosing spondylitis who has radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis and who has not received any PBS-subsidised treatment with either adalimumab, etanercept, golimumab or infliximab in this treatment cycle; AND

(a) who has at least 2 of the following:

- (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or
- (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI) [for further information on the BASMI please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; or
- (iii) limitation of chest expansion relative to normal values for age and gender [for chest expansion normal values please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; AND

(b) who has failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months.

The application must include details of the NSAIDs trialled, their doses and duration of treatment. If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance. Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Medicare Australia website at www.medicareaustralia.gov.au.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au] which must include the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form [www.medicareaustralia.gov.au]; and
 - (iii) a completed Exercise Program Self Certification Form included in the supporting information form; and
 - (iv) a signed patient acknowledgment form.

The assessment of the patient's response to the initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A maximum of 16 weeks of treatment with etanercept will be approved under this criterion.

Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial etanercept after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
<u>Authority required</u>							
Initial 2 (change or re-commencement for all patients)							
Initial PBS-subsidised treatment with etanercept, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who, in this treatment cycle, has received prior PBS-subsidised TNF-alfa antagonist treatment for this condition and is eligible to receive further TNF-alfa antagonist therapy, and has not failed PBS-subsidised therapy with etanercept in the current treatment cycle.							
Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised TNF-alfa antagonist therapy or, under this restriction, for patients who have received previous PBS-subsidised TNF-alfa antagonist therapy) the patient must have been assessed for response to that course following a minimum of 12 weeks of treatment. These assessments must be provided to Medicare Australia no later than 4 weeks from the date the course was ceased. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.							
Where the most recent course of PBS-subsidised etanercept treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.							
Authority applications must be made in writing and must include:							
(a) a completed authority prescription form; and							
(b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au].							
A maximum of 16 weeks of treatment with etanercept will be approved under this criterion.							
Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.							
Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial etanercept after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised.							
8778B	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	3	..	*1829.00	34.20	Enbrel WX
9085E	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	3	..	1774.37	34.20	Enbrel WX
9455P	Injection 50 mg in 1 mL single use auto-injector, 4	1	3	..	1774.37	34.20	Enbrel WX

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH ACTIVE ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept, golimumab and infliximab for adult patients with active ankylosing spondylitis. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab, etanercept, golimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 4 TNF-alfa antagonists at any 1 time.

From 1 March 2007, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised TNF-alfa antagonists without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 March 2007 is considered to be in their first cycle as of 1 March 2007.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than once. A patient who, prior to 1 March 2007, was authorised to receive PBS-subsidised initial treatment for ankylosing spondylitis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2007.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and 18 weeks of treatment for infliximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap to an alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with at least 1 NSAID, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the BASDAI, ESR and/or CRP levels are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for active ankylosing spondylitis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with golimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with golimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Continuing treatment for all patients

Continuing PBS-subsidised treatment, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who:

- (a) has demonstrated an adequate response to treatment with etanercept; and
- (b) whose most recent course of PBS-subsidised therapy in this treatment cycle was with etanercept.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au].

All measurements provided must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with etanercept will be authorised under this criterion.

Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.

All applications for continuing treatment with etanercept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment following an initial treatment course it must be made following a minimum of 12 weeks of treatment with etanercept. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Patients who fail to demonstrate a response to treatment with etanercept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial etanercept after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.							
<u>Note</u> No applications for increased maximum quantities and/or repeats will be authorised.							
8779C	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	5	..	*1829.00	34.20	Enbrel WX
9086F	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	5	..	1774.37	34.20	Enbrel WX
9456Q	Injection 50 mg in 1 mL single use auto-injector, 4	1	5	..	1774.37	34.20	Enbrel WX

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, etanercept, infliximab and ustekinumab, for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, infliximab and ustekinumab.

From 1 March 2010, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial adalimumab, etanercept, infliximab or ustekinumab without having to meet the initial treatment criteria, that is they will not need to experience a disease flare when swapping to an alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

A patient who received PBS-subsidised biological agent treatment for chronic plaque psoriasis prior to 1 March 2010 is considered to be in their first Cycle as of 1 March 2010.

Patients are eligible for PBS-subsidised treatment with only 1 biological agent at any 1 time.

Within the same Treatment Cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for a PBS-subsidised biological agent, they must change to an alternate agent if they wish to continue PBS-subsidised biological treatment. A patient who, prior to 1 March 2010, was authorised to receive PBS-subsidised initial treatment for chronic plaque psoriasis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2010.

Patients must be assessed for response to each course of continuing treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a Treatment Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological agent therapy before they are eligible to commence the next Cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological agent treatment in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Treatment Cycle.

Patients for whom a break in PBS-subsidised therapy of less than 5 years duration has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

How to prescribe biological agents for the treatment of severe chronic plaque psoriasis after 1 March 2010.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made in the following situations:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); or
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under '(4) Swapping therapy' below]; or
- (iii) patients who wish to re-commence treatment following a break in PBS-subsidised therapy with that agent (Initial 2).

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of treatment in the case of adalimumab and etanercept, 22 weeks of treatment in the case of infliximab and 28 weeks of treatment in the case of ustekinumab.

Grandfather patients (adalimumab and ustekinumab only).

Applications for patients who commenced treatment with adalimumab prior to 1 March 2009 or ustekinumab prior to 26 November 2009, may be made for initial PBS-subsidised treatment as continuing therapy under the initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with a biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment. Approval will be based on the criteria included in the relevant restriction.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological agent, a PASI assessment must be conducted after at least 12 weeks of treatment. This assessment must be submitted to Medicare Australia within 1 month of the completion of this initial treatment course. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Application for continuing treatment.

Following the completion of an initial treatment course of a biological agent to which an adequate response has been demonstrated, patients may qualify to receive up to 24 weeks of continuing treatment with that biological agent. Patients are eligible to continue to receive continuous treatment with 24 week courses providing they continue to sustain a response.

For second and subsequent courses of PBS-subsidised treatment with adalimumab, etanercept, infliximab or ustekinumab it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to sustain a response to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

(4) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate agent within the same Treatment Cycle without having to requalify with respect to disease severity (i.e. a PASI score of greater than 15), or prior treatment requirements.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

Patients may trial an alternate biological agent at any time, regardless of whether they are receiving therapy with a biological agent at the time of the application or not. However, they cannot swap to a particular agent if they have failed to respond to treatment with that particular agent within the same Cycle.

Patients who commenced treatment with adalimumab prior to 1 June 2009 or ustekinumab prior to 1 March 2010 access these interchangeability arrangements in the same way as patients who have not.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the agent being ceased.

(5) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a Treatment Cycle and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatment applications.

(6) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent Biological Treatment Cycle, following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment according to the criteria of the relevant restriction and index of disease severity. Patients must have had at least 1 prior treatment, as listed in the criteria, for a minimum of 6 weeks, and must have a PASI assessment conducted preferably whilst still on treatment, but no later than 1 month following cessation of treatment. The PASI assessment must be no older than 1 month at the time of application.

Authority required

Initial treatment [Initial 1, Whole body (New patients — No prior biological agent)]

Initial treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis; and
- (b) have not received any prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (whole body); and
- (d) have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 3 of the following 4 treatments:
 - (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or
 - (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or
 - (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or
 - (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities including severity, associated with phototherapy, methotrexate, cyclosporin and acitretin, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 16 weeks of treatment with etanercept will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 16 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare

Antineoplastic and immunomodulating agents

							Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium		\$	\$	Brand Name and Manufacturer	
	Australia on 1800 700 270 to discuss.								

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised etanercept treatment.

Authority required

Initial or re-Treatment [Initial 2, Whole body (Received prior biological agent under PBS)]

Treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have a documented history of severe chronic plaque psoriasis; and
- (b) have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have not failed PBS-subsidised therapy with etanercept for the treatment of this condition in the current Treatment Cycle.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)];
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

Applications for patients who have demonstrated a response to PBS-subsidised etanercept treatment within this Treatment Cycle and who wish to re-commence etanercept treatment within the same Cycle following a break in therapy, will only be approved where evidence of the patient's response to their most recent course of PBS-subsidised etanercept treatment has been submitted to Medicare Australia within 1 month of cessation of treatment.

A maximum of 16 weeks of treatment with etanercept will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 16 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised etanercept treatment.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Initial treatment [Initial 1, Face, hand, foot (New patients — No prior biological agent)]

Initial treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- (b) have not received any prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (face, hand, foot); and
- (d) have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 3 of the following 4 treatments:
 - (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or
 - (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or
 - (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or
 - (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities including severity, associated with phototherapy, methotrexate, cyclosporin and acitretin, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

application:

- (a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 16 weeks of treatment with etanercept will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 16 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised etanercept treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Authority required

Initial or re-Treatment [Initial 2, Face, hand, foot (Received prior biological agent under PBS)]

Treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot; and
- (b) have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have not failed PBS-subsidised therapy with etanercept for the treatment of this condition in the current Treatment Cycle.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

Applications for patients who have demonstrated a response to PBS-subsidised etanercept treatment within this Treatment Cycle and who wish to re-commence etanercept treatment within the same Cycle following a break in therapy, will only be approved where evidence of the patient's response to their most recent course of PBS-subsidised etanercept treatment has been submitted to Medicare Australia within 1 month of cessation of treatment.

A maximum of 16 weeks of treatment with etanercept will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 16 weeks.

A PASI assessment of the patient's response to this course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised etanercept treatment. The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline. Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle. <u>Note</u> No applications for increased maximum quantities and/or repeats will be authorised. <u>Note</u> Special Pricing Arrangements apply.							
9037P	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	3	..	*1829.00	34.20	Enbrel WX
9091L	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	3	..	1774.37	34.20	Enbrel WX
9461Y	Injection 50 mg in 1 mL single use auto-injector, 4	1	3	..	1774.37	34.20	Enbrel WX

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, etanercept, infliximab and ustekinumab, for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, infliximab and ustekinumab.

From 1 March 2010, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial adalimumab, etanercept, infliximab or ustekinumab without having to meet the initial treatment criteria, that is they will not need to experience a disease flare when swapping to an alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

A patient who received PBS-subsidised biological agent treatment for chronic plaque psoriasis prior to 1 March 2010 is considered to be in their first Cycle as of 1 March 2010.

Patients are eligible for PBS-subsidised treatment with only 1 biological agent at any 1 time.

Within the same Treatment Cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for a PBS-subsidised biological agent, they must change to an alternate agent if they wish to continue PBS-subsidised biological treatment. A patient who, prior to 1 March 2010, was authorised to receive PBS-subsidised initial treatment for chronic plaque psoriasis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2010.

Patients must be assessed for response to each course of continuing treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a Treatment Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological agent therapy before they are eligible to commence the next Cycle. The 5-year break

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

is measured from the date of the last approval for PBS-subsidised biological agent treatment in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Treatment Cycle.

Patients for whom a break in PBS-subsidised therapy of less than 5 years duration has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe chronic plaque psoriasis after 1 March 2010.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made in the following situations:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); or
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under '(4) Swapping therapy' below]; or
- (iii) patients who wish to re-commence treatment following a break in PBS-subsidised therapy with that agent (Initial 2).

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of treatment in the case of adalimumab and etanercept, 22 weeks of treatment in the case of infliximab and 28 weeks of treatment in the case of ustekinumab.

Grandfather patients (adalimumab and ustekinumab only).

Applications for patients who commenced treatment with adalimumab prior to 1 March 2009 or ustekinumab prior to 26 November 2009, may be made for initial PBS-subsidised treatment as continuing therapy under the initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with a biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment. Approval will be based on the criteria included in the relevant restriction.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological agent, a PASI assessment must be conducted after at least 12 weeks of treatment. This assessment must be submitted to Medicare Australia within 1 month of the completion of this initial treatment course. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Application for continuing treatment.

Following the completion of an initial treatment course of a biological agent to which an adequate response has been demonstrated, patients may qualify to receive up to 24 weeks of continuing treatment with that biological agent. Patients are eligible to continue to receive continuous treatment with 24 week courses providing they continue to sustain a response.

For second and subsequent courses of PBS-subsidised treatment with adalimumab, etanercept, infliximab or ustekinumab it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to sustain a response to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

(4) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate agent within the same Treatment Cycle without having to requalify with respect to disease severity (i.e. a PASI score of greater than 15), or prior treatment requirements.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	

Patients may trial an alternate biological agent at any time, regardless of whether they are receiving therapy with a biological agent at the time of the application or not. However, they cannot swap to a particular agent if they have failed to respond to treatment with that particular agent within the same Cycle.

Patients who commenced treatment with adalimumab prior to 1 June 2009 or ustekinumab prior to 1 March 2010 access these interchangeability arrangements in the same way as patients who have not.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the agent being ceased.

(5) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a Treatment Cycle and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatment applications.

(6) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent Biological Treatment Cycle, following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment according to the criteria of the relevant restriction and index of disease severity. Patients must have had at least 1 prior treatment, as listed in the criteria, for a minimum of 6 weeks, and must have a PASI assessment conducted preferably whilst still on treatment, but no later than 1 month following cessation of treatment. The PASI assessment must be no older than 1 month at the time of application.

Authority required

Continuing treatment (Whole body)

Continuing PBS-subsidised treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over:

- (a) who have a documented history of severe chronic plaque psoriasis; and
- (b) whose most recent course of PBS-subsidised biological treatment for this condition in this Treatment Cycle was with etanercept; and
- (c) who have demonstrated an adequate response to their most recent course of treatment with etanercept.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-biological treatment baseline value for this Treatment Cycle.

This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with etanercept, the assessment of response must be after a minimum of 12 weeks of treatment with an initial course.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet along with the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with etanercept.

A maximum of 24 weeks of treatment with etanercept will be authorised under this restriction.

Where fewer than 5 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for continuing authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks.

A PASI assessment of the patient's response must be conducted within 4 weeks prior to completion of this course of treatment. This assessment must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised etanercept treatment.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle. Authority required Continuing treatment (Face, hand, foot) Continuing PBS-subsidised treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over: (a) who have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot; and (b) whose most recent course of PBS-subsidised biological treatment for this condition in this Treatment Cycle was with etanercept; and (c) who have demonstrated an adequate response to treatment with etanercept. An adequate response to etanercept treatment is defined as the plaque or plaques assessed prior to biological treatment showing: (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the pre-biological treatment baseline values; or (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the pre-biological treatment baseline value. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with etanercept, the assessment of response must be after a minimum of 12 weeks of treatment with an initial course. Applications for authorisation must be made in writing and must include: (a) a completed authority prescription form; and (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following: (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams along with the date of the assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]. The most recent PASI assessment must be no more than 1 month old at the time of application. A maximum of 24 weeks of treatment with etanercept will be authorised under this restriction. Where fewer than 5 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for continuing authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks. A PASI assessment of the patient's response must be conducted within 4 weeks prior to completion of this course of treatment. This assessment must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss. It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised etanercept treatment. The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline. Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle. Note No applications for increased maximum quantities and/or repeats will be authorised. Note Special Pricing Arrangements apply.								
9429G	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	5	..	*1829.00	34.20	Enbrel	WX
9431J	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	5	..	1774.37	34.20	Enbrel	WX
9462B	Injection 50 mg in 1 mL single use auto-injector, 4	1	5	..	1774.37	34.20	Enbrel	WX

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and etanercept for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term bDMARD appears in the following NOTES and restrictions, it refers to adalimumab and etanercept only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 bDMARDs at any one time.

From 1 November 2010, a patient receiving PBS-subsidised bDMARD therapy is considered to be in a treatment cycle where they may swap to the alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy, and
- fail to respond, or to sustain a response to one PBS-subsidised bDMARD twice and the other PBS-subsidised bDMARD once only.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised bDMARD therapy before they are eligible to receive further PBS-subsidised bDMARD therapy. The length of a treatment break is measured from the date of the last approval for PBS-subsidised bDMARD therapy in the last treatment cycle and the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

A patient who was receiving PBS-subsidised bDMARD treatment immediately prior to 1 November 2010 is considered to be in their first cycle as of 1 November 2010. A patient who has had a break in bDMARD treatment of at least 12 months immediately prior to making a new application, on or after 1 November 2010, will commence a new treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle. The length of the break in therapy is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment was stopped to the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 November 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

A patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to the alternate bDMARD without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months.

A patient may trial the alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug twice within the current treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a bDMARD. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to the revised baseline measurement.

(4) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 12 months, must requalify for treatment under the Initial 1 treatment restriction.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for severe active juvenile idiopathic arthritis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must qualify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 12 month break in PBS-subsidised therapy' above for further details.

Authority required

Initial 1 (new patient or patient recommencing after a break of more than 12 months).

Initial treatment, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of a patient aged 18 years or older who:

- (a) has a documented history of juvenile idiopathic arthritis with onset prior to the age of 18 years; AND
- (b) has received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) has failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	— hydroxychloroquine at a dose of at least 200 mg daily; or — leflunomide at a dose of at least 10 mg daily; or — sulfasalazine at a dose of at least 2 g daily.						

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg per day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) an active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 16 weeks of treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
	time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment.							
	Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept.							
	If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial etanercept after a minimum of 5 years have elapsed between the date of the last approval for PBS-subsidised bDMARD therapy in the last treatment cycle and the date of the first application under a new treatment cycle.							
	Authority required							
	Initial 2 (change or re-commencement after break of less than 12 months).							
	Initial PBS-subsidised treatment with etanercept by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of a patient aged 18 years or older who:							
	(a) has a documented history of juvenile idiopathic arthritis with onset prior to the age of 18 years; AND							
	(b) in this treatment cycle, has received prior PBS-subsidised treatment with adalimumab or etanercept for this condition; and							
	(c) has not failed PBS-subsidised therapy with etanercept for this condition more than once in the current treatment cycle.							
	The authority application must be made in writing and must include:							
	(a) a completed authority prescription form; and							
	(b) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].							
	Applications for a patient who has received PBS-subsidised treatment with etanercept in this treatment cycle and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised etanercept treatment, within the timeframes specified below.							
	A maximum of 16 weeks of treatment will be authorised under this restriction.							
	Where fewer than 3 repeats are requested at the time of the initial authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with etanercept may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).							
	Where the most recent course of PBS-subsidised etanercept treatment was approved under either of the initial 1 or 2 treatment restrictions, the patient must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.							
	Where the most recent course of PBS-subsidised etanercept treatment was approved under the continuing treatment criteria, the patient must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.							
	Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to that particular course of bDMARD.							
	If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial etanercept after a minimum of 5 years have elapsed between the date of the last approval for PBS-subsidised bDMARD therapy in the last treatment cycle and the date of the first application under a new treatment cycle.							
3445H	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	3	..	*1829.00	34.20	Enbrel	WX
3446J	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	3	..	1774.37	34.20	Enbrel	WX
3447K	Injection 50 mg in 1 mL single use auto-injector, 4	1	3	..	1774.37	34.20	Enbrel	WX

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	
	HOBART TAS 7001						

;

Note

TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and etanercept for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term bDMARD appears in the following NOTES and restrictions, it refers to adalimumab and etanercept only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 bDMARDs at any one time.

From 1 November 2010, a patient receiving PBS-subsidised bDMARD therapy is considered to be in a treatment cycle where they may swap to the alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy, and
- fail to respond, or to sustain a response to one PBS-subsidised bDMARD twice and the other PBS-subsidised bDMARD once only.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised bDMARD therapy before they are eligible to receive further PBS-subsidised bDMARD therapy. The length of a treatment break is measured from the date of the last approval for PBS-subsidised bDMARD therapy in the last treatment cycle and the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

A patient who was receiving PBS-subsidised bDMARD treatment immediately prior to 1 November 2010 is considered to be in their first cycle as of 1 November 2010. A patient who has had a break in bDMARD treatment of at least 12 months immediately prior to making a new application, on or after 1 November 2010, will commence a new treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle. The length of the break in therapy is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment was stopped to the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 November 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

A patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to the alternate bDMARD without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months.

A patient may trial the alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug twice within the current treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a bDMARD. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to the revised baseline measurement.

(4) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 12 months, must requalify for treatment under the Initial 1 treatment restriction.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for severe active juvenile idiopathic arthritis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must qualify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 12 month break in PBS-subsidised therapy' above for further details.

Authority required

Continuing treatment.

Continuing PBS-subsidised treatment, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of a patient aged 18 years or older:

- (a) who has a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years; and
- (b) who has demonstrated an adequate response to treatment with etanercept; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with etanercept.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;
AND either of the following:

- (i) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (ii) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (iii) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
A maximum of 24 weeks of treatment will be approved under this restriction.							
Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).							
All applications for continuing treatment with etanercept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with etanercept, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.							
If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial etanercept after a minimum of 5 years have elapsed between the date of the last approval for PBS-subsidised bDMARD therapy in the last treatment cycle and the date of the first application under a new treatment cycle.							
3448L	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	2	5	..	*1829.00	34.20	Enbrel WX
3449M	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	5	..	1774.37	34.20	Enbrel WX
3450N	Injection 50 mg in 1 mL single use auto-injector, 4	1	5	..	1774.37	34.20	Enbrel WX

GOLIMUMAB

Note

Any queries concerning the arrangements to prescribe golimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe golimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF- α antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (new patient or patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with golimumab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 16 weeks of treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with golimumab.

Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with golimumab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

- (a) have a documented history of severe active rheumatoid arthritis; and
(b) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with golimumab and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised golimumab treatment, within the timeframes specified below.

A maximum of 16 weeks of treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised golimumab treatment was approved under either of the initial 1 or 2 treatment restrictions, patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised golimumab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

3426H	Injection 50 mg in 0.5 mL single use pre-filled syringe	1	3	..	1777.29	34.20	Simponi	SH
3427J	Injection 50 mg in 0.5 mL single use pre-filled pen	1	3	..	1777.29	34.20	Simponi	SH

GOLIMUMAB

Note

Any queries concerning the arrangements to prescribe golimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe golimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 3 ('grandfather' patients)

Initial PBS-subsidised supply for continuing treatment with golimumab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of an adult who:

- (a) has a documented history of severe active rheumatoid arthritis; and
- (b) was receiving treatment with golimumab prior to 1 March 2010; and
- (c) has demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with golimumab; and
- (d) is receiving treatment with golimumab at the time of application.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au]; and
- (3) a signed patient acknowledgement.

The same indices of disease severity used to establish baseline at the commencement of treatment with a bDMARD must be used for assessment of all continuing applications.

The assessment of the patient's response to a continuing course of therapy must be made within 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled in order to ensure continuity of treatment for those patients who meet the continuation criterion.

A maximum of 24 weeks of treatment with golimumab will be approved under this criterion.

Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with golimumab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have a documented history of severe active rheumatoid arthritis; and
- (b) who have demonstrated an adequate response to treatment with golimumab; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with golimumab.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with golimumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with golimumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

3428K	Injection 50 mg in 0.5 mL single use pre-filled syringe	1	5	..	1777.29	34.20	Simponi	SH
3429L	Injection 50 mg in 0.5 mL single use pre-filled pen	1	5	..	1777.29	34.20	Simponi	SH

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

GOLIMUMAB

Note

Any queries concerning the arrangements to prescribe golimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe golimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents (adalimumab, etanercept, golimumab and infliximab) for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time. Where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, golimumab and infliximab.

From 1 August 2006, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis after 1 August 2010.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients — golimumab only.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Applications for patients who commenced treatment with golimumab prior to 1 March 2010 may apply for initial PBS-subsidised treatment as continuing therapy under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialed it on the PBS; or
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment Cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Authority required

Initial 1

Initial PBS-subsidised treatment with golimumab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults who:

- (1) have severe active psoriatic arthritis; and
- (2) have received no prior PBS-subsidised biological treatment for this condition in this Treatment Cycle; and
- (3) have failed to achieve an adequate response to:
 - (a) methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months; and
 - (b) sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; or
 - (c) leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities, including severity, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

(i) an active joint count of at least 20 active (swollen and tender) joints; or

(ii) at least 4 active joints from the following list of major joints:

— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and

(3) a signed patient acknowledgement.

A maximum of 16 weeks treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

The assessment of the patient's response to the initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial golimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Initial 2

Initial PBS-subsidised treatment with golimumab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults who:

(1) have a documented history of severe active psoriatic arthritis; and

(2) have received prior PBS-subsidised biological treatment for this condition in this Treatment Cycle and are eligible to receive further biological therapy; and

(3) have not failed treatment with golimumab during the current Treatment Cycle.

Applications for patients who have received PBS-subsidised treatment with golimumab within this Treatment Cycle and who wish to re-commence therapy with this drug within this same Cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised golimumab treatment, within the timeframes specified below.

A maximum of 16 weeks treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised golimumab treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised biological therapy or, under this restriction, for patients who have received previous PBS-subsidised biological therapy), patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised golimumab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial golimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.								
<u>Note</u> No applications for increased maximum quantities and/or repeats will be authorised. Applications for treatment with golimumab where the dosing frequency exceeds 50 mg every 4 weeks will not be approved.								
3430M	Injection 50 mg in 0.5 mL single use pre-filled syringe	1	3	..	1777.29	34.20	Simponi	SH
3431N	Injection 50 mg in 0.5 mL single use pre-filled pen	1	3	..	1777.29	34.20	Simponi	SH

GOLIMUMAB

Note

Any queries concerning the arrangements to prescribe golimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe golimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents (adalimumab, etanercept, golimumab and infliximab) for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time. Where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, golimumab and infliximab.

From 1 August 2006, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis after 1 August 2010.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

therapy' below]; and

(iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients — golimumab only.

Applications for patients who commenced treatment with golimumab prior to 1 March 2010 may apply for initial PBS-subsidised treatment as continuing therapy under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; or
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment Cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.						
	Authority required Initial 3 — grandfather golimumab patients Initial PBS-subsidised supply for continuing treatment with golimumab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults who: (1) have a documented history of severe active psoriatic arthritis; and (2) were receiving treatment with golimumab prior to 1 March 2010; and (3) have demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with golimumab; and (4) are receiving treatment with golimumab at the time of application. The authority application must be made in writing and must include: (1) a completed authority prescription form; and (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and (3) a signed patient acknowledgement. A maximum of 24 weeks of treatment with golimumab will be authorised under this restriction. Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial golimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle. Patients may qualify for PBS-subsidised treatment under this restriction once only.						
	Authority required Continuing treatment Continuing PBS-subsidised treatment with golimumab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults: (1) who have a documented history of severe active psoriatic arthritis; and (2) whose most recent course of PBS-subsidised biological agent for this condition in the current Treatment Cycle was with golimumab; and (3) who, at the time of application, demonstrate an adequate response to treatment with golimumab. An adequate response to treatment with golimumab is defined as: an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; AND either of the following: (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%: — elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or — shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth). The authority application must be made in writing and must include: (1) a completed authority prescription form; and (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]. A maximum of 24 weeks of treatment will be approved under this restriction. Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). All applications for continuing treatment with golimumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with golimumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course. Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial golimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.						
	Note No applications for increased maximum quantities and/or repeats will be authorised. Applications for treatment with golimumab where the dosing frequency exceeds 50 mg every 4 weeks will not be approved.						
3432P	Injection 50 mg in 0.5 mL single use pre-filled	1	5	..	1777.29	34.20	Simponi SH

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
3433Q	syringe Injection 50 mg in 0.5 mL single use pre-filled pen	1	5	..	1777.29	34.20	Simponi SH

GOLIMUMAB

Note

Any queries concerning the arrangements to prescribe golimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe golimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH ACTIVE ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept, golimumab and infliximab for adult patients with active ankylosing spondylitis. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab, etanercept, golimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 4 TNF-alfa antagonists at any 1 time.

From 1 March 2007, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised TNF-alfa antagonists without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 March 2007 is considered to be in their first cycle as of 1 March 2007.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than once. A patient who, prior to 1 March 2007, was authorised to receive PBS-subsidised initial treatment for ankylosing spondylitis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2007.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and 18 weeks of treatment for infliximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap to an alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with at least 1 NSAID, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the BASDAI, ESR and/or CRP levels are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for active ankylosing spondylitis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with golimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with golimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial 1 (new patients)

Initial PBS-subsidised treatment with golimumab, by a rheumatologist, of an adult with active ankylosing spondylitis who has radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis and who has not received any PBS-subsidised treatment with either adalimumab, etanercept, golimumab or infliximab in this treatment cycle; AND

(a) who has at least 2 of the following:

- (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or
- (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI) [for further information on the BASMI please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; or
- (iii) limitation of chest expansion relative to normal values for age and gender [for chest expansion normal values please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; AND

(b) who has failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months.

The application must include details of the NSAIDs trialled, their doses and duration of treatment. If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance. Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Medicare Australia website at www.medicareaustralia.gov.au.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au] which must include the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form [www.medicareaustralia.gov.au]; and
 - (iii) a completed Exercise Program Self Certification Form included in the supporting information form; and
 - (iv) a signed patient acknowledgment form.

The assessment of the patient's response to the initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A maximum of 16 weeks of treatment with golimumab will be approved under this criterion.

Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial golimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.

Authority required

Initial 2 (change or re-commencement for all patients)

Initial PBS-subsidised treatment with golimumab, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who, in this treatment cycle, has received prior PBS-subsidised TNF-alfa antagonist treatment for this condition and is eligible to receive further TNF-alfa antagonist therapy, and has not failed PBS-subsidised therapy with golimumab in the current treatment cycle.

Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised TNF-alfa antagonist therapy or, under this restriction, for patients who have received previous PBS-

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
subsidised TNF-alfa antagonist therapy) the patient must have been assessed for response to that course following a minimum of 12 weeks of treatment. These assessments must be provided to Medicare Australia no later than 4 weeks from the date the course was ceased. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.							
Where the most recent course of PBS-subsidised golimumab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.							
Authority applications must be made in writing and must include: (a) a completed authority prescription form; and (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au].							
A maximum of 16 weeks of treatment with golimumab will be approved under this criterion.							
Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.							
Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial golimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.							
Note No applications for increased maximum quantities and/or repeats will be authorised. Applications for treatment with golimumab where the dosing frequency exceeds 50 mg every 4 weeks will not be approved.							
3434R	Injection 50 mg in 0.5 mL single use pre-filled syringe	1	3	..	1777.29	34.20	Simponi SH
3435T	Injection 50 mg in 0.5 mL single use pre-filled pen	1	3	..	1777.29	34.20	Simponi SH

GOLIMUMAB

Note

Any queries concerning the arrangements to prescribe golimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe golimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH ACTIVE ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept, golimumab and infliximab for adult patients with active ankylosing spondylitis. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab, etanercept, golimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 4 TNF-alfa antagonists at any 1 time.

From 1 March 2007, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised TNF-alfa antagonists without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 March 2007 is considered to be in their first cycle as of 1 March 2007.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than once. A patient who, prior to 1 March 2007, was authorised to receive PBS-subsidised initial treatment for ankylosing spondylitis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2007.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and 18 weeks of treatment for infliximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap to an alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty	Safety Net	
					\$	\$	
For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.							

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with at least 1 NSAID, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the BASDAI, ESR and/or CRP levels are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for active ankylosing spondylitis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with golimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with golimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial ('grandfather' patients)

Initial PBS-subsidised supply for continuing treatment with golimumab, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who has radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis and who was receiving treatment with golimumab prior to 1 March 2010; and

- (a) has demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with golimumab; and
- (b) is receiving treatment with golimumab at the time of application.

The BASDAI assessment and ESR and/or CRP measurements provided must be no more than 1 month old at the time of application. Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au] which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form [www.medicareaustralia.gov.au]; and
 - (iii) a signed patient acknowledgment form.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of the course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is due in order to ensure continuity of treatment for those patients who meet the continuation criteria. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A patient ceasing treatment or swapping to an alternate agent and wishing to demonstrate a response to treatment, must be assessed no earlier than 12 weeks from the commencement of PBS-subsidised treatment. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A maximum of 24 weeks of treatment with golimumab will be authorised under this criterion.

Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.

Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial golimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.

Patients may only qualify for PBS-subsidised treatment under this criterion once.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
<u>Authority required</u>							
Continuing treatment for all patients							
Continuing PBS-subsidised treatment, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who:							
(a) has demonstrated an adequate response to treatment with golimumab; and							
(b) whose most recent course of PBS-subsidised therapy in this treatment cycle was with golimumab.							
An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:							
(a) an ESR measurement no greater than 25 mm per hour; or							
(b) a CRP measurement no greater than 10 mg per L; or							
(c) an ESR or CRP measurement reduced by at least 20% from baseline.							
Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.							
Authority applications must be made in writing and must include:							
(a) a completed authority prescription form; and							
(b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au].							
All measurements provided must be no more than 1 month old at the time of application.							
A maximum of 24 weeks of treatment with golimumab will be authorised under this criterion.							
Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.							
All applications for continuing treatment with golimumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment following an initial treatment course it must be made following a minimum of 12 weeks of treatment with golimumab. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.							
Patients who fail to demonstrate a response to treatment with golimumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial golimumab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised. Applications for treatment with golimumab where the dosing frequency exceeds 50 mg every 4 weeks will not be approved.							
3436W	Injection 50 mg in 0.5 mL single use pre-filled syringe	1	5	..	1777.29	34.20	Simponi SH
3437X	Injection 50 mg in 0.5 mL single use pre-filled pen	1	5	..	1777.29	34.20	Simponi SH

Interleukin inhibitors

USTEKINUMAB

Note

Any queries concerning the arrangements to prescribe ustekinumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe ustekinumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, etanercept, infliximab and ustekinumab, for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, infliximab and ustekinumab.

From 1 March 2010, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial adalimumab, etanercept, infliximab or ustekinumab without having to meet the initial treatment criteria, that is they will not need to experience a disease flare when swapping to an alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
A patient who received PBS-subsidised biological agent treatment for chronic plaque psoriasis prior to 1 March 2010 is considered to be in their first Cycle as of 1 March 2010.							

Patients are eligible for PBS-subsidised treatment with only 1 biological agent at any 1 time.

Within the same Treatment Cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for a PBS-subsidised biological agent, they must change to an alternate agent if they wish to continue PBS-subsidised biological treatment. A patient who, prior to 1 March 2010, was authorised to receive PBS-subsidised initial treatment for chronic plaque psoriasis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2010.

Patients must be assessed for response to each course of continuing treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a Treatment Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological agent therapy before they are eligible to commence the next Cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological agent treatment in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Treatment Cycle.

Patients for whom a break in PBS-subsidised therapy of less than 5 years duration has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe chronic plaque psoriasis after 1 March 2010.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made in the following situations:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); or
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under '(4) Swapping therapy' below]; or
- (iii) patients who wish to re-commence treatment following a break in PBS-subsidised therapy with that agent (Initial 2).

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of treatment in the case of adalimumab and etanercept, 22 weeks of treatment in the case of infliximab and 28 weeks of treatment in the case of ustekinumab.

Grandfather patients (adalimumab and ustekinumab only).

Applications for patients who commenced treatment with adalimumab prior to 1 March 2009 or ustekinumab prior to 26 November 2009, may be made for initial PBS-subsidised treatment as continuing therapy under the initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with a biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment. Approval will be based on the criteria included in the relevant restriction.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological agent, a PASI assessment must be conducted after at least 12 weeks of treatment. This assessment must be submitted to Medicare Australia within 1 month of the completion of this initial treatment course. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Application for continuing treatment.

Following the completion of an initial treatment course of a biological agent to which an adequate response has been demonstrated, patients may qualify to receive up to 24 weeks of continuing treatment with that biological agent. Patients are eligible to continue to receive continuous treatment with 24 week courses providing they continue to sustain a response.

For second and subsequent courses of PBS-subsidised treatment with adalimumab, etanercept, infliximab or ustekinumab it is recommended that a

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.							

Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to sustain a response to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

(4) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate agent within the same Treatment Cycle without having to requalify with respect to disease severity (i.e. a PASI score of greater than 15), or prior treatment requirements.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

Patients may trial an alternate biological agent at any time, regardless of whether they are receiving therapy with a biological agent at the time of the application or not. However, they cannot swap to a particular agent if they have failed to respond to treatment with that particular agent within the same Cycle.

Patients who commenced treatment with adalimumab prior to 1 June 2009 or ustekinumab prior to 1 March 2010 access these interchangeability arrangements in the same way as patients who have not.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the agent being ceased.

(5) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a Treatment Cycle and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatment applications.

(6) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent Biological Treatment Cycle, following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment according to the criteria of the relevant restriction and index of disease severity. Patients must have had at least 1 prior treatment, as listed in the criteria, for a minimum of 6 weeks, and must have a PASI assessment conducted preferably whilst still on treatment, but no later than 1 month following cessation of treatment. The PASI assessment must be no older than 1 month at the time of application.

Authority required

Initial treatment [Initial 1, Whole body (New patients — No prior biological agent)]

Initial treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis; and
- (b) have not received any prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (whole body); and
- (d) have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 3 of the following 4 treatments:
 - (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or
 - (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or
 - (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or
 - (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities including severity, associated with phototherapy, methotrexate, cyclosporin and acitretin, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

month following cessation of the most recent prior treatment.

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]]; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
(iii) the signed patient and prescriber acknowledgements.

A maximum of 28 weeks of treatment with ustekinumab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 28 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 28 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with ustekinumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised ustekinumab treatment.

Note

No applications for increased repeats will be authorised.

Authority required

Initial or re-Treatment [Initial 2, Whole body (Received prior biological agent under PBS)]

Treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

(a) have a documented history of severe chronic plaque psoriasis; and

(b) have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and

(c) have not failed PBS-subsidised therapy with ustekinumab for the treatment of this condition in the current Treatment Cycle.

Applications for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:

(i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]]; and

(ii) details of prior biological treatment, including dosage, date and duration of treatment.

Applications for patients who have demonstrated a response to PBS-subsidised ustekinumab treatment within this Treatment Cycle and who wish to re-commence ustekinumab treatment within the same Cycle following a break in therapy, will only be approved where evidence of the patient's response to their most recent course of PBS-subsidised ustekinumab treatment has been submitted to Medicare Australia within 1 month of cessation of treatment.

A maximum of 28 weeks of treatment with ustekinumab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 28 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 28 weeks.

A PASI assessment of the patient's response to this course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with ustekinumab.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.							

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised ustekinumab treatment.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

No applications for increased repeats will be authorised.

Authority required

Initial treatment [Initial 1, Face, hand, foot (New patients — No prior biological agent)]

Initial treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- (b) have not received any prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (face, hand, foot); and
- (d) have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 3 of the following 4 treatments:
 - (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or
 - (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or
 - (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or
 - (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities including severity, associated with phototherapy, methotrexate, cyclosporin and acitretin, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 28 weeks of treatment with ustekinumab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 28 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 28 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
treatment with ustekinumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.							

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised ustekinumab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Note

No applications for increased repeats will be authorised.

Authority required

Initial or re-Treatment [Initial 2, Face, hand, foot (Received prior biological agent under PBS)]

Treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot; and
- (b) have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have not failed PBS-subsidised therapy with ustekinumab for the treatment of this condition in the current Treatment Cycle.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

Applications for patients who have demonstrated a response to PBS-subsidised ustekinumab treatment within this Treatment Cycle and who wish to re-commence ustekinumab treatment within the same Cycle following a break in therapy, will only be approved where evidence of the patient's response to their most recent course of PBS-subsidised ustekinumab treatment has been submitted to Medicare Australia within 1 month of cessation of treatment.

A maximum of 28 weeks of treatment with ustekinumab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 28 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 28 weeks.

A PASI assessment of the patient's response to this course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with ustekinumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised ustekinumab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

No applications for increased repeats will be authorised.

Note

Special Pricing Arrangements apply.

9304Q	Injection 45 mg in 0.5 mL	1	2	..	4601.42	34.20	Stelara	JC

USTEKINUMAB

Note

Any queries concerning the arrangements to prescribe ustekinumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe ustekinumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, etanercept, infliximab and ustekinumab, for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, infliximab and ustekinumab.

From 1 March 2010, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial adalimumab, etanercept, infliximab or ustekinumab without having to meet the initial treatment criteria, that is they will not need to experience a disease flare when swapping to an alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

A patient who received PBS-subsidised biological agent treatment for chronic plaque psoriasis prior to 1 March 2010 is considered to be in their first Cycle as of 1 March 2010.

Patients are eligible for PBS-subsidised treatment with only 1 biological agent at any 1 time.

Within the same Treatment Cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for a PBS-subsidised biological agent, they must change to an alternate agent if they wish to continue PBS-subsidised biological treatment. A patient who, prior to 1 March 2010, was authorised to receive PBS-subsidised initial treatment for chronic plaque psoriasis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2010.

Patients must be assessed for response to each course of continuing treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a Treatment Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological agent therapy before they are eligible to commence the next Cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological agent treatment in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Treatment Cycle.

Patients for whom a break in PBS-subsidised therapy of less than 5 years duration has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe chronic plaque psoriasis after 1 March 2010.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made in the following situations:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); or
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under '(4) Swapping therapy' below]; or
- (iii) patients who wish to re-commence treatment following a break in PBS-subsidised therapy with that agent (Initial 2).

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of treatment in the case of adalimumab and etanercept, 22 weeks of treatment in the case of infliximab and 28 weeks of treatment in the case of ustekinumab.

Grandfather patients (adalimumab and ustekinumab only).

Applications for patients who commenced treatment with adalimumab prior to 1 March 2009 or ustekinumab prior to 26 November 2009, may be made for initial PBS-subsidised treatment as continuing therapy under the initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with a biological agent prior to PBS listing of that

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment. Approval will be based on the criteria included in the relevant restriction.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological agent, a PASI assessment must be conducted after at least 12 weeks of treatment. This assessment must be submitted to Medicare Australia within 1 month of the completion of this initial treatment course. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Application for continuing treatment.

Following the completion of an initial treatment course of a biological agent to which an adequate response has been demonstrated, patients may qualify to receive up to 24 weeks of continuing treatment with that biological agent. Patients are eligible to continue to receive continuous treatment with 24 week courses providing they continue to sustain a response.

For second and subsequent courses of PBS-subsidised treatment with adalimumab, etanercept, infliximab or ustekinumab it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to sustain a response to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

(4) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate agent within the same Treatment Cycle without having to requalify with respect to disease severity (i.e. a PASI score of greater than 15), or prior treatment requirements.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

Patients may trial an alternate biological agent at any time, regardless of whether they are receiving therapy with a biological agent at the time of the application or not. However, they cannot swap to a particular agent if they have failed to respond to treatment with that particular agent within the same Cycle.

Patients who commenced treatment with adalimumab prior to 1 June 2009 or ustekinumab prior to 1 March 2010 access these interchangeability arrangements in the same way as patients who have not.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the agent being ceased.

(5) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a Treatment Cycle and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatment applications.

(6) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent Biological Treatment Cycle, following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment according to the criteria of the relevant restriction and index of disease severity. Patients must have had at least 1 prior treatment, as listed in the criteria, for a minimum of 6 weeks, and must have a PASI assessment conducted preferably whilst still on treatment, but no later than 1 month following cessation of treatment. The PASI assessment must be no older than 1 month at the time of application.

Authority required

Initial treatment [Initial 3, Whole body (Grandfather patients)]

Initial PBS-subsidised supply for continuing treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

and over who:

- (a) have a documented history of severe chronic plaque psoriasis and were receiving treatment with ustekinumab prior to 26 November 2009; and
- (b) had a Psoriasis Area and Severity Index (PASI) score of greater than 15 prior to commencing treatment with ustekinumab; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (whole body); and
- (d) have demonstrated a response as specified in the criterion included in the restriction for continuing PBS-subsidised treatment with ustekinumab (whole body).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition at baseline (prior to initiation of ustekinumab therapy) and the most recent PASI assessment [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 24 weeks of treatment with ustekinumab will be authorised under this restriction.

Where fewer than 1 repeat is requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks.

A PASI assessment of the patient's response to this initial PBS-subsidised course of therapy must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with ustekinumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The most recent PASI assessment must be no more than 1 month old at the time of application.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised ustekinumab treatment.

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Note

No applications for increased repeats will be authorised.

Authority required

Continuing treatment (Whole body)

Continuing PBS-subsidised treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over:

- (a) who have a documented history of severe chronic plaque psoriasis; and
- (b) whose most recent course of PBS-subsidised biological treatment for this condition in this Treatment Cycle was with ustekinumab; and
- (c) who have demonstrated an adequate response to their most recent course of treatment with ustekinumab.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-biological treatment baseline value for this Treatment Cycle.

This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with ustekinumab, the assessment of response must be after a minimum of 12 weeks of treatment with an initial course.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet along with the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with ustekinumab.

A maximum of 24 weeks of treatment with ustekinumab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

provide sufficient for a single injection. Up to a maximum of 1 repeat will be authorised.

Where fewer than 1 repeat is requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for continuing authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks.

A PASI assessment of the patient's response must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with ustekinumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised ustekinumab treatment.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

No applications for increased repeats will be authorised.

Authority required

Initial treatment [Initial 3, Face, hand, foot (Grandfather patients)]

Initial PBS-subsidised supply for continuing treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over:

- (a) who have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot and were receiving treatment with ustekinumab prior to 26 November 2009; and
- (b) whose disease, prior to treatment with ustekinumab, was of a severity as defined in the initiation criterion included in the initial treatment restriction (Initial 1, New patients — face, hand, foot); and
- (c) who have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (face, hand, foot); and
- (d) who have demonstrated a response as specified in the criterion included in the restriction for continuing PBS-subsidised treatment with ustekinumab (face, hand, foot).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams along with the date of the assessment of the patient's condition at baseline (prior to initiation of ustekinumab therapy) and the most recent PASI assessment [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 24 weeks of treatment with ustekinumab will be authorised under this restriction.

Where fewer than 1 repeat is requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 24 weeks.

A PASI assessment of the patient's response to this initial PBS-subsidised course of therapy must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with ustekinumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The most recent PASI assessment must be no more than 1 month old at the time of application.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised ustekinumab treatment.

The PASI assessment must be performed on the same affected area as assessed at baseline or prior to initiation of ustekinumab treatment.

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
<u>Note</u> No applications for increased repeats will be authorised. <u>Authority required</u> Continuing treatment (Face, hand, foot) Continuing PBS-subsidised treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over: (a) who have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot; and (b) whose most recent course of PBS-subsidised biological treatment for this condition in this Treatment Cycle was with ustekinumab; and (c) who have demonstrated an adequate response to treatment with ustekinumab. An adequate response to ustekinumab treatment is defined as the plaque or plaques assessed prior to biological treatment showing: (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the pre-biological treatment baseline values; or (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the pre-biological treatment baseline value. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with ustekinumab, the assessment of response must be after a minimum of 12 weeks of treatment with an initial course. Applications for authorisation must be made in writing and must include: (a) a completed authority prescription form; and (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following: (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams along with the date of the assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]. The most recent PASI assessment must be no more than 1 month old at the time of application. A maximum of 24 weeks of treatment with ustekinumab will be authorised under this restriction. At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 1 repeat will be authorised. Where fewer than 1 repeat is requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for continuing authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks. A PASI assessment of the patient's response must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with ustekinumab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss. It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised ustekinumab treatment. The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline. Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle. <u>Note</u> No applications for increased repeats will be authorised. <u>Note</u> Special Pricing Arrangements apply.							
9305R	Injection 45 mg in 0.5 mL	1	1	..	4601.42	34.20	Stelara JC

Calcineurin inhibitors

CYCLOSPORIN

Caution

Careful monitoring of patients is mandatory.

Authority required

Maintenance therapy, following initiation and stabilisation of treatment with cyclosporin, of:

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
(a) patients with organ or tissue transplants. Therapy must remain under the supervision and direction of the transplant unit reviewing the patient. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application; (b) patients with severe atopic dermatitis for whom other systemic therapies are ineffective or inappropriate. Therapy must remain under the supervision and direction of a dermatologist, clinical immunologist or specialised unit reviewing the patient. The name of the dermatologist, clinical immunologist or specialised unit reviewing treatment and the date of the latest review must be included in the authority application; (c) patients with severe psoriasis for whom other systemic therapies are ineffective or inappropriate and in whom the disease has caused significant interference with quality of life. Therapy must remain under the supervision and direction of a dermatologist or specialised unit reviewing the patient. The name of the dermatologist or specialised unit reviewing treatment and the date of the latest review must be included in the authority application; (d) patients with nephrotic syndrome in whom steroids and cytostatic drugs have failed or are not tolerated or are considered inappropriate and in whom renal function is unimpaired. Therapy must remain under the supervision and direction of a nephrologist or specialised unit reviewing the patient. The name of the nephrologist or specialised unit reviewing treatment and the date of the latest review must be included in the authority application; (e) patients with severe active rheumatoid arthritis for whom classical slow-acting anti-rheumatic agents (including methotrexate) are ineffective or inappropriate. Therapy must remain under the supervision and direction of a rheumatologist, clinical immunologist or specialised unit reviewing the patient. The name of the rheumatologist, clinical immunologist or specialised unit reviewing treatment and the date of the latest review must be included in the authority application; Management (which includes initiation, stabilisation and review of therapy) by dermatologists or clinical immunologists of patients with severe atopic dermatitis for whom other systemic therapies are ineffective or inappropriate; Management (which includes initiation, stabilisation and review of therapy) by dermatologists of patients with severe psoriasis for whom other systemic therapies are ineffective or inappropriate and in whom the disease has caused significant interference with quality of life; Management (which includes initiation, stabilisation and review of therapy) by rheumatologists or clinical immunologists of patients with severe active rheumatoid arthritis for whom classical slow-acting anti-rheumatic agents (including methotrexate) are ineffective or inappropriate.							
8657P	Capsule 10 mg	120	3	..	*94.42	34.20	Neoral 10 NV
8658Q	Capsule 25 mg	60	3	..	*99.08	34.20	^a Cicloral SZ
				^B 2.20	*101.28	34.20	^a Neoral 25 NV
8659R	Capsule 50 mg	60	3	..	*199.22	34.20	^a Cicloral SZ
				^B 2.38	*201.60	34.20	^a Neoral 50 NV
8660T	Capsule 100 mg	60	3	..	*381.58	34.20	^a Cicloral SZ
				^B 2.16	*383.74	34.20	^a Neoral 100 NV
8661W	Oral liquid 100 mg per mL, 50 mL	2	3	..	*712.66	34.20	Neoral NV

TACROLIMUS

Caution

Careful monitoring of patients is mandatory.

Authority required

Maintenance therapy, following initiation and stabilisation of treatment with tacrolimus, of patients with organ or tissue transplants. Therapy must remain under the supervision and direction of the transplant unit reviewing the patient. The name of the specialised transplant unit reviewing treatment and the date of the latest review at the specialised transplant unit must be included in the authority application.

5299X	Capsule 0.5 mg (once daily prolonged release)	30	3	..	64.59	34.20	Prograf XL	JC
5300Y	Capsule 1 mg (once daily prolonged release)	60	3	..	235.91	34.20	Prograf XL	JC
5451X	Capsule 5 mg (once daily prolonged release)	30	3	..	556.32	34.20	Prograf XL	JC
8646C	Capsule 500 micrograms	100	3	..	200.30	34.20	^a Prograf	JC
						^a Tacrolimus Sandoz	SZ	
8647D	Capsule 1 mg	100	3	..	376.91	34.20	^a Prograf	JC
						^a Tacrolimus Sandoz	SZ	
8648E	Capsule 5 mg	50	3	..	922.44	34.20	^a Prograf	JC
						^a Tacrolimus Sandoz	SZ	

Other immunosuppressants

AZATHIOPRINE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2687K NP	Tablet 50 mg	100	2	..	56.50	34.20	^a Azamun GM

Antineoplastic and immunomodulating agents

					Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	\$	\$	Brand Name and Manufacturer	
2688L <i>NP</i>	Tablet 25 mg	100	2	..	37.18	34.20	^a Azapin	SI
							^a Azathioprine Sandoz	SZ
							^a GenRx Azathioprine	GX
							^a Imuran	AS
							^a Thioprine	AF
							^a Azathioprine Sandoz	SZ
							^a Imuran	AS
METHOTREXATE								
1622J	Tablet 2.5 mg	30	5	..	13.26	14.33	^a Hospira Pty Limited	HH
2272N	Tablet 10 mg	15	1	..	22.15	23.22	^a Methoblastin	PF
							Methoblastin	PF
METHOTREXATE								
<u>Restricted benefit</u>								
For patients requiring doses greater than 20 mg per week.								
1623K	Tablet 10 mg	50	2	..	45.98	34.20	Methoblastin	PF

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer

Musculo-skeletal system

Antiinflammatory and antirheumatic products

Antiinflammatory and antirheumatic products, non-steroids

Acetic acid derivatives and related substances

1302M NP,MW	DICLOFENAC SODIUM Suppository 100 mg	40	3	..	*24.92	25.99	Voltaren 100	NV
	DICLOFENAC SODIUM <u>Restricted benefit</u> Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.							
1299J NP	Tablet 25 mg (enteric coated)	100	3	..	*13.06	14.13	^a APO-Diclofenac	TX
							^a Chem mart	CH
							^a Diclofenac	
							^a Clonac 25	SI
							^a Diclofenac-GA	GM
							^a Diclofenac Sandoz	SZ
							^a Fenac 25	AF
							^a Terry White Chemists	TW
							^a Diclofenac	
				^B 1.96	*15.02	14.13	^a Voltaren 25	NV
1300K NP	Tablet 50 mg (enteric coated)	50	3	..	11.05	12.12	^a APO-Diclofenac	TX
							^a Chem mart	CH
							^a Diclofenac	
							^a Clonac 50	SI
							^a Diclofenac-GA	GM
							^a Diclofenac Sandoz	SZ
							^a Fenac	AF
							^a Terry White Chemists	TW
							^a Diclofenac	
				^B 1.96	13.01	12.12	^a Voltaren 50	NV
	INDOMETHACIN Suppository 100 mg	40	3	..	*22.50	23.57	Indocid	AS
2757D NP								
	INDOMETHACIN <u>Restricted benefit</u> Chronic arthropathies (including osteoarthritis) with an inflammatory component; Bone pain due to malignant disease.							
2454E NP	Capsule 25 mg	100	3	..	*12.08	13.15	^a Arthrexin	AF
				^B 2.16	*14.24	13.15	^a Indocid	AS

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
	Bone pain due to malignant disease.							
2047R NP	Tablet 100 mg	100	3	..	*16.34	17.41	Aclin	AF
2048T NP	Tablet 200 mg	50	3	..	15.28	16.35	Aclin 200	AF
Oxicams								
MELOXICAM								
Note								
The use of meloxicam for the treatment of the following conditions is not subsidised through the PBS:								
(a) acute pain;								
(b) soft tissue injury;								
(c) arthrosis without an inflammatory component.								
Restricted benefit								
Symptomatic treatment of osteoarthritis;								
Symptomatic treatment of rheumatoid arthritis.								
8561N NP	Tablet 7.5 mg	30	3	..	19.02	20.09	^a Chem mart Meloxicam 7.5 mg	CH
							^a GenRx Meloxicam	GX
							^a Meloxicam	BF
							^a Meloxicam-GA	GM
							^a Meloxicam Ranbaxy	RA
							^a Meloxicam Sandoz	SZ
							^a Meloxicam Winthrop	WA
							^a Movalis 7.5	SI
							^a Moxicam 7.5	AF
							^a Pharmacor Meloxicam 7.5	CR
							^a Terry White Chemists Meloxicam 7.5 mg	TW
				^B 1.30	20.32	20.09	^a Mobic	BY
8562P NP	Tablet 15 mg	30	3	..	24.81	25.88	^a Chem mart Meloxicam 15 mg	CH
							^a GenRx Meloxicam	GX
							^a Meloxicam	BF
							^a Meloxicam-GA	GM
							^a Meloxicam Ranbaxy	RA
							^a Meloxicam Sandoz	SZ
							^a Meloxicam Winthrop	WA
							^a Movalis 15	SI
							^a Moxicam 15	AF
							^a Pharmacor Meloxicam 15	CR
							^a Terry White Chemists Meloxicam 15 mg	TW
				^B 1.30	26.11	25.88	^a Mobic	BY
8887R NP	Capsule 7.5 mg	30	3	..	19.02	20.09	^a Mobic	BY
							^a Movalis 7.5	SI
8888T NP	Capsule 15 mg	30	3	..	24.81	25.88	^a Mobic	BY

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
							^a Movalis 15	SI
PIROXICAM								
<u>Restricted benefit</u>								
Chronic arthropathies (including osteoarthritis) with an inflammatory component.								
1895R NP	Dispersible tablet 10 mg	50	3	..	12.50	13.57	Mobilis D-10	AF
1896T NP	Dispersible tablet 20 mg	25	3	..	12.20	13.27	^a Mobilis D-20	AF
				^B 2.64	14.84	13.27	^a Feldene-D	PF
1897W NP	Capsule 10 mg	50	3	..	12.50	13.57	^a Chem mart Piroxicam	CH
							^a GenRx Piroxicam	GX
							^a Mobilis 10	AF
							^a Terry White Chemists Piroxicam	TW
				^B 2.66	15.16	13.57	^a Feldene	PF
1898X NP	Capsule 20 mg	25	3	..	12.20	13.27	^a Chem mart Piroxicam	CH
							^a GenRx Piroxicam	GX
							^a Mobilis 20	AF
							^a Terry White Chemists Piroxicam	TW
				^B 2.64	14.84	13.27	^a Feldene	PF
Propionic acid derivatives								
IBUPROFEN								
3192B NP,MW	Tablet 400 mg	30	..	..	9.19	10.26	Brufen	AB
IBUPROFEN								
<u>Restricted benefit</u>								
Chronic arthropathies (including osteoarthritis) with an inflammatory component;								
Bone pain due to malignant disease.								
3190X NP	Tablet 400 mg	90	3	..	*14.73	15.80	Brufen	AB
KETOPROFEN								
1588N NP	Suppository 100 mg	40	3	..	*25.30	26.37	Orudis	SW
KETOPROFEN								
<u>Restricted benefit</u>								
Chronic arthropathies (including osteoarthritis) with an inflammatory component.								
1590Q NP	Capsule 200 mg (sustained release)	28	3	..	19.37	20.44	^a Oruvail SR	AV
				^B 2.21	21.58	20.44	^a Orudis SR 200	SW
NAPROXEN								
<u>Restricted benefit</u>								
Chronic arthropathies (including osteoarthritis) with an inflammatory component;								
Bone pain due to malignant disease.								
1614Y NP	Tablet 750 mg (sustained release)	28	3	..	12.38	13.45	^a Proxen SR 750	MD

Musculo-skeletal system

					Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	\$	\$	Brand Name and Manufacturer	
				^B 1.27	13.65	13.45	^a	Naprosyn SR750 RO
1615B NP	Tablet 1 g (sustained release)	28	3	..	14.37	15.44	^a	Proxen SR 1000 MD
				^B 1.34	15.71	15.44	^a	Naprosyn SR1000 RO
1659H NP	Tablet 500 mg	50	3	..	12.91	13.98	^a	Inza 500 AF
				^B 1.36	14.27	13.98	^a	Naprosyn RO
1674D NP	Tablet 250 mg	100	3	..	*13.72	14.79	^a	Inza 250 AF
				^B 2.34	*16.06	14.79	^a	Naprosyn RO

NAPROXEN

Authority required (STREAMLINED)

2270

Chronic arthropathies (including osteoarthritis) with an inflammatory component in patients unable to take a solid dose form of a non-steroidal anti-inflammatory agent;

2271

Bone pain due to malignant disease in patients unable to take a solid dose form of a non-steroidal anti-inflammatory agent.

1658G NP	Oral suspension 125 mg per 5 mL, 474 mL	±1	3	..	78.17	34.20		Naprosyn	RO

NAPROXEN SODIUM

Restricted benefit

Chronic arthropathies (including osteoarthritis) with an inflammatory component;

Bone pain due to malignant disease.

Note

Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.

1795L NP	Tablet 550 mg	50	3	..	13.10	14.17	^a	Crysanal	MD
				^B 2.29	15.39	14.17	^a	Anaprox 550	RO

TIAPROFENIC ACID

Caution

Cystitis and other urinary disorders have been reported with this drug.

Note

The recommended maximum dose is 600 mg per day.

Restricted benefit

Chronic arthropathies (including osteoarthritis) with an inflammatory component.

2103Q NP	Tablet 300 mg	60	3	..	17.58	18.65		Surgam	SW

Fenamates

MEFENAMIC ACID

Restricted benefit

Dysmenorrhoea;

Menorrhagia.

1824B NP	Capsule 250 mg	50	2	..	18.16	19.23		Ponstan	PD

Coxibs

CELECOXIB

Note

The use of celecoxib for the treatment of the following conditions is not subsidised through the PBS:

- (a) acute pain;
- (b) soft tissue injury;
- (c) arthrosis without an inflammatory component.

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty	Safety Net	
					\$	\$	
<u>Restricted benefit</u>							
	Symptomatic treatment of osteoarthritis;						
	Symptomatic treatment of rheumatoid arthritis.						
8439E NP	Capsule 100 mg	60	3	..	32.31	33.38	Celebrex PF
8440F NP	Capsule 200 mg	30	3	..	32.31	33.38	Celebrex PF

Specific antirheumatic agents

Quinolines

HYDROXYCHLOROQUINE SULFATE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1512N NP	Tablet 200 mg	100	1	..	37.59	34.20	Plaquenil SW

Gold preparations

AURANOFIN

Caution

Regular blood and urine checks are essential.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1095P NP	Tablet 3 mg	60	5	..	63.55	34.20	Ridaura GH

SODIUM AUROTHIOMALATE

Caution

Regular blood and urine checks are essential.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2016D NP	Injection 10 mg	10	..	..	67.03	34.20	Myocrisin SW
2017E NP	Injection 20 mg	10	1	..	102.67	34.20	Myocrisin SW
2018F NP	Injection 50 mg	10	1	..	152.47	34.20	Myocrisin SW

Penicillamine and similar agents

PENICILLAMINE

Caution

Regular blood and urine checks are essential.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2721F NP	Tablet 125 mg	100	1	..	31.63	32.70	D-Penamine AL
2838J NP	Tablet 250 mg	100	1	..	43.51	34.20	D-Penamine AL

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Other specific antirheumatic agents

LEFLUNOMIDE

Caution

Leflunomide is a category X drug and must not be given to pregnant women. Pregnancy should be avoided for two years after cessation of therapy, unless special wash-out procedures are carried out.

Authority required (STREAMLINED)

2643

Initial treatment of severe active rheumatoid arthritis where other disease modifying anti-rheumatic drugs (including methotrexate) are ineffective and/or inappropriate. Treatment must be initiated by a physician;

2681

Initial treatment of severe active psoriatic arthritis where other disease modifying anti-rheumatic drugs (including methotrexate) are ineffective and/or inappropriate. Treatment must be initiated by a physician.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8373Q	Pack containing 3 tablets leflunomide 100 mg and 30 tablets leflunomide 20 mg	±1	..	..	211.31	34.20	Arava	SW

LEFLUNOMIDE

Caution

Leflunomide is a category X drug and must not be given to pregnant women. Pregnancy should be avoided for two years after cessation of therapy, unless special wash-out procedures are carried out.

Authority required (STREAMLINED)

2644

Treatment of severe active rheumatoid arthritis where other disease modifying anti-rheumatic drugs (including methotrexate) are ineffective and/or inappropriate. Treatment must be initiated by a physician;

2682

Treatment of severe active psoriatic arthritis where other disease modifying anti-rheumatic drugs (including methotrexate) are ineffective and/or inappropriate. Treatment must be initiated by a physician.

8374R	Tablet 10 mg	30	5	..	91.91	34.20	^a Arabloc ^a Arava	AV SW
8375T	Tablet 20 mg	30	5	..	136.59	34.20	^a Arabloc ^a Arava	AV SW

Muscle relaxants

Muscle relaxants, centrally acting agents

Other centrally acting agents

BACLOFEN

2729P NP	Tablet 10 mg	100	5	..	32.08	33.15	^a Chem mart Baclofen ^a Clofen 10 ^a GenRx Baclofen ^a Stelax 10 ^a Terry White Chemists Baclofen	CH AF GX SI TW
				^B 2.26	34.34	33.15	^a Lioresal 10	NV
2730Q NP	Tablet 25 mg	100	5	..	60.09	34.20	^a Chem mart Baclofen ^a Clofen 25 ^a GenRx Baclofen ^a Stelax 25 ^a Terry White Chemists Baclofen	CH AF GX SI TW
				^B 2.18	62.27	34.20	^a Lioresal 25	NV

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	

Muscle relaxants, directly acting agents

Dantrolene and derivatives

DANTROLENE SODIUM

Restricted benefit

Treatment of chronic spasticity.

1779P NP	Capsule 25 mg	100	2	..	72.04	34.20	Dantrium	PF
1780Q NP	Capsule 50 mg	100	2	..	81.81	34.20	Dantrium	PF

Antigout preparations

Antigout preparations

Preparations inhibiting uric acid production

ALLOPURINOL

Note

The dose should be adjusted in accordance with renal function.

2600W NP	Tablet 100 mg	200	2	..	13.19	14.26	^a Allopurinol Sandoz	SZ
							^a Allosig	FM
							^a Chem mart	CH
							^a Allopurinol	
							^a GenRx Allopurinol	GX
							^a Terry White	TW
							^a Chemists	
				^B 3.01	16.20	14.26	^a Allopurinol	
					*13.20	14.27	^a Zyloprim	SI
							^a Progout 100	AF
2604C NP	Tablet 300 mg	60	2	..	10.42	11.49	^a Allopurinol Sandoz	SZ
							^a Allosig	FM
							^a Chem mart	CH
							^a Allopurinol	
							^a GenRx Allopurinol	GX
							^a Progout 300	AF
							^a Terry White	TW
							^a Chemists	
				^B 3.00	13.42	11.49	^a Allopurinol	
							^a Zyloprim	SI

Preparations increasing uric acid excretion

PROBENECID

1940D NP	Tablet 500 mg	100	5	..	75.69	34.20	Pro-Cid	PL

Preparations with no effect on uric acid metabolism

COLCHICINE

3410L NP	Tablet 500 micrograms	30	2	..	11.09	12.16	^a Lengout	LN
				^B 0.87	11.96	12.16	^a Colgout	AS

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
Drugs for treatment of bone diseases							

Drugs affecting bone structure and mineralization

Bisphosphonates

ALENDRONATE SODIUM

Authority required (STREAMLINED)

3070

Treatment as the sole PBS-subsidised anti-resorptive agent for corticosteroid-induced osteoporosis in a patient currently on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy with a Bone Mineral Density (BMD) T-score of -1.5 or less.

The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

2645

Treatment as the sole PBS-subsidised anti-resorptive agent for osteoporosis in a patient aged 70 years of age or older with a Bone Mineral Density (BMD) T-score of -3.0 or less.

The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

2646

Treatment as the sole PBS-subsidised anti-resorptive agent for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Note

Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.

8511Y NP	Tablet equivalent to 70 mg alendronic acid	4	5	..	45.86	34.20	^a	Adronat	AF
							^a	Alendrobell 70mg	BF
							^a	Alendronate-GA	GM
							^a	Alendronate Sandoz	SZ
							^a	Alendro Once Weekly	SI
							^a	APO-Alendronate	TX
							^a	Chem mart Alendronate 70mg	CH
							^a	Ossmax 70mg	RA
							^a	Terry White Chemists Alendronate 70mg	TW
				^B 2.00	47.86	34.20	^a	Fosamax Once Weekly	MK

ALENDRONATE SODIUM

Authority required (STREAMLINED)

3256

Symptomatic Paget disease of bone.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8090T NP	Tablet equivalent to 40 mg alendronic acid	30	5	..	133.35	34.20		Fosamax 40 mg	MK

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
DISODIUM ETIDRONATE								
<u>Authority required (STREAMLINED)</u>								
3257								
Symptomatic Paget disease of bone when calcitonin has been found to be unsatisfactory due to lack of efficacy;								
3258								
Symptomatic Paget disease of bone when calcitonin has been found to be unsatisfactory due to unacceptable side effects;								
1153								
Heterotopic ossification.								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
2920Q NP	Tablet 200 mg	60	5	..	115.17	34.20	Didronel	PF
DISODIUM PAMIDRONATE								
<u>Authority required (STREAMLINED)</u>								
3256								
Symptomatic Paget disease of bone.								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
<u>Note</u>								
The concentrated injection 15 mg and powder for I.V. infusion 15 mg (after reconstitution) are bioequivalent.								
8208B NP	Injection set containing 4 vials powder for I.V. infusion 15 mg and 4 ampoules solvent 5 mL	1	..	..	254.71	34.20	^a Aredia 15 mg	NV
8461H NP	Concentrated injection 15 mg in 5 mL	4	..	..	*254.74	34.20	^a Pamisol	HH
DISODIUM PAMIDRONATE								
<u>Authority required (STREAMLINED)</u>								
3256								
Symptomatic Paget disease of bone.								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
<u>Note</u>								
The concentrated injection 30 mg and powder for I.V. infusion 30 mg (after reconstitution) are bioequivalent.								
8209C NP	Injection set containing 2 vials powder for I.V. infusion 30 mg and 2 ampoules solvent 10 mL	1	..	..	254.71	34.20	^a Aredia 30 mg	NV
8462J NP	Concentrated injection 30 mg in 10 mL	2	..	..	*254.72	34.20	^a Pamisol	HH
DISODIUM PAMIDRONATE								
<u>Authority required (STREAMLINED)</u>								
3256								
Symptomatic Paget disease of bone.								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8463K NP	Concentrated injection 60 mg in 10 mL	1	..	..	254.71	34.20	Pamisol	HH

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
IBANDRONIC ACID							
<u>Restricted benefit</u>							
Bone metastases from breast cancer.							
<u>Note</u>							
Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
9357L NP	Tablet 50 mg (as ibandronate sodium monohydrate)	28	2	..	342.34	34.20	Bondronat HH
RISEDRONATE SODIUM							
<u>Authority required (STREAMLINED)</u>							
3070							
Treatment as the sole PBS-subsidised anti-resorptive agent for corticosteroid-induced osteoporosis in a patient currently on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy with a Bone Mineral Density (BMD) T-score of -1.5 or less.							
The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.							
<u>Authority required (STREAMLINED)</u>							
2645							
Treatment as the sole PBS-subsidised anti-resorptive agent for osteoporosis in a patient aged 70 years of age or older with a Bone Mineral Density (BMD) T-score of -3.0 or less.							
The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.							
<u>Authority required (STREAMLINED)</u>							
2646							
Treatment as the sole PBS-subsidised anti-resorptive agent for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.							
A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.							
<u>Note</u>							
Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.							
8481J NP	Tablet 5 mg	28	5	..	53.34	34.20	Actonel SW
8621R NP	Tablet 35 mg	4	5	..	53.34	34.20	Actonel Once-a-Week SW
9391G NP	Tablet 150 mg	1	5	..	56.98	34.20	Actonel Once-a-Month SW
RISEDRONATE SODIUM							
<u>Authority required (STREAMLINED)</u>							
3256							
Symptomatic Paget disease of bone.							
<u>Note</u>							
Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
8482K NP	Tablet 30 mg	28	1	..	304.62	34.20	Actonel SW
SODIUM CLODRONATE TETRAHYDRATE							
<u>Restricted benefit</u>							
Maintenance treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy;							
Multiple myeloma;							
Bone metastases from breast cancer.							

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Note							
Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
8132B NP	Capsule equivalent to 400 mg sodium clodronate	100	2	..	334.08	34.20	Bonefos SC
8265B NP	Tablet equivalent to 800 mg sodium clodronate	60	2	..	391.44	34.20	Bonefos 800 mg SC

TILUDRONATE DISODIUM

Authority required (STREAMLINED)

3256

Symptomatic Paget disease of bone.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8267D NP	Tablet equivalent to 200 mg tiludronic acid	56	2	..	304.62	34.20	Skelid SW

ZOLEDRONIC ACID

Authority required

Treatment as the sole PBS-subsidised anti-resorptive agent for corticosteroid-induced osteoporosis in a patient currently on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy with a Bone Mineral Density (BMD) T-score of -1.5 or less.

The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Only 1 treatment each year for 3 years per patient in a lifetime will be PBS-subsidised.

Authority required

Treatment as the sole PBS-subsidised anti-resorptive agent for osteoporosis in a patient aged 70 years of age or older with a Bone Mineral Density (BMD) T-score of -3.0 or less.

The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Only 1 treatment each year for 3 years per patient in a lifetime will be PBS-subsidised.

Authority required

Treatment as the sole PBS-subsidised anti-resorptive agent for established osteoporosis in a patient with fracture due to minimal trauma.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

In all cases, the fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.

Only 1 treatment each year for 3 years per patient in a lifetime will be PBS-subsidised.

Note

Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.

9288W	Solution for I.V. infusion 5 mg (as monohydrate) in 100 mL	1	..	..	589.17	34.20	Aclasta NV

ZOLEDRONIC ACID

Authority required

Symptomatic Paget disease of bone.

Only 1 treatment each year per patient will be PBS-subsidised.

9350D	Solution for I.V. infusion 5 mg (as monohydrate) in 100 mL	1	..	..	589.17	34.20	Aclasta NV

Bisphosphonates, combinations

ALENDRONATE SODIUM with COLECALCIFEROL

Authority required (STREAMLINED)

3070

Treatment as the sole PBS-subsidised anti-resorptive agent for corticosteroid-induced osteoporosis in a patient currently on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy with a Bone Mineral Density (BMD) T-score of -1.5 or

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	
	less. The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.						
	Authority required (STREAMLINED) 2645 Treatment as the sole PBS-subsidised anti-resorptive agent for osteoporosis in a patient aged 70 years of age or older with a Bone Mineral Density (BMD) T-score of -3.0 or less. The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.						
	Authority required (STREAMLINED) 2646 Treatment as the sole PBS-subsidised anti-resorptive agent for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.						
	Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.						
	Note Fosamax Plus provides a supplemental intake of vitamin D. The amount of colecalciferol present in Fosamax Plus is not sufficient to use as the sole treatment for correction of vitamin D deficiency.						
9012H NP	Tablet equivalent to 70 mg alendronic acid with 70 micrograms colecalciferol	4	5	..	45.86	34.20	Fosamax Plus MK

ALENDRONATE SODIUM with COLECALCIFEROL

Authority required (STREAMLINED)

3070

Treatment as the sole PBS-subsidised anti-resorptive agent for corticosteroid-induced osteoporosis in a patient currently on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy with a Bone Mineral Density (BMD) T-score of -1.5 or less.

The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

2645

Treatment as the sole PBS-subsidised anti-resorptive agent for osteoporosis in a patient aged 70 years of age or older with a Bone Mineral Density (BMD) T-score of -3.0 or less.

The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

2646

Treatment as the sole PBS-subsidised anti-resorptive agent for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Note

Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.

9183H NP	Tablet equivalent to 70 mg alendronic acid with 140 micrograms colecalciferol	4	5	..	45.86	34.20	^a Dronalen Plus	GM
							^a Fosamax Plus 70 mg/140 mcg	MK

ALENDRONATE SODIUM with COLECALCIFEROL and CALCIUM CARBONATE

Authority required (STREAMLINED)

3070

Treatment as the sole PBS-subsidised anti-resorptive agent for corticosteroid-induced osteoporosis in a patient currently on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy with a Bone Mineral Density (BMD) T-score of -1.5 or less.

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.							
<u>Authority required (STREAMLINED)</u>							
2645							
Treatment as the sole PBS-subsidised anti-resorptive agent for osteoporosis in a patient aged 70 years of age or older with a Bone Mineral Density (BMD) T-score of -3.0 or less.							
The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.							
<u>Authority required (STREAMLINED)</u>							
2646							
Treatment as the sole PBS-subsidised anti-resorptive agent for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.							
A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.							
<u>Note</u>							
Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.							
9351E NP	Pack containing 4 tablets containing the equivalent of 70 mg alendronic acid with 140 micrograms colecalciferol and 48 tablets calcium carbonate 1.25 g (equivalent to 500 mg calcium)	£1	5	..	45.86	34.20 ^a	Dronalen Plus D-Cal FR
						^a	Fosamax Plus D-Cal MK

DISODIUM ETIDRONATE and CALCIUM CARBONATE

Authority required (STREAMLINED)

2646

Treatment as the sole PBS-subsidised anti-resorptive agent for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Note

Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8056B NP	Pack containing 28 tablets disodium etidronate 200 mg and 76 tablets calcium carbonate 1.25 g (equivalent to 500 mg calcium)	£1	1	..	70.69	34.20	Didrocal PF

RISEDRONATE SODIUM and CALCIUM CARBONATE

Authority required (STREAMLINED)

3070

Treatment as the sole PBS-subsidised anti-resorptive agent for corticosteroid-induced osteoporosis in a patient currently on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy with a Bone Mineral Density (BMD) T-score of -1.5 or less.

The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

2645

Treatment as the sole PBS-subsidised anti-resorptive agent for osteoporosis in a patient aged 70 years of age or older with a Bone Mineral Density (BMD) T-score of -3.0 or less.

The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

2646

Treatment as the sole PBS-subsidised anti-resorptive agent for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.						
	Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.						
8899J NP	Pack containing 4 tablets risedronate sodium 35 mg and 24 tablets calcium carbonate 1.25 g (equivalent to 500 mg calcium)	₹1	5	..	53.34	34.20	Actonel Combi SW

RISEDRONATE SODIUM and CALCIUM CARBONATE with COLECALCIFEROL

Authority required (STREAMLINED)

3070

Treatment as the sole PBS-subsidised anti-resorptive agent for corticosteroid-induced osteoporosis in a patient currently on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy with a Bone Mineral Density (BMD) T-score of -1.5 or less.

The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

2645

Treatment as the sole PBS-subsidised anti-resorptive agent for osteoporosis in a patient aged 70 years of age or older with a Bone Mineral Density (BMD) T-score of -3.0 or less.

The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

2646

Treatment as the sole PBS-subsidised anti-resorptive agent for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Note

Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.

9147K NP	Pack containing 4 tablets risedronate sodium 35 mg and 24 sachets containing granules of calcium carbonate 2.5 g (equivalent to 1 g calcium) with colecalciferol 22 micrograms	₹1	5	..	53.34	34.20	Actonel Combi D SW

Other drugs affecting bone structure and mineralization

CALCITRIOL

Authority required (STREAMLINED)

1165

Hypocalcaemia due to renal disease;

1166

Hypoparathyroidism;

1167

Hypophosphataemic rickets;

1467

Vitamin D-resistant rickets;

2636

Treatment for established osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

2502Q NP	Capsule 0.25 microgram	100	3	..	43.25	34.20	^a	Calcitriol-DP	GN
							^a	Calcitriol-GA	GM
							^a	Calcitriol Sandoz	SZ
							^a	GenRx Calcitriol	GX

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
							^a Kosteo	SI
							^a Rocaltrol	RO
							^a Sical	AF

DENOSUMAB

Authority required

Treatment as the sole PBS-subsidised anti-resorptive agent for osteoporosis in a woman aged 70 years of age or older with a Bone Mineral Density (BMD) T-score of -3.0 or less.

The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated;

Treatment as the sole PBS-subsidised anti-resorptive agent for established post-menopausal osteoporosis in a woman with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Note

Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.

5457F	Injection 60 mg in 1 mL pre-filled syringe	1	..	..	304.87	34.20	Prolia	AN

RALOXIFENE HYDROCHLORIDE

Authority required (STREAMLINED)

2647

Treatment as the sole PBS-subsidised anti-resorptive agent for established post-menopausal osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Note

Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.

8363E NP	Tablet 60 mg	28	5	..	57.87	34.20	Evista	LY

STRONTIUM RANELATE

Authority required (STREAMLINED)

2758

Treatment as the sole PBS-subsidised anti-resorptive agent for osteoporosis in a woman aged 70 years or older with a bone mineral density (BMD) T-score of -3.0 or less.

The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

2647

Treatment as the sole PBS-subsidised anti-resorptive agent for established post-menopausal osteoporosis in patients with fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or CT-scan or MRI scan must be documented in the patient's medical records when treatment is initiated.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Note

Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, disodium etidronate, raloxifene hydrochloride, strontium ranelate and zoledronic acid.

3036T NP	Sachet containing granules for oral suspension 2 g	28	5	..	53.34	34.20	Protos 2 g	SE

TERIPARATIDE

Note

Any queries concerning the arrangements to prescribe teriparatide may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Written applications for authority to prescribe teriparatide should be forwarded to: Medicare Australia Prior Written Approval of Specialised Drugs Reply Paid 9826 GPO Box 9826 HOBART TAS 7001 Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au. Authority required Initial treatment, as the sole PBS-subsidised agent, by a specialist or consultant physician, for severe, established osteoporosis in a patient with a very high risk of fracture who: (a) has a bone mineral density (BMD) T-score of -3.0 or less; and (b) has had 2 or more fractures due to minimal trauma; and (c) has experienced at least 1 symptomatic new fracture after at least 12 months continuous therapy with an anti-resorptive agent at adequate doses. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body. If treatment with anti-resorptive therapy is contraindicated according to the relevant TGA-approved Product Information, details of the contraindication must be provided at the time of application. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the relevant period of use of one anti-resorptive agent, alternate anti-resorptive agents must be trialled so that the patient achieves the minimum requirement of 12 months continuous therapy. Details of accepted toxicities including severity can be found on the Medicare Australia website at www.medicareaustralia.gov.au and must be provided at the time of application. Anti-resorptive therapies for osteoporosis and their adequate doses which will be accepted for the purposes of administering this restriction are alendronate sodium 10 mg per day or 70 mg once weekly, risedronate sodium 5 mg per day or 35 mg once weekly or 150 mg once monthly, raloxifene hydrochloride 60 mg per day (women only), denosumab 60 mg once every 6 months, disodium etidronate 200 mg with calcium carbonate 1.25 g per day, strontium ranelate 2 g per day and zoledronic acid 5 mg per annum. Authority applications must be made in writing and must include: Details of prior anti-resorptive therapy, fracture history including the date(s), site(s), the symptoms associated with the fracture(s) which developed during the course of anti-resorptive therapy and the score of the qualifying BMD measurement. Note No applications for increased maximum quantities and/or repeats will be authorised. Authority required Initial treatment, as the sole PBS-subsidised agent, by a specialist or consultant physician, for severe, established osteoporosis in a patient with a very high risk of fracture who was receiving treatment with teriparatide prior to 1 May 2009. The authority application must be made in writing and the commencement date of treatment and the number of doses the patient has received of teriparatide must be provided with the application. The patient is eligible to receive a maximum of 18 months therapy of combined PBS-subsidised and non-PBS-subsidised therapy. Patients may qualify for PBS-subsidised treatment under this restriction once only. Note No applications for increased maximum quantities and/or repeats will be authorised. Authority required Continuing treatment for severe established osteoporosis where the patient has previously been issued with an authority prescription for this drug. Teriparatide must only be used for a lifetime maximum of 18 months therapy (18 pens). Up to a maximum of 18 pens will be reimbursed through the PBS. Authority applications for continuing treatment may be made by telephone to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Note No applications for increased maximum quantities and/or repeats will be authorised. Note Special Pricing Arrangements apply.							
9411H	Injection 250 micrograms per mL, 2.4 mL in multi-dose pre-filled pen	1	5	..	438.37	34.20	Forteo LY

Musculo-skeletal system

Name, Restriction, Manner of Administration and Form		Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Nervous system

Analgesics

Opioids

Natural opium alkaloids

CODEINE PHOSPHATE

1214X	Tablet 30 mg	20	..	..	16.87	17.94	Fawns and McAllan Proprietary Limited	FM

CODEINE PHOSPHATE with PARACETAMOL

Note

Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of codeine phosphate with paracetamol below.

1215Y NP	Tablet 30 mg-500 mg	20	..	..	8.08	9.15	^a APO- Paracetamol/Co deine 500/30	TX
							^a Codalgin Forte	FM
							^a Codapane Forte	AL
							^a Comfarol Forte	SZ
							^a Dolaforte	CO
							^a Prodeine Forte	AV
				^B 2.80	10.88	9.15	^a Panadeine Forte	SW

CODEINE PHOSPHATE with PARACETAMOL

Authority required

Severe disabling pain not responding to non-narcotic analgesics.

Note

Each authority approval will be limited to no more than 240 tablets per month for no more than 6 months.

8785J NP	Tablet 30 mg-500 mg	60	..	..	*11.40	12.47	^a APO- Paracetamol/Co deine 500/30	TX
							^a Codalgin Forte	FM
							^a Codapane Forte	AL
							^a Comfarol Forte	SZ
							^a Dolaforte	CO
							^a Prodeine Forte	AV
				^B 8.40	*19.80	12.47	^a Panadeine Forte	SW

HYDROMORPHONE HYDROCHLORIDE

Caution

The risk of drug dependence is high.

8420E NP	Injection 2 mg in 1 mL	5	..	..	22.84	23.91	Dilaudid	MF
8421F NP	Injection 10 mg in 1 mL	5	..	..	28.97	30.04	Dilaudid-HP	MF
8422G NP	Injection 50 mg in 5 mL	5	..	..	52.00	34.20	Dilaudid-HP	MF
8423H NP	Injection 500 mg in 50 mL	1	..	..	75.41	34.20	Dilaudid-HP	MF

HYDROMORPHONE HYDROCHLORIDE

The risk of drug dependence is high.

Severe disabling pain not responding to non-narcotic analgesics.

NOTE Authorities for increased maximum quantities and/or repeats will be granted only for:

- (i) severe disabling pain associated with proven malignant neoplasia; or
- (ii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or
- (iii) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or
- (iv) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.

MORPHINE HYDROCHLORIDE

Caution
The risk of drug dependence is high.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Restricted benefit							
Severe disabling pain not responding to non-narcotic analgesics.							
Note							
Authorities for increased maximum quantities and/or repeats will be granted only for:							
(i) severe disabling pain associated with proven malignant neoplasia; or							
(ii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or							
(iii) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or							
(iv) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.							
2122Q NP	Oral solution 2 mg per mL, 200 mL	1	..	..	18.16	19.23	Ordine 2 MF
2123R NP	Oral solution 5 mg per mL, 200 mL	1	..	..	20.78	21.85	Ordine 5 MF
2124T NP	Oral solution 10 mg per mL, 200 mL	1	..	..	24.92	25.99	Ordine 10 MF
MORPHINE SULFATE							
Caution							
The risk of drug dependence is high.							
1644M NP,MW	Injection 10 mg in 1 mL	5	..	..	14.10	15.17	Hospira Pty Limited HH
1645N NP,MW	Injection 15 mg in 1 mL	5	..	..	14.45	15.52	Hospira Pty Limited HH
1647Q NP	Injection 30 mg in 1 mL	5	..	..	15.90	16.97	Hospira Pty Limited HH
MORPHINE SULFATE							
Caution							
The risk of drug dependence is high.							
Restricted benefit							
Severe disabling pain due to cancer not responding to non-narcotic analgesics.							
8669G NP	Tablet 10 mg	20	..	..	14.41	15.48	Sevredol MF
8670H NP	Tablet 20 mg	20	..	..	15.39	16.46	Sevredol MF
MORPHINE SULFATE							
Caution							
The risk of drug dependence is high.							
Restricted benefit							
Severe disabling pain not responding to non-narcotic analgesics.							
Note							
Authorities for increased maximum quantities and/or repeats will be granted only for:							
(i) severe disabling pain associated with proven malignant neoplasia; or							
(ii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or							
(iii) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or							

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
	(iv) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.						
1646P NP	Tablet 30 mg	20	..	..	14.13	15.20	Anamorph FM
MORPHINE SULFATE							
Caution							
The risk of drug dependence is high.							
Restricted benefit							
Chronic severe disabling pain not responding to non-narcotic analgesics.							
Note							
Authorities for increased maximum quantities and/or repeats will be granted only for:							
(i) chronic severe disabling pain associated with proven malignant neoplasia; or							
(ii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or							
(iii) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or							
(iv) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.							
1653B NP	Tablet 10 mg (controlled release)	20	..	..	17.08	18.15	^a Momex SR 10 SI
						^a	MS Contin MF
1654C NP	Tablet 30 mg (controlled release)	20	..	..	28.62	29.69	^a Momex SR 30 SI
						^a	MS Contin MF
1655D NP	Tablet 60 mg (controlled release)	20	..	..	43.37	34.20	^a Momex SR 60 SI
						^a	MS Contin MF
1656E NP	Tablet 100 mg (controlled release)	20	..	..	55.63	34.20	^a Momex SR 100 SI
						^a	MS Contin MF
2839K NP	Capsule 20 mg (containing sustained release pellets)	20	..	..	20.70	21.77	Kapanol GK
2840L NP	Capsule 50 mg (containing sustained release pellets)	20	..	..	34.04	34.20	Kapanol GK
2841M NP	Capsule 100 mg (containing sustained release pellets)	20	..	..	54.27	34.20	Kapanol GK
8035X NP	Tablet 5 mg (controlled release)	20	..	..	15.29	16.36	MS Contin MF
8146R NP	Sachet containing controlled release granules for oral suspension, 30 mg per sachet	20	..	..	48.71	34.20	MS Contin Suspension 30 mg MF
8305D NP	Sachet containing controlled release granules for oral suspension, 60 mg per sachet	20	..	..	53.87	34.20	MS Contin Suspension 60 mg MF
8306E NP	Sachet containing controlled release granules for oral suspension, 100 mg per sachet	20	..	..	65.42	34.20	MS Contin Suspension 100 mg MF
8349K NP	Capsule 10 mg (containing sustained release pellets)	20	..	..	17.08	18.15	Kapanol GK
8489T NP	Tablet 15 mg (controlled release)	20	..	..	20.12	21.19	MS Contin MF
8490W NP	Sachet containing controlled release granules for oral suspension, 20 mg per sachet	20	..	..	47.53	34.20	MS Contin Suspension 20 mg MF
8491X NP	Capsule 30 mg (controlled release)	10	..	..	20.12	21.19	MS Mono MF

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8492Y NP	Capsule 60 mg (controlled release)	10	..	..	28.62	29.69	MS Mono	MF
8493B NP	Capsule 90 mg (controlled release)	10	..	..	32.67	33.74	MS Mono	MF
8494C NP	Capsule 120 mg (controlled release)	10	..	..	43.37	34.20	MS Mono	MF

MORPHINE SULFATE

Caution

The risk of drug dependence is high.

Authority required

Chronic severe disabling pain due to cancer.

8453X NP	Tablet 200 mg (controlled release)	20	..	..	91.28	34.20	MS Contin	MF
8454Y NP	Sachet containing controlled release granules for oral suspension, 200 mg per sachet	20	..	..	121.81	34.20	MS Contin Suspension 200 mg	MF

MORPHINE TARTRATE

Caution

The risk of drug dependence is high.

1607N NP	Injection 120 mg in 1.5 mL	5	..	..	31.11	32.18	Hospira Pty Limited	HH

OXYCODONE

Caution

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

Note

Authorities for increased maximum quantities and/or repeats will be granted only for:

- (i) severe disabling pain associated with proven malignant neoplasia; or
- (ii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or
- (iii) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or
- (iv) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.

2481N NP	Suppository 30 mg	12	..	..	43.66	34.20	Proladone	PL

OXYCODONE HYDROCHLORIDE

Caution

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

Note

Authorities for increased maximum quantities and/or repeats will be granted only for:

- (i) severe disabling pain associated with proven malignant neoplasia; or
- (ii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or
- (iii) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or							
(iv) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.							
2622B NP	Tablet 5 mg	20	..	..	12.30	13.37	Endone SI
8464L NP	Capsule 5 mg	20	..	..	12.30	13.37	OxyNorm MF
8501K NP	Capsule 10 mg	20	..	..	15.42	16.49	OxyNorm MF
8502L NP	Capsule 20 mg	20	..	..	20.15	21.22	OxyNorm MF
8644Y NP	Oral solution 5 mg per 5 mL, 250 mL	1	..	..	20.72	21.79	OxyNorm Liquid 5mg/5mL MF

OXYCODONE HYDROCHLORIDE

Caution

The risk of drug dependence is high.

Restricted benefit

Chronic severe disabling pain not responding to non-narcotic analgesics.

Note

Authorities for increased maximum quantities and/or repeats will be granted only for:

(i) chronic severe disabling pain associated with proven malignant neoplasia; or

(ii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or

(iii) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or

(iv) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.

8385H NP	Tablet 10 mg (controlled release)	20	..	..	21.95	23.02	OxyContin MF
8386J NP	Tablet 20 mg (controlled release)	20	..	..	31.87	32.94	OxyContin MF
8387K NP	Tablet 40 mg (controlled release)	20	..	..	48.35	34.20	OxyContin MF
8388L NP	Tablet 80 mg (controlled release)	20	..	..	71.70	34.20	OxyContin MF
8681X NP	Tablet 5 mg (controlled release)	20	..	..	21.18	22.25	OxyContin MF
9399Q NP	Tablet 15 mg (controlled release)	20	..	..	27.93	29.00	OxyContin MF
9400R NP	Tablet 30 mg (controlled release)	20	..	..	41.22	34.20	OxyContin MF

Phenylpiperidine derivatives

FENTANYL

Caution

The risk of drug dependence is high.

Restricted benefit

Chronic severe disabling pain not responding to non-narcotic analgesics.

Note

Authorities for increased maximum quantities and/or repeats will be granted only for:

(i) chronic severe disabling pain associated with proven malignant neoplasia; or

(ii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
(iii) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or (iv) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.							
Note							
Durogesic is not recommended in opioid naive patients with non-cancer pain, because of a high incidence of adverse events in these patients. Patients with cancer pain may be initiated on the lowest strength patch (12 micrograms per hour).							
8878G NP	Transdermal patch 2.1 mg (releasing approximately 12 micrograms per hour)	5	..	..	47.16	34.20	Durogesic 12 JC
8891Y NP	Transdermal patch 4.2 mg (releasing approximately 25 micrograms per hour)	5	..	..	56.28	34.20	Durogesic 25 JC
8892B NP	Transdermal patch 8.4 mg (releasing approximately 50 micrograms per hour)	5	..	..	95.38	34.20	Durogesic 50 JC
8893C NP	Transdermal patch 12.6 mg (releasing approximately 75 micrograms per hour)	5	..	..	127.28	34.20	Durogesic 75 JC
8894D NP	Transdermal patch 16.8 mg (releasing approximately 100 micrograms per hour)	5	..	..	155.76	34.20	Durogesic 100 JC

Diphenylpropylamine derivatives

METHADONE HYDROCHLORIDE

Caution

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

Note

Authorities for increased maximum quantities and/or repeats will be granted only for:

- (i) severe disabling pain associated with proven malignant neoplasia; or
- (ii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or
- (iii) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or
- (iv) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1606M NP	Injection 10 mg in 1 mL	5	..	..	49.31	34.20	Physeptone	SI
1609Q NP	Tablet 10 mg	20	..	..	15.35	16.42	Physeptone	SI

Oripavine derivatives

BUPRENORPHINE

Restricted benefit

Chronic severe disabling pain not responding to non-narcotic analgesics.

Note

Authorities for increased maximum quantities and/or repeats will be granted only for:

- (i) chronic severe disabling pain associated with proven malignant neoplasia; or
- (ii) chronic severe disabling pain not responding to non-narcotic analgesics where the total duration of narcotic analgesic treatment is less than 12 months; or

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	
	(iii) first application for treatment beyond 12 months of chronic severe disabling pain not responding to non-narcotic analgesics where the patient's pain management has been reviewed through consultation by the patient with another medical practitioner, and the clinical need for continuing narcotic analgesic treatment has been confirmed. The date of the consultation must be no more than 3 months prior to the application for a PBS authority. The full name of the medical practitioner consulted and the date of consultation are to be provided at the time of application; or						

(iv) subsequent application for treatment of chronic severe disabling pain not responding to non-narcotic analgesics where a PBS authority prescription for treatment beyond 12 months has previously been issued for this patient.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Caution

The risk of drug dependence is high.

8865N NP	Transdermal patch 5 mg (releasing approximately 5 micrograms per hour)	2	..	..	26.70	27.77	Norspan	MF
8866P NP	Transdermal patch 10 mg (releasing approximately 10 micrograms per hour)	2	..	..	40.77	34.20	Norspan	MF
8867Q NP	Transdermal patch 20 mg (releasing approximately 20 micrograms per hour)	2	..	..	56.08	34.20	Norspan	MF

Other opioids

TRAMADOL HYDROCHLORIDE

Restricted benefit

For acute pain where aspirin and/or paracetamol alone are inappropriate or have failed.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8455B NP	Capsule 50 mg	20	..	..	9.16	10.23	^a APO-Tramadol	TX
							^a Chem mart Tramadol	CH
							^a GA Tramadol 50mg	GM
							^a GenRx Tramadol	GX
							^a Lodam 50	ZP
							^a Terry White Chemists Tramadol	TW
							^a Tramadol Sandoz	SZ
							^a Tramedo	AF
							^a Zydol	SI
				^B 2.42	11.58	10.23	^a Tramal	CS

TRAMADOL HYDROCHLORIDE

Restricted benefit

For dosage titration in chronic pain where aspirin and/or paracetamol alone are inappropriate or have failed.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8611F NP	Capsule 50 mg	20	2	..	9.16	10.23	^a APO-Tramadol	TX
							^a Chem mart Tramadol	CH
							^a GA Tramadol 50mg	GM
							^a GenRx Tramadol	GX
							^a Lodam 50	ZP
							^a Terry White Chemists Tramadol	TW
							^a Tramadol Sandoz	SZ
							^a Tramedo	AF

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
				^B 2.42	11.58	10.23	^a Zydol	SI
							^a Tramal	CS
TRAMADOL HYDROCHLORIDE								
<u>Restricted benefit</u>								
For pain where aspirin and/or paracetamol alone are inappropriate or have failed.								
<u>Note</u>								
Authorities for increased maximum quantities and/or repeats will be granted only for severe disabling pain not responding to non-narcotic analgesics.								
2527B NP	Tablet 50 mg (twice daily sustained release)	20	..	..	11.63	12.70	Tramal SR 50	CS
8523N NP	Tablet 100 mg (twice daily sustained release)	20	..	..	13.86	14.93	^a APO-Tramadol SR	TX
							^a Chem mart Tramadol SR	CH
							^a GA Tramadol SR 100mg	GM
							^a Lodam SR 100	ZP
							^a Terry White Chemists Tramadol SR	TW
							^a TramaHexal SR	SZ
							^a Tramedo SR 100	AF
							^a Zydol SR 100	SI
				^B 4.51	18.37	14.93	^a Tramal SR 100	CS
8524P NP	Tablet 150 mg (twice daily sustained release)	20	..	..	16.45	17.52	^a APO-Tramadol SR	TX
							^a Chem mart Tramadol SR	CH
							^a GA Tramadol SR 150mg	GM
							^a Lodam SR 150	ZP
							^a Terry White Chemists Tramadol SR	TW
							^a Tramadol Sandoz SR	SZ
							^a Tramedo SR 150	AF
							^a Zydol SR 150	SI
				^B 5.37	21.82	17.52	^a Tramal SR 150	CS
8525Q NP	Tablet 200 mg (twice daily sustained release)	20	..	..	18.63	19.70	^a APO-Tramadol SR	TX
							^a Chem mart Tramadol SR	CH
							^a GA Tramadol SR 200mg	GM
							^a Lodam SR 200	ZP
							^a Terry White Chemists Tramadol SR	TW
							^a TramaHexal SR	SZ
							^a Tramedo SR 200	AF
							^a Zydol SR 200	SI
				^B 6.09	24.72	19.70	^a Tramal SR 200	CS
8843K NP	Oral drops 100 mg per mL, 10 mL	≠1	..	..	13.71	14.78	Tramal	CS
9199E NP	Tablet 100 mg (once a day extended release)	10	..	..	13.37	14.44	Durotram XR	IA
9200F NP	Tablet 200 mg (once a day extended release)	10	..	..	16.34	17.41	Durotram XR	IA
9201G	Tablet 300 mg (once a day extended release)	10	..	..	19.79	20.86	Durotram XR	IA

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
NP								
TRAMADOL HYDROCHLORIDE								
<u>Restricted benefit</u>								
Short-term treatment of acute pain.								
<u>Note</u>								
No applications for increased maximum quantities and/or repeats will be authorised.								
8582Q	Injection 100 mg in 2 mL	5	..	..	14.30	15.37	^a Tramahexal	SZ
NP							^a Tramal 100	CS
Other analgesics and antipyretics								
<i>Salicylic acid and derivatives</i>								
ASPIRIN								
1010E	Tablet 300 mg (dispersible)	96	1	..	8.55	9.62	Solprin	RC
NP								
<i>Anilides</i>								
PARACETAMOL								
1746X	Tablet 500 mg	100	1	..	8.42	9.49	^a APO-Paracetamol	TX
NP							^a Chem mart Paracetamol	CH
							^a Febridol	GM
							^a Generic Health Pty Ltd	GQ
							^a Panamax	SW
							^a Paracetamol Sandoz	SZ
							^a Paralgin	FM
							^a Pharmacy Choice Paracetamol	YM
							^a Terry White Chemists Paracetamol	TW
1747Y	Oral liquid 120 mg per 5 mL, 100 mL	‡1	2	..	9.38	10.45	Panamax	SW
NP								
1770E	Oral liquid 240 mg per 5 mL, 200 mL	‡1	2	..	10.68	11.75	Panamax 240 Elixir	SW
NP								
PARACETAMOL								
<u>Restricted benefit</u>								
Chronic arthropathies.								
8784H	Tablet 500 mg	300	4	..	*12.42	13.49	^a APO-Paracetamol	TX
NP							^a Chem mart Paracetamol	CH
							^a Febridol	GM
							^a Generic Health Pty Ltd	GQ
							^a Panamax	SW
							^a Paracetamol Sandoz	SZ
							^a Paralgin	FM
							^a Pharmacy Choice Paracetamol	YM
							^a Terry White Chemists Paracetamol	TW

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
PARACETAMOL								
<u>Restricted benefit</u>								
Relief of persistent pain associated with osteoarthritis.								
8814X NP	Tablet 665 mg (modified release)	192	5	..	*16.64	17.71	Panadol Osteo	GC
Antimigraine preparations								
<i>Ergot alkaloids</i>								
DIHYDROERGOTAMINE MESYLATE								
1323P	Injection 1 mg in 1 mL	5	..	..	17.06	18.13	Dihydergot	NV
METHYSERGIDE								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
2826R NP	Tablet 1 mg	100	2	..	*44.96	34.20	Deseril	LM
Selective 5HT₁ -receptor agonists								
ELETRIPTAN								
<u>Caution</u>								
Eletriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.								
<u>Authority required (STREAMLINED)</u>								
3233								
Migraine attack in a patient where attacks in the past have usually failed to respond to analgesics.								
<u>Note</u>								
No applications for increased maximum quantities and/or repeats will be authorised.								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
5290K NP	Tablet 40 mg (as hydrobromide)	4	5	..	24.75	25.82	Relpax	PF
5291L NP	Tablet 80 mg (as hydrobromide)	4	5	..	24.75	25.82	Relpax	PF
RIZATRIPTAN								
<u>Caution</u>								
Rizatriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.								
<u>Authority required (STREAMLINED)</u>								
3233								
Migraine attack in a patient where attacks in the past have usually failed to respond to analgesics.								
<u>Note</u>								
No applications for increased maximum quantities and/or repeats will be authorised.								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
9313E NP	Wafer 10 mg (as benzoate)	4	5	..	*25.12	26.19	Maxalt	MK

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
SUMATRIPTAN								
<u>Caution</u>								
Sumatriptan is contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.								
<u>Authority required (STREAMLINED)</u>								
3233								
Migraine attack in a patient where attacks in the past have usually failed to respond to analgesics.								
<u>Note</u>								
No applications for increased maximum quantities and/or repeats will be authorised.								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8144P NP	Tablet 50 mg (as succinate)	4	5	..	24.75	25.82	^a Pharmacor Sumatriptan 50	CR
					*24.76	25.83	^a APO-Sumatriptan	TX
							^a Chem mart Sumatriptan	CH
							^a Imigran	GK
							^a Sumagran 50	SI
							^a Sumatab	AF
							^a Terry White Chemists Sumatriptan	TW
8341B NP	Nasal spray 20 mg in 0.1 mL single dose unit	2	5	..	19.52	20.59	Imigran	GK
8885P NP	Tablet (fast disintegrating) 50 mg (as succinate)	4	5	..	*24.76	25.83	Imigran FDT	GK

Other antimigraine preparations

CYPROHEPTADINE HYDROCHLORIDE

Restricted benefit

Prevention of migraine.

Note

Cyproheptadine hydrochloride is not PBS-subsidised for use in hay fever or atopy.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1798P NP	Tablet 4 mg	100	2	..	14.19	15.26	Periactin	AS

PIZOTIFEN MALATE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

3074T NP	Tablet 500 micrograms (base)	100	2	..	21.75	22.82	Sandomigran 0.5	NV

Antiepileptics

Antiepileptics

Barbiturates and derivatives

PHENOBARBITONE

Restricted benefit

Epilepsy.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Note**Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1850J NP	Tablet 30 mg	200	4	..	16.60	17.67	Sigma Pharmaceuticals (Australia) Pty Ltd	SI

PHENOBARBITONE SODIUM**Restricted benefit**

Epilepsy.

Note**Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1853M NP	Injection 200 mg in 1 mL	5	..	..	39.02	34.20	Fawns and McAllan Proprietary Limited	FM

PRIMIDONE**Note****Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1939C NP	Tablet 250 mg	200	2	..	83.49	34.20	Mysoline	LM

Hydantoin derivatives

PHENYTOIN**Note****Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1249R NP	Tablet 50 mg	200	2	..	38.16	34.20	Dilantin Infatabs	PF
2692Q NP	Paediatric oral suspension 30 mg per 5 mL, 500 mL	£1	3	..	26.40	27.47	Dilantin	PF

PHENYTOIN SODIUM**Note****Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1873N NP	Capsule 30 mg	200	2	..	29.18	30.25	Dilantin Sodium	PF
1874P NP	Capsule 100 mg	200	2	..	30.12	31.19	Dilantin Sodium	PF

Succinimide derivatives

ETHOSUXIMIDE**Note****Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1413J NP	Capsule 250 mg	200	2	..	54.10	34.20	Zarontin	PF
1414K NP	Paediatric syrup 250 mg per 5 mL, 200 mL	£1	5	..	25.29	26.36	Zarontin	PF

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Benzodiazepine derivatives							
CLONAZEPAM							
<u>Authority required</u>							
Neurologically proven epilepsy.							
<u>Caution</u>							
Abuse of clonazepam has been reported. Refer to the current product information.							
<u>Note</u>							
Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
1805B NP	Tablet 500 micrograms	200	2	..	*20.20	21.27 ^a	Paxam 0.5 AF
				^B 3.58	*23.78	21.27 ^a	Rivotril RO
1806C NP	Tablet 2 mg	200	2	..	*32.34	33.41 ^a	Paxam 2 AF
				^B 4.08	*36.42	33.41 ^a	Rivotril RO
1808E NP	Oral liquid 2.5 mg per mL, 10 mL	2	..	..	*15.04	16.11	Rivotril RO

CLONAZEPAM

Restricted benefit

Epilepsy.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1807D NP	Injection 1 mg in 2 mL (set containing solution 1 mg in 1 mL and 1 mL diluent)	5	..	..	18.58	19.65	Rivotril RO

NITRAZEPAM

Authority required

Myoclonic epilepsy;

Malignant neoplasia (late stage);

For use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2732T NP	Tablet 5 mg	50	5	..	*9.36	10.43 ^a	Alodorm AF
				^B 3.06	*12.42	10.43 ^a	Mogadon VT

Carboxamide derivatives

CARBAMAZEPINE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2419H NP	Tablet 200 mg	200	2	..	30.20	31.27 ^a	Carbamazepine Sandoz SZ
						^a	Teril AF
				^B 2.74	*32.94	31.27 ^a	Tegretol 200 NV
2422L NP	Tablet 100 mg	200	2	..	19.14	20.21 ^a	Carbamazepine Sandoz SZ

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
2426Q	Tablet 200 mg (controlled release)	200	2	^B 2.56	*21.70	20.21	^a Tegretol 100	NV
NP				..	30.69	31.76	Tegretol CR 200	NV
2427R	Oral suspension 100 mg per 5 mL, 300 mL	±1	5	..	21.35	22.42	Tegretol Liquid	NV
NP								
2431Y	Tablet 400 mg (controlled release)	200	2	..	51.02	34.20	Tegretol CR 400	NV
NP								

OXCARBAZEPINE

Authority required (STREAMLINED)

1587

Treatment of partial epileptic seizures and primary generalised tonic-clonic seizures, which are not controlled satisfactorily by other anti-epileptic drugs.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8584T	Tablet 150 mg	100	5	..	72.27	34.20	Trileptal	NV
NP								
8585W	Tablet 300 mg	100	5	..	115.08	34.20	Trileptal	NV
NP								
8586X	Tablet 600 mg	100	5	..	187.98	34.20	Trileptal	NV
NP								
8588B	Oral suspension 60 mg per mL, 250 mL	2	5	..	*138.12	34.20	Trileptal	NV
NP								

Fatty acid derivatives

SODIUM VALPROATE

Caution

There are reports of fatal hepatotoxicity, particularly in children.

There is increasing evidence of dose-related teratogenesis from this drug.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2289L	Tablet 200 mg (enteric coated)	200	2	..	*32.78	33.85	^a Sodium Valproate	SZ
NP							Sandoz	
							^a Valprease 200	SI
							^a Valpro 200	AF
							^a Valproate	WA
							Winthrop EC 200	
				^B 1.42	*34.20	33.85	^a Epilim EC	SW
2290M	Tablet 500 mg (enteric coated)	200	2	..	*56.34	34.20	^a Sodium Valproate	SZ
NP							Sandoz	
							^a Valprease 500	SI
							^a Valpro 500	AF
							^a Valproate	WA
							Winthrop EC 500	
				^B 1.40	*57.74	34.20	^a Epilim EC	SW
2293Q	Oral liquid 200 mg per 5 mL, 300 mL	2	2	..	*34.92	34.20	Epilim Liquid	SW
NP								
2294R	Crushable tablet 100 mg	200	2	..	*32.00	33.07	Epilim	SW
NP								
2295T	Syrup 200 mg per 5 mL, 300 mL	2	2	..	*34.92	34.20	Epilim Syrup	SW
NP								

TIAGABINE HYDROCHLORIDE

Authority required (STREAMLINED)

2664

Treatment of partial epileptic seizures which are not controlled satisfactorily by other anti-epileptic drugs.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty	Safety Net	
<u>Note</u>							
Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
8221Q NP	Tablet 5 mg (base)	100	5	..	*72.64	34.20	Gabitril OA
8222R NP	Tablet 10 mg (base)	100	5	..	*138.84	34.20	Gabitril OA
8223T NP	Tablet 15 mg (base)	100	5	..	*196.88	34.20	Gabitril OA

VIGABATRIN

Caution

Visual field defects have been reported with this drug.

Authority required (STREAMLINED)

1426

Treatment of epileptic seizures which are not controlled satisfactorily by other anti-epileptic drugs.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2667J NP	Tablet 500 mg	100	5	..	100.93	34.20	Sabril SW
2668K NP	Oral powder, sachet 500 mg	60	5	..	67.70	34.20	Sabril SW

Other antiepileptics

GABAPENTIN

Authority required (STREAMLINED)

2664

Treatment of partial epileptic seizures which are not controlled satisfactorily by other anti-epileptic drugs.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1834M NP	Capsule 300 mg	100	5	..	62.01	34.20	^a DBL Gabapentin	HH
							^a Gabapentin 300	CR
							^a Gabapentin-GA	GM
							^a Gabapentin Sandoz	SZ
							^a Gabatine 300	SI
							^a Gantin	AW
							^a GenRx Gabapentin	GX
							^a Nupentin 300	AF
							^a Neurontin	PF
							^a DBL Gabapentin	HH
1835N NP	Capsule 400 mg	100	5	..	82.24	34.20	^a Douglas Gabapentin 400mg	GN
							^a Gabapentin 400	CR
							^a Gabapentin Sandoz	SZ
							^a Gabatine 400	SI
							^a Gantin	AW
							^a GenRx Gabapentin	GX
							^a Nupentin 400	AF
							^a Neurontin	PF
							^a Gabaran	RA
8389M NP	Tablet 800 mg	100	5	..	166.86	34.20		

2848X NP	Tablet 25 mg	56	5	..	29.40	30.47	^a	APO-Lamotrigine	TX
							^a	GenRx Lamotrigine	GX
							^a	Lamidus	RA
							^a	Lamogine	AF

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
2849Y NP	Tablet 50 mg	56	5	..	44.24	34.20	^a Lamotrigine-DP GM
							^a Lamotrigine-GA GN
							^a Lamotrigine generichealth GQ
							^a Lamotrigine Sandoz SZ
							^a Lamotruster 25 MI
							^a Seaze 25 SI
							^a Lamictal GK
							^a APO-Lamotrigine TX
							^a GenRx Lamotrigine GX
							^a Lamidus RA
							^a Lamogine AF
							^a Lamotrigine-DP GM
							^a Lamotrigine-GA GN
							^a Lamotrigine generichealth GQ
							^a Lamotrigine Sandoz SZ
2850B NP	Tablet 100 mg	56	5	..	67.49	34.20	^a Lamotruster 50 MI
							^a Seaze 50 SI
							^a Lamictal GK
							^a APO-Lamotrigine TX
							^a GenRx Lamotrigine GX
							^a Lamidus RA
							^a Lamogine AF
							^a Lamotrigine-DP GM
							^a Lamotrigine-GA GN
							^a Lamotrigine generichealth GQ
							^a Lamotrigine Sandoz SZ
							^a Lamotruster 100 MI
							^a Seaze 100 SI
							^a Lamictal GK
							^a APO-Lamotrigine TX
2851C NP	Tablet 200 mg	56	5	..	109.01	34.20	^a GenRx Lamotrigine GX
							^a Lamidus RA
							^a Lamogine AF
							^a Lamotrigine-DP GM
							^a Lamotrigine-GA GN
							^a Lamotrigine generichealth GQ
							^a Lamotrigine Sandoz SZ
							^a Lamotruster 200 MI
							^a Seaze 200 SI
							^a Lamictal GK
							^a APO-Lamotrigine TX
							^a GenRx Lamotrigine GX
							^a Lamidus RA
							^a Lamogine AF
							^a Lamotrigine-DP GM
8063J NP	Tablet 5 mg	56	5	..	16.75	17.82	^a Lamotrigine generichealth GQ
							^a Lamotrigine Sandoz SZ
							^a Lamotruster 200 MI
							^a Seaze 200 SI
							^a Lamictal GK
8063J NP	Tablet 5 mg	56	5	..	16.75	17.82	^a Lamogine AF
							^a Seaze 5 SI
							^a Lamictal GK

LEVETIRACETAM

Authority required (STREAMLINED)

2664

Treatment of partial epileptic seizures which are not controlled satisfactorily by other anti-epileptic drugs.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
Note								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8654L NP	Tablet 250 mg	60	5	..	55.46	34.20	^a	APO-Levetiracetam TX
							^a	Chem mart CH
							^a	Levetiracetam
							^a	Kepcet GM
							^a	Keppra UC
							^a	Kevtam AF
							^a	Levecetam 250 RZ
							^a	Levetiracetam GQ
							^a	generichealth
							^a	Levetiracetam SZ
8655M NP	Tablet 500 mg	60	5	..	88.13	34.20	^a	Levitam 250 SI
							^a	Terry White TW
							^a	Chemists
							^a	Levetiracetam
							^a	APO-Levetiracetam TX
							^a	Chem mart CH
							^a	Levetiracetam
							^a	Kepcet GM
							^a	Keppra UC
							^a	Kevtam AF
8656N NP	Tablet 1 g	60	5	..	142.55	34.20	^a	Levecetam 500 RZ
							^a	Levetiracetam GQ
							^a	generichealth
							^a	Levetiracetam SZ
							^a	Levitam 500 SI
							^a	Terry White TW
							^a	Chemists
							^a	Levetiracetam
							^a	APO-Levetiracetam TX
							^a	Chem mart CH
							^a	Levetiracetam
							^a	Kepcet GM
							^a	Keppra UC
							^a	Kevtam AF
							^a	Levecetam 1000 RZ
							^a	Levetiracetam GQ
							^a	generichealth
							^a	Levetiracetam SZ
							^a	Levitam 1000 SI
							^a	Terry White TW
							^a	Chemists
							^a	Levetiracetam
							^a	Levetiracetam

LEVETIRACETAM

Authority required (STREAMLINED)

3291

Treatment of partial epileptic seizures, which are not controlled satisfactorily by other anti-epileptic drugs in a patient unable to take a solid dose form of levetiracetam.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
9169N NP	Oral solution 100 mg per mL, 300 mL	1	5	..	111.42	34.20	Keppra	UC

SULTHIAME

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2099L NP	Tablet 50 mg	200	2	..	46.12	34.20	Ospolot	PL
2100M NP	Tablet 200 mg	200	2	..	98.57	34.20	Ospolot	PL

TOPIRAMATE

Authority required (STREAMLINED)

2797

Treatment of partial epileptic seizures, primary generalised tonic-clonic epileptic seizures and seizures of the Lennox-Gastaut syndrome, which are not controlled satisfactorily by other anti-epileptic drugs.

Authority required (STREAMLINED)

2799

Prophylaxis of migraine in a patient who has experienced an average of 3 or more migraines per month over a period of at least 6 months, and who:

- (a) has a contraindication to beta-blockers, as described in the relevant TGA-approved Product Information; OR
- (b) has experienced intolerance of a severity necessitating permanent withdrawal during treatment with a beta-blocker;
- AND
- (c) has a contraindication to pizotifen because the weight gain associated with this drug poses an unacceptable risk; OR
- (d) has experienced intolerance of a severity necessitating permanent withdrawal during treatment with pizotifen.

Details of the contraindication and/or intolerance(s) must be documented in the patient's medical records when treatment is initiated.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8163P NP	Tablet 25 mg	60	5	..	41.08	34.20	^a APO-Topiramate	TX
							^a Epiramax 25	SI
							^a RBX Topiramate	RA
							^a Tamate	AF
							^a Topamax	JC
							^a Topiramate-GA	GM
							^a Topiramate Sandoz	SZ
8164Q NP	Tablet 50 mg	60	5	..	60.88	34.20	^a APO-Topiramate	TX
							^a Epiramax 50	SI
							^a RBX Topiramate	RA
							^a Tamate	AF
							^a Topamax	JC
							^a Topiramate-GA	GM
							^a Topiramate Sandoz	SZ

TOPIRAMATE

Authority required (STREAMLINED)

2797

Treatment of partial epileptic seizures, primary generalised tonic-clonic epileptic seizures and seizures of the Lennox-Gastaut syndrome, which are not controlled satisfactorily by other anti-epileptic drugs.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8165R NP	Tablet 100 mg	60	5	..	92.11	34.20	^a APO-Topiramate	TX
							^a Epiramax 100	SI
							^a RBX Topiramate	RA
							^a Tamate	AF
							^a Topamax	JC
							^a Topiramate-GA	GM
							^a Topiramate Sandoz	SZ
8166T NP	Tablet 200 mg	60	5	..	150.37	34.20	^a APO-Topiramate	TX
							^a Epiramax 200	SI
							^a RBX Topiramate	RA
							^a Tamate	AF
							^a Topamax	JC
							^a Topiramate-GA	GM
							^a Topiramate Sandoz	SZ

TOPIRAMATE

Authority required (STREAMLINED)

2798

Treatment of partial epileptic seizures, primary generalised tonic-clonic epileptic seizures and seizures of the Lennox-Gastaut syndrome, which are not controlled satisfactorily by other anti-epileptic drugs in patients unable to take a solid dose form of topiramate.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8371N NP	Capsule 15 mg	60	5	..	31.66	32.73	Topamax Sprinkle	JC
8372P NP	Capsule 25 mg	60	5	..	40.53	34.20	Topamax Sprinkle	JC
8520K NP	Capsule 50 mg	60	5	..	60.83	34.20	Topamax Sprinkle	JC

Anti-Parkinson drugs

Anticholinergic agents

Tertiary amines

BENZHEXOL HYDROCHLORIDE

1109J NP	Tablet 2 mg	200	2	..	15.32	16.39	Artane	SI
1110K NP	Tablet 5 mg	200	1	..	22.01	23.08	Artane	SI

BIPERIDEN HYDROCHLORIDE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2544X NP	Tablet 2 mg	200	2	..	*20.88	21.95	Akineton	LM

Ethers of tropine or tropine derivatives

BENZTROPINE MESYLATE

2362H NP	Tablet 2 mg	60	2	..	12.76	13.83	Benztrop	PL
3038X NP	Injection 2 mg in 2 mL	5	..	..	103.59	34.20	Cogentin	FK

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Dopaminergic agents								
<i>Dopa and dopa derivatives</i>								
LEVODOPA with BENSERAZIDE								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
2225D NP	Capsule 100 mg-25 mg	100	5	..	38.92	34.20	Madopar 125	RO
2226E NP	Capsule 200 mg-50 mg	100	5	..	50.01	34.20	Madopar	RO
2227F NP	Capsule 50 mg-12.5 mg	100	5	..	23.00	24.07	Madopar 62.5	RO
2228G NP	Tablet 200 mg-50 mg	100	5	..	50.01	34.20	Madopar	RO
2229H NP	Tablet 100 mg-25 mg	100	5	..	38.92	34.20	Madopar 125	RO
2231K NP	Capsule 100 mg-25 mg (sustained release)	100	5	..	42.00	34.20	Madopar HBS	RO
8218M NP	Dispersible tablet 50 mg-12.5 mg	100	5	..	23.00	24.07	Madopar Rapid 62.5	RO
8219N NP	Dispersible tablet 100 mg-25 mg	100	5	..	38.92	34.20	Madopar Rapid 125	RO
LEVODOPA with CARBIDOPA								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
1242J NP	Tablet 100 mg-25 mg	100	5	..	38.94	34.20 ^a	Kinson	AF
				^B 5.20	44.14	34.20 ^a	Sinemet 100/25	MK
1245M NP	Tablet 250 mg-25 mg	100	5	..	45.79	34.20 ^a	Levo/Carbidopa Sandoz	SZ
				^B 2.98	48.77	34.20 ^a	Sinemet	MK
LEVODOPA with CARBIDOPA								
<u>Authority required (STREAMLINED)</u>								
1257								
Parkinson's disease where fluctuations in motor function are not adequately controlled by frequent dosing with conventional formulations of levodopa with decarboxylase inhibitor.								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
1255C NP	Tablet 200 mg-50 mg (modified release)	100	5	..	67.87	34.20	Sinemet CR	MK
LEVODOPA with CARBIDOPA and ENTACAPONE								
<u>Authority required (STREAMLINED)</u>								
3305								
Parkinson disease in patients being treated with levodopa—decarboxylase inhibitor combinations who are experiencing fluctuations in motor function due to end-of-dose effect;								
3306								
Parkinson disease in patients stabilised on concomitant treatment with levodopa—decarboxylase inhibitor combinations and entacapone.								
<u>Note</u>								
Continuing Therapy Only:								
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8797B	Tablet 50 mg-12.5 mg-200 mg	200	4	..	*311.88	34.20	Stalevo	NV

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
NP							50/12.5/200mg	
8798C	Tablet 100 mg-25 mg-200 mg	200	4	..	*341.92	34.20	Stalevo	NV
NP							100/25/200mg	
8799D	Tablet 150 mg-37.5 mg-200 mg	200	4	..	*371.96	34.20	Stalevo	NV
NP							150/37.5/200mg	
9292C	Tablet 200 mg-50 mg-200 mg	200	4	..	*399.62	34.20	Stalevo	NV
NP							200/50/200mg	
9344T	Tablet 75 mg-18.75 mg-200 mg	200	4	..	*325.12	34.20	Stalevo	NV
NP							75/18.75/200mg	
9345W	Tablet 125 mg-31.25 mg-200 mg	200	4	..	*353.96	34.20	Stalevo	NV
NP							125/31.25/200mg	

Adamantane derivatives

AMANTADINE HYDROCHLORIDE

Restricted benefit

Parkinson's disease which is not drug induced.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

3016R	Capsule 100 mg	100	5	..	44.30	34.20	Symmetrel 100	NV
NP								

Dopamine agonists

BROMOCRIPTINE MESYLATE

Restricted benefit

Acromegaly;

Parkinson's disease;

Pathological hyperprolactinaemia where surgery is not indicated;

Pathological hyperprolactinaemia where surgery has already been used with incomplete resolution;

Pathological hyperprolactinaemia where radiotherapy is not indicated;

Pathological hyperprolactinaemia where radiotherapy has already been used with incomplete resolution.

Note

Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.

1443Y	Tablet 2.5 mg (base)	60	5	..	32.73	33.80	^a Krypton 2.5	AF
				^B 2.92	35.65	33.80	^a Parlodel	NV
1445C	Capsule 10 mg (base)	100	5	..	155.94	34.20	^a Krypton 10	AF
				^B 3.08	159.02	34.20	^a Parlodel	NV
1446D	Capsule 5 mg (base)	60	5	..	50.25	34.20	^a Krypton 5	AF
				^B 2.91	53.16	34.20	^a Parlodel	NV

CABERGOLINE

Restricted benefit

Parkinson's disease.

Note

Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8393R	Tablet 1 mg	30	5	..	60.87	34.20	^a Bergoline 1	SI
NP							^a Cabaser	PF
8394T	Tablet 2 mg	30	5	..	79.41	34.20	^a Bergoline 2	SI
NP							^a Cabaser	PF

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
PERGOLIDE MESYLATE							
<u>Restricted benefit</u>							
Parkinson's disease as adjunctive therapy in patients being treated with levodopa—decarboxylase inhibitor combinations.							
<u>Note</u>							
Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.							
<u>Note</u>							
Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
2808T NP	Tablet 50 micrograms (base)	100	..	..	52.93	34.20	Permax AS
2809W NP	Tablet 250 micrograms (base)	100	5	..	66.18	34.20	Permax AS
2810X NP	Tablet 1 mg (base)	100	5	..	241.69	34.20	Permax AS
PRAMIPEXOLE HYDROCHLORIDE							
<u>Caution</u>							
Episodes of sudden onset of sleep without warning, during activity, have been reported with this drug.							
<u>Note</u>							
Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.							
<u>Restricted benefit</u>							
Parkinson disease.							
<u>Note</u>							
Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
9151P NP	Tablet 125 micrograms	30	..	..	11.74	12.81	Sifrol BY
9152Q NP	Tablet 250 micrograms	100	5	..	41.27	34.20	Sifrol BY
9153R NP	Tablet 1 mg	100	5	..	152.14	34.20	Sifrol BY
PRAMIPEXOLE HYDROCHLORIDE							
<u>Caution</u>							
Episodes of sudden onset of sleep without warning, during activity, have been reported with this drug.							
<u>Note</u>							
Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.							
<u>Restricted benefit</u>							
Treatment of severe primary Restless Legs Syndrome in a patient who manifests all 4 diagnostic criteria below and whose baseline International Restless Legs Syndrome Rating Scale (IRLSRS) score is greater than or equal to 21 points prior to initiation of pramipexole.							
The date and IRLSRS score must be documented in the patient's medical records at the time pramipexole treatment is initiated.							
The diagnostic criteria for Restless Legs Syndrome are:							
(a) An urge to move the legs usually accompanied or caused by unpleasant sensations in the legs; and							
(b) The urge to move or unpleasant sensations begin or worsen during periods of rest or inactivity such as lying or sitting; and							
(c) The urge to move or unpleasant sensations are partially or totally relieved by movement, such as walking or stretching, at least as long as the activity continues; and							
(d) The urge to move or unpleasant sensations are worse in the evening or night than during the day or only occur during the evening or night.							
Pramipexole is not PBS-subsidised for Restless Legs Syndrome secondary to other causes.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised.							
<u>Note</u>							
Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
9393J NP	Tablet 125 micrograms	30	2	..	11.74	12.81	Sifrol BY

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$		
9394K NP	Tablet 250 micrograms	100	2	..	41.27	34.20	Sifrol	BY

PRAMIPEXOLE HYDROCHLORIDE

Caution

Episodes of sudden onset of sleep without warning, during activity, have been reported with this drug.

Note

Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.

Restricted benefit

Parkinson disease.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

No applications for increased maximum quantities and/or repeats will be approved for extended release pramipexole formulations.

3418X NP	Tablet 0.375 mg (extended release)	30	..	..	22.39	23.46	Sifrol ER	BY
3419Y NP	Tablet 0.75 mg (extended release)	30	5	..	37.84	34.20	Sifrol ER	BY
3420B NP	Tablet 1.5 mg (extended release)	30	5	..	66.51	34.20	Sifrol ER	BY
3421C NP	Tablet 3 mg (extended release)	30	5	..	137.56	34.20	Sifrol ER	BY
3422D NP	Tablet 4.5 mg (extended release)	30	5	..	203.13	34.20	Sifrol ER	BY

Monoamine oxidase type B inhibitors

SELEGILINE HYDROCHLORIDE

Restricted benefit

Late stage Parkinson's disease as adjunctive therapy in patients being treated with levodopa—decarboxylase inhibitor combinations.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1973W NP	Tablet 5 mg	100	5	..	53.82	34.20 ^a	Eldepryl	AS
						^a	Selgene	AF

Other dopaminergic agents

ENTACAPONE

Authority required (STREAMLINED)

2067

Parkinson's disease as adjunctive therapy in patients being treated with levodopa—decarboxylase inhibitor combinations who are experiencing fluctuations in motor function due to end-of-dose effect.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8367J NP	Tablet 200 mg	200	4	..	*281.82	34.20	Comtan	NV

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
Psycholeptics							

Antipsychotics

Phenothiazine with aliphatic side-chain

CHLORPROMAZINE HYDROCHLORIDE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1195X NP	Injection 50 mg in 2 mL	10	..	..	20.48	21.55	Largactil	SW
1196Y NP	Tablet 10 mg	100	5	..	10.49	11.56	Largactil	SW
1197B NP	Tablet 25 mg	100	5	..	11.09	12.16	Largactil	SW
1199D NP	Tablet 100 mg	100	5	..	17.44	18.51	Largactil	SW
1201F NP	Mixture 25 mg per 5 mL, 100 mL	±1	5	..	12.57	13.64	Largactil	SW

Phenothiazine with piperazine structure

FLUPHENAZINE DECANOATE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1001Q NP	Injection 50 mg in 2 mL	5	..	..	37.65	34.20	Modectate	BQ
1046C NP	Injection 12.5 mg in 0.5 mL	5	..	..	19.22	20.29	Modectate	BQ
3098C NP	Injection 25 mg in 1 mL	5	..	..	26.38	27.45	Modectate	BQ

TRIFLUOPERAZINE HYDROCHLORIDE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2185B NP	Tablet 1 mg (base)	100	5	..	13.31	14.38	Stelazine	GH
2186C NP	Tablet 5 mg (base)	100	5	..	13.86	14.93	Stelazine	GH
2386N NP	Tablet 2 mg (base)	100	5	..	13.48	14.55	Stelazine	GH

Phenothiazines with piperidine structure

PERICYAZINE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

3052P NP	Tablet 2.5 mg	100	5	..	10.41	11.48	Neulactil	SW
3053Q NP	Tablet 10 mg	100	5	..	14.46	15.53	Neulactil	SW

Butyrophenone derivatives

HALOPERIDOL

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
2761H NP	Tablet 500 micrograms	100	5	..	10.05	11.12	Serenace	SI
2763K NP	Oral liquid 2 mg per mL, 100 mL	1	5	..	17.47	18.54	Serenace	SI
2767P NP	Tablet 1.5 mg	100	5	..	10.53	11.60	Serenace	SI
2768Q NP	Injection 5 mg in 1 mL	10	..	..	22.28	23.35	Serenace	SI
2770T NP	Tablet 5 mg	50	5	..	10.54	11.61	Serenace	SI

HALOPERIDOL DECANOATE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2765M NP	I.M. injection equivalent to 50 mg haloperidol in 1 mL	5	..	..	26.67	27.74	Haldol decanoate	JC
2766N NP	I.M. injection equivalent to 150 mg haloperidol in 3 mL	5	..	..	46.23	34.20	Haldol decanoate	JC

Indole derivatives

ZIPRASIDONE HYDROCHLORIDE

Authority required (STREAMLINED)

1589

Schizophrenia.

Authority required (STREAMLINED)

3084

Monotherapy, for up to 6 months, of an episode of acute mania or mixed episodes associated with bipolar I disorder.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9070J NP	Capsule 20 mg (base)	60	5	..	90.61	34.20	Zeldox	PF
9071K NP	Capsule 40 mg (base)	60	5	..	175.11	34.20	Zeldox	PF
9072L NP	Capsule 60 mg (base)	60	5	..	253.63	34.20	Zeldox	PF
9073M NP	Capsule 80 mg (base)	60	5	..	330.43	34.20	Zeldox	PF

Thioxanthene derivatives

FLUPENTHIXOL DECANOATE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2255Q NP	Oily I.M. injection 20 mg in 1 mL	5	..	..	19.22	20.29	Fluanxol Depot	LU
2256R NP	Oily I.M. injection 40 mg in 2 mL	5	..	..	26.38	27.45	Fluanxol Depot	LU
2257T NP	Oily I.M. injection 100 mg in 1 mL	5	..	..	44.87	34.20	Fluanxol Concentrated Depot	LU

ZUCLOPENTHIXOL DECANOATE

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8097E NP	Oily I.M. injection 200 mg in 1 mL	5	..	..	25.34	26.41	Clopixol Depot	LU

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
<i>Diazepines, oxazepines, thiazepines and oxepines</i>								
OLANZAPINE								
<u>Authority required (STREAMLINED)</u>								
1589								
Schizophrenia;								
2044								
Maintenance treatment of bipolar I disorder.								
<u>Note</u>								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8170B NP	Tablet 2.5 mg	28	5	..	53.55	34.20	Zyprexa	LY
8185T NP	Tablet 5 mg	28	5	..	99.27	34.20	Zyprexa	LY
8186W NP	Tablet 7.5 mg	28	5	..	147.11	34.20	Zyprexa	LY
8187X NP	Tablet 10 mg	28	5	..	194.00	34.20	Zyprexa	LY
8433W NP	Wafer 5 mg	28	5	..	99.27	34.20	Zyprexa Zydys	LY
8434X NP	Wafer 10 mg	28	5	..	194.00	34.20	Zyprexa Zydys	LY
OLANZAPINE								
<u>Authority required (STREAMLINED)</u>								
1589								
Schizophrenia.								
<u>Caution</u>								
Monitor for post-injection syndrome for at least three hours after each injection.								
<u>Note</u>								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
<u>Note</u>								
Special Pricing Arrangements apply.								
9294E NP	Powder for injection 210 mg (as pamoate monohydrate) with diluent	2	5	..	*499.78	34.20	Zyprexa Relprevv	LY
9295F NP	Powder for injection 300 mg (as pamoate monohydrate) with diluent	2	5	..	*809.26	34.20	Zyprexa Relprevv	LY
9303P NP	Powder for injection 405 mg (as pamoate monohydrate) with diluent	1	5	..	499.78	34.20	Zyprexa Relprevv	LY
QUETIAPINE								
<u>Authority required (STREAMLINED)</u>								
1589								
Schizophrenia;								
2765								
Monotherapy, for up to 6 months, of an episode of acute mania associated with bipolar I disorder;								
2044								
Maintenance treatment of bipolar I disorder.								
<u>Note</u>								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
5458G NP	Tablet (modified release) 150 mg (as fumarate)	60	5	..	139.47	34.20	Seroquel XR	AP
8456C NP	Tablet 25 mg (as fumarate)	60	5	..	53.66	34.20	Seroquel	AP
8457D NP	Tablet 100 mg (as fumarate)	90	5	..	139.47	34.20	Seroquel	AP

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8458E NP	Tablet 200 mg (as fumarate)	60	5	..	187.70	34.20	Seroquel	AP
8580N NP	Tablet 300 mg (as fumarate)	60	5	..	266.23	34.20	Seroquel	AP
9202H NP	Tablet (modified release) 50 mg (as fumarate)	60	5	..	100.45	34.20	Seroquel XR	AP
9203J NP	Tablet (modified release) 200 mg (as fumarate)	60	5	..	187.70	34.20	Seroquel XR	AP
9204K NP	Tablet (modified release) 300 mg (as fumarate)	60	5	..	266.23	34.20	Seroquel XR	AP
9205L NP	Tablet (modified release) 400 mg (as fumarate)	60	5	..	354.02	34.20	Seroquel XR	AP

Benzamides

AMISULPRIDE

Authority required (STREAMLINED)

1589

Schizophrenia.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8594H NP	Tablet 100 mg	30	5	..	33.17	34.20	^a Amisulpride 100	WA
							Winthrop	
							^a Amisulpride Sandoz	SZ
							^a Solian 100	SW
8595J NP	Tablet 200 mg	60	5	..	116.64	34.20	^a Sulprix	AF
							^a Amisulpride 200	WA
							Winthrop	
							^a Amisulpride Sandoz	SZ
8596K NP	Tablet 400 mg	60	5	..	206.11	34.20	^a Solian 200	SW
							^a Sulprix	AF
							^a Amipride 400	SI
							^a Amisulpride 400	WA
8736T NP	Oral solution 100 mg per mL, 60 mL	2	5	..	*148.74	34.20	Winthrop	
							^a Amisulpride Sandoz	SZ
							^a Solian 400	SW
							^a Sulprix	AF
							Solian Solution	SW

Lithium

LITHIUM CARBONATE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

3059B NP	Tablet 250 mg	200	2	..	16.89	17.96	Lithicarb	AS
8290H NP	Tablet 450 mg (slow release)	200	2	..	*34.30	34.20	Quilonum SR	GK

Other antipsychotics

ARIPIRAZOLE

Authority required (STREAMLINED)

1589

Schizophrenia.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty	Safety Net		
\$								
\$								
Note								
Shared Care Model:								
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8717T	Tablet 10 mg	30	5	..	152.25	34.20	Abilify	BQ
NP								
8718W	Tablet 15 mg	30	5	..	212.56	34.20	Abilify	BQ
NP								
8719X	Tablet 20 mg	30	5	..	253.43	34.20	Abilify	BQ
NP								
8720Y	Tablet 30 mg	30	5	..	303.49	34.20	Abilify	BQ
NP								

PALIPERIDONE

Authority required (STREAMLINED)

1589

Schizophrenia.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

Special Pricing Arrangements apply to 3 mg and 6 mg strengths.

9140C NP	Tablet 3 mg (prolonged release)	28	5	..	161.07	34.20	Invega JC
9141D NP	Tablet 6 mg (prolonged release)	28	5	..	169.68	34.20	Invega JC

PALIPERIDONE

Authority required (STREAMLINED)

1589

Schizophrenia.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9142E NP	Tablet 9 mg (prolonged release)	28	5	..	226.01	34.20	Invega JC

RISPERIDONE

Authority required (STREAMLINED)

2061

Behavioural disturbances characterised by psychotic symptoms and aggression in patients with dementia where non-pharmacological methods have been unsuccessful.

Caution

In placebo controlled trials in elderly patients with dementia there was a significantly higher incidence of cerebrovascular adverse events, such as stroke (including fatalities) and transient ischaemic attacks, in patients treated with risperidone compared with patients treated with placebo.

Authority required (STREAMLINED)

3083

Treatment under the supervision of a paediatrician or psychiatrist, in combination with non-pharmacological measures, of severe behavioural disturbances in a patient aged less than 18 years with autism.

Continuing PBS-subsidised treatment under the supervision of a paediatrician or psychiatrist, in combination with non-pharmacological measures, of severe behavioural disturbances in a patient 18 years of age or older with autism who was commenced on PBS-subsidised treatment with risperidone prior to turning 18 years of age.

Behaviour disturbances are defined as severe aggression and injuries to self or others where non-pharmacological methods alone have been unsuccessful.

The diagnosis of autism must be made based on the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) or ICD-10 international classification of mental and behavioural disorders.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Note Shared Care Model: For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
8787L NP	Tablet 0.5 mg	60	2	..	30.82	31.89	^a Ozidal RA
							^a Resdone 0.5 CR
							^a Rispa SI
							^a Risperidone-DRLA RZ
							^a Risperidone-GA GM
							^a Rixadone AF
							^a APO-Risperidone TX
8788M NP	Tablet 0.5 mg (orally disintegrating)	56	2	..	*30.84	31.91	^a Risperdal JC
							^a Risperdal Quicklet JC
8789N NP	Tablet 1 mg	60	2	..	51.89	34.20	^a APO-Risperidone TX
							^a Ozidal RA
							^a Resdone 1 CR
							^a Rispa SI
							^a Risperdal JC
							^a Risperidone-DRLA RZ
							^a Risperidone-GA GM
							^a Risperidone generichealth GQ
							^a Rixadone AF
8790P NP	Tablet 1 mg (orally disintegrating)	56	2	..	*58.72	34.20	Risperdal Quicklet JC
9293D NP	Oral solution 1 mg per mL, 100 mL	‡1	2	..	118.03	34.20	Risperdal JC

RISPERIDONE

Authority required (STREAMLINED)

1589

Schizophrenia.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8869T NP	Tablet 0.5 mg	60	5	..	30.82	31.89	^a	Ozidal RA
							^a	Resdone 0.5 CR
							^a	Rispa SI
							^a	Risperidone-DRLA RZ
							^a	Risperidone-GA GM
							^a	Rixadone AF
							^a	APO-Risperidone TX
8870W NP	Tablet 0.5 mg (orally disintegrating)	56	5	..	*33.74	34.20	^a	Risperdal JC
							^a	Risperdal Quicklet JC

RISPERIDONE

Authority required (STREAMLINED)

1589

Schizophrenia.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
<u>Authority required (STREAMLINED)</u>							
2272							
Adjunctive therapy to mood stabilisers for up to 6 months, of an episode of acute mania associated with bipolar I disorder.							
<u>Note</u>							
Shared Care Model:							
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
3169T NP	Tablet 1 mg	60	5	..	51.89	34.20	^a APO-Risperidone TX
							^a Ozidal RA
							^a Resdone 1 CR
							^a Rispa SI
							^a Risperdal JC
							^a Risperidone-DRLA RZ
							^a Risperidone-GA GM
							^a Risperidone generichealth GQ
							^a Rixadone AF
3170W NP	Tablet 2 mg	60	5	..	109.48	34.20	^a APO-Risperidone TX
							^a Ozidal RA
							^a Resdone 2 CR
							^a Rispa SI
							^a Risperdal JC
							^a Risperidone-DRLA RZ
							^a Risperidone-GA GM
							^a Risperidone generichealth GQ
							^a Rixadone AF
3171X NP	Tablet 3 mg	60	5	..	165.34	34.20	^a APO-Risperidone TX
							^a Ozidal RA
							^a Resdone 3 CR
							^a Rispa SI
							^a Risperdal JC
							^a Risperidone-DRLA RZ
							^a Risperidone-GA GM
							^a Risperidone generichealth GQ
							^a Rixadone AF
3172Y NP	Tablet 4 mg	60	5	..	219.45	34.20	^a APO-Risperidone TX
							^a Ozidal RA
							^a Resdone 4 CR
							^a Rispa SI
							^a Risperdal JC
							^a Risperidone-DRLA RZ
							^a Risperidone-GA GM
							^a Risperidone generichealth GQ
							^a Rixadone AF
8100H NP	Oral solution 1 mg per mL, 100 mL	‡1	5	..	118.03	34.20	Risperdal JC
8792R NP	Tablet 1 mg (orally disintegrating)	56	5	..	*58.72	34.20	Risperdal Quicklet JC
8794W NP	Tablet 2 mg (orally disintegrating)	56	5	..	*111.48	34.20	Risperdal Quicklet JC
9075P NP	Tablet 3 mg (orally disintegrating)	56	5	..	*162.54	34.20	Risperdal Quicklet JC

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
9076Q NP	Tablet 4 mg (orally disintegrating)	56	5	..	*213.56	34.20	Risperdal Quicklet	JC

RISPERIDONE

Authority required (STREAMLINED)

1589

Schizophrenia.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

Special Pricing Arrangements apply for the I.M. injections.

8780D NP	Powder for I.M. injection 25 mg (modified release) with 2 mL diluent in pre-filled syringe	2	5	..	*309.38	34.20	Risperdal Consta	JC
8781E NP	Powder for I.M. injection 37.5 mg (modified release) with 2 mL diluent in pre-filled syringe	2	5	..	*395.12	34.20	Risperdal Consta	JC
8782F NP	Powder for I.M. injection 50 mg (modified release) with 2 mL diluent in pre-filled syringe	2	5	..	*480.22	34.20	Risperdal Consta	JC

RISPERIDONE

Authority required (STREAMLINED)

3083

Treatment under the supervision of a paediatrician or psychiatrist, in combination with non-pharmacological measures, of severe behavioural disturbances in a patient aged less than 18 years with autism.

Continuing PBS-subsidised treatment under the supervision of a paediatrician or psychiatrist, in combination with non-pharmacological measures, of severe behavioural disturbances in a patient 18 years of age or older with autism who was commenced on PBS-subsidised treatment with risperidone prior to turning 18 years of age.

Behaviour disturbances are defined as severe aggression and injuries to self or others where non-pharmacological methods alone have been unsuccessful.

The diagnosis of autism must be made based on the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) or ICD-10 international classification of mental and behavioural disorders.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9079W NP	Tablet 2 mg	60	2	..	109.48	34.20	^a APO-Risperidone	TX
							^a Ozidal	RA
							^a Resdone 2	CR
							^a Rispa	SI
							^a Risperdal	JC
							^a Risperidone-DRLA	RZ
							^a Risperidone-GA	GM
							^a Risperidone generichealth	GQ
							^a Rixadone	AF
9080X NP	Tablet 2 mg (orally disintegrating)	56	2	..	*111.48	34.20	Risperdal Quicklet	JC

Anxiolytics

Benzodiazepine derivatives

ALPRAZOLAM

Authority required

Panic disorder where other treatments have failed or are inappropriate.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
2130D NP	Tablet 250 micrograms	50	..	..	9.55	10.62	^a	Alprax 0.25 SI
							^a	Alprazolam Sandoz SZ
							^a	Kalma 0.25 AF
				^B 1.04	10.59	10.62	^a	Xanax PF
2131E NP	Tablet 500 micrograms	50	..	..	11.48	12.55	^a	Alprax 0.5 SI
							^a	Alprazolam Sandoz SZ
							^a	Kalma 0.5 AF
				^B 1.12	12.60	12.55	^a	Xanax PF
2132F NP	Tablet 1 mg	50	2	..	15.23	16.30	^a	Alprax 1 SI
							^a	Alprazolam-GA GM
							^a	Alprazolam Sandoz SZ
							^a	Chem mart CH
							^a	Alprazolam GenRx Alprazolam GX
							^a	Kalma 1 AF
							^a	Terry White Chemists TW
				^B 1.32	16.55	16.30	^a	Alprazolam Xanax PF
8118G NP	Tablet 2 mg	50	2	..	20.06	21.13	^a	Alprax 2 SI
							^a	Alprazolam-GA GM
							^a	Alprazolam Sandoz SZ
							^a	Chem mart CH
							^a	Alprazolam GenRx Alprazolam GX
							^a	Kalma 2 AF
							^a	Terry White Chemists TW
				^B 1.60	21.66	21.13	^a	Alprazolam Xanax Tri-Score PF

DIAZEPAM

Note

Authorities for increased maximum quantities and/or repeats for the oral forms of diazepam will be granted only for

(i) the treatment of disabling spasticity; or

(ii) malignant neoplasia (late stage); or

(iii) use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past six months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal; or

(iv) use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past six months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.
Up to six months' treatment (i.e. one month's treatment with five repeats) may be requested.

2558P NP	Injection 10 mg in 2 mL	5	..	..	12.29	13.36		Hospira Pty Limited HH
3161J NP	Tablet 2 mg	50	..	..	7.79	8.86	^a	Antenex 2 AF
							^a	Valpam 2 SI
				^B 0.86	8.65	8.86	^a	Valium RO
3162K NP	Tablet 5 mg	50	..	..	7.93	9.00	^a	Antenex 5 AF
							^a	Diazepam-GA GM
							^a	Ranzepam RA
							^a	Valpam 5 SI
				^B 0.88	8.81	9.00	^a	Valium RO

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
OXAZEPAM								
Note								
Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of oxazepam below.								
3132W NP	Tablet 15 mg	25	..	..	7.51	8.58	^a	Alepam 15 AF
				^B 2.75	10.26	8.58	^a	Serepax SI
3133X NP	Tablet 30 mg	25	..	..	7.67	8.74	^a	Alepam 30 AF
							^a	APO-Oxazepam TX
							^a	Murelax FM
				^B 2.75	10.42	8.74	^a	Serepax SI

OXAZEPAM

Authority required

Malignant neoplasia (late stage);

For use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.

3134Y NP	Tablet 15 mg	50	5	..	*8.60	9.67	^a	Alepam 15 AF
				^B 5.50	*14.10	9.67	^a	Serepax SI
3135B NP	Tablet 30 mg	50	5	..	*8.92	9.99	^a	Alepam 30 AF
							^a	APO-Oxazepam TX
							^a	Murelax FM
				^B 5.50	*14.42	9.99	^a	Serepax SI

Other anxiolytics

CLOMIPRAMINE HYDROCHLORIDE

Restricted benefit

Cataplexy associated with narcolepsy;

Obsessive-compulsive disorder;

Phobic disorders in adults.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1561E NP	Tablet 25 mg	50	2	..	16.83	17.90	^a	Chem mart CH
							^a	Clomipramine
							^a	GenRx GX
							^a	Clomipramine
							^a	Placil AF
							^a	Terry White TW
							^a	Chemists
				^B 3.26	20.09	17.90	^a	Clomipramine
							^a	Anafranil 25 NV

Hypnotics and sedatives

Benzodiazepine derivatives

NITRAZEPAM

Note

Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of nitrazepam below.

2723H	Tablet 5 mg	25	..	..	7.89	8.96	^a	Alodorm AF

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
NP				^B 1.53	9.42	8.96	^a Mogadon	VT

NITRAZEPAM

Authority required

Myoclonic epilepsy;

Malignant neoplasia (late stage);

For use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2732T NP	Tablet 5 mg	50	5	..	*9.36	10.43	^a Alodorm	AF
				^B 3.06	*12.42	10.43	^a Mogadon	VT

TEMAZEPAM

Note

Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of temazepam below.

2089Y NP	Tablet 10 mg	25	..	..	7.71	8.78	^a APO-Temazepam	TX
							^a Temaze	AF
							^a Temtabs	FM
				^B 1.50	9.21	8.78	^a Normison	SI

TEMAZEPAM

Authority required

Malignant neoplasia (late stage);

For use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal;

For use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past 6 months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2088X NP	Tablet 10 mg	50	5	..	*9.00	10.07	^a APO-Temazepam	TX
							^a Temaze	AF
							^a Temtabs	FM
				^B 3.00	*12.00	10.07	^a Normison	SI

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	
Psychoanaleptics							

Antidepressants

Non-selective monoamine reuptake inhibitors

AMITRIPTYLINE HYDROCHLORIDE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2417F NP	Tablet 10 mg	50	2	..	8.44	9.51	Endep 10	AF
2418G NP	Tablet 25 mg	50	2	..	8.56	9.63	Endep 25	AF
2429W NP	Tablet 50 mg	50	2	..	8.89	9.96	Endep 50	AF

CLOMIPRAMINE HYDROCHLORIDE

Restricted benefit

Cataplexy associated with narcolepsy;

Obsessive-compulsive disorder;

Phobic disorders in adults.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1561E	Tablet 25 mg	50	2	..	16.83	17.90	^a Chem mart	CH
NP							Clomipramine	
							^a GenRx	GX
							Clomipramine	
							^a Placil	AF
							^a Terry White	TW
							Chemists	
							Clomipramine	
				^B 3.26	20.09	17.90	^a Anafranil 25	NV

DOTHIEPIN HYDROCHLORIDE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1357K NP	Capsule 25 mg	50	2	..	8.96	10.03	^a	Dothep 25	AF
				^B 1.58	10.54	10.03	^a	Prothiaden	AB
1358L NP	Tablet 75 mg	30	2	..	8.96	10.03	^a	Dothep 75	AF
				^B 0.81	9.77	10.03	^a	Prothiaden	AB

DOXEPIN HYDROCHLORIDE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1011F NP	Capsule 10 mg (base)	50	2	..	8.97	10.04	Deptran 10	AF
				^B 1.87	10.84	10.04	Sinequan	PF
1012G NP	Tablet 50 mg (base)	50	2	..	9.71	10.78	Deptran 50	AF
1013H NP	Capsule 25 mg (base)	50	2	..	9.40	10.47	Deptran 25	AF
				^B 1.57	10.97	10.47	Sinequan	PF

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
IMIPRAMINE HYDROCHLORIDE							
Note							
Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
2420J NP	Tablet 10 mg	50	2	..	8.58	9.65 ^a	Toleraide 10 PQ
				^B 2.85	11.43	9.65 ^a	Tofranil 10 LM
2421K NP	Tablet 25 mg	50	2	..	8.58	9.65 ^a	Toleraide 25 PQ
				^B 2.85	11.43	9.65 ^a	Tofranil 25 LM
NORTRIPTYLINE HYDROCHLORIDE							
Restricted benefit							
Major depression where other antidepressant therapy has failed;							
Major depression where other antidepressant therapy is contraindicated.							
Note							
Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
2522R NP	Tablet 10 mg (base)	50	2	..	13.32	14.39	Allegron AS
2523T NP	Tablet 25 mg (base)	50	2	..	15.10	16.17	Allegron AS
Selective serotonin reuptake inhibitors							
CITALOPRAM HYDROBROMIDE							
Restricted benefit							
Major depressive disorders.							
8220P NP	Tablet 20 mg (base)	28	5	..	22.95	24.02 ^a	APO-Citalopram TX
						^a	Celapram AF
						^a	Celica RA
						^a	Chem mart CH
						^a	Citalopram
						^a	Ciazil GM
						^a	Citalobell BF
						^a	Citalopram 20 CR
						^a	Citalopram GQ
						^a	generichealth
						^a	Citalopram Sandoz SZ
						^a	GenRx Citalopram GX
						^a	Talam SI
						^a	Terry White TW
						^a	Chemists
						^a	Citalopram
				^B 4.45	27.40	24.02 ^a	Cipramil LU
8702B NP	Tablet 10 mg (base)	28	5	..	17.31	18.38	Celapram AF
8703C NP	Tablet 40 mg (base)	28	5	..	34.39	34.20 ^a	APO-Citalopram TX
						^a	Celapram AF
						^a	Citalopram Sandoz SZ
						^a	GenRx Citalopram GX
ESCITALOPRAM OXALATE							
Restricted benefit							
Major depressive disorders.							
8700X	Tablet 10 mg (base)	28	5	..	28.50	29.57 ^a	APO-Escitalopram TX

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
8701Y NP	Tablet 20 mg (base)	28	5	B 4.69	33.19	29.57	a Chem mart CH
							Escitalopram
							a Esipram GM
							a Esitalo SZ
							a Lexam 10 SI
							a LoxaLate AF
				B 4.69	33.19	29.57	a Terry White TW
							Chemists
							Escitalopram
							a Lexapro LU
							a APO-Escitalopram TX
							a Chem mart CH
8701Y NP	Tablet 20 mg (base)	28	5	B 4.69	33.19	29.57	Escitalopram
							a Esipram GM
							a Esitalo SZ
							a Lexam 20 SI
							a LoxaLate AF
							a Terry White TW
				B 6.86	35.50	29.71	Chemists
							Escitalopram
							a Lexapro LU
							a Chem mart CH
							Escitalopram
							a Esipram GM

ESCITALOPRAM OXALATE

Restricted benefit

Moderate to severe generalised anxiety disorder (GAD), as defined by Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) criteria, in a patient who has not responded to non-pharmacological therapy and:

- (a) for whom a GP Mental Health Care Plan, as described under item 2710 of the Medicare Benefits Schedule, has been prepared; or
- (b) who has been assessed by a psychiatrist;

Continuing PBS-subsidised treatment, for moderate to severe generalised anxiety disorder (GAD), of a patient commenced on escitalopram prior to 1 March 2008.

Restricted benefit

Moderate to severe social anxiety disorder (social phobia, SAD), as described by Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) criteria, in a patient who has not responded to non-pharmacological therapy and:

- (a) for whom a GP Mental Health Care Plan, as described under item 2710 of the Medicare Benefits Schedule, has been prepared; or
- (b) who has been assessed by a psychiatrist;

Continuing PBS-subsidised treatment, for moderate to severe social anxiety disorder (social phobia, SAD), of a patient commenced on escitalopram prior to 1 March 2008.

9432K NP	Tablet 10 mg (base)	28	5	..	28.50	29.57	a Esipram	GM
				B 4.69	33.19	29.57	a Lexapro	LU
9433L NP	Tablet 20 mg (base)	28	5	..	28.64	29.71	a Esipram	GM
				B 6.86	35.50	29.71	a Lexapro	LU

ESCITALOPRAM OXALATE

Restricted benefit

Major depressive disorders.

Restricted benefit

Moderate to severe generalised anxiety disorder (GAD), as defined by Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) criteria, in a patient who has not responded to non-pharmacological therapy and:

- (a) for whom a GP Mental Health Care Plan, as described under item 2710 of the Medicare Benefits Schedule, has been prepared; or
- (b) who has been assessed by a psychiatrist;

Continuing PBS-subsidised treatment, for moderate to severe generalised anxiety disorder (GAD), of a patient commenced on escitalopram prior to 1 November 2008.

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
	<u>Restricted benefit</u> Moderate to severe social anxiety disorder (social phobia, SAD), as described by Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) criteria, in a patient who has not responded to non-pharmacological therapy and: (a) for whom a GP Mental Health Care Plan, as described under item 2710 of the Medicare Benefits Schedule, has been prepared; or (b) who has been assessed by a psychiatrist; Continuing PBS-subsidised treatment, for moderate to severe social anxiety disorder (social phobia, SAD), of a patient commenced on escitalopram prior to 1 November 2008.							
8849R NP	Oral solution 10 mg (base) per mL, 28 mL	#1	5	..	34.30	34.20		Lexapro LU
	FLUOXETINE HYDROCHLORIDE <u>Restricted benefit</u> Major depressive disorders; Obsessive-compulsive disorder.							
1434L NP	Capsule 20 mg (base)	28	5	..	20.81	21.88	^a	Auscap SI ^a Chem mart CH Fluoxetine ^a Fluohexal SZ ^a Fluoxebell BF ^a Fluoxetine 20 CR ^a Fluoxetine-GA GM ^a Fluoxetine generichealth GQ ^a GenRx Fluoxetine GX ^a Lovan AL ^a Terry White Chemists TW Fluoxetine ^a Zactin AF
8270G NP	Tablet 20 mg (base) (dispersible)	28	5	B4.13 .. B4.13	24.94 20.81 24.94	21.88 21.88 21.88	^a	Prozac 20 LY Lovan 20 Tab AL Prozac Tab LY
	FLUVOXAMINE MALEATE <u>Restricted benefit</u> Major depressive disorders; Obsessive-compulsive disorder.							
8174F NP	Tablet 100 mg	30	5	..	27.55	28.62	^a	APO-Fluvoxamine TX ^a Faverin 100 SI ^a Movox 100 AF ^a Voxam SZ ^a Luvox SM
8512B NP	Tablet 50 mg	30	5	B2.95 .. B2.96	30.50 20.37 23.33	28.62 21.44 21.44	^a	Luvanex TX ^a APO-Fluvoxamine TX ^a Faverin 50 SI ^a Movox 50 AL ^a Voxam SZ ^a Luvon SM
	PAROXYTINE <u>Restricted benefit</u> Major depressive disorders; Obsessive-compulsive disorder; Panic disorder.							

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Note Bioequivalence has been demonstrated between paroxetine tablet 20 mg (as hydrochloride) and paroxetine tablet 20 mg (as mesilate).							
2242B NP	Tablet 20 mg (as hydrochloride)	30	5	..	25.48	26.55	^a Chem mart Paroxetine ^a Extine 20 ^a GenRx Paroxetine ^a Paroxetine 20 ^a Paroxetine-DP ^a Paroxetine-GA ^a Paroxetine Sandoz ^a Paxtine ^a Terry White Chemists Paroxetine ^B 0.86 26.34 26.55 ^a Aropax GK
9197C NP	Tablet 20 mg (as mesilate)	30	5	..	25.48	26.55	^a Paroxetine generichealth ^a Pharmacor Paroxo 20 CR
SERTRALINE HYDROCHLORIDE Restricted benefit Major depressive disorders.							
2236Q NP	Tablet 50 mg (base)	30	5	..	24.66	25.73	^a Chem mart Sertraline ^a Concorz ^a Eleva 50 ^a GenRx Sertraline ^a Sertra 50 ^a Sertraline 50 ^a Sertraline-GA ^a Sertraline generichealth ^a Sertraline Winthrop ^a Setrona ^a Terry White Chemists Sertraline ^a Xydep 50 GN
2237R NP	Tablet 100 mg (base)	30	5	..	24.66	25.73	^B 1.49 26.15 25.73 ^a Zoloft PF ^a Chem mart Sertraline ^a Concorz ^a Eleva 100 ^a GenRx Sertraline ^a Sertra 100 ^a Sertraline 100 ^a Sertraline-GA ^a Sertraline generichealth ^a Setrona ^a Terry White Chemists Sertraline ^a Xydep 100 GN
				^B 1.49	26.15	25.73	^a Zoloft PF

Monoamine oxidase inhibitors, non-selective

Monoamine oxidase type A inhibitors

1900B	Tablet 150 mg	60	5	..	18.76	19.83	^a	Amira 150	AF
<i>NP</i>								Chem mart Moclobemide	CH
							^a	Clobemix	GM
							^a	GenRx Moclobemide	GX
							^a	Moclobemide Sandoz	SZ
							^a	Mohexal	HX
							^a	Terry White Chemists Moclobemide	TW
				^b 0.72	19.48	19.83	^a	Aurorix	RO
8003F	Tablet 300 mg	60	5	..	30.18	31.25	^a	Amira 300	AF
<i>NP</i>								Chem mart Moclobemide	CH
							^a	Clobemix	GM
							^a	GenRx Moclobemide	GX
							^a	Moclobemide Sandoz	SZ

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
							^a Terry White Chemists	TW
				^B 1.45	31.63	31.25	^a Moclobemide Aurorix 300 mg	RO

Other antidepressants

DESVENLAFAXINE SUCCINATE

Restricted benefit

Major depressive disorders.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9366Y NP	Tablet 50 mg (base) (extended release)	28	5	..	43.31	34.20	Pristiq	WX
9367B NP	Tablet 100 mg (base) (extended release)	28	5	..	50.42	34.20	Pristiq	WX

DULOXETINE HYDROCHLORIDE

Restricted benefit

Major depressive disorders.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9155W NP	Capsule 30 mg (base)	28	..	..	38.22	34.20	Cymbalta	LY
9156X NP	Capsule 60 mg (base)	28	5	..	50.42	34.20	Cymbalta	LY

LITHIUM CARBONATE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

3059B NP	Tablet 250 mg	200	2	..	16.89	17.96	Lithicarb	AS
8290H NP	Tablet 450 mg (slow release)	200	2	..	*34.30	34.20	Quilonum SR	GK

MIANSERIN HYDROCHLORIDE

Caution

Neutropenia and agranulocytosis are more frequent in the elderly, especially in the early months of therapy.

Restricted benefit

Severe depression.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1627P NP	Tablet 10 mg	50	5	..	15.56	16.63	^a Lumin 10	AF
				^B 1.91	17.47	16.63	^a Tolvon	SH
1628Q NP	Tablet 20 mg	50	5	..	25.73	26.80	^a Lumin 20	AF
				^B 2.85	28.58	26.80	^a Tolvon	SH

MIRTAZAPINE

Restricted benefit

Major depressive disorders.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8513C NP	Tablet 30 mg	30	5	..	27.34	28.41	^a	Axit 30 AF
							^a	Chem mart CH
							^a	Mirtazapine GX
							^a	GenRx Mirtazapine GM
							^a	Mirtazapine-DP SZ
							^a	Mirtazapine Sandoz SI
							^a	Mirtazon Terry White Chemists TW
				^B 3.09	30.43	28.41	^a	Avanza SH
8855C NP	Tablet 15 mg (orally disintegrating)	30	5	..	20.37	21.44		Avanza SolTab SH
8856D NP	Tablet 30 mg (orally disintegrating)	30	5	..	27.34	28.41		Avanza SolTab SH
8857E NP	Tablet 45 mg (orally disintegrating)	30	5	..	41.24	34.20		Avanza SolTab SH
8883M NP	Tablet 45 mg	30	5	..	41.24	34.20	^a	APO-Mirtazapine TX
							^a	Chem mart CH
							^a	Mirtazapine SZ
							^a	Mirtazapine Sandoz SI
							^a	Mirtazon Terry White Chemists TW
				^B 2.91	44.15	34.20	^a	Avanza SH
9365X NP	Tablet 15 mg	30	5	..	20.37	21.44		Axit 15 AF
REBOXETINE MESILATE Restricted benefit Major depressive disorders.								
Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8583R NP	Tablet 4 mg (base)	60	5	..	38.76	34.20		Edronax PF
VENLAFAXINE HYDROCHLORIDE Restricted benefit Major depressive disorders.								
Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.								
8301X NP	Capsule 75 mg (base) (modified release)	28	5	..	43.31	34.20		Efexor-XR WX
8302Y NP	Capsule 150 mg (base) (modified release)	28	5	..	50.42	34.20		Efexor-XR WX
8868R NP	Capsule 37.5 mg (base) (modified release)	28	..	..	27.53	28.60		Efexor-XR WX

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Psychostimulants, agents used for ADHD and nootropics

Centrally acting sympathomimetics

ATOMOXETINE HYDROCHLORIDE

Authority required

Initial sole PBS-subsidised treatment of attention-deficit hyperactivity disorder (ADHD) diagnosed between the ages of 6 and 18 years inclusive, by a paediatrician or psychiatrist according to the DSM-IV criteria, where:

(a) treatment with dexamphetamine sulfate or methylphenidate hydrochloride poses an unacceptable medical risk due to the following contraindications as specified in the TGA-approved product information:

- (1) The patient has a history of substance abuse or misuse (other than alcohol); and/or
- (2) The patient has comorbid motor tics or Tourette's Syndrome; and/or
- (3) The patient has comorbid severe anxiety diagnosed according to the DSM-IV; or

(b) treatment with dexamphetamine sulfate or methylphenidate hydrochloride has resulted in the development or worsening of a comorbid mood disorder (diagnosed according to the DSM-IV criteria i.e. anxiety disorder, obsessive compulsive disorder, depressive disorder) of a severity necessitating permanent stimulant treatment withdrawal; or where the combination of stimulant treatment with another agent would pose an unacceptable medical risk of a severity necessitating permanent stimulant treatment withdrawal; or

(c) treatment with dexamphetamine sulfate AND methylphenidate hydrochloride has resulted in the development of adverse reactions of a severity necessitating permanent treatment withdrawal:

- (1) Adverse effects on growth and weight; and/or
- (2) Adverse effects on sleep including insomnia; and/or
- (3) Adverse effects on appetite including anorexia.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Authority required

Continuing sole PBS-subsidised treatment where the patient has previously been issued with an authority prescription for this drug.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9092M	Capsule 10 mg (base)	56	5	..	*221.18	34.20	Strattera	LY
9093N	Capsule 18 mg (base)	56	5	..	*221.18	34.20	Strattera	LY
9094P	Capsule 25 mg (base)	56	5	..	*221.18	34.20	Strattera	LY
9095Q	Capsule 40 mg (base)	56	5	..	*221.18	34.20	Strattera	LY
9096R	Capsule 60 mg (base)	56	5	..	*221.18	34.20	Strattera	LY
9289X	Capsule 80 mg (base)	28	5	..	147.11	34.20	Strattera	LY
9290Y	Capsule 100 mg (base)	28	5	..	147.11	34.20	Strattera	LY

DEXAMPHETAMINE SULFATE

Note

Care must be taken to comply with the provisions of State/Territory law when prescribing dexamphetamine.

Authority required

Use in attention deficit hyperactivity disorder, in accordance with State/Territory law;

Narcolepsy.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1165H NP	Tablet 5 mg	100	5	..	18.19	19.26	Sigma Pharmaceuticals (Australia) Pty Ltd	SI

METHYLPHENIDATE HYDROCHLORIDE

Note

Care must be taken to comply with the provisions of State/Territory law when prescribing methylphenidate hydrochloride.

Authority required

Use in attention deficit hyperactivity disorder, in accordance with State/Territory law.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8839F NP	Tablet 10 mg	100	5	..	17.30	18.37	Ritalin 10	NV

METHYLPHENIDATE HYDROCHLORIDE

Note

Care must be taken to comply with the provisions of State/Territory law when prescribing methylphenidate hydrochloride.

Authority required

Treatment of attention deficit hyperactivity disorder (ADHD) in a patient diagnosed between the ages of 6 and 18 years inclusive, who has demonstrated a response to immediate release methylphenidate hydrochloride with no emergence of serious adverse events, and who requires continuous coverage over 12 hours.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2172H NP	Tablet 27 mg (extended release)	30	5	..	57.66	34.20	Concerta	JC
2387P NP	Tablet 18 mg (extended release)	30	5	..	53.30	34.20	Concerta	JC
2388Q NP	Tablet 36 mg (extended release)	30	5	..	62.36	34.20	Concerta	JC
2432B NP	Tablet 54 mg (extended release)	30	5	..	72.95	34.20	Concerta	JC

METHYLPHENIDATE HYDROCHLORIDE

Note

Care must be taken to comply with the provisions of State/Territory law when prescribing methylphenidate hydrochloride.

Authority required

Treatment of attention deficit hyperactivity disorder (ADHD) in a patient diagnosed between the ages of 6 and 18 years inclusive, who has demonstrated a response to immediate release methylphenidate hydrochloride with no emergence of serious adverse events, and who requires continuous coverage over 8 hours.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2276T NP	Capsule 20 mg (modified release)	30	5	..	46.20	34.20	Ritalin LA	NV
2280B NP	Capsule 30 mg (modified release)	30	5	..	54.05	34.20	Ritalin LA	NV
2283E NP	Capsule 40 mg (modified release)	30	5	..	56.71	34.20	Ritalin LA	NV
3440C NP	Capsule 10 mg (modified release)	30	5	..	35.35	34.20	Ritalin LA	NV

MODAFINIL

Note

Modafinil is not PBS-subsidised when used in combination with PBS-subsidised dexamphetamine sulfate.

Note

Any queries concerning the arrangements to prescribe modafinil may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe modafinil should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
<u>Authority required</u>							
Initial treatment, by a qualified sleep medicine practitioner or neurologist, of patients with narcolepsy where:							
(i) therapy with dexamphetamine sulfate poses an unacceptable medical risk; or							
(ii) intolerance to dexamphetamine sulfate of a severity necessitating treatment withdrawal develops.							
The presence of any 1 of the following indicates treatment with dexamphetamine sulfate poses an unacceptable medical risk:							
(a) a psychiatric disorder;							
(b) a cardiovascular disorder;							
(c) a history of substance abuse;							
(d) glaucoma;							
(e) any other absolute contraindication to dexamphetamine sulfate as specified in the TGA-approved Product Information.							
Patients must meet the following definition of narcolepsy:							
Excessive daytime sleepiness, recurrent naps or lapses into sleep occurring almost daily for at least 3 months and:							
(i) a definite history of cataplexy;							
or							
a mean sleep latency less than or equal to 10 minutes on a Multiple Sleep Latency Test (MSLT). The MSLT must be preceded by nocturnal polysomnography. Sleep prior to the MSLT must be at least 6 hours in duration;							
or							
an electroencephalographic (EEG) recording showing the pathologically rapid development of REM sleep; and							
(ii) absence of any medical or psychiatric disorder that could otherwise account for the hypersomnia.							
The authority application must be made in writing and must include the following:							
(a) a completed authority prescription form; and							
(b) a completed Modafinil (Modavigil) PBS Authority Application for Use in the Treatment of Narcolepsy - Supporting Information Form [www.medicareaustralia.gov.au]; and							
(c) details of the contraindication or intolerance to dexamphetamine sulfate; and							
(d) either:							
(i) the result and date of the polysomnography test and MSLT conducted by, or under the supervision of, a qualified sleep medicine practitioner; or							
(ii) the result and date of the EEG, conducted by, or under the supervision of, a neurologist.							
The polysomnography, MSLT or EEG test reports must be provided with the authority application.							
<u>Authority required</u>							
Continuing treatment of narcolepsy, where the patient has previously been issued with an authority prescription for this drug.							
8816B	Tablet 100 mg	120	5	..	*346.98	34.20	Modavigil CS

Anti-dementia drugs

Anticholinesterases

DONEPEZIL HYDROCHLORIDE

Authority required

INITIAL APPLICATION FOR THE TREATMENT OF MILD TO MODERATELY SEVERE ALZHEIMER'S DISEASE — Patients with an (S)MMSE of 10 or more. Initial treatment, as the sole PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist).

The authority application must include the result of the baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE). This baseline (S)MMSE must be a score of 10 or more. If this score is 25 - 30 points, the result of a baseline Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) may also be specified.

If an ADAS-Cog score is not supplied with the initial application, this scale cannot be used for the purpose of fulfilling the criteria for continued PBS supply.

This application must be made in writing, but initial supply may be sought by telephone.

For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. This telephone application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised;

CONTINUING TREATMENT — (S)MMSE or ADAS-Cog improvement.

Continuing treatment, as the sole PBS-subsidised therapy, following initial PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in cognitive function as measured by:

- (a) for patients with a baseline (S)MMSE score of 10 or more and less than 25, an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE);
- (b) for patients with a baseline (S)MMSE score of at least 25 points, a decrease of at least 4 points from baseline on the Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) or an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE).

The initial authority application for continuing treatment must include the relevant result from the (S)MMSE or the ADAS-Cog and must be in writing.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	

Subsequent applications for continuing treatment can be made by telephone.

Authority required

INITIAL APPLICATION FOR THE TREATMENT OF MILD TO MODERATELY SEVERE ALZHEIMER'S DISEASE — Patients with an (S)MMSE of 9 or less who require a clinician's assessment.

Initial treatment, as the sole PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease of patients with a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, who are unable to register a score of 10 or more for reasons other than their Alzheimer's disease, as specified below. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist).

Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

This application must be made in writing, but initial supply may be sought by telephone.

For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. This telephone application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment;

CONTINUING TREATMENT — Clinician assessed improvement.

Continuing treatment, as the sole PBS-subsidised therapy, following initial PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in function, based on a rating of "very much improved" or "much improved" on the Clinicians Interview Based Impression of Change (CIBIC) scale, which must be assessed by the same clinician who initiated treatment.

The initial authority application for continuing treatment must state the improvement achieved on the CIBIC scale and must be in writing.

Subsequent applications for continuing treatment can be made by telephone.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8495D NP	Tablet 5 mg	28	5	..	155.45	34.20	Aricept	PF
8496E NP	Tablet 10 mg	28	5	..	155.45	34.20	Aricept	PF

GALANTAMINE HYDROBROMIDE

Authority required

INITIAL APPLICATION FOR THE TREATMENT OF MILD TO MODERATELY SEVERE ALZHEIMER'S DISEASE — Patients with an (S)MMSE of 10 or more.

Initial treatment, as the sole PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist).

The authority application must include the result of the baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE). This baseline (S)MMSE must be a score of 10 or more. If this score is 25 - 30 points, the result of a baseline Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) may also be specified.

If an ADAS-Cog score is not supplied with the initial application, this scale cannot be used for the purpose of fulfilling the criteria for continued PBS supply.

This application must be made in writing, but initial supply may be sought by telephone.

For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. This telephone application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised;

CONTINUING TREATMENT — (S)MMSE or ADAS-Cog improvement.

Continuing treatment, as the sole PBS-subsidised therapy, following initial PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in cognitive function as measured by:

- (a) for patients with a baseline (S)MMSE score of 10 or more and less than 25, an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE);
- (b) for patients with a baseline (S)MMSE score of at least 25 points, a decrease of at least 4 points from baseline on the Alzheimer's Disease

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	Assessment Scale, cognitive sub-scale (ADAS-Cog) or an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE).						

The initial authority application for continuing treatment must include the relevant result from the (S)MMSE or the ADAS-Cog and must be in writing.

Subsequent applications for continuing treatment can be made by telephone.

Authority required

INITIAL APPLICATION FOR THE TREATMENT OF MILD TO MODERATELY SEVERE ALZHEIMER'S DISEASE — Patients with an (S)MMSE of 9 or less who require a clinician's assessment.

Initial treatment, as the sole PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease of patients with a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, who are unable to register a score of 10 or more for reasons other than their Alzheimer's disease, as specified below. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist).

Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

This application must be made in writing, but initial supply may be sought by telephone.

For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. This telephone application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment;

CONTINUING TREATMENT — Clinician assessed improvement.

Continuing treatment, as the sole PBS-subsidised therapy, following initial PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in function, based on a rating of "very much improved" or "much improved" on the Clinicians Interview Based Impression of Change (CIBIC) scale, which must be assessed by the same clinician who initiated treatment.

The initial authority application for continuing treatment must state the improvement achieved on the CIBIC scale and must be in writing.

Subsequent applications for continuing treatment can be made by telephone.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8770N NP	Capsule 8 mg (base) (prolonged release)	28	5	..	115.27	34.20	^a	Galantyl	AF
							^a	Reminyl	JC
8771P NP	Capsule 16 mg (base) (prolonged release)	28	5	..	139.58	34.20	^a	Galantyl	AF
							^a	Reminyl	JC
8772Q NP	Capsule 24 mg (base) (prolonged release)	28	5	..	165.10	34.20	^a	Galantyl	AF
							^a	Reminyl	JC

RIVASTIGMINE

Authority required

INITIAL APPLICATION FOR THE TREATMENT OF MILD TO MODERATELY SEVERE ALZHEIMER'S DISEASE — Patients with an (S)MMSE of 10 or more.

Initial treatment, as the sole PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist).

The authority application must include the result of the baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE). This baseline (S)MMSE must be a score of 10 or more. If this score is 25 - 30 points, the result of a baseline Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) may also be specified.

If an ADAS-Cog score is not supplied with the initial application, this scale cannot be used for the purpose of fulfilling the criteria for continued PBS supply.

This application must be made in writing, but initial supply may be sought by telephone.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
	For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. This telephone application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised; CONTINUING TREATMENT — (S)MMSE or ADAS-Cog improvement. Continuing treatment, as the sole PBS-subsidised therapy, following initial PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in cognitive function as measured by: (a) for patients with a baseline (S)MMSE score of 10 or more and less than 25, an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE); (b) for patients with a baseline (S)MMSE score of at least 25 points, a decrease of at least 4 points from baseline on the Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) or an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE). The initial authority application for continuing treatment must include the relevant result from the (S)MMSE or the ADAS-Cog and must be in writing. Subsequent applications for continuing treatment can be made by telephone. Authority required INITIAL APPLICATION FOR THE TREATMENT OF MILD TO MODERATELY SEVERE ALZHEIMER'S DISEASE — Patients with an (S)MMSE of 9 or less who require a clinician's assessment. Initial treatment, as the sole PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease of patients with a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, who are unable to register a score of 10 or more for reasons other than their Alzheimer's disease, as specified below. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist). Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs. This application must be made in writing, but initial supply may be sought by telephone. For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. This telephone application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised. Patients who qualify under this criterion are from 1 or more of the following groups: (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background; (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate; (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test; (4) Intellectual (developmental or acquired) disability, eg Down's syndrome; (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test; (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment; CONTINUING TREATMENT — Clinician assessed improvement. Continuing treatment, as the sole PBS-subsidised therapy, following initial PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in function, based on a rating of "very much improved" or "much improved" on the Clinicians Interview Based Impression of Change (CIBIC) scale, which must be assessed by the same clinician who initiated treatment. The initial authority application for continuing treatment must state the improvement achieved on the CIBIC scale and must be in writing. Subsequent applications for continuing treatment can be made by telephone. Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
9161E NP	Transdermal patch 9 mg (releasing approximately 4.6 mg per 24 hours)	30	5	..	166.09	34.20	Exelon Patch 5	NV
9162F NP	Transdermal patch 18 mg (releasing approximately 9.5 mg per 24 hours)	30	5	..	166.09	34.20	Exelon Patch 10	NV

RIVASTIGMINE HYDROGEN TARTRATE

Authority required

INITIAL APPLICATION FOR THE TREATMENT OF MILD TO MODERATELY SEVERE ALZHEIMER'S DISEASE — Patients with an (S)MMSE of 10 or more.

Initial treatment, as the sole PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist).

The authority application must include the result of the baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE). This baseline (S)MMSE must be a score of 10 or more. If this score is 25 - 30 points, the result of a baseline Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) may also be specified.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	
If an ADAS-Cog score is not supplied with the initial application, this scale cannot be used for the purpose of fulfilling the criteria for continued PBS supply.							

This application must be made in writing, but initial supply may be sought by telephone.

For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. This telephone application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised;

CONTINUING TREATMENT — (S)MMSE or ADAS-Cog improvement.

Continuing treatment, as the sole PBS-subsidised therapy, following initial PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in cognitive function as measured by:

- (a) for patients with a baseline (S)MMSE score of 10 or more and less than 25, an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE);
- (b) for patients with a baseline (S)MMSE score of at least 25 points, a decrease of at least 4 points from baseline on the Alzheimer's Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) or an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE).

The initial authority application for continuing treatment must include the relevant result from the (S)MMSE or the ADAS-Cog and must be in writing.

Subsequent applications for continuing treatment can be made by telephone.

Authority required

INITIAL APPLICATION FOR THE TREATMENT OF MILD TO MODERATELY SEVERE ALZHEIMER'S DISEASE — Patients with an (S)MMSE of 9 or less who require a clinician's assessment.

Initial treatment, as the sole PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease of patients with a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, who are unable to register a score of 10 or more for reasons other than their Alzheimer's disease, as specified below. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist).

Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

This application must be made in writing, but initial supply may be sought by telephone.

For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. This telephone application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment;

CONTINUING TREATMENT — Clinician assessed improvement.

Continuing treatment, as the sole PBS-subsidised therapy, following initial PBS-subsidised therapy, of mild to moderately severe Alzheimer's disease in patients with demonstrated improvement in function, based on a rating of "very much improved" or "much improved" on the Clinicians Interview Based Impression of Change (CIBIC) scale, which must be assessed by the same clinician who initiated treatment.

The initial authority application for continuing treatment must state the improvement achieved on the CIBIC scale and must be in writing.

Subsequent applications for continuing treatment can be made by telephone.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8497F NP	Capsule 1.5 mg (base)	56	5	..	155.45	34.20	Exelon	NV
8498G NP	Capsule 3 mg (base)	56	5	..	155.45	34.20	Exelon	NV
8499H NP	Capsule 4.5 mg (base)	56	5	..	155.45	34.20	Exelon	NV
8500J NP	Capsule 6 mg (base)	56	5	..	155.45	34.20	Exelon	NV
8563Q NP	Oral solution 2 mg (base) per mL, 120 mL	¥1	5	..	155.45	34.20	Exelon	NV

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	

Other anti-dementia drugs

MEMANTINE HYDROCHLORIDE

Authority required

INITIAL APPLICATION FOR THE TREATMENT OF MODERATELY SEVERE ALZHEIMER'S DISEASE — Patients with an (S)MMSE of 10 to 14.

Initial treatment, as the sole PBS-subsidised therapy, of moderately severe Alzheimer's disease. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist).

The authority application must include the result of the baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE). This baseline (S)MMSE must be a score of 10 to 14.

This application must be made in writing, but initial supply may be sought by telephone.

For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. This telephone application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised;

CONTINUING TREATMENT — (S)MMSE improvement.

Continuing treatment, as the sole PBS-subsidised therapy, following initial PBS-subsidised therapy, of moderately severe Alzheimer's disease in patients with demonstrated improvement in cognitive function as measured by an increase of at least 2 points from baseline on the Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE).

The initial authority application for continuing treatment must include the relevant result from the (S)MMSE and must be in writing.

Subsequent applications for continuing treatment can be made by telephone.

Authority required

INITIAL APPLICATION FOR THE TREATMENT OF MODERATELY SEVERE ALZHEIMER'S DISEASE — Patients with an (S)MMSE of 9 or less who require a clinician's assessment.

Initial treatment, as the sole PBS-subsidised therapy, of moderately severe Alzheimer's disease of patients with a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, who are unable to register a score of 10 to 14 for reasons other than their Alzheimer's disease, as specified below. Confirmation of this diagnosis must be made by a specialist/consultant physician (including a psychiatrist).

Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

This application must be made in writing, but initial supply may be sought by telephone.

For telephone applications, up to a maximum of 2 months' initial therapy will be authorised. This telephone application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment. For written applications where no prior telephone approval has been issued, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment;

CONTINUING TREATMENT — Clinician assessed improvement.

Continuing treatment, as the sole PBS-subsidised therapy, following initial PBS-subsidised therapy, of moderately severe Alzheimer's disease in patients with demonstrated improvement in function, based on a rating of "very much improved" or "much improved" on the Clinicians Interview Based Impression of Change (CIBIC) scale, which must be assessed by the same clinician who initiated treatment.

The initial authority application for continuing treatment must state the improvement achieved on the CIBIC scale and must be in writing.

Subsequent applications for continuing treatment can be made by telephone.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1956Y NP	Tablet 10 mg	56	5	..	109.00	34.20	^a	APO-Memantine	TX
							^a	Ebixa	LU
							^a	Memantina	SI
2059J NP	Oral drops 10 mg per g, 50 g	1	5	..	98.02	34.20		Ebixa	LU
9306T	Tablet 20 mg	28	5	..	109.00	34.20		Ebixa	LU

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer	
					Max. Qty \$	Value for Safety Net \$		
NP								

Other nervous system drugs

Parasympathomimetics

Anticholinesterases

PYRIDOSTIGMINE BROMIDE

1959D	Tablet 60 mg	150	5	..	62.76	34.20	Mestinon	VT
2608G	Tablet 180 mg (modified release)	100	5	..	*123.64	34.20	Mestinon	VT
2724J	Tablet 10 mg	100	5	..	19.67	20.74	Timespan Mestinon	VT

Choline esters

BETHANECHOL CHLORIDE

1062X NP	Tablet 10 mg	100	2	..	21.03	22.10	Uro-Carb	HA

Drugs used in addictive disorders

Drugs used in nicotine dependence

BUPROPION HYDROCHLORIDE

Note

Only one course of PBS-subsidised bupropion hydrochloride will be authorised per 12 months. The period between commencing a course of bupropion hydrochloride and varenicline tartrate must be at least 6 months. A course of treatment with bupropion hydrochloride is 9 weeks. No increased maximum quantities or repeats will be authorised. Clinical review is recommended within 2 to 3 weeks of the original prescription being requested.

Authority required

Commencement of short-term, sole PBS-subsidised, therapy as an aid to achieving abstinence in a patient who has indicated they are ready to cease smoking and:

(a) who has entered a comprehensive support and counselling program; or

(b) who is entering a comprehensive support and counselling program during the consultation at which this authority is requested.

Details of the program must be specified in the authority application.

8465M NP	Tablet 150 mg (sustained release)	30	..	..	74.46	34.20	^a Prexaton	AF
				^B 0.81	75.27	34.20	^a Zyban	GK

BUPROPION HYDROCHLORIDE

Note

Only one course of PBS-subsidised bupropion hydrochloride will be authorised per 12 months. The period between commencing a course of bupropion hydrochloride and varenicline tartrate must be at least 6 months. A course of treatment with bupropion hydrochloride is 9 weeks. No increased maximum quantities or repeats will be authorised. Clinical review is recommended within 2 to 3 weeks of the original prescription being requested.

Authority required

Completion of short-term, sole PBS-subsidised, therapy as an aid to achieving abstinence in a patient who has previously been issued with an authority prescription for this drug and who is enrolled in a comprehensive support and counselling program.

8710K NP	Tablet 150 mg (sustained release)	90	..	..	162.10	34.20	^a Prexaton	AF
				^B 0.84	162.94	34.20	^a Zyban	GK

NICOTINE

Authority required

Nicotine dependence in an Aboriginal or a Torres Strait Islander person as the sole PBS-subsidised therapy.

Note

Only 2 courses of PBS-subsidised nicotine replacement therapy will be authorised per year.

No applications for increased maximum quantities and/or repeats will be authorised.

Benefit is improved if used in conjunction with a comprehensive support and counselling program.

9198D NP	Transdermal patch releasing approximately 15 mg per 16 hours	28	2	..	55.22	34.20	Nicorette Patch	JT

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
VARENICLINE							
<u>Note</u>							
Only one course of PBS-subsidised varenicline tartrate will be authorised per year. The period between commencing a course of varenicline tartrate and bupropion hydrochloride must be at least 6 months. A course of treatment with varenicline tartrate is 12 weeks. No increased maximum quantities or repeats will be authorised. Clinical review is recommended within 2 to 3 weeks of the original prescription being requested.							
<u>Authority required</u>							
Commencement of short-term, sole PBS-subsidised, therapy as an aid to achieving abstinence in a patient who has indicated they are ready to cease smoking and:							
(a) who has entered a comprehensive support and counselling program; or							
(b) who is entering a comprehensive support and counselling program during the consultation at which this authority is requested.							
Details of the program must be specified in the authority application.							
9128K NP	Box containing 11 tablets 0.5 mg (as tartrate) and 14 tablets 1 mg (as tartrate) in the first pack and 28 tablets 1 mg (as tartrate) in the second pack	\$1	..	..	103.76	34.20	Champix PF

VARENICLINE

Note

Only one course of PBS-subsidised varenicline tartrate will be authorised per year. The period between commencing a course of varenicline tartrate and bupropion hydrochloride must be at least 6 months. A course of treatment with varenicline tartrate is 12 weeks. No increased maximum quantities or repeats will be authorised. Clinical review is recommended within 2 to 3 weeks of the original prescription being requested.

Authority required

Completion of short-term, sole PBS-subsidised, therapy as an aid to achieving abstinence in a patient who has previously been issued with an authority prescription for this drug and who is enrolled in a comprehensive support and counselling program.

9129L NP	Tablet 1 mg (as tartrate)	112	..	..	*233.06	34.20	Champix PF

Drugs used in alcohol dependence

ACAMPROSATE CALCIUM

Authority required (STREAMLINED)

2665

For use within a comprehensive treatment program for alcohol dependence with the goal of maintaining abstinence.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8357W NP	Tablet 333 mg (enteric coated)	180	1	..	166.58	34.20	Campral AF

NALTREXONE HYDROCHLORIDE

Caution

Naltrexone hydrochloride is contraindicated in patients receiving opioid drugs.

Authority required

For use within a comprehensive treatment program for alcohol dependence with the goal of maintaining abstinence.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8370M NP	Tablet 50 mg	30	1	..	138.31	34.20	^a Naltrexone GQ
						^a generichealth	
						^a Naltrexone QP	XF
						^a ReVia	BQ

Other nervous system drugs

Other nervous system drugs

RILUZOLE

Authority required

Initial treatment of amyotrophic lateral sclerosis, as diagnosed by a neurologist, in patients with disease duration of 5 years or less and who have at least 60 percent of predicted forced vital capacity within 2 months prior to commencing riluzole therapy and who:

(1) are ambulatory, and

(a) have not undergone tracheostomy, and

(b) have not experienced respiratory failure; OR

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
	(2) are not ambulatory, and (a) have not undergone tracheostomy, and (b) have not experienced respiratory failure, and (c) are either able to use upper limbs or able to swallow. The date of diagnosis and the date and results of spirometry (in terms of percent of predicted forced vital capacity) must be supplied with the initial authority application.						
	Authority required Continuing treatment of amyotrophic lateral sclerosis in patients who have previously been issued with an authority prescription for this drug and who: (1) are ambulatory, and (a) have not undergone tracheostomy, and (b) have not experienced respiratory failure; OR (2) are not ambulatory, and (a) have not undergone tracheostomy, and (b) have not experienced respiratory failure, and (c) are either able to use upper limbs or able to swallow.						
	Note Continuing Therapy Only: For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.						
8664B NP	Tablet 50 mg	56	5	..	662.00	34.20	Rilutek SW

TETRABENAZINE

Authority required (STREAMLINED)

1161

Hyperkinetic extrapyramidal disorders.

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1330B NP	Tablet 25 mg	112	5	..	337.55	34.20	Orphan Australia Pty Ltd	OA

Antiparasitic products, insecticides and repellents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Antiparasitic products, insecticides and repellents

Antiprotozoals

Agents against amoebiasis and other protozoal diseases

Nitroimidazole derivatives

METRONIDAZOLE								
1626N NP	Tablet 400 mg	5	2	..	7.88	8.95	Metrogyl 400	AF
1636D NP	Tablet 200 mg	21	1	..	7.96	9.03 ^a	Metrogyl 200	AF
							^a Metronide 200	AV
				^B 2.30	10.26	9.03 ^a	Flagyl	SW
1642K NP	Suppositories 500 mg, 10	1	..	..	23.16	24.23	Flagyl	SW
METRONIDAZOLE								
<u>Restricted benefit</u>								
Treatment of anaerobic infections.								
1621H NP	Tablet 400 mg	21	1	..	10.03	11.10 ^a	Metrogyl 400	AF
							^a Metronide 400	AV
				^B 2.30	12.33	11.10 ^a	Flagyl	SW
METRONIDAZOLE BENZOATE								
1630T NP	Oral suspension 320 mg per 5 mL (equivalent to 200 mg metronidazole in 5 mL), 100 mL	1	..	..	18.82	19.89	Flagyl S	SW
TINIDAZOLE								
1465D NP	Tablet 500 mg	4	..	..	10.88	11.95 ^a	Simplotan	GP
				^B 2.46	13.34	11.95 ^a	Fasigyn	PF

Other agents against amoebiasis and other protozoal diseases

ATOVAQUONE

Authority required (STREAMLINED)

1433

Treatment of mild to moderate *Pneumocystis carinii* pneumonia in adult patients who are intolerant of trimethoprim/sulfamethoxazole therapy.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

8300W NP	Oral suspension 750 mg per 5 mL, 210 mL	1	..	..	1034.57	34.20	Wellvone	GK
PYRIMETHAMINE								
1966L NP	Tablet 25 mg	50	..	..	16.37	17.44	Daraprim	GK

Antimalarials

Biguanides

ATOVAQUONE with PROGUANIL HYDROCHLORIDE

Authority required

Treatment of suspected or confirmed *Plasmodium falciparum* malaria in a patient aged 3 years or older where quinine containing regimens are inappropriate.

Antiparasitic products, insecticides and repellents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty	Safety Net	
<u>Note</u>							
Atovaquone with proguanil hydrochloride is not PBS-subsidised for the prophylaxis of malaria.							
<u>Note</u>							
Shared Care Model:							
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
9439T	Tablet 250 mg-100 mg	12	..	..	67.00	34.20	Malarone
NP							GK

Methanolquinolines

QUININE SULFATE

Caution

Severe thrombocytopenia has been reported with this drug.

Authority required (STREAMLINED)

2142

Malaria.

1975Y NP	Tablet 300 mg	50	2	..	14.30	15.37	Quinate AS

Artemisinin and derivatives

ARTEMETHER with LUMEFANTRINE

Authority required

Treatment of suspected or confirmed malaria due to Plasmodium falciparum.

Note

Artemether with lumefantrine is not PBS-subsidised for prophylaxis of malaria.

9498X	Tablet 20 mg-120 mg	24	..	..	96.90	34.20	Riamet NV

ARTEMETHER with LUMEFANTRINE

Authority required

Treatment of suspected or confirmed malaria due to Plasmodium falciparum in a patient unable to swallow a solid dosage form of artemether with lumefantrine.

Note

Artemether with lumefantrine is not PBS-subsidised for prophylaxis of malaria.

5296R	Tablet (dispersible) 20 mg-120 mg	18	..	..	96.90	34.20	Riamet 20mg/120mg Dispersible NV

Anthelmintics

Antitrepatodals

Quinoline derivatives and related substances

PRAZIQUANTEL

Authority required (STREAMLINED)

3147

Schistosomiasis.

9447F NP	Tablet 600 mg	8	..	..	40.85	34.20	Biltricide BN

Antinematodal agents

Benzimidazole derivatives

ALBENDAZOLE

Authority required (STREAMLINED)

2446

Treatment of whipworm infestation in an Aboriginal or a Torres Strait Islander person;

1388

Strongyloidiasis;

Antiparasitic products, insecticides and repellents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
	3241 Treatment of hookworm infestation.							
9047E NP	Tablet 200 mg	6	..	..	33.10	34.17	Zentel	GK
	ALBENDAZOLE <u>Authority required (STREAMLINED)</u> 1525 Treatment of tapeworm infestation.							
8503M NP	Tablet 200 mg	6	1	..	33.10	34.17	Zentel	GK
	ALBENDAZOLE <u>Authority required (STREAMLINED)</u> 1496 For the treatment of hydatid disease in conjunction with surgery or when a surgical cure cannot be achieved or where surgery cannot be used.							
8459F	Tablet 400 mg	60	2	..	185.25	34.20	Eskazole	GK

Tetrahydropyrimidine derivatives

PYRANTEL EMBONATE								
3047J NP	Tablet 125 mg (base)	6	..	..	8.41	9.48	Anthel 125	AF
3048K NP	Tablet 250 mg (base)	6	..	..	9.48	10.55	Anthel 250	AF

Avermectines

IVERMECTIN								
<u>Authority required (STREAMLINED)</u>								
1242								
Onchocerciasis;								
1388								
Strongyloidiasis.								
8359Y NP	Tablet 3 mg	4	..	..	31.31	32.38	Stromectol	MK

Ectoparasiticides, incl. scabicides, insecticides and repellents

Ectoparasiticides, incl. scabicides

Pyrethrines, incl. synthetic compounds

PERMETHRIN								
3054R NP	Cream 50 mg per g (5%), 30 g	±1	1	..	16.77	17.84	Lyclear	JT

Respiratory system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Respiratory system

Nasal preparations

Decongestants and other nasal preparations for topical use

*Other nasal preparations***MUPIROCIN****Authority required (STREAMLINED)****3136**

Nasal colonisation with Staphylococcus aureus in an Aboriginal or a Torres Strait Islander person.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9440W NP	Nasal ointment 20 mg (as calcium) per g (2%), 3 g	£1	..	..	20.63	21.70	Bactroban	GK

Drugs for obstructive airway diseases

Adrenergics, inhalants

*Selective beta-2-adrenoceptor agonists***EFORMOTEROL FUMARATE DIHYDRATE****Restricted benefit**

Patients with frequent episodes of asthma who are currently receiving treatment with oral corticosteroids;

Patients with frequent episodes of asthma who are currently receiving treatment with optimal doses of inhaled corticosteroids.

8136F NP	Capsule containing powder for oral inhalation 12 micrograms (for use in Foradile Aerolizer)	60	5	..	37.33	34.20	Foradile	NV
8239P NP	Powder for oral inhalation in breath actuated device 6 micrograms per dose (60 doses)	£1	5	..	26.38	27.45	Oxis Turbuhaler	AP
8240Q NP	Powder for oral inhalation in breath actuated device 12 micrograms per dose (60 doses)	£1	5	..	36.44	34.20	Oxis Turbuhaler	AP

SALBUTAMOL SULFATE

1099W NP	Capsule containing powder for oral inhalation 200 micrograms (base) (for use in Ventolin Rotahaler)	200	5	..	*18.50	19.57	Ventolin Rotacaps	GK
8288F NP	Oral pressurised inhalation 100 micrograms (base) per dose (200 doses), CFC-free formulation	2	5	..	*15.22	16.29	^a Airomir	IA
							^a Asmol CFC-free	AL
				^B 1.18	*16.40	16.29	^a Ventolin CFC-free	GK

SALBUTAMOL SULFATE**Restricted benefit**

Patients unable to achieve co-ordinated use of other metered dose inhalers containing this drug.

8354Q NP	Oral pressurised inhalation in breath actuated device 100 micrograms (base) per dose (200 doses), CFC-free formulation	2	5	..	*38.60	34.20	Airomir Autohaler	IA

SALBUTAMOL SULFATE**Restricted benefit**

Asthma in patients unable to use this drug delivered from an oral pressurised inhalation device via a spacer;

Chronic obstructive pulmonary disease in patients unable to use this drug delivered from an oral pressurised inhalation device via a spacer.

2000G NP	Nebuliser solution single dose units 2.5 mg (base) in 2.5 mL, 30	2	5	..	*18.94	20.01	^a Asmol 2.5 uni-dose	AF
							^a Butamol 2.5	SI
							^a GenRx Salbutamol	GX

Respiratory system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
							^a Pfizer Australia Pty Ltd PF
							^a Pharmacor CR
							^a Salbutamol 2.5 Salbutamol-GA GM
							^a Salbutamol Sandoz SZ
				^B 1.46	*20.40	20.01	^a Ventolin Nebules GK
2001H NP	Nebuliser solution single dose units 5 mg (base) in 2.5 mL, 30	2	5	..	*19.64	20.71	^a Asmol 5 uni-dose AF
							^a Butamol 5 SI
							^a GenRx Salbutamol GX
							^a Pharmacor CR
							^a Salbutamol 5 Salbutamol-GA GM
							^a Salbutamol Sandoz SZ
				^B 1.46	*21.10	20.71	^a Ventolin Nebules GK
2003K NP	Nebuliser solution 5 mg (base) per mL (0.5%), 30 mL	2	2	..	*19.64	20.71	Pfizer Australia Pty Ltd PF

SALMETEROL XINAFOATE**Restricted benefit**

Patients with frequent episodes of asthma who are currently receiving treatment with oral corticosteroids;

Patients with frequent episodes of asthma who are currently receiving treatment with optimal doses of inhaled corticosteroids.

8141L NP	Powder for oral inhalation in breath actuated device 50 micrograms (base) per dose (60 doses)	1	5	..	37.33	34.20	Serevent Accuhaler GK

TERBUTALINE SULFATE

1252X NP	Powder for oral inhalation in breath actuated device 500 micrograms per dose (200 doses)	1	5	..	17.83	18.90	Bricanyl Turbuhaler AP

Adrenergics and other drugs for obstructive airway diseases**BUDESONIDE with EFORMOTEROL FUMARATE DIHYDRATE****Restricted benefit**

Patients who previously had frequent episodes of asthma while receiving treatment with oral corticosteroids and who have been stabilised on concomitant inhaled eformoterol fumarate dihydrate and budesonide;

Patients who previously had frequent episodes of asthma while receiving treatment with optimal doses of inhaled corticosteroids and who have been stabilised on concomitant inhaled eformoterol fumarate dihydrate and budesonide;

For single maintenance and reliever therapy in a patient who experiences frequent asthma symptoms while receiving treatment with oral corticosteroids;

For single maintenance and reliever therapy in a patient who experiences frequent asthma symptoms while receiving treatment with inhaled corticosteroids;

For maintenance and reliever therapy in a patient who experiences frequent asthma symptoms while receiving treatment with a combination of an inhaled corticosteroid and a long-acting beta-2 agonist.

8625Y NP	Powder for oral inhalation in breath actuated device 200 micrograms-6 micrograms per dose (120 doses)	1	5	..	58.77	34.20	Symbicort Turbuhaler 200/6 AP
8796Y NP	Powder for oral inhalation in breath actuated device 100 micrograms-6 micrograms per dose (120 doses)	1	5	..	54.47	34.20	Symbicort Turbuhaler 100/6 AP

BUDESONIDE with EFORMOTEROL FUMARATE DIHYDRATE**Restricted benefit**

Patients who previously had frequent episodes of asthma while receiving treatment with oral corticosteroids and who have been stabilised on concomitant inhaled eformoterol fumarate dihydrate and budesonide;

Patients who previously had frequent episodes of asthma while receiving treatment with optimal doses of inhaled corticosteroids and who have been stabilised on concomitant inhaled eformoterol fumarate dihydrate and budesonide.

Respiratory system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer

Note

Symbicort 400/12 is not recommended nor PBS-subsidised for use as 'maintenance and reliever' therapy.

8750M NP	Powder for oral inhalation in breath actuated devices 400 micrograms-12 micrograms per dose (60 doses), 2	1	5	..	86.89	34.20	Symbicort Turbuhaler 400/12	AP

FLUTICASONE PROPIONATE with SALMETEROL XINAFOATE**Restricted benefit**

Patients who previously had frequent episodes of asthma while receiving treatment with oral corticosteroids and who have been stabilised on concomitant inhaled salmeterol xinafoate and fluticasone propionate;

Patients who previously had frequent episodes of asthma while receiving treatment with optimal doses of inhaled corticosteroids and who have been stabilised on concomitant inhaled salmeterol xinafoate and fluticasone propionate.

8430Q NP	Powder for oral inhalation in breath actuated device 100 micrograms-50 micrograms (base) per dose (60 doses)	1	5	..	47.20	34.20	Seretide Accuhaler 100/50	GK
8431R NP	Powder for oral inhalation in breath actuated device 250 micrograms-50 micrograms (base) per dose (60 doses)	1	5	..	59.31	34.20	Seretide Accuhaler 250/50	GK
8517G NP	Oral pressurised inhalation 50 micrograms-25 micrograms (base) per dose (120 doses), CFC-free formulation	1	5	..	47.20	34.20	Seretide MDI 50/25	GK
8518H NP	Oral pressurised inhalation 125 micrograms-25 micrograms (base) per dose (120 doses), CFC-free formulation	1	5	..	59.31	34.20	Seretide MDI 125/25	GK

FLUTICASONE PROPIONATE with SALMETEROL XINAFOATE**Restricted benefit**

Patients who previously had frequent episodes of asthma while receiving treatment with oral corticosteroids and who have been stabilised on concomitant inhaled salmeterol xinafoate and fluticasone propionate;

Patients who previously had frequent episodes of asthma while receiving treatment with optimal doses of inhaled corticosteroids and who have been stabilised on concomitant inhaled salmeterol xinafoate and fluticasone propionate;

Symptomatic treatment of chronic obstructive pulmonary disease (COPD), where the FEV1 is less than 50% predicted normal and there is a history of repeated exacerbations with significant symptoms despite regular beta-2 agonist bronchodilator therapy.

Note

Seretide is not indicated for the initiation of bronchodilator therapy in COPD.

8432T NP	Powder for oral inhalation in breath actuated device 500 micrograms-50 micrograms (base) per dose (60 doses)	1	5	..	78.67	34.20	Seretide Accuhaler 500/50	GK
8519J NP	Oral pressurised inhalation 250 micrograms-25 micrograms (base) per dose (120 doses), CFC-free formulation	1	5	..	78.67	34.20	Seretide MDI 250/25	GK

Other drugs for obstructive airway diseases, inhalants**Glucocorticoids****BECLOMETHASONE DIPROPIONATE**

8406K NP	Oral pressurised inhalation 50 micrograms per dose (200 doses), CFC-free formulation	1	5	..	19.29	20.36	Qvar 50	IA
8407L NP	Oral pressurised inhalation 100 micrograms per dose (200 doses), CFC-free formulation	1	5	..	33.46	34.20	Qvar 100	IA

BECLOMETHASONE DIPROPIONATE**Restricted benefit**

Patients unable to achieve co-ordinated use of other metered dose inhalers containing this drug.

8408M NP	Oral pressurised inhalation in breath actuated device 50 micrograms per dose (200 doses), CFC-free formulation	1	5	..	27.87	28.94	Qvar 50 Autohaler	IA
8409N NP	Oral pressurised inhalation in breath actuated device 100 micrograms per dose (200 doses), CFC-free formulation	1	5	..	39.13	34.20	Qvar 100 Autohaler	IA

Respiratory system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
BUDESONIDE								
2070Y NP	Powder for oral inhalation in breath actuated device 100 micrograms per dose (200 doses)	‡1	5	..	23.34	24.41	Pulmicort Turbuhaler	AP
2071B NP	Powder for oral inhalation in breath actuated device 200 micrograms per dose (200 doses)	‡1	5	..	31.08	32.15	Pulmicort Turbuhaler	AP
2072C NP	Powder for oral inhalation in breath actuated device 400 micrograms per dose (200 doses)	‡1	5	..	45.84	34.20	Pulmicort Turbuhaler	AP
BUDESONIDE								
<u>Authority required (STREAMLINED)</u>								
1351								
Severe chronic asthma in patients who require long-term steroid therapy and who are unable to use other forms of inhaled steroid therapy.								
2065Q NP	Nebuliser suspension single dose units 500 micrograms in 2 mL, 30	‡1	5	..	37.86	34.20	Pulmicort Respules	AP
2066R NP	Nebuliser suspension single dose units 1 mg in 2 mL, 30	‡1	5	..	49.00	34.20	Pulmicort Respules	AP
CICLESONIDE								
8853Y NP	Oral pressurised inhalation 80 micrograms per dose (120 doses), CFC-free formulation	‡1	5	..	26.15	27.22	Alvesco 80	NQ
8854B NP	Oral pressurised inhalation 160 micrograms per dose (120 doses), CFC-free formulation	‡1	5	..	42.25	34.20	Alvesco 160	NQ
FLUTICASONE PROPIONATE								
8147T NP	Powder for oral inhalation in breath actuated device 100 micrograms per dose (60 doses)	‡1	5	..	17.09	18.16	Flixotide Junior Accuhaler	GK
8148W NP	Powder for oral inhalation in breath actuated device 250 micrograms per dose (60 doses)	‡1	5	..	30.66	31.73	Flixotide Accuhaler	GK
8149X NP	Powder for oral inhalation in breath actuated device 500 micrograms per dose (60 doses)	‡1	1	..	49.72	34.20	Flixotide Accuhaler	GK
8345F NP	Oral pressurised inhalation 125 micrograms per dose (120 doses), CFC-free formulation	‡1	5	..	30.66	31.73	Flixotide	GK
8346G NP	Oral pressurised inhalation 250 micrograms per dose (120 doses), CFC-free formulation	‡1	1	..	49.72	34.20	Flixotide	GK
8516F NP	Oral pressurised inhalation 50 micrograms per dose (120 doses), CFC-free formulation	‡1	5	..	17.09	18.16	Flixotide Junior	GK
Anticholinergics								
IPRATROPIUM BROMIDE								
8671J NP	Oral pressurised inhalation 21 micrograms per dose (200 doses), CFC-free formulation	2	5	..	*35.28	34.20	Atrovent	BY
IPRATROPIUM BROMIDE								
<u>Restricted benefit</u>								
Asthma in patients unable to use this drug delivered from an oral pressurised inhalation device via a spacer;								
Chronic obstructive pulmonary disease in patients unable to use this drug delivered from an oral pressurised inhalation device via a spacer.								
1542E NP	Nebuliser solution single dose units 250 micrograms (anhydrous) in 1 mL, 30	2	5	..	*37.28	34.20	^a Aeron 250	SI
							^a APO-Ipratropium	TX
							^a Ipratrin	AF
							^a Ipravent	PF
				^B 0.72	*38.00	34.20	^a Atrovent	BY
8238N NP	Nebuliser solution single dose units 500 micrograms (anhydrous) in 1 mL, 30	2	5	..	*42.64	34.20	^a Aeron 500	SI
							^a APO-Ipratropium	TX
							^a Ipratrin Adult	AF
							^a Ipravent	PF
				^B 0.62	*43.26	34.20	^a Atrovent Adult	BY

Respiratory system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer	
					Max. Qty \$	Value for Safety Net \$		

TIOTROPIUM BROMIDE MONOHYDRATE**Restricted benefit**

For the long-term maintenance treatment of bronchospasm and dyspnoea associated with chronic obstructive pulmonary disease.

8626B NP	Capsule containing powder for oral inhalation 18 micrograms (base) (for use in HandiHaler)	30	5	..	76.89	34.20	Spiriva	BY

Antiallergic agents, excl. corticosteroids**NEDOCROMIL SODIUM**

8365G NP	Oral pressurised inhalation 2 mg per dose (112 doses), CFC-free formulation	¥1	5	..	37.69	34.20	Tilade CFC-Free	SW

SODIUM CROMOGLYCATE

2878L NP	Capsule containing powder for oral inhalation 20 mg (for use in Intal Spinhaler or Intal Haleromatic)	100	5	..	31.92	32.99	Intal Spincaps	GM
8334P NP	Oral pressurised inhalation 5 mg per dose (112 doses), CFC-free formulation	¥1	5	..	36.44	34.20	Intal Forte CFC-Free	SW
8767K NP	Oral pressurised inhalation 1 mg per dose (200 doses), CFC-free formulation	¥1	5	..	30.78	31.85	Intal CFC-Free	SW

Adrenergics for systemic use***Alpha- and beta-adrenoceptor agonists*****ADRENALINE**

1016L NP	Injection 1 mg in 1 mL (1 in 1,000)	5	1	..	20.34	21.41	AstraZeneca Pty Ltd	AP

ADRENALINE**Authority required**

Initial sole PBS-subsidised supply for anticipated emergency treatment of acute allergic reactions with anaphylaxis in a patient who:

(a) has been assessed to be at significant risk of anaphylaxis by, or in consultation with, a clinical immunologist, allergist, paediatrician or respiratory physician. The name of the specialist consulted must be provided at the time of application for initial supply; or

(b) has been discharged from hospital or an emergency department after treatment with adrenaline for acute allergic reaction with anaphylaxis;

Continuing sole PBS-subsidised supply for anticipated emergency treatment of acute allergic reactions with anaphylaxis, where the patient has previously been issued with an authority prescription for this drug.

Note

The auto-injector should be provided in the framework of a comprehensive anaphylaxis prevention program and an emergency action plan including training in recognition of the symptoms of anaphylaxis and the use of the auto-injector device. (For further information see the Australasian Society of Clinical Immunology and Allergy website at www.allergy.org.au.)

Note

Authority approvals will be limited to a maximum quantity of 2 auto-injectors (Anapen or EpiPen) at any one time.

No repeats will be issued.

Caution

EpiPen and Anapen products have different administration techniques and should not be prescribed to the same patient without training in their use.

3408J NP	I.M. injection 150 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	106.00	34.20	Anapen Junior	LM
3409K NP	I.M. injection 300 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	106.00	34.20	Anapen	LM
8697R NP	I.M. injection 150 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	106.00	34.20	EpiPen Jr.	AL
8698T NP	I.M. injection 300 micrograms in 0.3 mL single dose syringe auto-injector	1	..	..	106.00	34.20	EpiPen	AL

Selective beta-2-adrenoceptor agonists**SALBUTAMOL SULFATE**

1103C NP	Syrup 2 mg (base) per 5 mL, 150 mL	2	5	..	*22.20	23.27	Ventolin	GK

Respiratory system

					Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	\$	\$	Brand Name and Manufacturer	
TERBUTALINE SULFATE								
1034K NP	Injection 500 micrograms in 1 mL	5	..	..	30.59	31.66	Bricanyl	AP

Other systemic drugs for obstructive airway diseases

Xanthines**THEOPHYLLINE****Caution**

Because of variable effects of food on absorption of sustained release theophylline preparations, patients stabilised on one brand should not be changed to another without appropriate monitoring.

Note**Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

2614N NP	Syrup 133.3 mg per 25 mL, 500 mL	1	5	..	12.31	13.38	Nuelin	IA
2634P NP	Tablet 250 mg (sustained release)	100	5	..	13.32	14.39	Nuelin-SR 250	IA
8230E NP	Tablet 200 mg (sustained release)	100	5	..	12.16	13.23	Nuelin-SR 200	IA
8231F NP	Tablet 300 mg (sustained release)	100	5	..	14.70	15.77	Nuelin-SR 300	IA

Leukotriene receptor antagonists**MONTELUKAST SODIUM****Authority required (STREAMLINED)****2617**

First-line preventer medication, as the single preventer agent for children aged 2 to 5 years with frequent intermittent or mild persistent asthma, as an alternative to sodium cromoglycate or nedocromil sodium.

Note

Montelukast sodium is not PBS-subsidised for use in a child aged 2 to 5 years with moderate to severe asthma. It is not intended as an alternative for a child aged 2 to 5 years who requires a corticosteroid as a preventer medication.

Montelukast sodium is not subsidised in a child aged 2 to 5 years for use in combination with other preventer medications. PBS subsidy for montelukast sodium will therefore cease for a child aged 2 to 5 years who requires a preventer medication in addition to montelukast sodium.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8627C NP	Chewable tablet 4 mg (base)	28	5	..	47.93	34.20	Singulair	MK

MONTELUKAST SODIUM**Authority required (STREAMLINED)****2618**

First-line preventer medication, as the single preventer agent for children aged 6 to 14 years with frequent intermittent or mild persistent asthma, as an alternative to sodium cromoglycate or nedocromil sodium.

Authority required (STREAMLINED)**3217**

Prevention of exercise-induced asthma, as an alternative to adding salmeterol xinafoate or formoterol fumarate, in a child aged 6 to 14 years whose asthma is otherwise well controlled while receiving optimal dose inhaled corticosteroid, but who requires short-acting beta-2 agonist 3 or more times per week for prevention or relief of residual exercise-related symptoms.

Note

Montelukast sodium is not PBS-subsidised for use in a patient aged 15 years or older, or for use in addition to a long-acting beta-agonist in any age group, or for use as a single second line preventer, as an alternative to corticosteroids, in a child aged 6 to 14 years with moderate to severe asthma.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8628D NP	Chewable tablet 5 mg (base)	28	5	..	45.71	34.20	Singulair	MK

Respiratory system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	
Cough and cold preparations							

Cough suppressants, excl. combinations with expectorants

Opium alkaloids and derivatives

CODEINE PHOSPHATE							
1214X	Tablet 30 mg	20	..	..	16.87	17.94	Fawns and McAllan Proprietary Limited FM

Antihistamines for systemic use							

Antihistamines for systemic use

Phenothiazine derivatives

PROMETHAZINE HYDROCHLORIDE							
1948M NP	Injection 50 mg in 2 mL	10	..	..	*22.32	23.39	Hospira Pty Limited HH

Sensory organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer	
					Max. Qty \$	Value for Safety Net \$		

Sensory organs

Ophthalmologicals

Antiinfectives

Antibiotics

AZITHROMYCIN

Restricted benefit

Trachoma.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

8201P NP	Powder for oral suspension 200 mg per 5 mL, 15 mL	£1	..	..	#21.35	22.76	Zithromax	PF
8336R NP	Tablet 500 mg	2	2	..	21.39	22.46	^a Azithromycin Sandoz	SZ
							^a Zithromax	PF

CHLORAMPHENICOL

1171P NP,MW	Eye ointment 10 mg per g (1%), 4 g	£1	..	..	9.76	10.83	Chloromycetin	PF
							Chlorsig	SI
2360F NP,MW	Eye drops 5 mg per mL (0.5%), 10 mL	£1	2	..	11.00	12.07	Chloromycetin	PF
							Chlorsig	SI

GENTAMICIN SULFATE

Restricted benefit

Invasive ocular infection;

Perioperative use in ophthalmic surgery;

Suspected pseudomonal eye infection.

1441W	Eye drops 3 mg (base) per mL (0.3%), 5 mL	£1	2	..	18.53	19.60	Genoptic	AG

TOBRAMYCIN

Restricted benefit

Invasive ocular infection;

Perioperative use in ophthalmic surgery;

Suspected pseudomonal eye infection.

2328M	Eye drops 3 mg per mL (0.3%), 5 mL	£1	2	..	19.28	20.35	Tobrex	AQ
2329N	Eye ointment 3 mg per g (0.3%), 3.5 g	£1	..	..	22.38	23.45	Tobrex	AQ

Sulfonamides

SULFACETAMIDE SODIUM

2063N NP	Eye drops 100 mg per mL (10%), 15 mL	£1	2	..	14.96	16.03	Bleph 10	AG

Antivirals

ACICLOVIR

Restricted benefit

Herpes simplex keratitis.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1002R NP	Eye ointment 30 mg per g (3%), 4.5 g	£1	..	..	35.06	34.20	Zovirax	GK

Sensory organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Other antiinfectives								
CIPROFLOXACIN								
<u>Authority required</u>								
Bacterial keratitis.								
1217C	Eye drops 3 mg per mL (0.3%), 5 mL	2	..	.. ^B 2.06	*29.62 *31.68	30.69 30.69	^a CiloQuin ^a Ciloxan	IQ AQ
OFLOXACIN								
<u>Authority required</u>								
Bacterial keratitis.								
8383F	Eye drops 3 mg per mL (0.3%), 5 mL	2	..	..	*32.14	33.21	Ocuflox	AG
Antiinflammatory agents								
Corticosteroids, plain								
DEXAMETHASONE								
1288T NP	Eye drops 1 mg per mL (0.1%), 5 mL	‡1	2	..	10.61	11.68	Maxidex	AQ
FLUOROMETHOLONE								
1204J NP	Eye drops 1 mg per mL (0.1%), 5 mL	‡1	5	..	10.61	11.68	Flucon FML Liquifilm	AQ AG
FLUOROMETHOLONE ACETATE								
1438Q NP	Eye drops 1 mg per mL (0.1%), 5 mL	‡1	2	..	10.61	11.68	Flarex	AQ
HYDROCORTISONE ACETATE								
1497T NP	Eye ointment 5 mg per g (0.5%), 5 g	‡1	..	..	11.57	12.64	Hycor	SI
2441L NP	Eye ointment 10 mg per g (1%), 5 g	‡1	..	..	12.81	13.88	Hycor	SI
Corticosteroids and mydriatics in combination								
PREDNISOLONE ACETATE with PHENYLEPHRINE HYDROCHLORIDE								
<u>Restricted benefit</u>								
Corneal grafts; Uveitis.								
3112T NP	Eye drops 10 mg-1.2 mg per mL (1%-0.12%), 10 mL	‡1	2	..	24.16	25.23	Prednefrin Forte	AG
Antiinflammatory agents, non-steroids								
FLURBIPROFEN SODIUM								
8699W NP	Eye drops 300 micrograms per mL (0.03%), single dose units 0.4 mL, 5	1	..	..	15.37	16.44	Ocufen	AG
Antiglaucoma preparations and miotics								
Sympathomimetics in glaucoma therapy								
APRACLOPIDINE HYDROCHLORIDE								
<u>Restricted benefit</u>								
Short-term reduction of intra-ocular pressure in patients already on maximally tolerated anti-glaucoma therapy.								
8083K	Eye drops 5 mg (base) per mL (0.5%), 10 mL	‡1	2	..	41.77	34.20	Iopidine 0.5%	AQ
BRIMONIDINE TARTRATE								
5298W	Eye drops 1.5 mg per mL (0.15%), 5 mL	‡1	5	..	20.42	21.49	Alphagan P 1.5	AG

Sensory organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8351M	Eye drops 2 mg per mL (0.2%), 5 mL	‡1	5	..	20.42	21.49 ^a	Enidin	PE
				^B 1.66	22.08	21.49 ^a	Alphagan	AG

BRIMONIDINE TARTRATE with TIMOLOL MALEATE

Restricted benefit

Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;

Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.

8826M	Eye drops 2 mg-5 mg (base) per mL (0.2%-0.5%), 5 mL	‡1	5	..	26.61	27.68	Combigan	AG

Parasympathomimetics

PILOCARPINE HYDROCHLORIDE

2595N	Eye drops 10 mg per mL (1%), 15 mL	‡1	5	..	12.65	13.72	Isopto Carpine	AQ
2596P	Eye drops 20 mg per mL (2%), 15 mL	‡1	5	..	13.93	15.00	Isopto Carpine	AQ
2598R	Eye drops 40 mg per mL (4%), 15 mL	‡1	5	..	16.84	17.91	Isopto Carpine	AQ

Carbonic anhydrase inhibitors

ACETAZOLAMIDE

Note

Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

1004W NP	Tablet 250 mg	100	3	..	23.79	24.86	Diamox	SI

BRINZOLAMIDE

8483L	Eye drops 10 mg per mL (1%), 5 mL	‡1	5	..	23.11	24.18 ^a	BrinzoQuin	IQ
				^B 1.18	24.29	24.18 ^a	Azopt	AQ

BRINZOLAMIDE with TIMOLOL MALEATE

Restricted benefit

Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;

Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.

3438Y	Eye drops 10 mg-5 mg (base) per mL (1%-0.5%), 5 mL	‡1	5	..	27.49	28.56	Azarga	AQ

DORZOLAMIDE HYDROCHLORIDE

8488R	Eye drops 20 mg (base) per mL (2%), 5 mL	‡1	5	..	21.29	22.36	Trusopt	MK

DORZOLAMIDE HYDROCHLORIDE with TIMOLOL MALEATE

Restricted benefit

Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;

Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.

8567X	Eye drops 20 mg (base)-5 mg (base) per mL (2%-0.5%), 5 mL	‡1	5	..	27.49	28.56	Cosopt	MK

Beta blocking agents

BETAXOLOL HYDROCHLORIDE

2811Y	Eye drops, suspension, 2.5 mg (base) per mL (0.25%), 5 mL	‡1	5	..	14.94	16.01	Betoptic S	AQ
2825Q	Eye drops, solution, 5 mg (base) per mL (0.5%), 5 mL	‡1	5	..	14.94	16.01 ^a	BetoQuin	IQ
				^B 2.09	17.03	16.01 ^a	Betoptic	AQ

Sensory organs

					Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net			
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	\$	\$	Brand Name and Manufacturer		
TIMOLOL MALEATE									
1278G	Eye drops 2.5 mg (base) per mL (0.25%), 5 mL	‡1	5	.. B3.20	11.80 15.00	12.87 12.87	a a	Tenopt Timoptol	SI FR
1279H	Eye drops 5 mg (base) per mL (0.5%), 5 mL	‡1	5	.. B3.20	12.62 15.82	13.69 13.69	a a	Tenopt Timoptol	SI FR
1925H	Eye drops (gellan gum solution) 2.5 mg (base) per mL (0.25%), 2.5 mL	‡1	5	..	11.80	12.87		Timoptol XE	MK
1926J	Eye drops (gellan gum solution) 5 mg (base) per mL (0.5%), 2.5 mL	‡1	5	..	12.62	13.69		Timoptol XE	MK
8803H	Eye gel 1 mg (base) per g (0.1%), 5 g	‡1	5	..	13.22	14.29		Nyogel	NV

Prostaglandin analogues

BIMATOPROST								
8620Q	Eye drops 300 micrograms per mL (0.03%), 3 mL	‡1	5	..	42.14	34.20	Lumigan	AG
BIMATOPROST with TIMOLOL MALEATE								
<u>Restricted benefit</u>								
Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;								
Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.								
9464D	Eye drops 300 micrograms-5 mg (base) per mL (0.03%-0.5%), 3 mL	‡1	5	..	46.82	34.20	Ganfort 0.3/5	AG
LATANOPROST								
8243W	Eye drops 50 micrograms per mL (0.005%), 2.5 mL	‡1	5	..	42.14	34.20	Xalatan	PF
LATANOPROST with TIMOLOL MALEATE								
<u>Restricted benefit</u>								
Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;								
Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.								
8895E	Eye drops 50 micrograms-5 mg (base) per mL (0.005%-0.5%), 2.5 mL	‡1	5	..	46.82	34.20	Xalacom	PF
TRAVOPROST								
8597L	Eye drops 40 micrograms per mL (0.004%), 2.5 mL	‡1	5	..	42.14	34.20	Travatan	AQ
TRAVOPROST with TIMOLOL MALEATE								
<u>Restricted benefit</u>								
Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;								
Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.								
9057Q	Eye drops 40 micrograms-5 mg (base) per mL (0.004%-0.5%), 2.5 mL	‡1	5	..	46.82	34.20	Duotrav	AQ

Mydriatics and cycloplegics

Anticholinergics

ATROPINE SULFATE								
1093M NP	Eye drops 10 mg per mL (1%), 15 mL	‡1	2	..	21.77	22.84	Atropt	SI
HOMATROPINE HYDROBROMIDE								
2541R NP	Eye drops 20 mg per mL (2%), 15 mL	‡1	2	..	17.97	19.04	Isopto Homatropine	AQ

Sensory organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
Decongestants and antiallergics								
Other antiallergics								
SODIUM CROMOGLYCATE								
Restricted benefit								
Vernal kerato-conjunctivitis.								
1127H	Eye drops 20 mg per mL (2%), 10 mL	1	5	..	14.37	15.44	a	Cromolux
NP							a	Opticrom
								SW

Ocular vascular disorder agents

Antineovascularisation agents

RANIBIZUMAB**Authority required**

Initial treatment by an ophthalmologist, as the sole PBS-subsidised therapy, of subfoveal choroidal neovascularisation (CNV) due to age-related macular degeneration (AMD), as diagnosed by fluorescein angiography.

Where a fluorescein angiogram cannot be performed due to a contraindication as listed in the TGA-approved product information, details of the contraindication must be provided. A copy of the report of an alternative method of diagnosis must be included in the application, for example, optical coherence tomography (OCT) or red free photography.

Authority approvals will be administered by the PBS and Specialised Drugs Branch of Medicare Australia.

The first authority application for each eye must be made in writing, and must include:

- (a) a completed authority prescription form; and
- (b) a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form [www.medicareaustralia.gov.au]; and
- (c) a copy of the fluorescein angiogram or alternative method of diagnosis where applicable.

Written applications for authority to prescribe ranibizumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Alternatively, the first authority application may be faxed to Medicare Australia on (03) 6215 5474 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Medicare Australia will then contact the prescriber by telephone. The original documentation must be posted to the above address after approval has been gained.

Authority required

Continuing treatment by an ophthalmologist, as the sole PBS-subsidised therapy, of subfoveal choroidal neovascularisation (CNV) due to age-related macular degeneration (AMD) where the patient has previously been granted an authority prescription for the same eye.

Authority approvals will be administered by the PBS and Specialised Drugs Branch of Medicare Australia. Authority applications for continuing treatment in the same eye may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

1382R	Solution for intravitreal injection 2.3 mg in 0.23 mL	1	2	..	1976.36	34.20		Lucentis NV

VERTEPORFIN**Authority required**

Initial treatment by an ophthalmologist, as the sole PBS-subsidised therapy, of predominantly (greater than or equal to 50%) classic, subfoveal choroidal neovascularisation (CNV) due to age-related macular degeneration (AMD), as diagnosed by fluorescein angiography, in a patient with a baseline visual acuity equal to or better than 6/60 (20/200).

Authority approvals will be administered by the PBS and Specialised Drugs Branch of Medicare Australia.

The first authority application for each eye must be made in writing, and must include:

- (a) a completed authority prescription form; and
- (b) a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form [www.medicareaustralia.gov.au]; and
- (c) a copy of the fluorescein angiogram demonstrating that the CNV is predominantly classic (greater than or equal to 50%).

Written applications for authority to prescribe verteporfin should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs

Sensory organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
	Reply Paid 9826 GPO Box 9826 HOBART TAS 7001						

Alternatively, the first authority application may be faxed to Medicare Australia on (03) 6215 5474 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Medicare Australia will then contact the prescriber by telephone. The original documentation must be posted to the above address after approval has been gained.

No more than 15 treatments (1 initial and 14 continuing) per eye will be authorised.

Medicare Australia should be notified if treatment is abandoned prior to completion of the laser activation step but after infusion of verteporfin. Telephone 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). The reason treatment is abandoned must be provided. Where such notification has been made, the treatment so affected will not count towards the maximum.

Authority required

Initial PBS-subsidised treatment by an ophthalmologist, as the sole PBS-subsidised therapy, of predominantly (greater than or equal to 50%) classic, subfoveal choroidal neovascularisation (CNV) due to macular degeneration where the patient has been authorised by the Angiogram Review Panel to receive treatment with verteporfin in the same eye under the MBS Visudyne Therapy Program.

Authority approvals will be administered by the PBS and Specialised Drugs Branch of Medicare Australia.

The first authority application for each eye must be made in writing, and must include:

- (a) a completed authority prescription form; and
- (b) a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form [www.medicareaustralia.gov.au], which includes the date of review by the Angiogram Review Panel and the number of treatments administered in that eye under the MBS Visudyne Therapy Program; and
- (c) a copy of the fluorescein angiogram demonstrating that the CNV is predominantly classic (greater than or equal to 50%).

Written applications for authority to prescribe verteporfin should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Alternatively, the first authority application may be faxed to Medicare Australia on (03) 6215 5474 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Medicare Australia will then contact the prescriber by telephone. The original documentation must be posted to the above address after approval has been gained.

A patient is eligible for a total of 15 subsidised treatments per eye. This maximum includes treatments administered under the MBS Visudyne Therapy Program and the PBS.

Medicare Australia should be notified if treatment is abandoned prior to completion of the laser activation step but after infusion of verteporfin. Telephone 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). The reason treatment is abandoned must be provided. Where such notification has been made, the treatment so affected will not count towards the maximum.

Authority required

Continuing treatment by an ophthalmologist, as the sole PBS-subsidised therapy, of predominantly (greater than or equal to 50%) classic, subfoveal choroidal neovascularisation (CNV) due to macular degeneration where the patient has previously been granted an authority prescription for the same eye.

A patient is eligible for a total of 15 subsidised treatments per eye. This maximum includes treatments administered under the MBS Visudyne Therapy Program and the PBS.

Medicare Australia should be notified if treatment is abandoned prior to completion of the laser activation step but after infusion of verteporfin. Telephone 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). The reason treatment is abandoned must be provided. Where such notification has been made, the treatment so affected will not count towards the maximum.

Authority approvals will be administered by the PBS and Specialised Drugs Branch of Medicare Australia. Authority applications for continuing treatment in the same eye may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

1349B	Powder for I.V. infusion 15 mg	1	..	..	2246.36	34.20	Visudyne	NV

Other ophthalmologicals

Other ophthalmologicals

CARBOMER

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome.

Sensory organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8384G NP	Eye gel 2 mg per g (0.2%), 10 g	1	5	..	10.35	11.42	GelTears	BU
							^a PAA	NM
				^B 0.96	11.31	11.42	^a Viscotears	NV

CARBOMER

Restricted benefit

For use in patients who have severe dry eye syndrome, including Sjogren's syndrome, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9210R	Eye gel 2 mg per g (0.2%), 10 g	1	11	..	10.35	11.42	GelTears	BU
							^a PAA	NM
				^B 0.96	11.31	11.42	^a Viscotears	NV

CARBOMER

Authority required (STREAMLINED)

1359

Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.

8578L NP	Eye gel 2 mg per g (0.2%), single dose units 0.6 mL, 30	3	5	..	*36.72	34.20	Viscotears Gel PF	NV

CARBOMER 974

Authority required (STREAMLINED)

1359

Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.

8514D NP	Ocular lubricating gel 3 mg per g (0.3%), single dose units 0.5 g, 30	3	5	..	*36.06	34.20	Poly Gel	AQ

CARMELLOSE SODIUM

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome.

8548X NP	Eye drops 5 mg per mL (0.5%), 15 mL	1	5	..	10.59	11.66	Refresh Tears Plus	AG
8593G NP	Eye drops 10 mg per mL (1%), 15 mL	1	5	..	10.59	11.66	Refresh Liquigel	AG

CARMELLOSE SODIUM

Restricted benefit

For use in patients who have severe dry eye syndrome, including Sjogren's syndrome, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9211T	Eye drops 5 mg per mL (0.5%), 15 mL	1	11	..	10.59	11.66	Refresh Tears Plus	AG
9212W	Eye drops 10 mg per mL (1%), 15 mL	1	11	..	10.59	11.66	Refresh Liquigel	AG

CARMELLOSE SODIUM

Authority required (STREAMLINED)

1359

Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.

2324H NP	Eye drops 10 mg per mL (1%), single dose units 0.4 mL, 30	3	5	..	*36.06	34.20	Celluvisc	AG

Sensory organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
2338C NP	Eye drops 5 mg per mL (0.5%), single dose units 0.4 mL, 30	3	5	..	*36.06	34.20	Cellufresh	AG
8823J NP	Eye drops 2.5 mg per mL (0.25%), single dose units 0.6 mL, 24	4	5	..	*40.42	34.20	TheraTears	CX
8824K NP	Ocular lubricating gel 10 mg per mL (1%), single dose units 0.6 mL, 28	3	5	..	*34.08	34.20	TheraTears	CX

CARMELLOSE SODIUM with GLYCERIN

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome.

Note

The in-use shelf life of Optive is 6 months from the date of opening.

9355J NP	Eye drops 5 mg-9 mg per mL (0.5%-0.9%), 15 mL	£1	3	..	10.59	11.66	Optive	AG

CARMELLOSE SODIUM with GLYCERIN

Restricted benefit

For use in patients who have severe dry eye syndrome, including Sjogren's syndrome, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

The in-use shelf life of Optive is 6 months from the date of opening.

9356K	Eye drops 5 mg-9 mg per mL (0.5%-0.9%), 15 mL	£1	7	..	10.59	11.66	Optive	AG

CARMELLOSE SODIUM with GLYCERIN

Authority required (STREAMLINED)

1359

Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.

9307W NP	Eye drops 5 mg-9 mg per mL (0.5%-0.9%), single dose units 0.4 mL, 30	3	5	..	*36.06	34.20	Optive	AG

HYPROMELLOSE

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome.

2956N NP	Eye drops 5 mg per mL (0.5%), 15 mL	£1	5	..	10.35	11.42	Methopt	SI
8287E NP	Eye drops 3 mg per mL (0.3%), 15 mL (contains sodium perborate as preservative)	£1	5	..	10.35	11.42 ^a	In a Wink Moisturising	NM
				^B 1.79	12.14	11.42 ^a	Genteal	NV

HYPROMELLOSE

Restricted benefit

For use in patients who have severe dry eye syndrome, including Sjogren's syndrome, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9213X	Eye drops 3 mg per mL (0.3%), 15 mL (contains sodium perborate as preservative)	£1	11	..	10.35	11.42 ^a	In a Wink Moisturising	NM
				^B 1.79	12.14	11.42 ^a	Genteal	NV
9214Y	Eye drops 5 mg per mL (0.5%), 15 mL	£1	11	..	10.35	11.42	Methopt	SI

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer	
HYPROMELLOSE with CARBOMER 980									
<u>Restricted benefit</u>									
Severe dry eye syndrome, including Sjogren's syndrome.									
8564R NP	Ocular lubricating gel 3 mg-2 mg per g (0.3%-0.2%), 10 g	‡1	5	..	10.35	11.42	^a	HPMC PAA	NM
				^B 1.79	12.14	11.42	^a	Genteal gel	NV
HYPROMELLOSE with CARBOMER 980									
<u>Restricted benefit</u>									
For use in patients who have severe dry eye syndrome, including Sjogren's syndrome, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.									
<u>Note</u>									
No applications for increased maximum quantities and/or repeats will be authorised.									
9215B	Ocular lubricating gel 3 mg-2 mg per g (0.3%-0.2%), 10 g	‡1	11	..	10.35	11.42	^a	HPMC PAA	NM
				^B 1.79	12.14	11.42	^a	Genteal gel	NV
HYPROMELLOSE with DEXTRAN									
<u>Restricted benefit</u>									
Severe dry eye syndrome, including Sjogren's syndrome.									
1509K NP	Eye drops 3 mg-1 mg per mL (0.3%-0.1%), 15 mL	‡1	5	..	10.57	11.64	^a	Poly-Tears	IQ
				^B 1.77	12.34	11.64	^a	Tears Naturale	AQ
HYPROMELLOSE with DEXTRAN									
<u>Restricted benefit</u>									
For use in patients who have severe dry eye syndrome, including Sjogren's syndrome, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.									
<u>Note</u>									
No applications for increased maximum quantities and/or repeats will be authorised.									
9216C	Eye drops 3 mg-1 mg per mL (0.3%-0.1%), 15 mL	‡1	11	..	10.57	11.64	^a	Poly-Tears	IQ
				^B 1.77	12.34	11.64	^a	Tears Naturale	AQ
HYPROMELLOSE with DEXTRAN									
<u>Authority required (STREAMLINED)</u>									
1359									
Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.									
8299T NP	Eye drops 3 mg-1 mg per mL (0.3%-0.1%), single dose units 0.4 mL, 28	3	5	..	*35.64	34.20		Bion Tears	AQ
PARAFFIN									
1750D NP	Pack containing 2 tubes compound eye ointment 3.5 g	‡1	5	..	20.89	21.96		Poly Visc	IQ
				^B 2.16	23.05	21.96	^a	Ircal	PE
							^a	Lacri-Lube	AG
1754H NP	Compound eye ointment 3.5 g	2	5	..	*21.54	22.61	^a	Poly Visc	IQ
				^B 2.16	*23.70	22.61	^a	Duratears	AQ
PARAFFIN									
<u>Restricted benefit</u>									
For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.									

Sensory organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer		
Note									
No applications for increased maximum quantities and/or repeats will be authorised.									
9217D	Compound eye ointment 3.5 g	2	11	.. ^B 2.16	*21.54 *23.70	22.61 22.61	^a ^a	Poly Visc Duratears	IQ AQ
9218E	Pack containing 2 tubes compound eye ointment 3.5 g	1	11	.. ^B 2.16	20.89 23.05	21.96 21.96	 ^a ^a	Poly Visc Ircal Lacri-Lube	IQ PE AG

POLYETHYLENE GLYCOL 400

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome.

Note

The in-use shelf life of Blink Intensive Tears multi-dose formulation is 45 days from the date of opening.

9491M NP	Eye drops 2.5 mg per mL (0.25%), 15 mL	1	5	..	10.59	11.66		Blink Intensive Tears	AO

POLYETHYLENE GLYCOL 400

Restricted benefit

For use in patients who have severe dry eye syndrome, including Sjogren's syndrome, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Note

The in-use shelf life of Blink Intensive Tears multi-dose formulation is 45 days from the date of opening.

9492N	Eye drops 2.5 mg per mL (0.25%), 15 mL	1	11	..	10.59	11.66		Blink Intensive Tears	AO

POLYETHYLENE GLYCOL 400

Authority required (STREAMLINED)

1359

Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.

9493P NP	Eye drops 2.5 mg per mL (0.25%), single dose units 0.4 mL, 20	5	5	..	*39.37	34.20		Blink Intensive Tears	AO

POLYETHYLENE GLYCOL 400 with PROPYLENE GLYCOL

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome.

8676P NP	Eye drops 4 mg-3 mg per mL (0.4%-0.3%), 15 mL	1	5	..	10.59	11.66		Systane	AQ

POLYETHYLENE GLYCOL 400 with PROPYLENE GLYCOL

Restricted benefit

For use in patients who have severe dry eye syndrome, including Sjogren's syndrome, and who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9219F	Eye drops 4 mg-3 mg per mL (0.4%-0.3%), 15 mL	1	11	..	10.59	11.66		Systane	AQ

Otologicals

Antiinfectives

Antiinfectives

CHLORAMPHENICOL

1172Q	Ear drops (aqueous) 5 mg per mL (0.5%), 5 mL	#1	2	..	11.05	12.12	Chloromycetin	PF

CIPROFLOXACIN

Authority required

Treatment of chronic suppurative otitis media in an Aboriginal or a Torres Strait Islander person aged 1 month or older;

Sensory organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
	Treatment of chronic suppurative otitis media in a patient less than 18 years of age with perforation of the tympanic membrane; Treatment of chronic suppurative otitis media in a patient less than 18 years of age with a grommet in situ.						
2480M NP	Ear drops 3 mg per mL (0.3%), 5 mL	1	1	..	19.28	20.35	Ciloxan AQ
NEOMYCIN UNDECENOATE with BACITRACIN ZINC							
2296W NP	Ear ointment 12 mg (3.5 mg base)-400 units per g, 10 g	1	..	..	8.66	9.73	Nemdyn HA
Corticosteroids and antiinfectives in combination Corticosteroids and antiinfectives in combination							
DEXAMETHASONE with FRAMYCETIN SULFATE and GRAMICIDIN							
2781J NP	Ear drops 500 micrograms-5 mg-50 micrograms per mL, 8 mL	1	2	.. B1.91	9.44 11.35	10.51 10.51	a a Otodex Sofradex AV SW
TRIAMCINOLONE ACETONIDE with NEOMYCIN SULFATE, GRAMICIDIN and NYSTATIN							
2971J NP	Ear drops 1 mg-2.5 mg (base)- 250 micrograms-100,000 units per g (0.1%-0.25%-0.025%-100,000 units per g), 7.5 mL	1	2	.. B2.00	11.18 13.18	12.25 12.25	a a Otocomb Otic Kenacomb Otic SI FM
2974M NP	Ear ointment 1 mg-2.5 mg (base)- 250 micrograms-100,000 units per g (0.1%-0.25%-0.025%-100,000 units per g), 5 g	1	2	.. B2.01	8.21 10.22	9.28 9.28	a a Otocomb Otic Kenacomb Otic SI FM

Ophthalmological and otological preparations

Antiinfectives

Antiinfectives

FRAMYCETIN SULFATE								
1440T NP,MW	Eye and ear drops 5 mg per mL (0.5%), 8 mL	1	2	..	10.11	11.18	Soframycin	SW

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer

Various

Allergens

Allergens

Allergen extracts

INSECT ALLERGEN EXTRACT—HONEY BEE VENOM							
2886X	Injection set containing 550 micrograms	1	..	..	238.38	34.20	Albey Bee Venom HL
INSECT ALLERGEN EXTRACT—PAPER WASP VENOM							
Note							
Paper wasp venom is not European wasp venom.							
2918N	Injection set containing 550 micrograms	1	..	..	238.38	34.20	Albey Paper Wasp Venom HL
INSECT ALLERGEN EXTRACT—YELLOW JACKET VENOM							
2883R	Injection set containing 550 micrograms	1	..	..	238.38	34.20	Albey Yellow Jacket Venom HL

All other therapeutic products

All other therapeutic products

Antidotes

NALOXONE HYDROCHLORIDE							
1753G NP	Injection 2 mg in 5 mL	1	..	..	43.49	34.20	Naloxone Min-I-Jet CS

Drugs for treatment of hyperkalemia and hyperphosphatemia

LANTHANUM

Authority required (STREAMLINED)

3546

Maintenance therapy, following initiation and stabilisation of treatment with lanthanum carbonate, of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where serum phosphate is greater than 1.6 mmol per L at the commencement of therapy;

3547

Maintenance therapy, following initiation and stabilisation of treatment with lanthanum carbonate, of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where the serum calcium times phosphate product is greater than 4.0 at the commencement of therapy.

Note

Not to be used in combination with sevelamer.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

9403X NP	Tablet, chewable, 500 mg (as carbonate hydrate)	90	5	..	305.87	34.20	Fosrenol	ZI
9404Y NP	Tablet, chewable, 750 mg (as carbonate hydrate)	90	5	..	449.42	34.20	Fosrenol	ZI
9405B NP	Tablet, chewable, 1000 mg (as carbonate hydrate)	90	5	..	504.03	34.20	Fosrenol	ZI

SEVELAMER HYDROCHLORIDE

Authority required (STREAMLINED)

3548

Maintenance therapy, following initiation and stabilisation of treatment with sevelamer hydrochloride, of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where serum phosphate is greater than 1.6 mmol per L at the commencement of therapy;

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
3549							
Maintenance therapy, following initiation and stabilisation of treatment with sevelamer hydrochloride, of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where the serum calcium times phosphate product is greater than 4.0 at the commencement of therapy.							
<u>Note</u>							
Not to be used in combination with lanthanum.							
<u>Note</u>							
Shared Care Model:							
For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
2142R NP	Tablet 800 mg	180	5	..	357.73	34.20	Renagel GZ

Detoxifying agents for antineoplastic treatment

CALCIUM FOLINATE							
8740B NP	Injection equivalent to 50 mg folinic acid in 5 mL	5	5	..	*148.92	34.20 ^a	Leucovorin Calcium (Hospira Pty Limited) HH
					*148.94	34.20 ^a	Leucovorin Calcium (Pfizer Australia Pty Ltd) PF
					148.95	34.20 ^a	Calcium Folate Ebewe IT
8812T NP	Injection equivalent to 100 mg folinic acid in 10 mL	10	1	..	*263.52	34.20 ^a	Calcium Folate Ebewe IT
					263.56	34.20 ^a	Leucovorin Calcium (Pfizer Australia Pty Ltd) PF
9041W NP	Injection equivalent to 300 mg folinic acid in 30 mL	4	1	..	*304.10	34.20	Leucovorin Calcium (Hospira Pty Limited) HH
CALCIUM FOLINATE							
Restricted benefit							
Antidote to folic acid antagonists.							
2308L NP	Tablet equivalent to 15 mg folinic acid	10	..	..	98.15	34.20	Leucovorin Calcium (Hospira Pty Limited) HH
MESNA							
Restricted benefit							
Adjunctive therapy for use with ifosfamide or high dose cyclophosphamide.							
8078E	Solution for I.V. injection 400 mg in 4 mL	15	5	..	103.28	34.20	Uromitexan BX
8079F	Solution for I.V. injection 1 g in 10 mL	15	5	..	223.81	34.20	Uromitexan BX

Drugs for treatment of hypercalcemia

SODIUM ACID PHOSPHATE							
Authority required (STREAMLINED)							
1099							
Familial hypophosphataemia;							
1157							
Hypercalcaemia;							
1167							
Hypophosphataemic rickets;							
1467							
Vitamin D-resistant rickets.							
2946C NP	Compound effervescent tablet containing elemental phosphorus 500 mg, sodium 469 mg (20.4 mmol), potassium 123 mg (3.1 mmol)	100	5	..	81.63	34.20	Phosphate Sandoz NV

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Other therapeutic products

POLY-L-LACTIC ACID

Note

Authority applications to prescribe poly-l-lactic acid may be made by telephone to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial PBS-subsidised treatment, for facial administration only, of severe facial lipoatrophy caused by therapy for HIV infection.

Accreditation following completion of injection administration training with Sanofi-Aventis is required to prescribe poly-l-lactic acid under the PBS. Patients must be referred from the HIV physician to the accredited injector.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9475Q	Powder for injection 150 mg	2	4	..	*446.46	34.20	Sculptra	SW

POLY-L-LACTIC ACID

Note

Authority applications to prescribe poly-l-lactic acid may be made by telephone to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Maintenance PBS-subsidised treatment, for facial administration only, of severe facial lipoatrophy caused by therapy for HIV infection.

Accreditation following completion of injection administration training with Sanofi-Aventis is required to prescribe poly-l-lactic acid under the PBS. Patients must be referred from the HIV physician to the accredited injector.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Maintenance treatment is limited to one re-treatment (maximum 2 vials) every 2 years.

9476R	Powder for injection 150 mg	2	..	..	*446.46	34.20	Sculptra	SW

Diagnostic agents

Urine tests

COPPER SULFATE

1228P NP	Diagnostic compound tablets, 36	2	3	..	*71.48	34.20	Clinitest	BN

COPPER SULFATE

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9251X	Diagnostic compound tablets, 36	2	6	..	*71.48	34.20	Clinitest	BN

GLUCOSE and KETONE INDICATOR—URINE

3106L NP	Test strips, 50	2	2	..	*17.30	18.37	Keto-Diabur- Test 5000	RD

3107M NP	Test strips, 50	2	2	..	*17.42	18.49	Keto-Diastix	BN

GLUCOSE and KETONE INDICATOR—URINE

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Note No applications for increased maximum quantities and/or repeats will be authorised.							
9254C	Test strips, 50	2	4	..	*17.30	18.37	Keto-Diabur- Test 5000 RD
9255D	Test strips, 50	2	4	..	*17.42	18.49	Keto-Diastix BN
GLUCOSE INDICATOR—URINE							
2352T NP	Test strips, 50	2	2	..	*19.82	20.89	Clinistix BN
3104J NP	Test strips, 50	2	2	..	*18.84	19.91	Diastix BN

GLUCOSE INDICATOR—URINE

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

9252Y	Test strips, 50	2	4	..	*19.82	20.89	Clinistix BN
9253B	Test strips, 50	2	4	..	*18.84	19.91	Diastix BN

Other diagnostic agents

Tests for diabetes

GLUCOSE INDICATOR—BLOOD

2263D NP	Test strips, 50	2	5	..	*53.18	34.20	Optium Omega MS
2860M NP	Test strips, 50	2	5	..	*53.18	34.20	Betachek G5 NA
2890D NP	Test strips, 50	2	5	..	*53.18	34.20	Betachek NA
2891E NP	Test strips, 50	2	5	..	*53.18	34.20	Advantage II RD
2914J NP	Test strips, 50	2	5	..	*45.90	34.20	Glucoflex-R NA
2917M NP	Test strips, 50	2	5	..	*45.90	34.20	GlucoStix BN
2979T NP	Test strips, 100	‡1	5	..	53.16	34.20	Accu-Chek Performa RD
3406G NP	Test strips, 50	2	5	..	*53.18	34.20	CareSens N LB
3411M NP	Test strips, 100	‡1	5	..	53.16	34.20	Accu-Chek Advantage/Sens or Comfort RD
3441D NP	Test strips, 50	2	5	..	*53.18	34.20	OneTouch Verio JJ
8190C NP	Test strips, 100	‡1	5	..	53.16	34.20	Accu-Chek Active RD
8522M NP	Test strips, 100	‡1	5	..	53.16	34.20	Optium glucose MS
8682Y NP	Test strips, 50	2	5	..	*53.18	34.20	MWD Pen Sensor Strips WF
8723D NP	Test strips, 50	2	5	..	*53.18	34.20	Omnitest EZ BR
8739Y NP	Test strips, 50	2	5	..	*53.18	34.20	Accu-Chek Go RD
8749L NP	Test strips, 50	2	5	..	*53.18	34.20	GlucoOz OZ
8759B NP	Test strips, 50	2	5	..	*53.18	34.20	CareSens LB
8795X NP	Test strips, 50	2	5	..	*53.18	34.20	SensoCard PX
8806L NP	Test strips, 51	2	5	..	*53.18	34.20	Accu-Chek Integra RD

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8825L NP	Test strips, 50	2	5	..	*53.18	34.20	TrueTrack	NX
8890X NP	Test strips, 50	2	5	..	*53.18	34.20	Freestyle Papillon	MS
9013J NP	Test strips, 50	2	5	..	*53.18	34.20	Glucocard 01 Sensor	OZ
9154T NP	Test strips, 100	‡1	5	..	53.16	34.20	FreeStyle Lite	MS
9193W NP	Test strips, 25	4	5	..	*53.18	34.20	On-Call Plus	PZ
9298J NP	Test strips, 50	2	5	..	*53.18	34.20	Bionime Rightest	QB
9324R NP	Test strips, 50	2	5	..	*53.18	34.20	WaveSense Jazz	HE
9471L NP	Test strips, 50	2	5	..	*53.18	34.20	MyGlucoHealth	EH
9485F NP	Test strips, 50	2	5	..	*53.18	34.20	Lifeline Attest	OI

GLUCOSE INDICATOR—BLOOD

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

3407H	Test strips, 50	2	11	..	*53.18	34.20	CareSens N	LB
3412N	Test strips, 100	‡1	11	..	53.16	34.20	Accu-Chek Advantage/Sens or Comfort	RD
3442E	Test strips, 50	2	11	..	*53.18	34.20	OneTouch Verio	JJ
9256E	Test strips, 25	4	11	..	*53.18	34.20	On-Call Plus	PZ
9257F	Test strips, 100	‡1	11	..	53.16	34.20	Accu-Chek Performa	RD
9258G	Test strips, 50	2	11	..	*53.18	34.20	Advantage II	RD
9259H	Test strips, 50	2	11	..	*53.18	34.20	Freestyle Papillon	MS
9261K	Test strips, 50	2	11	..	*53.18	34.20	Glucocard 01 Sensor	OZ
9263M	Test strips, 50	2	11	..	*53.18	34.20	GlucoOz	OZ
9264N	Test strips, 50	2	11	..	*53.18	34.20	MWD Pen Sensor Strips	WF
9265P	Test strips, 50	2	11	..	*53.18	34.20	Omnitest EZ	BR
9267R	Test strips, 50	2	11	..	*53.18	34.20	Optium Omega	MS
9268T	Test strips, 50	2	11	..	*53.18	34.20	TrueTrack	NX
9269W	Test strips, 100	‡1	11	..	53.16	34.20	FreeStyle Lite	MS
9270X	Test strips, 100	‡1	11	..	53.16	34.20	Optium glucose	MS
9273C	Test strips, 100	‡1	11	..	53.16	34.20	Accu-Chek Active	RD
9274D	Test strips, 50	2	11	..	*53.18	34.20	Accu-Chek Go	RD
9275E	Test strips, 51	2	11	..	*53.18	34.20	Accu-Chek Integra	RD
9276F	Test strips, 50	2	11	..	*53.18	34.20	Betachek	NA
9277G	Test strips, 50	2	11	..	*53.18	34.20	Betachek G5	NA
9278H	Test strips, 50	2	11	..	*53.18	34.20	CareSens	LB
9279J	Test strips, 50	2	11	..	*45.90	34.20	Glucoflex-R	NA
9280K	Test strips, 50	2	11	..	*45.90	34.20	Glucostix	BN
9281L	Test strips, 50	2	11	..	*53.18	34.20	SensoCard	PX
9297H	Test strips, 50	2	11	..	*53.18	34.20	Bionime Rightest	QB
9325T	Test strips, 50	2	11	..	*53.18	34.20	WaveSense Jazz	HE
9472M	Test strips, 50	2	11	..	*53.18	34.20	MyGlucoHealth	EH
9486G	Test strips, 50	2	11	..	*53.18	34.20	Lifeline Attest	OI

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
GLUCOSE INDICATOR—BLOOD							
<u>Restricted benefit</u>							
For use in patients on insulin therapy.							
9300L NP	Test strips, 100	1	5	..	53.16	34.20	Accu-Chek Mobile RD
GLUCOSE INDICATOR—BLOOD							
<u>Restricted benefit</u>							
For use in patients on insulin therapy who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.							
<u>Note</u>							
No applications for increased maximum quantities and/or repeats will be authorised.							
9301M	Test strips, 100	1	11	..	53.16	34.20	Accu-Chek Mobile RD

General nutrients

Other nutrients

TRIGLYCERIDES, MEDIUM CHAIN

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Authority required

Chylous ascites;

Chyllothorax;

Fat malabsorption due to liver disease, short gut syndrome, cystic fibrosis and gastrointestinal disorders;

Hyperlipoproteinaemia type 1;

Intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect, requiring a ketogenic diet;

Long chain fatty acid oxidation disorders.

3128P NP	Oil 500 mL	2	5	..	*52.38	34.20	MCT Oil	SB
9327X NP	Emulsion 250 mL	8	5	..	*214.42	34.20	Liquigen	SB

Fat/carbohydrates/proteins/minerals/vitamins, combinations

AMINO ACIDS—SYNTHETIC, FORMULA

Authority required

Initial treatment for up to 3 months, by a clinical immunologist, suitably qualified allergist or gastroenterologist in a patient 18 years of age or less with eosinophilic oesophagitis who requires an amino acid based formula as a component of a dietary elimination programme. Treatment with oral steroids should not be commenced during the period of initial treatment.

Eosinophilic oesophagitis is demonstrated by the following criteria:

- (i) Chronic symptoms of reflux that persisted despite a 2-month trial of a proton pump inhibitor or chronic dysphagia; and
- (ii) A lack of demonstrable anatomic abnormality with the exception of stricture, which can be attributable to eosinophilic oesophagitis; and
- (iii) Eosinophilic infiltration of the oesophagus, demonstrated by oesophageal biopsy specimens obtained by endoscopy and where the most densely involved oesophageal biopsy had 20 or more eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies.

The date of birth of the patient must be included in the authority.

Authority required

Continuing treatment by a clinical immunologist, suitably qualified allergist or gastroenterologist in a patient 18 years of age or less with eosinophilic oesophagitis who has responded to an initial course of PBS-subsidised treatment. Response to initial treatment is demonstrated by oesophageal biopsy specimens obtained by endoscopy, where the most densely involved oesophageal biopsy had 5 or less eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies. The response criteria will not be deemed to have been met if oral steroids were commenced during initial treatment.

Note

Authorities for increased maximum quantities, up to a maximum of 52, may be authorised.

2250K NP	Compound powder 400 g	12	5	..	*531.66	34.20	EleCare	AB

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

AMINO ACIDS—SYNTHETIC, FORMULA

Authority required

Initial treatment, for up to 3 months, for combined intolerance (not infant colic) to cows' milk protein, soy protein and protein hydrolysate formulae in a child up to the age of 2 years. Combined intolerance is demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula. The date of birth of the patient must be included in the authority application;

Initial treatment, in consultation with a paediatric gastroenterologist or specialist allergist, for up to 3 months, of a child up to the age of 2 years with severe intolerance (not infant colic) to cows' milk protein. The date of birth of the patient must be included in the authority application.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

2244D NP	Compound powder 400 g	8	5	..	*361.14	34.20	Neocate Advance Tropical Flavour	SB
8443J NP	Compound powder 400 g	8	5	..	*361.14	34.20	Neocate	SB
8574G NP	Compound powder 400 g	8	5	..	*361.14	34.20	EleCare	AB
8754R NP	Compound powder 400 g	8	5	..	*361.14	34.20	Neocate Advance	SB

AMINO ACIDS—SYNTHETIC, FORMULA

Authority required

Continuing treatment for combined intolerance (not infant colic) to cows' milk protein, soy protein and protein hydrolysate formulae in a child up to the age of 2 years, where the child has been assessed by a suitably qualified allergist or paediatrician. The date of birth of the patient must be included in the authority application;

Treatment for combined intolerance (not infant colic) to cows' milk protein, soy protein and protein hydrolysate formulae in a child aged 2 years and over, where the child is assessed by a suitably qualified allergist or paediatrician at intervals not greater than 6 months. The date of birth of the patient must be included in the authority application;

Continuing treatment for severe intolerance (not infant colic) to cows' milk protein in a child up to the age of 2 years, where the child has been assessed by a paediatric gastroenterologist or specialist allergist and soy protein and protein hydrolysate formulae are not tolerated or not likely to be tolerated. The date of birth of the patient must be included in the authority application;

Treatment for severe intolerance (not infant colic) to cows' milk protein in a child aged 2 years and over, where the child is assessed by a paediatric gastroenterologist or specialist allergist at intervals not greater than 6 months. The date of birth of the patient must be included in the authority application;

Severe intestinal malabsorption including short bowel syndrome where protein hydrolysate formulae have failed;

Severe intestinal malabsorption including short bowel syndrome where the patient has been receiving parenteral nutrition.

Note

Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

2553J NP	Compound powder 400 g	8	5	..	*361.14	34.20	Neocate Advance Tropical Flavour	SB
3066J NP	Compound powder 400 g	8	5	..	*361.14	34.20	Neocate	SB
8575H NP	Compound powder 400 g	8	5	..	*361.14	34.20	EleCare	AB
8755T NP	Compound powder 400 g	8	5	..	*361.14	34.20	Neocate Advance	SB

AMINO ACID SYNTHETIC FORMULA supplemented with LONG CHAIN POLYUNSATURATED FATTY ACIDS

Authority required

Initial treatment, for up to 3 months, for combined intolerance (not infant colic) to cows' milk protein, soy protein and protein hydrolysate formulae in a child up to the age of 2 years. Combined intolerance is demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula. The date of birth of the patient must be included in the authority application;

Initial treatment, in consultation with a paediatric gastroenterologist or specialist allergist, for up to 3 months, of a child up to the age of 2 years with severe intolerance (not infant colic) to cows' milk protein. The date of birth of the patient must be included in the authority application.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

2246F NP	Compound powder 400 g	8	5	..	*367.86	34.20	Neocate LCP	SB

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
9339M NP	Compound powder 400 g	8	5	..	*367.86	34.20	EleCare LCP AB

AMINO ACID SYNTHETIC FORMULA supplemented with LONG CHAIN POLYUNSATURATED FATTY ACIDS

Authority required

Continuing treatment for combined intolerance (not infant colic) to cows' milk protein, soy protein and protein hydrolysate formulae in a child up to the age of 2 years, where the child has been assessed by a suitably qualified allergist or paediatrician. The date of birth of the patient must be included in the authority application;

Treatment for combined intolerance (not infant colic) to cows' milk protein, soy protein and protein hydrolysate formulae in a child aged 2 years and over, where the child is assessed by a suitably qualified allergist or paediatrician at intervals not greater than 6 months. The date of birth of the patient must be included in the authority application;

Continuing treatment for severe intolerance (not infant colic) to cows' milk protein in a child up to the age of 2 years, where the child has been assessed by a paediatric gastroenterologist or specialist allergist and soy protein and protein hydrolysate formulae are not tolerated or not likely to be tolerated. The date of birth of the patient must be included in the authority application;

Treatment for severe intolerance (not infant colic) to cows' milk protein in a child aged 2 years and over, where the child is assessed by a paediatric gastroenterologist or specialist allergist at intervals not greater than 6 months. The date of birth of the patient must be included in the authority application;

Severe intestinal malabsorption including short bowel syndrome where protein hydrolysate formulae have failed;

Severe intestinal malabsorption including short bowel syndrome where the patient has been receiving parenteral nutrition.

Note

Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

2560R NP	Compound powder 400 g	8	5	..	*367.86	34.20	Neocate LCP	SB
9340N NP	Compound powder 400 g	8	5	..	*367.86	34.20	EleCare LCP	AB

PROTEIN HYDROLYSATE FORMULA with MEDIUM CHAIN TRIGLYCERIDES

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Authority required

Initial treatment, for up to 3 months, for intolerance (not infant colic) to both cows' milk protein and soy protein in a child up to the age of 2 years. Intolerance is demonstrated when the child has failed to respond to a strict cows' milk protein free diet with a soy protein as the principal formula. The date of birth of the patient must be included in the authority application;

Continuing treatment for intolerance (not infant colic) to both cows' milk protein and soy protein in a child up to the age of 2 years, where clinical improvement has been demonstrated with the protein hydrolysate formula with medium chain triglycerides. The date of birth of the patient must be included in the authority application;

Continuing treatment for intolerance (not infant colic) to both cows' milk protein and soy protein in a child aged 2 years and over, where the child has been assessed by a suitably qualified allergist or paediatrician. The date of birth of the patient must be included in the authority application.

Authority required

Initial treatment, in consultation with a paediatric gastroenterologist or specialist allergist, for up to 3 months, of a child up to the age of 2 years with severe intolerance (not infant colic) to cows' milk protein. The date of birth of the patient must be included in the authority application;

Continuing treatment for severe intolerance (not infant colic) to cows' milk protein in a child up to the age of 2 years, where clinical improvement has been demonstrated with the protein hydrolysate formula with medium chain triglycerides and soy protein is not tolerated or is likely not to be tolerated. The date of birth of the patient must be included in the authority application;

Continuing treatment for severe intolerance (not infant colic) to cows' milk protein in a child aged 2 years and over, where the child has been assessed by a paediatric gastroenterologist or specialist allergist. The date of birth of the patient must be included in the authority application.

Authority required

Biliary atresia;

Chronic liver failure with fat malabsorption;

Chylous ascites;

Chylothorax;

Cystic fibrosis;

Enterokinase deficiency;

Proven fat malabsorption;

Severe diarrhoea of greater than 2 weeks' duration in an infant aged less than 4 months. The date of birth of the patient must be included in the authority application;

Severe intestinal malabsorption including short bowel syndrome.

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
2676W NP	Compound powder 400 g	8	5	..	*171.94	34.20	Alfaré NT

PROTEIN HYDROLYSATE FORMULA with MEDIUM CHAIN TRIGLYCERIDES

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Authority required

Initial treatment, for up to 3 months, for intolerance (not infant colic) to both cows' milk protein and soy protein in a child up to the age of 2 years. Intolerance is demonstrated when the child has failed to respond to a strict cows' milk protein free diet with a soy protein as the principal formula. The date of birth of the patient must be included in the authority application;

Continuing treatment for intolerance (not infant colic) to both cows' milk protein and soy protein in a child up to the age of 2 years, where clinical improvement has been demonstrated with the protein hydrolysate formula with medium chain triglycerides. The date of birth of the patient must be included in the authority application;

Continuing treatment for intolerance (not infant colic) to both cows' milk protein and soy protein in a child aged 2 years and over, where the child has been assessed by a suitably qualified allergist or paediatrician. The date of birth of the patient must be included in the authority application.

Authority required

Initial treatment, in consultation with a paediatric gastroenterologist or specialist allergist, for up to 3 months, of a child up to the age of 2 years with severe intolerance (not infant colic) to cows' milk protein. The date of birth of the patient must be included in the authority application;

Continuing treatment for severe intolerance (not infant colic) to cows' milk protein in a child up to the age of 2 years, where clinical improvement has been demonstrated with the protein hydrolysate formula with medium chain triglycerides and soy protein is not tolerated or is likely not to be tolerated. The date of birth of the patient must be included in the authority application;

Continuing treatment for severe intolerance (not infant colic) to cows' milk protein in a child aged 2 years and over, where the child has been assessed by a paediatric gastroenterologist or specialist allergist. The date of birth of the patient must be included in the authority application.

Authority required

Biliary atresia;

Chronic liver failure with fat malabsorption;

Chylous ascites;

Cystic fibrosis;

Enterokinase deficiency;

Proven fat malabsorption;

Severe diarrhoea of greater than 2 weeks' duration in an infant aged less than 4 months. The date of birth of the patient must be included in the authority application;

Severe intestinal malabsorption including short bowel syndrome.

8259Q NP	Compound powder 450 g	8	5	..	*109.86	34.20	Pepti-Junior Gold NU

TRIGLYCERIDES—MEDIUM CHAIN, FORMULA

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Restricted benefit

Chylous ascites;

Chylothorax;

Fat malabsorption due to liver disease, short gut syndrome, cystic fibrosis and gastrointestinal disorders;

Hyperlipoproteinaemia type 1;

Long chain fatty acid oxidation disorders.

Note

Monogen is not indicated for the treatment of intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect requiring a ketogenic diet.

8478F NP	Compound powder 400 g	8	5	..	*421.30	34.20	Monogen SB

TRIGLYCERIDES—MEDIUM CHAIN, FORMULA

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
<u>Restricted benefit</u>							
Chylous ascites;							
Chylothorax;							
Fat malabsorption due to liver disease, short gut syndrome, cystic fibrosis and gastrointestinal disorders.							
<u>Note</u>							
Caprilon is not indicated for the treatment of intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect requiring a ketogenic diet, long chain fatty acid oxidation disorders or hyperlipoproteinaemia type 1.							
8629E NP	Compound powder 420 g	8	5	..	*467.46	34.20	Caprilon SB

Carbohydrates

AMYLOPECTIN, MODIFIED LONG CHAIN

Restricted benefit

Glycogen storage disease.

9386B NP	Sachets 60 g, 30	4	5	..	*752.30	34.20	Glycosade VF

Milk substitutes

MILK POWDER—LACTOSE FREE FORMULA

Authority required

Acute lactose intolerance in infants up to the age of 12 months. The date of birth of the patient must be included in the authority application.

Note

No applications for increased maximum quantities and/or repeats will be authorised. No more than 1 application per patient will be authorised.

2350Q NP	Lactose-predigested powder infant formula 900 g	5	..	..	*88.92	34.20	Karicare De-Lact NU
8282X NP	Infant formula powder 900 g	5	..	..	*112.87	34.20	S-26 LF WX

MILK POWDER—LACTOSE FREE FORMULA

Authority required

Proven chronic lactose intolerance in infants up to the age of 12 months. The date of birth of the patient must be included in the authority application. Lactose intolerance must have been proven by either:

- (a) relief of symptoms on supervised withdrawal of lactose from the diet for 3 or 4 days and subsequent re-emergence of symptoms on rechallenge with lactose containing formulae or milk or food; or
- (b) not less than 0.5% reducing substance in stool exudate tested with copper sulfate diagnostic compound tablet; or
- (c) hydrogen breath test.

Note

No applications for increased maximum quantities and/or repeats will be authorised.

2349P NP	Lactose-predigested powder infant formula 900 g	5	5	..	*88.92	34.20	Karicare De-Lact NU
8283Y NP	Infant formula powder 900 g	5	5	..	*112.87	34.20	S-26 LF WX

MILK POWDER—LACTOSE MODIFIED

Authority required

Acute lactose intolerance in children aged 1 year and over. The date of birth of the patient must be included in the authority application.

Note

No applications for increased maximum quantities and/or repeats will be authorised. No more than 1 application per patient will be authorised.

2358D NP	Lactose-predigested powder 900 g	3	1	..	*72.81	34.20	Digestelact SJ

MILK POWDER—LACTOSE MODIFIED

Authority required

Proven chronic lactose intolerance in children aged 1 year and over who are significantly malnourished. The date of birth of the patient must be included in the authority application. Lactose intolerance must have been proven by either:

- (a) relief of symptoms on supervised withdrawal of lactose from the diet for 3 or 4 days and subsequent re-emergence of symptoms on rechallenge with lactose containing formulae or milk or food; or

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty	Safety Net	
					\$	\$	
	(b) not less than 0.5% reducing substance in stool exudate tested with copper sulfate diagnostic compound tablet; or (c) hydrogen breath test.						

Note

No applications for increased maximum quantities and/or repeats will be authorised.

2357C NP	Lactose-predigested powder 900 g	3	10	..	*72.81	34.20	Digestelact	SJ

MILK POWDER—SYNTHETIC

Note

No applications for increased maximum quantities and/or repeats will be authorised.

Authority required

Hypercalcaemia in children under the age of 4 years.

3092R NP	Low calcium compound powder 400 g	8	5	..	*381.38	34.20	Locasol	SB

Other combinations of nutrients

AMINO ACID FORMULA with FAT, CARBOHYDRATE, VITAMINS, MINERALS, and TRACE ELEMENTS, without METHIONINE and supplemented with DOCOSAHEXANOIC ACID

Restricted benefit

Pyridoxine non-responsive homocystinuria.

3417W NP	Oral liquid 125 mL, 36	4	5	..	*2507.98	34.20	HCU Anamix junior LQ	SB

AMINO ACID FORMULA with FAT, CARBOHYDRATE, VITAMINS, MINERALS and TRACE ELEMENTS without PHENYLALANINE and TYROSINE, and supplemented with DOCOSAHEXANOIC ACID

Restricted benefit

Tyrosinaemia.

9330C NP	Oral liquid 125 mL, 36	4	5	..	*2507.98	34.20	TYR Anamix junior LQ	SB

AMINO ACID FORMULA without PHENYLALANINE

Restricted benefit

Phenylketonuria.

2347M NP	Sachets containing powder 20 g, 30	7	5	..	*1462.91	34.20	Phlexy-10 Drink Mix	SB
8554F NP	Capsules 500 mg, 200	16	5	..	*1276.34	34.20	Phlexy-10	SB
8678R NP	Tablets 1 g, 75	24	5	..	*1426.98	34.20	Phlexy-10	SB

AMINO ACID FORMULA with VITAMINS, MINERALS and LONG CHAIN POLYUNSATURATED FATTY ACIDS without PHENYLALANINE

Restricted benefit

Phenylketonuria.

8479G NP	Infant formula, powder 400 g	8	5	..	*703.62	34.20	PKU Anamix infant	SB

AMINO ACID FORMULA with VITAMINS and MINERALS without LYSINE and low in TRYPTOPHAN

Restricted benefit

A child aged from 6 months up to 10 years with proven glutaric aciduria type 1.

9438R NP	Sachets 20 g, 30	4	5	..	*1797.30	34.20	GA gel	VF

AMINO ACID FORMULA with VITAMINS and MINERALS without LYSINE and low in TRYPTOPHAN

Restricted benefit

An infant or young child with proven glutaric aciduria type 1.

2650L NP	Infant formula, powder 400 g	8	5	..	*769.30	34.20	GA1 Anamix infant	SB

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
AMINO ACID FORMULA with VITAMINS and MINERALS without LYSINE and low in TRYPTOPHAN								
<u>Restricted benefit</u>								
A child aged less than 9 years with proven glutaric aciduria type 1.								
2646G NP	Powder 500 g	8	5	..	*1784.74	34.20	XLYS, LOW TRY Maxamaid	SB
AMINO ACID FORMULA with VITAMINS and MINERALS without METHIONINE								
<u>Restricted benefit</u>								
For infants and very young children with pyridoxine non-responsive homocystinuria.								
8417B NP	Infant formula, powder 400 g	8	5	..	*769.30	34.20	HCU Anamix infant	SB
AMINO ACID FORMULA with VITAMINS and MINERALS without METHIONINE								
<u>Restricted benefit</u>								
Pyridoxine non-responsive homocystinuria.								
8328H NP	Powder 500 g	8	5	..	*1784.74	34.20	XMET Maxamaid	SB
8416Y NP	Powder 500 g	8	5	..	*2704.74	34.20	XMET Maxamum	SB
8677Q NP	Sachets 20 g, 30	4	5	..	*1798.42	34.20	HCU gel	VF
8744F NP	Sachets 25 g, 30	4	5	..	*3098.38	34.20	HCU express	VF
9133Q NP	Oral liquid 130 mL, 30	4	5	..	*3098.38	34.20	HCU Cooler	VF
AMINO ACID FORMULA with VITAMINS and MINERALS without METHIONINE, THREONINE and VALINE and low in ISOLEUCINE								
<u>Restricted benefit</u>								
Methylmalonic acidemia;								
Propionic acidemia.								
3443F NP	Sachets 25 g, 30	4	5	..	*3098.38	34.20	MMA/PA express	VF
3444G NP	Sachets 20 g, 30	4	5	..	*1798.42	34.20	MMA/PA gel	VF
8058D NP	Infant formula, powder 400 g	8	5	..	*769.30	34.20	MMA/PA Anamix infant	SB
8059E NP	Powder 500 g	8	5	..	*1784.74	34.20	XMTVI Maxamaid	SB
8061G NP	Powder 500 g	8	5	..	*2704.74	34.20	XMTVI Maxamum	SB
AMINO ACID FORMULA with VITAMINS and MINERALS without PHENYLALANINE								
<u>Restricted benefit</u>								
Phenylketonuria.								
1411G NP	Sachets 18.2 g, 60	3	5	..	*1640.07	34.20	add-ins	SB
2382J NP	Oral liquid 87 mL, 30	4	5	..	*1034.78	34.20	PKU Cooler 10	VF
2474F NP	Oral liquid 174 mL, 30	4	5	..	*2054.02	34.20	PKU Cooler 20	VF
2738D NP	Powder 500 g	8	5	..	*884.02	34.20	XP Maxamaid	SB
2739E NP	Powder 500 g	8	5	..	*1352.42	34.20	XP Maxamum	SB
8545R NP	Powder 400 g	8	5	..	*848.58	34.20	Phenex-2	AB
8555G NP	Sachets 20 g, 30	4	5	..	*891.02	34.20	PKU-gel	VF
8591E NP	Sachets 25 g, 30	4	5	..	*1549.14	34.20	PKU-Express	VF

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
8613H NP	Sachets 29 g, 30	4	5	..	*892.10	34.20	PKU Anamix Junior	SB
8727H NP	Sachets 50 g, 30	3	5	..	*1512.06	34.20	XP Maxamum	SB
8746H NP	Oral liquid 250 mL	90	5	..	*1313.27	34.20	Easiphen	SB
8804J NP	Sachets 27.8 g, 30	3	5	..	*1549.44	34.20	Lophlex	SB
8846N NP	Oral liquid 130 mL, 30	4	5	..	*1548.34	34.20	PKU Cooler 15	VF
9021T NP	Oral liquid 125 mL, 30	3	5	..	*1549.44	34.20	PKU Lophlex LQ 20	SB
9396M NP	Oral liquid 125 mL, 36	4	5	..	*1269.86	34.20	PKU Anamix Junior LQ	SB
9397N NP	Oral liquid 62.5 mL, 60	2	5	..	*1059.36	34.20	PKU Lophlex LQ 10	SB

AMINO ACID FORMULA with VITAMINS and MINERALS without PHENYLALANINE and TYROSINE

Restricted benefit

Tyrosinaemia.

3078B NP	Powder 500 g	8	5	..	*2704.74	34.20	XPhen, Tyr Maxamum	SB
8445L NP	Infant formula, powder 400 g	8	5	..	*769.30	34.20	TYR Anamix infant	SB
8446M NP	Powder 500 g	8	5	..	*1784.74	34.20	XPhen, Tyr Maxamaid	SB
8631G NP	Sachets 20 g, 30	4	5	..	*1798.42	34.20	TYR gel	VF
8667E NP	Sachets 25 g, 30	4	5	..	*3098.38	34.20	TYR Express	VF
9132P NP	Oral liquid 130 mL, 30	4	5	..	*3098.38	34.20	TYR Cooler	VF
9395L NP	Sachets 29 g, 30	4	5	..	*1800.46	34.20	TYR Anamix Junior	SB

AMINO ACID FORMULA with VITAMINS and MINERALS without VALINE, LEUCINE and ISOLEUCINE

Restricted benefit

Maple syrup urine disease.

2375B NP	Oral liquid 130 mL, 30	4	5	..	*3098.38	34.20	MSUD Cooler	VF
2380G NP	Infant formula, powder 400 g	8	5	..	*769.30	34.20	MSUD Anamix infant	SB
8057C NP	Powder 500 g	8	5	..	*2704.74	34.20	MSUD Maxamum	SB
8260R NP	Powder 500 g	8	5	..	*1784.74	34.20	MSUD Maxamaid	SB
8310J NP	Powder 500 g	4	5	..	*2671.98	34.20	MSUD AID III	SB
8592F NP	Sachets 20 g, 30	4	5	..	*1798.42	34.20	MSUD-gel	VF
8632H NP	Sachets 25 g, 30	4	5	..	*3098.38	34.20	MSUD Express	VF
8745G NP	Sachets 29 g, 30	4	5	..	*1800.46	34.20	MSUD Anamix Junior	SB

AMINO ACID FORMULA with VITAMINS and MINERALS without VALINE, LEUCINE and ISOLEUCINE with FAT, CARBOHYDRATE and TRACE ELEMENTS and supplemented with DOCOSAHEXANOIC ACID

Restricted benefit

Maple syrup urine disease.

9499Y NP	Oral liquid 125 mL, 36	4	5	..	*2507.98	34.20	MSUD Anamix Junior LQ	SB

ARGININE with CARBOHYDRATE

Restricted benefit

Urea cycle disorders.

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Note								
Arginine with carbohydrate is not indicated for the treatment of arginase deficiency and other inborn errors of protein metabolism.								
9437Q NP	Sachets 4 g containing 500 mg arginine, 30	4	5	..	*516.02	34.20	Arginine Amino Acid Supplement	VF
CARBOHYDRATE, FAT, VITAMINS, MINERALS and TRACE ELEMENTS								
Restricted benefit								
Patients with proven inborn errors of protein metabolism who are unable to meet their energy requirements with permitted food and formulae.								
8369L NP	Powder 400 g	8	5	..	*291.46	34.20	Energivit	SB
CYSTINE with CARBOHYDRATE								
Restricted benefit								
Pyridoxine non-responsive homocystinuria.								
9164H NP	Sachets 4 g containing 500 mg cystine, 30	4	5	..	*516.02	34.20	Cystine Amino Acid Supplement	VF
ESSENTIAL AMINO ACIDS FORMULA								
Restricted benefit								
Gyrate atrophy of the choroid and retina;								
Urea cycle disorders.								
9329B NP	Powder 200 g, 2	3	5	..	*1200.57	34.20	Essential Amino Acid Mix	SB
ESSENTIAL AMINO ACIDS FORMULA with MINERALS and VITAMIN C								
Restricted benefit								
Gyrate atrophy of the choroid and retina;								
Urea cycle disorders.								
2027Q NP	Powder 400 g	5	5	..	*634.17	34.20	Dialamine	SB
ESSENTIAL AMINO ACIDS FORMULA with VITAMINS and MINERALS								
Restricted benefit								
Gyrate atrophy of the choroid and retina;								
Urea cycle disorders.								
9385Y NP	Sachets 12.5 g, 50	4	5	..	*1516.50	34.20	EAA Supplement	VF
HIGH FAT FORMULA with VITAMINS, MINERALS and TRACE ELEMENTS and low in PROTEIN and CARBOHYDRATE								
Restricted benefit								
Patients with intractable seizures requiring treatment with a ketogenic diet;								
Glucose transport protein defects;								
Pyruvate dehydrogenase deficiency.								
Note								
KetoCal should only be used under strict supervision of a dietician, together with a metabolic physician and/or neurologist.								
Note								
Authorities for increased maximum quantities, up to a maximum of 48, may be authorised.								
9446E NP	Powder 300 g	24	5	..	*988.26	34.20	KetoCal	SB
ISOLEUCINE with CARBOHYDRATE								
Restricted benefit								
Maple syrup urine disease.								
9134R NP	Sachets 4 g containing 50 mg isoleucine, 30	4	5	..	*516.02	34.20	Isoleucine Amino Acid Supplement	VF
9436P NP	Sachets 4 g containing 1 g isoleucine, 30	4	5	..	*566.98	34.20	Isoleucine 1000 Amino Acid	VF

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	Supplement
MILK PROTEIN and FAT FORMULA with VITAMINS and MINERALS—CARBOHYDRATE FREE								
<u>Restricted benefit</u>								
Patients with intractable seizures requiring treatment with a ketogenic diet;								
Glucose transport protein defects;								
Pyruvate dehydrogenase deficiency;								
Infants and young children with glucose-galactose intolerance and multiple monosaccharide intolerance.								
8630F NP	Powder 225 g	24	5	..	*648.42	34.20	Carbohydrate Free Mixture	SB
PHENYLALANINE with CARBOHYDRATE								
<u>Restricted benefit</u>								
Tyrosinaemia.								
9384X NP	Sachets 4 g containing 50 mg phenylalanine, 30	4	5	..	*516.02	34.20	Phenylalanine Amino Acid Supplement	VF
SOY PROTEIN and FAT FORMULA with VITAMINS and MINERALS—CARBOHYDRATE FREE								
<u>Restricted benefit</u>								
Patients with intractable seizures requiring treatment with a ketogenic diet;								
Glucose transport protein defects;								
Pyruvate dehydrogenase deficiency;								
Infants and young children with glucose-galactose intolerance and multiple monosaccharide intolerance.								
8577K NP	Liquid 384 mL	120	5	..	*670.02	34.20	RCF	AB
TRIGLYCERIDES, LONG CHAIN with GLUCOSE POLYMER								
<u>Restricted benefit</u>								
Patients with proven inborn errors of protein metabolism who are unable to meet their energy requirements with permitted food and formulae.								
9308X NP	Oral liquid 250 mL, 18	6	5	..	*339.78	34.20	ProZero	VF
9309Y NP	Oral liquid 1 L, 6	4	5	..	*304.02	34.20	ProZero	VF
TRIGLYCERIDES, MEDIUM CHAIN and LONG CHAIN with GLUCOSE POLYMER								
<u>Restricted benefit</u>								
Patients with proven inborn errors of protein metabolism who are unable to meet their energy requirements with permitted food and formulae.								
3136C NP	Compound powder 400 g	8	5	..	*295.54	34.20	Duocal	SB
TRIGLYCERIDES—MEDIUM CHAIN, FORMULA								
<u>Note</u>								
No applications for increased maximum quantities and/or repeats will be authorised.								
<u>Authority required</u>								
Chylous ascites;								
Chylothorax;								
Fat malabsorption due to liver disease, short gut syndrome, cystic fibrosis and gastrointestinal disorders;								
Hyperlipoproteinaemia type 1;								
Long chain fatty acid oxidation disorders.								
<u>Note</u>								
MCT Pro-Cal is not indicated for the treatment of intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect requiring a ketogenic diet.								
9383W NP	Sachets 16 g, 30	4	5	..	*253.62	34.20	MCT Pro-Cal	VF

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
TYROSINE with CARBOHYDRATE								
<u>Restricted benefit</u>								
Phenylketonuria.								
9165J NP	Sachets 4 g containing 1 g tyrosine, 30	4	5	..	*516.02	34.20	Tyrosine Amino Acid Supplement	VF
VALINE with CARBOHYDRATE								
<u>Restricted benefit</u>								
Maple syrup urine disease.								
9135T NP	Sachets 4 g containing 50 mg valine, 30	4	5	..	*516.02	34.20	Valine Amino Acid Supplement	VF
9434M NP	Sachets 4 g containing 1 g valine, 30	4	5	..	*566.98	34.20	Valine 1000 Amino Acid Supplement	VF
VITAMINS, MINERALS and TRACE ELEMENTS with CARBOHYDRATE								
<u>Authority required</u>								
Infants and children whose vitamin and mineral intake is insufficient due to a specific diagnosis requiring a highly restrictive therapeutic diet, and whose vitamin, mineral and trace element needs cannot be adequately met with other proprietary vitamin and mineral preparations.								
<u>Note</u>								
Paediatric Seravit should only be used under strict supervision of a dietitian and a paediatrician.								
9328Y NP	Powder 200 g	6	5	..	*390.42	34.20	Paediatric Seravit	SB
WHEY PROTEIN FORMULA supplemented with AMINO ACIDS, LONG CHAIN POLYUNSATURATED FATTY ACIDS, VITAMINS and MINERALS, and low in PROTEIN, PHOSPHATE, POTASSIUM and LACTOSE								
<u>Authority required</u>								
Infants and young children with chronic renal failure requiring treatment with a low protein and a low phosphorus diet, or a low protein, a low phosphorus and a low potassium diet.								
9382T NP	Sachets 100 g, 10	9	5	..	*1485.57	34.20	RenaStart	VF
WHEY PROTEIN FORMULA supplemented with AMINO ACIDS, VITAMINS and MINERALS, and low in PROTEIN, PHOSPHATE, POTASSIUM and LACTOSE								
<u>Authority required</u>								
Infants and young children with chronic renal failure requiring treatment with a low protein and a low phosphorus diet, or a low protein, a low phosphorus and a low potassium diet.								
8587Y NP	Powder 400 g	16	5	..	*1065.94	34.20	Kindergen	SB

All other non-therapeutic products

All other non-therapeutic products

Solvents and diluting agents, incl. irrigating solutions

SODIUM CHLORIDE								
2026P NP	Injection 9 mg per mL (0.9%), 10 mL	5	1	..	16.48	17.55	Pfizer Australia Pty Ltd	PF

PREPARATIONS WHICH MAY BE PRESCRIBED FOR PATIENTS RECEIVING PALLIATIVE CARE

The prescribing of medications listed in this section is in accordance with the requirements for General Pharmaceutical Benefits in the Schedule unless otherwise detailed in the listing for the item.

In addition, certain additional principles have been applied by the Pharmaceutical Benefits Advisory Committee (PBAC) in recommending for whom these medications may be prescribed, and the number of repeats that may be approved by Medicare Australia. These principles have been encompassed in the listings for the items, and further details are provided below to help doctors prescribing under this section.

For the purposes of this section a patient receiving palliative care is defined as:

- *A patient with an active, progressive, far-advanced disease for whom the prognosis is limited and the focus of care is the quality of life.*

The provision for increased maximum quantities and up to 3 repeats on the *initial* authority prescription is intended to provide up to 4 months' therapy in total. Where *continuing* treatment is required the provision of repeats is subject to confirmation by the prescriber that a palliative care physician or palliative care service has been consulted regarding the care of the patient.

Prescribers must heed State/Territory laws when prescribing drugs listed as narcotic, specified or restricted and must notify, or receive approval from, the appropriate health authority.

When a Palliative Care authority application is for a drug of addiction, the following guidelines apply:

- the maximum quantity authorised is generally for 1 month's therapy;
- where supply for a longer period is warranted, quantities are for up to 3 months' therapy;
- telephone approvals are limited to 1 month's therapy.

Doctors should also state (on the prescription) the interval of repeat where repeats are called for, and ensure State/Territory health authorities are notified about ongoing treatment.

Prescribers should be aware that patients receiving palliative care may also access PBS items included in the general part of the Schedule of Pharmaceutical Benefits including narcotic preparations, according to the restrictions that apply to individual items and the requirements that apply to the general part of the Schedule.

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$		

Alimentary tract and metabolism

Stomatological preparations

Stomatological preparations

Other agents for local oral treatment

BENZYDAMINE HYDROCHLORIDE

Authority required (STREAMLINED)

3634

Initial supply, for up to 4 months, for a palliative care patient where a painful mouth is a problem.

5385K NP	Mouth and throat rinse 22.5 mg per 15 mL, 500 mL	£1	3	..	22.26	23.33	Diffiam	IA

BENZYDAMINE HYDROCHLORIDE

Authority required (STREAMLINED)

3635

Continuing supply for a palliative care patient where a painful mouth is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5386L NP	Mouth and throat rinse 22.5 mg per 15 mL, 500 mL	£1	..	..	22.26	23.33	Diffiam	IA

CARMELLOSE SODIUM

Authority required (STREAMLINED)

3636

Initial supply, for up to 4 months, for a palliative care patient where dry mouth is a symptom.

5333Q NP	Mouth spray 10 mg per mL, 25 mL	£1	3	..	10.79	11.86	Aquae	HA
5334R NP	Mouth spray 10 mg per mL, 100 mL	£1	3	..	12.46	13.53	Aquae	HA

CARMELLOSE SODIUM

Authority required (STREAMLINED)

3637

Continuing supply for a palliative care patient where dry mouth is a symptom.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5335T NP	Mouth spray 10 mg per mL, 25 mL	£1	..	..	10.79	11.86	Aquae	HA
5336W NP	Mouth spray 10 mg per mL, 100 mL	£1	..	..	12.46	13.53	Aquae	HA

HYPROMELLOSE

Authority required (STREAMLINED)

3636

Initial supply, for up to 4 months, for a palliative care patient where dry mouth is a symptom.

5421H NP	Oral gel 20 mg per g, 100 g	£1	3	..	12.78	13.85	Aquae Gel	HA

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
HYPROMELLOSE							
<u>Authority required (STREAMLINED)</u>							
3637							
Continuing supply for a palliative care patient where dry mouth is a symptom.							
<u>Note</u>							
Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.							
5422J NP	Oral gel 20 mg per g, 100 g	‡1	..	..	12.78	13.85	Aquae Gel HA

Drugs for functional gastrointestinal disorders

Belladonna and derivatives, plain

Belladonna alkaloids semisynthetic, quaternary ammonium compounds

HYOSCINE BUTYLBROMIDE

Authority required (STREAMLINED)

3638

Initial supply, for up to 4 months, for a palliative care patient where colicky pain is a symptom.

5317W NP	Injection 20 mg in 1 mL	30	3	..	*108.54	34.20	Buscopan	BY

HYOSCINE BUTYLBROMIDE

Authority required (STREAMLINED)

3639

Continuing supply for a palliative care patient where colicky pain is a symptom.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5318X NP	Injection 20 mg in 1 mL	30	..	..	*108.54	34.20	Buscopan	BY

Antiemetics and antinauseants

Antiemetics and antinauseants

Other antiemetics

PROMETHAZINE HYDROCHLORIDE

Authority required (STREAMLINED)

3640

Initial supply, for up to 4 months, for a palliative care patient where nausea and/or vomiting is a problem.

5325G NP	Tablet 10 mg	50	3	..	14.67	15.74	Phenergan	SW
5326H NP	Tablet 25 mg	50	3	..	16.76	17.83	Phenergan	SW
5327J NP	Oral liquid 5 mg per 5 mL, 100 mL	‡1	3	..	15.34	16.41	Phenergan	SW

PROMETHAZINE HYDROCHLORIDE

Authority required (STREAMLINED)

3641

Continuing supply for a palliative care patient where nausea and/or vomiting is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5328K NP	Tablet 10 mg	50	..	..	14.67	15.74	Phenergan	SW
5329L	Tablet 25 mg	50	..	..	16.76	17.83	Phenergan	SW

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
5330M <i>NP</i>	Oral liquid 5 mg per 5 mL, 100 mL	‡1	..	..	15.34	16.41	Phenergan	SW

Laxatives

Laxatives

Contact laxatives

BISACODYL

Authority required (STREAMLINED)

3642

Initial supply, for up to 4 months, for a palliative care patient where constipation is a problem.

5301B <i>NP</i>	Tablet 5 mg	200	3	..	14.27	15.34	Bisalax	AS
							Lax-Tab	AE
5303D <i>NP</i>	Suppositories 10 mg, 10	3	3	..	*21.27	22.34 ^a	Petrus Bisacodyl Suppositories	PP
				^B 1.08	*22.35	22.34 ^a	Dulcolax	BY
5304E <i>NP</i>	Suppositories 10 mg, 12	3	3	..	*18.57	19.64	Petrus Bisacodyl Suppositories	PP

BISACODYL

Authority required (STREAMLINED)

3643

Continuing supply for a palliative care patient where constipation is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5305F <i>NP</i>	Tablet 5 mg	200	..	..	14.27	15.34	Bisalax	AS
							Lax-Tab	AE
5307H <i>NP</i>	Suppositories 10 mg, 10	3	..	..	*21.27	22.34 ^a	Petrus Bisacodyl Suppositories	PP
				^B 1.08	*22.35	22.34 ^a	Dulcolax	BY
5308J <i>NP</i>	Suppositories 10 mg, 12	3	..	..	*18.57	19.64	Petrus Bisacodyl Suppositories	PP

Bulk producers

STERCULIA with FRANGULA BARK

Authority required (STREAMLINED)

3642

Initial supply, for up to 4 months, for a palliative care patient where constipation is a problem.

5322D <i>NP</i>	Granules 620 mg-80 mg per g (62%-8%), 500 g	‡1	3	..	24.95	26.02	Normacol Plus	NE

STERCULIA with FRANGULA BARK

Authority required (STREAMLINED)

3643

Continuing supply for a palliative care patient where constipation is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5324F <i>NP</i>	Granules 620 mg-80 mg per g (62%-8%), 500 g	‡1	..	..	24.95	26.02	Normacol Plus	NE

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
<i>Osmotically acting laxatives</i>								
LACTULOSE								
<u>Authority required (STREAMLINED)</u>								
3642								
Initial supply, for up to 4 months, for a palliative care patient where constipation is a problem.								
5387M NP	Mixture 3.34 g per 5 mL, 500 mL	3	3	..	*29.85	30.92	^a	Actilax AF
							^a	Genlac SI
							^a	GenRx Lactulose GX
							^a	Lac-Dol GM
							^a	Lactocur SZ
				^B 5.01	*34.86	30.92	^a	Duphalac SM
LACTULOSE								
<u>Authority required (STREAMLINED)</u>								
3643								
Continuing supply for a palliative care patient where constipation is a problem.								
<u>Note</u>								
Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.								
5388N NP	Mixture 3.34 g per 5 mL, 500 mL	3	..	..	*29.85	30.92	^a	Actilax AF
							^a	Genlac SI
							^a	GenRx Lactulose GX
							^a	Lac-Dol GM
							^a	Lactocur SZ
				^B 5.01	*34.86	30.92	^a	Duphalac SM
MACROGOL 3350								
<u>Authority required (STREAMLINED)</u>								
3642								
Initial supply, for up to 4 months, for a palliative care patient where constipation is a problem.								
5389P NP	Sachets containing powder for solution 13.125 g with electrolytes, 30	2	3	..	*34.68	34.20		Movicol NE
5426N NP	Powder for oral solution 510 g	2	3	..	*34.68	34.20	^a	MediHealth ClearLax ON
							^a	OsmoLax KY
MACROGOL 3350								
<u>Authority required (STREAMLINED)</u>								
3643								
Continuing supply for a palliative care patient where constipation is a problem.								
<u>Note</u>								
Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.								
5390Q NP	Sachets containing powder for solution 13.125 g with electrolytes, 30	2	..	..	*34.68	34.20		Movicol NE
5427P NP	Powder for oral solution 510 g	2	..	..	*34.68	34.20	^a	MediHealth ClearLax ON
							^a	OsmoLax KY

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$		

Enemas

BISACODYL

Authority required (STREAMLINED)

3642

Initial supply, for up to 4 months, for a palliative care patient where constipation is a problem.

5302C NP	Enemas 10 mg in 5 mL, 25	£1	3	..	38.59	34.20	Bisalax	AS

BISACODYL

Authority required (STREAMLINED)

3643

Continuing supply for a palliative care patient where constipation is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5306G NP	Enemas 10 mg in 5 mL, 25	£1	..	..	38.59	34.20	Bisalax	AS

SORBITOL with SODIUM CITRATE and SODIUM LAURYL SULFOACETATE

Authority required (STREAMLINED)

3642

Initial supply, for up to 4 months, for a palliative care patient where constipation is a problem.

5331N NP	Enemas 3.125 g-450 mg-45 mg in 5 mL, 12	2	3	..	*32.80	33.87	^a Micolette	AE
							^a Microlax	JT

SORBITOL with SODIUM CITRATE and SODIUM LAURYL SULFOACETATE

Authority required (STREAMLINED)

3643

Continuing supply for a palliative care patient where constipation is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5332P NP	Enemas 3.125 g-450 mg-45 mg in 5 mL, 12	2	..	..	*32.80	33.87	^a Micolette	AE
							^a Microlax	JT

Peripheral opioid receptor antagonists

METHYLNALTREXONE

Authority required

Initial supply, in combination with oral laxatives, for a palliative care patient with opioid-induced constipation who has failed to respond to laxatives.

Note

No applications for repeats will be authorised.

Note

Special Pricing Arrangements apply.

5423K NP	Solution for injection containing methylnaltrexone bromide 12 mg in 0.6 mL	3	..	..	*130.59	34.20	Relistor	WX

METHYLNALTREXONE

Authority required

Continuing supply, in combination with oral laxatives, for a palliative care patient with opioid-induced constipation who has demonstrated a response to methylnaltrexone.

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Note

For first continuing supply, applications for increased repeats may be authorised.

Where consultation with a palliative care specialist or service has occurred, applications for increased repeats may be authorised.

Note

Special Pricing Arrangements apply.

5424L NP	Solution for injection containing methylnaltrexone bromide 12 mg in 0.6 mL	7	..	..	287.84	34.20	Relistor	WX

Other laxatives**GLYCEROL****Authority required (STREAMLINED)****3642**

Initial supply, for up to 4 months, for a palliative care patient where constipation is a problem.

5311M NP	Suppositories 700 mg (for infants), 12	3	3	..	*18.84	19.91	Petrus Pharmaceuticals Pty Ltd	PP
5312N NP	Suppositories 1.4 g (for children), 12	3	3	..	*19.26	20.33	Petrus Pharmaceuticals Pty Ltd	PP
5313P NP	Suppositories 2.8 g (for adults), 12	3	3	..	*19.74	20.81	Petrus Pharmaceuticals Pty Ltd	PP

GLYCEROL**Authority required (STREAMLINED)****3643**

Continuing supply for a palliative care patient where constipation is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5314Q NP	Suppositories 700 mg (for infants), 12	3	..	..	*18.84	19.91	Petrus Pharmaceuticals Pty Ltd	PP
5315R NP	Suppositories 1.4 g (for children), 12	3	..	..	*19.26	20.33	Petrus Pharmaceuticals Pty Ltd	PP
5316T NP	Suppositories 2.8 g (for adults), 12	3	..	..	*19.74	20.81	Petrus Pharmaceuticals Pty Ltd	PP

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Musculo-skeletal system

Antiinflammatory and antirheumatic products

Antiinflammatory and antirheumatic products, non-steroids

Acetic acid derivatives and related substances

DICLOFENAC SODIUM

Authority required (STREAMLINED)

3645

Initial supply, for up to 4 months, for a palliative care patient where severe pain is a problem.

5361E NP	Tablet 25 mg (enteric coated)	100	3	..	*13.06	14.13	^a	APO-Diclofenac	TX
							^a	Chem mart	CH
							^a	Diclofenac	SI
							^a	Clonac 25	GM
							^a	Diclofenac-GA	SZ
							^a	Diclofenac Sandoz	AF
							^a	Fenac 25	TW
5362F NP	Tablet 50 mg (enteric coated)	50	3	..	*15.02	14.13	^a	Terry White Chemists	Diclofenac
							^a	Diclofenac	NV
							^a	Voltaren 25	TX
							^a	APO-Diclofenac	CH
							^a	Chem mart	SI
							^a	Diclofenac	GM
							^a	Clonac 50	SZ
							^a	Diclofenac-GA	AF
							^a	Diclofenac Sandoz	TW
							^a	Fenac	TX
				..	11.05	12.12	^a	Terry White Chemists	Diclofenac
							^a	Diclofenac	NV
				..	13.01	12.12	^a	Voltaren 50	NV

DICLOFENAC SODIUM

Authority required

Initial supply, for up to 4 months, for a palliative care patient where severe pain is a problem.

5363G NP	Suppository 100 mg	40	3	..	*24.92	25.99		Voltaren 100	NV

DICLOFENAC SODIUM

Authority required (STREAMLINED)

3646

Continuing supply for a palliative care patient where severe pain is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5364H NP	Tablet 25 mg (enteric coated)	100	..	..	*13.06	14.13	^a	APO-Diclofenac	TX
							^a	Chem mart	CH
							^a	Diclofenac	SI
							^a	Clonac 25	GM
							^a	Diclofenac-GA	SZ

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
5365J NP	Tablet 50 mg (enteric coated)	50	..	..	11.05	12.12	^a Fenac 25 AF
							^a Terry White Chemists Diclofenac TW
							^B 1.96 *15.02 14.13 ^a Voltaren 25 NV
							^a APO-Diclofenac TX
							^a Chem mart Diclofenac CH
							^a Clonac 50 SI
							^a Diclofenac-GA GM
							^a Diclofenac Sandoz SZ
							^a Fenac AF
							^a Terry White Chemists Diclofenac TW
				^B 1.96	13.01	12.12	^a Voltaren 50 NV

DICLOFENAC SODIUM**Authority required**

Continuing supply for a palliative care patient where severe pain is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5366K NP	Suppository 100 mg	40	..	..	*24.92	25.99	Voltaren 100	NV

INDOMETHACIN**Authority required (STREAMLINED)****3645**

Initial supply, for up to 4 months, for a palliative care patient where severe pain is a problem.

5377B NP	Capsule 25 mg	100	3	..	*12.08	13.15	^a Arthrexin	AF
					^B 2.16 *14.24	13.15	^a Indocid	AS

INDOMETHACIN**Authority required**

Initial supply, for up to 4 months, for a palliative care patient where severe pain is a problem.

5378C NP	Suppository 100 mg	40	3	..	*22.50	23.57	Indocid	AS

INDOMETHACIN**Authority required (STREAMLINED)****3646**

Continuing supply for a palliative care patient where severe pain is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5379D NP	Capsule 25 mg	100	..	..	*12.08	13.15	^a Arthrexin	AF
					^B 2.16 *14.24	13.15	^a Indocid	AS

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
INDOMETHACIN								
<u>Authority required</u>								
Continuing supply for a palliative care patient where severe pain is a problem.								
<u>Note</u>								
Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.								
5380E NP	Suppository 100 mg	40	..	..	*22.50	23.57	Indocid	AS
SULINDAC								
<u>Authority required (STREAMLINED)</u>								
3645								
Initial supply, for up to 4 months, for a palliative care patient where severe pain is a problem.								
5381F NP	Tablet 100 mg	100	3	..	*16.34	17.41	Aclin	AF
5382G NP	Tablet 200 mg	50	3	..	15.28	16.35	Aclin 200	AF
SULINDAC								
<u>Authority required (STREAMLINED)</u>								
3646								
Continuing supply for a palliative care patient where severe pain is a problem.								
<u>Note</u>								
Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.								
5383H NP	Tablet 100 mg	100	..	..	*16.34	17.41	Aclin	AF
5384J NP	Tablet 200 mg	50	..	..	15.28	16.35	Aclin 200	AF
Propionic acid derivatives								
IBUPROFEN								
<u>Authority required</u>								
Initial supply, for up to 4 months, for a palliative care patient where severe pain is a problem.								
5368M NP	Tablet 400 mg	90	3	..	*14.73	15.80	Brufen	AB
IBUPROFEN								
<u>Authority required</u>								
Continuing supply for a palliative care patient where severe pain is a problem.								
<u>Note</u>								
Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.								
5370P NP	Tablet 400 mg	90	..	..	*14.73	15.80	Brufen	AB
NAPROXEN								
<u>Authority required (STREAMLINED)</u>								
3645								
Initial supply, for up to 4 months, for a palliative care patient where severe pain is a problem.								
5345H NP	Tablet 250 mg	100	3	..	*13.72	14.79 ^a	Inza 250	AF
				^B 2.34	*16.06	14.79 ^a	Naprosyn	RO
5346J NP	Tablet 500 mg	50	3	..	12.91	13.98 ^a	Inza 500	AF
				^B 1.36	14.27	13.98 ^a	Naprosyn	RO

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer
5347K NP	Tablet 750 mg (sustained release)	28	3	..	12.38	13.45	^a	Proxen SR 750 MD
				^B 1.27	13.65	13.45	^a	Naprosyn SR750 RO
5348L NP	Tablet 1 g (sustained release)	28	3	..	14.37	15.44	^a	Proxen SR 1000 MD
				^B 1.34	15.71	15.44	^a	Naprosyn SR1000 RO

NAPROXEN**Authority required (STREAMLINED)****3647**

Initial supply, for up to 4 months, for a palliative care patient where severe pain is a problem in patients unable to take a solid dose form of a non-steroidal anti-inflammatory agent.

5397C NP	Oral suspension 125 mg per 5 mL, 474 mL	£1	3	..	78.17	34.20		Naprosyn RO

NAPROXEN**Authority required (STREAMLINED)****3648**

Continuing supply for a palliative care patient where severe pain is a problem in patients unable to take a solid dose form of a non-steroidal anti-inflammatory agent.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5398D NP	Oral suspension 125 mg per 5 mL, 474 mL	£1	..	..	78.17	34.20		Naprosyn RO

NAPROXEN**Authority required (STREAMLINED)****3646**

Continuing supply for a palliative care patient where severe pain is a problem.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5349M NP	Tablet 250 mg	100	..	..	*13.72	14.79	^a	Inza 250 AF
				^B 2.34	*16.06	14.79	^a	Naprosyn RO
5350N NP	Tablet 500 mg	50	..	..	12.91	13.98	^a	Inza 500 AF
				^B 1.36	14.27	13.98	^a	Naprosyn RO
5351P NP	Tablet 750 mg (sustained release)	28	..	..	12.38	13.45	^a	Proxen SR 750 MD
				^B 1.27	13.65	13.45	^a	Naprosyn SR750 RO
5352Q NP	Tablet 1 g (sustained release)	28	..	..	14.37	15.44	^a	Proxen SR 1000 MD
				^B 1.34	15.71	15.44	^a	Naprosyn SR1000 RO

NAPROXEN SODIUM**Authority required (STREAMLINED)****3645**

Initial supply, for up to 4 months, for a palliative care patient where severe pain is a problem.

Note

Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.

5353R NP	Tablet 550 mg	50	3	..	13.10	14.17	^a	Crysanal MD
				^B 2.29	15.39	14.17	^a	Anaprox 550 RO

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
NAPROXEN SODIUM							
<u>Authority required (STREAMLINED)</u>							
3646							
Continuing supply for a palliative care patient where severe pain is a problem.							
<u>Note</u>							
Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.							
<u>Note</u>							
Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.							
5354T	Tablet 550 mg	50	..	..	13.10	14.17 ^a	Crysanal MD
NP				^B 2.29	15.39	14.17 ^a	Anaprox 550 RO

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Nervous system

Analgesics

Opioids

Natural opium alkaloids

MORPHINE SULFATE

Caution

The risk of drug dependence is high.

Authority required

Initial supply, for up to 3 months, for a palliative care patient with severe disabling pain not responding to non-narcotic analgesics.

Note

Telephone approvals are limited to 1 month's therapy.

5393W NP	Tablet 10 mg	20	2	..	14.41	15.48	Sevredol	MF
5394X NP	Tablet 20 mg	20	2	..	15.39	16.46	Sevredol	MF

MORPHINE SULFATE

Caution

The risk of drug dependence is high.

Authority required

Continuing supply for a palliative care patient with severe disabling pain not responding to non-narcotic analgesics.

Note

Where consultation with a palliative care specialist or service has occurred, applications for increased repeats for up to 3 months' supply may be authorised.

Telephone approvals are limited to 1 month's therapy.

5395Y NP	Tablet 10 mg	20	..	..	14.41	15.48	Sevredol	MF
5396B NP	Tablet 20 mg	20	..	..	15.39	16.46	Sevredol	MF

MORPHINE SULFATE

Caution

The risk of drug dependence is high.

Authority required

Initial supply, for up to 3 months, for a palliative care patient with chronic severe disabling pain not responding to non-narcotic analgesics.

Note

Telephone approvals are limited to 1 month's therapy.

5391R NP	Tablet 200 mg (controlled release)	20	2	..	91.28	34.20	MS Contin	MF

MORPHINE SULFATE

Caution

The risk of drug dependence is high.

Authority required

Continuing supply for a palliative care patient with chronic severe disabling pain not responding to non-narcotic analgesics.

Note

Where consultation with a palliative care specialist or service has occurred, applications for increased repeats for up to 3 months' supply may be authorised.

Telephone approvals are limited to 1 month's therapy.

5392T NP	Tablet 200 mg (controlled release)	20	..	..	91.28	34.20	MS Contin	MF

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Phenylpiperidine derivatives

FENTANYL

Caution

The risk of drug dependence is high.

Authority required

Initial supply for dose titration for breakthrough pain in a palliative care patient with cancer who is receiving opioids for their persistent pain and where further escalation in the dose of morphine for breakthrough pain results in intolerable adverse effects.

Note

No applications for increased repeats will be authorised.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

Special Pricing Arrangements apply.

5401G NP	Lozenges 200 micrograms (as citrate), 3	3	..	..	*115.60	34.20	Actiq	OA
5402H NP	Lozenges 400 micrograms (as citrate), 3	3	..	..	*115.60	34.20	Actiq	OA
5403J NP	Lozenges 600 micrograms (as citrate), 3	3	..	..	*115.60	34.20	Actiq	OA
5404K NP	Lozenges 800 micrograms (as citrate), 3	3	..	..	*115.60	34.20	Actiq	OA
5405L NP	Lozenges 1200 micrograms (as citrate), 3	3	..	..	*115.60	34.20	Actiq	OA
5406M NP	Lozenges 1600 micrograms (as citrate), 3	3	..	..	*115.60	34.20	Actiq	OA

FENTANYL

Caution

The risk of drug dependence is high.

Authority required

Continuing supply for breakthrough pain in a palliative care patient with cancer who is receiving opioids for their persistent pain and where further escalation in the dose of morphine for breakthrough pain results in intolerable adverse effects.

Note

For first continuing supply, applications for increased repeats for up to 3 months' supply may be authorised.

Where consultation with a palliative care specialist or service has occurred, applications for increased repeats for up to 3 months' supply may be authorised.

Telephone approvals are limited to 1 month's therapy.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note

Special Pricing Arrangements apply.

5407N NP	Lozenges 200 micrograms (as citrate), 3	20	..	..	*680.13	34.20	Actiq	OA
5408P NP	Lozenges 400 micrograms (as citrate), 3	20	..	..	*680.13	34.20	Actiq	OA
5409Q NP	Lozenges 600 micrograms (as citrate), 3	20	..	..	*680.13	34.20	Actiq	OA
5410R NP	Lozenges 800 micrograms (as citrate), 3	20	..	..	*680.13	34.20	Actiq	OA
5411T NP	Lozenges 1200 micrograms (as citrate), 3	20	..	..	*680.13	34.20	Actiq	OA
5412W NP	Lozenges 1600 micrograms (as citrate), 3	20	..	..	*680.13	34.20	Actiq	OA

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Diphenylpropylamine derivatives

METHADONE HYDROCHLORIDE

Caution

The risk of drug dependence is high.

Authority required

Initial supply, for up to 3 months, for a palliative care patient with chronic severe disabling pain not responding to non-narcotic analgesics.

Note

Telephone approvals are limited to 1 month's therapy.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

5399E NP	Oral liquid 25 mg per 5 mL, 200 mL	1	2	..	18.92	19.99	Sigma Pharmaceuticals (Australia) Pty Ltd	SI

METHADONE HYDROCHLORIDE

Caution

The risk of drug dependence is high.

Authority required

Continuing supply for a palliative care patient with chronic severe disabling pain not responding to non-narcotic analgesics.

Note

Where consultation with a palliative care specialist or service has occurred, applications for increased repeats for up to 3 months' supply may be authorised.

Telephone approvals are limited to 1 month's therapy.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

5400F NP	Oral liquid 25 mg per 5 mL, 200 mL	1	..	..	18.92	19.99	Sigma Pharmaceuticals (Australia) Pty Ltd	SI

Other analgesics and antipyretics

Anilides

PARACETAMOL

Authority required (STREAMLINED)

3649

Initial supply, for up to 4 months, for a palliative care patient for analgesia or fever where alternative therapy cannot be tolerated.

5319Y NP	Suppositories 500 mg, 24	4	3	..	*84.46	34.20	Panadol	GC
5343F NP	Tablet 665 mg (modified release)	192	3	..	*16.64	17.71	Panadol Osteo	GC

PARACETAMOL

Authority required (STREAMLINED)

3650

Continuing supply for a palliative care patient for analgesia or fever where alternative therapy cannot be tolerated.

Note

Written or telephone authority applications for increased repeats may be approved where consultation with a palliative care specialist or service has occurred.

5320B NP	Suppositories 500 mg, 24	4	..	..	*84.46	34.20	Panadol	GC
5344G	Tablet 665 mg (modified release)	192	..	..	*16.64	17.71	Panadol Osteo	GC

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum		Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$		

NP

Antiepileptics

Antiepileptics

Benzodiazepine derivatives

CLONAZEPAM

Authority required

Initial supply, for up to 4 months, for a palliative care patient for the prevention of epilepsy.

Note

No applications for increased repeats will be authorised.

5337X NP	Tablet 500 micrograms	100	3	..	13.31	14.38	^a	Paxam 0.5	AF
				^B 1.79	15.10	14.38	^a	Rivotril	RO
5338Y NP	Tablet 2 mg	100	3	..	19.38	20.45	^a	Paxam 2	AF
				^B 2.04	21.42	20.45	^a	Rivotril	RO
5339B NP	Oral liquid 2.5 mg per mL, 10 mL	2	3	..	*15.04	16.11		Rivotril	RO

CLONAZEPAM

Authority required

Continuing supply for a palliative care patient for the prevention of epilepsy.

Note

Where consultation with a palliative care specialist or service has occurred, applications for increased repeats may be authorised.

5340C NP	Tablet 500 micrograms	100	..	..	13.31	14.38	^a	Paxam 0.5	AF
				^B 1.79	15.10	14.38	^a	Rivotril	RO
5341D NP	Tablet 2 mg	100	..	..	19.38	20.45	^a	Paxam 2	AF
				^B 2.04	21.42	20.45	^a	Rivotril	RO
5342E NP	Oral liquid 2.5 mg per mL, 10 mL	2	..	..	*15.04	16.11		Rivotril	RO

Psycholeptics

Anxiolytics

Benzodiazepine derivatives

DIAZEPAM

Authority required

Initial supply, for up to 4 months, for a palliative care patient where anxiety is a problem.

Note

No applications for increased repeats will be authorised.

5355W NP	Tablet 2 mg	50	3	..	7.79	8.86	^a	Antenex 2	AF
							^a	Valpam 2	SI
				^B 0.86	8.65	8.86	^a	Valium	RO
5356X NP	Tablet 5 mg	50	3	..	7.93	9.00	^a	Antenex 5	AF
							^a	Diazepam-GA	GM
							^a	Ranzepam	RA
							^a	Valpam 5	SI
				^B 0.88	8.81	9.00	^a	Valium	RO

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
DIAZEPAM							
Authority required							
Continuing supply for a palliative care patient where anxiety is a problem.							
Note							
Where consultation with a palliative care specialist or service has occurred, applications for increased repeats may be authorised.							
5357Y NP	Tablet 2 mg	50	..	..	7.79	8.86 ^a	Antenex 2 AF
						^a	Valpam 2 SI
				^B 0.86	8.65	8.86 ^a	Valium RO
5358B NP	Tablet 5 mg	50	..	..	7.93	9.00 ^a	Antenex 5 AF
						^a	Diazepam-GA GM
						^a	Ranzepam RA
						^a	Valpam 5 SI
				^B 0.88	8.81	9.00 ^a	Valium RO

OXAZEPAM**Authority required**

Initial supply, for up to 4 months, for a palliative care patient where anxiety is a problem.

Note

No applications for increased repeats will be authorised.

5371Q NP	Tablet 15 mg	50	3	..	*8.60	9.67 ^a	Alepam 15 AF
				^B 5.50	*14.10	9.67 ^a	Serepax SI
5372R NP	Tablet 30 mg	50	3	..	*8.92	9.99 ^a	Alepam 30 AF
						^a	APO-Oxazepam TX
						^a	Murelax FM
				^B 5.50	*14.42	9.99 ^a	Serepax SI

OXAZEPAM**Authority required**

Continuing supply for a palliative care patient where anxiety is a problem.

Note

Where consultation with a palliative care specialist or service has occurred, applications for increased repeats may be authorised.

5373T NP	Tablet 15 mg	50	..	..	*8.60	9.67 ^a	Alepam 15 AF
				^B 5.50	*14.10	9.67 ^a	Serepax SI
5374W NP	Tablet 30 mg	50	..	..	*8.92	9.99 ^a	Alepam 30 AF
						^a	APO-Oxazepam TX
						^a	Murelax FM
				^B 5.50	*14.42	9.99 ^a	Serepax SI

Hypnotics and sedatives**Benzodiazepine derivatives****NITRAZEPAM****Authority required**

Initial supply, for up to 4 months, for a palliative care patient where insomnia is a problem.

Note

No applications for increased repeats will be authorised.

5359C NP	Tablet 5 mg	50	3	..	*9.36	10.43 ^a	Alodorm AF
				^B 3.06	*12.42	10.43 ^a	Mogadon VT

**PREPARATIONS WHICH MAY BE PRESCRIBED FOR
PATIENTS RECEIVING PALLIATIVE CARE**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum		Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$		

NITRAZEPAM**Authority required**

Continuing supply for a palliative care patient where insomnia is a problem.

Note

Where consultation with a palliative care specialist or service has occurred, applications for increased repeats may be authorised.

5360D NP	Tablet 5 mg	50	..	..	*9.36	10.43	^a	Alodorm	AF
					^B 3.06	*12.42	^a	Mogadon	VT

TEMAZEPAM**Authority required**

Initial supply, for up to 4 months, for a palliative care patient where insomnia is a problem.

Note

No applications for increased repeats will be authorised.

5375X NP	Tablet 10 mg	50	3	..	*9.00	10.07	^a	APO-Temazepam	TX
							^a	Temaze	AF
							^a	Temtabs	FM
					^B 3.00	*12.00	^a	Normison	SI

TEMAZEPAM**Authority required**

Continuing supply for a palliative care patient where insomnia is a problem.

Note

Where consultation with a palliative care specialist or service has occurred, applications for increased repeats may be authorised.

5376Y NP	Tablet 10 mg	50	..	..	*9.00	10.07	^a	APO-Temazepam	TX
							^a	Temaze	AF
							^a	Temtabs	FM
					^B 3.00	*12.00	^a	Normison	SI

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	

Alimentary tract and metabolism

Stomatological preparations

Stomatological preparations

Antiinfectives and antiseptics for local oral treatment

AMPHOTERICIN								
3306B	Lozenge 10 mg	20	..	..	12.03	13.10	Fungilin	SI
NYSTATIN								
3343Y	Oral suspension 100,000 units per mL, 24 mL	‡1	..	..	10.85	11.92	Mycostatin	FM
							Nilstat	SI

Other agents for local oral treatment

BENZYDAMINE HYDROCHLORIDE								
<u>Restricted benefit</u>								
Radiation induced mucositis.								
5032W	Mouth and throat rinse 22.5 mg per 15 mL, 500 mL	‡1	..	..	22.26	23.33	Diffiam	IA

Drugs for functional gastrointestinal disorders

Propulsives

Propulsives

METOCLOPRAMIDE HYDROCHLORIDE								
5151D	Tablet 10 mg	25	..	..	8.20	9.27	Pramin	AF
				^B 3.02	11.22	9.27	Maxolon	VT
5153F	Injection 10 mg in 2 mL	10	..	..	12.99	14.06	Maxolon	VT

Antiemetics and antinauseants

Antiemetics and antinauseants

Other antiemetics

PROCHLORPERAZINE								
<u>Caution</u>								
Prochlorperazine may be associated with parkinsonism and tardive dyskinesia and should be used for short-term treatment only.								
5205Y	Tablet containing prochlorperazine maleate 5 mg	25	..	..	9.51	10.58	^a APO-Prochlorperazine	TX
							^a ProCalm	SI
							^a Prochlorperazine-GA	GM
							^a Stemzine	AV
				^B 3.45	12.96	10.58	^a Stemetil	SW
5206B	Injection containing prochlorperazine mesylate 12.5 mg in 1 mL	10	..	..	16.82	17.89	Stemetil	SW
5208D	Suppositories containing prochlorperazine equivalent to 25 mg prochlorperazine maleate, 5	‡1	..	..	19.93	21.00	Stemetil	SW
PROMETHAZINE HYDROCHLORIDE								
3374N	Injection 50 mg in 2 mL	10	..	..	*22.32	23.39	Hospira Pty Limited	HH

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	
Antidiarrheals, intestinal antiinflammatory/ antiinfective agents							

Intestinal antiinfectives

Antibiotics

NYSTATIN

3342X	Tablet 500,000 units	50	..	..	17.98	19.05	Nilstat	SI
3345C	Capsule 500,000 units	50	..	..	17.98	19.05	Nilstat	SI

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Blood and blood forming organs

Blood substitutes and perfusion solutions

I.V. solutions

Solutions for parenteral nutrition

GLUCOSE

5005K	I.V. infusion 139 mmol (anhydrous) per 500 mL (5%), 500 mL	5	..	..	*18.07	19.14	^a B. Braun Australia Pty Ltd	BR
							^a Fresenius Kabi Australia Pty Limited	PK
5106R	I.V. infusion 278 mmol (anhydrous) per L (5%), 1 L	5	..	..	*23.17	24.24	^a B. Braun Australia Pty Ltd	BR
							^a Baxter Healthcare Pty Ltd	BX
							^a Fresenius Kabi Australia Pty Limited	PK

Solutions affecting the electrolyte balance

SODIUM CHLORIDE

5021G	I.V. infusion 77 mmol per 500 mL (0.9%), 500 mL	5	..	..	*18.07	19.14	^a B. Braun Australia Pty Ltd	BR
							^a Fresenius Kabi Australia Pty Limited	PK
5212H	I.V. infusion 154 mmol per L (0.9%), 1 L	5	..	..	*23.17	24.24	^a B. Braun Australia Pty Ltd	BR
							^a Baxter Healthcare Pty Ltd	BX
							^a Fresenius Kabi Australia Pty Limited	PK
5213J	I.V. infusion 513 mmol per L (3%), 1 L	2	..	..	*16.54	17.61	Baxter Healthcare Pty Ltd	BX

SODIUM CHLORIDE with GLUCOSE

5214K	I.V. infusion 31 mmol-222 mmol (anhydrous) per L (0.18%-4%), 1 L	5	..	..	*23.52	24.59	Baxter Healthcare Pty Ltd	BX
5215L	I.V. infusion 19 mmol-104 mmol (anhydrous) per 500 mL (0.225%-3.75%), 500 mL	5	..	..	*28.77	29.84	Baxter Healthcare Pty Ltd	BX
5216M	I.V. infusion 39 mmol-69 mmol (anhydrous) per 500 mL (0.45%-2.5%), 500 mL	5	..	..	*28.77	29.84	Baxter Healthcare Pty Ltd	BX

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer	
					for Max. Qty \$	Recordable Value for Safety Net \$		

Cardiovascular system

Cardiac therapy

Antiarrhythmics, class I and III

Antiarrhythmics, class IB

LIGNOCAINE HYDROCHLORIDE

5142P	Injection 100 mg in 5 mL	5	..	..	37.33	34.20	Pfizer Australia Pty Ltd	PF

Cardiac stimulants excl. cardiac glycosides

Adrenergic and dopaminergic agents

ADRENALINE

5004J	Injection 1 mg in 1 mL (1 in 1,000)	5	..	..	20.34	21.41	AstraZeneca Pty Ltd	AP

Vasodilators used in cardiac diseases

Organic nitrates

GLYCERYL TRINITRATE

5108W	Tablets 600 micrograms, 100	±1	..	..	15.00	16.07	Lycinate	FM
				^B 3.00	18.00	16.07		SI

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum		Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$		

Dermatologicals

Corticosteroids, dermatological preparations

Corticosteroids, plain

Corticosteroids, weak (group I)

HYDROCORTISONE ACETATE

Restricted benefit

Treatment of corticosteroid-responsive dermatoses.

5111B	Cream 10 mg per g (1%), 30 g	‡1	..	..	8.94	10.01	^a	Cortic-DS 1%	FM
				^B 2.75	11.69	10.01	^a	Sigmacort	SI
5112C	Topical ointment 10 mg per g (1%), 30 g	‡1	..	..	8.94	10.01	^a	Cortic-DS 1%	FM
				^B 2.75	11.69	10.01	^a	Sigmacort	SI
5113D	Cream 10 mg per g (1%), 50 g	‡1	..	..	8.61	9.68	^a	Cortic-DS 1%	FM
				^B 0.08	8.69	9.68		Cortef	VT
				^B 2.76	11.37	9.68	^a	Sigmacort	SI
5114E	Topical ointment 10 mg per g (1%), 50 g	‡1	..	..	8.61	9.68	^a	Cortic-DS 1%	FM
				^B 2.76	11.37	9.68	^a	Sigmacort	SI

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer	
					for Max. Qty \$	Recordable Value for Safety Net \$		

Systemic hormonal preparations, excl. sex hormones and insulins

Corticosteroids for systemic use

Corticosteroids for systemic use, plain *Glucocorticoids*

BETAMETHASONE ACETATE with BETAMETHASONE SODIUM PHOSPHATE

Restricted benefit

For local intra-articular or peri-articular infiltration;

Keloid;

Lichen planus hypertrophic.

5034Y	Injection 3 mg-3.9 mg (equivalent to 5.7 mg betamethasone) in 1 mL	5	..	..	25.38	26.45	Celestone Chronodose	SH

HYDROCORTISONE SODIUM SUCCINATE

Restricted benefit

For use in a hospital.

5118J	Injection equivalent to 100 mg hydrocortisone with 2 mL solvent	6	..	..	*37.32	34.20	Solu-Cortef	PF
5119K	Injection equivalent to 250 mg hydrocortisone with 2 mL solvent	6	..	..	*59.82	34.20	Solu-Cortef	PF

METHYLPREDNISOLONE ACETATE

Restricted benefit

For local intra-articular or peri-articular infiltration.

5148Y	Injection 40 mg in 1 mL	5	..	..	24.60	25.67 ^a	Depo-Nisolone	KR
				^B 0.73	25.33	25.67 ^a	Depo-Medrol	PF

TRIAMCINOLONE ACETONIDE

Restricted benefit

For local intra-articular or peri-articular infiltration;

Keloid;

Lichen planus hypertrophic.

5233K	Injection 10 mg in 1 mL	5	..	..	25.38	26.45	Kenacort-A10	SI

Pancreatic hormones

Glycogenolytic hormones *Glycogenolytic hormones*

GLUCAGON HYDROCHLORIDE

5105Q	Injection set containing 1 mg (1 i.u.) and 1 mL solvent in disposable syringe	1	..	..	45.63	34.20	GlucaGen Hypokit	NO

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	
					for Max. Qty \$	Recordable Value for Safety Net \$	
Brand Name and Manufacturer							

Antiinfectives for systemic use

Antibacterials for systemic use

Tetracyclines

Tetracyclines

DOXYCYCLINE

Note

Bioequivalence has been demonstrated between doxycycline tablet 100 mg (as hydrochloride) and doxycycline tablet 100 mg (as monohydrate).

3321T	Tablet 100 mg (as hydrochloride)	7	..	..	8.47	9.54	^a	Doxsig	SI
							^a	Doxy-100	GM
							^a	Doxylin 100	AF
5082L	Tablet 100 mg (as monohydrate)	7	..	..	8.47	9.54	^B	1.19	
							^a	Vibramycin	PF
							^a	Chem mart	CH
							^a	Doxycycline	
							^a	Doxyhexal	SZ
							^a	GenRx Doxycycline	GX
							^a	Terry White	TW
								Chemists	
								Doxycycline	

DOXYCYCLINE

3322W	Capsule 100 mg (as hydrochloride)	7	..	..	8.47	9.54	^a	Mayne Pharma	YT
							^B	1.15	
					9.62	9.54	^a	Doryx	YN

Beta-lactam antibacterials, penicillins

Penicillins with extended spectrum

AMOXYCILLIN

3300Q	Capsule 500 mg	20	..	..	10.65	11.72	^a	Alphamox 500	AF
							^a	Amoxycillin-GA	GM
							^a	Amoxycillin	RA
								Ranbaxy	
							^a	Amoxycillin Sandoz	SZ
							^a	APO-Amoxycillin	TX
							^a	Chem mart	CH
								Amoxycillin	
							^a	Cilamox	SI
							^a	GenRx Amoxycillin	GX
3301R	Capsule 250 mg	20	..	..	8.55	9.62	^a	Terry White	TW
								Chemists	
								Amoxycillin	
							^B	0.80	
							^a	Amoxil	GK
							^a	Alphamox 250	AF
							^a	Amoxycillin-GA	GM
							^a	Amoxycillin	RA
								Ranbaxy	
							^a	Amoxycillin Sandoz	SZ
							^a	APO-Amoxycillin	TX
							^a	Chem mart	CH
								Amoxycillin	
							^a	Cilamox	SI
							^a	GenRx Amoxycillin	GX
							^a	Terry White	TW
								Chemists	
								Amoxycillin	
							^a	Cilamox	SI
							^a	GenRx Amoxycillin	GX

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
				^B 0.79	9.34	9.62	^a Amoxycillin
3302T	Powder for syrup 125 mg per 5 mL, 100 mL	‡1	..	..	#10.85	12.26	^a Amoxil GK ^a Alphamox 125 AF ^a Amoxycillin Sandoz SZ ^a Bgramin GM ^a Chem mart CH ^a Amoxycillin ^a GenRx Amoxycillin GX ^a Ranmoxy RA ^a Terry White Chemists TW ^a Amoxycillin
3309E	Sachet containing oral powder 3 g	1	..	..	9.11	10.18	^a Amoxil GK
3393N	Powder for syrup 250 mg per 5 mL, 100 mL	‡1	..	..	#11.69	13.10	^a Alphamox 250 AF ^a Amoxycillin Sandoz SZ ^a Bgramin GM ^a Chem mart CH ^a Amoxycillin ^a Cilamox SI ^a GenRx Amoxycillin GX ^a Ranmoxy RA ^a Terry White Chemists TW ^a Amoxycillin
5225B	Powder for oral suspension 500 mg per 5 mL, 100 mL	‡1	..	..	#12.48 #14.70	13.10 16.11	^a Amoxil Forte GK Maxamox SZ
AMPICILLIN							
3313J	Powder for injection 500 mg	5	..	..	11.08	12.15	^a Austrapen LN ^a Ibimicyn TS
3314K	Powder for injection 1 g	5	..	..	14.07	15.14	^a Aspen Ampicyn AS ^a Austrapen LN ^a Ibimicyn TS
Beta-lactamase sensitive penicillins							
BENZATHINE BENZYL PENICILLIN							
5027N	Injection 900 mg in 2.3 mL single use pre-filled syringe	10	..	..	293.11	34.20	Bicillin L-A AS
BENZYL PENICILLIN							
3398W	Powder for injection 600 mg	10	..	..	*42.92	34.20	BenPen CS
3399X	Powder for injection 3 g	10	..	..	*66.92	34.20	BenPen CS
PHENOXYMETHYL PENICILLIN							
3360W	Tablet 250 mg	50	..	..	*11.58	12.65	Abbocillin-VK Filmtab SI
3361X	Tablet 500 mg	50	..	..	*14.06	15.13	Abbocillin-VK Filmtab SI
3363B	Capsule 250 mg	50	..	..	11.41	12.48	^a Cilicaine VK FM ^a Cilopen VK GM LPV AS
3364C	Capsule 500 mg	50	..	..	13.84	14.91	^a Cilicaine VK FM ^a Cilopen VK GM LPV AS
5012T	Oral suspension 150 mg (as benzathine) per	2	..	..	*22.40	23.47	^a Cilicaine V FM

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
	5 mL, 100 mL			^B 2.00	*24.40	23.47 ^a	Abbocillin-V SI

3371K	PROCAINE PENICILLIN Injection 1.5 g	5	..	..	92.22	34.20	Cilicaine SI

Beta-lactamase resistant penicillins

DICLOXACILLIN

Restricted benefit

Serious staphylococcal infections.

5096F	Capsule 250 mg	24	..	..	11.45	12.52 ^a	Dicloxsig SI
						^a Distaph 250	AF
5097G	Capsule 500 mg	24	..	..	16.94	18.01 ^a	Diclocil BQ
						^a Dicloxsig	SI
						^a Distaph 500	AF

FLUCLOXACILLIN

Caution

Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

5094D	Powder for injection 500 mg	5	..	..	15.49	16.56 ^a	Flubiclox TS
						^a Flucil	AS
5095E	Powder for injection 1 g	5	..	..	20.66	21.73 ^a	Flubiclox TS
						^a Flucil	AS
						^a Hospira Pty Limited	HH

FLUCLOXACILLIN

Caution

Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

Restricted benefit

Serious staphylococcal infections.

5090X	Capsule 250 mg (as sodium)	24	..	..	11.45	12.52	^a Flopen	AS
						^a Staphylex 250	AF	
5091Y	Capsule 500 mg (as sodium)	24	..	..	16.94	18.01	^a Flopen	AS
						^a Staphylex 500	AF	
5257Q	Powder for oral liquid 125 mg (as sodium) per 5 mL, 100 mL	#1	..	..	#16.37	17.78	Aspen Pharmacare Australia Pty Limited	LN
5258R	Powder for oral liquid 250 mg (as sodium) per 5 mL, 100 mL	#1	..	..	#20.08	21.49	Aspen Pharmacare Australia Pty Limited	LN

Combinations of penicillins, incl. beta-lactamase inhibitors

AMOXYCILLIN with CLAVULANIC ACID

Caution

Hepatotoxicity has been reported with this drug.

Restricted benefit

Infections where resistance to amoxycillin is suspected;

Infections where resistance to amoxycillin is proven.

5006L	Tablet 875 mg-125 mg	10	..	..	14.60	15.67	^a Amoxicillin/ Clavulanic Acid 875/125 generichealth	GQ
							^a Chem mart	CH

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

					Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$					
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium			Brand Name and Manufacturer				
3318P	Capsule 500 mg	20	..	..	10.77	11.84	^a Cephalixin generichealth	GQ			
							^a Cephatrust 250	MI			
							^a Chem mart Cephalexin	CH			
							^a Cilex	GM			
							^a GenRx Cephalexin	GX			
							^a Ialex	LN			
							^a Ibilex 250	AF			
							^a Rancef	RA			
							^a Terry White Chemists Cephalexin	TW			
							^B 3.30	12.15	9.92	^a Keflex	AS
							^a Cefalexin Sandoz	SZ			
							^a Cephabell	BF			
							^a Cephalixin generichealth	GQ			
							^a Cephatrust 500	MI			
							^a Chem mart Cephalexin	CH			
							^a Cilex	GM			
							^a GenRx Cephalexin	GX			
							^a Ialex	LN			
							^a Ibilex 500	AF			
^a Rancef	RA										
^a Terry White Chemists Cephalexin	TW										
3319Q	Granules for syrup 125 mg per 5 mL, 100 mL	#1	..	..	#11.84	13.25	^B 4.41	15.18	11.84	^a Keflex	AS
							^a APO-Cephalexin	TX			
							^a Cefalexin Sandoz	SZ			
							^a Chem mart Cephalexin	CH			
							^a Cilex	GM			
							^a GenRx Cephalexin	GX			
							^a Ialex	LN			
							^a Ibilex 125	AF			
							^a Terry White Chemists Cephalexin	TW			
							^B 3.55	#15.39	13.25	^a Keflex	AS
3320R	Granules for syrup 250 mg per 5 mL, 100 mL	#1	..	..	#13.24	14.65	^a APO-Cephalexin	TX			
							^a Cefalexin Sandoz	SZ			
							^a Chem mart Cephalexin	CH			
							^a Cilex	GM			
							^a GenRx Cephalexin	GX			
							^a Ialex	LN			
							^a Ibilex 250	AF			
							^a Terry White Chemists Cephalexin	TW			
							^B 4.38	#17.62	14.65	^a Keflex	AS

Second-generation cephalosporins

CEFACLOR

Caution

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Serum sickness-like reactions have been reported with this drug, especially in children.							
5045M	Tablet 375 mg (sustained release)	10	..	..	12.89	13.96	^a Cefaclor-GA GN ^a Chem mart CH ^a Cefaclor CD ^a Douglas Cefaclor-CD GM ^a GenRx Cefaclor CD GX ^a Karlor CD LN ^a Keflor CD AF ^a Ozcef RA ^a Terry White Chemists TW ^a Cefaclor CD
				^B 5.20	18.09	13.96	^a Ceclor CD AS
5046N	Powder for oral suspension 125 mg per 5 mL, 100 mL	‡1	..	..	#13.50	14.91	^a Aclor 125 SI ^a Cefaclor Sandoz SZ ^a Chem mart CH ^a Cefaclor ^a GenRx Cefaclor GX ^a Keflor AF ^a Ozcef RA ^a Terry White Chemists TW ^a Cefaclor
				^B 4.18	#17.68	14.91	^a Ceclor AS
5047P	Powder for oral suspension 250 mg per 5 mL, 75 mL	‡1	..	..	#13.83	15.24	^a Aclor 250 SI ^a Cefaclor Sandoz SZ ^a Chem mart CH ^a Cefaclor ^a GenRx Cefaclor GX ^a Keflor AF ^a Ozcef RA ^a Terry White Chemists TW ^a Cefaclor
				^B 4.37	#18.20	15.24	^a Ceclor AS
CEFUROXIME AXETIL							
5052X	Tablet 250 mg (base)	14	..	..	18.62	19.69	Zinnat GK

Third-generation cephalosporins

CEFOTAXIME

Restricted benefit

Infections where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent.

5048Q	Powder for injection 1 g	10	..	..	*26.82	27.89	^a Cefotaxime Sandoz SZ ^a Hospira Pty Limited HH
					26.86	27.93	
5049R	Powder for injection 2 g	10	..	..	*43.62	34.20	^a Cefotaxime Sandoz SZ ^a Hospira Pty Limited HH
					43.68	34.20	

Sulfonamides and trimethoprim

Combinations of sulfonamides and trimethoprim, incl. derivatives

TRIMETHOPRIM with SULFAMETHOXAZOLE

Caution

There is an increased risk of severe adverse reactions with this combination in the elderly.

3389J	Tablet 80 mg-400 mg	10	..	..	8.67	9.74	Resprim AF

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer	
3390K	Tablet 160 mg-800 mg	10	..	..	9.39	10.46	^a	Bactrim DS	RO
							^a	Resprim Forte	AF
				^B 1.54	10.93	10.46	^a	Septrin Forte	SI
3391L	Oral suspension 40 mg-200 mg per 5 mL, 100 mL	‡1	..	..	9.05	10.12		Bactrim	RO
				^B 1.90	10.95	10.12		Septrin	SI

Macrolides, lincosamides and streptogramins

Macrolides

ERYTHROMYCIN

3325B	Capsule 250 mg	25	..	..	9.43	10.50	^a	Mayne Pharma	YT
				^B 1.35	10.78	10.50	^a	Erythromycin Eryc	YN

ERYTHROMYCIN ETHYL SUCCINATE

3334L	Powder for oral liquid 200 mg (base) per 5 mL, 100 mL	‡1	..	..	#12.32	13.73	^a	E-Mycin 200	AF
				^B 2.86	#15.18	13.73	^a	E.E.S. 200	LM
3336N	Tablet 400 mg (base)	25	..	..	10.92	11.99	^a	E-Mycin	AF
				^B 2.80	13.72	11.99	^a	E.E.S. 400 Filmtab	LM
3337P	Powder for oral liquid 400 mg (base) per 5 mL, 100 mL	‡1	..	..	#13.40	14.81	^a	E-Mycin 400	AF
				^B 2.89	#16.29	14.81	^a	E.E.S. Granules	LM

ERYTHROMYCIN LACTOBIONATE

5088T	Powder for I.V. infusion 1 g (base)	5	..	..	*88.92	34.20		Erythrocin-I.V.	LM

ROXITHROMYCIN

5259T	Tablet for oral suspension 50 mg	10	..	..	12.89	13.96		Rulide D	SW
5260W	Tablet 150 mg	10	..	..	11.76	12.83	^a	APO-Roxithromycin	TX
							^a	Biaxsig	AV
							^a	Chem mart Roxithromycin	CH
							^a	Roxar 150	SI
							^a	Roxide	SZ
							^a	Roximycin	AF
							^a	Roxithromycin-GA	GM
							^a	Terry White Chemists Roxithromycin	TW
				^B 2.60	14.36	12.83	^a	Rulide	SW
5261X	Tablet 300 mg	5	..	..	11.76	12.83	^a	APO-Roxithromycin	TX
							^a	Biaxsig	AV
							^a	Chem mart Roxithromycin	CH
							^a	Roxar 300	SI
							^a	Roxide	SZ
							^a	Roximycin	AF
							^a	Roxithromycin-GA	GM
							^a	Terry White Chemists Roxithromycin	TW
				^B 2.60	14.36	12.83	^a	Rulide	SW

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		Brand Name and Manufacturer	
Lincosamides									
CLINDAMYCIN									
<u>Restricted benefit</u>									
Gram-positive coccal infections where these cannot be safely and effectively treated with a penicillin.									
5057E	Capsule 150 mg	24	..	.. ^B 1.40	20.02 21.42	21.09 21.09	^a ^a	Cleocin Dalacin C	KR PF
LINCOMYCIN									
5144R	Injection 600 mg in 2 mL	5	..	..	33.74	34.20		Lincocin	PF
Other antibacterials									
<i>Glycopeptide antibacterials</i>									
VANCOMYCIN									
<u>Restricted benefit</u>									
Prophylaxis of endocarditis in patients hypersensitive to penicillin.									
3323X	Powder for injection 500 mg (500,000 i.u.) vancomycin activity	2	..	..	*17.96	19.03	^a ^a ^a	Hospira Pty Limited Vancocin CP Vancomycin Sandoz	HH AS SZ
5083M	Powder for injection 1 g (1,000,000 i.u.) vancomycin activity	1	..	..	17.95	19.02	^a ^a	Hospira Pty Limited Vancomycin Sandoz	HH SZ
Imidazole derivatives									
METRONIDAZOLE									
3339R	Tablet 200 mg	21	..	.. ^B 2.30	7.96 10.26	9.03 9.03	^a ^a ^a	Metrogyl 200 Metronide 200 Flagyl	AF AV SW
5157K	Suppositories 500 mg, 10	‡1	..	..	23.16	24.23		Flagyl	SW
5159M	Tablet 400 mg	5	..	..	7.88	8.95		Metrogyl 400	AF
METRONIDAZOLE									
<u>Restricted benefit</u>									
Treatment of anaerobic infections.									
5155H	Tablet 400 mg	21	..	.. ^B 2.30	10.03 12.33	11.10 11.10	^a ^a ^a	Metrogyl 400 Metronide 400 Flagyl	AF AV SW
METRONIDAZOLE									
<u>Restricted benefit</u>									
Treatment, in a hospital, of acute anaerobic sepsis.									
5154G	I.V. infusion 500 mg in 100 mL	5	..	..	*31.97 *32.00	33.04 33.07	^a ^a ^a	Baxter Healthcare Pty Ltd DBL Metronidazole Intravenous Infusion Metronidazole Sandoz	BX HH SZ
METRONIDAZOLE BENZOATE									
3341W	Oral suspension 320 mg per 5 mL (equivalent to	‡1	..	..	18.82	19.89		Flagyl S	SW

PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY

					Dispensed Price		Maximum		
					for		Recordable		
					Max. Qty		Value for		
					\$		Safety Net		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium					
	200 mg metronidazole in 5 mL), 100 mL							Brand Name and Manufacturer	

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

Musculo-skeletal system

Antiinflammatory and antirheumatic products

Antiinflammatory and antirheumatic products, non-steroids

Acetic acid derivatives and related substances

DICLOFENAC SODIUM								
5079H	Suppository 100 mg	40	..	..	*24.92	25.99	Voltaren 100	NV
DICLOFENAC SODIUM								
<u>Restricted benefit</u>								
Chronic arthropathies (including osteoarthritis) with an inflammatory component;								
Bone pain due to malignant disease.								
5076E	Tablet 25 mg (enteric coated)	100	..	..	*13.06	14.13	^a APO-Diclofenac	TX
							^a Chem mart	CH
							^a Diclofenac	
							^a Clonac 25	SI
							^a Diclofenac-GA	GM
							^a Diclofenac Sandoz	SZ
							^a Fenac 25	AF
							^a Terry White	TW
							^a Chemists	
							^a Diclofenac	
				^B 1.96	*15.02	14.13	^a Voltaren 25	NV
5077F	Tablet 50 mg (enteric coated)	50	..	..	11.05	12.12	^a APO-Diclofenac	TX
							^a Chem mart	CH
							^a Diclofenac	
							^a Clonac 50	SI
							^a Diclofenac-GA	GM
							^a Diclofenac Sandoz	SZ
							^a Fenac	AF
							^a Terry White	TW
							^a Chemists	
							^a Diclofenac	
				^B 1.96	13.01	12.12	^a Voltaren 50	NV
INDOMETHACIN								
5128X	Suppository 100 mg	40	..	..	*22.50	23.57	Indocid	AS
INDOMETHACIN								
<u>Restricted benefit</u>								
Chronic arthropathies (including osteoarthritis) with an inflammatory component;								
Bone pain due to malignant disease.								
5126T	Capsule 25 mg	100	..	..	*12.08	13.15	^a Arthrexin	AF
				^B 2.16	*14.24	13.15	^a Indocid	AS
SULINDAC								
<u>Restricted benefit</u>								
Chronic arthropathies (including osteoarthritis) with an inflammatory component;								
Bone pain due to malignant disease.								
5217N	Tablet 100 mg	100	..	..	*16.34	17.41	Aclin	AF

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
5218P	Tablet 200 mg	50	..	..	15.28	16.35	Acilin 200 AF

Oxicams

PIROXICAM

Restricted benefit

Chronic arthropathies (including osteoarthritis) with an inflammatory component.

5201R	Dispersible tablet 10 mg	50	..	..	12.50	13.57	Mobilis D-10	AF
5202T	Dispersible tablet 20 mg	25	..	..	12.20	13.27	^a Mobilis D-20	AF
				^B 2.64	14.84	13.27	^a Feldene-D	PF
5203W	Capsule 10 mg	50	..	..	12.50	13.57	^a Chem mart	CH
							^a Piroxicam	
							^a GenRx Piroxicam	GX
							^a Mobilis 10	AF
							^a Terry White	TW
							^a Chemists	
				^B 2.66	15.16	13.57	^a Piroxicam	
							^a Feldene	PF
5204X	Capsule 20 mg	25	..	..	12.20	13.27	^a Chem mart	CH
							^a Piroxicam	
							^a GenRx Piroxicam	GX
							^a Mobilis 20	AF
							^a Terry White	TW
							^a Chemists	
				^B 2.64	14.84	13.27	^a Piroxicam	
							^a Feldene	PF

Propionic acid derivatives

IBUPROFEN

5124Q	Tablet 400 mg	30	..	..	9.19	10.26	Brufen	AB

IBUPROFEN

Restricted benefit

Chronic arthropathies (including osteoarthritis) with an inflammatory component;

Bone pain due to malignant disease.

5123P	Tablet 400 mg	90	..	..	*14.73	15.80	Brufen	AB

KETOPROFEN

5139L	Suppository 100 mg	40	..	..	*25.30	26.37	Orudis	SW

KETOPROFEN

Restricted benefit

Chronic arthropathies (including osteoarthritis) with an inflammatory component.

5136H	Capsule 200 mg (sustained release)	28	..	..	19.37	20.44	^a Oruvail SR	AV
				^B 2.21	21.58	20.44	^a Orudis SR 200	SW

NAPROXEN

Restricted benefit

Chronic arthropathies (including osteoarthritis) with an inflammatory component;

Bone pain due to malignant disease.

5176K	Tablet 250 mg	100	..	..	*13.72	14.79	^a Inza 250	AF
				^B 2.34	*16.06	14.79	^a Naprosyn	RO
5177L	Tablet 500 mg	50	..	..	12.91	13.98	^a Inza 500	AF

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
				^B 1.36	14.27	13.98 ^a	Naprosyn RO
5178M	Tablet 750 mg (sustained release)	28	..	..	12.38	13.45 ^a	Proxen SR 750 MD
				^B 1.27	13.65	13.45 ^a	Naprosyn SR750 RO
5179N	Tablet 1 g (sustained release)	28	..	..	14.37	15.44 ^a	Proxen SR 1000 MD
				^B 1.34	15.71	15.44 ^a	Naprosyn SR1000 RO

NAPROXEN SODIUM

Restricted benefit

Chronic arthropathies (including osteoarthritis) with an inflammatory component;

Bone pain due to malignant disease.

Note

Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.

5186Y	Tablet 550 mg	50	..	..	13.10	14.17 ^a	Crysanal MD
				^B 2.29	15.39	14.17 ^a	Anaprox 550 RO

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer	
					for Max. Qty \$	Recordable Value for Safety Net \$		

Nervous system

Analgesics

Opioids

Natural opium alkaloids

CODEINE PHOSPHATE

Note

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

5063L	Tablet 30 mg	20	..	..	16.87	17.94	Fawns and McAllan Proprietary Limited	FM
CODEINE PHOSPHATE with PARACETAMOL								
3316M	Tablet 30 mg-500 mg	20	..	..	8.08	9.15	^a APO- Paracetamol/Co deine 500/30	TX
							^a Codalgin Forte	FM
							^a Codapane Forte	AL
							^a Comfarol Forte	SZ
							^a Dolaforte	CO
							^a Prodeine Forte	AV
				^B 2.80	10.88	9.15	^a Panadeine Forte	SW

HYDROMORPHONE HYDROCHLORIDE

Caution

The risk of drug dependence is high.

Note

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

5129Y	Injection 2 mg in 1 mL	5	..	..	22.84	23.91	Dilaudid	MF
5130B	Injection 10 mg in 1 mL	5	..	..	28.97	30.04	Dilaudid-HP	MF
5131C	Injection 50 mg in 5 mL	5	..	..	52.00	34.20	Dilaudid-HP	MF

HYDROMORPHONE HYDROCHLORIDE

Caution

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

Note

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

5115F	Tablet 2 mg	20	..	..	17.10	18.17	Dilaudid	MF
5116G	Tablet 4 mg	20	..	..	19.85	20.92	Dilaudid	MF
5117H	Tablet 8 mg	20	..	..	30.03	31.10	Dilaudid	MF
5132D	Oral liquid 1 mg per mL, 473 mL	1	..	..	63.70	34.20	Dilaudid	MF

HYDROMORPHONE HYDROCHLORIDE

Caution

The risk of drug dependence is high.

Restricted benefit

Chronic severe disabling pain not responding to non-narcotic analgesics.

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Note							
Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.							
3357Q	Tablet 8 mg (modified release)	14	..	..	36.41	34.20	Jurnista JC
3358R	Tablet 16 mg (modified release)	14	..	..	52.82	34.20	Jurnista JC
3367F	Tablet 32 mg (modified release)	14	..	..	88.70	34.20	Jurnista JC
3368G	Tablet 64 mg (modified release)	14	..	..	149.38	34.20	Jurnista JC
5023J	Tablet 4 mg (modified release)	14	..	..	30.95	32.02	Jurnista JC

MORPHINE HYDROCHLORIDE**Caution**

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

Note

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

5237P	Oral solution 2 mg per mL, 200 mL	1	..	..	18.16	19.23	Ordine 2	MF
5238Q	Oral solution 5 mg per mL, 200 mL	1	..	..	20.78	21.85	Ordine 5	MF
5239R	Oral solution 10 mg per mL, 200 mL	1	..	..	24.92	25.99	Ordine 10	MF

MORPHINE SULFATE**Caution**

The risk of drug dependence is high.

Note

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

5168B	Injection 10 mg in 1 mL	5	..	..	14.10	15.17	Hospira Pty Limited	HH
5169C	Injection 15 mg in 1 mL	5	..	..	14.45	15.52	Hospira Pty Limited	HH
5170D	Injection 30 mg in 1 mL	5	..	..	15.90	16.97	Hospira Pty Limited	HH

MORPHINE SULFATE**Caution**

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

Note

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

5163R	Tablet 30 mg	20	..	..	14.13	15.20	Anamorph	FM

MORPHINE SULFATE**Caution**

The risk of drug dependence is high.

Restricted benefit

Chronic severe disabling pain not responding to non-narcotic analgesics.

Note

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

5064M	Capsule 30 mg (controlled release)	10	..	..	20.12	21.19	MS Mono	MF
5065N	Capsule 60 mg (controlled release)	10	..	..	28.62	29.69	MS Mono	MF
5066P	Capsule 90 mg (controlled release)	10	..	..	32.67	33.74	MS Mono	MF
5067Q	Capsule 120 mg (controlled release)	10	..	..	43.37	34.20	MS Mono	MF
5161P	Tablet 15 mg (controlled release)	20	..	..	20.12	21.19	MS Contin	MF
5162Q	Tablet 5 mg (controlled release)	20	..	..	15.29	16.36	MS Contin	MF

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
5164T	Tablet 10 mg (controlled release)	20	..	..	17.08	18.15	^a Momex SR 10	SI
						^a	MS Contin	MF
5165W	Tablet 30 mg (controlled release)	20	..	..	28.62	29.69	^a Momex SR 30	SI
						^a	MS Contin	MF
5166X	Tablet 60 mg (controlled release)	20	..	..	43.37	34.20	^a Momex SR 60	SI
						^a	MS Contin	MF
5167Y	Tablet 100 mg (controlled release)	20	..	..	55.63	34.20	^a Momex SR 100	SI
						^a	MS Contin	MF
5171E	Sachet containing controlled release granules for oral suspension, 20 mg per sachet	20	..	..	47.53	34.20	MS Contin Suspension 20 mg	MF
5240T	Capsule 20 mg (containing sustained release pellets)	20	..	..	20.70	21.77	Kapanol	GK
5241W	Capsule 50 mg (containing sustained release pellets)	20	..	..	34.04	34.20	Kapanol	GK
5242X	Capsule 100 mg (containing sustained release pellets)	20	..	..	54.27	34.20	Kapanol	GK
5243Y	Sachet containing controlled release granules for oral suspension, 30 mg per sachet	20	..	..	48.71	34.20	MS Contin Suspension 30 mg	MF
5244B	Sachet containing controlled release granules for oral suspension, 60 mg per sachet	20	..	..	53.87	34.20	MS Contin Suspension 60 mg	MF
5245C	Sachet containing controlled release granules for oral suspension, 100 mg per sachet	20	..	..	65.42	34.20	MS Contin Suspension 100 mg	MF
5246D	Capsule 10 mg (containing sustained release pellets)	20	..	..	17.08	18.15	Kapanol	GK

OXYCODONE**Caution**

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

Note

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

5194J	Suppository 30 mg	12	..	..	43.66	34.20	Proladone	PL

OXYCODONE HYDROCHLORIDE**Caution**

The risk of drug dependence is high.

Restricted benefit

Severe disabling pain not responding to non-narcotic analgesics.

Note

Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

5190E	Oral solution 5 mg per 5 mL, 250 mL	1	..	..	20.72	21.79	OxyNorm Liquid 5mg/5mL	MF
5191F	Capsule 5 mg	20	..	..	12.30	13.37	OxyNorm	MF
5195K	Tablet 5 mg	20	..	..	12.30	13.37	Endone	SI
5197M	Capsule 10 mg	20	..	..	15.42	16.49	OxyNorm	MF
5198N	Capsule 20 mg	20	..	..	20.15	21.22	OxyNorm	MF

OXYCODONE HYDROCHLORIDE**Caution**

The risk of drug dependence is high.

Restricted benefit

Chronic severe disabling pain not responding to non-narcotic analgesics.

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	
Note							
Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.							
5015Y	Tablet 15 mg (controlled release)	20	..	..	27.93	29.00	OxyContin MF
5016B	Tablet 30 mg (controlled release)	20	..	..	41.22	34.20	OxyContin MF
5227D	Tablet 5 mg (controlled release)	20	..	..	21.18	22.25	OxyContin MF
5247E	Tablet 10 mg (controlled release)	20	..	..	21.95	23.02	OxyContin MF
5248F	Tablet 20 mg (controlled release)	20	..	..	31.87	32.94	OxyContin MF
5249G	Tablet 40 mg (controlled release)	20	..	..	48.35	34.20	OxyContin MF
5250H	Tablet 80 mg (controlled release)	20	..	..	71.70	34.20	OxyContin MF

Other opioids

TRAMADOL HYDROCHLORIDE

Restricted benefit

For acute pain where aspirin and/or paracetamol alone are inappropriate or have failed;

For dosage titration in chronic pain where aspirin and/or paracetamol alone are inappropriate or have failed.

5232J	Capsule 50 mg	20	..	..	9.16	10.23	^a APO-Tramadol TX
							^a Chem mart CH
							Tramadol
							^a GA Tramadol 50mg GM
							^a GenRx Tramadol GX
							^a Lodam 50 ZP
							^a Terry White TW
							Chemists
							Tramadol
							^a Tramadol Sandoz SZ
							^a Tramedo AF
							^a Zydol SI
				^B 2.42	11.58	10.23	^a Tramal CS

TRAMADOL HYDROCHLORIDE

Restricted benefit

For pain where aspirin and/or paracetamol alone are inappropriate or have failed.

3338Q	Tablet 50 mg (twice daily sustained release)	20	..	..	11.63	12.70	Tramal SR 50 CS
5001F	Tablet 100 mg (once a day extended release)	10	..	..	13.37	14.44	Durotram XR IA
5002G	Tablet 200 mg (once a day extended release)	10	..	..	16.34	17.41	Durotram XR IA
5003H	Tablet 300 mg (once a day extended release)	10	..	..	19.79	20.86	Durotram XR IA
5150C	Oral drops 100 mg per mL, 10 mL	±1	..	..	13.71	14.78	Tramal CS
5234L	Tablet 100 mg (twice daily sustained release)	20	..	..	13.86	14.93	^a APO-Tramadol SR TX
							^a Chem mart CH
							Tramadol SR
							^a GA Tramadol SR GM
							100mg
							^a Lodam SR 100 ZP
							^a Terry White TW
							Chemists
							Tramadol SR
							^a Tramahexal SR SZ
							^a Tramedo SR 100 AF
							^a Zydol SR 100 SI
				^B 4.51	18.37	14.93	^a Tramal SR 100 CS
5235M	Tablet 150 mg (twice daily sustained release)	20	..	..	16.45	17.52	^a APO-Tramadol SR TX
							^a Chem mart CH
							Tramadol SR
							^a GA Tramadol SR GM
							150mg

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
							^a Lodam SR 150	ZP
							^a Terry White Chemists Tramadol SR	TW
							^a Tramadol Sandoz SR	SZ
							^a Tramedo SR 150	AF
							^a Zydol SR 150	SI
				^B 5.37	21.82	17.52	^a Tramal SR 150	CS
5236N	Tablet 200 mg (twice daily sustained release)	20	..	..	18.63	19.70	^a APO-Tramadol SR	TX
							^a Chem mart Tramadol SR	CH
							^a GA Tramadol SR 200mg	GM
							^a Lodam SR 200	ZP
							^a Terry White Chemists Tramadol SR	TW
							^a Tramahexal SR	SZ
							^a Tramedo SR 200	AF
							^a Zydol SR 200	SI
				^B 6.09	24.72	19.70	^a Tramal SR 200	CS

TRAMADOL HYDROCHLORIDE

Restricted benefit

Short-term treatment of acute pain.

5231H	Injection 100 mg in 2 mL	5	..	..	14.30	15.37	^a Tramahexal	SZ
							^a Tramal 100	CS

Other analgesics and antipyretics
Salicylic acid and derivatives

ASPIRIN

5018D	Tablet 300 mg (dispersible)	96	..	..	8.55	9.62	Solprin	RC

Anilides

PARACETAMOL

3348F	Oral liquid 120 mg per 5 mL, 100 mL	‡1	..	..	9.38	10.45	Panamax	SW
3349G	Oral liquid 240 mg per 5 mL, 200 mL	‡1	..	..	10.68	11.75	Panamax 240 Elixir	SW
5196L	Tablet 500 mg	100	..	..	8.42	9.49	^a APO-Paracetamol	TX
							^a Chem mart Paracetamol	CH
							^a Febridol	GM
							^a Generic Health Pty Ltd	GQ
							^a Panamax	SW
							^a Paracetamol Sandoz	SZ
							^a Paralgin	FM
							^a Pharmacy Choice Paracetamol	YM
							^a Terry White Chemists Paracetamol	TW

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for		Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$			
PARACETAMOL									
<u>Restricted benefit</u>									
Chronic arthropathies.									
5224Y	Tablet 500 mg	300	..	..	*12.42	13.49	^a	APO-Paracetamol	TX
							^a	Chem mart	CH
								Paracetamol	
							^a	Febridol	GM
							^a	Generic Health Pty Ltd	GQ
							^a	Panamax	SW
							^a	Paracetamol	SZ
								Sandoz	
							^a	Paralgin	FM
							^a	Pharmacy Choice	YM
	Paracetamol								
							^a	Terry White	TW
								Chemists	
								Paracetamol	

Antiepileptics

Antiepileptics

Carboxamide derivatives

CARBAMAZEPINE								
5037D	Tablet 400 mg (controlled release)	200	..	..	51.02	34.20	Tegretol CR 400	NV
5038E	Tablet 200 mg (controlled release)	200	..	..	30.69	31.76	Tegretol CR 200	NV
5039F	Tablet 100 mg	200	..	..	19.14	20.21	^a Carbamazepine	SZ
							Sandoz	
				^B 2.56	*21.70	20.21	^a Tegretol 100	NV
5040G	Tablet 200 mg	200	..	..	30.20	31.27	^a Carbamazepine	SZ
							Sandoz	
							^a Teril	AF
				^B 2.74	*32.94	31.27	^a Tegretol 200	NV
5041H	Oral suspension 100 mg per 5 mL, 300 mL	‡1	..	..	21.35	22.42	Tegretol Liquid	NV

Anti-Parkinson drugs

Anticholinergic agents

Ethers of tropine or tropine derivatives

BENZTROPINE MESYLATE								
5031T	Injection 2 mg in 2 mL	5	..	..	103.59	34.20	Cogentin	FK

Psycholeptics

Anxiolytics

Benzodiazepine derivatives

DIAZEPAM								
5071X	Tablet 2 mg	50	..	..	7.79	8.86	^a Antenex 2	AF
							^a Valpam 2	SI
				^B 0.86	8.65	8.86	^a Valium	RO
5072Y	Tablet 5 mg	50	..	..	7.93	9.00	^a Antenex 5	AF
							^a Diazepam-GA	GM
							^a Ranzepam	RA
							^a Valpam 5	SI
				^B 0.88	8.81	9.00	^a Valium	RO

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
5073B	Injection 10 mg in 2 mL	5	..	..	12.29	13.36	Hospira Pty Limited	HH
OXAZEPAM								
5192G	Tablet 15 mg	25	..	..	7.51	8.58 ^a	Alepam 15	AF
				^B 2.75	10.26	8.58 ^a	Serepax	SI
5193H	Tablet 30 mg	25	..	..	7.67	8.74 ^a	Alepam 30	AF
						^a	APO-Oxazepam	TX
						^a	Murelax	FM
				^B 2.75	10.42	8.74 ^a	Serepax	SI
Hypnotics and sedatives								
<i>Benzodiazepine derivatives</i>								
NITRAZEPAM								
5189D	Tablet 5 mg	25	..	..	7.89	8.96 ^a	Alodorm	AF
				^B 1.53	9.42	8.96 ^a	Mogadon	VT
TEMAZEPAM								
5221T	Tablet 10 mg	25	..	..	7.71	8.78 ^a	APO-Temazepam	TX
						^a	Temaze	AF
						^a	Temtabs	FM
				^B 1.50	9.21	8.78 ^a	Normison	SI

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

Respiratory system

Drugs for obstructive airway diseases

Adrenergics for systemic use

Alpha- and beta-adrenoceptor agonists

ADRENALINE							
5004J	Injection 1 mg in 1 mL (1 in 1,000)	5	..	..	20.34	21.41	AstraZeneca Pty Ltd AP

Sensory organs

Ophthalmologicals

Antiinfectives

Antibiotics

CHLORAMPHENICOL

5055C	Eye drops 5 mg per mL (0.5%), 10 mL	±1	..	..	11.00	12.07	Chloromycetin Chlorsig	PF SI

**PREPARATIONS WHICH MAY BE PRESCRIBED BY PARTICIPATING
DENTAL PRACTITIONERS FOR DENTAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

Various

All other therapeutic products

All other therapeutic products

Antidotes

NALOXONE HYDROCHLORIDE							
5175J	Injection 2 mg in 5 mL	1	..	..	43.49	34.20	Naloxone Min-I-Jet CS

All other non-therapeutic products

All other non-therapeutic products

Solvents and diluting agents, incl. irrigating solutions

SODIUM CHLORIDE							
5211G	Injection 9 mg per mL (0.9%), 10 mL	5	..	..	16.48	17.55	Pfizer Australia Pty Ltd PF

PREPARATIONS WHICH MAY BE PRESCRIBED BY AUTHORISED OPTOMETRISTS FOR OPTOMETRICAL TREATMENT ONLY

From 1 January 2008, optometrists accredited to prescribe under State or Territory legislation can apply for approval as PBS prescribers (*authorised optometrists*). Information for optometrists on becoming a PBS prescriber is available on the Medicare Australia website at: www.medicareaustralia.gov.au

The medications listed in this section are for prescribing by authorised optometrists only. Optometrists must not write PBS prescriptions for medicines listed elsewhere in the PBS Schedule.

The Pharmaceutical Benefits Advisory Committee (PBAC) is responsible for making recommendations regarding preparations for inclusion in the optometrist section.

Some products are included in more than one section of the Schedule. For a prescription to be eligible for subsidy, prescribers must ensure that they prescribe under the PBS only those medicines, and in accordance with the restrictions, listed for their practitioner type. Listing details for the same product may differ between sections and different PBS item codes apply for each prescriber type.

Optometrist PBS prescriptions are identifiable by colour, and include the words 'PBS/RPBS optometrist'. Prescriptions must include the optometrist's PBS prescriber number. The same optometrist prescription form is used to prescribe unrestricted, restricted or authority items. Only one item is allowed per form. Optometrist PBS prescriptions may include repeats.

Regulation 24 does not apply for optometrist prescribing. An optometrist cannot direct that original and repeat supplies of pharmaceutical benefits be supplied at the one time.

Authority prescriptions: Authority prescriptions for authority required items, or for increased quantities or repeats, require prior approval from Medicare Australia or the DVA for each prescription. (Refer to details in the Explanatory Notes section Authority required PBS Prescriptions, for more information on authority prescriptions.) DVA approval for non-Schedule items is not available for optometrist prescribing.

RPBS: Optometrists approved as PBS prescribers may write prescriptions for supply under the RPBS. The list of optometrist medicines under the RPBS is the same as the PBS. There are no optometrist listings in the Repatriation Schedule for prescribing for veterans only. There is no provision for optometrist prescribers to request approval to prescribe items that are not included in the PBS optometrist list (non-Schedule items).

State and Territory requirements: Optometrists may prescribe medications as private prescriptions according to their State/Territory prescribing accreditation. The medicines which can be prescribed differ between States and Territories. It is the optometrist's responsibility to ensure adherence to State/Territory law for all prescriptions (PBS and private) and additionally to all PBS requirements for PBS/RPBS prescriptions.

GENERAL STATEMENT FOR TOPICAL ANTI-GLAUCOMA DRUGS PRESCRIBED BY AUTHORISED OPTOMETRISTS AS PHARMACEUTICAL BENEFITS

Use the following guidelines to determine patient eligibility for subsidisation under the PBS for the following drugs prescribed by authorised optometrists:

- Betaxolol hydrochloride
- Bimatoprost
- Brimonidine tartrate
- Brimonidine tartrate with timolol maleate
- Brinzolamide
- Dorzolamide hydrochloride
- Dorzolamide hydrochloride with timolol maleate
- Latanoprost
- Latanoprost with timolol maleate
- Pilocarpine hydrochloride
- Timolol maleate
- Travoprost
- Travoprost with timolol maleate

By writing a PBS prescription, the prescriber is certifying the criteria set out in these guidelines are satisfied, and use is in accordance with the registered indications – refer to the current Product Information for details.

GUIDELINES FOR SHARED CARE OF GLAUCOMA PATIENTS

Under these guidelines, authorised optometrists who are approved to use therapeutic drugs in their practices and who have adequate professional indemnity cover, will be able to co-manage glaucoma patients in a shared care arrangement with an ophthalmologist.

Initial Referral to Ophthalmologist

An authorised optometrist who makes a provisional diagnosis of glaucoma is to refer the patient to an ophthalmologist for confirmation of the diagnosis and the development of a management plan.

Where clinically important delays are expected before the patient's first review by an ophthalmologist, the optometrist should seek interim advice on the patient's management from the ophthalmologist by telephone (or alternate means).

The patient's consent is to be obtained by the ophthalmologist and optometrist for all aspects of the management plan, including the sharing of care between the two practitioners, and the communication of clinical information to the patient's nominated general practitioner.

Patients being considered for anti-glaucoma therapy with a beta blocking agent should be assessed for any potential cardiovascular or respiratory risk by a medical practitioner (often the patient's general practitioner), prior to initiating therapy. This assessment should be repeated if a change in dose of the beta blocker is proposed.

Once the diagnosis of glaucoma is confirmed by the ophthalmologist and a treatment plan is in place for the patient, the optometrist can perform ongoing reviews to monitor the patient and prescribe topical drugs under the PBS providing that:

- Periodic review demonstrates the treatment to be effective, and
- Changes to management are only initiated following consultation between treating practitioners.

Patient Management Plan

The management plan must be in writing and specify the following:

1. All the agreed components of treatment including any drug therapy;
2. Target pressures and action to be taken if these are not achieved within a specified time frame;
3. An agreed approach to monitoring visual fields and optic disc imaging and action to be taken following changes in visual fields;
4. Triggers for referral for more immediate ophthalmological and general practitioner review;
5. Likely side effects from agreed treatment and the action to be taken to address these;
6. An agreed schedule for patient review by both practitioners;
7. Who is responsible for performing each of the required tests and the required frequency for performing them;
8. An agreed method for timely communication of clinical findings and patient management between the two practitioners and the patient's nominated general practitioner.

Ophthalmologists must be available for consultation by the treating optometrist and for consultation by the patient where that consultation has been recommended or requested by the optometrist.

The involvement of a pharmacist to provide medicines information, advice relating to administration and techniques to limit systemic absorption and side effects of ophthalmic medications is recommended.

**PREPARATIONS WHICH MAY BE PRESCRIBED BY AUTHORISED
OPTOMETRISTS FOR OPTOMETRICAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$		

Sensory organs

Ophthalmologicals

Antiinfectives

Antibiotics

CHLORAMPHENICOL

5511C	Eye ointment 10 mg per g (1%), 4 g	£1	..	..	9.76	10.83	Chloromycetin	PF
							Chlorsig	SI
5512D	Eye drops 5 mg per mL (0.5%), 10 mL	£1	2	..	11.00	12.07	Chloromycetin	PF
							Chlorsig	SI

Sulfonamides

SULFACETAMIDE SODIUM

5530C	Eye drops 100 mg per mL (10%), 15 mL	£1	2	..	14.96	16.03	Bleph 10	AG

Antivirals

ACICLOVIR

Restricted benefit

Herpes simplex keratitis.

5501M	Eye ointment 30 mg per g (3%), 4.5 g	£1	..	..	35.06	34.20	Zovirax	GK

Antiinflammatory agents

Corticosteroids, plain

FLUOROMETHOLONE

Note

No applications for increased maximum quantities and/or repeats will be authorised.

5513E	Eye drops 1 mg per mL (0.1%), 5 mL	£1	..	..	10.61	11.68	Flucon	AQ
							FML Liquifilm	AG

FLUOROMETHOLONE ACETATE

Note

No applications for increased maximum quantities and/or repeats will be authorised.

5533F	Eye drops 1 mg per mL (0.1%), 5 mL	£1	..	..	10.61	11.68	Flarex	AQ

HYDROCORTISONE ACETATE

Note

No applications for increased maximum quantities and/or repeats will be authorised.

5515G	Eye ointment 5 mg per g (0.5%), 5 g	£1	..	..	11.57	12.64	Hycor	SI
5516H	Eye ointment 10 mg per g (1%), 5 g	£1	..	..	12.81	13.88	Hycor	SI

Antiinflammatory agents, non-steroids

FLURBIPROFEN SODIUM

5514F	Eye drops 300 micrograms per mL (0.03%), single dose units 0.4 mL, 5	1	..	..	15.37	16.44	Ocufen	AG

Antiglaucoma preparations and miotics

Sympathomimetics in glaucoma therapy

BRIMONIDINE TARTRATE

5534G	Eye drops 2 mg per mL (0.2%), 5 mL	£1	5	..	20.42	21.49 ^a	Enidin	PE
				^B 1.66	22.08	21.49 ^a	Alphagan	AG

**PREPARATIONS WHICH MAY BE PRESCRIBED BY AUTHORISED
OPTOMETRISTS FOR OPTOMETRICAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
5563T	Eye drops 1.5 mg per mL (0.15%), 5 mL	1	5	..	20.42	21.49	Alphagan P 1.5	AG

BRIMONIDINE TARTRATE with TIMOLOL MALEATE

Restricted benefit

Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;

Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.

5535H	Eye drops 2 mg-5 mg (base) per mL (0.2%-0.5%), 5 mL	1	5	..	26.61	27.68	Combigan	AG

Parasympathomimetics

PILOCARPINE HYDROCHLORIDE

5536J	Eye drops 10 mg per mL (1%), 15 mL	1	5	..	12.65	13.72	Isopto Carpine	AQ
5537K	Eye drops 20 mg per mL (2%), 15 mL	1	5	..	13.93	15.00	Isopto Carpine	AQ
5538L	Eye drops 40 mg per mL (4%), 15 mL	1	5	..	16.84	17.91	Isopto Carpine	AQ

Carbonic anhydrase inhibitors

BRINZOLAMIDE

5540N	Eye drops 10 mg per mL (1%), 5 mL	1	5	..	23.11	24.18 ^a	BrinzoQuin	IQ
				^B 1.18	24.29	24.18 ^a	Azopt	AQ

BRINZOLAMIDE with TIMOLOL MALEATE

Restricted benefit

Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;

Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.

5562R	Eye drops 10 mg-5 mg (base) per mL (1%-0.5%), 5 mL	1	5	..	27.49	28.56	Azarga	AQ

DORZOLAMIDE HYDROCHLORIDE

5541P	Eye drops 20 mg (base) per mL (2%), 5 mL	1	5	..	21.29	22.36	Trusopt	MK

DORZOLAMIDE HYDROCHLORIDE with TIMOLOL MALEATE

Restricted benefit

Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;

Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.

5542Q	Eye drops 20 mg (base)-5 mg (base) per mL (2%-0.5%), 5 mL	1	5	..	27.49	28.56	Cosopt	MK

Beta blocking agents

BETAXOLOL HYDROCHLORIDE

5543R	Eye drops, suspension, 2.5 mg (base) per mL (0.25%), 5 mL	1	5	..	14.94	16.01	Betoptic S	AQ
5544T	Eye drops, solution, 5 mg (base) per mL (0.5%), 5 mL	1	5	..	14.94	16.01 ^a	BetoQuin	IQ
				^B 2.09	17.03	16.01 ^a	Betoptic	AQ

TIMOLOL MALEATE

5546X	Eye gel 1 mg (base) per g (0.1%), 5 g	1	5	..	13.22	14.29	Nyogel	NV
5547Y	Eye drops 2.5 mg (base) per mL (0.25%), 5 mL	1	5	..	11.80	12.87 ^a	Tenopt	SI
				^B 3.20	15.00	12.87 ^a	Timoptol	FR
5548B	Eye drops 5 mg (base) per mL (0.5%), 5 mL	1	5	..	12.62	13.69 ^a	Tenopt	SI
				^B 3.20	15.82	13.69 ^a	Timoptol	FR
5549C	Eye drops (gellan gum solution) 2.5 mg (base) per mL (0.25%), 2.5 mL	1	5	..	11.80	12.87	Timoptol XE	MK
5550D	Eye drops (gellan gum solution) 5 mg (base) per mL (0.5%), 2.5 mL	1	5	..	12.62	13.69	Timoptol XE	MK

**PREPARATIONS WHICH MAY BE PRESCRIBED BY AUTHORISED
OPTOMETRISTS FOR OPTOMETRICAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Prostaglandin analogues								
BIMATOPROST								
5551E	Eye drops 300 micrograms per mL (0.03%), 3 mL	1	5	..	42.14	34.20	Lumigan	AG
BIMATOPROST with TIMOLOL MALEATE								
<u>Restricted benefit</u>								
Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;								
Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.								
5558M	Eye drops 300 micrograms-5 mg (base) per mL (0.03%-0.5%), 3 mL	1	5	..	46.82	34.20	Ganfort 0.3/5	AG
LATANOPROST								
5552F	Eye drops 50 micrograms per mL (0.005%), 2.5 mL	1	5	..	42.14	34.20	Xalatan	PF
LATANOPROST with TIMOLOL MALEATE								
<u>Restricted benefit</u>								
Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;								
Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.								
5553G	Eye drops 50 micrograms-5 mg (base) per mL (0.005%-0.5%), 2.5 mL	1	5	..	46.82	34.20	Xalacom	PF
TRAVOPROST								
5554H	Eye drops 40 micrograms per mL (0.004%), 2.5 mL	1	5	..	42.14	34.20	Travatan	AQ
TRAVOPROST with TIMOLOL MALEATE								
<u>Restricted benefit</u>								
Reduction of elevated intra-ocular pressure in a patient with open-angle glaucoma that is not adequately controlled with monotherapy;								
Reduction of elevated intra-ocular pressure in a patient with ocular hypertension that is not adequately controlled with monotherapy.								
5555J	Eye drops 40 micrograms-5 mg (base) per mL (0.004%-0.5%), 2.5 mL	1	5	..	46.82	34.20	Duotrav	AQ
Decongestants and antiallergics								
<i>Other antiallergics</i>								
SODIUM CROMOGLYCATE								
<u>Restricted benefit</u>								
Vernal kerato-conjunctivitis.								
5529B	Eye drops 20 mg per mL (2%), 10 mL	1	5	..	14.37	15.44	^a Cromolux ^a Opticrom	AE SW
Other ophthalmologicals								
<i>Other ophthalmologicals</i>								
CARBOMER								
<u>Restricted benefit</u>								
Severe dry eye syndrome, including Sjogren's syndrome.								
5503P	Eye gel 2 mg per g (0.2%), 10 g	1	5	..	10.35	11.42	GelTears	BU
							^a PAA	NM
				^B 0.96	11.31	11.42	^a Viscotears	NV

**PREPARATIONS WHICH MAY BE PRESCRIBED BY AUTHORISED
OPTOMETRISTS FOR OPTOMETRICAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
CARBOMER								
<u>Authority required</u>								
Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.								
5504Q	Eye gel 2 mg per g (0.2%), single dose units 0.6 mL, 30	3	5	..	*36.72	34.20	Viscotears Gel PF	NV
CARBOMER 974								
<u>Authority required</u>								
Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.								
5502N	Ocular lubricating gel 3 mg per g (0.3%), single dose units 0.5 g, 30	3	5	..	*36.06	34.20	Poly Gel	AQ
CARMELLOSE SODIUM								
<u>Restricted benefit</u>								
Severe dry eye syndrome, including Sjogren's syndrome.								
5507W	Eye drops 5 mg per mL (0.5%), 15 mL	‡1	5	..	10.59	11.66	Refresh Tears Plus	AG
5508X	Eye drops 10 mg per mL (1%), 15 mL	‡1	5	..	10.59	11.66	Refresh Liquigel	AG
CARMELLOSE SODIUM								
<u>Authority required</u>								
Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.								
5505R	Eye drops 10 mg per mL (1%), single dose units 0.4 mL, 30	3	5	..	*36.06	34.20	Celluvisc	AG
5506T	Eye drops 5 mg per mL (0.5%), single dose units 0.4 mL, 30	3	5	..	*36.06	34.20	Cellufresh	AG
5509Y	Eye drops 2.5 mg per mL (0.25%), single dose units 0.6 mL, 24	4	5	..	*40.42	34.20	TheraTears	CX
5510B	Ocular lubricating gel 10 mg per mL (1%), single dose units 0.6 mL, 28	3	5	..	*34.08	34.20	TheraTears	CX
CARMELLOSE SODIUM with GLYCERIN								
<u>Restricted benefit</u>								
Severe dry eye syndrome, including Sjogren's syndrome.								
<u>Note</u>								
The in-use shelf life of Optive is 6 months from the date of opening.								
5556K	Eye drops 5 mg-9 mg per mL (0.5%-0.9%), 15 mL	‡1	3	..	10.59	11.66	Optive	AG
CARMELLOSE SODIUM with GLYCERIN								
<u>Authority required</u>								
Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.								
5561Q	Eye drops 5 mg-9 mg per mL (0.5%-0.9%), single dose units 0.4 mL, 30	3	5	..	*36.06	34.20	Optive	AG
HYPROMELLOSE								
<u>Restricted benefit</u>								
Severe dry eye syndrome, including Sjogren's syndrome.								
5517J	Eye drops 5 mg per mL (0.5%), 15 mL	‡1	5	..	10.35	11.42	Methopt	SI
5518K	Eye drops 3 mg per mL (0.3%), 15 mL (contains sodium perborate as preservative)	‡1	5	..	10.35	11.42 ^a	In a Wink Moisturising	NM
				^B 1.79	12.14	11.42 ^a	Genteal	NV
HYPROMELLOSE with CARBOMER 980								
<u>Restricted benefit</u>								
Severe dry eye syndrome, including Sjogren's syndrome.								

Severe dry eye syndrome, including Sjogren's syndrome.

**PREPARATIONS WHICH MAY BE PRESCRIBED BY AUTHORISED
OPTOMETRISTS FOR OPTOMETRICAL TREATMENT ONLY**

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
5525T	Eye drops 30 mg per mL (3%), 15 mL	‡1	5	.. ^B 5.71	10.35 16.06	11.42 11.42	^a PVA Forte ^a Liquifilm Forte	PE AG
5526W	Eye drops 14 mg per mL (1.4%), 15 mL	‡1	5	.. ^B 1.64	10.35 11.99	11.42 11.42	^a PVA Tears ^a Liquifilm Tears	PE AG
5527X	Eye drops 14 mg per mL (1.4%), 15 mL (contains sodium chlorite/hydrogen peroxide as preservative)	‡1	5	..	10.35	11.42	Vistil	AE
5528Y	Eye drops 30 mg per mL (3%), 15 mL (contains sodium chlorite/hydrogen peroxide as preservative)	‡1	5	..	10.35	11.42	Vistil Forte	AE

SOY LECITHIN**Authority required**

Severe dry eye syndrome in patients who are sensitive to preservatives in multi-dose eye drops.

5545W	Eye spray 10 mg per mL (1%), 10 mL	2	5	..	*36.06	34.20	tearsagain	RB

Ophthalmological and otological preparations

Antiinfectives***Antiinfectives*****FRAMYCETIN SULFATE**

5557L	Eye and ear drops 5 mg per mL (0.5%), 8 mL	‡1	2	..	10.11	11.18	Soframycin	SW

Section 100 – Items Available under Special Arrangement

In addition to the drugs and medicinal preparations available under normal PBS arrangements listed in this Schedule, a number of drugs are also available as pharmaceutical benefits but are distributed under alternative arrangements where these are considered more appropriate.

These alternative arrangements are provided for under section 100 of the National Health Act 1953. Several programs exist for the provision of drugs as pharmaceutical benefits in this way and this section lists those drugs which are available under the following programs:

- **Highly Specialised Drugs Program**
- **Botulinum Toxin Program**
- **Human Growth Hormone Program**
- **IVF/GIFT Program**
- **Opiate Dependence Treatment Program**
- **Special Authority Program (Public)**
- **Special Authority Program (Private)**

Complete details concerning the availability of drugs as benefits under these programs may be obtained by telephoning the relevant contact number(s) shown in each section, or in certain cases, by referring to the telephone number provided for individual drugs listings.

Section 100 – Highly Specialised Drugs Program

The Australian Government provides funding for certain specialised medications under the Highly Specialised Drugs Program. Highly Specialised Drugs are medicines for the treatment of chronic conditions which, because of their clinical use or other special features, are restricted to supply through public and private hospitals having access to appropriate specialist facilities. To prescribe these drugs as pharmaceutical benefit items, medical practitioners are required to be affiliated with these specialist hospital units. A general practitioner or non-specialist hospital doctor may only prescribe Highly Specialised Drugs to provide maintenance therapy under the guidance of the treating specialist.

Benefits are available for the listed clinical indications only. There is no facility for individual patient approval for indications outside those listed.

To gain access to a Commonwealth funded drug under this program, a patient must attend a participating hospital and be a day admitted patient, a non-admitted patient or a patient on discharge, be under appropriate specialist medical care, meet the specific medical criteria and be an Australian resident in Australia (or other eligible person).

A patient will be required to pay a contribution for each supply of a highly specialised drug at a similar rate to the Pharmaceutical Benefits Scheme. Commonwealth subsidy is not available for hospital in-patients.

Reciprocal Health Care Agreement – Where a patient is entitled to be treated as an eligible person as a visitor from a country with which Australia has entered into a Reciprocal Health Care Agreement, the supply will be limited to the original prescription only. Repeat prescriptions for these patients are not permitted.

Private Hospitals – **In addition to the above requirements**, for Highly Specialised Drugs prescribed through private hospitals, claiming and approval of authority prescriptions is administered by Medicare Australia. Highly Specialised Drugs are authority required items. Medical practitioners must seek approval to prescribe these items as pharmaceutical benefits prior to their dispensing under the PBS. Approval of authority prescriptions by Medicare Australia may be obtained either by posting an Authority Prescription Form to Medicare Australia, or by using Medicare Australia's Authority Freecall service (1800 888 333). **Prescribers must quote the provider number of the hospital when applying.** Not more than two months' supply (one month's supply in the case of Clozapine), with provision for up to 5 repeats, will be authorised. Prescriptions for Highly Specialised Drugs can be dispensed by an approved private hospital's dispensary or by a community pharmacy.

The remuneration rates for Highly Specialised Drugs prescribed through private hospitals comprise the normal PBS ready-prepared dispensing fee plus a mark-up ascertained as follows:

- 10% for drugs with a price ex-manufacturer of less than \$40;
- \$4 for drugs with a price ex-manufacturer of between \$40 and \$100;
- 4% for drugs with a price ex-manufacturer of between \$100.01 and \$1000;
- \$40 for drugs with a price ex-manufacturer of greater than \$1000.

Public Hospitals – For Highly Specialised Drugs prescribed through public hospitals, claiming and access to the program is administered by the States/Territories Health Departments. Prescriptions for Highly Specialised Drugs can be dispensed by public hospital pharmacies.

If you would like further information about the Highly Specialised Drugs Program, please contact your pharmacy, Medicare Australia (Ph: 132 290) or the Australian Government adviser, the Highly Specialised Drugs Working Party Secretariat (Ph: (02) 6289 2331).

- Highly Specialised Drugs Program - Summary of Changes
- Highly Specialised Drugs Program - (Public Hospital)
- Highly Specialised Drugs Program - (Private Hospital)

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Blood and blood forming organs

Antianemic preparations

Other antianemic preparations

Other antianemic preparations

DARBEPOETIN ALFA

Authority required

Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.

6320P	Injection 10 micrograms in 0.4 mL pre-filled syringe	8	5	..	*376.74	Aranesp	AN
6321Q	Injection 20 micrograms in 0.5 mL pre-filled syringe	8	5	..	*703.86	Aranesp	AN
6322R	Injection 30 micrograms in 0.3 mL pre-filled syringe	8	5	..	*960.58	Aranesp	AN
6323T	Injection 40 micrograms in 0.4 mL pre-filled syringe	8	5	..	*1160.02	Aranesp	AN
6324W	Injection 50 micrograms in 0.5 mL pre-filled syringe	8	5	..	*1423.20	Aranesp	AN
6325X	Injection 60 micrograms in 0.3 mL pre-filled syringe	8	5	..	*1663.08	Aranesp	AN
6326Y	Injection 100 micrograms in 0.5 mL pre-filled syringe	8	5	..	*2666.92	Aranesp	AN
6365B	Injection 150 micrograms in 0.3 mL pre-filled syringe	8	5	..	*3950.92	Aranesp	AN
6438W	Injection 80 micrograms in 0.4 mL pre-filled syringe	8	5	..	*2174.42	Aranesp	AN
6488L	Injection 20 micrograms in 0.5 mL pre-filled injection pen	8	5	..	*703.86	Aranesp SureClick	AN
6489M	Injection 40 micrograms in 0.4 mL pre-filled injection pen	8	5	..	*1160.02	Aranesp SureClick	AN
6490N	Injection 60 micrograms in 0.3 mL pre-filled injection pen	8	5	..	*1663.06	Aranesp SureClick	AN
6491P	Injection 80 micrograms in 0.4 mL pre-filled injection pen	8	5	..	*2174.42	Aranesp SureClick	AN
6492Q	Injection 100 micrograms in 0.5 mL pre-filled injection pen	8	5	..	*2666.90	Aranesp SureClick	AN
6493R	Injection 150 micrograms in 0.3 mL pre-filled injection pen	8	5	..	*3950.90	Aranesp SureClick	AN

EPOETIN ALFA

Authority required

Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.

6204M	Injection 2,000 units in 0.5 mL pre-filled syringe	12	5	..	*543.90	Eprex 2000	JC
6205N	Injection 3,000 units in 0.3 mL pre-filled syringe	12	5	..	*700.00	Eprex 3000	JC
6206P	Injection 4,000 units in 0.4 mL pre-filled syringe	12	5	..	*889.70	Eprex 4000	JC
6207Q	Injection 10,000 units in 1 mL pre-filled syringe	12	5	..	*2016.72	Eprex 10000	JC
6251B	Injection 1,000 units in 0.5 mL pre-filled syringe	12	5	..	*296.90	Eprex 1000	JC
6302Q	Injection 5,000 units in 0.5 mL pre-filled syringe	12	5	..	*1103.76	Eprex 5000	JC
6303R	Injection 6,000 units in 0.6 mL pre-filled syringe	12	5	..	*1301.56	Eprex 6000	JC
6305W	Injection 8,000 units in 0.8 mL pre-filled syringe	12	5	..	*1674.34	Eprex 8000	JC
6339P	Injection 40,000 units in 1 mL pre-filled syringe	2	5	..	*1300.42	Eprex 40,000	JC
6434P	Injection 20,000 units in 0.5 mL pre-filled syringe	12	5	..	*3922.42	Eprex 20,000	JC
9623L	Injection 30,000 units in 0.75 mL pre-filled syringe	12	5	..	*5774.92	Eprex 30,000	JC

EPOETIN BETA

Authority required

Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.

6480C	Injection 2,000 units in 0.3 mL pre-filled syringe	12	5	..	*543.90	NeoRecormon	RO
6481D	Injection 3,000 units in 0.3 mL pre-filled syringe	12	5	..	*700.00	NeoRecormon	RO
6482E	Injection 4,000 units in 0.3 mL pre-filled syringe	12	5	..	*889.70	NeoRecormon	RO
6483F	Injection 5,000 units in 0.3 mL pre-filled syringe	12	5	..	*1103.78	NeoRecormon	RO

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
6484G	Injection 6,000 units in 0.3 mL pre-filled syringe	12	5	..	*1301.56		NeoRecormon RO
6485H	Injection 10,000 units in 0.6 mL pre-filled syringe	12	5	..	*2016.72		NeoRecormon RO
6486J	Injection 20,000 units in 0.6 mL pre-filled syringe	12	5	..	*3922.42		NeoRecormon RO

EPOETIN LAMBDA

Note

Epoetin lambda should only be administered by the intravenous route.

Authority required

Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.

9588P	Injection 5,000 units in 0.5 mL pre-filled syringe	12	5	..	*1048.12		Novicrit NV
9590R	Injection 6,000 units in 0.6 mL pre-filled syringe	12	5	..	*1235.50		Novicrit NV
9593X	Injection 8,000 units in 0.8 mL pre-filled syringe	12	5	..	*1588.66		Novicrit NV
9595B	Injection 10,000 units in 1 mL pre-filled syringe	12	5	..	*1913.02		Novicrit NV
9685R	Injection 1,000 units in 0.5 mL pre-filled syringe	12	5	..	*281.60		Novicrit NV
9686T	Injection 2,000 units in 1 mL pre-filled syringe	12	5	..	*515.60		Novicrit NV
9687W	Injection 3,000 units in 0.3 mL pre-filled syringe	12	5	..	*663.50		Novicrit NV
9688X	Injection 4,000 units in 0.4 mL pre-filled syringe	12	5	..	*843.20		Novicrit NV

METHOXY POLYETHYLENE GLYCOL-EPOETIN BETA

Authority required

Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.

9574X	Injection 30 micrograms in 0.3 mL pre-filled syringe	2	5	..	*390.36		Mircera RO
9575Y	Injection 50 micrograms in 0.3 mL pre-filled syringe	2	5	..	*646.34		Mircera RO
9576B	Injection 75 micrograms in 0.3 mL pre-filled syringe	2	5	..	*938.28		Mircera RO
9577C	Injection 100 micrograms in 0.3 mL pre-filled syringe	2	5	..	*1205.24		Mircera RO
9578D	Injection 120 micrograms in 0.3 mL pre-filled syringe	2	5	..	*1388.06		Mircera RO
9579E	Injection 200 micrograms in 0.3 mL pre-filled syringe	2	5	..	*1970.72		Mircera RO
9580F	Injection 360 micrograms in 0.6 mL pre-filled syringe	2	5	..	*3372.94		Mircera RO

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Cardiovascular system

Antihypertensives

Other antihypertensives

Other antihypertensives

AMBRISENTAN

Caution

Ambrisentan is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of treatment with this drug.

Note

Any queries concerning the arrangements to prescribe ambrisentan may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

subsidised treatment with an alternate PAH agent; AND

— primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND

(c) epoprostenol sodium, of:

— primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-

subsidised treatment with an alternate PAH agent; AND

— primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND

(d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND

(e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND

(f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (5) RHC composite assessment only;
(6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent. All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with ambrisentan of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with ambrisentan of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (c) WHO Functional Class IV primary pulmonary hypertension; OR
- (d) WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (i) RHC composite assessment; and
- (ii) ECHO composite assessment; and
- (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (grandfather patients)

Application for initial PBS-subsidised treatment with ambrisentan of patients who were receiving treatment with ambrisentan prior to 1 December 2009 and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease; OR
- (c) WHO Functional Class IV primary pulmonary hypertension; OR
- (d) WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) for patients who have received less than 6 months of ambrisentan treatment at the time of application — a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results of the following 3 tests, where available, at the time treatment with ambrisentan was commenced:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) the date of commencement of ambrisentan treatment; and
- (4) a signed patient acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. The number of repeats authorised will be dependent on the duration of prior ambrisentan therapy. Where patients have received less than 6 months of non-PBS-subsidised treatment with ambrisentan, sufficient repeats to allow the patient to complete a total of 6 months of combined PBS-subsidised and non-PBS-subsidised therapy may be requested. Where fewer than the maximum allowable number of repeats are requested at the time of application, authority approvals for the remainder of the allowable repeats may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients)

Application for initial treatment with ambrisentan of patients with one of the following:

- (a) primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who wish to re-commence PBS-subsidised ambrisentan after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with ambrisentan; OR
- (b) primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with an alternate PAH agent other than ambrisentan.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with ambrisentan of patients who have received approval for initial PBS-subsidised treatment with ambrisentan and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of ambrisentan treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats will be authorised. Where fewer than 5 repeats are initially requested under this criterion, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note

Special Pricing Arrangements apply.

9648T	Tablet 5 mg	30	..	..	4081.42	Volibris	GK
9649W	Tablet 10 mg	30	..	..	4081.42	Volibris	GK

BOSENTAN MONOHYDRATE

Caution

Bosentan monohydrate is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of treatment with this drug.

Note

Any queries concerning the arrangements to prescribe bosentan monohydrate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
- primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

- (a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

- (b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

- (a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent.

All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new adult patients)

Application for initial PBS-subsidised treatment with bosentan monohydrate of adult patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to scleroderma and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new adult patients)

Application for initial PBS-subsidised treatment with bosentan monohydrate of adult patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to scleroderma and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (c) WHO Functional Class IV primary pulmonary hypertension; OR
- (d) WHO Functional Class IV pulmonary arterial hypertension secondary to scleroderma.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients under 18 years of age)

Application for initial PBS-subsidised treatment with bosentan monohydrate of patients aged less than 18 years who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

WHO Functional Class III primary pulmonary hypertension and either a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate prior vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a patient and prescriber acknowledgment, signed by the parent or authorised guardian, indicating that they understand and acknowledge that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients under 18 years of age)

Application for initial PBS-subsidised treatment with bosentan monohydrate of patients aged less than 18 years who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and either a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class IV primary pulmonary hypertension.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a patient and prescriber acknowledgment, signed by the parent or authorised guardian, indicating that they understand and acknowledge that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with bosentan monohydrate of a patient who has been assessed by a physician from a designated hospital to have WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment (and signed by the parent or authorised guardian for patients under 18 years of age) indicating that the patient understands and acknowledges that PBS-subsidised treatment with bosentan monohydrate will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for adult patients)

Application for initial treatment with bosentan monohydrate of adult patients with one of the following:

- (a) primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), who wish to re-commence PBS-subsidised bosentan monohydrate after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with bosentan monohydrate; OR
- (b) primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma and whose most recent course of PBS-subsidised

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

treatment was with an alternate PAH agent other than bosentan monohydrate.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for patients under 18 years of age)

Application for initial treatment with bosentan monohydrate of patients aged less than 18 years with one of the following:

- (a) primary pulmonary hypertension, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), who wish to re-commence PBS-subsidised bosentan monohydrate after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with bosentan monohydrate; OR
- (b) primary pulmonary hypertension and whose most recent course of PBS-subsidised treatment was with a PAH agent other than bosentan monohydrate.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with bosentan monohydrate of patients who have received approval for initial PBS-subsidised treatment with bosentan monohydrate and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of bosentan monohydrate treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats will be authorised.

Where fewer than 5 repeats are initially requested under this criterion, authority approvals for sufficient repeats to complete a maximum of 6

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Cessation of treatment (all patients)

Final PBS-subsidised supply for patients with WHO Functional Class III or IV primary pulmonary hypertension or WHO Functional Class III or IV pulmonary arterial hypertension secondary to scleroderma or WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), who have not responded to bosentan monohydrate therapy [see Note for definition of response], to allow for gradual cessation of treatment.

Applications for authorisation under this criterion should be made on the telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) [see Note on authority approval requirements].

Approval will only be granted for the 62.5 mg tablet strength. The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment.

Under no circumstances will telephone approvals be granted for treatment that would extend the final treatment period beyond 1 month.

Note

Special Pricing Arrangements apply.

6429J	Tablet 62.5 mg (base)	60	..	..	4081.42	Tracleer	AT
6430K	Tablet 125 mg (base)	60	..	..	4081.42	Tracleer	AT

EPOPROSTENOL SODIUM

Note

Any queries concerning the arrangements to prescribe epoprostenol sodium may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

(a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND

(b) iloprost trometamol, of:

— primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND

— primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND

— drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND

(c) epoprostenol sodium, of:

— primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND

— primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND

(d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND

(e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND

(f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent.

All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new adult patients)

Application for initial PBS-subsidised treatment with epoprostenol sodium of adult patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:
WHO Functional Class IV primary pulmonary hypertension.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients under 18 years of age)

Application for initial PBS-subsidised treatment with epoprostenol sodium of patients aged less than 18 years who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:
WHO Functional Class IV primary pulmonary hypertension.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a patient acknowledgment, signed by the parent or authorised guardian and the prescriber, indicating that they understand and acknowledge

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

that PBS-subsidised treatment with PAH agents will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats will be authorised under this criterion. Where fewer than 5 repeats are initially requested, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all adult patients)

Application for initial PBS-subsidised treatment with epoprostenol sodium of adult patients with one of the following:

- (a) primary pulmonary hypertension who wish to re-commence PBS-subsidised epoprostenol sodium after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with epoprostenol sodium; OR
- (b) WHO Functional Class IV primary pulmonary hypertension and who have received prior treatment with a PBS-subsidised PAH agent other than epoprostenol sodium; OR
- (c) WHO Functional Class III primary pulmonary hypertension and who have failed to respond to a prior PBS-subsidised PAH agent.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (5) for WHO Functional Class III patients, where this is the first application for epoprostenol sodium, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients under 18 years of age)

Application for initial PBS-subsidised treatment with epoprostenol sodium of patients aged less than 18 years with one of the following:

- (a) primary pulmonary hypertension who wish to re-commence PBS-subsidised epoprostenol sodium after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with epoprostenol sodium; OR
- (b) WHO Functional Class IV primary pulmonary hypertension and who have received prior treatment with a PBS-subsidised PAH agent other than epoprostenol sodium; OR
- (c) WHO Functional Class III primary pulmonary hypertension and who have failed to respond to a prior PBS-subsidised PAH agent.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (5) for WHO Functional Class III patients, where this is the first application for epoprostenol sodium, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with epoprostenol sodium of patients who have received approval for initial PBS-subsidised treatment with epoprostenol sodium, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of epoprostenol sodium treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
(1) a completed authority prescription form; and (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available: (i) RHC composite assessment; and (ii) ECHO composite assessment; and (iii) 6MWT. The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application. The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).						
6477X	Powder for I.V. infusion 500 micrograms (base) with diluent	1	..	..	52.11	Flolan GK
6478Y	Powder for I.V. infusion 1.5 mg (base) with diluent	1	..	..	93.79	Flolan GK

ILOPROST TROMETAMOL

Note

Any queries concerning the arrangements to prescribe iloprost trometamol may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent. All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with iloprost trometamol of patients who have not received prior PBS-subsidised treatment with iloprost and who have been assessed by a physician from a designated hospital to have:

WHO Functional Class III drug-induced pulmonary arterial hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with iloprost trometamol of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III drug-induced pulmonary arterial hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class IV primary pulmonary hypertension; OR
- (c) WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- (d) WHO Functional Class IV drug-induced pulmonary arterial hypertension.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients)

Application for initial PBS-subsidised treatment with iloprost trometamol of patients with one of the following:

- (a) primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who wish to re-commence PBS-subsidised iloprost trometamol after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with iloprost trometamol; OR
- (b) WHO Functional Class IV primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease and who have received prior treatment with a PBS-subsidised PAH agent other than iloprost trometamol; OR
- (c) WHO Functional Class III primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease and who have failed to respond to a prior PBS-subsidised PAH agent.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (5) for WHO Functional Class III patients, where this is the first application for iloprost trometamol, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with iloprost trometamol of patients who have received approval for initial PBS-subsidised treatment with iloprost trometamol, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of iloprost trometamol treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note

Special Pricing Arrangements apply.

6456T	Solution for inhalation 20 micrograms (base) in 2 mL	30	..	..	1122.42	Ventavis	SC

SILDENAFIL CITRATE

Note

Any queries concerning the arrangements to prescribe sildenafil citrate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent. All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with sildenafil citrate of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with sildenafil citrate of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients)

Application for initial PBS-subsidised treatment with sildenafil citrate of patients with one of the following:

- (a) WHO Functional Class III primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who wish to re-commence PBS-subsidised sildenafil citrate after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with sildenafil citrate; OR
- (b) WHO Functional Class III primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with a PAH agent other than sildenafil citrate.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with sildenafil citrate of patients who have received approval for initial PBS-subsidised treatment with sildenafil citrate, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of sildenafil citrate treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

9605M	Tablet 20 mg (base)	90	..	..	940.79	Revatio	PF

SITAXENTAN SODIUM

Caution

Sitaxentan sodium is a category X drug and must not be given to pregnant women. Pregnancy must be excluded before the start of treatment and avoided during treatment with this drug.

Note

Any queries concerning the arrangements to prescribe sitaxentan sodium may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent. All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with sitaxentan sodium of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with sitaxentan sodium of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients)

Application for initial PBS-subsidised treatment with sitaxentan sodium of patients with one of the following:

- (a) WHO Functional Class III primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who wish to re-commence PBS-subsidised sitaxentan sodium after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with sitaxentan sodium; OR

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(b) WHO Functional Class III primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with a PAH agent other than sitaxentan sodium.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with sitaxentan sodium of patients who have received approval for initial PBS-subsidised treatment with sitaxentan sodium, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of sitaxentan sodium treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

9622K	Tablet 100 mg	30	..	..	2790.22	Thelin	PF

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Systemic hormonal preparations, excl. sex hormones and insulins

Pituitary and hypothalamic hormones and analogues

Hypothalamic hormones

Antigrowth hormone

LANREOTIDE ACETATE

Authority required

Active acromegaly in a patient with persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre AND

- (a) after failure of other therapy including dopamine agonists; or
- (b) as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; or
- (c) if the patient is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated.

In a patient treated with radiotherapy, treatment must cease if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (6 weeks after the last dose). Lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Treatment must cease if IGF1 is not lower after 3 months treatment.

6332G	Powder for suspension for injection 30 mg (base) with diluent ampoule	2	11	..	*1546.42	Somatuline LA	IS

LANREOTIDE ACETATE

Authority required

Active acromegaly in a patient with persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre AND

- (a) after failure of other therapy including dopamine agonists; or
- (b) as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; or
- (c) if the patient is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated.

In a patient treated with radiotherapy, treatment must cease if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose). Lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Treatment must cease if IGF1 is not lower after 3 months treatment;

Functional carcinoid tumour causing intractable symptoms. The patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, and surgery or antineoplastic therapy must have failed or be inappropriate.

Treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 120 mg every 28 days. Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

6423C	Injection 60 mg (base) in single dose pre-filled syringe	2	11	..	*2736.42	Somatuline Autogel	IS
6424D	Injection 90 mg (base) in single dose pre-filled syringe	2	11	..	*3626.42	Somatuline Autogel	IS
6425E	Injection 120 mg (base) in single dose pre-filled syringe	2	11	..	*4526.42	Somatuline Autogel	IS

OCTREOTIDE

Authority required

Active acromegaly in a patient with persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre AND

- (a) after failure of other therapy including dopamine agonists; or
- (b) as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; or
- (c) if the patient is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated.

In a patient treated with radiotherapy, treatment must cease if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks. Octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Treatment must cease if IGF1 is not lower after 3 months treatment at a dose of 100 micrograms 3 times daily;

Functional carcinoid tumour or vasoactive intestinal peptide secreting tumour (VIPoma) causing intractable symptoms. The patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, and surgery or antineoplastic therapy must have failed or be inappropriate.

Treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months' therapy. Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

6227R	Injection 50 micrograms (as acetate) in 1 mL	90	11	..	*663.24 ^a	Hospira Pty Limited	HH

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
6228T	Injection 100 micrograms (as acetate) in 1 mL	90	11	..	*1309.44	^a	Octreotide MaxRx XF
						^a	Sandostatin 0.05 NV
						^a	Hospira Pty Limited HH
						^a	Octreotide MaxRx XF
6229W	Injection 500 micrograms (as acetate) in 1 mL	90	11	..	*6368.70	^a	Sandostatin 0.1 NV
						^a	Hospira Pty Limited HH
						^a	Octreotide MaxRx XF
						^a	Sandostatin 0.5 NV

OCTREOTIDE

Authority required

Acromegaly in a patient controlled on Sandostatin subcutaneous injections.

In a patient treated with radiotherapy, treatment must cease if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose). Octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Treatment must cease if IGF1 is not lower after 3 months of treatment;

Functional carcinoid tumour or vasoactive intestinal peptide secreting tumour (VIPoma) with symptom control on Sandostatin subcutaneous injections.

Treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with Sandostatin subcutaneous injections. Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

6426F	Injection (modified release) 10 mg (as acetate), vial and diluent syringe	1	11	..	1381.38	Sandostatin LAR	NV
6427G	Injection (modified release) 20 mg (as acetate), vial and diluent syringe	1	11	..	1823.16	Sandostatin LAR	NV
6428H	Injection (modified release) 30 mg (as acetate), vial and diluent syringe	1	11	..	2269.75	Sandostatin LAR	NV

Calcium homeostasis

Anti-parathyroid agents

Other anti-parathyroid agents

CINACALCET HYDROCHLORIDE

Authority required

Management, including initiation and stabilisation, by a nephrologist, of a patient with chronic kidney disease on dialysis who has sustained secondary hyperparathyroidism with iPTH of at least 50 pmol per L, not responding to conventional therapy.

Note

During the titration phase, intact PTH should be monitored 4 weekly (measured at least 12 hours post dose) and dose titrated until an appropriate iPTH concentration is achieved. During the titration phase, approval will be limited to sufficient supply for 4 weeks treatment at a time, with doses between 30 and 180 mg per day according to the patient's response and tolerability.

During the maintenance phase, approval will be limited to provide sufficient quantity for 4 weeks treatment up to a maximum of 6 months supply for doses between 30 and 180 mg per day according to the patient's response and tolerability. Intact PTH should be monitored quarterly (measured at least 12 hours post dose) and dose adjusted as necessary to maintain an appropriate iPTH concentration.

"Sustained" means the abnormality was detected on at least 2 blood samples collected over a period of 2 to 4 months.

Authority required

Management, including initiation and stabilisation, by a nephrologist, of a patient with chronic kidney disease on dialysis who has sustained secondary hyperparathyroidism with iPTH of at least 15 pmol per L and less than 50 pmol per L AND an (adjusted) serum calcium concentration at least 2.6 mmol per L, not responding to conventional treatment.

Note

During the titration phase, intact PTH should be monitored 4 weekly (measured at least 12 hours post dose) and dose titrated until an appropriate iPTH concentration is achieved. During the titration phase, approval will be limited to sufficient supply for 4 weeks treatment at a time, with doses between 30 and 180 mg per day according to the patient's response and tolerability.

During the maintenance phase, approval will be limited to provide sufficient quantity for 4 weeks treatment up to a maximum of 6 months supply for doses between 30 and 180 mg per day according to the patient's response and tolerability. Intact PTH should be monitored quarterly (measured at least 12 hours post dose) and dose adjusted as necessary to maintain an appropriate iPTH concentration.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

"Sustained" means the abnormality was detected on at least 2 blood samples collected over a period of 2 to 4 months.

Note

Special Pricing Arrangements apply.

9625N	Tablet 30 mg (base)	56	5	..	*623.88	Sensipar	AN
9626P	Tablet 60 mg (base)	56	5	..	*1233.86	Sensipar	AN
9627Q	Tablet 90 mg (base)	56	5	..	*1827.58	Sensipar	AN

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Antiinfectives for systemic use

Antibacterials for systemic use

Macrolides, lincosamides and streptogramins

Macrolides

AZITHROMYCIN

Authority required

Prophylaxis against Mycobacterium avium complex infections in HIV-positive patients with CD4 cell counts of less than 75 per cubic millimetre.

6221K	Tablet 600 mg	16	5	..	*127.38	Zithromax	PF

CLARITHROMYCIN

Authority required

Treatment of Mycobacterium avium complex infections.

6151R	Tablet 250 mg	100	2	..	46.15	Klacid	AB
6152T	Tablet 500 mg	100	2	..	82.66	Klacid	AB

Antimycobacterials

Drugs for treatment of tuberculosis

Antibiotics

RIFABUTIN

Authority required

Treatment of Mycobacterium avium complex infections in HIV-positive patients;

Prophylaxis against Mycobacterium avium complex infections in HIV-positive patients with CD4 cell counts of less than 75 per cubic millimetre.

6195C	Capsule 150 mg	120	5	..	*617.94	Mycobutin	PF

Antivirals for systemic use

Direct acting antivirals

Nucleosides and nucleotides excl. reverse transcriptase inhibitors

CIDOFOVIR

Authority required

Treatment of cytomegalovirus retinitis in patients with AIDS.

6247T	Solution for I.V. infusion 375 mg (anhydrous) in 5 mL single use vial	4	3	..	*3646.42	Vistide	GI

GANCICLOVIR

Authority required

Cytomegalovirus retinitis in severely immunocompromised patients;

Prophylaxis of cytomegalovirus disease in bone marrow transplant patients at risk of cytomegalovirus disease;

Prophylaxis of cytomegalovirus disease in solid organ transplant patients at risk of cytomegalovirus disease.

6136Y	Powder for I.V. infusion 500 mg (as sodium)	10	1	..	*588.82	Cymevene	RO

VALACICLOVIR HYDROCHLORIDE

Authority required

Prophylaxis of cytomegalovirus (CMV) infection and disease following renal transplantation in patients at risk of CMV disease.

6280M	Tablet 500 mg (base)	500	2	..	*2162.32	Valtrex	GK

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer	
					Max. Qty \$		
VALGANCICLOVIR HYDROCHLORIDE							
<u>Authority required</u>							
Cytomegalovirus retinitis in patients with acquired immunodeficiency syndrome;							
Prophylaxis of cytomegalovirus infection and disease in solid organ transplant patients at risk of cytomegalovirus disease.							
6357N	Tablet 450 mg (base)	120	5	..	*4538.02	Valcyte	RO
9675F	Powder for oral solution 50 mg (base) per mL, 100 mL	11	5	..	*#4623.7	Valcyte	RO

Phosphonic acid derivatives

FOSCARNET SODIUM

Authority required

Treatment of cytomegalovirus retinitis in patients with AIDS;

Treatment of aciclovir-resistant herpes simplex virus infection in immunocompromised patients with HIV infection.

6134W	I.V. infusion 24 mg per mL, 250 mL	6	1	..	417.22	Foscavir	AP

Protease inhibitors

ATAZANAVIR

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6451M	Capsule 150 mg (as sulfate)	120	5	..	*1090.24	Reyataz	BQ
6452N	Capsule 200 mg (as sulfate)	120	5	..	*1438.18	Reyataz	BQ
9614B	Capsule 300 mg (as sulfate)	60	5	..	*1090.24	Reyataz	BQ
9646Q	Capsule 100 mg (as sulfate)	120	5	..	*730.14	Reyataz	BQ

DARUNAVIR

Authority required

Treatment of HIV infection, in addition to optimised background therapy in combination with other antiretroviral agents, and co-administered with 100 mg ritonavir twice daily in an antiretroviral experienced patient who, after at least one antiretroviral regimen, has experienced virological failure or clinical failure or genotypic resistance.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

9581G	Tablet 150 mg (as ethanolate)	240	5	..	1095.13	Prezista	JC
9616D	Tablet 300 mg (as ethanolate)	240	5	..	*2143.84	Prezista	JC

FOSAMPRENAVIR

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6453P	Tablet 700 mg (as calcium)	120	5	..	*795.08	Telzir	GK
6454Q	Oral liquid 50 mg (as calcium) per mL, 225 mL	8	5	..	*851.38	Telzir	VI

INDINAVIR

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6202K	Capsule 400 mg (as sulfate)	360	5	..	*952.82	Crixivan 400 mg	MK

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
RITONAVIR						
<u>Authority required</u>						
Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;						
Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.						
6494T	Oral solution 600 mg per 7.5 mL (80 mg per mL), 90 mL	10	5	..	*952.82	Norvir AB
9677H	Tablet 100 mg	720	5	..	*952.98	Norvir AB

SAQUINAVIR

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6498B	Tablet 500 mg (as mesylate)	240	5	..	*1057.54		Invirase RO

TIPRANAVIR

Authority required

Treatment of HIV infection, in addition to optimised background therapy in combination with other antiretroviral agents, and co-administered with 200 mg ritonavir twice daily in an antiretroviral experienced patient who, after each of at least three different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes, has experienced virological failure or clinical failure or genotypic resistance.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

Note

Special Pricing Arrangements apply.

9610T	Capsule 250 mg	240	5	..	*2188.42		Aptivus BY

TIPRANAVIR

Authority required

Treatment of HIV infection, in addition to optimised background therapy in combination with other antiretroviral agents, and co-administered with ritonavir in an antiretroviral experienced patient who, after each of at least three different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes, has experienced virological failure or clinical failure or genotypic resistance.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

Note

Special Pricing Arrangements apply.

9676G	Oral liquid 100 mg per mL, 95 mL	7	5	..	*2420.44		Aptivus BY

Nucleoside and nucleotide reverse transcriptase inhibitors

ABACAVIR

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6264Q	Tablet 300 mg (as sulfate)	120	5	..	*592.98		Ziagen VI
6265R	Oral solution 20 mg (as sulfate) per mL, 240 mL	8	5	..	*689.86		Ziagen VI

ADEFOVIR DIPIVOXIL

Authority required

Chronic hepatitis B in a patient who has failed antihepadnaviral therapy and who satisfies all of the following criteria:

(1)(a) Repeatedly elevated serum ALT levels while on concurrent antihepadnaviral therapy of greater than or equal to 6 months duration in conjunction with documented chronic hepatitis B infection; or

(b) Repeatedly elevated HBV DNA levels one log greater than the nadir value or failure to achieve a 1 log reduction in HBV DNA within 3 months,

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

whilst on previous antihepadnaviral therapy except in patients with evidence of poor compliance;

(2) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

Note

Patients should have undergone a liver biopsy at some point since initial diagnosis to obtain histological evidence of chronic hepatitis.

Patients may receive treatment in combination with lamivudine but not with other PBS-subsidised antihepadnaviral therapy.

6450L	Tablet 10 mg	60	5	..	*1296.42	Hepsera	GI

DIDANOSINE

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6298L	Capsule 125 mg (containing enteric coated beadlets)	60	5	..	*298.52	Videx EC	BQ
6299M	Capsule 200 mg (containing enteric coated beadlets)	60	5	..	*346.30	Videx EC	BQ
6300N	Capsule 250 mg (containing enteric coated beadlets)	60	5	..	*431.24	Videx EC	BQ
6301P	Capsule 400 mg (containing enteric coated beadlets)	60	5	..	*686.14	Videx EC	BQ

EMTRICITABINE

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6137B	Capsule 200 mg	60	5	..	*592.98	Emtriva	GI

ENTECAVIR MONOHYDRATE

Authority required

Patients with chronic hepatitis B who satisfy all of the following criteria:

(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);

(2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or

(b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;

(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

Note

PBS-subsidised entecavir monohydrate must be used as monotherapy.

9602J	Tablet 0.5 mg	60	5	..	*805.76	Baraclude	BQ

ENTECAVIR MONOHYDRATE

Authority required

Patients with chronic hepatitis B who have failed lamivudine therapy and who satisfy all of the following criteria:

(1)(a) Repeatedly elevated serum ALT levels while on concurrent antihepadnaviral therapy of greater than or equal to 6 months duration in conjunction with documented chronic hepatitis B infection; or

(b) Repeatedly elevated HBV DNA levels one log greater than the nadir value or failure to achieve a 1 log reduction in HBV DNA within 3 months, whilst on previous antihepadnaviral therapy except in patients with evidence of poor compliance;

(2) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

Note

Patients should have undergone a liver biopsy at some point since initial diagnosis to obtain histological evidence of chronic hepatitis.

PBS-subsidised entecavir monohydrate must be used as monotherapy.

9603K	Tablet 1 mg	60	5	..	*1296.42	Baraclude	BQ

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

LAMIVUDINE

Authority required

Patients with chronic hepatitis B who satisfy all of the following criteria:

(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);

(2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or

(b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;

(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

6257H	Tablet 100 mg	56	5	..	*317.08	Zeffix	GK
6271C	Oral solution 5 mg per mL, 240 mL	5	5	..	*369.97	Zeffix	GK

LAMIVUDINE

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6193Y	Tablet 150 mg	120	5	..	*592.98	3TC	VI
6194B	Oral solution 10 mg per mL, 240 mL	8	5	..	*725.94	3TC	VI
6435Q	Tablet 300 mg	60	5	..	*592.98	3TC	VI

STAVUDINE

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6186N	Capsule 20 mg	120	5	..	*588.82	Zerit	BQ
6189R	Capsule 30 mg	120	5	..	*700.48	Zerit	BQ
6190T	Capsule 40 mg	120	5	..	*931.82	Zerit	BQ
6250Y	Powder for oral solution 1 mg per mL, 200 mL	24	5	..	*#2849.3 5	Zerit	BQ

TELBIVUDINE

Authority required

Treatment, as sole PBS-subsidised therapy, in a patient with chronic hepatitis B who is nucleoside analogue naive and satisfies all of the following criteria:

(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);

(2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or

(b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;

(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

9630W	Tablet 600 mg	56	5	..	*528.26	Sebivo	NV

TENOFOVIR

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Authority required

Treatment, as sole PBS-subsidised therapy, of chronic hepatitis B in a patient who is nucleoside analogue naive and satisfies all of the following criteria:

(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);

(2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or

(b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;

(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy;

Chronic hepatitis B in a patient who has failed antihepadnaviral therapy and who satisfies all of the following criteria:

(1)(a) Repeatedly elevated serum ALT levels while on concurrent antihepadnaviral therapy of greater than or equal to 6 months duration in conjunction with documented chronic hepatitis B infection; or

(b) Repeatedly elevated HBV DNA levels one log greater than the nadir value or failure to achieve a 1 log reduction in HBV DNA within 3 months, whilst on previous antihepadnaviral therapy except in patients with evidence of poor compliance;

(2) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

Note

Patients should have undergone a liver biopsy at some point since initial diagnosis to obtain histological evidence of chronic hepatitis.

Patients may receive tenofovir treatment in combination with lamivudine but not with other PBS-subsidised antihepadnaviral therapy.

6358P	Tablet containing tenofovir disoproxil fumarate 300 mg	60	5	..	*1011.26	Viread	GI

ZIDOVUDINE

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6153W	Capsule 100 mg	400	5	..	*861.14	Retrovir	GK
6154X	Capsule 250 mg	240	5	..	*1279.18	Retrovir	GK
6155Y	Syrup 10 mg per mL, 200 mL	15	5	..	*706.62	Retrovir	GK

Non-nucleoside reverse transcriptase inhibitors

EFAVIRENZ

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

Note

Special Pricing Arrangements apply.

6356M	Tablet 600 mg	60	5	..	*947.92	Stocrin	MK
6372J	Oral solution 30 mg per mL, 180 mL	7	5	..	*994.96	Stocrin	MK
9618F	Tablet 200 mg	180	5	..	*947.92	Stocrin	MK

ETRAVIRINE

Authority required

Treatment of HIV infection, in addition to optimised background therapy in combination with other antiretroviral agents in an antiretroviral experienced patient who, after each of at least three different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes, has experienced virological failure or clinical failure or genotypic resistance.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

9639H	Tablet 100 mg	240	5	..	*1279.42	Intelence	JC

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
NEVIRAPINE						
<u>Authority required</u>						
Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;						
Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.						
6215D	Tablet 200 mg	120	5	..	*571.30	Viramune BY
9571R	Oral suspension 50 mg (as hemihydrate) per 5 mL, 240 mL	10	5	..	*1396.42	Viramune BY

Antivirals for treatment of HIV infections, combinations

ABACAVIR SULFATE with LAMIVUDINE and ZIDOVUDINE

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient over 12 years of age, weighing 40 kg or more, with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient over 12 years of age, weighing 40 kg or more, has previously received PBS-subsidised therapy for HIV infection.

6327B	Tablet containing abacavir 300 mg (as sulfate) with lamivudine 150 mg and zidovudine 300 mg	120	5	..	*1750.42		Trizivir VI

ABACAVIR with LAMIVUDINE

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient over 12 years of age, weighing 40 kg or more, with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient over 12 years of age, weighing 40 kg or more, has previously received PBS-subsidised therapy for HIV infection.

6458X	Tablet containing abacavir 600 mg (as sulfate) with lamivudine 300 mg	60	5	..	*1174.42		Kivexa VI

LAMIVUDINE with ZIDOVUDINE

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6234D	Tablet 150 mg-300 mg	120	5	..	*1203.62		Combivir VI

LOPINAVIR with RITONAVIR

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6341R	Oral liquid 400 mg-100 mg per 5 mL, 60 mL	10	5	..	*1336.42		Kaletra AB
6495W	Tablet 200 mg-50 mg	240	5	..	*1416.42		Kaletra AB
9633B	Tablet 100 mg-25 mg	120	5	..	*362.62		Kaletra AB

TENOFOVIR with EMTRICITABINE

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

6468K	Tablet containing tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg	60	5	..	*1576.62		Truvada GI

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

TENOFOVIR with EMTRICITABINE and EFAVIRENZ

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

9650X	Tablet containing tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg and efavirenz 600 mg	60	5	..	*2481.90	Atripla	GI

Other antirvirals

ENFUVIRTIDE

Authority required

Treatment of HIV infection, in addition to optimised background therapy in combination with other antiretroviral agents in an antiretroviral experienced patient who, after each of at least three different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes, has experienced virological failure or clinical failure or genotypic resistance.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

6455R	Pack containing 60 vials powder for injection 90 mg with 60 vials water for injections 1.1 mL (with syringes and swabs)	2	5	..	*4472.42	Fuzeon	RO

MARAVIROC

Authority required

Treatment, in addition to optimised background therapy in combination with other antiretroviral agents, of an antiretroviral experienced patient infected with only CCR5-tropic HIV-1, who, after each of at least three different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes, has experienced virological failure or clinical failure or genotypic resistance. A tropism assay to determine CCR5 only strain status is required prior to initiation. Individuals with CXCR4 tropism demonstrated at any time point are not eligible.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

9572T	Tablet 150 mg	120	5	..	*1881.82	Celsentri	PF
9573W	Tablet 300 mg	120	5	..	*1881.82	Celsentri	PF

RALTEGRAVIR

Authority required

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

9629T	Tablet 400 mg (as potassium)	120	5	..	*1377.52	Isentress	MK

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Antineoplastic and immunomodulating agents

Antineoplastic agents

Cytotoxic antibiotics and related substances

Anthracyclines and related substances

DOXORUBICIN HYDROCHLORIDE, PEGYLATED LIPOSOMAL

Authority required

Treatment of AIDS-related Kaposi's sarcoma in patients with CD4 cell counts of less than 200 per cubic millimetre and extensive mucocutaneous involvement;

Treatment of AIDS-related Kaposi's sarcoma in patients with CD4 cell counts of less than 200 per cubic millimetre and extensive visceral involvement.

6249X	Suspension for I.V. infusion 20 mg in 10 mL	4	5	..	*2538.38	Caelyx	SH

Immunostimulants

Immunostimulants

Colony stimulating factors

FILGRASTIM

Authority required

For use in a patient undergoing induction and consolidation therapy for acute myeloid leukaemia;

Mobilisation of peripheral blood progenitor cells to facilitate harvest of such cells for autologous transplantation into a patient with a non-myeloid malignancy who has had myeloablative or myelosuppressive therapy;

Mobilisation of peripheral blood progenitor cells, in a normal volunteer, for use in allogeneic transplantation;

A patient receiving marrow-ablative chemotherapy and subsequent bone marrow transplantation;

A patient with a non-myeloid malignancy receiving marrow-ablative chemotherapy and subsequent autologous peripheral blood progenitor cell transplantation;

A patient with breast cancer receiving standard dose adjuvant chemotherapy who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

A patient receiving chemotherapy for B-cell chronic lymphocytic leukaemia with fludarabine and cyclophosphamide who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

A patient receiving first-line chemotherapy for Hodgkin disease who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

A patient receiving chemotherapy for myeloma who has had a prior episode of febrile neutropenia, and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

A patient with severe congenital neutropenia (absolute neutrophil count of less than 100 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, and in whom a bone marrow examination has shown evidence of maturational arrest of the neutrophil lineage);

A patient with severe chronic neutropenia (absolute neutrophil count of less than 1,000 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, or evidence of neutrophil dysfunction, and, either having experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics in the previous 12 months, or having recurrent clinically significant infections (a minimum of 3 in the previous 12 months));

A patient with chronic cyclic neutropenia (absolute neutrophil count of less than 500 million cells per litre lasting for 3 days per cycle, measured over 3 separate cycles, and, either having experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics, or having recurrent clinically significant infections (a minimum of 3 in the previous 12 months));

A patient with inoperable Stage III, IVa or IVb squamous cell carcinoma of the oral cavity, larynx, oropharynx or hypopharynx receiving neoadjuvant treatment with docetaxel in combination with cisplatin and fluorouracil who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
<u>Authority required</u>						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in acute lymphoblastic leukaemia;						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in breast cancer (adjuvant chemotherapy with docetaxel in combination with an anthracycline and cyclophosphamide);						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in germ cell tumours;						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in infants and children with CNS tumours;						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in neuroblastoma;						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in non-Hodgkin lymphoma (aggressive grades; or low grade receiving an anthracycline-containing regimen);						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in relapsed Hodgkin disease;						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in sarcoma.						
6126K	Injection 300 micrograms in 1 mL	20	11	..	*3054.42	Neupogen AN
6127L	Injection 480 micrograms in 1.6 mL	20	11	..	*4860.42	Neupogen AN
6291D	Injection 300 micrograms in 0.5 mL single use pre-filled syringe	20	11	..	*3054.42	Neupogen AN
6292E	Injection 480 micrograms in 0.5 mL single use pre-filled syringe	20	11	..	*4860.42	Neupogen AN

LENOGRASTIM

Authority required

Mobilisation of peripheral blood progenitor cells to facilitate harvest of such cells for reinfusion into patients with non-myeloid malignancies who have had myeloablative or myelosuppressive therapy;

Mobilisation of peripheral blood progenitor cells, in normal volunteers, for use in allogeneic transplantation to facilitate harvest of such cells in healthy donors;

Patients with non-myeloid malignancies receiving marrow-ablative chemotherapy and subsequent peripheral blood progenitor cell or bone marrow transplantation;

Patients with breast cancer receiving standard dose adjuvant chemotherapy who have had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

Patients receiving first-line chemotherapy for Hodgkin's disease who have had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned.

Authority required

Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in acute lymphoblastic leukaemia;

Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in Ewing's sarcoma;

Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in germ cell tumours;

Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in infants and children with CNS tumours;

Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in neuroblastoma;

Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in non-Hodgkin's lymphoma (intermediate or high grade);

Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in osteosarcoma;

Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in relapsed Hodgkin's disease;

Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in rhabdomyosarcoma.

6337M	Powder for injection 13,400,000 i.u. (105 micrograms)	20	11	..	*1071.42	Granocyte 13	HH
6338N	Powder for injection 33,600,000 i.u. (263 micrograms)	20	11	..	*2613.62	Granocyte 34	HH

PEGFILGRASTIM

Authority required

For use in a patient undergoing induction and consolidation therapy for acute myeloid leukaemia;

A patient with breast cancer receiving standard dose adjuvant chemotherapy who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for Max. Qty		Brand Name and Manufacturer
					\$	\$	
	therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;						
	A patient receiving chemotherapy for B-cell chronic lymphocytic leukaemia with fludarabine and cyclophosphamide who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;						
	A patient receiving first-line chemotherapy for Hodgkin disease who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;						
	A patient receiving chemotherapy for myeloma who has had a prior episode of febrile neutropenia, and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;						
	A patient with inoperable Stage III, IVa or IVb squamous cell carcinoma of the oral cavity, larynx, oropharynx or hypopharynx receiving neoadjuvant treatment with docetaxel in combination with cisplatin and fluorouracil who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned.						
	<u>Authority required</u>						
	A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in acute lymphoblastic leukaemia;						
	A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in breast cancer (adjuvant chemotherapy with docetaxel in combination with an anthracycline and cyclophosphamide);						
	A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in germ cell tumours;						
	A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in infants and children with CNS tumours;						
	A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in neuroblastoma;						
	A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in non-Hodgkin lymphoma (aggressive grades; or low grade receiving an anthracycline-containing regimen);						
	A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in relapsed Hodgkin disease;						
	A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in sarcoma.						
6363X	Injection 6 mg in 0.6 mL single use pre-filled syringe	1	11	..	1971.42	Neulasta	AN

Interferons

INTERFERON ALFA-2a

Caution

Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required

Use in the treatment of Philadelphia chromosome positive myelogenous leukaemia in the chronic phase;

Patients with chronic hepatitis B who satisfy all of the following criteria:

- (1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);
- (2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or
- (b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;
- (3) Are not persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L);
- (4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

6210W	Injection 3,000,000 i.u. in 0.5 mL single dose pre-filled syringe	30	5	..	*936.12	Roferon-A	RO
6211X	Injection 4,500,000 i.u. in 0.5 mL single dose pre-filled syringe	30	5	..	*1387.32	Roferon-A	RO
6212Y	Injection 6,000,000 i.u. in 0.5 mL single dose pre-filled syringe	30	5	..	*1833.72	Roferon-A	RO
6213B	Injection 9,000,000 i.u. in 0.5 mL single dose pre-filled syringe	30	5	..	*2727.72	Roferon-A	RO

INTERFERON ALFA-2b

Caution

Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
<u>Authority required</u>							
Adjunctive therapy of malignant melanoma following surgery in patients with nodal involvement;							
Use in the treatment of Philadelphia chromosome positive myelogenous leukaemia in the chronic phase;							
Patients with chronic hepatitis B who satisfy all of the following criteria:							
(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);							
(2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or							
(b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;							
(3) Are not persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L);							
(4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.							
6218G	Solution for injection 18,000,000 i.u. in 3 mL single dose vial	15	5	..	*2727.57	Intron A	SH
6219H	Solution for injection 25,000,000 i.u. in 2.5 mL single dose vial	15	5	..	*3770.22	Intron A	SH
6246R	Solution for injection 10,000,000 i.u. in 1 mL single dose vial	15	5	..	*1535.91	Intron A	SH
6253D	Solution for injection 18,000,000 i.u. in 1.2 mL multi-dose injection pen	2	5	..	*378.20	Intron A Redipen	SH
6254E	Solution for injection 30,000,000 i.u. in 1.2 mL multi-dose injection pen	2	5	..	*626.06	Intron A Redipen	SH
6255F	Solution for injection 60,000,000 i.u. in 1.2 mL multi-dose injection pen	2	5	..	*1238.02	Intron A Redipen	SH
INTERFERON GAMMA-1b							
<u>Authority required</u>							
Treatment of chronic granulomatous disease in patients with frequent and severe infections despite adequate prophylaxis with antimicrobial agents							
6148N	Injection 2,000,000 i.u. in 0.5 mL	12	11	..	*2768.22	Imukin	BY
PEGINTERFERON ALFA-2a							
<u>Caution</u>							
Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.							
<u>Authority required</u>							
Monotherapy in patients with chronic hepatitis B and compensated liver disease who satisfy all of the following criteria:							
(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);							
(2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or							
(b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;							
(3) Have received no prior peginterferon alfa therapy for the treatment of hepatitis B;							
(4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception;							
(5) Are not persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L).							
Treatment is limited to 1 course of treatment for a duration of up to 48 weeks;							
Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa or peginterferon alfa treatment for hepatitis C and have a contraindication to ribavirin, who satisfy all of the following criteria:							
(1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);							
(2) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.							
The treatment course is limited to up to 48 weeks.							
Patients may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop.							
<u>Note</u>							
Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:							
(a) a nurse educator/counsellor for patients; and							
(b) 24 hour access by patients to medical advice; and							
(c) an established liver clinic; and							
(d) facilities for safe liver biopsy.							
6439X	Injection 135 micrograms in 0.5 mL single use pre-filled syringe	8	5	..	*2378.22	Pegasys	RO
6449K	Injection 180 micrograms in 0.5 mL single use pre-filled syringe	8	5	..	*2746.88	Pegasys	RO

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
PEGINTERFERON ALFA-2b							
Caution							
Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.							
Authority required							
Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa or peginterferon alfa treatment for hepatitis C and have a contraindication to ribavirin, who satisfy all of the following criteria:							
(1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);							
(2) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.							
The treatment course is limited to up to 48 weeks.							
Patients may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop.							
Note							
Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:							
(a) a nurse educator/counsellor for patients; and							
(b) 24 hour access by patients to medical advice; and							
(c) an established liver clinic; and							
(d) facilities for safe liver biopsy.							
6411K	Powder for injection 50 micrograms with diluent in single use injection pen	8	5	..	*1886.42	PEG-Intron Redipen	SH
6412L	Powder for injection 80 micrograms with diluent in single use injection pen	8	5	..	*2990.42	PEG-Intron Redipen	SH
6413M	Powder for injection 100 micrograms with diluent in single use injection pen	8	5	..	*3726.42	PEG-Intron Redipen	SH
6414N	Powder for injection 120 micrograms with diluent in single use injection pen	8	5	..	*4462.42	PEG-Intron Redipen	SH
6415P	Powder for injection 150 micrograms with diluent in single use injection pen	8	5	..	*5566.42	PEG-Intron Redipen	SH

RIBAVIRIN and PEGINTERFERON ALFA-2a

Caution

Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Caution

Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.

Authority required

Patients naïve to interferon based therapies (non-pegylated or pegylated)

Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa or peginterferon alfa treatment for hepatitis C and who satisfy all of the following criteria:

(1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);

(2) Female patients of child-bearing age are not pregnant, not breast-feeding, and both patient and their partner are using effective forms of contraception (one for each partner). Male patients and their partners are using effective forms of contraception (one for each partner). Female partners of male patients are not pregnant.

For patients with genotype 2 or 3 hepatitis C without hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 24 weeks. For hepatitis C patients with genotype 1, 4, 5 or 6 and those genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 48 weeks.

Patients with genotype 1, 4, 5 or 6 who are eligible for 48 weeks of treatment may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop. (An HCV RNA assay at week 12 is unnecessary for genotype 2 and 3 patients because of the high likelihood of early viral response by week 12).

Patients with genotype 1, 4, 5 or 6 who are viral positive at week 12 but have attained at least a 2 log drop in viral load may only continue treatment after the first 24 weeks of treatment if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. Similarly, genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis may only continue treatment after the first 24 weeks if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. An HCV RNA qualitative assay at week 24 is unnecessary for those patients with genotype 1, 4, 5 or 6 who became viral negative at week 12.

Note

Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:

(a) a nurse educator/counsellor for patients; and

(b) 24 hour access by patients to medical advice; and

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
(c) an established liver clinic; and (d) facilities for safe liver biopsy. Authority required Patients who have failed one prior attempt at interferon based therapies (non-pegylated or pegylated) Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no more than one prior treatment with interferon alfa or peginterferon alfa for hepatitis C and who satisfy all of the following criteria: (1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive); (2) Female patients of child-bearing age are not pregnant, not breast-feeding, and both patient and their partner are using effective forms of contraception (one for each partner). Male patients and their partners are using effective forms of contraception (one for each partner). Female partners of male patients are not pregnant. The treatment course is limited to 48 weeks. Patients may only continue treatment after the first 12 weeks of treatment if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 12. Note Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C: (a) a nurse educator/counsellor for patients; and (b) 24 hour access by patients to medical advice; and (c) an established liver clinic; and (d) facilities for safe liver biopsy.							
6392K	Pack containing 168 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 135 micrograms	2	5	..	*3119.26	Pegasys RBV	RO
6394M	Pack containing 112 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 180 micrograms	2	5	..	*3131.70	Pegasys RBV	RO
6395N	Pack containing 140 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 180 micrograms	2	5	..	*3292.24	Pegasys RBV	RO
6396P	Pack containing 168 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 180 micrograms	2	5	..	*3452.78	Pegasys RBV	RO

RIBAVIRIN and PEGINTERFERON ALFA-2b

Caution

Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Caution

Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.

Authority required

Patients naïve to interferon based therapies (non-pegylated or pegylated)

Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa or peginterferon alfa treatment for hepatitis C and who satisfy all of the following criteria:

- (1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);
- (2) Female patients of child-bearing age are not pregnant, not breast-feeding, and both patient and their partner are using effective forms of contraception (one for each partner). Male patients and their partners are using effective forms of contraception (one for each partner). Female partners of male patients are not pregnant.

For patients with genotype 2 or 3 hepatitis C without hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 24 weeks. For hepatitis C patients with genotype 1, 4, 5 or 6 and those genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 48 weeks.

Patients with genotype 1, 4, 5 or 6 who are eligible for 48 weeks of treatment may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop. (An HCV RNA assay at week 12 is unnecessary for genotype 2 and 3 patients because of the high likelihood of early viral response by week 12).

Patients with genotype 1, 4, 5 or 6 who are viral positive at week 12 but have attained at least a 2 log drop in viral load may only continue treatment after the first 24 weeks of treatment if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. Similarly, genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis may only continue treatment after the first 24 weeks if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. An HCV RNA qualitative assay at week 24 is unnecessary for those patients with genotype 1, 4, 5 or 6 who became viral negative at week 12.

Note

Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:

- (a) a nurse educator/counsellor for patients; and
- (b) 24 hour access by patients to medical advice; and

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
(c) an established liver clinic; and (d) facilities for safe liver biopsy. Authority required Patients who have failed one prior attempt at interferon based therapies (non-pegylated or pegylated) Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no more than one prior treatment with interferon alfa or peginterferon alfa for hepatitis C and who satisfy all of the following criteria: (1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive); (2) Female patients of child-bearing age are not pregnant, not breast-feeding, and both patient and their partner are using effective forms of contraception (one for each partner). Male patients and their partners are using effective forms of contraception (one for each partner). Female partners of male patients are not pregnant. The treatment course is limited to 48 weeks. Patients may only continue treatment after the first 12 weeks of treatment if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 12. Note Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C: (a) a nurse educator/counsellor for patients; and (b) 24 hour access by patients to medical advice; and (c) an established liver clinic; and (d) facilities for safe liver biopsy.							
6399T	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 50 micrograms with diluent	2	5	..	*1881.18	Pegatron	SH
6400W	Pack containing 112 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 50 micrograms with diluent	2	5	..	*2166.16	Pegatron	SH
6401X	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 80 micrograms with diluent	2	5	..	*2469.14	Pegatron	SH
6402Y	Pack containing 140 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 80 micrograms with diluent	2	5	..	*2754.08	Pegatron	SH
6403B	Pack containing 168 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 80 micrograms with diluent	2	5	..	*2754.08	Pegatron	SH
6404C	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 100 micrograms with diluent	2	5	..	*2861.08	Pegatron	SH
6405D	Pack containing 112 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 100 micrograms with diluent	2	5	..	*3146.04	Pegatron	SH
6406E	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 120 micrograms with diluent	2	5	..	*3253.04	Pegatron	SH
6407F	Pack containing 140 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 120 micrograms with diluent	2	5	..	*3538.00	Pegatron	SH
6408G	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	2	5	..	*3840.98	Pegatron	SH
6409H	Pack containing 140 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	2	5	..	*4125.94	Pegatron	SH
6410J	Pack containing 168 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	2	5	..	*4125.94	Pegatron	SH
9634C	Pack containing 196 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	2	5	..	*4410.90	Pegatron	SH

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Immunosuppressants

Immunosuppressants

Selective immunosuppressants

ABATACEPT

Note

Any queries concerning the arrangements to prescribe abatacept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Further prescribing information (including Authority Application Forms) is on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe abatacept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (new patient or patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with abatacept, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

[www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

(i) a total active joint count of at least 20 active (swollen and tender) joints; or

(ii) at least 4 active joints from the following list of major joints:

— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and

(3) a signed patient acknowledgement.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion. Up to a maximum of 4 repeats may be authorised.

Where fewer than 4 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with abatacept.

Patients who fail to demonstrate a response to treatment with abatacept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with abatacept, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

(a) have a documented history of severe active rheumatoid arthritis; and

(b) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with abatacept and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised abatacept treatment, within the timeframes

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

specified below.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion. Up to a maximum of 4 repeats may be authorised.

Where fewer than 4 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised abatacept treatment was approved under either of the initial 1 or 2 treatment restrictions, patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised abatacept treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Patients who fail to demonstrate a response to treatment with abatacept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with abatacept, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have a documented history of severe active rheumatoid arthritis; and
- (b) who have demonstrated an adequate response to treatment with abatacept; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with abatacept.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion. Up to a maximum of 5 repeats may be authorised.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with abatacept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with abatacept, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Patients who fail to demonstrate a response to treatment with abatacept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

9621J	Powder for I.V. infusion 250 mg	1	..	..	531.03	Orencia	BQ

EVEROLIMUS

Caution

Careful monitoring of patients is mandatory.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
<u>Authority required</u>						
Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required;						
Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of cardiac allograft rejection. Management includes initiation, stabilisation and review of therapy as required.						
6459Y	Tablet 0.25 mg	120	5	..	*506.24	Certican NV
6460B	Tablet 0.5 mg	120	5	..	*1006.06	Certican NV
6461C	Tablet 0.75 mg	240	5	..	*2930.02	Certican NV
9582H	Tablet 1 mg	240	5	..	*3891.22	Certican NV

MYCOPHENOLATE MOFETIL

Caution

Careful monitoring of patients is mandatory.

Authority required

Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required;

Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of cardiac allograft rejection. Management includes initiation, stabilisation and review of therapy as required.

6208R	Capsule 250 mg	600	5	..	*1157.82	CellCept	RO
6209T	Tablet 500 mg	300	5	..	*1157.82	CellCept	RO
6364Y	Powder for oral suspension 1 g per 5 mL, 165 mL	2	5	..	*517.53	CellCept	RO

MYCOPHENOLATE SODIUM

Caution

Careful monitoring of patients is mandatory.

Authority required

Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required.

6369F	Tablet (enteric coated) 180 mg (mycophenolic acid)	240	5	..	*468.76	Myfortic	NV
6370G	Tablet (enteric coated) 360 mg (mycophenolic acid)	240	5	..	*931.10	Myfortic	NV

NATALIZUMAB

Caution

Progressive multifocal leukoencephalopathy has been reported with this drug.

Note

Neurologists prescribing natalizumab under the PBS listing must be registered with the Tysabri Australian Prescribing Program.

Authority required

Initial treatment, as monotherapy, by a neurologist, of clinically definite relapsing-remitting multiple sclerosis in an ambulatory (without assistance or support) patient 18 years of age or older, who has experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years.

The diagnosis must be confirmed by magnetic resonance imaging of the brain and/or spinal cord and the date of the scan included in the authority application, unless the authority application is accompanied by written certification provided by a radiologist that an MRI scan is contraindicated because of the risk of physical (not psychological) injury to the patient.

Authority required

Continuing treatment, as monotherapy, of clinically definite relapsing-remitting multiple sclerosis in a patient previously issued with an authority prescription for this drug who does not show continuing progression of disability while on treatment with this drug, and who has demonstrated compliance with, and an ability to tolerate, this therapy.

Note

Special Pricing Arrangements apply.

9624M	Solution concentrate for I.V. infusion 300 mg in 15 mL	1	5	..	2084.88	Tysabri	BD

SIROLIMUS

Caution

Careful monitoring of patients is mandatory.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
<u>Authority required</u>						
Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required.						
6436R	Tablet 1 mg	200	5	..	*1493.08	Rapamune WX
6437T	Oral solution 1 mg per mL, 60 mL	2	5	..	*979.86	Rapamune WX
6457W	Tablet 2 mg	200	5	..	*2939.76	Rapamune WX

Tumor necrosis factor alpha (TNF-alpha) inhibitors

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and etanercept for a patient who has severe active juvenile idiopathic arthritis. Where the term bDMARD appears in the following NOTES and restrictions, it refers to adalimumab and etanercept only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 bDMARDs at any 1 time.

From 1 November 2010, a patient receiving PBS-subsidised bDMARD therapy is considered to be in a treatment cycle where they may swap to the alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy, and
- fail to respond, or to sustain a response to one PBS-subsidised bDMARD twice and the other PBS-subsidised bDMARD once only.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological therapy before they are eligible to receive further PBS-subsidised bDMARD therapy. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment was stopped to the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

A patient who was receiving PBS-subsidised bDMARD treatment immediately prior to 1 November 2010 is considered to be in their first cycle as of 1 November 2010. A patient who has had a break in bDMARD treatment of at least 12 months immediately prior to making a new application, on or after 1 November 2010, will commence a new treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 November 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to the alternate bDMARD without having to requalify with respect to the indices of disease severity (joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months.

A patient may trial the alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug twice within the current treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a bDMARD. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to the revised baseline measurement.

(4) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 12 months, must requalify for treatment under the Initial 1 treatment restriction.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for severe active juvenile idiopathic arthritis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must qualify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 12 month break in PBS-subsidised therapy' above for further details.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with bDMARDs should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Medicare Australia at the time treatment is ceased.

Authority required

Initial 1 (new patient or patient recommencing after a break of more than 12 months).

Initial treatment by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of a patient under 18 years:

- (a) who has severe active juvenile idiopathic arthritis; AND
- (b) whose parent or authorised guardian has signed a patient acknowledgement; AND
- (c) who has not received PBS-subsidised treatment with adalimumab or etanercept for this condition in the previous 12 months; AND
- (d) who has demonstrated either:
 - (i) severe intolerance of, or toxicity due to, methotrexate (see below for definition of severe intolerance and toxicity); or
 - (ii) failure to achieve an adequate response to 1 or more of the following treatment regimens:
 - oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or
 - oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other DMARD, alone or in combination with corticosteroids, for a minimum of 3 months. (Note: use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.)

Severe intolerance is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant NSAIDs on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, please provide details at time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of this toxicity at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; OR
- (b) at least 4 active joints from the following list:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment should be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) an acknowledgement signed by a parent or authorised guardian.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of authority application, medical practitioners should request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 4 weeks from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial adalimumab after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Authority required

Initial 2 (change or re-commencement after break of less than 12 months).

Initial PBS-subsidised treatment with adalimumab by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of a patient under 18 years who:

- (a) has a documented history of severe active juvenile idiopathic arthritis; and
- (b) in this treatment cycle, has received prior PBS-subsidised treatment with adalimumab or etanercept for this condition; and
- (c) has not failed PBS-subsidised therapy with adalimumab for this condition more than once in the current treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for a patient who has received PBS-subsidised treatment with adalimumab in this treatment cycle and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised adalimumab treatment, within the timeframes specified below.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of authority application, medical practitioners should request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised adalimumab treatment was approved under either of the Initial 1 or 2 treatment restrictions, the patient must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised adalimumab treatment was approved under the continuing treatment criteria, the patient must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to that particular course of bDMARD.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial adalimumab after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

Authority required

Initial 3 ('grandfather' patients).

Initial PBS-subsidised supply for continuing treatment with adalimumab, by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of a patient under 18 years who:

- (a) has a documented history of severe active juvenile idiopathic arthritis; and
- (b) was receiving treatment with adalimumab prior to 1 March 2010; and
- (c) has demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with adalimumab; and
- (d) is receiving treatment with adalimumab at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) an acknowledgement signed by a parent or authorised guardian.

A maximum of 24 weeks of treatment will be authorised under this restriction.

At the time of authority application, medical practitioners should request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 5 repeats will be authorised.

Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

The assessment of the patient's response to this initial course of PBS-subsidised therapy must be made within the 4 weeks prior to completion of the course in order to ensure continuity of treatment.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

A patient ceasing treatment or swapping to an alternate agent and wishing to demonstrate a response to treatment, must be assessed no earlier than 12 weeks from the commencement of PBS-subsidised treatment. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A patient may only qualify for PBS-subsidised treatment under this restriction once.

Authority required

Continuing treatment.

Continuing PBS-subsidised treatment with adalimumab, by a rheumatologist or under the supervision of a paediatric rheumatology treatment centre, of a patient:

- (a) who has a documented history of severe active juvenile idiopathic arthritis; and
- (b) who has demonstrated an adequate response to treatment with adalimumab; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment in this treatment cycle was with adalimumab.

An adequate response to treatment is defined as:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

At the time of authority application, medical practitioners should request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 5 repeats will be authorised.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with adalimumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with adalimumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial adalimumab after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

9678J	Injection 20 mg in 0.4 mL pre-filled syringe	2	..	..	1676.42	Humira	AB
9679K	Injection 40 mg in 0.8 mL pre-filled syringe	2	..	..	1676.42	Humira	AB
9680L	Injection 40 mg in 0.8 mL pre-filled pen	2	..	..	1676.42	Humira	AB

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

;

Note

TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and etanercept for a patient who has severe active juvenile idiopathic arthritis. Where the term bDMARD appears in the following NOTES and restrictions, it refers to adalimumab and etanercept only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 bDMARDs at any 1 time.

From 1 November 2010, a patient receiving PBS-subsidised bDMARD therapy is considered to be in a treatment cycle where they may swap to the alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy, and
- fail to respond, or to sustain a response to one PBS-subsidised bDMARD twice and the other PBS-subsidised bDMARD once only.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological therapy before they are eligible to receive further PBS-subsidised bDMARD therapy. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment was stopped to the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

A patient who was receiving PBS-subsidised bDMARD treatment immediately prior to 1 November 2010 is considered to be in their first cycle as of 1 November 2010. A patient who has had a break in bDMARD treatment of at least 12 months immediately prior to making a new application, on or after 1 November 2010, will commence a new treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 November 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to the alternate bDMARD without having to requalify with respect to the indices of disease severity (joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months.

A patient may trial the alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug twice within the current treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a bDMARD. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to the revised baseline measurement.

(4) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 12 months, must requalify for treatment under the Initial 1 treatment restriction.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for severe active juvenile idiopathic arthritis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must qualify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 12 month break in PBS-subsidised therapy' above for further details.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with bDMARDs should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Medicare Australia at the time treatment is ceased.

Authority required

Initial 1 (new patient or patient recommencing after a break of more than 12 months).

Initial treatment by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of a patient under 18 years:

- (a) who has severe active juvenile idiopathic arthritis; AND
- (b) whose parent or authorised guardian has signed a patient acknowledgement; AND
- (c) who has not received PBS-subsidised treatment with adalimumab or etanercept for this condition in the previous 12 months; AND
- (d) who has demonstrated either:

- (i) severe intolerance of, or toxicity due to, methotrexate (see below for definition of severe intolerance and toxicity); or
- (ii) failure to achieve an adequate response to 1 or more of the following treatment regimens:

— oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or

— oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other DMARD, alone or in combination with corticosteroids, for a minimum of 3 months. (Note: use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.)

Severe intolerance is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant NSAIDs on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

divided doses over 24 hours.

Toxicity is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, please provide details at time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of this toxicity at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; OR
- (b) at least 4 active joints from the following list:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment should be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) an acknowledgement signed by a parent or authorised guardian.

A maximum of 16 weeks of treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 4 weeks from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial etanercept after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months).

Initial PBS-subsidised treatment with etanercept by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of a patient under 18 years who:

- (a) has a documented history of severe active juvenile idiopathic arthritis; and
- (b) in this treatment cycle, has received prior PBS-subsidised treatment with adalimumab or etanercept for this condition; and
- (c) has not failed PBS-subsidised therapy with etanercept for this condition more than once in the current treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for a patient who has received PBS-subsidised treatment with etanercept in this treatment cycle and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised etanercept treatment, within the timeframes specified below.

A maximum of 16 weeks of treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with etanercept may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised etanercept treatment was approved under either of the Initial 1 or 2 treatment restrictions, the patient must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised etanercept treatment was approved under the continuing treatment criteria, the patient must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to that particular course of bDMARD.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial etanercept after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

Authority required

Continuing treatment.

Continuing PBS-subsidised treatment with etanercept, by a rheumatologist or under the supervision of a paediatric rheumatology treatment centre, of a patient:

- (a) who has a documented history of severe active juvenile idiopathic arthritis; and
- (b) who has demonstrated an adequate response to treatment with etanercept; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment in this treatment cycle was with etanercept.

An adequate response to treatment is defined as:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with etanercept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with etanercept, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial etanercept after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

6367D	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	1	..	..	854.02	Enbrel	WX

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

;

Note

TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and etanercept for a patient who has severe active juvenile idiopathic arthritis. Where the term bDMARD appears in the following NOTES and restrictions, it refers to adalimumab and etanercept only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 bDMARDs at any 1 time.

From 1 November 2010, a patient receiving PBS-subsidised bDMARD therapy is considered to be in a treatment cycle where they may swap to the alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy, and
- fail to respond, or to sustain a response to one PBS-subsidised bDMARD twice and the other PBS-subsidised bDMARD once only.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological therapy before they are eligible to receive further PBS-subsidised bDMARD therapy. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment was stopped to the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

A patient who was receiving PBS-subsidised bDMARD treatment immediately prior to 1 November 2010 is considered to be in their first cycle as of 1 November 2010. A patient who has had a break in bDMARD treatment of at least 12 months immediately prior to making a new application, on or after 1 November 2010, will commence a new treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 November 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to the alternate bDMARD without having to requalify with respect to the indices of disease severity (joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months.

A patient may trial the alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug twice within the current treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a bDMARD. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to the revised baseline measurement.

(4) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 12 months, must requalify for treatment under the Initial 1 treatment restriction.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for severe active juvenile idiopathic arthritis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must qualify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 12 month break in PBS-subsidised therapy' above for further details.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with bDMARDs should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Medicare Australia at the time treatment is ceased.

Authority required

Continuing treatment.

Continuing PBS-subsidised treatment with etanercept, by a rheumatologist or under the supervision of a paediatric rheumatology treatment centre, of a patient 18 years or older:

- (a) who has a documented history of severe active juvenile idiopathic arthritis; and
- (b) who has demonstrated an adequate response to treatment with etanercept; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment in this treatment cycle was with etanercept.

An adequate response to treatment is defined as:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].						
A maximum of 24 weeks of treatment will be approved under this restriction.						
Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).						
All applications for continuing treatment with etanercept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with etanercept, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.						
If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle.						
Where a patient with severe active juvenile idiopathic arthritis continues treatment with etanercept and is 18 years or older, etanercept 50 mg may be prescribed.						
9615C	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	..	..	1676.43	Enbrel WX
9641K	Injection 50 mg in 1 mL single use auto-injector, 4	1	..	..	1676.43	Enbrel WX

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH ACTIVE ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept, golimumab and infliximab for adult patients with active ankylosing spondylitis. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab, etanercept, golimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 4 TNF-alfa antagonists at any 1 time.

From 1 March 2007, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised TNF-alfa antagonists without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 March 2007 is considered to be in their first cycle as of 1 March 2007.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than once. A patient who, prior to 1 March 2007, was authorised to receive PBS-subsidised initial treatment for ankylosing spondylitis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2007.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and 18 weeks of treatment for infliximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap to an alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with at least 1 NSAID, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the BASDAI, ESR and/or CRP levels are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for active ankylosing spondylitis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with golimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with golimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial 1 (new patients)

Initial PBS-subsidised treatment with infliximab, by a rheumatologist, of an adult with active ankylosing spondylitis who has radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis and who has not received any PBS-subsidised treatment with either adalimumab, etanercept, golimumab or infliximab in this treatment cycle; AND

(a) who has at least 2 of the following:

- (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or
 - (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI) [for further information on the BASMI please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; or
 - (iii) limitation of chest expansion relative to normal values for age and gender [for chest expansion normal values please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; AND
- (b) who has failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months.

The application must include details of the NSAIDs trialled, their doses and duration of treatment. If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance. Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Medicare Australia website at www.medicareaustralia.gov.au.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au] which must include the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (ii) a completed BASDAI Assessment Form [www.medicareaustralia.gov.au]; and
- (iii) a completed Exercise Program Self Certification Form included in the supporting information form; and
- (iv) a signed patient acknowledgment form.

The assessment of the patient's response to the initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A maximum of 18 weeks of treatment with infliximab will be approved under this criterion.

At the time of the authority application, the doctor should request the appropriate number of vials, based on the weight of the patient, to provide for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 18 weeks of treatment may be requested by telephone.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF- α antagonist was approved in this cycle and the date of the first application under a new cycle.

Authority required

Initial 2 (change or re-commencement for all patients)

Initial PBS-subsidised treatment with infliximab, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who, in this treatment cycle, has received prior PBS-subsidised TNF- α antagonist treatment for this condition and is eligible to receive further TNF- α antagonist therapy, and has not failed PBS-subsidised therapy with infliximab in the current treatment cycle.

Where the most recent course of PBS-subsidised TNF- α antagonist treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised TNF- α antagonist therapy or, under this restriction, for patients who have received previous PBS-subsidised TNF- α antagonist therapy) the patient must have been assessed for response to that course following a minimum of 12 weeks of treatment. These assessments must be provided to Medicare Australia no later than 4 weeks from the date the course was ceased. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Where the most recent course of PBS-subsidised infliximab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au].

A maximum of 18 weeks of treatment with infliximab will be approved under this criterion.

At the time of the authority application, the doctor should request the appropriate number of vials, based on the weight of the patient, to provide for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 18 weeks of treatment may be requested by telephone.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF- α antagonist was approved in this cycle and the date of the first application under a new cycle.

Authority required

Continuing treatment for all patients

Continuing PBS-subsidised treatment, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who:

- (a) has demonstrated an adequate response to treatment with infliximab; and
- (b) whose most recent course of PBS-subsidised therapy in this treatment cycle was with infliximab.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au].

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
All measurements provided must be no more than 1 month old at the time of application. A maximum of 24 weeks of treatment with infliximab will be authorised under this criterion. At the time of the authority application, the doctor should request the appropriate number of vials, based on the weight of the patient, to provide for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised. Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone. All applications for continuing treatment with infliximab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment following an initial treatment course it must be made following a minimum of 12 weeks of treatment with infliximab. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment. Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.							
6448J	Powder for I.V. infusion 100 mg	1	..	..	788.19	Remicade	SH

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (new patient or patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with infliximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 22 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 3 mg per kg. Up to a maximum of 3 repeats may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with infliximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have a documented history of severe active rheumatoid arthritis; and
- (b) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with infliximab and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised infliximab treatment, within the timeframes specified below.

A maximum of 22 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 3 mg per kg. Up to a maximum of 3 repeats may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised infliximab treatment was approved under either of the initial 1 or 2 treatment restrictions, patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised infliximab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with infliximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have a documented history of severe active rheumatoid arthritis; and
- (b) who have demonstrated an adequate response to treatment with infliximab; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with infliximab.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 3 mg per kg. Up to a maximum of 2 repeats may be authorised.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with infliximab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with infliximab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

6397Q	Powder for I.V. infusion 100 mg	1	..	..	788.19	Remicade	SH

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents (adalimumab, etanercept, golimumab and infliximab) for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time. Where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, golimumab and infliximab.

From 1 August 2006, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis after 1 August 2010.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients — golimumab only.

Applications for patients who commenced treatment with golimumab prior to 1 March 2010 may apply for initial PBS-subsidised treatment as continuing therapy under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; or
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment Cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Authority required

Initial 1

Initial PBS-subsidised treatment with infliximab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults who:

- (1) have severe active psoriatic arthritis; and
- (2) have received no prior PBS-subsidised biological treatment for this condition in this Treatment Cycle; and
- (3) have failed to achieve an adequate response to:
 - (a) methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months; and
 - (b) sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; or
 - (c) leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities, including severity, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) an active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 22 weeks treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

The assessment of the patient's response to the initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Initial 2

Initial PBS-subsidised treatment with infliximab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis,

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

of adults who:

- (1) have a documented history of severe active psoriatic arthritis; and
- (2) have received prior PBS-subsidised biological treatment for this condition in this Treatment Cycle and are eligible to receive further biological therapy; and
- (3) have not failed treatment with infliximab during the current Treatment Cycle.

Applications for patients who have received PBS-subsidised treatment with infliximab within this Treatment Cycle and who wish to re-commence therapy with this drug within this same Cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised infliximab treatment, within the timeframes specified below.

A maximum of 22 weeks treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised infliximab treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised biological therapy or, under this restriction, for patients who have received previous PBS-subsidised biological therapy), patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised infliximab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with infliximab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults:

- (1) who have a documented history of severe active psoriatic arthritis; and
- (2) whose most recent course of PBS-subsidised biological agent for this condition in the current Treatment Cycle was with infliximab; and
- (3) who, at the time of application, demonstrate an adequate response to treatment with infliximab.

An adequate response to treatment with infliximab is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats may be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with infliximab must include a measurement of response to the prior course of therapy. This assessment

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with infliximab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents (adalimumab, etanercept, golimumab and infliximab) for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time. Where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, golimumab and infliximab.

From 1 August 2006, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis after 1 August 2010.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients — golimumab only.

Applications for patients who commenced treatment with golimumab prior to 1 March 2010 may apply for initial PBS-subsidised treatment as continuing therapy under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialed it on the PBS; or
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment Cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

6496X	Powder for I.V. infusion 100 mg	1	..	..	788.19	Remicade	SH

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and infliximab for adult patients with severe refractory Crohn disease. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alfa antagonists at any 1 time.

From 1 August 2008, under the PBS, all patients will be able to commence a treatment cycle where they may trial each PBS-subsidised TNF-alfa antagonist without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 August 2008 is considered to be in their first cycle as of 1 August 2008.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than twice.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2008.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2008, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap if eligible to the alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, evidence of intestinal inflammation), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab or infliximab.

A patient who commenced treatment with adalimumab for severe refractory Crohn disease prior to 9 November 2007 or infliximab prior to 7 March 2007 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab or infliximab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab or infliximab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial 1 (new patients)

Initial treatment of Crohn disease in a patient assessed by CDAI.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist or a consultant physician as specified in the NOTE below, of a patient with severe refractory Crohn disease who satisfies the following criteria:

(a) has confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

diagnosis confirmed by a gastroenterologist or a consultant physician as specified in the NOTE below; and

(b) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and

(c) has failed to achieve an adequate response to prior systemic therapy including:

(i) a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; and

(ii) immunosuppressive therapy including:

- azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
- 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
- methotrexate at a dose of at least 15 mg weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

(a) have a severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as assessed.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

The most recent CDAI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the signed patient acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete the 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

A CDAI assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Initial 2

Change or re-commencement of treatment of Crohn disease in a patient assessed by CDAI.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist or a consultant physician as specified in the NOTE below of a patient who:

- (a) has a documented history of severe refractory Crohn disease; and
- (b) in this treatment cycle, has received prior PBS-subsidised treatment with infliximab or adalimumab for this condition; and
- (c) has not failed PBS-subsidised therapy with infliximab for this condition more than once in the current treatment cycle.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of TNF-alfa antagonist therapy within the timeframes specified in the relevant restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of TNF-alfa antagonist treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of TNF-alfa antagonist.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition; and
 - (ii) details of prior TNF alfa antagonist treatment including details of date and duration of treatment.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete the 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

A CDAI assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Continuing treatment of Crohn disease in a patient assessed by CDAI.

Continuing PBS-subsidised treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease; and
- (b) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition.

The CDAI assessment must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with infliximab, a CDAI assessment of the patient's response must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial 1

Initial treatment of Crohn disease in a patient with short gut syndrome or an ostomy patient.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist, or consultant physician as specified in the NOTE below of a patient who satisfies the following criteria:

- (a) has confirmed Crohn disease defined by standard clinical, endoscopic and/or imaging features, including histological evidence with the diagnosis confirmed by a gastroenterologist or consultant physician as specified in the NOTE below; and
- (b) has diagnostic imaging or surgical evidence of short gut syndrome or has an ileostomy or colostomy; and
- (c) has evidence of intestinal inflammation; and
- (d) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (e) has failed to achieve an adequate response to prior systemic drug therapy including:
 - (i) a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; and
 - (ii) immunosuppressive therapy including:
 - azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
 - 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
 - methotrexate at a dose of at least 15 mg weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) have evidence of intestinal inflammation, including:
 - (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; AND/OR
 - (ii) faeces: higher than normal lactoferrin or calprotectin level; AND/OR
 - (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery;
 AND/OR
- (b) be assessed clinically as being in a high faecal output state;
- (c) be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of infliximab.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (ii) reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iii) date of the most recent clinical assessment; and
 - (iv) the signed patient acknowledgement.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete the 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Initial 2

Change or re-commencement of treatment of Crohn disease in a patient with short gut syndrome, an ostomy patient or a patient with extensive small intestine disease.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist or a consultant physician as specified in the NOTE below of a patient who:

- (a) has a documented history of severe refractory Crohn disease; and
- (b) in this treatment cycle, has received prior PBS-subsidised treatment with infliximab or adalimumab for this condition; and
- (c) has not failed PBS-subsidised therapy with infliximab for this condition more than once in the current treatment cycle.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of TNF-alfa antagonist therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of TNF-alfa antagonist treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of TNF-alfa antagonist.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criteria, if relevant; and
 - (ii) details of prior TNF alfa antagonist treatment including details of date and duration of treatment.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Continuing treatment of Crohn disease in a patient with short gut syndrome or an ostomy patient.

Continuing PBS-subsidised treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (a) has a documented history of severe refractory Crohn disease with intestinal inflammation and with short gut syndrome or with an ileostomy or colostomy; and
- (b) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as:

(a) improvement of intestinal inflammation as demonstrated by:

- (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; AND/OR
- (ii) faeces: normalisation of lactoferrin or calprotectin level; AND/OR
- (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or
- (b) reversal of high faecal output state; or
- (c) avoidance of the need for surgery or total parenteral nutrition (TPN).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy or the date of clinical assessment.

The patient's assessment must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with infliximab, an assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial 1

Initial treatment of Crohn disease in a patient with extensive small intestine disease.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist or a consultant physician as specified in the NOTE below, of a patient with severe refractory Crohn disease who satisfies the following criteria:

- (a) has confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or consultant physician as specified in the NOTE below; and
- (b) has extensive small intestinal disease with radiological evidence of intestinal inflammation affecting more than 50 cm of the small intestine; and
- (c) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) has failed to achieve an adequate response to prior systemic therapy including:
 - (i) a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; and
 - (ii) immunosuppressive therapy including:
 - azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
 - 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
 - methotrexate at a dose of at least 15 mg weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) have severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; AND/OR
- (b) have evidence of active intestinal inflammation, including:
 - (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; AND/OR
 - (ii) faeces: higher than normal lactoferrin or calprotectin level; AND/OR
 - (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery; AND/OR
- (c) be assessed clinically as being in a high faecal output state; AND/OR
- (d) be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of infliximab.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (ii) (1) reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; or
 - (2) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the dates of assessment of the patient's condition, if relevant; and
 - (iii) date of the most recent clinical assessment; and
 - (iv) the signed patient acknowledgement.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete the 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Continuing treatment of Crohn disease in a patient with extensive small intestine disease.

Continuing PBS-subsidised treatment with infliximab by a gastroenterologist, or consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease with extensive intestinal inflammation affecting more than 50 cm of the small intestine; and
- (b) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

An adequate response to infliximab treatment is defined as:

- (a) a reduction in Crohn Disease Activity Index (CDAI) Score to no greater than 150; or
- (b) improvement of intestinal inflammation as demonstrated by:
 - (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; AND/OR
 - (ii) faeces: normalisation of lactoferrin or calprotectin level; AND/OR
 - (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or
- (c) reversal of high faecal output state; or
- (d) avoidance of the need for surgery or total parenteral nutrition (TPN).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition; or
 - (ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy; or
 - (iii) the date of clinical assessment.

All assessments must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with infliximab, an assessment of the patient's response must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial 3 (grandfather)

Initial PBS-subsidised treatment of Crohn disease in a patient assessed by CDAI who has previously received non-PBS-subsidised therapy with infliximab.

Initial PBS-subsidised supply for continuing treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below, or other consultant physician in consultation with a gastroenterologist of a patient who:

- (a) has a documented history of severe refractory Crohn disease and was receiving treatment with infliximab prior to 7 March 2007; and
- (b) had a Crohn Disease Activity Index (CDAI) Score of greater than or equal to 300 prior to commencing treatment with infliximab. Where a baseline CDAI assessment is not available, please call Medicare Australia on 1800 700 270 to discuss; and
- (c) has signed a patient acknowledgement indicating that they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) has demonstrated or sustained an adequate response to treatment with infliximab. For advice please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as a reduction in Crohn Disease Activity Index (CDAI) Score to no greater than 150.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and baseline Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition; and
 - (ii) the signed patient acknowledgement.

The current CDAI assessment must be no more than 1 month old at the time of application. The baseline CDAI assessment must be from

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

immediately prior to commencing treatment with infliximab.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Authority required

Initial 3

Initial PBS-subsidised treatment of Crohn disease in a patient with short gut syndrome, an ostomy patient, or a patient with extensive small intestine disease, who has previously received non-PBS-subsidised therapy with infliximab.

Initial PBS-subsidised supply for continuing treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below, or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease and was receiving treatment with infliximab prior to 7 March 2007; and
- (b) (1) has a history of extensive small intestinal disease with radiological evidence of intestinal inflammation affecting more than 50 cm of the small intestine; or
- (2) has diagnostic imaging or surgical evidence of short gut syndrome or has an ileostomy or colostomy with a documented history of intestinal inflammation; and
- (c) has signed a patient acknowledgement indicating that they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) has demonstrated or sustained an adequate response to treatment with infliximab according to the criteria included in the relevant continuation restriction. For advice please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

The same criteria used to determine an inadequate response to prior treatment at baseline must be used to determine response to treatment and eligibility for continuing therapy, according to the criteria included in the continuing treatment restriction.

An adequate response to infliximab treatment is defined as:

- (a) a reduction in Crohn Disease Activity Index (CDAI) Score to no greater than 150; or
- (b) improvement of intestinal inflammation as demonstrated by:
 - (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; AND/OR
 - (ii) faeces: normalisation of lactoferrin or calprotectin level; AND/OR
 - (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or
- (c) reversal of high faecal output state; or
- (d) avoidance of the need for surgery or total parenteral nutrition (TPN).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) (1) the completed current and baseline Crohn Disease Activity Index (CDAI) Score calculation sheet, where relevant, including the date of the assessment of the patient's condition; or
 - (2) the reports and dates of the current and baseline pathology or diagnostic imaging test(s) in order to assess response to therapy; or
 - (3) the date of clinical assessment(s); and
 - (ii) the signed patient acknowledgement.

The patient's assessment must be no more than 1 month old at the time of application. The baseline CDAI assessments must be from immediately prior to commencing treatment with infliximab. Where a baseline assessment is not available, please call Medicare Australia on 1800 700 270 to discuss.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and infliximab for adult patients with severe refractory Crohn disease. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alfa antagonists at any 1 time.

From 1 August 2008, under the PBS, all patients will be able to commence a treatment cycle where they may trial each PBS-subsidised TNF-alfa antagonist without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 August 2008 is considered to be in their first cycle as of 1 August 2008.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than twice.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2008.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2008, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap if eligible to the alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, evidence of intestinal inflammation), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab or infliximab.

A patient who commenced treatment with adalimumab for severe refractory Crohn disease prior to 9 November 2007 or infliximab prior to 7 March 2007 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab or infliximab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab or infliximab will be assessed under the continuing treatment restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
9613Y	Powder for I.V. infusion 100 mg	1	..	..	788.19		Remicade SH

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and infliximab for adult patients with severe refractory Crohn disease. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alfa antagonists at any 1 time.

From 1 August 2008, under the PBS, all patients will be able to commence a treatment cycle where they may trial each PBS-subsidised TNF-alfa antagonist without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 August 2008 is considered to be in their first cycle as of 1 August 2008.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than twice.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2008.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2008, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap if eligible to the alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, evidence of intestinal inflammation), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab or infliximab.

A patient who commenced treatment with adalimumab for severe refractory Crohn disease prior to 9 November 2007 or infliximab prior to 7 March

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

2007 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab or infliximab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab or infliximab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial treatment of Crohn disease in a paediatric patient.

Initial PBS-subsidised treatment by a gastroenterologist, paediatrician or consultant physician as specified in the NOTE below, of a patient aged 6 to 17 years inclusive with moderate to severe refractory Crohn disease who satisfies the following criteria:

- (a) has confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or consultant physician as specified in the NOTE below; and
- (b) whose parent or authorised guardian has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (c) has failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including:
 - (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period;
 - (ii) an 8 week course of enteral nutrition;
 - (iii) immunosuppressive therapy including:
 - azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
 - 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
 - methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) severity of disease activity which results in a Paediatric Crohn Disease Activity Index (PCDAI) Score greater than or equal to 30 as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) The most recent PCDAI assessment must be no more than 1 month old at the time of application.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet including the date of assessment of the patient's condition; and
 - (ii) details of previous systemic drug therapy [dosage, date of commencement and duration of therapy], or dates of enteral nutrition; and
 - (iii) the signed patient acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete the 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

A PCDAI assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Continuing treatment of Crohn disease in a patient initiated on PBS-subsidised treatment as a paediatric patient.

Continuing PBS-subsidised treatment with infliximab by a gastroenterologist, paediatrician, consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of moderate to severe refractory Crohn disease; and
- (b) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as a reduction in Paediatric Crohn Disease Activity Index (PCDAI) Score by at least 15 points as compared to baseline AND a total PCDAI score of 30 points or less.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet along with the date of the assessment of the patient's condition.

The PCDAI assessment must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with infliximab, a PCDAI assessment of the patient's response must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

Patients who fail to demonstrate or sustain a response to treatment with infliximab for Crohn disease as specified in the criteria for continuing treatment with infliximab, will not be eligible to receive PBS-subsidised treatment with this drug within 12 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial PBS-subsidised treatment of Crohn disease in a paediatric patient who has previously received non-PBS-subsidised therapy with infliximab.

Initial PBS-subsidised supply for continuing treatment with infliximab by a gastroenterologist, paediatrician, consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient aged 6 to 17 years inclusive who:

- (a) has a documented history of moderate to severe refractory Crohn disease and was receiving treatment with infliximab prior to 4 July 2007; and
- (b) had a Paediatric Crohn Disease Activity Index (PCDAI) Score of greater than 30 prior to commencing treatment with infliximab. Where a baseline CDAI assessment is not available, please call Medicare Australia on 1800 700 270 to discuss; and
- (c) whose parent or authorised guardian has signed a patient acknowledgement indicating that they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) has demonstrated or sustained an adequate response to treatment with infliximab. For advice please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as a reduction in Paediatric Crohn Disease Activity Index (PCDAI) Score by at least 15 points as compared to baseline AND a total PCDAI score of 30 points or less.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and baseline Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet along with the date of the assessment of the patient's condition; and
 - (ii) the signed patient acknowledgement.

The current PCDAI assessment must be no more than 1 month old at the time of application. The baseline PCDAI assessment must be from immediately prior to commencing treatment with infliximab.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

Patients who fail to demonstrate or sustain a response to treatment with infliximab for Crohn disease as specified in the criteria for continuing treatment with infliximab, will not be eligible to recommence PBS-subsidised treatment with this drug within 12 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

9612X	Powder for I.V. infusion 100 mg	1	..	..	788.19	Remicade	SH

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Authority required

Initial treatment of complex refractory FISTULISING CROHN DISEASE.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist or a consultant physician as specified in the NOTE below, of a patient with complex refractory fistulising Crohn disease who:

- (a) has confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician as specified in the NOTE below; and
- (b) has an externally draining enterocutaneous or rectovaginal fistula; and

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(c) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criteria for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) a completed current Fistula Assessment Form including the date of assessment of the patient's condition; and
 - (ii) a signed patient acknowledgement.

The most recent fistula assessment must be no more than 1 month old at the time of application.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6 will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

An assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (up to 6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

A patient who fails to respond to a course of PBS-subsidised infliximab for the treatment of complex refractory fistulising Crohn disease will not be eligible to receive further PBS-subsidised treatment with infliximab for this condition within 12 months of the date on which treatment was ceased.

Authority required

Re-commencement of treatment of complex refractory FISTULISING CROHN DISEASE.

Re-initiation of PBS-subsidised treatment of complex refractory fistulising Crohn disease by a gastroenterologist or a consultant physician as specified in the NOTE below, of a patient with complex refractory fistulising Crohn disease who:

- (a) has a documented history of complex refractory fistulising Crohn disease; and
- (b) has an externally draining enterocutaneous or rectovaginal fistula; and
- (c) has previously received PBS-subsidised infliximab treatment for a draining enterocutaneous or rectovaginal fistula; and EITHER
- (d) has demonstrated or sustained an adequate response to the most recent course of PBS-subsidised treatment with infliximab for this condition; or
- (e) has failed to demonstrate or sustain an adequate response to PBS-subsidised treatment with infliximab for this condition and 12 months have elapsed from the date on which treatment was ceased.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes a completed current Fistula Assessment Form including the date of assessment of the patient's condition.

The most recent fistula assessment must be no more than 1 month old at the time of application.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

An assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (up to 6 weeks following

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

the third dose) so that there is adequate time for a response to be demonstrated.

This assessment must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

A patient who fails to respond to a course of PBS-subsidised infliximab for the treatment of complex refractory fistulising Crohn disease will not be eligible to receive further PBS-subsidised treatment with infliximab for this condition within 12 months of the date on which treatment was ceased.

Authority required

Initial PBS-subsidised treatment of complex refractory FISTULISING CROHN DISEASE in a patient who has previously received non-PBS-subsidised therapy with infliximab.

Initial PBS-subsidised supply for continuing treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below, or other consultant physician in consultation with a gastroenterologist of a patient who satisfies the following criteria:

- (a) has a documented history of complex refractory fistulising Crohn disease and was receiving treatment with infliximab prior to 1 March 2010; and
- (b) had a draining enterocutaneous or rectovaginal fistula(e) prior to commencing treatment with infliximab; and
- (c) has signed a patient acknowledgement indicating that they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criteria for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) is receiving treatment with infliximab at the time of application; and
- (e) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as:

- (a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or
- (b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) a completed current Fistula Assessment form including the date of assessment of the patient's condition; and
 - (ii) a signed patient acknowledgement.

The current fistula assessment must be no more than 1 month old at the time of application.

The baseline fistula assessment must be from immediately prior to commencing treatment with infliximab.

An assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criteria.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

A patient who fails to respond to a course of PBS-subsidised infliximab for the treatment of complex refractory fistulising Crohn disease will not be eligible to receive further PBS-subsidised treatment with infliximab for this condition within 12 months of the date on which treatment was ceased.

Authority required

Continuing treatment of complex refractory FISTULISING CROHN DISEASE.

Continuing PBS-subsidised treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below or other

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

consultant physician in consultation with a gastroenterologist, of a patient who:
(a) has a documented history of complex refractory fistulising Crohn disease; and
(b) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response is defined as:

- (a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or
- (b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes a completed Fistula Assessment form including the date of the assessment of the patient's condition.

The fistula assessment must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with infliximab, an assessment of the patient's response must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated.

An assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criteria.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

A patient who fails to respond to a course of PBS-subsidised infliximab for the treatment of complex refractory fistulising Crohn disease will not be eligible to receive further PBS-subsidised treatment with infliximab for this condition within 12 months of the date on which treatment was ceased.

9674E	Powder for I.V. infusion 100 mg	1	..	..	788.19	Remicade	SH

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, etanercept, infliximab and ustekinumab, for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, infliximab and ustekinumab.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

From 1 March 2010, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial adalimumab, etanercept, infliximab or ustekinumab without having to meet the initial treatment criteria, that is they will not need to experience a disease flare when swapping to an alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

A patient who received PBS-subsidised biological agent treatment for chronic plaque psoriasis prior to 1 March 2010 is considered to be in their first Cycle as of 1 March 2010.

Patients are eligible for PBS-subsidised treatment with only 1 biological agent at any 1 time.

Within the same Treatment Cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for a PBS-subsidised biological agent, they must change to an alternate agent if they wish to continue PBS-subsidised biological treatment. A patient who, prior to 1 March 2010, was authorised to receive PBS-subsidised initial treatment for chronic plaque psoriasis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2010.

Patients must be assessed for response to each course of continuing treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a Treatment Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological agent therapy before they are eligible to commence the next Cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological agent treatment in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Treatment Cycle.

Patients for whom a break in PBS-subsidised therapy of less than 5 years duration has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe chronic plaque psoriasis after 1 March 2010.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made in the following situations:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); or
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under '(4) Swapping therapy' below]; or
- (iii) patients who wish to re-commence treatment following a break in PBS-subsidised therapy with that agent (Initial 2).

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of treatment in the case of adalimumab and etanercept, 22 weeks of treatment in the case of infliximab and 28 weeks of treatment in the case of ustekinumab.

Grandfather patients (adalimumab and ustekinumab only).

Applications for patients who commenced treatment with adalimumab prior to 1 March 2009 or ustekinumab prior to 26 November 2009, may be made for initial PBS-subsidised treatment as continuing therapy under the initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with a biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment. Approval will be based on the criteria included in the relevant restriction.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological agent, a PASI assessment must be conducted after at least 12 weeks of treatment. This assessment must be submitted to Medicare Australia within 1 month of the completion of this initial treatment course. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(3) Application for continuing treatment.

Following the completion of an initial treatment course of a biological agent to which an adequate response has been demonstrated, patients may qualify to receive up to 24 weeks of continuing treatment with that biological agent. Patients are eligible to continue to receive continuous treatment with 24 week courses providing they continue to sustain a response.

For second and subsequent courses of PBS-subsidised treatment with adalimumab, etanercept, infliximab or ustekinumab it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to sustain a response to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

(4) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate agent within the same Treatment Cycle without having to requalify with respect to disease severity (i.e. a PASI score of greater than 15), or prior treatment requirements.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

Patients may trial an alternate biological agent at any time, regardless of whether they are receiving therapy with a biological agent at the time of the application or not. However, they cannot swap to a particular agent if they have failed to respond to treatment with that particular agent within the same Cycle.

Patients who commenced treatment with adalimumab prior to 1 June 2009 or ustekinumab prior to 1 March 2010 access these interchangeability arrangements in the same way as patients who have not.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the agent being ceased.

(5) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a Treatment Cycle and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatment applications.

(6) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent Biological Treatment Cycle, following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment according to the criteria of the relevant restriction and index of disease severity. Patients must have had at least 1 prior treatment, as listed in the criteria, for a minimum of 6 weeks, and must have a PASI assessment conducted preferably whilst still on treatment, but no later than 1 month following cessation of treatment. The PASI assessment must be no older than 1 month at the time of application.

Authority required

Initial treatment [Initial 1, Whole body (New patients — No prior biological agent)]

Initial treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis; and
- (b) have not received any prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (whole body); and
- (d) have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 3 of the following 4 treatments:
 - (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or
 - (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or
 - (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or
 - (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or where

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

phototherapy is contraindicated, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities including severity, associated with phototherapy, methotrexate, cyclosporin and acitretin, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]]; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 22 weeks of treatment with infliximab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 22 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Initial or re-Treatment [Initial 2, Whole body (Received prior biological agent under PBS)]

Treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have a documented history of severe chronic plaque psoriasis; and
- (b) have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have not failed PBS-subsidised therapy with infliximab for the treatment of this condition in the current Treatment Cycle.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]]; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

Applications for patients who have demonstrated a response to PBS-subsidised infliximab treatment within this Treatment Cycle and who wish to re-commence infliximab treatment within the same Cycle following a break in therapy, will only be approved where evidence of the patient's response to their most recent course of PBS-subsidised infliximab treatment has been submitted to Medicare Australia within 1 month of cessation of treatment.

A maximum of 22 weeks of treatment with infliximab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 22 weeks.

A PASI assessment of the patient's response to this course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Continuing treatment (Whole body)

Continuing PBS-subsidised treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over:

- (a) who have a documented history of severe chronic plaque psoriasis; and
- (b) whose most recent course of PBS-subsidised biological treatment for this condition in this Treatment Cycle was with infliximab; and
- (c) who have demonstrated an adequate response to their most recent course of treatment with infliximab.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-biological treatment baseline value for this Treatment Cycle.

This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with infliximab, the assessment of response must be after a minimum of 12 weeks of treatment with an initial course.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet along with the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with infliximab.

A maximum of 24 weeks of treatment with infliximab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for continuing authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks.

A PASI assessment of the patient's response must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Authority required

Initial treatment [Initial 1, Face, hand, foot (New patients — No prior biological agent)]

Initial treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- (b) have not received any prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (face, hand, foot); and
- (d) have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 3 of the following 4 treatments:
 - (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or
 - (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or
 - (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or
 - (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities including severity, associated with phototherapy, methotrexate, cyclosporin and acitretin, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)];
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 22 weeks of treatment with infliximab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 22 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Authority required

Initial or re-Treatment [Initial 2, Face, hand, foot (Received prior biological agent under PBS)]

Treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot; and

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (b) have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
(c) have not failed PBS-subsidised therapy with infliximab for the treatment of this condition in the current Treatment Cycle.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

Applications for patients who have demonstrated a response to PBS-subsidised infliximab treatment within this Treatment Cycle and who wish to re-commence infliximab treatment within the same Cycle following a break in therapy, will only be approved where evidence of the patient's response to their most recent course of PBS-subsidised infliximab treatment has been submitted to Medicare Australia within 1 month of cessation of treatment.

A maximum of 22 weeks of treatment with infliximab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 22 weeks.

A PASI assessment of the patient's response to this course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Continuing treatment (Face, hand, foot)

Continuing PBS-subsidised treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over:

- (a) who have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot; and
- (b) whose most recent course of PBS-subsidised biological treatment for this condition in this Treatment Cycle was with infliximab; and
- (c) who have demonstrated an adequate response to treatment with infliximab.

An adequate response to infliximab treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the pre-biological treatment baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the pre-biological treatment baseline value.

This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with infliximab, the assessment of response must be after a minimum of 12 weeks of treatment with an initial course.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams along with the date of the assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

The most recent PASI assessment must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with infliximab will be authorised under this restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for continuing authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks.

A PASI assessment of the patient's response must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for further continuing treatment must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

No applications for increased repeats will be authorised.

9617E	Powder for I.V. infusion 100 mg	1	..	..	788.19	Remicade	SH

Interleukin inhibitors

TOCILIZUMAB

Note

Any queries concerning the arrangements to prescribe tocilizumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Further prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe tocilizumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alpha antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alpha antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

— once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (new patient or patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with tocilizumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

least 20 mg weekly and one of which must be:

- hydroxychloroquine at a dose of at least 200 mg daily; or
- leflunomide at a dose of at least 10 mg daily; or
- sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Up to a maximum of 3 repeats of each strength may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab.

Patients who fail to demonstrate a response to treatment with tocilizumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with tocilizumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have a documented history of severe active rheumatoid arthritis; and
- (b) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with tocilizumab and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised tocilizumab treatment, within the timeframes specified below.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats of each strength may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised tocilizumab treatment was approved under either of the initial 1 or 2 treatment restrictions, patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised tocilizumab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Patients who fail to demonstrate a response to treatment with tocilizumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 3 ('grandfather' patients)

Initial PBS-subsidised supply for continuing treatment with tocilizumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of an adult who:

- (a) has a documented history of severe active rheumatoid arthritis; and
- (b) was receiving treatment with tocilizumab prior to 1 July 2009; and
- (c) has demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with tocilizumab; and
- (d) is receiving treatment with tocilizumab at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au]; and
- (3) a signed patient acknowledgement.

The same indices of disease severity used to establish baseline at the commencement of treatment with a bDMARD must be used for assessment of

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

all continuing applications.

The assessment of the patient's response to a continuing course of therapy must be made within 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled in order to ensure continuity of treatment for those patients who meet the continuation criterion.

A maximum of 24 weeks of treatment with tocilizumab will be approved under this criterion.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 5 repeats of each strength may be authorised.

Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Patients who fail to demonstrate a response to treatment with tocilizumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with tocilizumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have a documented history of severe active rheumatoid arthritis; and
- (b) who have demonstrated an adequate response to treatment with tocilizumab; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with tocilizumab.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 5 repeats of each strength may be authorised.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with tocilizumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with tocilizumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Patients who fail to demonstrate a response to treatment with tocilizumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

9671B	Concentrate for injection 80 mg in 4 mL	1	..	..	200.78	Actemra	RO
9672C	Concentrate for injection 200 mg in 10 mL	1	..	..	492.31	Actemra	RO
9673D	Concentrate for injection 400 mg in 20 mL	1	..	..	978.20	Actemra	RO

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Calcineurin inhibitors

CYCLOSPORIN

Caution

Careful monitoring of patients is mandatory.

Authority required

For use by organ or tissue transplant recipients.

6109M	Solution concentrate for I.V. infusion 50 mg in 1 mL	10	..	..	64.52	Sandimmun	NV

CYCLOSPORIN

Caution

Careful monitoring of patients is mandatory.

Authority required

Management of rejection in patients following organ or tissue transplantation, under the supervision and direction of a transplant unit.

Management includes initiation, stabilisation and review of therapy as required;

Management (which includes initiation, stabilisation and review of therapy) by dermatologists or clinical immunologists of patients with severe atopic dermatitis for whom other systemic therapies are ineffective or inappropriate;

Management (which includes initiation, stabilisation and review of therapy) by dermatologists of patients with severe psoriasis for whom other systemic therapies are ineffective or inappropriate and in whom the disease has caused significant interference with quality of life;

Management (which includes initiation, stabilisation and review of therapy) by nephrologists of patients with nephrotic syndrome in patients in whom steroids and cytostatic drugs have failed or are not tolerated or are considered inappropriate and in whom renal function is unimpaired;

Management (which includes initiation, stabilisation and review of therapy) by rheumatologists or clinical immunologists of patients with severe active rheumatoid arthritis for whom classical slow-acting anti-rheumatic agents (including methotrexate) are ineffective or inappropriate.

6125J	Oral liquid 100 mg per mL, 50 mL	4	5	..	*1309.58	Neoral	NV
6232B	Capsule 10 mg	120	5	..	*84.82	Neoral 10	NV
6352H	Capsule 25 mg	120	5	..	*169.38	^a Cicloral	SZ
				^B 3.88	*173.26	^a Neoral 25	NV
6353J	Capsule 50 mg	120	5	..	*345.50	^a Cicloral	SZ
				^B 4.20	*349.70	^a Neoral 50	NV
6354K	Capsule 100 mg	120	5	..	*697.34	^a Cicloral	SZ
				^B 4.16	*701.50	^a Neoral 100	NV

TACROLIMUS

Caution

Careful monitoring of patients is mandatory.

Authority required

Management of rejection in patients following organ or tissue transplantation, under the supervision and direction of a transplant unit.

Management includes initiation, stabilisation and review of therapy as required.

6216E	Capsule 1 mg	200	5	..	*688.32	^a Prograf ^a Tacrolimus Sandoz	JC SZ
6217F	Capsule 5 mg	100	5	..	*1684.80	^a Prograf ^a Tacrolimus Sandoz	JC SZ
6328C	Capsule 500 micrograms	200	5	..	*347.38	^a Prograf ^a Tacrolimus Sandoz	JC SZ
9681M	Capsule 0.5 mg (once daily prolonged release)	60	5	..	*108.78	Prograf XL	JC
9682N	Capsule 1 mg (once daily prolonged release)	120	5	..	*415.56	Prograf XL	JC
9683P	Capsule 5 mg (once daily prolonged release)	60	5	..	*1029.30	Prograf XL	JC

Other immunosuppressants

LENALIDOMIDE

Note

Any queries concerning the arrangements to prescribe lenalidomide may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Any queries concerning patients who are enrolled on the Lenalidomide Compassionate program may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). These patients must demonstrate they met initial criteria prior to commencing treatment on the compassionate program and also demonstrate they do not have progressive disease. Baseline and current pathology reports must be submitted with the initial application.

Applications for authority to prescribe lenalidomide should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001.

Authority required

Initial PBS-subsidised treatment, as monotherapy or in combination with dexamethasone, of a patient with a histological diagnosis of multiple myeloma who has progressive disease after at least 1 prior therapy and who has undergone or is ineligible for a primary stem cell transplant. The patient must have experienced treatment failure after a trial of at least four (4) weeks of thalidomide at a dose of at least 100 mg daily or have failed to achieve at least a minimal response after eight (8) or more weeks of thalidomide-based therapy for progressive disease.

If the dosing requirement for thalidomide cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein and less than 200 mg per 24 hour Bence-Jones proteinuria.

Thalidomide treatment failure is defined as:

- (1) confirmed disease progression during thalidomide treatment or within 6 months of discontinuing thalidomide treatment; or
- (2) severe intolerance or toxicity unresponsive to clinically appropriate dose adjustment.

Severe intolerance due to thalidomide is defined as unacceptable somnolence or sedation interfering with activities of daily living.

Toxicity from thalidomide is defined as peripheral neuropathy (Grade 2 or greater, interfering with function), drug-related seizures, serious Grade 3 or 4 drug-related dermatological reactions, such as Stevens-Johnson Syndrome, or other Grade 3 or 4 toxicity.

Failure to achieve at least a minimal response after 8 or more weeks of thalidomide-based therapy for progressive disease is defined as:

- (1) less than a 25% reduction in serum or urine M protein; or
- (2) in oligo-secretory and non-secretory myeloma patients only, less than a 25% reduction in the difference between involved and uninvolved serum free light chain levels.

Lenalidomide will only be subsidised for patients with multiple myeloma who are not receiving concomitant PBS-subsidised bortezomib.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Multiple Myeloma Authority Application - Supporting Information Form, which includes details of the histological diagnosis of multiple myeloma, prior treatments including name(s) of drug(s) and date of most recent treatment cycle and record of prior stem cell transplant or ineligibility for prior stem cell transplant; details of thalidomide treatment failure; details of the basis of the diagnosis of progressive disease or failure to respond; and nomination of which disease activity parameters will be used to assess response.

To enable confirmation by Medicare Australia, current diagnostic reports of at least one of the following are required:

- (a) the level of serum monoclonal protein; or
- (b) Bence-Jones proteinuria — the results of 24-hour urinary light chain M protein excretion; or
- (c) the serum level of free kappa and lambda light chains; or
- (d) bone marrow aspirate or trephine; or
- (e) if present, the size and location of lytic bone lesions (not including compression fractures); or
- (f) if present, the size and location of all soft tissue plasmacytomas by clinical or radiographic examination i.e. MRI or CT-scan; or
- (g) if present, the level of hypercalcaemia, corrected for albumin concentration.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

As these parameters will be used to determine response, results for either (a) or (b) or (c) should be provided for all patients. Where the patient has oligo-secretory or non-secretory multiple myeloma, either (c) or (d) or if relevant (e), (f) or (g) should be provided. Where the prescriber plans to assess response in patients with oligo-secretory or non-secretory multiple myeloma with free light chain assays, evidence of the oligo-secretory or non-secretory nature of the multiple myeloma (either previous or current serum M protein less than 10 g per L and urinary Bence-Jones protein undetectable or less than 200 mg per 24 hours) must be provided; and
(3) duration of thalidomide and daily dose prescribed; and
(4) a signed patient acknowledgment.

Note

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

Authority required

Continuing PBS-subsidised treatment, as monotherapy or in combination with dexamethasone, of multiple myeloma in a patient who has previously been issued with an authority prescription for lenalidomide and who does not have progressive disease.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Authority applications for continuing treatment may be made by telephone to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

Note

Special Pricing Arrangements apply.

9642L	Capsule 5 mg	21	..	..	5438.80	Revlimid	CJ
9643M	Capsule 10 mg	21	..	..	5689.75	Revlimid	CJ
9644N	Capsule 15 mg	21	..	..	6628.03	Revlimid	CJ
9645P	Capsule 25 mg	21	..	..	6980.62	Revlimid	CJ

RITUXIMAB

Note

Any queries concerning the arrangements to prescribe rituximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Further prescribing information (including Authority Application Forms) is on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe rituximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alpha antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Authority required

Initial 1 (patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with rituximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have failed to respond to at least 1 PBS-subsidised TNF-alfa antagonist; and
- (c) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (d) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of two infusions will be authorised under this restriction.

Assessment of a patient's response to an initial course of treatment must be made at least 12 weeks after the first infusion so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia within 4 weeks of the date it was conducted.

Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with rituximab.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

Patients who fail to demonstrate a response to treatment with rituximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Patients who fail to demonstrate a response to rituximab treatment and who qualify to trial an alternate bDMARD according to the interchangeability arrangements for bDMARDs for the treatment of severe rheumatoid arthritis, may do so without having to have a 22 week treatment-free period.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with rituximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have a documented history of severe active rheumatoid arthritis; and
- (b) have failed to respond to at least 1 PBS-subsidised TNF-alfa antagonist; and
- (c) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with rituximab and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised rituximab treatment, within the timeframes specified below.

A maximum of two infusions will be authorised under this restriction.

Where the most recent course of PBS-subsidised rituximab treatment was approved under either of the initial 1 or 2 treatment restrictions patients must be assessed for response at least 12 weeks after the first infusion. This assessment must be provided to Medicare Australia no later than 4 weeks from the date of assessment.

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent provided they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The demonstration of response must be submitted to Medicare Australia within 4 weeks of assessment.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

Patients who fail to demonstrate a response to treatment with rituximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Patients who fail to demonstrate a response to rituximab treatment and who qualify to trial an alternate bDMARD according to the interchangeability arrangements for bDMARDs for the treatment of severe rheumatoid arthritis, may do so without having to have a 22 week treatment-free period.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with rituximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have a documented history of severe active rheumatoid arthritis; and
- (b) who have demonstrated an adequate response to treatment with rituximab; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with rituximab.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of two infusions will be authorised under this restriction.

Patients may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The demonstration of response must be submitted to Medicare Australia within 4 weeks of assessment.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

Patients who fail to demonstrate a response to treatment with rituximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

9611W	Solution for I.V. infusion 500 mg in 50 mL	1	..	..	2309.99	Mabthera	RO

THALIDOMIDE

Caution

Thalidomide is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and for 1 month after cessation of treatment.

Authority required

Multiple myeloma.

Note

Patients receiving thalidomide under the PBS listing must be registered in the i-access risk management program.

6469L	Capsule 50 mg	112	..	..	*1726.42	Thalomid	CJ
9684Q	Capsule 100 mg	56	..	..	*1726.42	Thalomid	CJ

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Musculo-skeletal system

Muscle relaxants

Muscle relaxants, centrally acting agents

Other centrally acting agents

BACLOFEN

Authority required

Severe chronic spasticity, where oral antispastic agents have failed or have caused unacceptable side effects, in patients with chronic spasticity of cerebral origin;

Severe chronic spasticity, where oral antispastic agents have failed or have caused unacceptable side effects, in patients with chronic spasticity due to multiple sclerosis;

Severe chronic spasticity, where oral antispastic agents have failed or have caused unacceptable side effects, in patients with chronic spasticity due to spinal cord injury;

Severe chronic spasticity, where oral antispastic agents have failed or have caused unacceptable side effects, in patients with chronic spasticity due to spinal cord disease.

6284R	Intrathecal injection 10 mg in 5 mL	10	..	..	*1530.12		Lioresal Intrathecal	NV

Drugs for treatment of bone diseases

Drugs affecting bone structure and mineralization

Bisphosphonates

DISODIUM PAMIDRONATE

Authority required

Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.

Note

The concentrated injection 15 mg and powder for I.V. infusion 15 mg (after reconstitution) are bioequivalent.

6286W	Concentrated injection 15 mg in 5 mL	4	2	..	*229.18	^a	Pamisol	HH
6290C	Injection set containing 4 vials powder for I.V. infusion 15 mg and 4 ampoules solvent 5 mL	1	2	..	229.18	^a	Aredia 15 mg	NV

DISODIUM PAMIDRONATE

Authority required

Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.

Note

The concentrated injection 30 mg and powder for I.V. infusion 30 mg (after reconstitution) are bioequivalent.

6279L	Injection set containing 2 vials powder for I.V. infusion 30 mg and 2 ampoules solvent 10 mL	1	2	..	229.18	^a	Aredia 30 mg	NV
6287X	Concentrated injection 30 mg in 10 mL	2	2	..	*229.18	^a	Pamisol	HH

DISODIUM PAMIDRONATE

Authority required

Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.

6288Y	Concentrated injection 60 mg in 10 mL	1	2	..	229.17		Pamisol	HH

DISODIUM PAMIDRONATE

Authority required

Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.

Authority required

Multiple myeloma;

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer	
					Max. Qty	\$		
	Bone metastases from breast cancer.							
	<u>Note</u>							
	The concentrated injection 90 mg and powder for I.V. infusion 90 mg (after reconstitution) are bioequivalent.							
6223M	Injection set containing 1 vial powder for I.V. infusion 90 mg and 1 ampoule solvent 10 mL	1	11	..	340.55	^a	Aredia 90 mg	NV
6289B	Concentrated injection 90 mg in 10 mL	1	11	..	340.55	^a	Pamisol	HH
	IBANDRONIC ACID							
	<u>Authority required</u>							
	Bone metastases from breast cancer.							
9619G	Concentrated injection for I.V. infusion 6 mg (as ibandronate sodium monohydrate) in 6 mL	1	11	..	361.43		Bondronat	HH
	ZOLEDRONIC ACID							
	<u>Authority required</u>							
	Multiple myeloma;							
	Bone metastases from breast cancer;							
	Bone metastases from hormone-resistant prostate cancer, with demonstration of biochemical progression of disease despite maximal therapy with hormonal treatments;							
	Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.							
	<u>Note</u>							
	Special Pricing Arrangements apply.							
6371H	Injection concentrate for I.V. infusion 4 mg (as monohydrate) in 5 mL	1	11	..	474.42		Zometa	NV

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Nervous system

Anti-Parkinson drugs

Dopaminergic agents

Dopamine agonists

APOMORPHINE HYDROCHLORIDE

Authority required

Parkinson's disease in patients severely disabled by motor fluctuations which do not respond to other therapy.

9607P	Injection 20 mg in 2 mL	5	..	..	88.28	Apomine	HH
9640J	Injection 50 mg in 5 mL	5	..	..	208.86	APO-go	HH
9647R	Solution for subcutaneous infusion 50 mg in 10 mL pre-filled syringe	5	..	..	208.86	Apomine PFS	HH

Psycholeptics

Antipsychotics

Diazepines, oxazepines, thiazepines and oxepines

CLOZAPINE

Authority required

Schizophrenia in patients who are non-responsive to other neuroleptic agents;

Schizophrenia in patients who are intolerant of other neuroleptic agents.

6101D	Tablet 25 mg	100	..	..	79.57	^a Clopine 25	HH
						^a Clozaril 25	NV
6102E	Tablet 100 mg	100	..	..	276.10	^a Clopine 100	HH
						^a Clozaril 100	NV
6417R	Tablet 50 mg	100	..	..	150.25	Clopine 50	HH
6418T	Tablet 200 mg	100	..	..	545.78	Clopine 200	HH
9632Y	Oral liquid 50 mg per mL, 100 mL	1	..	..	146.82	Clopine Suspension	HH

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Respiratory system

Cough and cold preparations

Expectorants, excl. combinations with cough suppressants

Mucolytics

DORNASE ALFA

Authority required

Use by cystic fibrosis patients who satisfy all of the following criteria:

- (1) are 5 years of age or older;
- (2) have a FVC greater than 40% predicted for age, gender and height;
- (3) have evidence of chronic suppurative lung disease (cough and sputum most days of the week, or greater than 3 respiratory tract infections of more than 2 weeks' duration in any 12 months, or objective evidence of obstructive airways disease);
- (4) are participating in a 4 week trial as detailed below or have achieved a 10% or greater improvement in FEV1 (compared to baseline established prior to dornase alfa treatment) after a 4 week trial.

In order for patients to be eligible for participation in the HSD program, the following conditions must be met:

- (1) Patients must be assessed at cystic fibrosis clinics/centres which are under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis and the prescribing of dornase alfa under the HSD program is limited to such physicians. If attendance at such units is not possible because of geographical isolation, management (including prescribing) may be by specialist physician or paediatrician in consultation with such a unit;
- (2) The measurement of lung function is to be conducted by independent (other than the treating doctor) experienced personnel at established lung function testing laboratories, unless this is not possible because of geographical isolation;
- (3) Prior to dornase alfa therapy, a baseline measurement of FEV1 must be undertaken during a stable period of the disease;
- (4) Initial therapy is limited to 4 weeks' treatment with dornase alfa at a dose of 2.5 mg daily;
- (5) At or towards the end of the initial 4 weeks' trial, patients must be reassessed and a further FEV1 measurement be undertaken (single test under conditions as above). Patients who achieve a 10% or greater improvement in FEV1 (compared to baseline established prior to dornase alfa treatment) are eligible for continued subsidy under the HSD program at a dose of 2.5 mg daily;
- (6) Patients who fail to meet a 10% or greater improvement in FEV1 after the initial 4 weeks' treatment at a dose of 2.5 mg daily, may have 1 further trial in the next 12 months but not before 3 months after the initial trial;
- (7) Following an initial 6 months' therapy, a global assessment must be undertaken involving the patient, the patient's family (in the case of paediatric patients) and the treating physician(s) to establish that all agree that dornase alfa treatment is continuing to produce worthwhile benefits. (Dornase alfa therapy should cease if there is not general agreement of benefit as there is always the possibility of harm from unnecessary use.) Further reassessments are to be undertaken at six-monthly intervals;
- (8) Other aspects of treatment, such as physiotherapy, must be continued;
- (9) Where there is documented evidence that a patient already receiving dornase alfa therapy would have met the criteria for subsidy (i.e. satisfied the criteria for the 4 week trial and achieved a 10% or greater improvement in FEV1) then the patient is eligible to continue treatment under the HSD program. Where such evidence is not available, patients will need to satisfy the initiation and continuation criteria as for new patients. (Four weeks is considered a suitable wash-out period).

Note

It is highly desirable that all patients be included in the national cystic fibrosis patient data-base.

Authority required

Treatment of cystic fibrosis in a patient less than 5 years of age who has:

- (1) A severe clinical course with frequent respiratory exacerbations or chronic respiratory symptoms (including chronic or recurrent cough, wheeze or tachypnoea) requiring frequent hospital admissions more frequently than 3 times per year; or
- (2) Significant bronchiectasis on chest high resolution computed tomography scan; or
- (3) Severe cystic fibrosis bronchiolitis with persistent wheeze non-responsive to conventional medicines; or
- (4) Severe physiological deficit measure by forced oscillation technique or multiple breath nitrogen washout and failure to respond to conventional therapy.

In order for the patient to be eligible for participation in the HSD program, the following conditions must be met:

- (1) The patient must be assessed at a cystic fibrosis clinic/centre which is under the supervision of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis, and the prescribing of dornase alfa under the HSD program is limited to such physicians. If attendance at such a unit is not possible because of geographical isolation, management (including prescribing) may be by specialist physician or paediatrician in consultation with such a unit;
- (2) Following an initial 6 months therapy, a comprehensive assessment must be undertaken and documented involving the patient, the patient's family, the treating physician and an additional independent member of the cystic fibrosis treatment team to establish agreement that dornase alfa treatment is continuing to produce worthwhile benefit. Treatment with dornase alfa should cease if there is not agreement of benefit as there is always the possibility of harm from unnecessary use. Further reassessments are to be undertaken and documented yearly.

Note

It is highly desirable that all patients be included in the national cystic fibrosis patient data-base.

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Authority required

Grandfather — continuing for patients five years or older

Continuation of treatment of cystic fibrosis in a patient 5 years of age or older, who initiated treatment with dornase alfa at an age of less than 5 years and for whom a comprehensive assessment, involving the patient's family, the treating physician and an additional independent member of the cystic fibrosis treatment team, documents agreement that dornase alfa treatment is continuing to produce worthwhile benefit. Further reassessments are to be undertaken and documented yearly. Treatment with dornase alfa should cease if there is not agreement of benefit as there is always the possibility of harm from unnecessary use.

Note

It is highly desirable that all patients be included in the national cystic fibrosis patient data-base.

Authority required

Grandfather — for patients less than five years of age who initiated dornase alfa prior to listing

Treatment of cystic fibrosis in a patient less than 5 years of age who initiated treatment with dornase alfa prior to 1 November 2009 and for whom a comprehensive assessment, involving the patient's family, the treating physician and an additional independent member of the cystic fibrosis treatment team, documents agreement that dornase alfa treatment is continuing to produce worthwhile benefit. Further reassessments are to be undertaken and documented yearly. Treatment with dornase alfa should cease if there is not agreement of benefit as there is always the possibility of harm from unnecessary use.

Note

It is highly desirable that all patients be included in the national cystic fibrosis patient data-base.

6120D	Solution for inhalation 2.5 mg (2,500 units) in 2.5 mL	60	5	..	*2406.42	Pulmozyme	RO

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Sensory organs

Ophthalmologicals

Antiinfectives

Antivirals

GANCICLOVIR

Authority required

Cytomegalovirus retinitis in severely immunocompromised patients.

6256G	Intravitreal implant 4.5 mg	1	..	..	6046.42	Vitrasert	BU

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Various

All other therapeutic products

All other therapeutic products

Iron chelating agents

DEFERASIROX

Authority required

Chronic iron overload in adults, adolescents and children 6 years and older associated with disorders of erythropoiesis;

Chronic iron overload in paediatric patients aged 2 to 5 years, associated with disorders of erythropoiesis, who are intolerant to desferrioxamine or in whom desferrioxamine has proven ineffective.

Note

Special Pricing Arrangements apply.

6499C	Tablet 125 mg (dispersible)	168	5	..	*1447.92	Exjade	NV
6500D	Tablet 250 mg (dispersible)	168	5	..	*2849.34	Exjade	NV
9600G	Tablet 500 mg (dispersible)	168	5	..	*5652.24	Exjade	NV

DEFERIPRONE

Authority required

Iron overload in patients with thalassaemia major who are unable to take desferrioxamine therapy;

Iron overload in patients with thalassaemia major in whom desferrioxamine therapy has proven ineffective.

6416Q	Tablet 500 mg	600	5	..	*2749.80	Ferriprox	OA
9638G	Oral solution 100 mg per mL, 250 mL	5	5	..	*1172.82	Ferriprox	OA

DEFERRIOXAMINE MESYLATE

Authority required

Disorders of erythropoiesis associated with treatment-related chronic iron overload.

6113R	Powder for injection 500 mg	400	5	..	*3849.62	^a Hospira Pty Limited	HH
				^B 314.80	*4164.42	^a Desferal 500 mg	NV
6270B	Powder for injection 2 g	60	5	..	*2328.42	^a Hospira Pty Limited	HH
				^B 23.40	*2351.82	^a Desferal 2 g	NV

Drugs for treatment of hyperkalemia and hyperphosphatemia

LANTHANUM

Authority required

Management of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where serum phosphate is greater than 1.6 mmol per L at the commencement of therapy.

Management includes initiation, stabilisation and review of therapy as required;

Management of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where the serum calcium times phosphate product is greater than 4.0 at the commencement of therapy.

Management includes initiation, stabilisation and review of therapy as required.

Note

Not to be used in combination with sevelamer.

9635D	Tablet, chewable, 500 mg (as carbonate hydrate)	180	5	..	*550.90	Fosrenol	ZI
9636E	Tablet, chewable, 750 mg (as carbonate hydrate)	180	5	..	*828.60	Fosrenol	ZI
9637F	Tablet, chewable, 1000 mg (as carbonate hydrate)	180	5	..	*932.04	Fosrenol	ZI

SEVELAMER HYDROCHLORIDE

Authority required

Management of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where serum phosphate is greater than 1.6 mmol per L at the commencement of therapy.

Management includes initiation, stabilisation and review of therapy as required;

HIGHLY SPECIALISED DRUGS PROGRAM (Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Management of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where the serum calcium times phosphate product is greater than 4.0 at the commencement of therapy.

Management includes initiation, stabilisation and review of therapy as required.

Note

Not to be used in combination with lanthanum.

9620H	Tablet 800 mg	360	5	..	*651.22	Renagel	GZ

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Blood and blood forming organs

Antianemic preparations

Other antianemic preparations

Other antianemic preparations

DARBEPOETIN ALFA

Authority required (STREAMLINED)

3334

Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.

5637Q	Injection 10 micrograms in 0.4 mL pre-filled syringe	8	5	..	*356.08	Aranesp	AN
5638R	Injection 20 micrograms in 0.5 mL pre-filled syringe	8	5	..	*670.62	Aranesp	AN
5639T	Injection 30 micrograms in 0.3 mL pre-filled syringe	8	5	..	*917.46	Aranesp	AN
5640W	Injection 40 micrograms in 0.4 mL pre-filled syringe	8	5	..	*1113.60	Aranesp	AN
5641X	Injection 50 micrograms in 0.5 mL pre-filled syringe	8	5	..	*1376.78	Aranesp	AN
5642Y	Injection 60 micrograms in 0.3 mL pre-filled syringe	8	5	..	*1616.66	Aranesp	AN
5643B	Injection 150 micrograms in 0.3 mL pre-filled syringe	8	5	..	*3904.50	Aranesp	AN
5644C	Injection 80 micrograms in 0.4 mL pre-filled syringe	8	5	..	*2128.00	Aranesp	AN
5645D	Injection 20 micrograms in 0.5 mL pre-filled injection pen	8	5	..	*670.64	Aranesp SureClick	AN
5646E	Injection 40 micrograms in 0.4 mL pre-filled injection pen	8	5	..	*1113.60	Aranesp SureClick	AN
5647F	Injection 60 micrograms in 0.3 mL pre-filled injection pen	8	5	..	*1616.64	Aranesp SureClick	AN
5648G	Injection 80 micrograms in 0.4 mL pre-filled injection pen	8	5	..	*2128.00	Aranesp SureClick	AN
5649H	Injection 100 micrograms in 0.5 mL pre-filled injection pen	8	5	..	*2620.48	Aranesp SureClick	AN
5650J	Injection 150 micrograms in 0.3 mL pre-filled injection pen	8	5	..	*3904.48	Aranesp SureClick	AN
5651K	Injection 100 micrograms in 0.5 mL pre-filled syringe	8	5	..	*2620.50	Aranesp	AN

EPOETIN ALFA

Authority required (STREAMLINED)

3334

Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.

5713Q	Injection 20,000 units in 0.5 mL pre-filled syringe	12	5	..	*3876.00	Eprex 20,000	JC
5714R	Injection 1,000 units in 0.5 mL pre-filled syringe	12	5	..	*279.30	Eprex 1000	JC
5715T	Injection 5,000 units in 0.5 mL pre-filled syringe	12	5	..	*1057.34	Eprex 5000	JC
5716W	Injection 6,000 units in 0.6 mL pre-filled syringe	12	5	..	*1255.14	Eprex 6000	JC
5717X	Injection 8,000 units in 0.8 mL pre-filled syringe	12	5	..	*1627.92	Eprex 8000	JC
5718Y	Injection 40,000 units in 1 mL pre-filled syringe	2	5	..	*1254.00	Eprex 40,000	JC
5719B	Injection 2,000 units in 0.5 mL pre-filled syringe	12	5	..	*516.80	Eprex 2000	JC
5720C	Injection 3,000 units in 0.3 mL pre-filled syringe	12	5	..	*666.90	Eprex 3000	JC
5721D	Injection 4,000 units in 0.4 mL pre-filled syringe	12	5	..	*849.30	Eprex 4000	JC
5722E	Injection 10,000 units in 1 mL pre-filled syringe	12	5	..	*1970.30	Eprex 10000	JC
5723F	Injection 30,000 units in 0.75 mL pre-filled syringe	12	5	..	*5728.50	Eprex 30,000	JC

EPOETIN BETA

Authority required (STREAMLINED)

3334

Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.

5724G	Injection 2,000 units in 0.3 mL pre-filled syringe	12	5	..	*516.80	NeoRecormon	RO
5725H	Injection 3,000 units in 0.3 mL pre-filled syringe	12	5	..	*666.90	NeoRecormon	RO
5726J	Injection 4,000 units in 0.3 mL pre-filled syringe	12	5	..	*849.30	NeoRecormon	RO

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
5727K	Injection 5,000 units in 0.3 mL pre-filled syringe	12	5	..	*1057.36		NeoRecormon RO
5728L	Injection 6,000 units in 0.3 mL pre-filled syringe	12	5	..	*1255.14		NeoRecormon RO
5729M	Injection 10,000 units in 0.6 mL pre-filled syringe	12	5	..	*1970.30		NeoRecormon RO
5730N	Injection 20,000 units in 0.6 mL pre-filled syringe	12	5	..	*3876.00		NeoRecormon RO

EPOETIN LAMBDA

Note

Epoetin lambda should only be administered by the intravenous route.

Authority required (STREAMLINED)

3334

Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.

9587N	Injection 4,000 units in 0.4 mL pre-filled syringe	12	5	..	*804.60		Novicrit NV
9589Q	Injection 5,000 units in 0.5 mL pre-filled syringe	12	5	..	*1001.70		Novicrit NV
9591T	Injection 6,000 units in 0.6 mL pre-filled syringe	12	5	..	*1189.08		Novicrit NV
9594Y	Injection 8,000 units in 0.8 mL pre-filled syringe	12	5	..	*1542.24		Novicrit NV
9596C	Injection 10,000 units in 1 mL pre-filled syringe	12	5	..	*1866.60		Novicrit NV
9668W	Injection 1,000 units in 0.5 mL pre-filled syringe	12	5	..	*264.60		Novicrit NV
9669X	Injection 2,000 units in 1 mL pre-filled syringe	12	5	..	*489.60		Novicrit NV
9670Y	Injection 3,000 units in 0.3 mL pre-filled syringe	12	5	..	*631.80		Novicrit NV

METHOXY POLYETHYLENE GLYCOL-EPOETIN BETA

Authority required (STREAMLINED)

3334

Treatment of anaemia requiring transfusion, defined as a haemoglobin level of less than 100 g per L, where intrinsic renal disease, as assessed by a nephrologist, is the primary cause of the anaemia.

5794Y	Injection 30 micrograms in 0.3 mL pre-filled syringe	2	5	..	*369.18		Mircera RO
5795B	Injection 50 micrograms in 0.3 mL pre-filled syringe	2	5	..	*615.30		Mircera RO
5796C	Injection 75 micrograms in 0.3 mL pre-filled syringe	2	5	..	*896.02		Mircera RO
5797D	Injection 100 micrograms in 0.3 mL pre-filled syringe	2	5	..	*1158.82		Mircera RO
5798E	Injection 120 micrograms in 0.3 mL pre-filled syringe	2	5	..	*1341.64		Mircera RO
5799F	Injection 200 micrograms in 0.3 mL pre-filled syringe	2	5	..	*1924.30		Mircera RO
5800G	Injection 360 micrograms in 0.6 mL pre-filled syringe	2	5	..	*3326.52		Mircera RO

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Cardiovascular system

Antihypertensives

Other antihypertensives

Other antihypertensives

AMBRISENTAN

Caution

Ambrisentan is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of treatment with this drug.

Note

Any queries concerning the arrangements to prescribe ambrisentan may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(c) epoprostenol sodium, of:

— primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND

— primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND

(d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND

(e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND

(f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or

(ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or

(iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments;

(2) RHC composite assessment plus 6MWT;

(3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

(1) ECHO composite assessment plus 6MWT;

(2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments plus 6MWT;

(2) RHC plus ECHO composite assessments;

(3) RHC composite assessment plus 6MWT;

(4) ECHO composite assessment plus 6MWT;

(5) RHC composite assessment only;

(6) ECHO composite assessment only.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent. All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with ambrisentan of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with ambrisentan of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (c) WHO Functional Class IV primary pulmonary hypertension; OR
- (d) WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (grandfather patients)

Application for initial PBS-subsidised treatment with ambrisentan of patients who were receiving treatment with ambrisentan prior to 1 December 2009 and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease; OR
- (c) WHO Functional Class IV primary pulmonary hypertension; OR
- (d) WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) for patients who have received less than 6 months of ambrisentan treatment at the time of application — a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results of the following 3 tests, where available, at the time treatment with ambrisentan was commenced:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) the date of commencement of ambrisentan treatment; and
- (4) a signed patient acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. The number of repeats authorised will be dependent on the duration of prior ambrisentan therapy. Where patients have received less than 6 months of non-PBS-subsidised treatment with ambrisentan, sufficient repeats to allow the patient to complete a total of 6 months of combined PBS-subsidised and non-PBS-subsidised therapy may be requested. Where fewer than the maximum allowable number of repeats are requested at the time of application, authority approvals for the remainder of the allowable repeats may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients)

Application for initial treatment with ambrisentan of patients with one of the following:

- (a) primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who wish to re-commence PBS-subsidised ambrisentan after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with ambrisentan; OR
- (b) primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with an alternate PAH agent other than ambrisentan.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with ambrisentan of patients who have received approval for initial PBS-subsidised treatment with ambrisentan and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of ambrisentan treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(1) a completed authority prescription form; and
 (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 (i) RHC composite assessment; and
 (ii) ECHO composite assessment; and
 (iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats will be authorised. Where fewer than 5 repeats are initially requested under this criterion, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note

Special Pricing Arrangements apply.

5607D	Tablet 5 mg	30	..	..	4035.00	Volibris	GK
5608E	Tablet 10 mg	30	..	..	4035.00	Volibris	GK

BOSENTAN MONOHYDRATE

Caution

Bosentan monohydrate is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of treatment with this drug.

Note

Any queries concerning the arrangements to prescribe bosentan monohydrate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
 Prior Written Approval of Specialised Drugs
 Reply Paid 9826
 GPO Box 9826
 HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (1) ECHO composite assessment plus 6MWT;
(2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent. All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new adult patients)

Application for initial PBS-subsidised treatment with bosentan monohydrate of adult patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to scleroderma and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new adult patients)

Application for initial PBS-subsidised treatment with bosentan monohydrate of adult patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to scleroderma and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (c) WHO Functional Class IV primary pulmonary hypertension; OR
- (d) WHO Functional Class IV pulmonary arterial hypertension secondary to scleroderma.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients under 18 years of age)

Application for initial PBS-subsidised treatment with bosentan monohydrate of patients aged less than 18 years who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

WHO Functional Class III primary pulmonary hypertension and either a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate prior vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a patient and prescriber acknowledgment, signed by the parent or authorised guardian, indicating that they understand and acknowledge that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients under 18 years of age)

Application for initial PBS-subsidised treatment with bosentan monohydrate of patients aged less than 18 years who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and either a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class IV primary pulmonary hypertension.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a patient and prescriber acknowledgment, signed by the parent or authorised guardian, indicating that they understand and acknowledge that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with bosentan monohydrate of a patient who has been assessed by a physician from a designated hospital to have WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment (and signed by the parent or authorised guardian for patients under 18 years of age) indicating that the patient understands and acknowledges that PBS-subsidised treatment with bosentan monohydrate will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for adult patients)

Application for initial treatment with bosentan monohydrate of adult patients with one of the following:

- (a) primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), who wish to re-commence PBS-subsidised bosentan monohydrate after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with bosentan monohydrate; OR
- (b) primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma and whose most recent course of PBS-subsidised treatment was with an alternate PAH agent other than bosentan monohydrate.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for patients under 18 years of age)

Application for initial treatment with bosentan monohydrate of patients aged less than 18 years with one of the following:

- (a) primary pulmonary hypertension, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), who wish to re-commence PBS-subsidised bosentan monohydrate after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with bosentan monohydrate; OR
- (b) primary pulmonary hypertension and whose most recent course of PBS-subsidised treatment was with a PAH agent other than bosentan monohydrate.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms [see Note for authority approval requirements]; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for the first authority prescription issued under this criterion [see Note for full details of authority approval requirements]. A maximum of 4 repeats will be authorised for the second authority prescription issued under this criterion. Where fewer than 4 repeats are initially requested with the second authority prescription, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with bosentan monohydrate of patients who have received approval for initial PBS-subsidised treatment with bosentan monohydrate and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of bosentan monohydrate treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats will be authorised.

Where fewer than 5 repeats are initially requested under this criterion, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Cessation of treatment (all patients)

Final PBS-subsidised supply for patients with WHO Functional Class III or IV primary pulmonary hypertension or WHO Functional Class III or IV pulmonary arterial hypertension secondary to scleroderma or WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), who have not responded to bosentan monohydrate therapy [see Note for definition of response], to allow for gradual cessation of treatment.

Applications for authorisation under this criterion should be made on the telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) [see Note on authority approval requirements].

Approval will only be granted for the 62.5 mg tablet strength. The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment.

Under no circumstances will telephone approvals be granted for treatment that would extend the final treatment period beyond 1 month.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
<u>Note</u>							
Special Pricing Arrangements apply.							
5618Q	Tablet 62.5 mg (base)	60	..	..	4035.00	Tracleer	AT
5619R	Tablet 125 mg (base)	60	..	..	4035.00	Tracleer	AT

EPOPROSTENOL SODIUM

Note

Any queries concerning the arrangements to prescribe epoprostenol sodium may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
(f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

- (a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

- (b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

- (a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

- (b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent. All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new adult patients)

Application for initial PBS-subsidised treatment with epoprostenol sodium of adult patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

WHO Functional Class IV primary pulmonary hypertension.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients under 18 years of age)

Application for initial PBS-subsidised treatment with epoprostenol sodium of patients aged less than 18 years who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

WHO Functional Class IV primary pulmonary hypertension.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a patient acknowledgment, signed by the parent or authorised guardian and the prescriber, indicating that they understand and acknowledge that PBS-subsidised treatment with PAH agents will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats will be authorised under this criterion. Where fewer than 5 repeats are initially requested, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all adult patients)

Application for initial PBS-subsidised treatment with epoprostenol sodium of adult patients with one of the following:

- (a) primary pulmonary hypertension who wish to re-commence PBS-subsidised epoprostenol sodium after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with epoprostenol sodium; OR
- (b) WHO Functional Class IV primary pulmonary hypertension and who have received prior treatment with a PBS-subsidised PAH agent other than epoprostenol sodium; OR
- (c) WHO Functional Class III primary pulmonary hypertension and who have failed to respond to a prior PBS-subsidised PAH agent.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (5) for WHO Functional Class III patients, where this is the first application for epoprostenol sodium, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients under 18 years of age)

Application for initial PBS-subsidised treatment with epoprostenol sodium of patients aged less than 18 years with one of the following:

- (a) primary pulmonary hypertension who wish to re-commence PBS-subsidised epoprostenol sodium after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with epoprostenol sodium; OR
- (b) WHO Functional Class IV primary pulmonary hypertension and who have received prior treatment with a PBS-subsidised PAH agent other than epoprostenol sodium; OR
- (c) WHO Functional Class III primary pulmonary hypertension and who have failed to respond to a prior PBS-subsidised PAH agent.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (5) for WHO Functional Class III patients, where this is the first application for epoprostenol sodium, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with epoprostenol sodium of patients who have received approval for initial PBS-subsidised treatment with epoprostenol sodium, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of epoprostenol sodium treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

5731P	Powder for I.V. infusion 500 micrograms (base) with diluent	1	..	..	41.69	Flolan	GK

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

EPOPROSTENOL SODIUM

Note

Any queries concerning the arrangements to prescribe epoprostenol sodium may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent. All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new adult patients)

Application for initial PBS-subsidised treatment with epoprostenol sodium of adult patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:
WHO Functional Class IV primary pulmonary hypertension.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients under 18 years of age)

Application for initial PBS-subsidised treatment with epoprostenol sodium of patients aged less than 18 years who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:
WHO Functional Class IV primary pulmonary hypertension.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a patient acknowledgment, signed by the parent or authorised guardian and the prescriber, indicating that they understand and acknowledge that PBS-subsidised treatment with PAH agents will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats will be authorised under this criterion. Where fewer than 5 repeats are initially requested, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all adult patients)

Application for initial PBS-subsidised treatment with epoprostenol sodium of adult patients with one of the following:

- (a) primary pulmonary hypertension who wish to re-commence PBS-subsidised epoprostenol sodium after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with epoprostenol sodium; OR
- (b) WHO Functional Class IV primary pulmonary hypertension and who have received prior treatment with a PBS-subsidised PAH agent other than epoprostenol sodium; OR
- (c) WHO Functional Class III primary pulmonary hypertension and who have failed to respond to a prior PBS-subsidised PAH agent.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (5) for WHO Functional Class III patients, where this is the first application for epoprostenol sodium, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients under 18 years of age)

Application for initial PBS-subsidised treatment with epoprostenol sodium of patients aged less than 18 years with one of the following:

- (a) primary pulmonary hypertension who wish to re-commence PBS-subsidised epoprostenol sodium after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with epoprostenol sodium; OR
- (b) WHO Functional Class IV primary pulmonary hypertension and who have received prior treatment with a PBS-subsidised PAH agent other than epoprostenol sodium; OR
- (c) WHO Functional Class III primary pulmonary hypertension and who have failed to respond to a prior PBS-subsidised PAH agent.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (5) for WHO Functional Class III patients, where this is the first application for epoprostenol sodium, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with epoprostenol sodium of patients who have received approval for initial PBS-subsidised treatment with epoprostenol sodium, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of epoprostenol sodium treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

5732Q	Powder for I.V. infusion 1.5 mg (base) with diluent	1	..	..	83.37	Flolan	GK

ILOPROST TROMETAMOL

Note

Any queries concerning the arrangements to prescribe iloprost trometamol may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
 (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent. All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with iloprost trometamol of patients who have not received prior PBS-subsidised treatment with iloprost and who have been assessed by a physician from a designated hospital to have:

WHO Functional Class III drug-induced pulmonary arterial hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with iloprost trometamol of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III drug-induced pulmonary arterial hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class IV primary pulmonary hypertension; OR
- (c) WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- (d) WHO Functional Class IV drug-induced pulmonary arterial hypertension.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients)

Application for initial PBS-subsidised treatment with iloprost trometamol of patients with one of the following:

- (a) primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who wish to re-commence PBS-subsidised iloprost trometamol after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with iloprost trometamol; OR
- (b) WHO Functional Class IV primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease and who have received prior treatment with a PBS-subsidised PAH agent other than iloprost trometamol; OR
- (c) WHO Functional Class III primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease and who have failed to respond to a prior PBS-subsidised PAH agent.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
 (5) for WHO Functional Class III patients, where this is the first application for iloprost trometamol, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with iloprost trometamol of patients who have received approval for initial PBS-subsidised treatment with iloprost trometamol, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of iloprost trometamol treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note

Special Pricing Arrangements apply.

5751Q	Solution for inhalation 20 micrograms (base) in 2 mL	30	..	..	1076.00	Ventavis	SC

SILDENAFIL CITRATE

Note

Any queries concerning the arrangements to prescribe sildenafil citrate may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
 Prior Written Approval of Specialised Drugs
 Reply Paid 9826
 GPO Box 9826
 HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND

- primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
- drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND

(c) epoprostenol sodium, of:

- primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
- primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND

(d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND

(e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND

(f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent.

All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with sildenafil citrate of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with sildenafil citrate of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

- (a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR
- (b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6MWT; and
- (3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients)

Application for initial PBS-subsidised treatment with sildenafil citrate of patients with one of the following:

- (a) WHO Functional Class III primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who wish to re-commence PBS-subsidised sildenafil citrate after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with sildenafil citrate; OR
- (b) WHO Functional Class III primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with a PAH agent other than sildenafil citrate.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and
- (3) the date of the first application for PBS-subsidised treatment with a PAH agent; and
- (4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with sildenafil citrate of patients who have received approval for initial PBS-subsidised treatment with sildenafil citrate, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of sildenafil citrate treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:
 - (i) RHC composite assessment; and

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
(ii) ECHO composite assessment; and (iii) 6MWT. The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application. The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).						
9547L	Tablet 20 mg (base)	90	..	..	898.43	Revatio PF

SITAXENTAN SODIUM

Caution

Sitaxentan sodium is a category X drug and must not be given to pregnant women. Pregnancy must be excluded before the start of treatment and avoided during treatment with this drug.

Note

Any queries concerning the arrangements to prescribe sitaxentan sodium may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe PAH agents should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of agents for primary pulmonary hypertension and pulmonary arterial hypertension. Where the term PAH agents appears in the following notes and restrictions it refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, sitaxentan sodium and ambrisentan.

Patients are eligible for PBS-subsidised treatment with only 1 of the above PAH agents at any 1 time. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with scleroderma or connective tissue disease, where the total lung capacity is less than 70% of that predicted.

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of adults with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class IV severity; AND
 - drug-induced pulmonary arterial hypertension, in patients with disease of WHO Functional Class III and IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease, in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, adult patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

The following provides some explanatory notes regarding the availability of PBS-subsidised treatment of patients under the age of 18 years with:

- (a) bosentan monohydrate, of primary pulmonary hypertension or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), in patients with disease of WHO Functional Class III or IV severity; AND
- (b) iloprost trometamol, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (c) epoprostenol sodium, of:
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class III severity and who have failed to respond to prior PBS-subsidised treatment with an alternate PAH agent; AND
 - primary pulmonary hypertension, in patients with disease of WHO Functional Class IV severity; AND
- (d) sildenafil citrate, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (e) sitaxentan sodium, of primary pulmonary hypertension in patients with disease of WHO Functional Class III severity; AND
- (f) ambrisentan, of primary pulmonary hypertension in patients with disease of WHO Functional Class III or IV severity.

From 1 December 2009, patients under the age of 18 years can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 6 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. (New baselines may be submitted where the patient has failed to respond to their current treatment.)

1. Definition of primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology).

Primary pulmonary hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease, including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary capillary wedge pressure (PCWP) less than 18 mmHg; or
- (ii) mPAP greater than 30 mmHg with exercise and PCWP less than 18 mmHg; or
- (iii) where a right heart catheter cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

2. Definition of WHO Functional Class III or IV disease severity.

(a) WHO Functional Class III disease severity is defined as follows:

Patients with pulmonary hypertension resulting in marked limitation of physical activity who are comfortable at rest and on ordinary physical activity experience dyspnoea or fatigue, chest pain or near syncope.

(b) WHO Functional Class IV disease severity is defined as follows:

Patients with the inability to carry out any physical activity without symptoms. These patients manifest signs of right heart failure. Dyspnoea and/or fatigue may even be present at rest. Discomfort is increased by any physical activity.

3. Designated hospitals.

Refer to the Medicare Australia website at www.medicareaustralia.gov.au for a list of designated hospitals.

Note

4. Test requirements to establish baseline for initiation of treatment and response to treatment for continuation of treatment.

(a) Initiation of treatment.

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment, plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted to Medicare Australia for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application.

(b) Continuation of treatment.

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test(s) results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

5. Definition of response to a PAH agent or prior vasodilator treatment.

For adult patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For adult patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least 1 of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

6. Authority approval requirements.

(a) Initiation of PBS-subsidised treatment with a PAH agent, where the patient has not received prior PBS-subsidised treatment with that agent. All applications for initial treatment must be made in writing, must include a completed authority prescription and must be submitted to Medicare Australia for authorisation. The total duration of initial PBS-subsidised treatment that will be approved with this first written application is up to 6 months, based on the dosage recommendations in the TGA-approved Product Information.

Bosentan only:

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA)-approved Product Information. No repeats will be authorised for this prescription. The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Where the 62.5 mg tablet strength is required, please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats. The approved second authority prescription will be returned to the prescriber by Medicare Australia 2 weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the 6 month initial treatment course. Medicare Australia will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Patients who received non-PBS-subsidised treatment with ambrisentan prior to 1 December 2009:

For patients with primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who were commenced on treatment with ambrisentan prior to 1 December 2009 and who have received less than 6 months treatment with ambrisentan at the time of application, the first application for PBS-subsidised treatment must include, where available, all 3 test results at the time that the patient commenced treatment with ambrisentan, bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate or sitaxentan sodium, whichever was initiated first.

(b) Continuation of treatment.

Written applications for continuing treatment for patients who have demonstrated an adequate response to their current treatment must be submitted to Medicare Australia for authorisation every 6 months. Approvals will be limited to provide sufficient supply for up to a maximum of 6 months of treatment, based on the dosage recommendations in the TGA-approved Product Information.

The assessment of the patient's response to the first and subsequent 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated. Applications for continuing treatment with a PAH agent should be made prior to the completion of the 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

(c) Swapping between PAH agents.

For eligible patients, applications to swap between these 6 drugs must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

It is important that patients are assessed for response to every course of treatment approved within the timeframes specified in the relevant

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

restriction, in order to maximise the choice of treatment.

To avoid confusion, applications for patients who wish to swap to an alternate treatment should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

(d) Cessation of treatment — bosentan patients only.

Patients who fail to demonstrate a response to PBS-subsidised bosentan monohydrate treatment at the time where an assessment is required must cease PBS-subsidised bosentan monohydrate therapy.

For patients ceasing treatment, approval will only be granted to provide sufficient supply of the 62.5 mg tablet strength to allow gradual dose reduction over a period of no more than 1 month duration. Prescribers should telephone Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) to receive authorisation for this final supply and to ensure no unintended break in treatment occurs.

7. Re-treatment with a PAH agent.

Patients who do not respond to treatment are not eligible to receive further PBS-subsidised treatment with that agent under any circumstances.

8. Further information.

A tabulated representation of the above information and the restriction can be obtained from the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with sitaxentan sodium of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

(a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR

(b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure of 8 mmHg or less, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Patients must have failed to respond [see Note for definition of response] to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6MWT; and

(3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details on the nature of the adverse event or contraindication according to the TGA-approved Product Information must also be provided with the application.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (new patients)

Application for initial PBS-subsidised treatment with sitaxentan sodium of patients who have not received prior PBS-subsidised treatment with a PAH agent and who have been assessed by a physician from a designated hospital to have:

(a) WHO Functional Class III primary pulmonary hypertension and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO; OR

(b) WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC, or, where a RHC cannot be performed on clinical grounds, right ventricular function as assessed by ECHO.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:

(i) RHC composite assessment; and

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(ii) ECHO composite assessment; and

(iii) 6MWT; and

(3) a signed patient and prescriber acknowledgment indicating that the patient understands and acknowledges that PBS-subsidised treatment with a PAH agent will cease if the treating physician determines that the patient has not achieved a response to treatment [see Note for definition of response].

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial (change or re-commencement for all patients)

Application for initial PBS-subsidised treatment with sitaxentan sodium of patients with one of the following:

(a) WHO Functional Class III primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease who wish to re-commence PBS-subsidised sitaxentan sodium after a break in therapy and who have demonstrated a response to their most recent course of PBS-subsidised treatment with sitaxentan sodium; OR

(b) WHO Functional Class III primary pulmonary hypertension or pulmonary arterial hypertension secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with a PAH agent other than sitaxentan sodium.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes the results on which approval for the first application for PBS-subsidised PAH agent was granted; and

(3) the date of the first application for PBS-subsidised treatment with a PAH agent; and

(4) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests (see requirement 2 above) are able to be performed on clinical grounds, a patient specific reason outlining why the particular test/s could not be conducted must be provided with the authority application [see Note for test requirements].

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Continuing treatment (all patients)

Continuing PBS-subsidised treatment with sitaxentan sodium of patients who have received approval for initial PBS-subsidised treatment with sitaxentan sodium, and who have been assessed by a physician from a designated hospital to have achieved a response to their most recent course of sitaxentan sodium treatment [see Note for definition of response].

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form [www.medicareaustralia.gov.au] which includes results from the 3 tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6MWT.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e. every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information. A maximum of 5 repeats may be requested. Where fewer than 5 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 6 months of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

9551Q	Tablet 100 mg	30	..	..	2743.80	Thelin	PF

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Systemic hormonal preparations, excl. sex hormones and insulins

Pituitary and hypothalamic hormones and analogues

Hypothalamic hormones

Antigrowth hormone

LANREOTIDE ACETATE

Authority required (STREAMLINED)

3387

Active acromegaly in a patient with persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre AND

(a) after failure of other therapy including dopamine agonists; or

(b) as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; or

(c) if the patient is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated.

In a patient treated with radiotherapy, treatment must cease if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (6 weeks after the last dose). Lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Treatment must cease if IGF1 is not lower after 3 months treatment.

5776B	Powder for suspension for injection 30 mg (base) with diluent ampoule	2	11	..	*1500.00	Somatuline LA	IS

LANREOTIDE ACETATE

Authority required (STREAMLINED)

3388

Active acromegaly in a patient with persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre AND

(a) after failure of other therapy including dopamine agonists; or

(b) as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; or

(c) if the patient is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated.

In a patient treated with radiotherapy, treatment must cease if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose). Lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Treatment must cease if IGF1 is not lower after 3 months treatment;

3389

Functional carcinoid tumour causing intractable symptoms. The patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, and surgery or antineoplastic therapy must have failed or be inappropriate.

Treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 120 mg every 28 days. Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

5777C	Injection 60 mg (base) in single dose pre-filled syringe	2	11	..	*2690.00	Somatuline Autogel	IS
5778D	Injection 90 mg (base) in single dose pre-filled syringe	2	11	..	*3580.00	Somatuline Autogel	IS
5779E	Injection 120 mg (base) in single dose pre-filled syringe	2	11	..	*4480.00	Somatuline Autogel	IS

OCTREOTIDE

Authority required (STREAMLINED)

3407

Active acromegaly in a patient with persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre AND

(a) after failure of other therapy including dopamine agonists; or

(b) as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; or

(c) if the patient is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated.

In a patient treated with radiotherapy, treatment must cease if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks. Octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Treatment must cease if IGF1 is not lower after 3 months treatment at a dose of 100 micrograms 3 times daily;

3408

Functional carcinoid tumour or vasoactive intestinal peptide secreting tumour (VIPoma) causing intractable symptoms. The patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, and surgery or antineoplastic therapy must have failed or be inappropriate.

Treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months'

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

					Dispensed Price for			
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Max. Qty \$	Brand Name and Manufacturer		
	therapy. Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.							
9508K	Injection 50 micrograms (as acetate) in 1 mL	90	11	..	*631.62	^a	Hospira Pty Limited	HH
						^a	Octreotide MaxRx	XF
						^a	Sandostatin 0.05	NV
9509L	Injection 100 micrograms (as acetate) in 1 mL	90	11	..	*1263.06	^a	Hospira Pty Limited	HH
						^a	Octreotide MaxRx	XF
						^a	Sandostatin 0.1	NV
9510M	Injection 500 micrograms (as acetate) in 1 mL	90	11	..	*6322.32	^a	Hospira Pty Limited	HH
						^a	Octreotide MaxRx	XF
						^a	Sandostatin 0.5	NV

OCTREOTIDE

Authority required (STREAMLINED)

3409

Acromegaly in a patient controlled on Sandostatin subcutaneous injections.

In a patient treated with radiotherapy, treatment must cease if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose). Octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Treatment must cease if IGF1 is not lower after 3 months of treatment;

3410

Functional carcinoid tumour or vasoactive intestinal peptide secreting tumour (VIPoma) with symptom control on Sandostatin subcutaneous injections.

Treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with Sandostatin subcutaneous injections. Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

9511N	Injection (modified release) 10 mg (as acetate), vial and diluent syringe	1	11	..	1334.96	Sandostatin LAR	NV
9512P	Injection (modified release) 20 mg (as acetate), vial and diluent syringe	1	11	..	1776.74	Sandostatin LAR	NV
9513Q	Injection (modified release) 30 mg (as acetate), vial and diluent syringe	1	11	..	2223.33	Sandostatin LAR	NV

Calcium homeostasis

Anti-parathyroid agents

Other anti-parathyroid agents

CINACALCET HYDROCHLORIDE

Authority required (STREAMLINED)

3323

Management, including initiation and stabilisation, by a nephrologist, of a patient with chronic kidney disease on dialysis who has sustained secondary hyperparathyroidism with iPTH of at least 50 pmol per L, not responding to conventional therapy.

Note

During the titration phase, intact PTH should be monitored 4 weekly (measured at least 12 hours post dose) and dose titrated until an appropriate iPTH concentration is achieved. During the titration phase, approval will be limited to sufficient supply for 4 weeks treatment at a time, with doses between 30 and 180 mg per day according to the patient's response and tolerability.

During the maintenance phase, approval will be limited to provide sufficient quantity for 4 weeks treatment up to a maximum of 6 months supply for doses between 30 and 180 mg per day according to the patient's response and tolerability. Intact PTH should be monitored quarterly (measured at least 12 hours post dose) and dose adjusted as necessary to maintain an appropriate iPTH concentration.

"Sustained" means the abnormality was detected on at least 2 blood samples collected over a period of 2 to 4 months.

Authority required (STREAMLINED)

3324

Management, including initiation and stabilisation, by a nephrologist, of a patient with chronic kidney disease on dialysis who has sustained secondary hyperparathyroidism with iPTH of at least 15 pmol per L and less than 50 pmol per L AND an (adjusted) serum calcium concentration at least 2.6 mmol per L, not responding to conventional treatment.

Note

During the titration phase, intact PTH should be monitored 4 weekly (measured at least 12 hours post dose) and dose titrated until an appropriate iPTH concentration is achieved. During the titration phase, approval will be limited to sufficient supply for 4 weeks treatment at a time, with doses

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

between 30 and 180 mg per day according to the patient's response and tolerability.

During the maintenance phase, approval will be limited to provide sufficient quantity for 4 weeks treatment up to a maximum of 6 months supply for doses between 30 and 180 mg per day according to the patient's response and tolerability. Intact PTH should be monitored quarterly (measured at least 12 hours post dose) and dose adjusted as necessary to maintain an appropriate iPTH concentration.

"Sustained" means the abnormality was detected on at least 2 blood samples collected over a period of 2 to 4 months.

Note

Special Pricing Arrangements apply.

5621W	Tablet 30 mg (base)	56	5	..	*593.72	Sensipar	AN
5622X	Tablet 60 mg (base)	56	5	..	*1187.44	Sensipar	AN
5623Y	Tablet 90 mg (base)	56	5	..	*1781.16	Sensipar	AN

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Antiinfectives for systemic use

Antibacterials for systemic use

Macrolides, lincosamides and streptogramins

Macrolides

AZITHROMYCIN

Authority required (STREAMLINED)

3317

Prophylaxis against Mycobacterium avium complex infections in HIV-positive patients with CD4 cell counts of less than 75 per cubic millimetre.

5616N	Tablet 600 mg	16	5	..	*116.30	Zithromax	PF

CLARITHROMYCIN

Authority required (STREAMLINED)

3325

Treatment of Mycobacterium avium complex infections.

5624B	Tablet 500 mg	100	2	..	72.24	Klacid	AB

5625C	Tablet 250 mg	100	2	..	36.12	Klacid	AB

Antimycobacterials

Drugs for treatment of tuberculosis

Antibiotics

RIFABUTIN

Authority required (STREAMLINED)

3415

Treatment of Mycobacterium avium complex infections in HIV-positive patients;

3317

Prophylaxis against Mycobacterium avium complex infections in HIV-positive patients with CD4 cell counts of less than 75 per cubic millimetre.

9541E	Capsule 150 mg	120	5	..	*588.00	Mycobutin	PF

Antivirals for systemic use

Direct acting antivirals

Nucleosides and nucleotides excl. reverse transcriptase inhibitors

CIDOFOVIR

Authority required (STREAMLINED)

3322

Treatment of cytomegalovirus retinitis in patients with AIDS.

5620T	Solution for I.V. infusion 375 mg (anhydrous) in 5 mL single use vial	4	3	..	*3600.00	Vistide	GI

GANCICLOVIR

Authority required (STREAMLINED)

3379

Cytomegalovirus retinitis in severely immunocompromised patients;

3380

Prophylaxis of cytomegalovirus disease in bone marrow transplant patients at risk of cytomegalovirus disease;

3381

Prophylaxis of cytomegalovirus disease in solid organ transplant patients at risk of cytomegalovirus disease.

5749N	Powder for I.V. infusion 500 mg (as sodium)	10	1	..	*560.00	Cymevene	RO

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

VALACICLOVIR HYDROCHLORIDE

Authority required (STREAMLINED)

3419

Prophylaxis of cytomegalovirus (CMV) infection and disease following renal transplantation in patients at risk of CMV disease.

9568N	Tablet 500 mg (base)	500	2	..	*2115.90	Valtrex	GK

VALGANCICLOVIR HYDROCHLORIDE

Authority required (STREAMLINED)

3420

Cytomegalovirus retinitis in patients with acquired immunodeficiency syndrome;

3421

Prophylaxis of cytomegalovirus infection and disease in solid organ transplant patients at risk of cytomegalovirus disease.

9569P	Tablet 450 mg (base)	120	5	..	*4491.60	Valcyte	RO
9655E	Powder for oral solution 50 mg (base) per mL, 100 mL	11	5	..	*4574.79	Valcyte	RO

Phosphonic acid derivatives

FOSCARNET SODIUM

Authority required (STREAMLINED)

3322

Treatment of cytomegalovirus retinitis in patients with AIDS;

3378

Treatment of aciclovir-resistant herpes simplex virus infection in immunocompromised patients with HIV infection.

5747L	I.V. infusion 24 mg per mL, 250 mL	6	1	..	395.00	Foscavir	AP

Protease inhibitors

ATAZANAVIR

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

5612J	Capsule 300 mg (as sulfate)	60	5	..	*1043.82	Reyataz	BQ
5613K	Capsule 150 mg (as sulfate)	120	5	..	*1043.82	Reyataz	BQ
5614L	Capsule 200 mg (as sulfate)	120	5	..	*1391.76	Reyataz	BQ
5615M	Capsule 100 mg (as sulfate)	120	5	..	*695.88	Reyataz	BQ

DARUNAVIR

Authority required (STREAMLINED)

3595

Treatment of HIV infection, in addition to optimised background therapy in combination with other antiretroviral agents, and co-administered with 100 mg ritonavir twice daily in an antiretroviral experienced patient who, after at least one antiretroviral regimen, has experienced virological failure or clinical failure or genotypic resistance.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

5652L	Tablet 300 mg (as ethanolate)	240	5	..	*2097.42	Prezista	JC
5653M	Tablet 150 mg (as ethanolate)	240	5	..	1048.71	Prezista	JC

FOSAMPRENAVIR

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty \$		
5745J	Oral liquid 50 mg (as calcium) per mL, 225 mL	8	5	..	*812.48	Telzir	VI
5746K	Tablet 700 mg (as calcium)	120	5	..	*758.32	Telzir	GK

INDINAVIR

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

5752R	Capsule 400 mg (as sulfate)	360	5	..	*910.00	Crixivan 400 mg	MK

RITONAVIR

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

9542F	Oral solution 600 mg per 7.5 mL (80 mg per mL), 90 mL	10	5	..	*910.00	Norvir	AB
9660K	Tablet 100 mg	720	5	..	*910.08	Norvir	AB

SAQUINAVIR

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

9545J	Tablet 500 mg (as mesylate)	240	5	..	*1011.12	Invirase	RO

TIPRANAVIR

Authority required (STREAMLINED)

3601

Treatment of HIV infection, in addition to optimised background therapy in combination with other antiretroviral agents, and co-administered with 200 mg ritonavir twice daily in an antiretroviral experienced patient who, after each of at least three different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes, has experienced virological failure or clinical failure or genotypic resistance. Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

Note

Special Pricing Arrangements apply.

9567M	Capsule 250 mg	240	5	..	*2142.00	Aptivus	BY

TIPRANAVIR

Authority required (STREAMLINED)

3603

Treatment of HIV infection, in addition to optimised background therapy in combination with other antiretroviral agents, and co-administered with ritonavir in an antiretroviral experienced patient who, after each of at least three different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes, has experienced virological failure or clinical failure or genotypic resistance.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

Note

Special Pricing Arrangements apply.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty \$		
9656F	Oral liquid 100 mg per mL, 95 mL	7	5	..	*2374.05		Aptivus BY

Nucleoside and nucleotide reverse transcriptase inhibitors

ABACAVIR

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

5601T	Tablet 300 mg (as sulfate)	120	5	..	*564.00	Ziagen	VI
5602W	Oral solution 20 mg (as sulfate) per mL, 240 mL	8	5	..	*657.12	Ziagen	VI

ADEFOVIR DIPIVOXIL

Authority required (STREAMLINED)

3313

Chronic hepatitis B in a patient who has failed antihepadnaviral therapy and who satisfies all of the following criteria:

- (1)(a) Repeatedly elevated serum ALT levels while on concurrent antihepadnaviral therapy of greater than or equal to 6 months duration in conjunction with documented chronic hepatitis B infection; or
 - (b) Repeatedly elevated HBV DNA levels one log greater than the nadir value or failure to achieve a 1 log reduction in HBV DNA within 3 months, whilst on previous antihepadnaviral therapy except in patients with evidence of poor compliance;
 - (2) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.
- Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

Note

Patients should have undergone a liver biopsy at some point since initial diagnosis to obtain histological evidence of chronic hepatitis. Patients may receive treatment in combination with lamivudine but not with other PBS-subsidised antihepadnaviral therapy.

5606C	Tablet 10 mg	60	5	..	*1250.00	Hepsera	GI

DIDANOSINE

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

5663C	Capsule 125 mg (containing enteric coated beadlets)	60	5	..	*280.86	Videx EC	BQ
5664D	Capsule 200 mg (containing enteric coated beadlets)	60	5	..	*326.80	Videx EC	BQ
5665E	Capsule 250 mg (containing enteric coated beadlets)	60	5	..	*408.48	Videx EC	BQ
5666F	Capsule 400 mg (containing enteric coated beadlets)	60	5	..	*653.58	Videx EC	BQ

EMTRICITABINE

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

5709L	Capsule 200 mg	60	5	..	*564.00	Emtriva	GI

ENTECAVIR MONOHYDRATE

Authority required (STREAMLINED)

3352

Patients with chronic hepatitis B who satisfy all of the following criteria:

- (1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

biopsy);

(2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or

(b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;

(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

Note

PBS-subsidised entecavir monohydrate must be used as monotherapy.

5711N	Tablet 0.5 mg	60	5	..	*768.60	Baraclude	BQ

ENTECAVIR MONOHYDRATE

Authority required (STREAMLINED)

3353

Patients with chronic hepatitis B who have failed lamivudine therapy and who satisfy all of the following criteria:

(1)(a) Repeatedly elevated serum ALT levels while on concurrent antihepadnaviral therapy of greater than or equal to 6 months duration in conjunction with documented chronic hepatitis B infection; or

(b) Repeatedly elevated HBV DNA levels one log greater than the nadir value or failure to achieve a 1 log reduction in HBV DNA within 3 months, whilst on previous antihepadnaviral therapy except in patients with evidence of poor compliance;

(2) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

Note

Patients should have undergone a liver biopsy at some point since initial diagnosis to obtain histological evidence of chronic hepatitis.

PBS-subsidised entecavir monohydrate must be used as monotherapy.

5712P	Tablet 1 mg	60	5	..	*1250.00	Baraclude	BQ

LAMIVUDINE

Authority required (STREAMLINED)

3386

Patients with chronic hepatitis B who satisfy all of the following criteria:

(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);

(2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or

(b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;

(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

5770Q	Tablet 100 mg	56	5	..	*298.72	Zeffix	GK

5771R	Oral solution 5 mg per mL, 240 mL	5	5	..	*349.55	Zeffix	GK

LAMIVUDINE

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

5772T	Tablet 150 mg	120	5	..	*564.00	3TC	VI

5773W	Oral solution 10 mg per mL, 240 mL	8	5	..	*691.84	3TC	VI

5774X	Tablet 300 mg	60	5	..	*564.00	3TC	VI

STAVUDINE

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
3589						
Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.						
9552R	Powder for oral solution 1 mg per mL, 200 mL	24	5	..	*2800.32	Zerit BQ
9553T	Capsule 20 mg	120	5	..	*560.00	Zerit BQ
9554W	Capsule 30 mg	120	5	..	*667.36	Zerit BQ
9556Y	Capsule 40 mg	120	5	..	*889.80	Zerit BQ

TELBIVUDINE

Authority required (STREAMLINED)

3416

Treatment, as sole PBS-subsidised therapy, in a patient with chronic hepatitis B who is nucleoside analogue naive and satisfies all of the following criteria:

(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);

(2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or

(b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;

(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

9562G	Tablet 600 mg	56	5	..	*501.76	Sebivo	NV

TENOFOVIR

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

Authority required (STREAMLINED)

3417

Treatment, as sole PBS-subsidised therapy, of chronic hepatitis B in a patient who is nucleoside analogue naive and satisfies all of the following criteria:

(1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);

(2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or

(b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;

(3) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy;

3313

Chronic hepatitis B in a patient who has failed antihepadnaviral therapy and who satisfies all of the following criteria:

(1)(a) Repeatedly elevated serum ALT levels while on concurrent antihepadnaviral therapy of greater than or equal to 6 months duration in conjunction with documented chronic hepatitis B infection; or

(b) Repeatedly elevated HBV DNA levels one log greater than the nadir value or failure to achieve a 1 log reduction in HBV DNA within 3 months, whilst on previous antihepadnaviral therapy except in patients with evidence of poor compliance;

(2) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

Note

Patients should have undergone a liver biopsy at some point since initial diagnosis to obtain histological evidence of chronic hepatitis.

Patients may receive tenofovir treatment in combination with lamivudine but not with other PBS-subsidised antihepadnaviral therapy.

9563H	Tablet containing tenofovir disoproxil fumarate 300 mg	60	5	..	*966.20	Viread	GI

ZIDOVUDINE

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
3589							
Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.							
9570Q	Syrup 10 mg per mL, 200 mL	15	5	..	*673.20	Retrovir	GK
9651Y	Capsule 100 mg	400	5	..	*821.84	Retrovir	GK
9652B	Capsule 250 mg	240	5	..	*1232.76	Retrovir	GK

Non-nucleoside reverse transcriptase inhibitors

EFAVIRENZ

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

Note

Special Pricing Arrangements apply.

5706H	Tablet 600 mg	60	5	..	*905.28	Stocrin	MK
5707J	Oral solution 30 mg per mL, 180 mL	7	5	..	*950.53	Stocrin	MK
5708K	Tablet 200 mg	180	5	..	*905.28	Stocrin	MK

ETRAVIRINE

Authority required (STREAMLINED)

3597

Treatment of HIV infection, in addition to optimised background therapy in combination with other antiretroviral agents in an antiretroviral experienced patient who, after each of at least three different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes, has experienced virological failure or clinical failure or genotypic resistance.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

5736X	Tablet 100 mg	240	5	..	*1233.00	Intelence	JC

NEVIRAPINE

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

9506H	Tablet 200 mg	120	5	..	*543.16	Viramune	BY
9507J	Oral suspension 50 mg (as hemihydrate) per 5 mL, 240 mL	10	5	..	*1350.00	Viramune	BY

Antivirals for treatment of HIV infections, combinations

ABACAVIR SULFATE with LAMIVUDINE and ZIDOVUDINE

Authority required (STREAMLINED)

3592

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient over 12 years of age, weighing 40 kg or more, with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3593

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient over 12 years of age, weighing 40 kg or more, has previously received PBS-subsidised therapy for HIV infection.

5604Y	Tablet containing abacavir 300 mg (as sulfate) with lamivudine 150 mg and zidovudine 300 mg	120	5	..	*1704.00	Trizivir	VI

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price	Brand Name and Manufacturer	
					for Max. Qty \$		
ABACAVIR with LAMIVUDINE							
<u>Authority required (STREAMLINED)</u>							
3592							
Initial treatment of HIV infection in combination with other antiretroviral agents in a patient over 12 years of age, weighing 40 kg or more, with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;							
3593							
Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient over 12 years of age, weighing 40 kg or more, has previously received PBS-subsidised therapy for HIV infection.							
5603X	Tablet containing abacavir 600 mg (as sulfate) with lamivudine 300 mg	60	5	..	*1128.00	Kivexa	VI
LAMIVUDINE with ZIDOVUDINE							
<u>Authority required (STREAMLINED)</u>							
3588							
Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;							
3589							
Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.							
5775Y	Tablet 150 mg-300 mg	120	5	..	*1157.20	Combivir	VI
LOPINAVIR with RITONAVIR							
<u>Authority required (STREAMLINED)</u>							
3588							
Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;							
3589							
Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.							
5789Q	Oral liquid 400 mg-100 mg per 5 mL, 60 mL	10	5	..	*1290.00	Kaletra	AB
5790R	Tablet 100 mg-25 mg	120	5	..	*342.50	Kaletra	AB
5791T	Tablet 200 mg-50 mg	240	5	..	*1370.00	Kaletra	AB
TENOFOVIR with EMTRICITABINE							
<u>Authority required (STREAMLINED)</u>							
3588							
Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;							
3589							
Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.							
9564J	Tablet containing tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg	60	5	..	*1530.20	Truvada	GI
TENOFOVIR with EMTRICITABINE and EFAVIRENZ							
<u>Authority required (STREAMLINED)</u>							
3588							
Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;							
3589							
Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.							
9565K	Tablet containing tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg and efavirenz 600 mg	60	5	..	*2435.48	Atripla	GI

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Other antiretrovirals

ENFUVIRTIDE

Authority required (STREAMLINED)

3597

Treatment of HIV infection, in addition to optimised background therapy in combination with other antiretroviral agents in an antiretroviral experienced patient who, after each of at least three different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes, has experienced virological failure or clinical failure or genotypic resistance.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

5710M	Pack containing 60 vials powder for injection 90 mg with 60 vials water for injections 1.1 mL (with syringes and swabs)	2	5	..	*4426.00	Fuzeon	RO

MARAVIROC

Authority required (STREAMLINED)

3599

Treatment, in addition to optimised background therapy in combination with other antiretroviral agents, of an antiretroviral experienced patient infected with only CCR5-tropic HIV-1, who, after each of at least three different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes, has experienced virological failure or clinical failure or genotypic resistance. A tropism assay to determine CCR5 only strain status is required prior to initiation. Individuals with CXCR4 tropism demonstrated at any time point are not eligible.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

5792W	Tablet 150 mg	120	5	..	*1835.40	Celsentri	PF
5793X	Tablet 300 mg	120	5	..	*1835.40	Celsentri	PF

RALTEGRAVIR

Authority required (STREAMLINED)

3588

Initial treatment of HIV infection in combination with other antiretroviral agents in a patient with a CD4 count of less than 500 per cubic millimetre or symptomatic HIV disease;

3589

Continuing treatment of HIV infection in combination with other antiretroviral agents where the patient has previously received PBS-subsidised therapy for HIV infection.

9523F	Tablet 400 mg (as potassium)	120	5	..	*1331.10	Isentress	MK

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Antineoplastic and immunomodulating agents

Antineoplastic agents

Cytotoxic antibiotics and related substances

Anthracyclines and related substances

DOXORUBICIN HYDROCHLORIDE, PEGYLATED LIPOSOMAL

Authority required (STREAMLINED)

3348

Treatment of AIDS-related Kaposi's sarcoma in patients with CD4 cell counts of less than 200 per cubic millimetre and extensive mucocutaneous involvement;

3349

Treatment of AIDS-related Kaposi's sarcoma in patients with CD4 cell counts of less than 200 per cubic millimetre and extensive visceral involvement.

5705G	Suspension for I.V. infusion 20 mg in 10 mL	4	5	..	*2491.96	Caelyx	SH

Immunostimulants

Immunostimulants

Colony stimulating factors

FILGRASTIM

Authority required (STREAMLINED)

3357

For use in a patient undergoing induction and consolidation therapy for acute myeloid leukaemia;

3358

Mobilisation of peripheral blood progenitor cells to facilitate harvest of such cells for autologous transplantation into a patient with a non-myeloid malignancy who has had myeloablative or myelosuppressive therapy;

3359

Mobilisation of peripheral blood progenitor cells, in a normal volunteer, for use in allogeneic transplantation;

3360

A patient receiving marrow-ablative chemotherapy and subsequent bone marrow transplantation;

3361

A patient with a non-myeloid malignancy receiving marrow-ablative chemotherapy and subsequent autologous peripheral blood progenitor cell transplantation;

3362

A patient with breast cancer receiving standard dose adjuvant chemotherapy who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

3363

A patient receiving chemotherapy for B-cell chronic lymphocytic leukaemia with fludarabine and cyclophosphamide who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

3364

A patient receiving first-line chemotherapy for Hodgkin disease who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

3365

A patient receiving chemotherapy for myeloma who has had a prior episode of febrile neutropenia, and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

3366

A patient with severe congenital neutropenia (absolute neutrophil count of less than 100 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, and in whom a bone marrow examination has shown evidence of maturational arrest of the neutrophil lineage);

3367

A patient with severe chronic neutropenia (absolute neutrophil count of less than 1,000 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, or evidence of neutrophil dysfunction, and, either having experienced a life-threatening infectious episode requiring

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer	
					Max. Qty \$		
hospitalisation and treatment with intravenous antibiotics in the previous 12 months, or having recurrent clinically significant infections (a minimum of 3 in the previous 12 months));							
3368							
A patient with chronic cyclic neutropenia (absolute neutrophil count of less than 500 million cells per litre lasting for 3 days per cycle, measured over 3 separate cycles, and, either having experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics, or having recurrent clinically significant infections (a minimum of 3 in the previous 12 months));							
3369							
A patient with inoperable Stage III, IVa or IVb squamous cell carcinoma of the oral cavity, larynx, oropharynx or hypopharynx receiving neoadjuvant treatment with docetaxel in combination with cisplatin and fluorouracil who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned.							
<u>Authority required (STREAMLINED)</u>							
3370							
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in acute lymphoblastic leukaemia;							
3371							
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in breast cancer (adjuvant chemotherapy with docetaxel in combination with an anthracycline and cyclophosphamide);							
3372							
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in germ cell tumours;							
3373							
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in infants and children with CNS tumours;							
3374							
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in neuroblastoma;							
3375							
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in non-Hodgkin lymphoma (aggressive grades; or low grade receiving an anthracycline-containing regimen);							
3376							
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in relapsed Hodgkin disease;							
3377							
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in sarcoma.							
5741E	Injection 300 micrograms in 1 mL	20	11	..	*3008.00	Neupogen	AN
5742F	Injection 300 micrograms in 0.5 mL single use pre-filled syringe	20	11	..	*3008.00	Neupogen	AN
5743G	Injection 480 micrograms in 1.6 mL	20	11	..	*4814.00	Neupogen	AN
5744H	Injection 480 micrograms in 0.5 mL single use pre-filled syringe	20	11	..	*4814.00	Neupogen	AN

LENOGRASTIM

Authority required (STREAMLINED)

3392

Mobilisation of peripheral blood progenitor cells to facilitate harvest of such cells for reinfusion into patients with non-myeloid malignancies who have had myeloablative or myelosuppressive therapy;

3393

Mobilisation of peripheral blood progenitor cells, in normal volunteers, for use in allogeneic transplantation to facilitate harvest of such cells in healthy donors;

3394

Patients with non-myeloid malignancies receiving marrow-ablative chemotherapy and subsequent peripheral blood progenitor cell or bone marrow transplantation;

3395

Patients with breast cancer receiving standard dose adjuvant chemotherapy who have had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

3396

Patients receiving first-line chemotherapy for Hodgkin's disease who have had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
<u>Authority required (STREAMLINED)</u>							
3397							
Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in acute lymphoblastic leukaemia;							
3398							
Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in Ewing's sarcoma;							
3399							
Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in germ cell tumours;							
3400							
Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in infants and children with CNS tumours;							
3401							
Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in neuroblastoma;							
3402							
Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in non-Hodgkin's lymphoma (intermediate or high grade);							
3403							
Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in osteosarcoma;							
3404							
Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in relapsed Hodgkin's disease;							
3405							
Patients being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in rhabdomyosarcoma.							
5787N	Powder for injection 13,400,000 i.u. (105 micrograms)	20	11	..	*1025.00	Granocyte 13	HH
5788P	Powder for injection 33,600,000 i.u. (263 micrograms)	20	11	..	*2567.20	Granocyte 34	HH

PEGFILGRASTIM

Authority required (STREAMLINED)

3357

For use in a patient undergoing induction and consolidation therapy for acute myeloid leukaemia;

3362

A patient with breast cancer receiving standard dose adjuvant chemotherapy who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

3363

A patient receiving chemotherapy for B-cell chronic lymphocytic leukaemia with fludarabine and cyclophosphamide who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

3364

A patient receiving first-line chemotherapy for Hodgkin disease who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

3365

A patient receiving chemotherapy for myeloma who has had a prior episode of febrile neutropenia, and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned;

3369

A patient with inoperable Stage III, IVa or IVb squamous cell carcinoma of the oral cavity, larynx, oropharynx or hypopharynx receiving neoadjuvant treatment with docetaxel in combination with cisplatin and fluorouracil who has had a prior episode of febrile neutropenia or prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), and for whom there is clinical justification for wishing to continue therapy with the same drug combination, dosage and treatment schedule, and for whom a good response to treatment is anticipated providing chemotherapy can be delivered as planned.

Authority required (STREAMLINED)

3370

A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in acute lymphoblastic leukaemia;

3371

A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in breast cancer (adjuvant chemotherapy with docetaxel in combination with an anthracycline and cyclophosphamide);

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
3372						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in germ cell tumours;						
3373						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in infants and children with CNS tumours;						
3374						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in neuroblastoma;						
3375						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in non-Hodgkin lymphoma (aggressive grades; or low grade receiving an anthracycline-containing regimen);						
3376						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in relapsed Hodgkin disease;						
3377						
A patient being treated with aggressive chemotherapy with the intention of achieving a cure or substantial remission in sarcoma.						
9514R	Injection 6 mg in 0.6 mL single use pre-filled syringe	1	11	..	1925.00	Neulasta AN

Interferons

INTERFERON ALFA-2a

Caution

Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required (STREAMLINED)

3382

Use in the treatment of Philadelphia chromosome positive myelogenous leukaemia in the chronic phase;

3383

Patients with chronic hepatitis B who satisfy all of the following criteria:

- (1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);
- (2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or
- (b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;
- (3) Are not persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L);
- (4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

5759D	Injection 3,000,000 i.u. in 0.5 mL single dose pre-filled syringe	30	5	..	*894.00	Roferon-A	RO
5760E	Injection 4,500,000 i.u. in 0.5 mL single dose pre-filled syringe	30	5	..	*1341.00	Roferon-A	RO
5761F	Injection 6,000,000 i.u. in 0.5 mL single dose pre-filled syringe	30	5	..	*1787.40	Roferon-A	RO
5762G	Injection 9,000,000 i.u. in 0.5 mL single dose pre-filled syringe	30	5	..	*2681.40	Roferon-A	RO

INTERFERON ALFA-2b

Caution

Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required (STREAMLINED)

3384

Adjunctive therapy of malignant melanoma following surgery in patients with nodal involvement;

3382

Use in the treatment of Philadelphia chromosome positive myelogenous leukaemia in the chronic phase;

3383

Patients with chronic hepatitis B who satisfy all of the following criteria:

- (1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);
- (2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or
- (b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;
- (3) Are not persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L);
- (4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

5763H	Solution for injection 18,000,000 i.u. in 1.2 mL multi-dose injection pen	2	5	..	*357.48	Intron A Redipen	SH
5764J	Solution for injection 30,000,000 i.u. in 1.2 mL multi-dose injection pen	2	5	..	*595.80	Intron A Redipen	SH

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty \$		
5765K	Solution for injection 60,000,000 i.u. in 1.2 mL multi-dose injection pen	2	5	..	*1191.60	Intron A Redipen	SH
5766L	Solution for injection 18,000,000 i.u. in 3 mL single dose vial	15	5	..	*2681.10	Intron A	SH
5767M	Solution for injection 25,000,000 i.u. in 2.5 mL single dose vial	15	5	..	*3723.75	Intron A	SH
5768N	Solution for injection 10,000,000 i.u. in 1 mL single dose vial	15	5	..	*1489.50	Intron A	SH

INTERFERON GAMMA-1b

Authority required (STREAMLINED)

3385

Treatment of chronic granulomatous disease in patients with frequent and severe infections despite adequate prophylaxis with antimicrobial agents.

5769P	Injection 2,000,000 i.u. in 0.5 mL	12	11	..	*2721.80	Imukin	BY

PEGINTERFERON ALFA-2a

Caution

Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required (STREAMLINED)

3411

Monotherapy in patients with chronic hepatitis B and compensated liver disease who satisfy all of the following criteria:

- (1) Histological evidence of chronic hepatitis on liver biopsy (except in patients with coagulation disorders considered severe enough to prevent liver biopsy);
- (2)(a) Abnormal serum ALT levels in conjunction with documented chronic hepatitis B infection; or
- (b) Elevated HBV DNA levels in conjunction with documented chronic hepatitis B infection;
- (3) Have received no prior peginterferon alfa therapy for the treatment of hepatitis B;
- (4) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception;
- (5) Are not persons with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L).

Treatment is limited to 1 course of treatment for a duration of up to 48 weeks;

3412

Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa or peginterferon alfa treatment for hepatitis C and have a contraindication to ribavirin, who satisfy all of the following criteria:

- (1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);
- (2) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

The treatment course is limited to up to 48 weeks.

Patients may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop.

Note

Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:

- (a) a nurse educator/counsellor for patients; and
- (b) 24 hour access by patients to medical advice; and
- (c) an established liver clinic; and
- (d) facilities for safe liver biopsy.

9515T	Injection 135 micrograms in 0.5 mL single use pre-filled syringe	8	5	..	*2331.80	Pegasys	RO
9516W	Injection 180 micrograms in 0.5 mL single use pre-filled syringe	8	5	..	*2700.46	Pegasys	RO

PEGINTERFERON ALFA-2b

Caution

Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required (STREAMLINED)

3412

Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa or peginterferon alfa treatment for hepatitis C and have a contraindication to ribavirin, who satisfy all of the following criteria:

- (1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);
- (2) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

The treatment course is limited to up to 48 weeks.

Patients may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer
					Max. Qty \$	
Note						
Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:						
(a) a nurse educator/counsellor for patients; and						
(b) 24 hour access by patients to medical advice; and						
(c) an established liver clinic; and						
(d) facilities for safe liver biopsy.						
9517X	Powder for injection 50 micrograms with diluent in single use injection pen	8	5	..	*1840.00	PEG-Intron Redipen SH
9518Y	Powder for injection 80 micrograms with diluent in single use injection pen	8	5	..	*2944.00	PEG-Intron Redipen SH
9520C	Powder for injection 100 micrograms with diluent in single use injection pen	8	5	..	*3680.00	PEG-Intron Redipen SH
9521D	Powder for injection 120 micrograms with diluent in single use injection pen	8	5	..	*4416.00	PEG-Intron Redipen SH
9522E	Powder for injection 150 micrograms with diluent in single use injection pen	8	5	..	*5520.00	PEG-Intron Redipen SH

RIBAVIRIN and PEGINTERFERON ALFA-2a

Caution

Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Caution

Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.

Authority required (STREAMLINED)

3413

Patients naive to interferon based therapies (non-pegylated or pegylated)

Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa or peginterferon alfa treatment for hepatitis C and who satisfy all of the following criteria:

- (1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);
- (2) Female patients of child-bearing age are not pregnant, not breast-feeding, and both patient and their partner are using effective forms of contraception (one for each partner). Male patients and their partners are using effective forms of contraception (one for each partner). Female partners of male patients are not pregnant.

For patients with genotype 2 or 3 hepatitis C without hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 24 weeks. For hepatitis C patients with genotype 1, 4, 5 or 6 and those genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 48 weeks.

Patients with genotype 1, 4, 5 or 6 who are eligible for 48 weeks of treatment may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop. (An HCV RNA assay at week 12 is unnecessary for genotype 2 and 3 patients because of the high likelihood of early viral response by week 12).

Patients with genotype 1, 4, 5 or 6 who are viral positive at week 12 but have attained at least a 2 log drop in viral load may only continue treatment after the first 24 weeks of treatment if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. Similarly, genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis may only continue treatment after the first 24 weeks if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. An HCV RNA qualitative assay at week 24 is unnecessary for those patients with genotype 1, 4, 5 or 6 who became viral negative at week 12.

Note

Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:

- (a) a nurse educator/counsellor for patients; and
- (b) 24 hour access by patients to medical advice; and
- (c) an established liver clinic; and
- (d) facilities for safe liver biopsy.

Authority required (STREAMLINED)

3414

Patients who have failed one prior attempt at interferon based therapies (non-pegylated or pegylated)

Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no more than one prior treatment with interferon alfa or peginterferon alfa for hepatitis C and who satisfy all of the following criteria:

- (1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);
- (2) Female patients of child-bearing age are not pregnant, not breast-feeding, and both patient and their partner are using effective forms of contraception (one for each partner). Male patients and their partners are using effective forms of contraception (one for each partner). Female partners of male patients are not pregnant.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer	
					Max. Qty \$		
The treatment course is limited to 48 weeks. Patients may only continue treatment after the first 12 weeks of treatment if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 12.							
Note							
Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:							
(a) a nurse educator/counsellor for patients; and							
(b) 24 hour access by patients to medical advice; and							
(c) an established liver clinic; and							
(d) facilities for safe liver biopsy.							
9524G	Pack containing 168 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 135 micrograms	2	5	..	*3072.84	Pegasys RBV	RO
9525H	Pack containing 112 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 180 micrograms	2	5	..	*3085.28	Pegasys RBV	RO
9526J	Pack containing 140 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 180 micrograms	2	5	..	*3245.82	Pegasys RBV	RO
9527K	Pack containing 168 tablets ribavirin 200 mg and 4 pre-filled syringes peginterferon alfa-2a injection 180 micrograms	2	5	..	*3406.36	Pegasys RBV	RO

RIBAVIRIN and PEGINTERFERON ALFA-2b

Caution

Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Caution

Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.

Authority required (STREAMLINED)

3413

Patients naive to interferon based therapies (non-pegylated or pegylated)

Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no prior interferon alfa or peginterferon alfa treatment for hepatitis C and who satisfy all of the following criteria:

- (1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);
- (2) Female patients of child-bearing age are not pregnant, not breast-feeding, and both patient and their partner are using effective forms of contraception (one for each partner). Male patients and their partners are using effective forms of contraception (one for each partner). Female partners of male patients are not pregnant.

For patients with genotype 2 or 3 hepatitis C without hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 24 weeks. For hepatitis C patients with genotype 1, 4, 5 or 6 and those genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis, the treatment course is limited to 48 weeks.

Patients with genotype 1, 4, 5 or 6 who are eligible for 48 weeks of treatment may only continue treatment after the first 12 weeks if the result of an HCV RNA quantitative assay (performed at the same laboratory using the same test) shows that the plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop. (An HCV RNA assay at week 12 is unnecessary for genotype 2 and 3 patients because of the high likelihood of early viral response by week 12).

Patients with genotype 1, 4, 5 or 6 who are viral positive at week 12 but have attained at least a 2 log drop in viral load may only continue treatment after the first 24 weeks of treatment if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. Similarly, genotype 2 or 3 patients with hepatic cirrhosis or bridging fibrosis may only continue treatment after the first 24 weeks if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 24. An HCV RNA qualitative assay at week 24 is unnecessary for those patients with genotype 1, 4, 5 or 6 who became viral negative at week 12.

Note

Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:

- a nurse educator/counsellor for patients; and
- 24 hour access by patients to medical advice; and
- an established liver clinic; and
- facilities for safe liver biopsy.

Authority required (STREAMLINED)

3414

Patients who have failed one prior attempt at interferon based therapies (non-pegylated or pegylated)

Treatment, managed by an accredited treatment centre, of chronic hepatitis C in patients 18 years or older who have compensated liver disease and who have received no more than one prior treatment with interferon alfa or peginterferon alfa for hepatitis C and who satisfy all of the following criteria:

- (1) Documented chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive);
- (2) Female patients of child-bearing age are not pregnant, not breast-feeding, and both patient and their partner are using effective forms of contraception (one for each partner). Male patients and their partners are using effective forms of contraception (one for each partner). Female partners of male patients are not pregnant.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price	Brand Name and Manufacturer	
					for Max. Qty \$		
The treatment course is limited to 48 weeks. Patients may only continue treatment after the first 12 weeks of treatment if plasma HCV RNA is not detectable by an HCV RNA qualitative assay at week 12.							
Note							
Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:							
(a) a nurse educator/counsellor for patients; and							
(b) 24 hour access by patients to medical advice; and							
(c) an established liver clinic; and							
(d) facilities for safe liver biopsy.							
9528L	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 50 micrograms with diluent	2	5	..	*1834.76	Pegatron	SH
9529M	Pack containing 112 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 50 micrograms with diluent	2	5	..	*2119.74	Pegatron	SH
9530N	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 80 micrograms with diluent	2	5	..	*2422.72	Pegatron	SH
9531P	Pack containing 140 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 80 micrograms with diluent	2	5	..	*2707.66	Pegatron	SH
9532Q	Pack containing 168 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 80 micrograms with diluent	2	5	..	*2707.66	Pegatron	SH
9533R	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 100 micrograms with diluent	2	5	..	*2814.66	Pegatron	SH
9534T	Pack containing 112 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 100 micrograms with diluent	2	5	..	*3099.62	Pegatron	SH
9535W	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 120 micrograms with diluent	2	5	..	*3206.62	Pegatron	SH
9536X	Pack containing 140 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 120 micrograms with diluent	2	5	..	*3491.58	Pegatron	SH
9537Y	Pack containing 84 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	2	5	..	*3794.56	Pegatron	SH
9538B	Pack containing 140 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	2	5	..	*4079.52	Pegatron	SH
9539C	Pack containing 168 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	2	5	..	*4079.52	Pegatron	SH
9540D	Pack containing 196 capsules ribavirin 200 mg and 4 single use injection pens containing peginterferon alfa-2b powder for injection 150 micrograms with diluent	2	5	..	*4364.48	Pegatron	SH

Immunosuppressants

Immunosuppressants

Selective immunosuppressants

ABATACEPT

Note

Any queries concerning the arrangements to prescribe abatacept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Further prescribing information (including Authority Application Forms) is on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe abatacept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty \$		

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (new patient or patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with abatacept, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion. Up to a maximum of 4 repeats may be authorised.

Where fewer than 4 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with abatacept.

Patients who fail to demonstrate a response to treatment with abatacept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with abatacept, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have a documented history of severe active rheumatoid arthritis; and
- (b) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with abatacept and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised abatacept treatment, within the timeframes specified below.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion. Up to a maximum of 4 repeats may be authorised.

Where fewer than 4 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised abatacept treatment was approved under either of the initial 1 or 2 treatment restrictions, patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised abatacept treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Patients who fail to demonstrate a response to treatment with abatacept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with abatacept, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (a) who have a documented history of severe active rheumatoid arthritis; and
 (b) who have demonstrated an adequate response to treatment with abatacept; and
 (c) whose most recent course of PBS-subsidised bDMARD treatment was with abatacept.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion. Up to a maximum of 5 repeats may be authorised.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with abatacept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with abatacept, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Patients who fail to demonstrate a response to treatment with abatacept under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

5605B	Powder for I.V. infusion 250 mg	1	..	..	504.43	Orencia	BQ

EVEROLIMUS

Caution

Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

3355

Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required;

3356

Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of cardiac allograft rejection. Management includes initiation, stabilisation and review of therapy as required.

5737Y	Tablet 1 mg	240	5	..	*3844.80	Certican	NV
5738B	Tablet 0.25 mg	120	5	..	*480.60	Certican	NV
5739C	Tablet 0.5 mg	120	5	..	*961.20	Certican	NV
5740D	Tablet 0.75 mg	240	5	..	*2883.60	Certican	NV

MYCOPHENOLATE MOFETIL

Caution

Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

3355

Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required;

3356

Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of cardiac allograft rejection. Management includes initiation, stabilisation and review of therapy as required.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for	Brand Name and Manufacturer	
					Max. Qty \$		
9500B	Powder for oral suspension 1 g per 5 mL, 165 mL	2	5	..	*489.02	CellCept	RO
9501C	Capsule 250 mg	600	5	..	*1111.38	CellCept	RO
9502D	Tablet 500 mg	300	5	..	*1111.38	CellCept	RO

MYCOPHENOLATE SODIUM

Caution

Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

3355

Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required.

9503E	Tablet (enteric coated) 180 mg (mycophenolic acid)	240	5	..	*444.56	Myfortic	NV
9504F	Tablet (enteric coated) 360 mg (mycophenolic acid)	240	5	..	*889.12	Myfortic	NV

NATALIZUMAB

Caution

Progressive multifocal leukoencephalopathy has been reported with this drug.

Note

Neurologists prescribing natalizumab under the PBS listing must be registered with the Tysabri Australian Prescribing Program.

Authority required (STREAMLINED)

3425

Treatment, as monotherapy, by a neurologist, of clinically definite relapsing-remitting multiple sclerosis in an ambulatory (without assistance or support) patient 18 years of age or older, who has experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years.

The diagnosis must be confirmed by magnetic resonance imaging of the brain and/or spinal cord and the date of the scan included in the patient's medical notes, unless written certification provided by a radiologist that an MRI scan is contraindicated because of the risk of physical (not psychological) injury to the patient is included in the patient's medical notes.

Natalizumab must be ceased if there is continuing progression of disability while on treatment with natalizumab. For continued treatment the patient must demonstrate compliance with, and an ability to tolerate, natalizumab.

Note

Special Pricing Arrangements apply.

9505G	Solution concentrate for I.V. infusion 300 mg in 15 mL	1	5	..	2038.46	Tysabri	BD

SIROLIMUS

Caution

Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

3355

Management of rejection, under the supervision and direction of a transplant unit, in patients receiving this drug for prophylaxis of renal allograft rejection. Management includes initiation, stabilisation and review of therapy as required.

9548M	Tablet 2 mg	200	5	..	*2893.34	Rapamune	WX
9549N	Tablet 1 mg	200	5	..	*1446.66	Rapamune	WX
9550P	Oral solution 1 mg per mL, 60 mL	2	5	..	*936.00	Rapamune	WX

Tumor necrosis factor alpha (TNF-alpha) inhibitors

ADALIMUMAB

Note

Any queries concerning the arrangements to prescribe adalimumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe adalimumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and etanercept for a patient who has severe active juvenile idiopathic arthritis. Where the term bDMARD appears in the following NOTES and restrictions, it refers to adalimumab and etanercept only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 bDMARDs at any 1 time.

From 1 November 2010, a patient receiving PBS-subsidised bDMARD therapy is considered to be in a treatment cycle where they may swap to the alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy, and
- fail to respond, or to sustain a response to one PBS-subsidised bDMARD twice and the other PBS-subsidised bDMARD once only.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological therapy before they are eligible to receive further PBS-subsidised bDMARD therapy. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment was stopped to the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

A patient who was receiving PBS-subsidised bDMARD treatment immediately prior to 1 November 2010 is considered to be in their first cycle as of 1 November 2010. A patient who has had a break in bDMARD treatment of at least 12 months immediately prior to making a new application, on or after 1 November 2010, will commence a new treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 November 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to the alternate bDMARD without having to requalify with respect to the indices of disease severity (joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months.

A patient may trial the alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug twice within the current treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a bDMARD. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to the revised baseline measurement.

(4) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 12 months, must requalify for treatment under the Initial 1 treatment restriction.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for severe active juvenile idiopathic arthritis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must qualify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 12 month break in PBS-subsidised therapy' above for further details.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with bDMARDs should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Medicare Australia at the time treatment is ceased.

Authority required

Initial 1 (new patient or patient recommencing after a break of more than 12 months).

Initial treatment by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of a patient under 18 years:

- (a) who has severe active juvenile idiopathic arthritis; AND
- (b) whose parent or authorised guardian has signed a patient acknowledgement; AND
- (c) who has not received PBS-subsidised treatment with adalimumab or etanercept for this condition in the previous 12 months; AND
- (d) who has demonstrated either:
 - (i) severe intolerance of, or toxicity due to, methotrexate (see below for definition of severe intolerance and toxicity); or
 - (ii) failure to achieve an adequate response to 1 or more of the following treatment regimens:
 - oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or
 - oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other DMARD, alone or in combination with corticosteroids, for a minimum of 3 months. (Note: use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.)

Severe intolerance is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant NSAIDs on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Toxicity is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, please provide details at time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of this toxicity at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; OR
- (b) at least 4 active joints from the following list:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment should be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) an acknowledgement signed by a parent or authorised guardian.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of authority application, medical practitioners should request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 4 weeks from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with adalimumab.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial adalimumab after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months).

Initial PBS-subsidised treatment with adalimumab by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of a patient under 18 years who:

- (a) has a documented history of severe active juvenile idiopathic arthritis; and
- (b) in this treatment cycle, has received prior PBS-subsidised treatment with adalimumab or etanercept for this condition; and
- (c) has not failed PBS-subsidised therapy with adalimumab for this condition more than once in the current treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for a patient who has received PBS-subsidised treatment with adalimumab in this treatment cycle and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised adalimumab treatment, within the timeframes specified below.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of authority application, medical practitioners should request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised adalimumab treatment was approved under either of the Initial 1 or 2 treatment restrictions, the patient must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised adalimumab treatment was approved under the continuing treatment criteria, the patient must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to that particular course of bDMARD.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial adalimumab after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

Authority required

Initial 3 ('grandfather' patients).

Initial PBS-subsidised supply for continuing treatment with adalimumab, by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of a patient under 18 years who:

- (a) has a documented history of severe active juvenile idiopathic arthritis; and
- (b) was receiving treatment with adalimumab prior to 1 March 2010; and
- (c) has demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with adalimumab; and
- (d) is receiving treatment with adalimumab at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) an acknowledgement signed by a parent or authorised guardian.

A maximum of 24 weeks of treatment will be authorised under this restriction.

At the time of authority application, medical practitioners should request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 5 repeats will be authorised.

Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

The assessment of the patient's response to this initial course of PBS-subsidised therapy must be made within the 4 weeks prior to completion of the course in order to ensure continuity of treatment.

A patient ceasing treatment or swapping to an alternate agent and wishing to demonstrate a response to treatment, must be assessed no earlier than 12 weeks from the commencement of PBS-subsidised treatment. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A patient may only qualify for PBS-subsidised treatment under this restriction once.

Authority required

Continuing treatment.

Continuing PBS-subsidised treatment with adalimumab, by a rheumatologist or under the supervision of a paediatric rheumatology treatment centre, of a patient:

- (a) who has a documented history of severe active juvenile idiopathic arthritis; and
- (b) who has demonstrated an adequate response to treatment with adalimumab; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment in this treatment cycle was with adalimumab.

An adequate response to treatment is defined as:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	
(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]. A maximum of 24 weeks of treatment will be approved under this restriction. At the time of authority application, medical practitioners should request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 5 repeats will be authorised. Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). All applications for continuing treatment with adalimumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with adalimumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course. If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial adalimumab after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.							
9661L	Injection 20 mg in 0.4 mL pre-filled syringe	2	..	..	1630.00	Humira	AB
9662M	Injection 40 mg in 0.8 mL pre-filled syringe	2	..	..	1630.00	Humira	AB
9663N	Injection 40 mg in 0.8 mL pre-filled pen	2	..	..	1630.00	Humira	AB

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and etanercept for a patient who has severe active juvenile idiopathic arthritis. Where the term bDMARD appears in the following NOTES and restrictions, it refers to adalimumab and etanercept only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 bDMARDs at any 1 time.

From 1 November 2010, a patient receiving PBS-subsidised bDMARD therapy is considered to be in a treatment cycle where they may swap to the alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy, and
- fail to respond, or to sustain a response to one PBS-subsidised bDMARD twice and the other PBS-subsidised bDMARD once only.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological therapy before they are eligible to receive further PBS-subsidised bDMARD therapy. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment was stopped to the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

A patient who was receiving PBS-subsidised bDMARD treatment immediately prior to 1 November 2010 is considered to be in their first cycle as of 1 November 2010. A patient who has had a break in bDMARD treatment of at least 12 months immediately prior to making a new application, on or after 1 November 2010, will commence a new treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 November 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to the alternate bDMARD without having to requalify with respect to the indices of disease severity (joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months.

A patient may trial the alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug twice within the current treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a bDMARD. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to the revised baseline measurement.

(4) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 12 months, must requalify for treatment under the Initial 1 treatment restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for severe active juvenile idiopathic arthritis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must qualify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 12 month break in PBS-subsidised therapy' above for further details.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with bDMARDs should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Medicare Australia at the time treatment is ceased.

Authority required

Initial 1 (new patient or patient recommencing after a break of more than 12 months).

Initial treatment by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of a patient under 18 years:

- (a) who has severe active juvenile idiopathic arthritis; AND
- (b) whose parent or authorised guardian has signed a patient acknowledgement; AND
- (c) who has not received PBS-subsidised treatment with adalimumab or etanercept for this condition in the previous 12 months; AND
- (d) who has demonstrated either:
 - (i) severe intolerance of, or toxicity due to, methotrexate (see below for definition of severe intolerance and toxicity); or
 - (ii) failure to achieve an adequate response to 1 or more of the following treatment regimens:
 - oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or
 - oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other DMARD, alone or in combination with corticosteroids, for a minimum of 3 months. (Note: use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.)

Severe intolerance is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant NSAIDs on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, please provide details at time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of this toxicity at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; OR
- (b) at least 4 active joints from the following list:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment should be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) an acknowledgement signed by a parent or authorised guardian.

A maximum of 16 weeks of treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	Max. Price	

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 4 weeks from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial etanercept after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months).

Initial PBS-subsidised treatment with etanercept by a paediatric rheumatologist, or under the supervision of a paediatric rheumatology treatment centre, of a patient under 18 years who:

- (a) has a documented history of severe active juvenile idiopathic arthritis; and
- (b) in this treatment cycle, has received prior PBS-subsidised treatment with adalimumab or etanercept for this condition; and
- (c) has not failed PBS-subsidised therapy with etanercept for this condition more than once in the current treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for a patient who has received PBS-subsidised treatment with etanercept in this treatment cycle and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised etanercept treatment, within the timeframes specified below.

A maximum of 16 weeks of treatment will be authorised under this restriction.

Where fewer than 3 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with etanercept may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised etanercept treatment was approved under either of the Initial 1 or 2 treatment restrictions, the patient must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised etanercept treatment was approved under the continuing treatment criteria, the patient must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to that particular course of bDMARD.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial etanercept after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

Authority required

Continuing treatment.

Continuing PBS-subsidised treatment with etanercept, by a rheumatologist or under the supervision of a paediatric rheumatology treatment centre, of a patient:

- (a) who has a documented history of severe active juvenile idiopathic arthritis; and
- (b) who has demonstrated an adequate response to treatment with etanercept; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment in this treatment cycle was with etanercept.

An adequate response to treatment is defined as:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

A maximum of 24 weeks of treatment will be approved under this restriction.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with etanercept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with etanercept, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle. A patient may re-trial etanercept after a minimum of 12 months have elapsed between the date the last PBS-subsidised bDMARD was stopped and the date of the first application under a new treatment cycle.

5734T	Injection set containing 4 vials powder for injection 25 mg and 4 pre-filled syringes solvent 1 mL	1	..	..	815.00	Enbrel	WX

ETANERCEPT

Note

Any queries concerning the arrangements to prescribe etanercept may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe etanercept should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and etanercept for a patient who has severe active juvenile idiopathic arthritis. Where the term bDMARD appears in the following NOTES and restrictions, it refers to adalimumab and etanercept only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 bDMARDs at any 1 time.

From 1 November 2010, a patient receiving PBS-subsidised bDMARD therapy is considered to be in a treatment cycle where they may swap to the alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy, and
- fail to respond, or to sustain a response to one PBS-subsidised bDMARD twice and the other PBS-subsidised bDMARD once only.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological therapy before they are eligible to receive further PBS-subsidised bDMARD therapy. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment was stopped to the date of the first application for initial treatment with a bDMARD under the new treatment cycle.

A patient who was receiving PBS-subsidised bDMARD treatment immediately prior to 1 November 2010 is considered to be in their first cycle as of 1 November 2010. A patient who has had a break in bDMARD treatment of at least 12 months immediately prior to making a new application, on or after 1 November 2010, will commence a new treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of a bDMARD in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 November 2010.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to the alternate bDMARD without having to requalify with respect to the indices of disease severity (joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months.

A patient may trial the alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug twice within the current treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a bDMARD. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to the revised baseline measurement.

(4) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised bDMARD therapy of at least 12 months, must requalify for treatment under the Initial 1 treatment restriction.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for severe active juvenile idiopathic arthritis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must qualify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 12 month break in PBS-subsidised therapy' above for further details.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with bDMARDs should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Medicare Australia at the time treatment is ceased.

Authority required

Continuing treatment.

Continuing PBS-subsidised treatment with etanercept, by a rheumatologist or under the supervision of a paediatric rheumatology treatment centre, of a patient 18 years or older:

- (a) who has a documented history of severe active juvenile idiopathic arthritis; and
- (b) who has demonstrated an adequate response to treatment with etanercept; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment in this treatment cycle was with etanercept.

An adequate response to treatment is defined as:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with etanercept must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with etanercept, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

If a patient fails to respond to treatment 3 times (twice with one agent and once with the other) they will not be eligible to receive further PBS-subsidised bDMARD therapy in this treatment cycle.

Where a patient with severe active juvenile idiopathic arthritis continues treatment with etanercept and is 18 years or older, etanercept 50 mg may be prescribed.

5733R	Injections 50 mg in 1 mL single use pre-filled syringes, 4	1	..	..	1630.01	Enbrel	WX
5735W	Injection 50 mg in 1 mL single use auto-injector, 4	1	..	..	1630.01	Enbrel	WX

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

;

Note

TREATMENT OF ADULT PATIENTS WITH ACTIVE ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept, golimumab and infliximab for adult patients with active ankylosing spondylitis. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab, etanercept, golimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 4 TNF-alfa antagonists at any 1 time.

From 1 March 2007, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised TNF-alfa antagonists without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 March 2007 is considered to be in their first cycle as of 1 March 2007.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than once. A patient who, prior to 1 March 2007, was authorised to receive PBS-subsidised initial treatment for ankylosing spondylitis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2007.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and 18 weeks of treatment for infliximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap to an alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with at least 1 NSAID, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the BASDAI, ESR and/or CRP levels are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for active ankylosing spondylitis prior to 1 March 2010 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with golimumab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with golimumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial 1 (new patients)

Initial PBS-subsidised treatment with infliximab, by a rheumatologist, of an adult with active ankylosing spondylitis who has radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis and who has not received any PBS-subsidised treatment with either adalimumab, etanercept, golimumab or infliximab in this treatment cycle; AND

(a) who has at least 2 of the following:

- (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or
- (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI) [for further information on the BASMI please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; or
- (iii) limitation of chest expansion relative to normal values for age and gender [for chest expansion normal values please refer to the Medicare Australia website at www.medicareaustralia.gov.au]; AND

(b) who has failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

The application must include details of the NSAIDs trialled, their doses and duration of treatment. If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance. Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Medicare Australia website at www.medicareaustralia.gov.au.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au] which must include the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form [www.medicareaustralia.gov.au]; and
 - (iii) a completed Exercise Program Self Certification Form included in the supporting information form; and
 - (iv) a signed patient acknowledgment form.

The assessment of the patient's response to the initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

A maximum of 18 weeks of treatment with infliximab will be approved under this criterion.

At the time of the authority application, the doctor should request the appropriate number of vials, based on the weight of the patient, to provide for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 18 weeks of treatment may be requested by telephone.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.

Authority required

Initial 2 (change or re-commencement for all patients)

Initial PBS-subsidised treatment with infliximab, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who, in this treatment cycle, has received prior PBS-subsidised TNF-alfa antagonist treatment for this condition and is eligible to receive further TNF-alfa antagonist therapy, and has not failed PBS-subsidised therapy with infliximab in the current treatment cycle.

Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised TNF-alfa antagonist therapy or, under this restriction, for patients who have received previous PBS-subsidised TNF-alfa antagonist therapy) the patient must have been assessed for response to that course following a minimum of 12 weeks of treatment. These assessments must be provided to Medicare Australia no later than 4 weeks from the date the course was ceased. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Where the most recent course of PBS-subsidised infliximab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au].

A maximum of 18 weeks of treatment with infliximab will be approved under this criterion.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

At the time of the authority application, the doctor should request the appropriate number of vials, based on the weight of the patient, to provide for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 18 weeks of treatment may be requested by telephone.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.

Authority required

Continuing treatment for all patients

Continuing PBS-subsidised treatment, by a rheumatologist, of an adult with a documented history of active ankylosing spondylitis who:

- (a) has demonstrated an adequate response to treatment with infliximab; and
- (b) whose most recent course of PBS-subsidised therapy in this treatment cycle was with infliximab.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au].

All measurements provided must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with infliximab will be authorised under this criterion.

At the time of the authority application, the doctor should request the appropriate number of vials, based on the weight of the patient, to provide for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone.

All applications for continuing treatment with infliximab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment following an initial treatment course it must be made following a minimum of 12 weeks of treatment with infliximab. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised TNF-alfa antagonist was approved in this cycle and the date of the first application under a new cycle.

5753T	Powder for I.V. infusion 100 mg	1	..	..	751.70	Remicade	SH

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (new patient or patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with infliximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 22 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 3 mg per kg. Up to a maximum of 3 repeats may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with infliximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have a documented history of severe active rheumatoid arthritis; and
- (b) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with infliximab and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised infliximab treatment, within the timeframes specified below.

A maximum of 22 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 3 mg per kg. Up to a maximum of 3 repeats may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised infliximab treatment was approved under either of the initial 1 or 2 treatment restrictions, patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised infliximab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with infliximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (a) who have a documented history of severe active rheumatoid arthritis; and
 (b) who have demonstrated an adequate response to treatment with infliximab; and
 (c) whose most recent course of PBS-subsidised bDMARD treatment was with infliximab.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 3 mg per kg. Up to a maximum of 2 repeats may be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with infliximab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with infliximab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

5757B	Powder for I.V. infusion 100 mg	1	..	..	751.70	Remicade	SH

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
 Prior Written Approval of Specialised Drugs
 Reply Paid 9826
 GPO Box 9826
 HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents (adalimumab, etanercept, golimumab and infliximab) for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time. Where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, golimumab and infliximab.

From 1 August 2006, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis after 1 August 2010.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients — golimumab only.

Applications for patients who commenced treatment with golimumab prior to 1 March 2010 may apply for initial PBS-subsidised treatment as continuing therapy under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; or
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment Cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Authority required

Initial 1

Initial PBS-subsidised treatment with infliximab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults who:

- (1) have severe active psoriatic arthritis; and
- (2) have received no prior PBS-subsidised biological treatment for this condition in this Treatment Cycle; and
- (3) have failed to achieve an adequate response to:
 - (a) methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months; and
 - (b) sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; or
 - (c) leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities, including severity, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) an active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 22 weeks treatment will be authorised under this restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

The assessment of the patient's response to the initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Initial 2

Initial PBS-subsidised treatment with infliximab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults who:

- (1) have a documented history of severe active psoriatic arthritis; and
- (2) have received prior PBS-subsidised biological treatment for this condition in this Treatment Cycle and are eligible to receive further biological therapy; and
- (3) have not failed treatment with infliximab during the current Treatment Cycle.

Applications for patients who have received PBS-subsidised treatment with infliximab within this Treatment Cycle and who wish to re-commence therapy with this drug within this same Cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised infliximab treatment, within the timeframes specified below.

A maximum of 22 weeks treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised infliximab treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised biological therapy or, under this restriction, for patients who have received previous PBS-subsidised biological therapy), patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised infliximab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with infliximab, by a rheumatologist or clinical immunologist with expertise in the management of psoriatic arthritis, of adults:

- (1) who have a documented history of severe active psoriatic arthritis; and
- (2) whose most recent course of PBS-subsidised biological agent for this condition in the current Treatment Cycle was with infliximab; and
- (3) who, at the time of application, demonstrate an adequate response to treatment with infliximab.

An adequate response to treatment with infliximab is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats may be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with infliximab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with infliximab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

Patients who fail to demonstrate a response to treatment with infliximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug, in this Treatment Cycle. Patients may re-trial infliximab after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents (adalimumab, etanercept, golimumab and infliximab) for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time. Where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, golimumab and infliximab.

From 1 August 2006, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The 5-year break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis after 1 August 2010.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 2).

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for all agents except for infliximab, for which a maximum of 22 weeks will be authorised. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients — golimumab only.

Applications for patients who commenced treatment with golimumab prior to 1 March 2010 may apply for initial PBS-subsidised treatment as continuing therapy under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; or
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a treatment Cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must re-qualify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

5756Y	Powder for I.V. infusion 100 mg	1	..	..	751.70	Remicade	SH

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and infliximab for adult patients with severe refractory Crohn disease. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alfa antagonists at any 1 time.

From 1 August 2008, under the PBS, all patients will be able to commence a treatment cycle where they may trial each PBS-subsidised TNF-alfa antagonist without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 August 2008 is considered to be in their first cycle as of 1 August 2008.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than twice.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2008.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2008, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap if eligible to the alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, evidence of intestinal inflammation), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab or infliximab.

A patient who commenced treatment with adalimumab for severe refractory Crohn disease prior to 9 November 2007 or infliximab prior to 7 March 2007 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab or infliximab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab or infliximab will be assessed under the continuing treatment restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial 1 (new patients)

Initial treatment of Crohn disease in a patient assessed by CDAI.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist or a consultant physician as specified in the NOTE below, of a patient with severe refractory Crohn disease who satisfies the following criteria:

- (a) has confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician as specified in the NOTE below; and
- (b) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (c) has failed to achieve an adequate response to prior systemic therapy including:
 - (i) a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; and
 - (ii) immunosuppressive therapy including:
 - azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
 - 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
 - methotrexate at a dose of at least 15 mg weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) have a severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as assessed.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

The most recent CDAI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the signed patient acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete the 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

A CDAI assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Initial 2

Change or re-commencement of treatment of Crohn disease in a patient assessed by CDAI.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist or a consultant physician as specified in the NOTE below of a patient who:

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (a) has a documented history of severe refractory Crohn disease; and
- (b) in this treatment cycle, has received prior PBS-subsidised treatment with infliximab or adalimumab for this condition; and
- (c) has not failed PBS-subsidised therapy with infliximab for this condition more than once in the current treatment cycle.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of TNF-alfa antagonist therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of TNF-alfa antagonist treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of TNF-alfa antagonist.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition; and
 - (ii) details of prior TNF alfa antagonist treatment including details of date and duration of treatment.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete the 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

A CDAI assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Continuing treatment of Crohn disease in a patient assessed by CDAI.

Continuing PBS-subsidised treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease; and
- (b) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition.

The CDAI assessment must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with infliximab, a CDAI assessment of the patient's response must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial 1

Initial treatment of Crohn disease in a patient with short gut syndrome or an ostomy patient.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist, or consultant physician as specified in the NOTE below of a patient who satisfies the following criteria:

- (a) has confirmed Crohn disease defined by standard clinical, endoscopic and/or imaging features, including histological evidence with the diagnosis confirmed by a gastroenterologist or consultant physician as specified in the NOTE below; and
- (b) has diagnostic imaging or surgical evidence of short gut syndrome or has an ileostomy or colostomy; and
- (c) has evidence of intestinal inflammation; and
- (d) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (e) has failed to achieve an adequate response to prior systemic drug therapy including:
 - (i) a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; and
 - (ii) immunosuppressive therapy including:
 - azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
 - 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
 - methotrexate at a dose of at least 15 mg weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) have evidence of intestinal inflammation, including:
 - (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; AND/OR
 - (ii) faeces: higher than normal lactoferrin or calprotectin level; AND/OR
 - (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery;
- AND/OR
- (b) be assessed clinically as being in a high faecal output state;
- AND/OR
- (c) be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of infliximab.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (ii) reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (iii) date of the most recent clinical assessment; and
- (iv) the signed patient acknowledgement.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete the 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Initial 2

Change or re-commencement of treatment of Crohn disease in a patient with short gut syndrome, an ostomy patient or a patient with extensive small intestine disease.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist or a consultant physician as specified in the NOTE below of a patient who:

- (a) has a documented history of severe refractory Crohn disease; and
- (b) in this treatment cycle, has received prior PBS-subsidised treatment with infliximab or adalimumab for this condition; and
- (c) has not failed PBS-subsidised therapy with infliximab for this condition more than once in the current treatment cycle.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of TNF-alfa antagonist therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised TNF-alfa antagonist treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of TNF-alfa antagonist treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of TNF-alfa antagonist.

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criteria, if relevant; and
 - (ii) details of prior TNF alfa antagonist treatment including details of date and duration of treatment.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Authority required

Continuing treatment of Crohn disease in a patient with short gut syndrome or an ostomy patient.

Continuing PBS-subsidised treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease with intestinal inflammation and with short gut syndrome or with an ileostomy or colostomy; and
- (b) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as:

(a) improvement of intestinal inflammation as demonstrated by:

- (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; AND/OR
- (ii) faeces: normalisation of lactoferrin or calprotectin level; AND/OR
- (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or
- (b) reversal of high faecal output state; or
- (c) avoidance of the need for surgery or total parenteral nutrition (TPN).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
- (i) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy or the date of clinical assessment.

The patient's assessment must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with infliximab, an assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial 1

Initial treatment of Crohn disease in a patient with extensive small intestine disease.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist or a consultant physician as specified in the NOTE below, of a patient with severe refractory Crohn disease who satisfies the following criteria:

- (a) has confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or consultant physician as specified in the NOTE below; and
- (b) has extensive small intestinal disease with radiological evidence of intestinal inflammation affecting more than 50 cm of the small intestine; and
- (c) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) has failed to achieve an adequate response to prior systemic therapy including:
 - (i) a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; and
 - (ii) immunosuppressive therapy including:
 - azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
 - 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
 - methotrexate at a dose of at least 15 mg weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) have severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; AND/OR
- (b) have evidence of active intestinal inflammation, including:
 - (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; AND/OR
 - (ii) faeces: higher than normal lactoferrin or calprotectin level; AND/OR
 - (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery; AND/OR
- (c) be assessed clinically as being in a high faecal output state; AND/OR
- (d) be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of infliximab.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (ii) (1) reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; or
 - (2) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the dates of assessment of the patient's condition, if relevant; and
 - (iii) date of the most recent clinical assessment; and
 - (iv) the signed patient acknowledgement.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete the 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Continuing treatment of Crohn disease in a patient with extensive small intestine disease.

Continuing PBS-subsidised treatment with infliximab by a gastroenterologist, or consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease with extensive intestinal inflammation affecting more than 50 cm of the small intestine; and
- (b) has demonstrated or sustained an adequate response to treatment with infliximab.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as:

- (a) a reduction in Crohn Disease Activity Index (CDAI) Score to no greater than 150; or
- (b) improvement of intestinal inflammation as demonstrated by:
 - (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; AND/OR
 - (ii) faeces: normalisation of lactoferrin or calprotectin level; AND/OR
 - (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or
- (c) reversal of high faecal output state; or
- (d) avoidance of the need for surgery or total parenteral nutrition (TPN).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition; or
 - (ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy; or
 - (iii) the date of clinical assessment.

All assessments must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with infliximab, an assessment of the patient's response must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial 3 (grandfather)

Initial PBS-subsidised treatment of Crohn disease in a patient assessed by CDAI who has previously received non-PBS-subsidised therapy with infliximab.

Initial PBS-subsidised supply for continuing treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below, or other consultant physician in consultation with a gastroenterologist of a patient who:

- (a) has a documented history of severe refractory Crohn disease and was receiving treatment with infliximab prior to 7 March 2007; and
- (b) had a Crohn Disease Activity Index (CDAI) Score of greater than or equal to 300 prior to commencing treatment with infliximab. Where a baseline CDAI assessment is not available, please call Medicare Australia on 1800 700 270 to discuss; and
- (c) has signed a patient acknowledgement indicating that they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) has demonstrated or sustained an adequate response to treatment with infliximab. For advice please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as a reduction in Crohn Disease Activity Index (CDAI) Score to no greater than 150.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and baseline Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition; and
 - (ii) the signed patient acknowledgement.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

The current CDAI assessment must be no more than 1 month old at the time of application. The baseline CDAI assessment must be from immediately prior to commencing treatment with infliximab.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Authority required

Initial 3

Initial PBS-subsidised treatment of Crohn disease in a patient with short gut syndrome, an ostomy patient, or a patient with extensive small intestine disease, who has previously received non-PBS-subsidised therapy with infliximab.

Initial PBS-subsidised supply for continuing treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below, or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of severe refractory Crohn disease and was receiving treatment with infliximab prior to 7 March 2007; and
- (b) (1) has a history of extensive small intestinal disease with radiological evidence of intestinal inflammation affecting more than 50 cm of the small intestine; or
- (2) has diagnostic imaging or surgical evidence of short gut syndrome or has an ileostomy or colostomy with a documented history of intestinal inflammation; and
- (c) has signed a patient acknowledgement indicating that they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) has demonstrated or sustained an adequate response to treatment with infliximab according to the criteria included in the relevant continuation restriction. For advice please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

The same criteria used to determine an inadequate response to prior treatment at baseline must be used to determine response to treatment and eligibility for continuing therapy, according to the criteria included in the continuing treatment restriction.

An adequate response to infliximab treatment is defined as:

- (a) a reduction in Crohn Disease Activity Index (CDAI) Score to no greater than 150; or
- (b) improvement of intestinal inflammation as demonstrated by:
 - (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; AND/OR
 - (ii) faeces: normalisation of lactoferrin or calprotectin level; AND/OR
 - (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or
- (c) reversal of high faecal output state; or
- (d) avoidance of the need for surgery or total parenteral nutrition (TPN).

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) (1) the completed current and baseline Crohn Disease Activity Index (CDAI) Score calculation sheet, where relevant, including the date of the assessment of the patient's condition; or
 - (2) the reports and dates of the current and baseline pathology or diagnostic imaging test(s) in order to assess response to therapy; or
 - (3) the date of clinical assessment(s); and
 - (ii) the signed patient acknowledgement.

The patient's assessment must be no more than 1 month old at the time of application. The baseline CDAI assessments must be from immediately prior to commencing treatment with infliximab. Where a baseline assessment is not available, please call Medicare Australia on 1800 700 270 to discuss.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and infliximab for adult patients with severe refractory Crohn disease. Where the term 'tumour necrosis factor (TNF) alpha antagonist' appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alpha antagonists at any 1 time.

From 1 August 2008, under the PBS, all patients will be able to commence a treatment cycle where they may trial each PBS-subsidised TNF-alpha antagonist without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alpha antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alpha antagonist treatment prior to 1 August 2008 is considered to be in their first cycle as of 1 August 2008.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alpha antagonist more than twice.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alpha antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alpha antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alpha antagonist under the new treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alpha antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alpha antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alpha antagonist therapy after 1 August 2008.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alpha antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alpha antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alpha antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2008, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap if eligible to the alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, evidence of intestinal inflammation), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab or infliximab.

A patient who commenced treatment with adalimumab for severe refractory Crohn disease prior to 9 November 2007 or infliximab prior to 7 March 2007 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab or infliximab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab or infliximab will be assessed under the continuing treatment restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

5754W	Powder for I.V. infusion 100 mg	1	..	..	751.70	Remicade	SH

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab and infliximab for adult patients with severe refractory Crohn disease. Where the term 'tumour necrosis factor (TNF) alfa antagonist' appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alfa antagonists at any 1 time.

From 1 August 2008, under the PBS, all patients will be able to commence a treatment cycle where they may trial each PBS-subsidised TNF-alfa antagonist without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a TNF-alfa antagonist while they continue to show a response to therapy.

A patient who received PBS-subsidised TNF-alfa antagonist treatment prior to 1 August 2008 is considered to be in their first cycle as of 1 August 2008.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alfa antagonist more than twice.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alfa antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alfa antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alfa antagonist under the new treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of TNF-alfa antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised TNF-alfa antagonist therapy after 1 August 2008.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised TNF-alfa antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) TNF-alfa antagonist therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific TNF-alfa antagonist following a break in PBS-subsidised therapy with that agent (Initial 2).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

infliximab.

From 1 August 2008, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific TNF-alfa antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alfa antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alfa antagonist supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap if eligible to the alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, evidence of intestinal inflammation), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate TNF-alfa antagonist at any time, regardless of whether they are receiving therapy (initial or continuing) with a TNF-alfa antagonist at the time of the application. However, they cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to the alternate TNF-alfa antagonist should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab or infliximab.

A patient who commenced treatment with adalimumab for severe refractory Crohn disease prior to 9 November 2007 or infliximab prior to 7 March 2007 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab or infliximab will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab or infliximab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Initial treatment of Crohn disease in a paediatric patient.

Initial PBS-subsidised treatment by a gastroenterologist, paediatrician or consultant physician as specified in the NOTE below, of a patient aged 6 to 17 years inclusive with moderate to severe refractory Crohn disease who satisfies the following criteria:

- (a) has confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or consultant physician as specified in the NOTE below; and
- (b) whose parent or authorised guardian has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (c) has failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including:
 - (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period;
 - (ii) an 8 week course of enteral nutrition;
 - (iii) immunosuppressive therapy including:
 - azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; or
 - 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; or
 - methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) severity of disease activity which results in a Paediatric Crohn Disease Activity Index (PCDAI) Score greater than or equal to 30 as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) The most recent PCDAI assessment must be no more than 1 month old at the time of application.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet including the date of assessment of the patient's condition; and
 - (ii) details of previous systemic drug therapy [dosage, date of commencement and duration of therapy], or dates of enteral nutrition; and
 - (iii) the signed patient acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete the 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

A PCDAI assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Continuing treatment of Crohn disease in a patient initiated on PBS-subsidised treatment as a paediatric patient.

Continuing PBS-subsidised treatment with infliximab by a gastroenterologist, paediatrician, consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of moderate to severe refractory Crohn disease; and
- (b) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as a reduction in Paediatric Crohn Disease Activity Index (PCDAI) Score by at least 15 points as compared to baseline AND a total PCDAI score of 30 points or less.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
- (i) the completed Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet along with the date of the assessment of the patient's condition.

The PCDAI assessment must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with infliximab, a PCDAI assessment of the patient's response must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

Patients who fail to demonstrate or sustain a response to treatment with infliximab for Crohn disease as specified in the criteria for continuing treatment with infliximab, will not be eligible to receive PBS-subsidised treatment with this drug within 12 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Initial PBS-subsidised treatment of Crohn disease in a paediatric patient who has previously received non-PBS-subsidised therapy with infliximab.

Initial PBS-subsidised supply for continuing treatment with infliximab by a gastroenterologist, paediatrician, consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient aged 6 to 17 years inclusive who:

- (a) has a documented history of moderate to severe refractory Crohn disease and was receiving treatment with infliximab prior to 4 July 2007; and
- (b) had a Paediatric Crohn Disease Activity Index (PCDAI) Score of greater than 30 prior to commencing treatment with infliximab. Where a baseline CDAI assessment is not available, please call Medicare Australia on 1800 700 270 to discuss; and
- (c) whose parent or authorised guardian has signed a patient acknowledgement indicating that they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) has demonstrated or sustained an adequate response to treatment with infliximab. For advice please contact Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as a reduction in Paediatric Crohn Disease Activity Index (PCDAI) Score by at least 15 points as compared to baseline AND a total PCDAI score of 30 points or less.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and baseline Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet along with the date of the assessment of the patient's condition; and
 - (ii) the signed patient acknowledgement.

The current PCDAI assessment must be no more than 1 month old at the time of application. The baseline PCDAI assessment must be from immediately prior to commencing treatment with infliximab.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

Patients who fail to demonstrate or sustain a response to treatment with infliximab for Crohn disease as specified in the criteria for continuing treatment with infliximab, will not be eligible to recommence PBS-subsidised treatment with this drug within 12 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

5755X	Powder for I.V. infusion 100 mg	1	..	..	751.70	Remicade	SH

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Authority required

Initial treatment of complex refractory FISTULISING CROHN DISEASE.

Initial PBS-subsidised treatment with infliximab by a gastroenterologist or a consultant physician as specified in the NOTE below, of a patient with complex refractory fistulising Crohn disease who:

- (a) has confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician as specified in the NOTE below; and
- (b) has an externally draining enterocutaneous or rectovaginal fistula; and
- (c) has signed a patient acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criteria for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

Authority applications must be made in writing and must include:

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (a) a completed authority prescription form; and
 (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 (i) a completed current Fistula Assessment Form including the date of assessment of the patient's condition; and
 (ii) a signed patient acknowledgement.

The most recent fistula assessment must be no more than 1 month old at the time of application.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6 will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

An assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (up to 6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

A patient who fails to respond to a course of PBS-subsidised infliximab for the treatment of complex refractory fistulising Crohn disease will not be eligible to receive further PBS-subsidised treatment with infliximab for this condition within 12 months of the date on which treatment was ceased.

Authority required

Re-commencement of treatment of complex refractory FISTULISING CROHN DISEASE.

Re-initiation of PBS-subsidised treatment of complex refractory fistulising Crohn disease by a gastroenterologist or a consultant physician as specified in the NOTE below, of a patient with complex refractory fistulising Crohn disease who:

- (a) has a documented history of complex refractory fistulising Crohn disease; and
 (b) has an externally draining enterocutaneous or rectovaginal fistula; and
 (c) has previously received PBS-subsidised infliximab treatment for a draining enterocutaneous or rectovaginal fistula; and EITHER
 (d) has demonstrated or sustained an adequate response to the most recent course of PBS-subsidised treatment with infliximab for this condition; or
 (e) has failed to demonstrate or sustain an adequate response to PBS-subsidised treatment with infliximab for this condition and 12 months have elapsed from the date on which treatment was ceased.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
 (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes a completed current Fistula Assessment Form including the date of assessment of the patient's condition.

The most recent fistula assessment must be no more than 1 month old at the time of application.

A maximum quantity and number of repeats to provide for an initial course of infliximab consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete 3 doses of infliximab may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

An assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (up to 6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab.

It is recommended that an application for continuing treatment is posted to Medicare Australia at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

A patient who fails to respond to a course of PBS-subsidised infliximab for the treatment of complex refractory fistulising Crohn disease will not be eligible to receive further PBS-subsidised treatment with infliximab for this condition within 12 months of the date on which treatment was ceased.

Authority required

Initial PBS-subsidised treatment of complex refractory FISTULISING CROHN DISEASE in a patient who has previously received non-PBS-subsidised therapy with infliximab.

Initial PBS-subsidised supply for continuing treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below, or other consultant physician in consultation with a gastroenterologist of a patient who satisfies the following criteria:

- (a) has a documented history of complex refractory fistulising Crohn disease and was receiving treatment with infliximab prior to 1 March 2010; and
- (b) had a draining enterocutaneous or rectovaginal fistula(e) prior to commencing treatment with infliximab; and
- (c) has signed a patient acknowledgement indicating that they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criteria for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment; and
- (d) is receiving treatment with infliximab at the time of application; and
- (e) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response to infliximab treatment is defined as:

- (a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or
- (b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) a completed current Fistula Assessment form including the date of assessment of the patient's condition; and
 - (ii) a signed patient acknowledgement.

The current fistula assessment must be no more than 1 month old at the time of application.

The baseline fistula assessment must be from immediately prior to commencing treatment with infliximab.

An assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criteria.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

A patient who fails to respond to a course of PBS-subsidised infliximab for the treatment of complex refractory fistulising Crohn disease will not be eligible to receive further PBS-subsidised treatment with infliximab for this condition within 12 months of the date on which treatment was ceased.

Authority required

Continuing treatment of complex refractory FISTULISING CROHN DISEASE.

Continuing PBS-subsidised treatment with infliximab by a gastroenterologist, a consultant physician as specified in the NOTE below or other consultant physician in consultation with a gastroenterologist, of a patient who:

- (a) has a documented history of complex refractory fistulising Crohn disease; and
- (b) has demonstrated or sustained an adequate response to treatment with infliximab.

NOTE: Prescribers must be gastroenterologists (code 87), consultant physicians [internal medicine specialising in gastroenterology (code 81)] or consultant physicians [general medicine specialising in gastroenterology (code 82)].

An adequate response is defined as:

- (a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or
- (b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Authority applications must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes a completed Fistula Assessment form including the date of the assessment of the patient's condition.

The fistula assessment must be no more than 1 month old at the time of application.

If the application is the first application for continuing treatment with infliximab, an assessment of the patient's response must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated.

An assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criteria.

Where an assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with infliximab.

Patients are eligible to receive continuing infliximab treatment in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised. No applications for increased repeats will be authorised.

Where fewer than 2 repeats are requested at the time of application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

A patient who fails to respond to a course of PBS-subsidised infliximab for the treatment of complex refractory fistulising Crohn disease will not be eligible to receive further PBS-subsidised treatment with infliximab for this condition within 12 months of the date on which treatment was ceased.

9654D	Powder for I.V. infusion 100 mg	1	..	..	751.70	Remicade	SH

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe infliximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, etanercept, infliximab and ustekinumab, for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological agents' appears in the following NOTES and restrictions, it only refers to adalimumab, etanercept, infliximab and ustekinumab.

From 1 March 2010, all patients will be able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial adalimumab, etanercept, infliximab or ustekinumab without having to meet the initial treatment criteria, that is they will not need to experience a disease flare when swapping to an alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

A patient who received PBS-subsidised biological agent treatment for chronic plaque psoriasis prior to 1 March 2010 is considered to be in their first Cycle as of 1 March 2010.

Patients are eligible for PBS-subsidised treatment with only 1 biological agent at any 1 time.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Within the same Treatment Cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for a PBS-subsidised biological agent, they must change to an alternate agent if they wish to continue PBS-subsidised biological treatment. A patient who, prior to 1 March 2010, was authorised to receive PBS-subsidised initial treatment for chronic plaque psoriasis with the same agent twice, is exempt from this condition in respect of applications approved prior to 1 March 2010.

Patients must be assessed for response to each course of continuing treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a Treatment Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological agent therapy before they are eligible to commence the next Cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological agent treatment in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Treatment Cycle.

Patients for whom a break in PBS-subsidised therapy of less than 5 years duration has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe chronic plaque psoriasis after 1 March 2010.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made in the following situations:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); or
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under '(4) Swapping therapy' below]; or
- (iii) patients who wish to re-commence treatment following a break in PBS-subsidised therapy with that agent (Initial 2).

All applications for initial treatment will be limited to provide for a maximum of 16 weeks of treatment in the case of adalimumab and etanercept, 22 weeks of treatment in the case of infliximab and 28 weeks of treatment in the case of ustekinumab.

Grandfather patients (adalimumab and ustekinumab only).

Applications for patients who commenced treatment with adalimumab prior to 1 March 2009 or ustekinumab prior to 26 November 2009, may be made for initial PBS-subsidised treatment as continuing therapy under the initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with a biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment. Approval will be based on the criteria included in the relevant restriction.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological agent, a PASI assessment must be conducted after at least 12 weeks of treatment. This assessment must be submitted to Medicare Australia within 1 month of the completion of this initial treatment course. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Application for continuing treatment.

Following the completion of an initial treatment course of a biological agent to which an adequate response has been demonstrated, patients may qualify to receive up to 24 weeks of continuing treatment with that biological agent. Patients are eligible to continue to receive continuous treatment with 24 week courses providing they continue to sustain a response.

For second and subsequent courses of PBS-subsidised treatment with adalimumab, etanercept, infliximab or ustekinumab it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Where a response assessment is not submitted to Medicare Australia within these timeframes, patients will be deemed to have failed to sustain a response to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

timeframes, please call Medicare Australia on 1800 700 270 to discuss.

(4) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate agent within the same Treatment Cycle without having to requalify with respect to disease severity (i.e. a PASI score of greater than 15), or prior treatment requirements.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

Patients may trial an alternate biological agent at any time, regardless of whether they are receiving therapy with a biological agent at the time of the application or not. However, they cannot swap to a particular agent if they have failed to respond to treatment with that particular agent within the same Cycle.

Patients who commenced treatment with adalimumab prior to 1 June 2009 or ustekinumab prior to 1 March 2010 access these interchangeability arrangements in the same way as patients who have not.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the agent being ceased.

(5) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment authority is submitted within a Treatment Cycle and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatment applications.

(6) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent Biological Treatment Cycle, following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment according to the criteria of the relevant restriction and index of disease severity. Patients must have had at least 1 prior treatment, as listed in the criteria, for a minimum of 6 weeks, and must have a PASI assessment conducted preferably whilst still on treatment, but no later than 1 month following cessation of treatment. The PASI assessment must be no older than 1 month at the time of application.

Authority required

Initial treatment [Initial 1, Whole body (New patients — No prior biological agent)]

Initial treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis; and
- (b) have not received any prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (whole body); and
- (d) have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 3 of the following 4 treatments:
 - (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or
 - (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or
 - (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or
 - (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities including severity, associated with phototherapy, methotrexate, cyclosporin and acitretin, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 22 weeks of treatment with infliximab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 22 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Authority required

Initial or re-Treatment [Initial 2, Whole body (Received prior biological agent under PBS)]

Treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have a documented history of severe chronic plaque psoriasis; and
- (b) have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have not failed PBS-subsidised therapy with infliximab for the treatment of this condition in the current Treatment Cycle.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

Applications for patients who have demonstrated a response to PBS-subsidised infliximab treatment within this Treatment Cycle and who wish to re-commence infliximab treatment within the same Cycle following a break in therapy, will only be approved where evidence of the patient's response to their most recent course of PBS-subsidised infliximab treatment has been submitted to Medicare Australia within 1 month of cessation of treatment.

A maximum of 22 weeks of treatment with infliximab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 22 weeks.

A PASI assessment of the patient's response to this course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Continuing treatment (Whole body)

Continuing PBS-subsidised treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over:

- (a) who have a documented history of severe chronic plaque psoriasis; and
- (b) whose most recent course of PBS-subsidised biological treatment for this condition in this Treatment Cycle was with infliximab; and
- (c) who have demonstrated an adequate response to their most recent course of treatment with infliximab.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-biological treatment baseline value for this Treatment Cycle.

This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with infliximab, the assessment of response must be after a minimum of 12 weeks of treatment with an initial course.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet along with the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with infliximab.

A maximum of 24 weeks of treatment with infliximab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for continuing authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks.

A PASI assessment of the patient's response must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Initial treatment [Initial 1, Face, hand, foot (New patients — No prior biological agent)]

Initial treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- (b) have not received any prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have signed a patient and prescriber acknowledgement indicating they understand and acknowledge that PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment (face, hand, foot); and
- (d) have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 3 of the following 4 treatments:
 - (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or
 - (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or
 - (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or
 - (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, or where

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

phototherapy is contraindicated, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity to necessitate permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of acceptable toxicities including severity, associated with phototherapy, methotrexate, cyclosporin and acitretin, can be found on the Medicare Australia website (www.medicareaustralia.gov.au).

The following initiation criterion indicates failure to achieve an adequate response and must be demonstrated in all patients at the time of the application:

- (a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)];
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the signed patient and prescriber acknowledgements.

A maximum of 22 weeks of treatment with infliximab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 22 weeks.

A PASI assessment of the patient's response to this initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Authority required

Initial or re-Treatment [Initial 2, Face, hand, foot (Received prior biological agent under PBS)]

Treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over who:

- (a) have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot; and
- (b) have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle; and
- (c) have not failed PBS-subsidised therapy with infliximab for the treatment of this condition in the current Treatment Cycle.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)];
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

Applications for patients who have demonstrated a response to PBS-subsidised infliximab treatment within this Treatment Cycle and who wish to re-commence infliximab treatment within the same Cycle following a break in therapy, will only be approved where evidence of the patient's response to their most recent course of PBS-subsidised infliximab treatment has been submitted to Medicare Australia within 1 month of cessation of treatment.

A maximum of 22 weeks of treatment with infliximab will be authorised under this restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period beyond 22 weeks.

A PASI assessment of the patient's response to this course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Authority required

Continuing treatment (Face, hand, foot)

Continuing PBS-subsidised treatment as systemic monotherapy (other than methotrexate) by a dermatologist for adults 18 years and over:

- (a) who have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot; and
- (b) whose most recent course of PBS-subsidised biological treatment for this condition in this Treatment Cycle was with infliximab; and
- (c) who have demonstrated an adequate response to treatment with infliximab.

An adequate response to infliximab treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the pre-biological treatment baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the pre-biological treatment baseline value.

This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with infliximab, the assessment of response must be after a minimum of 12 weeks of treatment with an initial course.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes the following:
 - (i) the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams along with the date of the assessment of the patient's condition [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

The most recent PASI assessment must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with infliximab will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats will be authorised.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Under no circumstances will telephone approvals be granted for continuing authority applications, or for treatment that would otherwise extend the treatment period beyond 24 weeks.

A PASI assessment of the patient's response must be conducted within 4 weeks prior to completion of this course of treatment. This assessment, which will be used to determine eligibility for further continuing treatment must be submitted to Medicare Australia no later than 1 month from the date of completion of this course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with infliximab. In circumstances where it is not possible to submit a response assessment within these timeframes, please call Medicare Australia on 1800 700 270 to discuss.

It is recommended that an application is posted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised infliximab treatment.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Patients who fail to demonstrate a response to treatment with 3 biological agents are deemed to have completed this Treatment Cycle and must cease PBS-subsidised therapy. These patients may re-commence a new Biological Treatment Cycle after a minimum of 5 years has elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note

No applications for increased repeats will be authorised.

5758C	Powder for I.V. infusion 100 mg	1	..	..	751.70	Remicade	SH

Interleukin inhibitors

TOCILIZUMAB

Note

Any queries concerning the arrangements to prescribe tocilizumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Further prescribing information (including Authority Application Forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe tocilizumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or

(iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (new patient or patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with tocilizumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (c) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least 3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested.

Up to a maximum of 3 repeats of each strength may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Assessment of a patient's response to an initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab.

Patients who fail to demonstrate a response to treatment with tocilizumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with tocilizumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have a documented history of severe active rheumatoid arthritis; and
- (b) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with tocilizumab and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised tocilizumab treatment, within the timeframes specified below.

A maximum of 16 weeks of treatment will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats of each strength may be authorised.

Where fewer than 3 repeats are requested at the time of the initial application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Where the most recent course of PBS-subsidised tocilizumab treatment was approved under either of the initial 1 or 2 treatment restrictions, patients must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised tocilizumab treatment was approved under the continuing treatment criteria, patients must have been assessed for response, and the assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Patients who fail to demonstrate a response to treatment with tocilizumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Initial 3 ('grandfather' patients)

Initial PBS-subsidised supply for continuing treatment with tocilizumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of an adult who:

- (a) has a documented history of severe active rheumatoid arthritis; and
- (b) was receiving treatment with tocilizumab prior to 1 July 2009; and
- (c) has demonstrated a response as specified in the criteria for continuing PBS-subsidised treatment with tocilizumab; and
- (d) is receiving treatment with tocilizumab at the time of application.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [www.medicareaustralia.gov.au]; and
- (3) a signed patient acknowledgement.

The same indices of disease severity used to establish baseline at the commencement of treatment with a bDMARD must be used for assessment of all continuing applications.

The assessment of the patient's response to a continuing course of therapy must be made within 4 weeks prior to completion of that course and posted to Medicare Australia no less than 2 weeks prior to the date the next dose is scheduled in order to ensure continuity of treatment for those patients who meet the continuation criterion.

A maximum of 24 weeks of treatment with tocilizumab will be approved under this criterion.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 5 repeats of each strength may be authorised.

Where fewer than 5 repeats are initially requested with the authority prescription, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Patients may qualify for PBS-subsidised treatment under this restriction once only.

Patients who fail to demonstrate a response to treatment with tocilizumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with tocilizumab, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have a documented history of severe active rheumatoid arthritis; and

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

- (b) who have demonstrated an adequate response to treatment with tocilizumab; and
(c) whose most recent course of PBS-subsidised bDMARD treatment was with tocilizumab.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

— elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

— shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of 24 weeks of treatment will be approved under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 5 repeats of each strength may be authorised.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

All applications for continuing treatment with tocilizumab must include a measurement of response to the prior course of therapy. This assessment must be provided to Medicare Australia no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with tocilizumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Patients who fail to demonstrate a response to treatment with tocilizumab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

9657G	Concentrate for injection 80 mg in 4 mL	1	..	..	186.88	Actemra	RO
9658H	Concentrate for injection 200 mg in 10 mL	1	..	..	467.20	Actemra	RO
9659J	Concentrate for injection 400 mg in 20 mL	1	..	..	934.40	Actemra	RO

Calcineurin inhibitors

CYCLOSPORIN

Caution

Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

3328

Management of rejection in patients following organ or tissue transplantation, under the supervision and direction of a transplant unit.

Management includes initiation, stabilisation and review of therapy as required;

3329

Management (which includes initiation, stabilisation and review of therapy) by dermatologists or clinical immunologists of patients with severe atopic dermatitis for whom other systemic therapies are ineffective or inappropriate;

3330

Management (which includes initiation, stabilisation and review of therapy) by dermatologists of patients with severe psoriasis for whom other systemic therapies are ineffective or inappropriate and in whom the disease has caused significant interference with quality of life;

3331

Management (which includes initiation, stabilisation and review of therapy) by nephrologists of patients with nephrotic syndrome in patients in whom steroids and cytostatic drugs have failed or are not tolerated or are considered inappropriate and in whom renal function is unimpaired;

3332

Management (which includes initiation, stabilisation and review of therapy) by rheumatologists or clinical immunologists of patients with severe active rheumatoid arthritis for whom classical slow-acting anti-rheumatic agents (including methotrexate) are ineffective or inappropriate.

5632K	Capsule 10 mg	120	5	..	*74.40	Neoral 10	NV
5633L	Oral liquid 100 mg per mL, 50 mL	4	5	..	*1263.16	Neoral	NV
5634M	Capsule 25 mg	120	5	..	*156.68 ^a	Cicloral	SZ

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer	
					Max. Qty \$			
5635N	Capsule 50 mg	120	5	^B 3.76 ..	*160.44 *326.04	^a ^a	Neoral 25 Cicloral	NV SZ
5636P	Capsule 100 mg	120	5	^B 4.04 ..	*330.08 *664.36	^a ^a	Neoral 50 Cicloral	NV SZ
				^B 4.00	*668.36	^a	Neoral 100	NV

CYCLOSPORIN

Caution

Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

3333

For use by organ or tissue transplant recipients.

5631J	Solution concentrate for I.V. infusion 50 mg in 1 mL	10	..	..	54.10		Sandimmun	NV

TACROLIMUS

Caution

Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

3328

Management of rejection in patients following organ or tissue transplantation, under the supervision and direction of a transplant unit. Management includes initiation, stabilisation and review of therapy as required.

9558C	Capsule 500 micrograms	200	5	..	*327.84	^a ^a	Prograf Tacrolimus Sandoz	JC SZ
9560E	Capsule 1 mg	200	5	..	*655.68	^a ^a	Prograf Tacrolimus Sandoz	JC SZ
9561F	Capsule 5 mg	100	5	..	*1638.38	^a ^a	Prograf Tacrolimus Sandoz	JC SZ
9664P	Capsule 0.5 mg (once daily prolonged release)	60	5	..	*98.36		Prograf XL	JC
9665Q	Capsule 1 mg (once daily prolonged release)	120	5	..	*393.40		Prograf XL	JC
9666R	Capsule 5 mg (once daily prolonged release)	60	5	..	*983.54		Prograf XL	JC

Other immunosuppressants

LENALIDOMIDE

Note

Any queries concerning the arrangements to prescribe lenalidomide may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms) is available on the Medicare Australia website at www.medicareaustralia.gov.au.

Any queries concerning patients who are enrolled on the Lenalidomide Compassionate program may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). These patients must demonstrate they met initial criteria prior to commencing treatment on the compassionate program and also demonstrate they do not have progressive disease. Baseline and current pathology reports must be submitted with the initial application.

Applications for authority to prescribe lenalidomide should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001.

Authority required

Initial PBS-subsidised treatment, as monotherapy or in combination with dexamethasone, of a patient with a histological diagnosis of multiple myeloma who has progressive disease after at least 1 prior therapy and who has undergone or is ineligible for a primary stem cell transplant. The patient must have experienced treatment failure after a trial of at least four (4) weeks of thalidomide at a dose of at least 100 mg daily or have failed to achieve at least a minimal response after eight (8) or more weeks of thalidomide-based therapy for progressive disease.

If the dosing requirement for thalidomide cannot be met, the application must state the reasons why this criterion cannot be satisfied.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein and less than 200 mg per 24 hour Bence-Jones proteinuria.

Thalidomide treatment failure is defined as:

- (1) confirmed disease progression during thalidomide treatment or within 6 months of discontinuing thalidomide treatment; or
- (2) severe intolerance or toxicity unresponsive to clinically appropriate dose adjustment.

Severe intolerance due to thalidomide is defined as unacceptable somnolence or sedation interfering with activities of daily living.

Toxicity from thalidomide is defined as peripheral neuropathy (Grade 2 or greater, interfering with function), drug-related seizures, serious Grade 3 or 4 drug-related dermatological reactions, such as Stevens-Johnson Syndrome, or other Grade 3 or 4 toxicity.

Failure to achieve at least a minimal response after 8 or more weeks of thalidomide-based therapy for progressive disease is defined as:

- (1) less than a 25% reduction in serum or urine M protein; or
- (2) in oligo-secretory and non-secretory myeloma patients only, less than a 25% reduction in the difference between involved and uninvolved serum free light chain levels.

Lenalidomide will only be subsidised for patients with multiple myeloma who are not receiving concomitant PBS-subsidised bortezomib.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Multiple Myeloma Authority Application - Supporting Information Form, which includes details of the histological diagnosis of multiple myeloma, prior treatments including name(s) of drug(s) and date of most recent treatment cycle and record of prior stem cell transplant or ineligibility for prior stem cell transplant; details of thalidomide treatment failure; details of the basis of the diagnosis of progressive disease or failure to respond; and nomination of which disease activity parameters will be used to assess response.

To enable confirmation by Medicare Australia, current diagnostic reports of at least one of the following are required:

- (a) the level of serum monoclonal protein; or
- (b) Bence-Jones proteinuria — the results of 24-hour urinary light chain M protein excretion; or
- (c) the serum level of free kappa and lambda light chains; or
- (d) bone marrow aspirate or trephine; or
- (e) if present, the size and location of lytic bone lesions (not including compression fractures); or
- (f) if present, the size and location of all soft tissue plasmacytomas by clinical or radiographic examination i.e. MRI or CT-scan; or
- (g) if present, the level of hypercalcaemia, corrected for albumin concentration.

As these parameters will be used to determine response, results for either (a) or (b) or (c) should be provided for all patients. Where the patient has oligo-secretory or non-secretory multiple myeloma, either (c) or (d) or if relevant (e), (f) or (g) should be provided. Where the prescriber plans to assess response in patients with oligo-secretory or non-secretory multiple myeloma with free light chain assays, evidence of the oligo-secretory or non-secretory nature of the multiple myeloma (either previous or current serum M protein less than 10 g per L and urinary Bence-Jones protein undetectable or less than 200 mg per 24 hours) must be provided; and

- (3) duration of thalidomide and daily dose prescribed; and
- (4) a signed patient acknowledgment.

Note

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

Authority required

Continuing PBS-subsidised treatment, as monotherapy or in combination with dexamethasone, of multiple myeloma in a patient who has previously been issued with an authority prescription for lenalidomide and who does not have progressive disease.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Authority applications for continuing treatment may be made by telephone to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

Note

Special Pricing Arrangements apply.

5783J	Capsule 5 mg	21	..	..	5392.38	Revlimid	CJ
5784K	Capsule 10 mg	21	..	..	5643.33	Revlimid	CJ
5785L	Capsule 15 mg	21	..	..	6581.61	Revlimid	CJ
5786M	Capsule 25 mg	21	..	..	6934.20	Revlimid	CJ

RITUXIMAB

Note

Any queries concerning the arrangements to prescribe rituximab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Further prescribing information (including Authority Application Forms) is on the Medicare Australia website at www.medicareaustralia.gov.au.

Written applications for authority to prescribe rituximab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001;

Note

TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying anti-rheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab) and the T-cell co-stimulation modulator (abatacept).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

PBS-subsidised abatacept, golimumab, infliximab and rituximab must be used in combination with methotrexate at a dose of at least 7.5 mg weekly. Where a patient cannot tolerate 7.5 mg of methotrexate weekly, they are eligible to receive PBS-subsidised adalimumab, certolizumab pegol, etanercept and tocilizumab.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact Medicare Australia on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 12 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 12 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 12 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 12 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab and tocilizumab, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to Medicare Australia within 4 weeks.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to Medicare Australia no later than 2 weeks prior to the patient completing their current treatment course.

Rituximab patients:

A further application may be submitted to Medicare Australia 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to Medicare Australia no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 12 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

Note

(3) Baseline measurements to determine response.

Medicare Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and Medicare Australia will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with certolizumab pegol, golimumab or tocilizumab.

From 1 August 2010, a patient who commenced treatment with certolizumab pegol or golimumab for severe rheumatoid arthritis prior to 1 March 2010 or tocilizumab for severe rheumatoid arthritis prior to 1 July 2009 and who was 'grandfathered' on to PBS-subsidised therapy, and who continues to receive treatment will have further applications for treatment with certolizumab pegol, golimumab or tocilizumab assessed under the continuing treatment restriction.

A patient may only qualify for PBS-subsidised treatment under the grandfather restriction (Initial 3 ('grandfather patients')) once. A maximum of 24 weeks of treatment with certolizumab pegol, golimumab or tocilizumab will be authorised under this restriction.

Authority required

Initial 1 (patient re-commencing after a break of more than 12 months)

Initial PBS-subsidised treatment with rituximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have severe active rheumatoid arthritis; and
- (b) have failed to respond to at least 1 PBS-subsidised TNF-alfa antagonist; and
- (c) have received no PBS-subsidised treatment with a bDMARD for this condition in the previous 12 months; and
- (d) have failed to achieve an adequate response to at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs), which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be:
 - hydroxychloroquine at a dose of at least 200 mg daily; or
 - leflunomide at a dose of at least 10 mg daily; or
 - sulfasalazine at a dose of at least 2 g daily.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, then the 6 months of intensive DMARD treatment must include at least 3 months continuous treatment with each of at least 2 of the DMARDs:

- hydroxychloroquine at a dose of at least 200 mg daily; and/or
- leflunomide at a dose of at least 10 mg daily; and/or
- sulfasalazine at a dose of at least 2 g daily.

The application must include details of the contraindication or intolerance to methotrexate. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose can be found on the Medicare Australia website [www.medicareaustralia.gov.au]. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved product information or cannot be tolerated at the doses specified above, then one or more of the following DMARDs may be used in place of these agents in order to satisfy the requirement for a trial of 6 months of intensive DMARD therapy with at least 2 DMARDs taken continuously for at least

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

3 months each:

- azathioprine at a dose of at least 1 mg/kg per day; and/or
- cyclosporin at a dose of at least 2 mg/kg/day; and/or
- sodium aurothiomalate at a dose of 50 mg weekly.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances. Details of the toxicities, including severity, which will be accepted as a reason for substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application. Details of the toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy can be found on the Medicare Australia website [www.medicareaustralia.gov.au].

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (i) a total active joint count of at least 20 active (swollen and tender) joints; or
- (ii) at least 4 active joints from the following list of major joints:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)]; and
- (3) a signed patient acknowledgement.

A maximum of two infusions will be authorised under this restriction.

Assessment of a patient's response to an initial course of treatment must be made at least 12 weeks after the first infusion so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to Medicare Australia within 4 weeks of the date it was conducted.

Where a response assessment is not undertaken and submitted to Medicare Australia within these timeframes, the patient will be deemed to have failed to respond to treatment with rituximab.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

Patients who fail to demonstrate a response to treatment with rituximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Patients who fail to demonstrate a response to rituximab treatment and who qualify to trial an alternate bDMARD according to the interchangeability arrangements for bDMARDs for the treatment of severe rheumatoid arthritis, may do so without having to have a 22 week treatment-free period.

Authority required

Initial 2 (change or re-commencement after break of less than 12 months)

Initial course of PBS-subsidised treatment with rituximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults who:

- (a) have a documented history of severe active rheumatoid arthritis; and
- (b) have failed to respond to at least 1 PBS-subsidised TNF-alfa antagonist; and

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

(c) have received prior PBS-subsidised bDMARD treatment for this condition and are eligible to receive further bDMARD therapy.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

Applications for patients who have received PBS-subsidised treatment with rituximab and who wish to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised rituximab treatment, within the timeframes specified below.

A maximum of two infusions will be authorised under this restriction.

Where the most recent course of PBS-subsidised rituximab treatment was approved under either of the initial 1 or 2 treatment restrictions patients must be assessed for response at least 12 weeks after the first infusion. This assessment must be provided to Medicare Australia no later than 4 weeks from the date of assessment.

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent provided they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The demonstration of response must be submitted to Medicare Australia within 4 weeks of assessment.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

Patients who fail to demonstrate a response to treatment with rituximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Patients who fail to demonstrate a response to rituximab treatment and who qualify to trial an alternate bDMARD according to the interchangeability arrangements for bDMARDs for the treatment of severe rheumatoid arthritis, may do so without having to have a 22 week treatment-free period.

Authority required

Continuing treatment

Continuing PBS-subsidised treatment with rituximab, in combination with methotrexate at a dose of at least 7.5 mg weekly, by a rheumatologist or clinical immunologist with expertise in the management of rheumatoid arthritis, of adults:

- (a) who have a documented history of severe active rheumatoid arthritis; and
- (b) who have demonstrated an adequate response to treatment with rituximab; and
- (c) whose most recent course of PBS-subsidised bDMARD treatment was with rituximab.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

A maximum of two infusions will be authorised under this restriction.

Patients may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The demonstration of response must be submitted to Medicare Australia within 4 weeks of assessment.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

Patients who fail to demonstrate a response to treatment with rituximab under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note

Special Pricing Arrangements apply.

9544H	Solution for I.V. infusion 500 mg in 50 mL	1	..	..	2263.57	Mabthera	RO

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

THALIDOMIDE

Caution

Thalidomide is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and for 1 month after cessation of treatment.

Authority required (STREAMLINED)

3342

Multiple myeloma.

Note

Patients receiving thalidomide under the PBS listing must be registered in the i-access risk management program.

9566L	Capsule 50 mg	112	..	..	*1680.00	Thalomid	CJ
9667T	Capsule 100 mg	56	..	..	*1680.00	Thalomid	CJ

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Musculo-skeletal system

Muscle relaxants

Muscle relaxants, centrally acting agents

Other centrally acting agents

BACLOFEN

Authority required (STREAMLINED)

3318

Severe chronic spasticity, where oral antispastic agents have failed or have caused unacceptable side effects, in patients with chronic spasticity of cerebral origin;

3319

Severe chronic spasticity, where oral antispastic agents have failed or have caused unacceptable side effects, in patients with chronic spasticity due to multiple sclerosis;

3320

Severe chronic spasticity, where oral antispastic agents have failed or have caused unacceptable side effects, in patients with chronic spasticity due to spinal cord injury;

3321

Severe chronic spasticity, where oral antispastic agents have failed or have caused unacceptable side effects, in patients with chronic spasticity due to spinal cord disease.

5617P	Intrathecal injection 10 mg in 5 mL	10	..	..	*1483.70		Lioresal Intrathecal	NV

Drugs for treatment of bone diseases

Drugs affecting bone structure and mineralization

Bisphosphonates

DISODIUM PAMIDRONATE

Authority required (STREAMLINED)

3341

Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.

Note

The concentrated injection 15 mg and powder for I.V. infusion 15 mg (after reconstitution) are bioequivalent.

5667G	Concentrated injection 15 mg in 5 mL	4	2	..	*214.20	^a	Pamisol	HH
5701C	Injection set containing 4 vials powder for I.V. infusion 15 mg and 4 ampoules solvent 5 mL	1	2	..	214.19	^a	Aredia 15 mg	NV

DISODIUM PAMIDRONATE

Authority required (STREAMLINED)

3341

Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.

Note

The concentrated injection 30 mg and powder for I.V. infusion 30 mg (after reconstitution) are bioequivalent.

5668H	Concentrated injection 30 mg in 10 mL	2	2	..	*214.20	^a	Pamisol	HH
5702D	Injection set containing 2 vials powder for I.V. infusion 30 mg and 2 ampoules solvent 10 mL	1	2	..	214.19	^a	Aredia 30 mg	NV

DISODIUM PAMIDRONATE

Authority required (STREAMLINED)

3341

Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.

5669J	Concentrated injection 60 mg in 10 mL	1	2	..	214.18		Pamisol	HH

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty \$		
DISODIUM PAMIDRONATE							
<u>Authority required (STREAMLINED)</u>							
3341							
Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.							
<u>Authority required (STREAMLINED)</u>							
3342							
Multiple myeloma;							
3343							
Bone metastases from breast cancer.							
<u>Note</u>							
The concentrated injection 90 mg and powder for I.V. infusion 90 mg (after reconstitution) are bioequivalent.							
5670K	Concentrated injection 90 mg in 10 mL	1	11	..	321.28	^a	Pamisol HH
5703E	Injection set containing 1 vial powder for I.V. infusion 90 mg and 1 ampoule solvent 10 mL	1	11	..	321.28	^a	Aredia 90 mg NV
IBANDRONIC ACID							
<u>Authority required (STREAMLINED)</u>							
3343							
Bone metastases from breast cancer.							
5750P	Concentrated injection for I.V. infusion 6 mg (as ibandronate sodium monohydrate) in 6 mL	1	11	..	341.36		Bondronat HH
ZOLEDRONIC ACID							
<u>Authority required (STREAMLINED)</u>							
3342							
Multiple myeloma;							
3343							
Bone metastases from breast cancer;							
3422							
Bone metastases from hormone-resistant prostate cancer, with demonstration of biochemical progression of disease despite maximal therapy with hormonal treatments;							
3341							
Treatment of hypercalcaemia of malignancy refractory to anti-neoplastic therapy.							
<u>Note</u>							
Special Pricing Arrangements apply.							
9653C	Injection concentrate for I.V. infusion 4 mg (as monohydrate) in 5 mL	1	11	..	450.00		Zometa NV

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Nervous system

Anti-Parkinson drugs

Dopaminergic agents

Dopamine agonists

APOMORPHINE HYDROCHLORIDE

Authority required (STREAMLINED)

3314

Parkinson's disease in patients severely disabled by motor fluctuations which do not respond to other therapy.

5609F	Injection 20 mg in 2 mL	5	..	..	77.86	Apomine	HH
5610G	Injection 50 mg in 5 mL	5	..	..	194.65	APO-go	HH
5611H	Solution for subcutaneous infusion 50 mg in 10 mL pre-filled syringe	5	..	..	194.65	Apomine PFS	HH

Psycholeptics

Antipsychotics

Diazepines, oxazepines, thiazepines and oxepines

CLOZAPINE

Authority required (STREAMLINED)

3326

Schizophrenia in patients who are non-responsive to other neuroleptic agents;

3327

Schizophrenia in patients who are intolerant of other neuroleptic agents.

5626D	Tablet 50 mg	100	..	..	138.30	Clopine 50	HH
5627E	Tablet 200 mg	100	..	..	518.62	Clopine 200	HH
5628F	Tablet 25 mg	100	..	..	69.15	^a Clopine 25	HH
						^a Clozaril 25	NV
5629G	Tablet 100 mg	100	..	..	259.31	^a Clopine 100	HH
						^a Clozaril 100	NV
5630H	Oral liquid 50 mg per mL, 100 mL	1	..	..	135.00	Clopine Suspension	HH

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Respiratory system

Cough and cold preparations

Expectorants, excl. combinations with cough suppressants

Mucolytics

DORNASE ALFA

Authority required (STREAMLINED)

3344

Use by cystic fibrosis patients who satisfy all of the following criteria:

- (1) are 5 years of age or older;
- (2) have a FVC greater than 40% predicted for age, gender and height;
- (3) have evidence of chronic suppurative lung disease (cough and sputum most days of the week, or greater than 3 respiratory tract infections of more than 2 weeks' duration in any 12 months, or objective evidence of obstructive airways disease);
- (4) are participating in a 4 week trial as detailed below or have achieved a 10% or greater improvement in FEV1 (compared to baseline established prior to dornase alfa treatment) after a 4 week trial.

In order for patients to be eligible for participation in the HSD program, the following conditions must be met:

- (1) Patients must be assessed at cystic fibrosis clinics/centres which are under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis and the prescribing of dornase alfa under the HSD program is limited to such physicians. If attendance at such units is not possible because of geographical isolation, management (including prescribing) may be by specialist physician or paediatrician in consultation with such a unit;
- (2) The measurement of lung function is to be conducted by independent (other than the treating doctor) experienced personnel at established lung function testing laboratories, unless this is not possible because of geographical isolation;
- (3) Prior to dornase alfa therapy, a baseline measurement of FEV1 must be undertaken during a stable period of the disease;
- (4) Initial therapy is limited to 4 weeks' treatment with dornase alfa at a dose of 2.5 mg daily;
- (5) At or towards the end of the initial 4 weeks' trial, patients must be reassessed and a further FEV1 measurement be undertaken (single test under conditions as above). Patients who achieve a 10% or greater improvement in FEV1 (compared to baseline established prior to dornase alfa treatment) are eligible for continued subsidy under the HSD program at a dose of 2.5 mg daily;
- (6) Patients who fail to meet a 10% or greater improvement in FEV1 after the initial 4 weeks' treatment at a dose of 2.5 mg daily, may have 1 further trial in the next 12 months but not before 3 months after the initial trial;
- (7) Following an initial 6 months' therapy, a global assessment must be undertaken involving the patient, the patient's family (in the case of paediatric patients) and the treating physician(s) to establish that all agree that dornase alfa treatment is continuing to produce worthwhile benefits. (Dornase alfa therapy should cease if there is not general agreement of benefit as there is always the possibility of harm from unnecessary use.) Further reassessments are to be undertaken at six-monthly intervals;
- (8) Other aspects of treatment, such as physiotherapy, must be continued;
- (9) Where there is documented evidence that a patient already receiving dornase alfa therapy would have met the criteria for subsidy (i.e. satisfied the criteria for the 4 week trial and achieved a 10% or greater improvement in FEV1) then the patient is eligible to continue treatment under the HSD program. Where such evidence is not available, patients will need to satisfy the initiation and continuation criteria as for new patients. (Four weeks is considered a suitable wash-out period).

Note

It is highly desirable that all patients be included in the national cystic fibrosis patient data-base.

Authority required (STREAMLINED)

3345

Treatment of cystic fibrosis in a patient less than 5 years of age who has:

- (1) A severe clinical course with frequent respiratory exacerbations or chronic respiratory symptoms (including chronic or recurrent cough, wheeze or tachypnoea) requiring frequent hospital admissions more frequently than 3 times per year; or
- (2) Significant bronchiectasis on chest high resolution computed tomography scan; or
- (3) Severe cystic fibrosis bronchiolitis with persistent wheeze non-responsive to conventional medicines; or
- (4) Severe physiological deficit measure by forced oscillation technique or multiple breath nitrogen washout and failure to respond to conventional therapy.

In order for the patient to be eligible for participation in the HSD program, the following conditions must be met:

- (1) The patient must be assessed at a cystic fibrosis clinic/centre which is under the supervision of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis, and the prescribing of dornase alfa under the HSD program is limited to such physicians. If attendance at such a unit is not possible because of geographical isolation, management (including prescribing) may be by specialist physician or paediatrician in consultation with such a unit;
- (2) Following an initial 6 months therapy, a comprehensive assessment must be undertaken and documented involving the patient, the patient's family, the treating physician and an additional independent member of the cystic fibrosis treatment team to establish agreement that dornase alfa treatment is continuing to produce worthwhile benefit. Treatment with dornase alfa should cease if there is not agreement of benefit as there is always the possibility of harm from unnecessary use. Further reassessments are to be undertaken and documented yearly.

Note

It is highly desirable that all patients be included in the national cystic fibrosis patient data-base.

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Authority required (STREAMLINED)

3346

Grandfather — continuing for patients five years or older

Continuation of treatment of cystic fibrosis in a patient 5 years of age or older, who initiated treatment with dornase alfa at an age of less than 5 years and for whom a comprehensive assessment, involving the patient's family, the treating physician and an additional independent member of the cystic fibrosis treatment team, documents agreement that dornase alfa treatment is continuing to produce worthwhile benefit. Further reassessments are to be undertaken and documented yearly. Treatment with dornase alfa should cease if there is not agreement of benefit as there is always the possibility of harm from unnecessary use.

Note

It is highly desirable that all patients be included in the national cystic fibrosis patient data-base.

Authority required (STREAMLINED)

3347

Grandfather — for patients less than five years of age who initiated dornase alfa prior to listing

Treatment of cystic fibrosis in a patient less than 5 years of age who initiated treatment with dornase alfa prior to 1 November 2009 and for whom a comprehensive assessment, involving the patient's family, the treating physician and an additional independent member of the cystic fibrosis treatment team, documents agreement that dornase alfa treatment is continuing to produce worthwhile benefit. Further reassessments are to be undertaken and documented yearly. Treatment with dornase alfa should cease if there is not agreement of benefit as there is always the possibility of harm from unnecessary use.

Note

It is highly desirable that all patients be included in the national cystic fibrosis patient data-base.

5704F	Solution for inhalation 2.5 mg (2,500 units) in 2.5 mL	60	5	..	*2360.00	Pulmozyme	RO

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Sensory organs

Ophthalmologicals

Antiinfectives

Antivirals

GANCICLOVIR
Authority required (STREAMLINED)
3379

Cytomegalovirus retinitis in severely immunocompromised patients.

5748M	Intravitreal implant 4.5 mg	1	..	..	6000.00	Vitrasert	BU

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price for		Brand Name and Manufacturer
					Max. Qty	\$	

Various

All other therapeutic products

All other therapeutic products

Iron chelating agents

DEFERASIROX

Authority required (STREAMLINED)

3336

Chronic iron overload in adults, adolescents and children 6 years and older associated with disorders of erythropoiesis;

3337

Chronic iron overload in paediatric patients aged 2 to 5 years, associated with disorders of erythropoiesis, who are intolerant to desferrioxamine or in whom desferrioxamine has proven ineffective.

Note

Special Pricing Arrangements apply.

5654N	Tablet 125 mg (dispersible)	168	5	..	*1401.48	Exjade	NV
5655P	Tablet 250 mg (dispersible)	168	5	..	*2802.90	Exjade	NV
5656Q	Tablet 500 mg (dispersible)	168	5	..	*5605.80	Exjade	NV

DEFERIPRONE

Authority required (STREAMLINED)

3338

Iron overload in patients with thalassaemia major who are unable to take desferrioxamine therapy;

3339

Iron overload in patients with thalassaemia major in whom desferrioxamine therapy has proven ineffective.

5657R	Tablet 500 mg	600	5	..	*2703.36	Ferriprox	OA
5658T	Oral solution 100 mg per mL, 250 mL	5	5	..	*1126.40	Ferriprox	OA

DEFERRIOXAMINE MESYLATE

Authority required (STREAMLINED)

3340

Disorders of erythropoiesis associated with treatment-related chronic iron overload.

5661Y	Powder for injection 2 g	60	5	..	*2281.80	^a Hospira Pty Limited	HH
				^b 23.40	*2305.20	^a Desferal 2 g	NV
5662B	Powder for injection 500 mg	400	5	..	*3803.20	^a Hospira Pty Limited	HH
				^b 314.80	*4118.00	^a Desferal 500 mg	NV

Drugs for treatment of hyperkalemia and hyperphosphatemia

LANTHANUM

Authority required (STREAMLINED)

3390

Management of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where serum phosphate is greater than 1.6 mmol per L at the commencement of therapy.

Management includes initiation, stabilisation and review of therapy as required;

3391

Management of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where the serum calcium times phosphate product is greater than 4.0 at the commencement of therapy.

Management includes initiation, stabilisation and review of therapy as required.

Note

Not to be used in combination with sevelamer.

5780F	Tablet, chewable, 500 mg (as carbonate hydrate)	180	5	..	*523.54	Fosrenol	ZI
5781G	Tablet, chewable, 750 mg (as carbonate hydrate)	180	5	..	*790.56	Fosrenol	ZI
5782H	Tablet, chewable, 1000 mg (as carbonate hydrate)	180	5	..	*890.02	Fosrenol	ZI

HIGHLY SPECIALISED DRUGS PROGRAM (Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium \$	Dispensed Price	Brand Name and Manufacturer
					for Max. Qty \$	
SEVELAMER HYDROCHLORIDE						
<u>Authority required (STREAMLINED)</u>						
3390						
Management of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where serum phosphate is greater than 1.6 mmol per L at the commencement of therapy.						
Management includes initiation, stabilisation and review of therapy as required;						
3391						
Management of hyperphosphataemia in a patient with chronic kidney disease on dialysis whose serum phosphate is not controlled on calcium and where the serum calcium times phosphate product is greater than 4.0 at the commencement of therapy.						
Management includes initiation, stabilisation and review of therapy as required.						
<u>Note</u>						
Not to be used in combination with lanthanum.						
9546K	Tablet 800 mg	360	5	..	*620.00	Renagel GZ

SECTION 100 (BOTULINUM TOXIN PROGRAM)

Code	Name, Restriction, Manner of Administration and Form	Pack Size	Price ex manufacturer		Brand Name and Manufacturer
			\$		

SECTION 100 (BOTULINUM TOXIN PROGRAM)

BOTULINUM TOXIN TYPE A PURIFIED NEUROTOXIN COMPLEX

Note

Arrangements to prescribe this item should be made by medical practitioners with Medicare Australia, contact telephone number 1800 700 270.

Criteria for availability

Treatment of blepharospasm associated with dystonia, including benign blepharospasm and VIIth nerve disorders (hemifacial spasm) in patients 12 years and older;

Treatment of dynamic equinus foot deformity due to spasticity in an ambulant paediatric cerebral palsy patient aged from 2 to 17 years inclusive;

Continuing PBS-subsidised treatment of dynamic equinus foot deformity due to spasticity in an ambulant cerebral palsy patient 18 years of age or older who was commenced on PBS-subsidised treatment with botulinum toxin type A purified neurotoxin complex as a paediatric patient;

Treatment of spasmodic torticollis, either as monotherapy or as adjunctive therapy to current standard care.

Criteria for availability

Treatment of moderate to severe spasticity of the upper limb in a cerebral palsy patient aged from 2 to 17 years inclusive;

Continuing PBS-subsidised treatment of moderate to severe spasticity of the upper limb in a cerebral palsy patient 18 years of age or older who was commenced on PBS-subsidised treatment with botulinum toxin type A purified neurotoxin complex as a paediatric patient.

Note

Contact Medicare Australia before commencing PBS-subsidised treatment in cerebral palsy patients who have been treated for moderate to severe spasticity of the upper limb with non-PBS-subsidised botulinum toxin prior to the age of 18.

Criteria for availability

Treatment of moderate to severe spasticity [defined as MAS greater than or equal to 3 using modified Ashworth scale] of the upper limb in adults following a stroke, as second line therapy when standard management has failed (e.g. physiotherapy and/or oral spasticity agents) or as an adjunct to physical therapy.

Maximum number of treatments to be authorised is 4 (total Botox and Dysport) per upper limb per lifetime. Treatment should not be initiated until 3 months post-stroke in patients who do not have established severe contracture. Treatment should be discontinued if the patient does not respond (decrease of MAS greater than 1 in at least one joint) after two treatments.

The date of the stroke must be provided.

Contraindications to treatment include established severe contracture and known sensitivity to botulinum toxin.

Note

The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

6103F	Lyophilised powder for I.M. injection 100 units	1	415.50	Botox	AG

CLOSTRIDIUM BOTULINUM TYPE A TOXIN—HAEMAGGLUTININ COMPLEX

Note

Arrangements to prescribe this item should be made by medical practitioners with Medicare Australia, contact telephone number 1800 700 270.

Criteria for availability

Treatment of dynamic equinus foot deformity due to spasticity in an ambulant paediatric cerebral palsy patient aged from 2 to 17 years inclusive;

Continuing PBS-subsidised treatment of dynamic equinus foot deformity due to spasticity in an ambulant cerebral palsy patient 18 years of age or older who was commenced on PBS-subsidised treatment with clostridium botulinum type A toxin-haemagglutinin complex as a paediatric patient;

Treatment of spasmodic torticollis, either as monotherapy or as adjunctive therapy to current standard care.

Criteria for availability

Treatment of moderate to severe spasticity [defined as MAS greater than or equal to 3 using modified Ashworth scale] of the upper limb in adults following a stroke, as second line therapy when standard management has failed (e.g. physiotherapy and/or oral spasticity agents) or as an adjunct to physical therapy.

Maximum number of treatments to be authorised is 4 (total Botox and Dysport) per upper limb per lifetime. Treatment should not be initiated until 3 months post-stroke in patients who do not have established severe contracture. Treatment should be discontinued if the patient does not respond (decrease of MAS greater than 1 in at least one joint) after two treatments.

The date of the stroke must be provided.

Contraindications to treatment include established severe contracture and known sensitivity to botulinum toxin.

Note

The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

6293F	Lyophilised powder for I.M. injection 500 units	1	650.00	Dysport	IS

SECTION 100 (HUMAN GROWTH HORMONE)

Code	Name, Restriction, Manner of Administration and Form	Pack Size	Price ex manufacturer		Brand Name and Manufacturer
			\$		

SECTION 100 (HUMAN GROWTH HORMONE)

SOMATROPIN (Recombinant human growth hormone)

Criteria for availability
Short stature in accordance with the 'Guidelines for the Availability of Human Growth Hormone (hGH) as a Pharmaceutical Benefit'.

Genotropin branded products (including MiniQuick) are also available for the treatment of Prader-Willi Syndrome in accordance with the 'Guidelines for the Availability of Human Growth Hormone (hGH) as a Pharmaceutical Benefit for the treatment of Prader-Willi Syndrome'.

Note
These guidelines may be obtained from the Department of Health and Ageing's internet site at <http://www.health.gov.au/hGH>, or from:

Growth Hormone Program
Access and Systems Branch
Department of Health and Ageing
GPO Box 9848
CANBERRA ACT 2601
Contact telephone number (02) 6289 7274

Note
Special Pricing Arrangements apply.

6465G	Solution for injection 5 mg (15 i.u.) in 1.5 mL cartridge (with preservative)	1	315.50	Norditropin Nordiflex	NO
6466H	Solution for injection 10 mg (30 i.u.) in 1.5 mL cartridge (with preservative)	1	631.00	Norditropin Nordiflex	NO
6467J	Solution for injection 15 mg (45 i.u.) in 1.5 mL cartridge (with preservative)	1	946.50	Norditropin Nordiflex	NO

SOMATROPIN (Recombinant human growth hormone)

Criteria for availability
Short stature in accordance with the 'Guidelines for the Availability of Human Growth Hormone (hGH) as a Pharmaceutical Benefit'.

Genotropin branded products (including MiniQuick) are also available for the treatment of Prader-Willi Syndrome in accordance with the 'Guidelines for the Availability of Human Growth Hormone (hGH) as a Pharmaceutical Benefit for the treatment of Prader-Willi Syndrome'.

Note
These guidelines may be obtained from the Department of Health and Ageing's internet site at <http://www.health.gov.au/hGH>, or from:

Growth Hormone Program
Access and Systems Branch
Department of Health and Ageing
GPO Box 9848
CANBERRA ACT 2601
Contact telephone number (02) 6289 7274

6169Q	Injection 18 i.u. (6 mg) cartridge with 3.15 mL diluent (with preservative)	1	297.00	Humatrope	LY
6170R	Injection 36 i.u. (12 mg) cartridge with 3.15 mL diluent (with preservative)	1	594.00	Humatrope	LY
6266T	Injection 4 mg (12 i.u.) vial with 3.5 mL diluent (with preservative)	1	198.00	Zomacton	FP
6295H	Solution for injection 5 mg (15 i.u.) in 1.5 mL cartridge (with preservative)	1	247.50	Norditropin SimpleXx	NO
6296J	Solution for injection 10 mg (30 i.u.) in 1.5 mL cartridge (with preservative)	1	495.00	Norditropin SimpleXx	NO
6297K	Solution for injection 15 mg (45 i.u.) in 1.5 mL cartridge (with preservative)	1	742.50	Norditropin SimpleXx	NO
6312F	Injection 12 mg (36 i.u.) in 1 mL cartridge (with preservative)	1	594.00	Genotropin	PF
6313G	Injection 0.8 mg (2.4 i.u.) with diluent in single use syringe (without preservative)	7	277.20	Genotropin MiniQuick	PF
6314H	Injection 1 mg (3 i.u.) with diluent in single use syringe (without preservative)	7	346.50	Genotropin MiniQuick	PF
6315J	Injection 1.2 mg (3.6 i.u.) with diluent in single use syringe (without preservative)	7	415.80	Genotropin MiniQuick	PF
6316K	Injection 1.4 mg (4.2 i.u.) with diluent in single use syringe (without preservative)	7	485.10	Genotropin MiniQuick	PF
6317L	Injection 1.6 mg (4.8 i.u.) with diluent in single use syringe (without preservative)	7	554.40	Genotropin MiniQuick	PF
6318M	Injection 1.8 mg (5.4 i.u.) with diluent in single use syringe (without preservative)	7	623.70	Genotropin MiniQuick	PF
6319N	Injection 2 mg (6 i.u.) with diluent in single use syringe (without preservative)	7	693.00	Genotropin	PF

SECTION 100 (HUMAN GROWTH HORMONE)

Code	Name, Restriction, Manner of Administration and Form	Pack Size	Price ex manufacturer		Brand Name and Manufacturer
			\$		
6329D	Injection 8 mg (24 i.u.) vial with 1.37 mL diluent cartridge (with preservative) (for use with one.click auto-injector)	1	396.00	MiniQuick Saizen 8 mg click.easy	SG
6330E	Injection 5 mg (15 i.u.) in 1 mL cartridge (with preservative)	1	247.50	Genotropin	PF
6345Y	Injection 72 i.u. (24 mg) cartridge with 3.15 mL diluent (with preservative)	1	1188.00	Humatrope	LY
6476W	Solution for injection 5 mg (15 i.u.) in 1.5 mL cartridge (with preservative)	1	247.50	Omnitrope	SZ
9585L	Powder for injection 5 mg (15 i.u.) with diluent in pre-filled pen (with preservative)	1	247.50	Genotropin GoQuick	PF
9586M	Powder for injection 12 mg (36 i.u.) with diluent in pre-filled pen (with preservative)	1	594.00	Genotropin GoQuick	PF
9604L	Solution for injection 10 mg (30 i.u.) in 2 mL cartridge (with preservative)	1	495.00	NutropinAq	IS
9628R	Injection 0.6 mg (1.8 i.u.) with diluent in single use syringe (without preservative)	7	207.90	Genotropin MiniQuick	PF

SECTION 100 (IVF/GIFT TREATMENT)

Code	Name, Restriction, Manner of Administration and Form	Pack Size	Price ex manufacturer		Brand Name and Manufacturer
			\$		

SECTION 100 (IVF/GIFT TREATMENT)

PROGESTERONE

Criteria for availability

For luteal phase support in patients who are receiving medical treatment as described in items 13200 or 13201 of the Medicare Benefits Schedule. The luteal phase is defined as the time span from embryo transfer until implantation confirmed by positive B-hCG measurement.

Note

Supply of these items is through an accredited IVF/GIFT clinic. For enquiries relating to the IVF/GIFT Program, medical practitioners should contact Medicare Australia on 1800 700 270.

Note

Special Pricing Arrangements apply.

6366C	Vaginal gel (prolonged release) 90 mg in single dose pre-filled applicator	15	148.50	Crinone 8%	SG

PROGESTERONE

Criteria for availability

For luteal phase support in patients who are receiving medical treatment as described in items 13200 or 13201 of the Medicare Benefits Schedule. The luteal phase is defined as the time span from embryo transfer until implantation confirmed by positive B-hCG measurement.

Note

Supply of these items is through an accredited IVF/GIFT clinic. For enquiries relating to the IVF/GIFT Program, medical practitioners should contact Medicare Australia on 1800 700 270.

9608Q	Pessary 100 mg	15	50.40	Orion Laboratories Pty Ltd	ON
9609R	Pessary 200 mg	15	55.60	Orion Laboratories Pty Ltd	ON

Gonadotropins and other ovulation stimulants

Gonadotropins

CHORIOGONADOTROPIN ALFA

Criteria for availability

Patients who are receiving medical treatment as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

Note

Supply of this item is through an accredited IVF/GIFT clinic. For enquiries relating to the IVF/GIFT Program, medical practitioners should contact Medicare Australia on 1800 700 270.

Note

Special Pricing Arrangements apply.

9631X	Solution for injection 250 micrograms in 0.5 mL pre-filled syringe	1	54.80	Ovidrel	SG

FOLLITROPIN ALFA

Criteria for availability

Patients who are receiving medical treatment as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

Note

Supply of these items is through an accredited IVF/GIFT clinic. For enquiries relating to the IVF/GIFT Program, medical practitioners should contact Medicare Australia on 1800 700 270.

6431L	Injection 300 i.u. in 0.5 mL multi-dose cartridge	1	144.00	Gonal-f Pen	SG
6432M	Injection 450 i.u. in 0.75 mL multi-dose cartridge	1	216.00	Gonal-f Pen	SG
6433N	Injection 900 i.u. in 1.5 mL multi-dose cartridge	1	432.00	Gonal-f Pen	SG

FOLLITROPIN BETA

Criteria for availability

Patients who are receiving medical treatment as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

Note

Supply of these items is through an accredited IVF/GIFT clinic. For enquiries relating to the IVF/GIFT Program, medical practitioners should contact Medicare Australia on 1800 700 270.

6335K	Solution for injection 300 i.u. in 0.36 mL multi-dose cartridge	1	144.04	Puregon 300 IU/0.36 mL	SH
6336L	Solution for injection 600 i.u. in 0.72 mL multi-dose cartridge	1	288.09	Puregon 600 IU/0.72 mL	SH

SECTION 100 (IVF/GIFT TREATMENT)

Code	Name, Restriction, Manner of Administration and Form	Pack Size	Price ex	Brand Name and Manufacturer	
			manufacturer \$		
6464F	Solution for injection 900 i.u. in 1.08 mL multi-dose cartridge	1	432.11	Puregon 900 IU/1.08 mL	SH

HUMAN CHORIONIC GONADOTROPHIN

Criteria for availability

Patients who are receiving medical treatment as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

Note

Supply of these items is through an accredited IVF/GIFT clinic. For enquiries relating to the IVF/GIFT Program, medical practitioners should contact Medicare Australia on 1800 700 270.

6178E	Injection set containing 3 ampoules powder for injection 1,500 units and 3 ampoules solvent 1 mL	1	39.57	Pregnyl	SH
6181H	Powder for injection 5,000 units with solvent	1	11.49	Pregnyl	SH

CETRORELIX

Criteria for availability

For the prevention of premature luteinisation and ovulation in patients undergoing controlled ovarian stimulation, followed by oocyte pick-up and assisted reproductive techniques as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

Note

Supply of these items is through an accredited IVF/GIFT clinic. For enquiries relating to the IVF/GIFT Program, medical practitioners should contact Medicare Australia on 1800 700 270.

9599F	Powder for injection 250 micrograms (as acetate) with diluent	1	46.08	Cetrotide	SG

GANIRELIX

Criteria for availability

For the prevention of premature luteinisation and ovulation in patients undergoing controlled ovarian stimulation, followed by oocyte pick-up and assisted reproductive techniques as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

Note

Supply of these items is through an accredited IVF/GIFT clinic. For enquiries relating to the IVF/GIFT Program, medical practitioners should contact Medicare Australia on 1800 700 270.

9583J	Injection 250 micrograms (as acetate) in 0.5 mL pre-filled syringe	1	46.08	Orgalutran	SH
9584K	Injection 250 micrograms (as acetate) in 0.5 mL pre-filled syringe	5	230.40	Orgalutran	SH

SECTION 100 (OPIATE DEPENDENCE TREATMENT PROGRAM)

Code	Name, Restriction, Manner of Administration and Form	Pack Size	Price ex manufacturer		Brand Name and Manufacturer
			\$		

SECTION 100 (OPIATE DEPENDENCE TREATMENT PROGRAM)

BUPRENORPHINE

Criteria for availability

Treatment of opiate dependence, including maintenance and detoxification (withdrawal), within a framework of medical, social and psychological treatment.

Note

Treatment must be in accordance with the law of the relevant State or Territory.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

6307Y NP	Tablet (sublingual) 400 micrograms (as hydrochloride)	7	6.16	Subutex	RC
6308B NP	Tablet (sublingual) 2 mg (as hydrochloride)	7	10.50	Subutex	RC
6309C NP	Tablet (sublingual) 8 mg (as hydrochloride)	7	30.10	Subutex	RC

BUPRENORPHINE with NALOXONE

Criteria for availability

Treatment of opiate dependence within a framework of medical, social and psychological treatment.

Note

Treatment must be in accordance with the law of the relevant State or Territory.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

6470M NP	Tablet (sublingual) 2 mg (as hydrochloride)-0.5 mg (as hydrochloride)	28	46.20	Suboxone	RC
6471N NP	Tablet (sublingual) 8 mg (as hydrochloride)-2 mg (as hydrochloride)	28	132.44	Suboxone	RC

METHADONE HYDROCHLORIDE

Caution

The risk of drug dependence is high.

Criteria for availability

Treatment of opiate dependence in accordance with the law of the relevant State or Territory.

Note

Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

6171T NP	Oral liquid 25 mg per 5 mL, 200 mL	1	7.91	^a Biodone Forte	MW
				^a Sigma Pharmaceuticals (Australia) Pty Ltd	SI
6172W NP	Oral liquid 25 mg per 5 mL, 1 L	1	33.88	^a Biodone Forte	MW
				^a Sigma Pharmaceuticals (Australia) Pty Ltd	SI

SECTION 100 (SPECIAL AUTHORITY ITEMS – Private Hospital)

Code	Name, Restriction, Manner of Administration and Form	Pack Size	Price ex manufacturer		Brand Name and Manufacturer
			\$		

SECTION 100 (SPECIAL AUTHORITY ITEMS – Private Hospital)

TRASTUZUMAB

Note

Any queries concerning the arrangements to prescribe trastuzumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe trastuzumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial treatment for HER2 positive early breast cancer commencing concurrently with adjuvant chemotherapy following surgery.

The total duration of PBS-subsidised treatment (initial plus continuing) that will be authorised is 52 weeks.

HER2 positivity must be demonstrated by in situ hybridisation (ISH).

Trastuzumab must not be used in patients with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure. Cardiac function must be tested by a suitable method including, for example, ECHO or MUGA, prior to seeking the initial authority approval and then at 3 monthly intervals during treatment.

Authority applications for initial treatment must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Early Breast Cancer - PBS Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes:
 - (i) a copy of the pathology report from an Approved Pathology Authority confirming the presence of HER2 gene amplification by in situ hybridisation (ISH); and
 - (ii) a copy of the signed patient acknowledgement form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

The medical practitioner should request sufficient quantity based on the weight of the patient to provide for a maximum of 3 weeks' treatment (equivalent to the loading dose for the 3 weekly regimen, and the loading dose and 2 weekly doses for the once weekly regimen).

Authority required

Continuing treatment for HER2 positive early breast cancer where the patient has previously received treatment with PBS-subsidised trastuzumab.

The patient is eligible to receive sufficient trastuzumab to complete 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy.

Trastuzumab must not be used in patients with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure. Cardiac function must be tested by a suitable method including, for example, ECHO or MUGA, at 3 monthly intervals during treatment.

Authority applications for continuing treatment may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

The medical practitioner should request sufficient quantity based on the weight of the patient for 3 weeks' supply (equivalent to 1 dose for the 3 weekly dosing regimen, or 3 doses for the once weekly dosing regimen). Up to a maximum of 3 repeats may be authorised.

Breaks in therapy.

Where a patient has a break in trastuzumab therapy of more than 1 week but less than 6 weeks from when the last dose was due, authority approval will be granted for a new loading dose. Authority applications for new loading doses may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

6497Y	Powder for I.V. infusion 150 mg	1	1076.63	Herceptin	RO

SECTION 100 (SPECIAL AUTHORITY ITEMS – Public Hospital)

Code	Name, Restriction, Manner of Administration and Form	Pack Size	Price ex	Brand Name and Manufacturer
			manufacturer \$	

SECTION 100 (SPECIAL AUTHORITY ITEMS – Public Hospital)

TRASTUZUMAB

Note

Any queries concerning the arrangements to prescribe trastuzumab may be directed to Medicare Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authority to prescribe trastuzumab should be forwarded to:

Medicare Australia
Prior Written Approval of Specialised Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Further prescribing information is on the Medicare Australia website at www.medicareaustralia.gov.au.

Authority required

Initial treatment for HER2 positive early breast cancer commencing concurrently with adjuvant chemotherapy following surgery.

The total duration of PBS-subsidised treatment (initial plus continuing) that will be authorised is 52 weeks.

HER2 positivity must be demonstrated by in situ hybridisation (ISH).

Trastuzumab must not be used in patients with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure. Cardiac function must be tested by a suitable method including, for example, ECHO or MUGA, prior to seeking the initial authority approval and then at 3 monthly intervals during treatment.

Authority applications for initial treatment must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Early Breast Cancer - PBS Supporting Information Form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)] which includes:
 - (i) a copy of the pathology report from an Approved Pathology Authority confirming the presence of HER2 gene amplification by in situ hybridisation (ISH); and
 - (ii) a copy of the signed patient acknowledgement form [may be downloaded from the Medicare Australia website (www.medicareaustralia.gov.au)].

The medical practitioner should request sufficient quantity based on the weight of the patient to provide for a maximum of 3 weeks' treatment (equivalent to the loading dose for the 3 weekly regimen, and the loading dose and 2 weekly doses for the once weekly regimen).

Authority required

Continuing treatment for HER2 positive early breast cancer where the patient has previously received treatment with PBS-subsidised trastuzumab.

The patient is eligible to receive sufficient trastuzumab to complete 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy.

Trastuzumab must not be used in patients with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure. Cardiac function must be tested by a suitable method including, for example, ECHO or MUGA, at 3 monthly intervals during treatment.

Authority applications for continuing treatment may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

The medical practitioner should request sufficient quantity based on the weight of the patient for 3 weeks' supply (equivalent to 1 dose for the 3 weekly dosing regimen, or 3 doses for the once weekly dosing regimen). Up to a maximum of 3 repeats may be authorised.

Breaks in therapy.

Where a patient has a break in trastuzumab therapy of more than 1 week but less than 6 weeks from when the last dose was due, authority approval will be granted for a new loading dose. Authority applications for new loading doses may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

9689Y	Powder for I.V. infusion 150 mg	1	1030.21	Herceptin	RO

Section 3 – Container Prices, Fees, Standard Packs and Prices for Ready Prepared Pharmaceutical Benefits

CONTAINER PRICES FOR QUANTITIES OF READY PREPARED BENEFITS LESS THAN THE STANDARD PACK:

Injectables	150 mL vial	\$0.78
Other Items	25 mL vial	\$0.31

(The 25 mL is the most commonly used size)

FEES:

Dispensing Fee for Ready Prepared Benefits	\$6.42
Dangerous Drug Fee	\$2.71
Additional Fee for Agreed Price Ready Prepared Benefits	\$1.07

NOTE -

Standard packs and prices (including mark-up, but without dispensing fee and dangerous drug fee) are for items against the price of which an asterisk () is shown in Section 2 of the Schedule.*

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
8048N	ABCIXIMAB	10 mg in 5 mL	1@ 482.23	LY
1003T	ACICLOVIR	200 mg	25@ 31.56	AF,SZ,GM,GK
2600W	ALLOPURINOL	100 mg	100@ 3.39	AF
2157M	ALUMINIUM HYDROXIDE with MAGNESIUM HYDROXIDE	200 mg-200 mg per 5 mL, 500 mL	1@ 4.65	JT
2159P	ALUMINIUM HYDROXIDE with MAGNESIUM TRISILICATE and MAGNESIUM HYDROXIDE	250 mg-120 mg-120 mg per 5 mL, 500 mL	1@ 5.64	FM
3109P	AMILORIDE HYDROCHLORIDE	5 mg	50@ 2.28	AF
3417W	AMINO ACID FORMULA with FAT, CARBOHYDRATE, VITAMINS, MINERALS, and TRACE ELEMENTS, without METHIONINE and supplemented with DOCOSAHEXANOIC ACID	125 mL, 36	1@ 625.39	SB
9330C	AMINO ACID FORMULA with FAT, CARBOHYDRATE, VITAMINS, MINERALS and TRACE ELEMENTS without PHENYLALANINE and TYROSINE, and supplemented with DOCOSAHEXANOIC ACID	125 mL, 36	1@ 625.39	SB
2347M	AMINO ACID FORMULA without PHENYLALANINE	20 g, 30	1@ 208.07	SB
8554F		500 mg, 200	1@ 79.37	SB
8678R		1 g, 75	1@ 59.19	SB
8479G	AMINO ACID FORMULA with VITAMINS, MINERALS and LONG CHAIN POLYUNSATURATED FATTY ACIDS without PHENYLALANINE	400 g	1@ 87.15	SB
2646G	AMINO ACID FORMULA with VITAMINS and MINERALS without LYSINE and low in TRYPTOPHAN	500 g	1@ 222.29	SB
2650L		400 g	1@ 95.36	SB
9438R		20 g, 30	1@ 447.72	VF
8328H	AMINO ACID FORMULA with VITAMINS and MINERALS without METHIONINE	500 g	1@ 222.29	SB
8416Y		500 g	1@ 337.29	SB
8417B		400 g	1@ 95.36	SB
8677Q		20 g, 30	1@ 448.00	VF
8744F		25 g, 30	1@ 772.99	VF
9133Q		130 mL, 30	1@ 772.99	VF
3443F	AMINO ACID FORMULA with VITAMINS and MINERALS without METHIONINE, THREONINE and VALINE and low in ISOLEUCINE	25 g, 30	1@ 772.99	VF
3444G		20 g, 30	1@ 448.00	VF
8058D		400 g	1@ 95.36	SB
8059E		500 g	1@ 222.29	SB
8061G		500 g	1@ 337.29	SB
1411G	AMINO ACID FORMULA with VITAMINS and MINERALS without PHENYLALANINE	18.2 g, 60	1@ 544.55	SB
2382J		87 mL, 30	1@ 257.09	VF
2474F		174 mL, 30	1@ 511.90	VF
2738D		500 g	1@ 109.70	SB
2739E		500 g	1@ 168.25	SB
8545R		400 g	1@ 105.27	AB
8555G		20 g, 30	1@ 221.15	VF
8591E		25 g, 30	1@ 385.68	VF
8613H		29 g, 30	1@ 221.42	SB
8727H		50 g, 30	1@ 501.88	SB
8746H		250 mL	18@ 261.37	SB
8804J		27.8 g, 30	1@ 514.34	SB
8846N		130 mL, 30	1@ 385.48	VF
9021T		125 mL, 30	1@ 514.34	SB
9396M		125 mL, 36	1@ 315.86	SB
9397N		62.5 mL, 60	1@ 526.47	SB
3078B	AMINO ACID FORMULA with VITAMINS and MINERALS without PHENYLALANINE and TYROSINE	500 g	1@ 337.29	SB
8445L		400 g	1@ 95.36	SB
8446M		500 g	1@ 222.29	SB
8631G		20 g, 30	1@ 448.00	VF
8667E		25 g, 30	1@ 772.99	VF
9132P		130 mL, 30	1@ 772.99	VF
9395L		29 g, 30	1@ 448.51	SB
2375B	AMINO ACID FORMULA with VITAMINS and MINERALS without VALINE, LEUCINE and ISOLEUCINE	130 mL, 30	1@ 772.99	VF

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
2380G		400 g	1@ 95.36	SB
8057C		500 g	1@ 337.29	SB
8260R		500 g	1@ 222.29	SB
8310J		500 g	1@ 666.39	SB
8592F		20 g, 30	1@ 448.00	VF
8632H		25 g, 30	1@ 772.99	VF
8745G		29 g, 30	1@ 448.51	SB
9499Y	AMINO ACID FORMULA with VITAMINS and MINERALS without VALINE, LEUCINE and ISOLEUCINE with FAT, CARBOHYDRATE and TRACE ELEMENTS and supplemented with DOCOSAHEXANOIC ACID	125 mL, 36	1@ 625.39	SB
2244D	AMINO ACIDS—SYNTHETIC, FORMULA	400 g	1@ 44.34	SB
2250K		400 g	1@ 43.77	AB
2553J		400 g	1@ 44.34	SB
3066J		400 g	1@ 44.34	SB
8443J		400 g	1@ 44.34	SB
8574G		400 g	1@ 44.34	AB
8575H		400 g	1@ 44.34	AB
8754R		400 g	1@ 44.34	SB
8755T		400 g	1@ 44.34	SB
2246F	AMINO ACID SYNTHETIC FORMULA supplemented with LONG CHAIN POLYUNSATURATED FATTY ACIDS	400 g	1@ 45.18	SB
2560R		400 g	1@ 45.18	SB
9339M		400 g	1@ 45.18	AB
9340N		400 g	1@ 45.18	AB
8736T	AMISULPRIDE	100 mg per mL, 60 mL	1@ 71.16	SW
9386B	AMYLOPECTIN, MODIFIED LONG CHAIN	60 g, 30	1@ 186.47	VF
9437Q	ARGININE with CARBOHYDRATE	4 g containing 500 mg arginine, 30	1@ 127.40	VF
9453M	ARSENIC TRIOXIDE	10 mg in 10 mL	10@ 4031.61	PL
9092M	ATOMOXETINE HYDROCHLORIDE	10 mg (base)	28@ 107.38	LY
9093N		18 mg (base)	28@ 107.38	LY
9094P		25 mg (base)	28@ 107.38	LY
9095Q		40 mg (base)	28@ 107.38	LY
9096R		60 mg (base)	28@ 107.38	LY
1140B	BCG IMMUNOTHERAPEUTIC (Bacillus Calmette-Guérin/ Connaught strain)	6.6 to 19.2 x 10 ⁸ CFU set	1@ 151.15	SW
3486L	BENZYPENICILLIN	600 mg	1@ 3.65	CS
1775K		600 mg	1@ 3.65	CS
2647H		3 g	1@ 6.05	CS
3398W		600 mg	1@ 3.65	CS
3399X		3 g	1@ 6.05	CS
2812B	BETAMETHASONE VALERATE	200 mcg (base) per g, 100 g	1@ 9.09	SI,SH,EX,FM
2820K		200 mcg (base) per g, 100 g	1@ 9.09	SH,EX
2544X	BIPERIDEN HYDROCHLORIDE	2 mg	100@ 7.23	LM
1258F	BISACODYL	10 mg, 12	1@ 4.05	PP
1260H		10 mg, 10	1@ 4.95	BY,PP
5303D		10 mg, 10	1@ 4.95	BY,PP
5304E		10 mg, 12	1@ 4.05	PP
5307H		10 mg, 10	1@ 4.95	BY,PP
5308J		10 mg, 12	1@ 4.05	PP
2315W	BLEOMYCIN SULFATE	15,000 i.u.	1@ 46.66	HH
9117W	BORTEZOMIB	3.5 mg	1@ 1748.99	JC
9118X		3.5 mg	1@ 1748.99	JC
6097X	BUPRENORPHINE		2@ 17.53	MF
6098Y			2@ 30.26	MF
7201B			2@ 46.70	MF
3116B	CALCIUM	500 mg	60@ 6.01	IA
8740B	CALCIUM FOLINATE	equiv. to 50 mg folinic acid in 5 mL	1@ 28.50	HH
8812T		equiv. to 100 mg folinic acid in 10 mL	1@ 25.71	IT
9041W		equiv. to 300 mg folinic acid in 30 mL	1@ 74.42	HH
2419H	CARBAMAZEPINE	200 mg	100@ 11.89	NV
2422L		100 mg	100@ 6.36	NV
5039F		100 mg	100@ 6.36	NV
5040G		200 mg	100@ 11.89	NV
1153Q	CARBIMAZOLE	5 mg	100@ 12.31	LM
8369L	CARBOHYDRATE, FAT, VITAMINS, MINERALS and TRACE ELEMENTS	400 g	1@ 35.63	SB
8578L	CARBOMER	2 mg per g, 0.6 mL, 30	1@ 10.10	NV
5504Q		2 mg per g, 0.6 mL, 30	1@ 10.10	NV
8514D	CARBOMER 974	3 mg per g, 0.5 g, 30	1@ 9.88	AQ
5502N		3 mg per g, 0.5 g, 30	1@ 9.88	AQ

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
1160C	CARBOPLATIN	50 mg in 5 mL	1@ 29.72	HH,PF,IT
1161D		150 mg in 15 mL	1@ 68.23	HH,PF,IT
1162E		450 mg in 45 mL	1@ 131.91	HH,PF,IT
2324H	CARMELLOSE SODIUM	10 mg per mL, 0.4 mL, 30	1@ 9.88	AG
2338C		5 mg per mL, 0.4 mL, 30	1@ 9.88	AG
8823J		2.5 mg per mL, 0.6 mL, 24	1@ 8.50	CX
8824K		10 mg per mL, 0.6 mL, 28	1@ 9.22	CX
5505R		10 mg per mL, 0.4 mL, 30	1@ 9.88	AG
5506T		5 mg per mL, 0.4 mL, 30	1@ 9.88	AG
5509Y		2.5 mg per mL, 0.6 mL, 24	1@ 8.50	CX
5510B		10 mg per mL, 0.6 mL, 28	1@ 9.22	CX
9307W	CARMELLOSE SODIUM with GLYCERIN	5 mg-9 mg per mL, 0.4 mL, 30	1@ 9.88	AG
5561Q		5 mg-9 mg per mL, 0.4 mL, 30	1@ 9.88	AG
8315P	CEFEPIME	1 g	1@ 15.52	BQ,OE
8316Q		2 g	1@ 28.68	BQ,OE
1085D	CEFOTAXIME	1 g	1@ 2.04	SZ
1086E		2 g	1@ 3.72	SZ
5048Q		1 g	1@ 2.04	SZ
5049R		2 g	1@ 3.72	SZ
1783W	CEFTRIAXONE	500 mg	1@ 3.91	PP
1784X		1 g	1@ 6.11	RO,HH,SZ,PP
1785Y		2 g	1@ 10.84	RO,HH,SZ,PP
1256D	CEPHAZOLIN	500 mg	5@ 16.73	HH
1257E		1 g	5@ 25.25	HH
9326W		2 g	1@ 9.78	SZ,AF
1163F	CHLORAMBUCIL	2 mg	25@ 32.89	AS
1585K	CHLORTHALIDONE	25 mg	50@ 3.61	LM
2967E	CHOLESTYRAMINE	4.7 g (equiv. to 4 g cholestyramine)	1@ 32.76	SI
9249T		4.7 g (equivalent to 4 g cholestyramine)	1@ 32.76	SI
1217C	CIPROFLOXACIN	3 mg per mL, 5 mL	1@ 11.60	AQ,IQ
1811H	CLADRIBINE	10 mg in 10 mL	1@ 660.45	JC
8800E		10 mg in 5 mL	1@ 660.45	OA
1805B	CLONAZEPAM	500 mcg	100@ 6.89	AF,RO
1806C		2 mg	100@ 12.96	AF,RO
1808E		2.5 mg per mL, 10 mL	1@ 4.31	RO
5339B		2.5 mg per mL, 10 mL	1@ 4.31	RO
5342E		2.5 mg per mL, 10 mL	1@ 4.31	RO
1017M	CLOTRIMAZOLE	10 mg per g, 20 g	1@ 2.42	AF
8785J	CODEINE PHOSPHATE with PARACETAMOL	30 mg-500 mg	20@ 1.66	FM,SW,AV,CO,AL,SZ,T X
1228P	COPPER SULFATE	Tablets, 36	1@ 32.53	BN
9251X		Tablets, 36	1@ 32.53	BN
1079T	CYCLOPHOSPHAMIDE	500 mg	1@ 17.14	BX
8657P	CYCLOSPORIN	10 mg	60@ 44.00	NV
8658Q		25 mg	30@ 46.33	NV,SZ
8659R		50 mg	30@ 96.40	NV,SZ
8660T		100 mg	30@ 187.58	NV,SZ
8661W		100 mg per mL, 50 mL	1@ 353.12	NV
1270W	CYPROTERONE ACETATE	50 mg	50@ 100.65	AF,GM,SC,SY,GX,SZ
9164H	CYSTINE with CARBOHYDRATE	4 g containing 500 mg cystine, 30	1@ 127.40	VF
2884T	CYTARABINE	100 mg in 5 mL	5@ 59.68	PF
9318K	DABIGATRAN ETEXILATE	75 mg (as mesilate)	10@ 37.37	BY
9319L		110 mg (as mesilate)	10@ 37.37	BY
9322P		75 mg (as mesilate)	10@ 37.37	BY
9323Q		110 mg (as mesilate)	10@ 37.37	BY
8641T	DALTEPARIN SODIUM (Low Molecular Weight Heparin Sodium—porcine mucous)	2,500 units (anti-Xa) in 0.2 mL	10@ 49.16	PF
8642W		5,000 units (anti-Xa) in 0.2 mL	10@ 51.23	PF
8643X		7,500 units (anti-Xa) in 0.75 mL	10@ 77.23	PF
2129C	DESMOPRESSIN ACETATE	100 mcg per mL, 2.5 mL	1@ 30.95	FP
8662X		200 mcg	30@ 57.83	FP
8711L		10 mcg per actuation, 60 actuations, 6 mL	1@ 77.31	FP
6033M	DIAZEPAM		50@ 1.37	AF,RO,SI
6034N			50@ 1.51	RO,AF,SI,RA,GM
1299J	DICLOFENAC SODIUM	25 mg (e.c.)	50@ 3.32	SZ,NV,CH,TW,AF,SI,G M,TX
1302M		100 mg	20@ 9.25	NV
5361E		25 mg (e.c.)	50@ 3.32	SZ,NV,CH,TW,AF,SI,G M,TX
5363G		100 mg	20@ 9.25	NV
5364H		25 mg (e.c.)	50@ 3.32	SZ,NV,CH,TW,AF,SI,G M,TX
5366K		100 mg	20@ 9.25	NV

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$		Manufacturer
5076E		25 mg (e.c.)	50@	3.32	SZ,NV,CH,TW,AF,SI,G M,TX
5079H		100 mg	20@	9.25	NV
3164M	DIGOXIN	50 mcg per mL, 60 mL	1@	11.13	SI
3463G	DIPHTHERIA and TETANUS VACCINE, ADSORBED, DILUTED FOR ADULT USE	0.5 mL	5@	67.15	CS
8461H	DISODIUM PAMIDRONATE	15 mg in 5 mL	1@	62.08	HH
8462J		30 mg in 10 mL	1@	124.15	HH
5463M	DOCETAXEL	20 mg in 1 mL	1@	322.37	SW
8071T		20 mg (anhydrous) set	1@	322.37	SW
1336H	DOXORUBICIN HYDROCHLORIDE	10 mg in 5 mL	1@	8.51	HH,PF,IT
1340M		20 mg in 10 mL	1@	14.54	PF
1342P		50 mg in 25 mL	1@	34.39	HH,PF,IT
2702F	DOXYCYCLINE	100 mg (as hydrochloride)	7@	2.05	GM,SI,AF,PF
2703G		100 mg (as hydrochloride)	7@	2.05	YT,YN
2714W		100 mg (as hydrochloride)	7@	2.05	GM,SI,AF,PF
9107H		100 mg (as monohydrate)	7@	2.05	SZ,CH,TW,GX
9108J		100 mg (as monohydrate)	7@	2.05	SZ,CH,TW
3199J	ELECTROLYTE REPLACEMENT SOLUTION	1 L	1@	7.77	BX
8510X	ENOXAPARIN SODIUM	40 mg (4,000 i.u. anti-Xa) in 0.4 mL	10@	51.23	SW
8558K		20 mg (2,000 i.u. anti-Xa) in 0.2 mL	10@	49.16	SW
8639Q		40 mg (4,000 i.u. anti-Xa) in 0.4 mL	10@	51.23	SW
8640R		60 mg (6,000 i.u. anti-Xa) in 0.6 mL	10@	73.26	SW
8716R		20 mg (2,000 i.u. anti-Xa) in 0.2 mL	10@	49.16	SW
9195Y		40 mg (4,000 i.u. anti-Xa) in 0.4 mL	10@	51.23	SW
9196B		40 mg (4,000 i.u. anti-Xa) in 0.4 mL	10@	51.23	SW
8367J	ENTACAPONE	200 mg	100@	137.70	NV
1375J	EPIRUBICIN HYDROCHLORIDE	10 mg in 5 mL	1@	51.40	PF,IT
1376K		20 mg in 10 mL	1@	95.03	PF
1377L		50 mg in 25 mL	1@	232.82	PF,HH,IT
8817C		100 mg in 50 mL	1@	459.39	HH,IT
8397Y	EPROSARTAN MESYLATE	400 mg (base)	28@	11.33	SM
8951D		400 mg (base)	28@	12.33	SM
8683B	EPTIFIBATIDE ACETATE	20 mg (base) in 10 mL	1@	128.06	SH
8684C		75 mg (base) in 100 mL	1@	337.98	SH
1397M	ERYTHROMYCIN LACTOBIONATE	1 g (base)	1@	16.50	LM
5088T		1 g (base)	1@	16.50	LM
9329B	ESSENTIAL AMINO ACIDS FORMULA	200 g, 2	1@	398.05	SB
2027Q	ESSENTIAL AMINO ACIDS FORMULA with MINERALS and VITAMIN C	400 g	1@	125.55	SB
9385Y	ESSENTIAL AMINO ACIDS FORMULA with VITAMINS and MINERALS	12.5 g, 50	1@	377.52	VF
3445H	ETANERCEPT	25 mg and 1 mL solvent, 4	1@	911.29	WX
3448L		25 mg and 1 mL solvent, 4	1@	911.29	WX
8637N		25 mg and 1 mL solvent, 4	1@	911.29	WX
8638P		25 mg and 1 mL solvent, 4	1@	911.29	WX
8778B		25 mg and 1 mL solvent, 4	1@	911.29	WX
8779C		25 mg and 1 mL solvent, 4	1@	911.29	WX
9035M		25 mg and 1 mL solvent, 4	1@	911.29	WX
9036N		25 mg and 1 mL solvent, 4	1@	911.29	WX
9037P		25 mg and 1 mL solvent, 4	1@	911.29	WX
9429G		25 mg and 1 mL solvent, 4	1@	911.29	WX
8748K	ETHACRYNIC ACID	25 mg	100@	95.44	FK
1390E	ETOPOSIDE	100 mg in 5 mL	1@	32.05	HH
8120J		100 mg (as phosphate)	1@	32.05	BQ
8842J	EVEROLIMUS	0.75 mg	60@	786.10	NV
9352F		1 mg	60@	1031.17	NV
7202C	FENTANYL		5@	36.88	JC
7203D			5@	46.92	JC
7204E			5@	86.25	JC
7205F			5@	116.41	JC
7206G			5@	142.30	JC
5401G		200 mcg, 3	1@	35.49	OA
5402H		400 mcg, 3	1@	35.49	OA
5403J		600 mcg, 3	1@	35.49	OA
5404K		800 mcg, 3	1@	35.49	OA
5405L		1200 mcg, 3	1@	35.49	OA
5406M		1600 mcg, 3	1@	35.49	OA
5407N		200 mcg, 3	1@	33.55	OA
5408P		400 mcg, 3	1@	33.55	OA
5409Q		600 mcg, 3	1@	33.55	OA
5410R		800 mcg, 3	1@	33.55	OA
5411T		1200 mcg, 3	1@	33.55	OA

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
5412W		1600 mcg, 3	1@ 33.55	OA
1473M	FLUCONAZOLE	100 mg in 50 mL	1@ 21.85	PF,SZ,AE
1474N		200 mg in 100 mL	1@ 39.93	PF,HX,SZ,AE,BX
9185K	FLUDARABINE PHOSPHATE	50 mg	1@ 305.88	GQ,WQ
1433K	FLUDROCORTISONE ACETATE	100 mcg	100@ 20.04	SI
2528C	FLUOROURACIL	500 mg in 10 mL	1@ 4.93	HH
9005Y		1000 mg in 20 mL	1@ 8.51	IT
1437P	FOLIC ACID	5 mg	100@ 3.80	AF
2958Q		500 mcg	100@ 3.68	AF
8713N	FOLLITROPIN ALFA	300 i.u.	1@ 185.67	SG
8714P		450 i.u.	1@ 278.50	SG
8715Q		900 i.u.	1@ 554.41	SG
8565T	FOLLITROPIN BETA	300 i.u. in 0.36 mL	1@ 185.67	SH
8566W		600 i.u. in 0.72 mL	1@ 371.33	SH
8871X		900 i.u. in 1.08 mL	1@ 554.40	SH
2414C	FRUSEMIDE	20 mg	50@ 1.30	SW,FM
8444K	GELATIN - SUCCINYLATED	20 g per 500 mL, 500 mL	1@ 13.11	BR
8049P	GEMCITABINE	200 mg	1@ 49.84	LY,IT,HH,ZP,GQ,PK,ZF
8050Q		1 g	1@ 232.91	LY,IT,HH,ZP,GQ,PK,ZF
9401T		200 mg in 20 mL	1@ 49.84	IT
9402W		1000 mg in 100 mL	1@ 232.91	IT
9463C		500 mg in 50 mL	1@ 116.46	IT
2824P	GENTAMICIN SULFATE	80 mg (base) in 2 mL	5@ 6.76	HH
2245E	GLUCOSE	278 mmol per L, 1 L	1@ 3.35	BR,PK,BX
9444C		139 mmol per 500 mL, 500 mL	1@ 2.33	BR,PK
9445D		278 mmol per 500 mL, 500 mL	1@ 2.33	PK
9474P		69.5 mmol per 250 mL, 250 mL	1@ 3.52	BR,PK
5005K		139 mmol per 500 mL, 500 mL	1@ 2.33	BR,PK
5106R		278 mmol per L, 1 L	1@ 3.35	BR,PK,BX
3106L	GLUCOSE and KETONE INDICATOR—URINE	Test strips, 50	1@ 5.44	RD
3107M		Test strips, 50	1@ 5.50	BN
9254C		Test strips, 50	1@ 5.44	RD
9255D		Test strips, 50	1@ 5.50	BN
2263D	GLUCOSE INDICATOR—BLOOD	Test strips, 50	1@ 23.38	MS
2860M		Test strips, 50	1@ 23.38	NA
2890D		Test strips, 50	1@ 23.38	NA
2891E		Test strips, 50	1@ 23.38	RD
2914J		Test strips, 50	1@ 19.74	NA
2917M		Test strips, 50	1@ 19.74	BN
3406G		Test strips, 50	1@ 23.38	LB
3407H		Test strips, 50	1@ 23.38	LB
3441D		Test strips, 50	1@ 23.38	JJ
3442E		Test strips, 50	1@ 23.38	JJ
8682Y		Test strips, 50	1@ 23.38	WF
8723D		Test strips, 50	1@ 23.38	BR
8739Y		Test strips, 50	1@ 23.38	RD
8749L		Test strips, 50	1@ 23.38	OZ
8759B		Test strips, 50	1@ 23.38	LB
8795X		Test strips, 50	1@ 23.38	PX
8806L		Test strips, 51	1@ 23.38	RD
8825L		Test strips, 50	1@ 23.38	NX
8890X		Test strips, 50	1@ 23.38	MS
9013J		Test strips, 50	1@ 23.38	OZ
9193W		Test strips, 25	1@ 11.69	PZ
9256E		Test strips, 25	1@ 11.69	PZ
9258G		Test strips, 50	1@ 23.38	RD
9259H		Test strips, 50	1@ 23.38	MS
9261K		Test strips, 50	1@ 23.38	OZ
9263M		Test strips, 50	1@ 23.38	OZ
9264N		Test strips, 50	1@ 23.38	WF
9265P		Test strips, 50	1@ 23.38	BR
9267R		Test strips, 50	1@ 23.38	MS
9268T		Test strips, 50	1@ 23.38	NX
9274D		Test strips, 50	1@ 23.38	RD
9275E		Test strips, 51	1@ 23.38	RD
9276F		Test strips, 50	1@ 23.38	NA
9277G		Test strips, 50	1@ 23.38	NA
9278H		Test strips, 50	1@ 23.38	LB
9279J		Test strips, 50	1@ 19.74	NA
9280K		Test strips, 50	1@ 19.74	BN
9281L		Test strips, 50	1@ 23.38	PX
9297H		Test strips, 50	1@ 23.38	QB
9298J		Test strips, 50	1@ 23.38	QB

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
9324R		Test strips, 50	1@ 23.38	HE
9325T		Test strips, 50	1@ 23.38	HE
9471L		Test strips, 50	1@ 23.38	EH
9472M		Test strips, 50	1@ 23.38	EH
9485F		Test strips, 50	1@ 23.38	OI
9486G		Test strips, 50	1@ 23.38	OI
2352T	GLUCOSE INDICATOR—URINE	Test strips, 50	1@ 6.70	BN
3104J		Test strips, 50	1@ 6.21	BN
9252Y		Test strips, 50	1@ 6.70	BN
9253B		Test strips, 50	1@ 6.21	BN
2555L	GLYCEROL	700 mg, 12	1@ 4.14	PP
2556M		1.4 g, 12	1@ 4.28	PP
2557N		2.8 g, 12	1@ 4.44	PP
5311M		700 mg, 12	1@ 4.14	PP
5312N		1.4 g, 12	1@ 4.28	PP
5313P		2.8 g, 12	1@ 4.44	PP
5314Q		700 mg, 12	1@ 4.14	PP
5315R		1.4 g, 12	1@ 4.28	PP
5316T		2.8 g, 12	1@ 4.44	PP
8728J	GRANISETRON HYDROCHLORIDE	2 mg (base)	1@ 26.28	HH
8729K		3 mg (base) in 3 mL	5@ 91.62	PK
8730L		3 mg (base) in 3 mL	5@ 91.62	PK
1076P	HEPARIN SODIUM	35,000 units in 35 mL	1@ 22.68	HH
9446E	HIGH FAT FORMULA with VITAMINS, MINERALS and TRACE ELEMENTS and low in PROTEIN and CARBOHYDRATE	300 g	1@ 40.91	SB
1639G	HYDRALAZINE HYDROCHLORIDE	50 mg	100@ 5.50	AF
1640H		25 mg	100@ 4.54	AF
1486F	HYDROCHLOROTHIAZIDE with AMILORIDE HYDROCHLORIDE	50 mg-5 mg	50@ 3.73	AS
1502C	HYDROCORTISONE ACETATE	21.1 g	1@ 15.64	AS
3470P	HYDROCORTISONE SODIUM SUCCINATE	100 mg with 2 mL solvent	1@ 5.15	PF
1501B		100 mg with 2 mL solvent	1@ 5.15	PF
1510L		100 mg with 2 mL solvent	1@ 5.15	PF
1511M		250 mg with 2 mL solvent	1@ 8.90	PF
5118J		100 mg with 2 mL solvent	1@ 5.15	PF
5119K		250 mg with 2 mL solvent	1@ 8.90	PF
6062C	HYDROMORPHONE HYDROCHLORIDE		1@ 54.57	MF
6075R			20@ 7.97	MF
6076T			20@ 10.72	MF
6077W			20@ 20.42	MF
7217W			14@ 26.09	JC
7218X			14@ 43.11	JC
7219Y			14@ 79.57	JC
7220B			14@ 136.50	JC
7221C			14@ 21.22	JC
9487H	HYDROXYETHYL STARCH 130/0.4	30 g per 500 mL, 500 mL	1@ 13.11	PK
5317W	HYOSCINE BUTYLBROMIDE	20 mg in 1 mL	5@ 17.02	BY
5318X		20 mg in 1 mL	5@ 17.02	BY
8299T	HYPROMELLOSE with DEXTRAN	3 mg-1 mg per mL, 0.4 mL, 28	1@ 9.74	AQ
5521N		3 mg-1 mg per mL, 0.4 mL, 28	1@ 9.74	AQ
3190X	IBUPROFEN	400 mg	30@ 2.77	AB
5368M		400 mg	30@ 2.77	AB
5370P		400 mg	30@ 2.77	AB
5123P		400 mg	30@ 2.77	AB
2446R	IDARUBICIN HYDROCHLORIDE	5 mg	1@ 80.23	PF
2448W		10 mg	1@ 148.40	PF
8076C	IFOSFAMIDE	1 g	1@ 67.20	BX
8077D		2 g	1@ 132.39	BX
2454E	INDOMETHACIN	25 mg	50@ 2.83	AF,AS
2757D		100 mg	20@ 8.04	AS
5377B		25 mg	50@ 2.83	AF,AS
5378C		100 mg	20@ 8.04	AS
5379D		25 mg	50@ 2.83	AF,AS
5380E		100 mg	20@ 8.04	AS
5126T		25 mg	50@ 2.83	AF,AS
5128X		100 mg	20@ 8.04	AS
8435Y	INSULIN ASPART	100 units per mL, 3 mL, 5	1@ 51.56	NO,NF
8571D		100 units per mL, 10 mL	1@ 30.57	NO
8609D	INSULIN ASPART—INSULIN ASPART PROTAMINE SUSPENSION	100 units (30 units-70 units) per mL, 3 mL, 5	1@ 51.56	NF,NO
9040T	INSULIN DETEMIR	100 units per mL, 3 mL, 5	1@ 85.26	NF,NO
9039R	INSULIN GLARGINE	100 units per mL, 3 mL, 5	1@ 85.26	SW,AV

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
1921D	INSULIN GLULISINE	100 units per mL, 3 mL, 5	1@ 51.56	AV,SW
9224L		100 units per mL, 10 mL	1@ 30.57	SW
1533Q	INSULIN ISOPHANE (N.P.H.)	100 units per mL, 10 mL	1@ 25.48	LY,NO
1711C		100 units per mL, 10 mL	1@ 33.12	AS
1761Q		100 units per mL, 3 mL, 5	1@ 43.58	NO,NL,NI,LY
8084L	INSULIN LISPRO	100 units per mL, 10 mL	1@ 30.57	LY
8212F		100 units per mL, 3 mL, 5	1@ 51.56	LY,KP
8390N	INSULIN LISPRO—INSULIN LISPRO PROTAMINE SUSPENSION	100 units (25 units-75 units) per mL, 3 mL, 5	1@ 51.56	LY,KP
8874C		100 units (50 units-50 units) per mL, 3 mL, 5	1@ 51.56	LY,KP
1531N	INSULIN NEUTRAL	100 units per mL, 10 mL	1@ 25.48	NO,LY
1713E		100 units per mL, 10 mL	1@ 33.12	AS
1762R		100 units per mL, 3 mL, 5	1@ 43.58	NO,LY
1426C	INSULIN NEUTRAL—INSULIN ISOPHANE (N.P.H.), (MIXED) (Biphasic Isophane)	100 units (30 units-70 units) per mL, 10 mL	1@ 25.48	LY
1763T		100 units (30 units-70 units) per mL, 3 mL, 5	1@ 43.58	LY,NO,NI
2062M		100 units (50 units-50 units) per mL, 3 mL, 5	1@ 43.58	NO
8180M	INTERFERON ALFA-2a	3,000,000 i.u. in 0.5 mL	1@ 33.32	RO
8181N		3,000,000 i.u. in 0.5 mL	1@ 33.32	RO
8182P		4,500,000 i.u. in 0.5 mL	1@ 51.66	RO
8183Q		6,000,000 i.u. in 0.5 mL	1@ 67.66	RO
8184R		9,000,000 i.u. in 0.5 mL	1@ 99.94	RO
8551C		4,500,000 i.u. in 0.5 mL	1@ 51.66	RO
8552D		6,000,000 i.u. in 0.5 mL	1@ 67.66	RO
8553E		9,000,000 i.u. in 0.5 mL	1@ 99.94	RO
8348J	INTERFERON ALFA-2b	18,000,000 i.u. in 1.2 mL	1@ 199.87	SH
8476D		30,000,000 i.u. in 1.2 mL	1@ 333.11	SH
8572E		18,000,000 i.u. in 1.2 mL	1@ 199.87	SH
1542E	IPRATROPIUM BROMIDE	250 mcg (anhydrous) in 1 mL, 30	1@ 15.43	AF,PF,SI,TX,BY
8238N		500 mcg (anhydrous) in 1 mL, 30	1@ 18.11	AF,BY,PF,SI,TX
8671J		21 mcg per dose (200 doses)	1@ 14.43	BY
8415X	IRINOTECAN HYDROCHLORIDE TRIHYDRATE	100 mg in 5 mL	1@ 310.80	HH,SZ,PF,OE,GQ,AF,IT, WQ
9134R	ISOLEUCINE with CARBOHYDRATE	4 g containing 50 mg isoleucine, 30	1@ 127.40	VF
9436P		4 g containing 1 g isoleucine, 30	1@ 140.14	VF
2587E	ISOSORBIDE DINITRATE	10 mg	100@ 3.70	AF
2588F		5 mg	100@ 4.15	SI
1588N	KETOPROFEN	100 mg	20@ 9.44	SW
5139L		100 mg	20@ 9.44	SW
5387M	LACTULOSE	3.34 g per 5 mL, 500 mL	1@ 7.81	AF,GM,SI,GX,SZ,SM
5388N		3.34 g per 5 mL, 500 mL	1@ 7.81	AF,GM,SI,GX,SZ,SM
9148L	LAPATINIB	250 mg (as ditosylate monohydrate)	70@ 1690.52	GK
8797B	LEVODOPA with CARBIDOPA and ENTACAPONE	50 mg-12.5 mg-200 mg	100@ 152.73	NV
8798C		100 mg-25 mg-200 mg	100@ 167.75	NV
8799D		150 mg-37.5 mg-200 mg	100@ 182.77	NV
9292C		200 mg-50 mg-200 mg	100@ 196.60	NV
9344T		75 mg-18.75 mg-200 mg	100@ 159.35	NV
9345W		125 mg-31.25 mg-200 mg	100@ 173.77	NV
8290H	LITHIUM CARBONATE	450 mg (s.r.)	100@ 13.94	GK
5389P	MACROGOL 3350	13.125 g, 30	1@ 14.13	NE
5390Q		13.125 g, 30	1@ 14.13	NE
5426N		510 g	1@ 14.13	KY,ON
5427P		510 g	1@ 14.13	KY,ON
1598D	MERCAPTOPYRINE	50 mg	25@ 61.38	AS
2214M	MESALAZINE	500 mg (p.r.)	100@ 145.51	FP
3413P		1 g (p.r.)	60@ 162.13	FP
8598M		500 mg	100@ 145.51	OA
8616L		2 g in 60 mL, 7	1@ 82.45	OA
8617M		4 g in 60 mL, 7	1@ 109.87	OA
8731M		500 mg (e.c.)	100@ 145.51	OA
8753Q		1 g in 100 mL, 7	1@ 82.45	FP
8768L		80 g	1@ 82.45	OA
6035P	METHADONE HYDROCHLORIDE		5@ 39.25	SI
6037R			20@ 6.22	SI
5423K	METHYLNALTREXONE	12 mg in 0.6 mL	1@ 41.39	WX
2826R	METHYSERGIDE	1 mg	50@ 19.27	LM
1638F	METRONIDAZOLE	500 mg in 100 mL	10@ 39.28	SZ,HH
5154G		500 mg in 100 mL	10@ 39.28	SZ,HH
9026C	MICONAZOLE NITRATE	20 mg per g, 15 g	1@ 4.74	JT
2349P	MILK POWDER—LACTOSE FREE FORMULA	900 g	1@ 16.50	NU
2350Q		900 g	1@ 16.50	NU
8282X		900 g	1@ 21.29	WX

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
8283Y		900 g	1@ 21.29	WX
2357C	MILK POWDER—LACTOSE MODIFIED	900 g	1@ 22.13	SJ
2358D		900 g	1@ 22.13	SJ
3092R	MILK POWDER—SYNTHETIC	400 g	1@ 46.87	SB
8630F	MILK PROTEIN and FAT FORMULA with VITAMINS and MINERALS—CARBOHYDRATE FREE	225 g	1@ 26.75	SB
8816B	MODAFINIL	100 mg	60@ 170.28	CS
6047G	MORPHINE HYDROCHLORIDE		1@ 9.03	MF
6048H			1@ 11.65	MF
6049J			1@ 15.79	MF
6038T	MORPHINE SULFATE		20@ 7.95	SI,MF
6039W			20@ 5.00	FM
6040X			20@ 19.20	SI,MF
6041Y			20@ 32.75	SI,MF
6042B			20@ 46.20	SI,MF
6050K			20@ 11.57	GK
6051L			20@ 23.91	GK
6052M			20@ 44.70	GK
6053N			20@ 6.16	MF
6054P			20@ 38.59	MF
6055Q			20@ 44.26	MF
6056R			20@ 56.29	MF
6057T			20@ 7.95	GK
6064E			20@ 10.99	MF
6065F			20@ 37.29	MF
6066G			10@ 10.99	MF
6067H			10@ 19.20	MF
6068J			10@ 22.72	MF
6069K			10@ 32.75	MF
8649F	MYCOPHENOLATE MOFETIL	250 mg	100@ 207.13	RO
8650G		500 mg	50@ 207.13	RO
3482G	NALOXONE HYDROCHLORIDE	2 mg in 5 mL	1@ 35.83	CS
1674D	NAPROXEN	250 mg	50@ 3.65	AF,RO
5345H		250 mg	50@ 3.65	AF,RO
5349M		250 mg	50@ 3.65	AF,RO
5176K		250 mg	50@ 3.65	AF,RO
8298R	NARATRIPTAN	2.5 mg (as hydrochloride)	2@ 9.74	GK
9734H		2.5 mg (as hydrochloride)	2@ 11.13	GK
9316H	NEBIVOLOL	1.25 mg (as hydrochloride)	28@ 22.10	CS
9171Q	NILOTINIB	200 mg (as hydrochloride monohydrate)	28@ 1371.00	NV
9285Q		200 mg (as hydrochloride monohydrate)	28@ 1371.00	NV
2732T	NITRAZEPAM	5 mg	25@ 1.47	AF,VT
5359C		5 mg	25@ 1.47	AF,VT
5360D		5 mg	25@ 1.47	AF,VT
1967M	NORETHISTERONE	350 mcg	1@ 2.55	KR,JC,PF
2772X	NORETHISTERONE with ETHINYLOESTRADIOL	500 mcg-35 mcg	1@ 2.55	PF
2773Y		1 mg-35 mcg	1@ 2.55	PF
2774B		Tablet-Pack	1@ 2.55	KR,PF
2775C		Tablet-Pack	1@ 2.55	KR,PF
2776D		Tablet-Pack	1@ 2.55	KR,PF
3176E	NORETHISTERONE with MESTRANOL	1 mg-50 mcg	1@ 2.55	PF
3179H		Tablet-Pack	1@ 2.55	PF
1698J	NYSTATIN	100,000 units per g, 15 g	1@ 6.07	FM
8383F	OFLOXACIN	3 mg per mL, 5 mL	1@ 12.86	AG
9294E	OLANZAPINE	210 mg	1@ 246.68	LY
9295F		300 mg	1@ 401.42	LY
3134Y	OXAZEPAM	15 mg	25@ 1.09	AF,SI
3135B		30 mg	25@ 1.25	TX,FM,SI,AF
5371Q		15 mg	25@ 1.09	AF,SI
5372R		30 mg	25@ 1.25	TX,FM,SI,AF
5373T		15 mg	25@ 1.09	AF,SI
5374W		30 mg	25@ 1.25	TX,FM,SI,AF
8588B	OXCARBAZEPINE	60 mg per mL, 250 mL	1@ 65.85	NV
6044D	OXYCODONE		12@ 33.03	PL
6043C	OXYCODONE HYDROCHLORIDE		20@ 3.17	SI
6058W			20@ 12.82	MF
6059X			20@ 22.02	MF
6060Y			20@ 38.19	MF
6061B			20@ 62.57	MF
6063D			20@ 3.17	MF
6070L			20@ 6.29	MF
6071M			20@ 11.02	MF

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
6078X			1@ 11.59	MF
6079Y			20@ 12.05	MF
7215R			20@ 18.60	MF
7216T			20@ 30.69	MF
3017T	PACLITAXEL	150 mg in 25 mL	1@ 894.95	HH,IT,GQ,WQ
3026G		30 mg in 5 mL	1@ 185.10	BQ,HH,GQ,WQ,PK
8018B		100 mg in 16.7 mL	1@ 613.97	BQ,HH,IT,GQ,WQ,PK
5453B	PANCREATIC EXTRACT	20 g	1@ 45.12	SM
5454C		20 g	1@ 45.12	SM
8020D		not less than 10,000 BP units lipase activity	100@ 32.87	SM
8021E		not less than 25,000 BP units lipase activity	100@ 65.74	SM
8556H		not less than 5,000 BP units lipase activity	100@ 22.56	SM
9225M		not less than 5,000 BP units lipase activity	100@ 22.56	SM
9226N		not less than 10,000 BP units lipase activity	100@ 32.87	SM
9227P		not less than 25,000 BP units lipase activity	100@ 65.74	SM
9412J		not less than 40,000 BP units lipase activity	100@ 104.60	SM
9413K		not less than 40,000 BP units lipase activity	100@ 104.60	SM
8366H	PANCRELIPASE	not less than 25,000 BP units lipase activity	100@ 65.74	TM
9229R		not less than 25,000 BP units lipase activity	100@ 65.74	TM
8784H	PARACETAMOL	500 mg	100@ 2.00	GM,TW,CH,SZ,YM,TX, GQ,SW,FM
8814X		665 mg (m.r.)	96@ 5.11	GC
5319Y		500 mg, 24	1@ 19.51	GC
5320B		500 mg, 24	1@ 19.51	GC
5343F		665 mg (m.r.)	96@ 5.11	GC
5344G		665 mg (m.r.)	96@ 5.11	GC
5224Y		500 mg	100@ 2.00	GM,TW,CH,SZ,YM,TX, GQ,SW,FM
1754H	PARAFFIN	3.5 g	1@ 7.56	IQ,AQ
9217D		3.5 g	1@ 7.56	IQ,AQ
5523Q		3.5 g	1@ 7.56	IQ,AQ
1166J	PHENOXYBENZAMINE HYDROCHLORIDE	10 mg, 30	1@ 66.16	GH
1703P	PHENOXYMETHYLPENICILLIN	250 mg	25@ 2.58	SI
1787C		250 mg	25@ 2.58	SI
3028J		500 mg	25@ 3.82	SI
9143F		150 mg per 5 mL, 100 mL	1@ 7.99	SI,FM
3360W		250 mg	25@ 2.58	SI
3361X		500 mg	25@ 3.82	SI
5012T		150 mg per 5 mL, 100 mL	1@ 7.99	SI,FM
9384X	PHENYLALANINE with CARBOHYDRATE	4 g containing 50 mg phenylalanine, 30	1@ 127.40	VF
9493P	POLYETHYLENE GLYCOL 400	2.5 mg per mL, single dose units 0.4 mL, 20	1@ 6.59	AO
5560P		2.5 mg per mL, single dose units 0.4 mL, 20	1@ 6.59	AO
9170P	POLYETHYLENE GLYCOL 400 with PROPYLENE GLYCOL	4 mg-3 mg per mL, single dose units 0.8 mL, 28	1@ 13.83	AQ
5532E		4 mg-3 mg per mL, single dose units 0.8 mL, 28	1@ 13.83	AQ
2334W	POLYGELINE	17.5 g per 500 mL, 500 mL	1@ 13.11	AE
9475Q	POLY-L-LACTIC ACID	150 mg	1@ 220.02	SW
9476R		150 mg	1@ 220.02	SW
2642C	POTASSIUM CHLORIDE	600 mg	100@ 3.30	NM,NV
1920C	PREDNISOLONE SODIUM PHOSPHATE	equiv. to 20 mg prednisolone in 100 mL	7@ 53.69	SI
2554K		equiv. to 5 mg prednisolone, 10	1@ 12.30	SI
3488N	PROMETHAZINE HYDROCHLORIDE	50 mg in 2 mL	5@ 7.95	HH
1948M		50 mg in 2 mL	5@ 7.95	HH
3374N		50 mg in 2 mL	5@ 7.95	HH
1953T	PROPANTHELINE BROMIDE	15 mg	100@ 10.02	SI
1955X	PROPYLTHIOURACIL	50 mg	100@ 21.61	PL
2676W	PROTEIN HYDROLYSATE FORMULA with MEDIUM CHAIN TRIGLYCERIDES	400 g	1@ 20.69	NT
8259Q		450 g	1@ 12.93	NU
2608G	PYRIDOSTIGMINE BROMIDE	180 mg (m.r.)	50@ 58.61	VT
8284B	RALTITREXED	2 mg	1@ 283.29	HH
1937Y	RANITIDINE HYDROCHLORIDE	150 mg (base), effervescent	30@ 5.74	GK
8162N		150 mg (base) per 10 mL, 300 mL	1@ 7.65	GK
8903N		150 mg (base), effervescent	30@ 7.37	GK
8905Q		150 mg (base) per 10 mL, 300 mL	1@ 8.75	GK
8780D	RISPERIDONE	25 mg	1@ 151.48	JC
8781E		37.5 mg	1@ 194.35	JC
8782F		50 mg	1@ 236.90	JC
8787L		0.5 mg	20@ 8.14	JC,TX
8788M		0.5 mg (orally disintegrating)	28@ 13.66	JC
8790P		1 mg (orally disintegrating)	28@ 26.15	JC
8792R		1 mg (orally disintegrating)	28@ 26.15	JC
8794W		2 mg (orally disintegrating)	28@ 52.53	JC

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
8869T		0.5 mg	20@ 8.14	JC,TX
8870W		0.5 mg (orally disintegrating)	28@ 13.66	JC
9075P		3 mg (orally disintegrating)	28@ 78.06	JC
9076Q		4 mg (orally disintegrating)	28@ 103.57	JC
9080X		2 mg (orally disintegrating)	28@ 52.53	JC
9313E	RIZATRIPTAN	10 mg (as benzoate)	2@ 9.35	MK
1099W	SALBUTAMOL SULFATE	200 mcg (base)	100@ 6.04	GK
1103C		2 mg (base) per 5 mL, 150 mL	1@ 7.89	GK
2000G		2.5 mg (base) in 2.5 mL, 30	1@ 6.26	AF,GX,SI,PF,CR,SZ,GM, GK
2001H		5 mg (base) in 2.5 mL, 30	1@ 6.61	AF,GX,SI,CR,SZ,GM,GK
2003K		5 mg (base) per mL, 30 mL	1@ 6.61	PF
8288F		100 mcg (base) per dose (200 doses)	1@ 4.40	AL,GK,IA
8354Q		100 mcg (base) per dose (200 doses)	1@ 16.09	IA
2995P	SALCATONIN	50 i.u. in 1 mL	5@ 33.54	NV
2997R		100 i.u. in 1 mL	5@ 51.57	NV
2014B	SODIUM ALGINATE with CALCIUM CARBONATE and SODIUM BICARBONATE	1 g-320 mg-534 mg in 20 mL, 500 mL	1@ 4.13	RC
2260Y	SODIUM CHLORIDE	513 mmol per L, 1 L	1@ 5.06	BX
2264E		154 mmol per L, 1 L	1@ 3.35	BR,PK,BX
9392H		77 mmol per 500 mL, 500 mL	1@ 2.33	BR,PK
9473N		38.5 mmol per 250 mL, 250 mL	1@ 3.52	BR,PK
5021G		77 mmol per 500 mL, 500 mL	1@ 2.33	BR,PK
5212H		154 mmol per L, 1 L	1@ 3.35	BR,PK,BX
5213J		513 mmol per L, 1 L	1@ 5.06	BX
2266G	SODIUM CHLORIDE COMPOUND	1 L	1@ 5.90	BX
2278X	SODIUM CHLORIDE with GLUCOSE	39 mmol-69 mmol per 500 mL, 500 mL	1@ 4.47	BX
2279Y		19 mmol-104 mmol per 500 mL, 500 mL	1@ 4.47	BX
2281C		31 mmol-222 mmol per L, 1 L	1@ 3.42	BX
5214K		31 mmol-222 mmol per L, 1 L	1@ 3.42	BX
5215L		19 mmol-104 mmol per 500 mL, 500 mL	1@ 4.47	BX
5216M		39 mmol-69 mmol per 500 mL, 500 mL	1@ 4.47	BX
2286H	SODIUM LACTATE COMPOUND	1 L	1@ 3.35	BR,PK,BX
9416N		500 mL	1@ 2.33	BR,PK
2289L	SODIUM VALPROATE	200 mg (e.c.)	100@ 13.18	AF,WA,SZ,SI,SW
2290M		500 mg (e.c.)	100@ 24.96	AF,WA,SZ,SI,SW
2293Q		200 mg per 5 mL, 300 mL	1@ 14.25	SW
2294R		100 mg	100@ 12.79	SW
2295T		200 mg per 5 mL, 300 mL	1@ 14.25	SW
9380Q	SORAFENIB	200 mg (as tosylate)	60@ 3225.33	BN
2091C	SORBITOL with SODIUM CITRATE and SODIUM LAURYL SULFOACETATE	3.125 g-450 mg-45 mg in 5 mL, 12	1@ 13.19	JT,AE
5331N		3.125 g-450 mg-45 mg in 5 mL, 12	1@ 13.19	JT,AE
5332P		3.125 g-450 mg-45 mg in 5 mL, 12	1@ 13.19	JT,AE
9448G	SOY LECITHIN	10 mg per mL, 10 mL	1@ 14.82	RB
5545W		10 mg per mL, 10 mL	1@ 14.82	RB
8577K	SOY PROTEIN and FAT FORMULA with VITAMINS and MINERALS—CARBOHYDRATE FREE	384 mL	1@ 5.53	AB
2093E	SULFASALAZINE	500 mg	100@ 22.33	PF
2096H		500 mg (e.c.)	100@ 24.35	KR,PF
9208P		500 mg	100@ 22.33	PF
9209Q		500 mg (e.c.)	100@ 24.35	KR,PF
2047R	SULINDAC	100 mg	50@ 4.96	AF
5381F		100 mg	50@ 4.96	AF
5383H		100 mg	50@ 4.96	AF
5217N		100 mg	50@ 4.96	AF
8144P	SUMATRIPTAN	50 mg (as succinate)	2@ 9.17	GK,AF,SI,TX,CH,TW
8885P		50 mg (as succinate) (fast disintegrating)	2@ 9.17	GK
2110C	TAMOXIFEN CITRATE	20 mg (base)	30@ 26.92	AP
2088X	TEMAZEPAM	10 mg	25@ 1.29	FM,AF,TX,SI
5375X		10 mg	25@ 1.29	FM,AF,TX,SI
5376Y		10 mg	25@ 1.29	FM,AF,TX,SI
8819E	TEMOZOLOMIDE	5 mg	5@ 67.21	SH
8820F		20 mg	5@ 187.17	SH
8821G		100 mg	5@ 794.29	SH
9361Q		140 mg	5@ 1086.56	SH
9160D	TERBINAFINE	10 mg per g, 15 g	1@ 15.47	NC
8098F	TESTOSTERONE	100 mg	1@ 33.86	SH
8099G		200 mg	1@ 67.71	SH
2101N	TESTOSTERONE ESTERS	250 mg	1@ 9.02	SH
2832C	TETRACOSACTRIN	1 mg in 1 mL	1@ 12.97	NV
2345K	THIOTEPA	15 mg	1@ 74.62	SI

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$		Manufacturer
8221Q	TIAGABINE HYDROCHLORIDE	5 mg (base)	50@	33.11	OA
8222R		10 mg (base)	50@	66.21	OA
8223T		15 mg (base)	50@	95.23	OA
1356J		80 mg (base) in 2 mL	5@	29.30	HH
8872Y	TOBRAMYCIN SULFATE	80 mg (base) in 2 mL (without preservative)	5@	29.30	PF
6072N			20@	7.44	CS,SI,SZ,AF,TX,CH,TW,ZP,GM
6073P	TRAMADOL HYDROCHLORIDE		20@	10.03	CS,SI,AF,TX,CH,TW,ZP,GM,SZ
6074Q			20@	12.21	CS,SI,SZ,AF,TX,CH,TW,ZP,GM
6096W	TRIAMCINOLONE ACETONIDE		1@	7.29	CS
7211M			20@	5.21	CS
7212N			10@	6.95	IA
7213P			10@	9.92	IA
7214Q			10@	13.37	IA
2117K		200 mcg per g, 100 g	1@	4.07	FM,SI
2118L	TRIGLYCERIDES, LONG CHAIN with GLUCOSE POLYMER	200 mcg per g, 100 g	1@	4.07	FM,SI
9308X		250 mL, 18	1@	55.56	VF
9309Y		1 L, 6	1@	74.40	VF
3128P	TRIGLYCERIDES, MEDIUM CHAIN	500 mL	1@	22.98	SB
9327X		250 mL	1@	26.00	SB
3136C	TRIGLYCERIDES, MEDIUM CHAIN and LONG CHAIN with GLUCOSE POLYMER	400 g	1@	36.14	SB
8478F	TRIGLYCERIDES—MEDIUM CHAIN, FORMULA	400 g	1@	51.86	SB
8629E		420 g	1@	57.63	SB
9383W		16 g, 30	1@	61.80	VF
9165J	TYROSINE with CARBOHYDRATE	4 g containing 1 g tyrosine, 30	1@	127.40	VF
8448P	URSODEOXYCHOLIC ACID	250 mg	100@	183.09	OA
8133C	VALACICLOVIR HYDROCHLORIDE	500 mg (base)	10@	49.68	GK
9135T	VALINE with CARBOHYDRATE	4 g containing 50 mg valine, 30	1@	127.40	VF
9434M		4 g containing 1 g valine, 30	1@	140.14	VF
2270L	VANCOMYCIN	1 g	1@	11.53	HH,SZ
3113W		125 mg	20@	115.04	AS
3114X		250 mg	20@	221.06	AS
3130R		500 mg	1@	5.77	HH,AS,SZ
3131T		500 mg	1@	5.77	HH,AS,SZ
3323X		500 mg	1@	5.77	HH,AS,SZ
9129L	VARENICLINE	1 mg (as tartrate)	56@	113.32	PF
2374Y	VINCISTINE SULFATE	1 mg in 1 mL	5@	74.39	HH,PF
8280T	VINORELBINE TARTRATE	10 mg (base) in 1 mL	1@	70.67	HH,FB,IT,PK
8281W		50 mg (base) in 5 mL	1@	294.90	HH,FB,IT,PK
9009E		20 mg (base)	1@	100.27	FB
9010F		30 mg (base)	1@	148.74	FB
9328Y	VITAMINS, MINERALS and TRACE ELEMENTS with CARBOHYDRATE	200 g	1@	64.00	SB
9382T	WHEY PROTEIN FORMULA supplemented with AMINO ACIDS, LONG CHAIN POLYUNSATURATED FATTY ACIDS, VITAMINS and MINERALS, and low in PROTEIN, PHOSPHATE, POTASSIUM and LACTOSE	100 g, 10	1@	164.35	VF
8587Y	WHEY PROTEIN FORMULA supplemented with AMINO ACIDS, VITAMINS and MINERALS, and low in PROTEIN, PHOSPHATE, POTASSIUM and LACTOSE	400 g	1@	66.22	SB
8266C	ZOLMITRIPTAN	2.5 mg	2@	9.71	AP
9736K		2.5 mg	2@	11.09	AP

Section 4

Drug Tariff

Container Prices

Standard Formulae Preparations

Table of Codes, Maximum Quantities, and Number of Repeats for Extemporaneously Prepared Pharmaceutical Benefits

Drug Tariff

Drug	Standard	Recovery Prices			
		0.1 g/mL	1 g/mL	10 g/mL	100 g/mL
		\$	\$	\$	\$
Acacia Mucilage (by weight)	APF 15	0.01	0.08	0.64	5.67
Acacia, powdered	BP	0.02	0.13	1.06	9.45
Acetic Acid (6 per cent)	BP	0.01	0.01	0.11	1.02
Acetic Acid (33 per cent)	BP	0.01	0.05	0.38	3.38
Acetone (use as additive only)	BP	0.01	0.09	0.71	6.33
Alum	BP	0.02	0.16	1.29	11.43
Aluminium Acetate Solution	BP	0.03	0.21	1.67	14.83
Anise Water Concentrated 1 in 40 (use as additive only)	BP	0.01	0.05	0.39	3.49
Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.01	0.07	0.59	5.23
Ascorbic Acid (for use only as an ingredient of ferrous sulfate mixtures)	BP	0.08	0.61	4.85	43.15
Aspirin	BP	0.06	0.49	3.88	34.50
Belladonna Tincture	BP	0.08	0.62	4.92	43.70
Benzocaine	BP	0.06	0.51	4.08	36.25
Benzoic Acid	BP	0.04	0.32	2.52	22.39
Benzoic Acid Compound Ointment	APF	0.02	0.16	1.31	11.61
Benzoic Acid Solution	BP	0.01	0.09	0.74	6.62
Benzoin Compound Tincture	BP	0.05	0.36	2.87	25.48
Boric Acid (use as additive only)	BP	0.01	0.05	0.39	3.45
Boric Acid, Olive Oil and Zinc Oxide Ointment	QHF	0.01	0.07	0.56	4.94
Calcium Hydroxide	BP	0.08	0.67	5.34	47.50
Calcium Hydroxide Solution	BP	0.01	0.01	0.07	0.66
Castor Oil (use as additive only)	BP	0.01	0.04	0.35	3.10
Cetomacrogol Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.01	0.03	0.24	2.10
Cetrimide Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.05	0.39	3.11	27.67
Chlorhexidine Acetate (use as additive only)	BP	0.53	4.21	33.65	299.12
Chlorhexidine Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.06	0.48	3.81	33.86
Chloroform (use as additive only)	BP	0.07	0.59	4.69	41.67
Chloroform Spirit	BP	0.01	0.03	0.25	2.26

Drug	Standard	Recovery Prices			
		0.1 g/mL	1 g/mL	10 g/mL	100 g/mL
		\$	\$	\$	\$
Chloroform Water Concentrated 1 in 40	APF 15	0.01	0.03	0.25	2.26
Citric Acid Monohydrate	BP	0.01	0.04	0.30	2.65
Coal Tar	BP	0.12	0.98	7.83	69.60
Coal Tar Solution	BP	0.02	0.12	0.95	8.44
Cocaine Hydrochloride	BP	6.36	50.86	406.90	3616.86
Coconut Oil	BP	0.01	0.10	0.80	7.16
Codeine Linctus	APF	0.01	0.06	0.45	4.01
Codeine Phosphate (may only be prescribed in linctuses, mixtures or mixtures for children)	BP	1.49	11.91	95.31	847.16
Collodion Flexible	BP	0.15	1.23	9.86	37.60
Dithranol	BP	3.97	31.78	254.22	2259.76
Emulsifying Ointment (for use only as a base combined with active ingredients)	BP	0.01	0.07	0.55	4.92
Ephedrine Hydrochloride (may only be prescribed in nasal instillations)	BP	0.89	7.09	56.70	504.00
Ethanol (90 per cent) (use as additive only)	BP	0.01	0.03	0.21	1.88
Ethanol (96 per cent) (use as additive only)	BP	0.01	0.03	0.25	2.25
Ether Solvent (use as additive only)	BP	0.01	0.11	0.90	8.01
Eucalyptus Oil (use as additive only)	BP	0.02	0.12	0.97	8.63
Ferrous Sulfate	BP	0.11	0.91	7.31	65.00
Formaldehyde Solution	BP	0.10	0.80	6.39	56.79
Gentian Alkaline Mixture	APF	0.01	0.06	0.51	4.56
Glycerol	BP	0.01	0.06	0.45	3.96
Honey Purified (use as additive only)	BP 1993	0.01	0.02	0.15	1.35
Hydroxybenzoate Compound Solution	APF	0.13	1.06	8.50	75.55
Iodine	BP	0.27	2.12	16.92	150.37
Iodine Alcoholic Solution	BP	0.02	0.15	1.20	10.65
Iodine Aqueous Oral Solution	BP	0.03	0.21	1.67	14.82
Kaolin Mixture	BPC 1968	0.01	0.07	0.54	4.83
Kaolin and Opium Mixture	APF 14	0.01	0.09	0.69	6.10
Lactic Acid	BP	0.06	0.47	3.72	33.07
Lavender Spike Oil	BPC 1968	0.09	0.68	5.44	48.35
Liquorice Liquid Extract	BP	0.02	0.12	0.95	8.43

Drug	Standard	Recovery Prices			
		0.1 g/mL	1 g/mL	10 g/mL	100 g/mL
		\$	\$	\$	\$
Magnesium Carbonate Light	BP	0.03	0.24	1.89	16.79
Magnesium Sulfate (may only be prescribed for other than oral use)	BP	0.01	0.02	0.12	1.10
Magnesium Trisilicate	BP	0.04	0.32	2.59	23.03
Menthol, Racemic or Levomenthol	BP	0.23	1.83	14.60	129.76
Methyl Hydroxybenzoate	BP	0.30	2.40	19.22	170.84
Methyl Hydroxybenzoate Solution	APF	0.03	0.23	1.81	16.09
Methylated Industrial Spirit (use as additive only)	BP	0.01	0.04	0.32	2.86
Olive Oil (use as additive only)	BP	0.02	0.12	0.95	8.41
Paraffin Hard	BP	0.02	0.19	1.52	13.50
Paraffin Liquid (may only be prescribed for other than oral use)	BP	0.01	0.04	0.34	3.06
Paraffin Light Liquid	BP	0.02	0.15	1.20	10.69
Paraffin Soft White	BP	0.01	0.05	0.41	3.69
Paraffin Soft Yellow	BP	0.02	0.19	1.51	13.43
Peppermint Oil (use as additive only)	BP	0.13	1.01	8.08	71.84
Peppermint Water Concentrated 1 in 40 (use as additive only)	APF 16	0.02	0.18	1.40	12.43
Phenobarbitone Sodium (may only be prescribed for the treatment of epilepsy)	BP	10.67	85.38	683.00	6071.11
Phenol Liquefied (not available for ear drops)	BP	0.17	1.37	10.94	97.20
Podophyllum Resin	BP	0.95	7.61	60.90	541.33
Potassium Citrate	BP	0.01	0.10	0.79	7.02
Potassium Iodide	BP	0.11	0.87	6.92	61.51
Potassium Permanganate	BP	0.01	0.10	0.82	7.30
Propyl Hydroxybenzoate	BP	0.25	2.02	16.12	143.29
Propylene Glycol	BP	0.01	0.06	0.48	4.31
Red Syrup	APF 15	0.02	0.13	1.02	9.05
Resorcinol	BP	0.19	1.49	11.90	105.78
Salicylic Acid	BP	0.03	0.21	1.67	14.87
Salicylic Acid Ointment	APF	0.01	0.11	0.87	7.78
Salicylic Acid Ointment	BP	0.01	0.11	0.87	7.78
Simple Ointment (white) (for use only as a base combined with active ingredients)	BP	0.02	0.14	1.14	10.12
Simple Ointment (yellow) (for use only as a base combined with active ingredients)	BP	0.02	0.16	1.25	11.10

Drug	Standard	Recovery Prices			
		0.1 g/mL	1 g/mL	10 g/mL	100 g/mL
		\$	\$	\$	\$
Sodium Bicarbonate	BP	0.01	0.08	0.60	5.31
Sodium Chloride	BP	0.02	0.13	1.03	9.12
Sodium Chloride Solution	BP	0.01	0.01	0.07	0.61
Sodium Citrate	BP	0.02	0.15	1.16	10.32
Sodium Thiosulfate (use as additive only)	BP	0.03	0.21	1.67	14.87
Starch	BP	0.02	0.14	1.11	9.89
Sulfur Ointment (for use only as a base combined with active ingredients)	BP 1980	0.02	0.15	1.19	10.55
Sulfur Precipitated	BP 1980	0.03	0.22	1.72	15.32
Syrup	BP	0.01	0.05	0.43	3.81
Talc Purified, sterilised	BP	0.02	0.15	1.19	10.56
Thymol	BP	0.23	1.81	14.50	128.89
Thymol Compound Mouth Wash	APF 15	0.01	0.04	0.35	3.12
Tragacanth Compound Powder	BP 1980	0.04	0.29	2.30	20.47
Tragacanth Mucilage	APF 13	0.01	0.03	0.24	2.10
Tragacanth Mucilage	BPC 1973	0.01	0.02	0.17	1.55
Tragacanth, powdered	BP	0.12	0.95	7.59	67.50
Trichloroacetic Acid	BP 1980	0.31	2.47	19.74	175.50
Triethanolamine	BP	0.05	0.41	3.26	29.00
Water For Injections, sterilised (b) (extemporaneously prepared eye drops and eye lotions)	BP	0	0	0	7.37
Water Purified	BP	0.01	0.01	0.05	0.49
Wool Alcohols Ointment (white) (for use only as a base combined with active ingredients)	BP	0.02	0.14	1.15	10.19
Wool Alcohols Ointment (yellow) (for use only as a base combined with active ingredients)	BP	0.02	0.14	1.15	10.19
Wool Fat	BP	0.02	0.17	1.37	12.14
Wool Fat Hydrous	BP	0.02	0.13	1.05	9.36
Zinc Compound Paste	BP	0.03	0.22	1.74	15.43
Zinc Cream (for use only as a base combined with active ingredients)	BP	0.01	0.07	0.55	4.86
Zinc Oxide	BP	0.01	0.11	0.84	7.45
Zinc and Salicylic Acid Paste	BP	0.02	0.16	1.24	11.06
Zinc Sulfate	BP	0.03	0.20	1.61	14.27

Container Prices

Container Prices

\$

DISPENSING BOTTLES –

25mL	0.55
50mL	0.56
100mL	0.57
200mL	0.84
500mL	1.34

POISON BOTTLES –

25mL	0.52
50mL	0.51
100mL	0.55
200mL	0.83
500mL	1.34

SCREW CAP JARS –

25g	0.64
50g	0.70
100g	0.84
200g	0.89
500g	1.29

DROPPER CONTAINERS –

15mL polythene	0.84
15mL glass	0.86

Dispensing Fee for Extemporaneously Prepared Benefits 8.46

Additional Fee for Agreed Price 1.41

Extemporaneously Prepared Benefits

Standard Formula Preparations

The following list is not intended to indicate in any way which particular formula an approved pharmacist should use in filling a prescription.

The prices shown in the column 'Dispensed Price for Max. Qty' are for the ingredients, the container and the dispensing fee. The prices shown in the column 'Maximum Recordable Value for Safety Net' are for the ingredients, the container and the dispensing fee and, where applicable, the additional fee for agreed price benefits.

KEY TO REFERENCES:

APF Australian Pharmaceutical Formulary

BP British Pharmacopoeia

BPC British Pharmaceutical Codex

QHF Queensland Hospital Formulary

Standard Formula Preparations

Code	Item	Reference	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net
			\$	\$
CREAMS (Maximum Quantity 100 g and 1 Repeat)				
7502W	Salicylic Acid and Sulfur Aqueous	APF	15.17	16.58
DUSTING POWDERS (Maximum Quantity 100 g and 1 Repeat)				
7458M	Zinc, Starch and Talc	APF 15 & BPC 1973	20.17	21.58
EAR DROPS (Maximum Quantity 15 mL and 2 Repeats)				
7642F	Aluminium Acetate	APF	10.97	12.38
7643G	Aluminium Acetate	BP	11.80	13.21
7314Y	Sodium Bicarbonate	APF & BP	9.66	11.07
7313X	Spirit	APF	10.91	12.32
INHALATIONS (Maximum Quantity 50 mL and 1 Repeat)				
7484X	Benzoin and Menthol	APF	25.13	26.54
7308P	Menthol	APF	11.86	13.27
7310R	Menthol and Eucalyptus	BP 1980	12.36	13.77
LINCTUSES CONTAINING CODEINE PHOSPHATE (Maximum Quantity 100 mL and 0 Repeats)				
7530H	Codeine	APF	13.04	14.45
LOTIONS (Maximum Quantity 200 mL and 2 Repeats)				
7709R	Aluminium Acetate Aqueous	APF	11.87	13.28
MIXTURES, OTHER (Maximum Quantity 200 mL and 4 Repeats)				
7604F	Gentian Alkaline	APF	18.41	19.82
7348R	Kaolin	BPC 1968	18.95	20.36
7301G	Kaolin and Opium	APF 14	21.49	22.90
7342K	Magnesium Trisilicate	BPC 1968	16.20	17.61
7343L	Magnesium Trisilicate and Belladonna	BPC 1968	21.01	22.42
MOUTH WASHES (Maximum Quantity 200 mL and 1 Repeat)				

Code	Item	Reference	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$
7457L	Thymol Compound	APF 15	15.53	16.94
	OINTMENTS (Maximum Quantity 100 g and 1 Repeat)			
7914M	Benzoic Acid Compound	APF	20.91	22.32
7914M	Benzoic Acid Compound (extemporaneous formula)	BP	20.91	22.32
7902X	Boric Acid, Olive Oil and Zinc Oxide	QHF	14.24	15.65
7926E	Salicylic Acid	APF	17.08	18.49
7928G	Salicylic Acid (extemporaneous formula)	BP	17.08	18.49
	PAINTS (Maximum Quantity 25 mL and 1 Repeat)			
7567G	Podophyllin Compound	APF 16 & BP	44.70	34.20
7568H	Salicylic Acid	APF	34.14	34.20
	PASTES, OTHER (Maximum Quantity 100 g and 1 Repeat)			
7558T	Zinc	APF	24.73	26.14
7558T	Zinc Compound (extemporaneous formula)	BP	24.73	26.14
	POWDER FOR INTERNAL USE (Maximum Quantity 100 g and 2 Repeats)			
7545D	Magnesium Trisilicate	BP	32.38	33.79

—CONTAINER RATES ARE INCLUDED—

**Table of Codes, Maximum Quantities, and Number of Repeats for
Extemporaneously Prepared Benefits**

Code	Preparation	Maximum Quantity	Number of Repeats
13Q	Creams	100 g	1
48M	Dusting Powders	100 g	1
15T	Ear Drops	15 mL	2
19B	Eye Drops containing Cocaine Hydrochloride	15 mL	..
22E	Eye Drops, Other	15 mL	5
23F	Eye Lotions	200 mL	2
29M	Inhalations	50 mL	1
64J	Linctuses containing Codeine Phosphate	100 mL	..
34T	Linctuses, Other	100 mL	2
39C	Lotions	200 mL	2
65K	Mixtures containing Codeine Phosphate	200 mL	..
40D	Mixtures, Other	200 mL	4
66L	Mixtures for Children containing Codeine Phosphate	100 mL	..
41E	Mixtures for Children, Other	100 mL	4
30N	Mouth Washes	200 mL	1
42F	Nasal Instillations	15 mL	2
43G	Ointments, Waxes	100 g	1
44H	Paints	25 mL	1
63H	Pastes containing Cocaine Hydrochloride	25 g	..
45J	Pastes, Other	100 g	1
49N	Powders for Internal Use	100 g	2
52R	Solutions	200 mL	2

Special Note: Purified Water BP is the minimum requirement for water in all PBS extemporaneous preparations.



Australian Government

Department of Health and Ageing

REPATRIATION SCHEDULE OF PHARMACEUTICAL BENEFITS

1 January 2011


The benefits listed in this Schedule may only be prescribed to Department of Veterans' Affairs beneficiaries holding a:

- Repatriation Health Card For All Conditions (gold); or
- Repatriation Health Card For Specific Conditions (white); or
- Repatriation Pharmaceutical Benefits Card (orange);

BENEFICIARIES' ENTITLEMENT CARDS AND ELIGIBILITY

The diagram below outlines the drug eligibility of Department of Veterans' Affairs

REPATRIATION PHARMACEUTICAL BENEFITS



FOR REPATRIATION PHARMACEUTICAL BENEFITS

beneficiaries in accordance with their treatment entitlement.

PBS Schedule listings written on the common PBS/RPBS prescription form, ticked as 'RPBS' (restrictions apply)	+	Repatriation Schedule listings written on the common PBS/RPBS prescription form, ticked as 'RPBS'	+	Items for which prescribing approval has been authorised on the common PBS/RPBS authority prescription form for: • PBS and Repatriation Schedules 'Authority required' items; • greater quantities/repeats of drugs listed in PBS and Repatriation Schedules; or • items not listed in PBS or Repatriation Schedules.

Only for disabilities that have been accepted for treatment by the Department of Veterans' Affairs

PBS Schedule listings written on the common PBS/RPBS prescription form, ticked as 'RPBS' (restrictions apply)	+	Repatriation Schedule listings written on the common PBS/RPBS prescription form, ticked as 'RPBS'	+	Items for which prescribing approval has been authorised on a common PBS/RPBS authority prescription form for: • PBS and Repatriation Schedules 'Authority required' items; • greater quantities/repeats of drugs listed in PBS and Repatriation Schedules; or • items not listed in PBS or Repatriation Schedules.

RPBS Explanatory Notes

Introduction

The Australian Repatriation System

- The Australian Repatriation system is based primarily on the principle of compensation to veterans and eligible dependants for injury or death related to war service. In certain cases, treatment is also provided for accepted injuries or conditions that are not service-related or have occurred as a result of other than war service.
- Through the *Veterans' Entitlements Act 1986* the Department of Veterans' Affairs provides programs of compensation, income support and treatment for eligible veterans and their dependants. One of the defined benefits for eligible veterans is the Repatriation Pharmaceutical Benefits Scheme. This range of medications and dressings is more comprehensive than is available through the Pharmaceutical Benefits Scheme.

RPBS prescribing provisions

- Unless otherwise stated, Repatriation Pharmaceutical Benefits Scheme (RPBS) prescriptions must conform with the requirements of Pharmaceutical Benefits Scheme (PBS) prescriptions, as detailed in Section 1 – Explanatory Notes in the *Schedule of Pharmaceutical Benefits* book. The prescriber shall ensure that a prescription contains the following details:
 - the category of benefit, i.e., RPBS, by placing a cross in the relevant box;
 - the patient's full name and address;
 - the prescription date;
 - the DVA file number of the patient as evidence of entitlement;
 - in the case of authority prescriptions, the Authority approval number or the four digit streamlined authority code;
 - the item, form, strength, quantity and directions;
 - the number of repeats, if applicable;
 - indicate when brand substitution is not permitted; and
 - the name, signature, the prescriber number and address of the prescriber.

Prior Approval Arrangements

- The prior approval of the Department is required to prescribe the following:
 - 'Authority required' items (excluding 'Authority required (STREAMLINED)' items) listed in either the PBS or RPBS Schedule;
 - increased quantities and/or repeats of items listed in either the PBS or RPBS Schedule;
 - items listed under section 100 of the *National Health Act 1953*; and
 - other items not listed in either Schedule (non-Schedule items).
- The above items are to be prescribed on the common PBS/RPBS authority prescription form in accordance with the directions stated in the Explanatory Notes in the *Schedule of Pharmaceutical Benefits* (See also information regarding dental prescribing and prescribing by optometrists under the RPBS in these Notes.)
- All Authority required prescriptions and requests for non-Schedule items must receive prior approval from the Department. This can be achieved by either:
 - using the Department's national free call number 1800 552 580; or
 - by mailing the written authority prescription to the Veterans' Affairs Pharmaceutical Advisory Centre (VAPAC) at the reply paid address shown at the end of these RPBS Explanatory Notes.

Prior approval is not required from DVA to prescribe an Authority required (STREAMLINED) item (except where increased quantities and/or repeats are required). Instead the authority prescription form must include a four digit streamlined authority code.

- Some requests for prior approval (including some non-Schedule items) need to be referred by VAPAC to the Repatriation Pharmaceutical Reference Committee for consideration. In such cases a VAPAC pharmacist will advise the prescriber to submit a request in writing that provides the following information:

- A current clinical report on the patient's condition (such as age, co-morbidities, renal, liver failure) and clinical reports including pathology, biochemistry, diagnostic and other investigations if appropriate.
 - Details of past and current therapy for the condition. Include details of PBS, RPBS and non-Schedule items utilised, and the results of those therapies.
 - Details of the proposed treatment regimen. Include intended dose and duration of treatment and objective measures of response.
 - When the proposed use of the item is outside the TGA-approved indications for use in Australia, provide copies of articles from peer reviewed publications supporting the proposed treatment.
 - Signed, informed patient consent where the item is to be used for a non-TGA-approved indication.
 - For items without Australian marketing approval, a copy of the TGA Special Access Scheme approval to prescribe the drug.
- Requests for prior approval to prescribe a non-Schedule (PBS or RPBS) item that is of the same therapeutic class (ATC level 3) as an item that is listed on the Schedule, will not be approved unless unequivocal clinical evidence is presented to demonstrate that the requested item is essential for effective treatment of the nominated patient.
 - A pharmacist should not supply an item prescribed on an RPBS Authority Prescription Form unless the form has been approved and stamped by VAPAC, or has been endorsed by the prescriber with a telephone Authority approval number provided by VAPAC. Medicare Australia will not accept RPBS Authority prescriptions that have not been approved by the Department of Veterans' Affairs for payment.

Palliative Care Drugs

- The following medications may be available, or made available in increased quantities or doses under prior approval arrangements for use only in the palliative care of terminal disease:
 - clonazepam
 - cyclizine
 - dexamethasone
 - disodium pamidronate
 - fentanyl
 - glycopyrrolate
 - hyoscine butylbromide
 - hyoscine hydrobromide
 - ketamine
 - midazolam
 - octreotide
- For further information telephone VAPAC on 1800 552 580.

Dental Prescribing

- Under Department of Veterans' Affairs arrangements, financial responsibility for pharmaceutical benefits prescribed by a Local Dental Officer (LDO) is limited to treatment to which holders of the following cards are entitled:
 - a Gold Repatriation Health Card – For All Conditions; or
 - a White Repatriation Health Card – For Specific Conditions; or
 - an Orange Repatriation Pharmaceutical Benefits Card.
- Where possible the LDO shall prescribe in accordance with the provisions governing dental prescribing under the Pharmaceutical Benefits Scheme (PBS).
- Prescriptions for PBS Dental Schedule items for Gold, White and Orange Card holders are to be dispensed at the PBS concessional rate. Claims for payment by the dispensing pharmacist are to be included with other Repatriation prescriptions. The card holder is required to meet the cost of any applicable brand premium.
- When a non-PBS Dental Schedule item is prescribed for an eligible card holder, the LDO's private prescription form should be used. The dispensing pharmacist may charge the patient the full cost of the prescription. The patient may claim a refund for the full cost of a non-Schedule item from the Department if an itemised receipt (not a cash register receipt) and a copy of the prescription are provided.

Prescribing by optometrists

- Optometrists approved as 'PBS prescribers' may write RPBS prescriptions as outlined in Section 1 for medicines listed in Section 2 of the PBS Schedule as pharmaceutical benefits for optometrical use.
- Medicines in the optometrist list include non-Authority and Authority required items. Procedures for obtaining VAPAC approval to prescribe 'Authority required' optometrist items or increased quantities and/or repeats of optometrist items under the RPBS are the same as indicated under prior approval arrangements above.
- The list of medicines for prescribing by optometrists under the RPBS is the same as applies under the PBS. There are no optometrist listings in the RPBS Schedule for prescribing for veterans only. There is no provision for optometrist prescribers to request approval to prescribe items that are not included in the PBS optometrist list (non-Schedule items).
- Optometrist PBS/RPBS prescription forms are for use for prescribing non-Authority or Authority required optometrist items under the RPBS with one item per form only.

Provisions governing pricing and payment for RPBS benefits

Introduction

- Unless otherwise stated, the pricing and payment principles and arrangements for approved pharmacists supplying pharmaceutical benefits under the RPBS will be the same as those arrangements applying under the PBS.
- Where a pharmaceutical benefit that is not listed on the PBS or RPBS Schedule is dispensed on an RPBS Authority prescription, a pharmacist will price the benefit and enter the serial number, prescription identifying number and price on the sticker or stamp imprint affixed to the prescription.

Pricing of Schedule Items

- Items supplied under the RPBS from the PBS Schedule, both ready-prepared and extemporaneously-prepared, will be paid on the same basis as benefits supplied under the PBS. Items supplied under the RPBS from the Repatriation Schedule, including wound dressings, will be paid on the basis of the price as given in the Repatriation Pharmaceutical Benefits section (Section 1 – RPBS Schedule, Drugs, Medicines and Dressings) of the *Schedule of Pharmaceutical Benefits*.

Pricing of Non-Schedule Ready Prepared Items

- Non-Schedule ready-prepared items are to be priced on the basis of the invoiced, GST-exclusive wholesale price to pharmacists plus the appropriate PBS mark-up and the PBS dispensing fee. Where the item price to pharmacists is greater than \$100.00, a copy of the invoice pertaining to the supply of that item is to be submitted together with the appropriate copy of the authority prescription as part of the claim for payment.

Pricing of Non-Schedule Extemporaneously Prepared Items

- When an ingredient drug is not listed in the PBS Drug Tariff, the recovery price will be based on the invoiced wholesale price to pharmacists, increased by a mark-up of 100%, calculated in accordance with the directions contained in the pricing instructions for pricing of PBS extemporaneously-prepared benefits in this Schedule. The price paid by the pharmacist for the commercial pack from which the ingredient is used shall be endorsed on the prescription form.

Miscellaneous Pricing Rules

- The price to pharmacists used as the basis of pricing will be the invoiced, GST-exclusive price from the wholesaler.
- If multiple quantities of a manufacturer's original pack are supplied, the PBS mark-up is applied to the price to pharmacist of each pack and then totalled. The PBS dispensing fee, and the PBS dangerous drug fee if applicable, are then added to the total of the marked-up prices.
- When the quantity prescribed corresponds with the quantity of a manufacturer's original pack, in no circumstances will the price payable for one pack exceed that payable for multiples or combinations of packs to supply the quantity prescribed.
- The list of ingredient drugs and prices included in the PBS Drug Tariff are common to both the PBS and RPBS. Certain restrictions apply regarding the prescribing and dispensing of some of these ingredient drugs as pharmaceutical benefits, e.g., use as additive only.
- For items prescribed generically, including non-Schedule and wound dressings, the pharmacist should indicate on the prescription the quantity and brand supplied. If prescriptions are not endorsed, the Department will pay the lowest priced acceptable product available.

General

Packaging Material, Postage or Freight

- Payment to a pharmacist for the costs of packaging materials, postage or freight required to supply a pharmaceutical benefit is to be paid by the patient, who may then claim reimbursement from the Department through the provision of a pharmacist's itemised receipt.

Payment for Items Supplied at Short Intervals

- For all items dispensed at specific short intervals of time, the Department will pay a separate PBS dispensing fee for each occasion that the drug is supplied and which is acknowledged on receipt by the patient or agent.
- The price payable on the items supplied will be based on the individual dose quantity supplied. Where applicable, a PBS dangerous drug fee and a minimum container charge will be payable for each supply.

Receipts for Patient Charges

- Where a charge is paid by a patient in any of the circumstances of paragraphs 13 or 24, the pharmacist is required to provide a printed receipt to the patient with the details of the items or services provided, the amount paid, date of supply and the patient's name and address. The patient may apply for reimbursement from the Department.

Special Patient Contributions

- The Special Patient Contribution for items listed as Special Pharmaceutical Benefits in the PBS Schedule is not payable by veterans entitled to pharmaceutical benefits under the RPBS. Eligible veterans receiving Special Pharmaceutical Benefits under the RPBS are required to pay only the concessional patient contribution and any applicable brand premium. If a Safety Net Entitlement card is held, the veteran should receive a Special Pharmaceutical Benefit free of charge, subject to any brand premium applicable. Medicare Australia will reimburse the dispensing pharmacist the total dispensed price, less the concessional patient contribution and/or brand premium if applicable.

Therapeutic Group Premiums — Authority Processing

- Items attracting a therapeutic group premium are dual listed. Dispensing pharmacists are therefore required to select the appropriate code for those items that are dual listed as authority and non-authority items, in order to correctly charge the patient and claim from Medicare Australia. Those authority prescriptions that grant exemption from a therapeutic group premium will have the letters 'TPX' at the beginning of the telephone Authority approval number, or, in the case of a written approval, will be stamped with the words "This prescription does not attract a therapeutic group premium".

DEPARTMENT OF VETERANS' AFFAIRS

Authority Prescription Applications

Applications for authority to prescribe under the Repatriation Pharmaceutical Benefits Scheme (RPBS) should be sent to the Veterans' Affairs Pharmaceutical Advisory Centre (VAPAC) using the free postal service:

REPLY PAID 9998

VAPAC (Veterans' Affairs Pharmaceutical Advisory Centre)

Department of Veterans' Affairs

GPO Box 9998

BRISBANE QLD 4001

For RPBS enquiries and telephone approvals 24 hours a day the Freecall number is:

1800 552 580

Departmental pharmacists answer applications for prior approval for non-Schedule items and Authority application calls.

REPATRIATION PHARMACETICAL BENEFITS

These changes to the Schedule of Pharmaceutical Benefits are effective from 1 January 2011. The Schedule is updated on the first day of each month and is available on the Internet at www.pbs.gov.au.

SUMMARY OF CHANGES

DELETION

Deletion – Brands

4657D **BANDAGE—COMPRESSION**, Bandage, high stretch, 10 cm x 3.5 m (*Eloflex 2480*)

ALTERATIONS

Alterations — Proprietary Name

From:

4727T **BANDAGE—RETENTION—COTTON CREPE**, Bandage 5 cm x 2.3 m (*Elastocrepe 36102520*)

To:

4727T **BANDAGE—RETENTION—COTTON CREPE**, Bandage 5 cm x 2.3 m (*Tensocrepe 36300501*)

From:

4728W **BANDAGE—RETENTION—COTTON CREPE**, Bandage 7.5 cm x 2.3 m (*Elastocrepe 36102420*)

To:

4728W **BANDAGE—RETENTION—COTTON CREPE**, Bandage 7.5 cm x 2.3 m (*Tensocrepe 36307501*)

From:

4729X **BANDAGE—RETENTION—COTTON CREPE**, Bandage 10 cm x 2.3 m (*Elastocrepe 36102320*)

To:

4729X **BANDAGE—RETENTION—COTTON CREPE**, Bandage 10 cm x 2.3 m (*Tensocrepe 36301001*)

Therapeutic Index for RPBS Schedule

Alimentary tract and metabolism 837*Stomatological preparations 837*

Stomatological preparations 837

Drugs for acid related disorders 837

Antacids 837

Drugs for functional gastrointestinal disorders 837

Drugs for functional bowel disorders 837

Belladonna and derivatives, plain 837

Laxatives 837

Laxatives 837

Antiobesity preparations, excl. diet products 838

Antiobesity preparations, excl. diet products 838

*Vitamins 839*Vitamin B₁, plain and in combination with vitamin B₆ and vitamin B₁₂ 839

Vitamin B-complex, incl. combinations 839

Mineral supplements 839

Calcium 839

Other mineral supplements 839

Blood and blood forming organs 840*Antithrombotic agents 840*

Antithrombotic agents 840

Blood substitutes and perfusion solutions 840

Irrigating solutions 840

Cardiovascular system 841*Vasoprotectives 841*

Agents for treatment of hemorrhoids and anal fissures for topical use 841

Dermatologicals 842*Antifungals for dermatological use 842*

Antifungals for topical use 842

Antifungals for systemic use 843

Emollients and protectives 843

Emollients and protectives 843

Protectives against UV-radiation 843

Antipruritics, incl. antihistamines, anesthetics, etc. 844

Antipruritics, incl. antihistamines, anesthetics, etc. 844

Antipsoriatics 844

Antipsoriatics for topical use 844

Antibiotics and chemotherapeutics for dermatological use 844

Antibiotics for topical use 844

Chemotherapeutics for topical use 844

Corticosteroids, dermatological preparations 845

Corticosteroids, plain 845

Corticosteroids, combinations with antibiotics 845

Antiseptics and disinfectants 845

Antiseptics and disinfectants 845

Other dermatological preparations 845

Other dermatological preparations 845

Genito urinary system and sex hormones 847*Gynecological antiinfectives and antiseptics 847*

Antiinfectives and antiseptics, excl. comb. with corticosteroids 847

Other gynecologicals 847

Other gynecologicals 847

Sex hormones and modulators of the genital system 847

Estrogens 847

Urologicals 847

Other urologicals, incl. antispasmodics 847

Drugs used in benign prostatic hypertrophy 848

Antiinfectives for systemic use 849*Antibacterials for systemic use 849*

Macrolides, lincosamides and streptogramins 849

Antineoplastic and immunomodulating agents 850*Antineoplastic agents 850*

Antimetabolites 850

Immunosuppressants 850

Immunosuppressants 850

Musculo-skeletal system 852*Antiinflammatory and antirheumatic products 852*

Antiinflammatory and antirheumatic products, non-steroids 852

Topical products for joint and muscular pain 852

Topical products for joint and muscular pain 852

Drugs for treatment of bone diseases 852

Drugs affecting bone structure and mineralization 852

Nervous system 853*Analgesics 853*

Opioids 853

Other analgesics and antipyretics 853

Psycholeptics 854

Anxiolytics 854

Hypnotics and sedatives 854

Other nervous system drugs 855

Drugs used in addictive disorders 855

Antiparasitic products, insecticides and repellents 856*Anthelmintics 856*

Antinematodal agents 856

Respiratory system 857*Nasal preparations 857*

Decongestants and other nasal preparations for topical use 857

Nasal decongestants for systemic use 857

Cough and cold preparations 857

Expectorants, excl. combinations with cough suppressants 857

Cough suppressants, excl. combinations with expectorants 858

Antihistamines for systemic use 858

Antihistamines for systemic use 858

Sensory organs 859*Ophthalmologicals 859*

Decongestants and antiallergics 859

Otologicals 859

Other otologicals 859

Various 860*All other therapeutic products 860*

All other therapeutic products 860

All other non-therapeutic products 863

All other non-therapeutic products 863

Section 1

Drugs, Medicines and Dressings

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer	
					Max. Qty \$	Value for Safety Net \$		

Alimentary tract and metabolism

Stomatological preparations

Stomatological preparations

Antiinfectives and antiseptics for local oral treatment

CHLORHEXIDINE GLUCONATE

4161B	Mouth wash 2 mg per mL (0.2%), 250 mL	₹1	..	..	11.89	5.60	Plaqacide	OB
4204G	Mouth wash 2 mg per mL (0.2%), 300 mL	₹1	..	..	15.28	5.60	Savacol Mouth and Throat Rinse	OM

Other agents for local oral treatment

CARMELLOSE SODIUM

4568K	Mouth spray 10 mg per mL, 25 mL	₹1	1	..	10.79	5.60	Aquae	HA
4569L	Mouth spray 10 mg per mL, 100 mL	₹1	..	..	12.46	5.60	Aquae	HA

Drugs for acid related disorders

Antacids

Calcium compounds

CALCIUM CARBONATE with GLYCINE

Note

For patients with chronic renal failure.

4055K	Tablet 420 mg-180 mg	200	5	..	*23.18	5.60	Titralac	MM

Combinations and complexes of aluminium, calcium and magnesium compounds

ALUMINIUM HYDROXIDE with MAGNESIUM HYDROXIDE and SIMETHICONE

4118R	Oral suspension 400 mg-400 mg-30 mg per 5 mL, 500 mL	2	5	..	*22.64	5.60	Mylanta Double Strength	JT
4453J	Tablet 400 mg-400 mg-40 mg	200	5	..	*46.12	5.60	Mylanta Double Strength	JT

Drugs for functional gastrointestinal disorders

Drugs for functional bowel disorders

Synthetic anticholinergics, esters with tertiary amino group

MEBEVERINE HYDROCHLORIDE

4328T	Tablet 135 mg	90	..	..	26.91	5.60	^a Colese	AF
					32.09	5.60	^a Colofac	SM

Belladonna and derivatives, plain

Belladonna alkaloids semisynthetic, quaternary ammonium compounds

HYOSCINE BUTYLBROMIDE

4279F	Injection 20 mg in 1 mL	5	..	..	24.21	5.60	Buscopan	BY

Laxatives

Laxatives

Softeners, emollients

DOCUSATE SODIUM

4200C	Tablet 50 mg	100	2	..	14.31	5.60	Coloxyl 50	FM

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Contact laxatives								
DOCUSATE SODIUM with SENNA								
4028B	Tablet 50 mg-8 mg	100	2	..	14.41	5.60	Soflax	GM
4198Y	Tablet 50 mg-8 mg	90	2	..	16.70	5.60	Coloxyl with Senna	FM
SENNA STANDARDISED								
4455L	Tablet 7.5 mg	100	1	..	13.86	5.60	Senokot	RC
Bulk producers								
ISPAGHULA HUSK								
4285M	Sachets 3.5 g, 30	£1	1	..	17.64	5.60	Fybogel	RC
PSYLLIUM HYDROPHILIC MUCILLOID								
4419N	Oral powder (orange-flavoured, sugar-free) 283 g	£1	1	..	21.67	5.60	Metamucil Smooth Texture Orange	PY
4422R	Oral powder (non-flavoured) 336 g	£1	1	..	21.67	5.60	Metamucil Regular	PY
PSYLLIUM HYDROPHILIC MUCILLOID with HIGH AMYLOSE MAIZE STARCH								
4416K	Oral powder 2.7 g-0.7 g per 7.5 g, 440 g	£1	1	..	21.27	5.60	Nucolox	SI
STERCULIA with FRANGULA BARK								
4558X	Granules 620 mg-80 mg per g (62%-8%), 500 g	£1	1	..	24.95	5.60	Normacol Plus	NE
Enemas								
SORBITOL with SODIUM CITRATE and SODIUM LAURYL SULFOACETATE								
4462W	Enemas 3.125 g-450 mg-45 mg in 5 mL, 4	£1	..	..	12.10	5.60	Microlax	JT
Other laxatives								
GLYCEROL								
<u>Restricted benefit</u>								
Short-term use when oral laxative therapy has failed or is inappropriate.								
4246L	Suppositories 2.8 g (for adults), 12	3	..	..	*19.74	5.60	Petrus Pharmaceuticals Pty Ltd	PP

Antibesity preparations, excl. diet products

Antibesity preparations, excl. diet products

Peripherally acting antiobesity products

ORLISTAT

Authority required

For the treatment of obese patients.

Total treatment will not exceed 12 months from initial application.

Patients are eligible for 1 continuous treatment in a lifetime.

The patient must be receiving, or enrolled to receive, professional dietetic and weight management advice (where this is available).

Initial treatment for patients who meet the following criteria to qualify:

(a) Body Mass Index (BMI) greater than or equal to 35 with no known co-morbidities; or

(b) BMI greater than or equal to 30 with 1 or more of the following co-morbidities:

(i) diabetes;

(ii) ischaemic heart disease;

(iii) psychiatric conditions;

(iv) hypertension.

The prescriber must provide the following:

(a) initial body weight; and

(b) BMI.

Alimentary tract and metabolism

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	
Continuing treatment for patients who have previously been issued with an authority prescription for orlistat. After 3 months and up to 6 months following commencement of orlistat treatment, patient's initial body weight must have been reduced by 2.5 kg or 2.5% (whichever is the lesser).							
Continuing treatment for patients who have previously been issued with an authority prescription for orlistat. After 6 months and up to 12 months following commencement of orlistat treatment, patient's initial body weight must have been reduced by 5 kg or 5% (whichever is the lesser).							
<u>Note</u> The patient should be ideally enrolled in an exercise program and be receiving supplemental vitamins.							
4570M	Capsule 120 mg	84	2	..	140.16	5.60	Xenical RO

Vitamins

Vitamin B₁, plain and in combination with vitamin B₆ and vitamin B₁₂

Vitamin B₁, plain

THIAMINE HYDROCHLORIDE

4043T	Tablet 100 mg	100	2	..	10.82	5.60	Betamin SW

Vitamin B-complex, incl. combinations

Vitamin B-complex, plain

VITAMIN B GROUP COMPLEX

4493L	Oral liquid 200 mL	£1	2	..	13.34	5.60	Accomin Adult Tonic WT

Mineral supplements

Calcium

Calcium

CALCIUM

Restricted benefit

Hypocalcaemia;

Osteoporosis;

Proven calcium malabsorption.

4082W	Tablet 600 mg (as carbonate)	120	1	..	14.31	5.60	CAL-600 PP
4333C	Tablet (chewable) 500 mg (as carbonate)	120	1	..	*18.44	5.60	Cal-Sup IA

CALCIUM

Restricted benefit

Hyperphosphataemia in chronic renal failure.

4094L	Tablet (chewable) 500 mg (as carbonate)	240	1	..	*30.46	5.60	Cal-Sup IA
4142B	Tablet 600 mg (as carbonate)	240	1	..	*22.20	5.60	CAL-600 PP

Other mineral supplements

Magnesium

MAGNESIUM ASPARTATE

Restricted benefit

Patients with documented hypomagnesaemia.

4321K	Tablet 500 mg	50	..	..	14.39	5.60	Magmin BB
							Mag-Sup PP

Blood and blood forming organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer	
					Max. Qty \$	Value for Safety Net \$		

Blood and blood forming organs

Antithrombotic agents

Antithrombotic agents

Platelet aggregation inhibitors excl. heparin

ASPIRIN

4076M	Tablet 100 mg (with glycine)	90	1	..	15.67	5.60	Cardiprin 100	RC

ASPIRIN

Note

The enteric coated preparations are for patients with a significant risk of gastrointestinal bleeding.

4077N	Tablet 100 mg (enteric coated)	84	1	..	13.71	5.60	Cartia	GC

4078P	Capsule 100 mg (containing enteric coated pellets)	84	1	..	14.62	5.60	Astrix	YN

CLOPIDOGREL

Authority required

For use in patients pre- and post-angioplasty.

4179Y	Tablet 75 mg (as hydrogen sulfate)	28	3	..	71.61	5.60	^a Iscover	BQ
						^a Plavix		SW

Blood substitutes and perfusion solutions

Irrigating solutions

Salt solutions

SODIUM CHLORIDE

4460R	Irrigation solution 9 mg per mL (0.9%), 500 mL	£1	2	..	10.33	5.60	Baxter Healthcare Pty Ltd	BX

4461T	Irrigation solution 9 mg per mL (0.9%), 1 L	£1	2	..	10.65	5.60	Baxter Healthcare Pty Ltd	BX

Cardiovascular system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

Cardiovascular system

Vasoprotectives

Agents for treatment of hemorrhoids and anal fissures for topical use

Corticosteroids

HYDROCORTISONE with CINCHOCAINE HYDROCHLORIDE

Caution

Long-term use may lead to skin atrophy.

4036K	Ointment 5 mg-5 mg per g (0.5%-0.5%), 30 g	‡1	..	..	22.53	5.60	Proctosedyl	SW
4038M	Suppositories 5 mg-5 mg, 12	‡1	..	..	21.24	5.60	Proctosedyl	SW

Other agents for treatment of hemorrhoids and anal fissures for topical use

ZINC OXIDE

4039N	Compound ointment 50 g	‡1	1	..	14.44	5.60	Anusol	JT
4040P	Compound suppositories, 12	‡1	1	..	13.35	5.60	Anusol	JT

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	

Dermatologicals

Antifungals for dermatological use

Antifungals for topical use

Antibiotics

NYSTATIN

4001N	Cream 100,000 units per g, 15 g	‡1	1	..	12.49	5.60	Mycostatin	FM

Imidazole and triazole derivatives

CLOTRIMAZOLE

4004R	Cream 10 mg per g (1%), 20 g	‡1	1	..	8.84	5.60	Clonea	AF

KETOCONAZOLE

Restricted benefit

Severe seborrhoeic dermatitis.

4007X	Shampoo 20 mg per g (2%), 100 mL	‡1	..	..	19.37	5.60	Sebizole	GM
4008Y	Shampoo 20 mg per g (2%), 60 mL	‡1	..	..	18.31	5.60	Nizoral 2%	JT

MICONAZOLE

4341L	Tincture 20 mg per mL (2%), 30 mL	‡1	1	..	19.47	5.60	Daktarin	JT

MICONAZOLE NITRATE

4454K	Cream 20 mg per g (2%), 30 g	‡1	1	..	14.79	5.60	Daktarin	JT

Other antifungals for topical use

AMOROLFINE HYDROCHLORIDE

Restricted benefit

Onychomycosis.

4010C	Nail treatment kit containing nail lacquer 50 mg (base) per mL (5%), 5 mL, 60 isopropyl alcohol cleaning pads, 10 spatulas and 30 nail files	‡1	1	..	96.14	5.60	Loceryl	GA

CICLOPIROX OLAMINE

Restricted benefit

Severe seborrhoeic dermatitis.

4106D	Shampoo 15 mg per g (1.5%), 60 mL	‡1	..	..	16.56	5.60	Stieprox Liquid	GK

TERBINAFINE

Restricted benefit

Tinea pedis.

4463X	Gel 10 mg per g (1%), 15 g	‡1	..	..	23.35	5.60	Lamisil DermGel	NC
4473K	Cream containing terbinafine hydrochloride 10 mg per g (1%), 15 g	‡1	1	..	21.89	5.60	Lamisil	NC

TOLNAFTATE

4481W	Spray aerosol 10 mg per g (1%), 100 g	‡1	..	..	15.09	5.60	Tinaderm	SH

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer
Antifungals for systemic use							
<i>Antifungals for systemic use</i>							
TERBINAFINE							
<u>Authority required</u>							
Onychomycosis due to dermatophyte infection proven by microscopy or culture and confirmed by an approved pathology provider.							
4011D	Tablet 250 mg (as hydrochloride)	42	1	..	102.82	5.60	^a Tamsil SI
							^a Terbihexal SZ
							^a Terbinafine-DP GN
							^a Zabel AF
					104.28	5.60	^a Lamisil NV

Emollients and protectives

Emollients and protectives

Silicone products

DIMETHICONE with GLYCEROL

Restricted benefit

For colostomy and ileostomy use;

For use by paraplegic and quadriplegic patients;

For use with surgical appliances.

4551M	Cream 150 mg-20 mg per g (15%-2%), 500 g	‡1	..	..	26.41	5.60	Silic 15	EO
4556T	Cream 150 mg-20 mg per g (15%-2%), 75 g	‡1	..	..	12.53	5.60	Silic 15	EO

Soft paraffin and fat products

WOOL ALCOHOLS

4041Q	Ointment 100 g	‡1	1	..	14.18	5.60	Eucerin	BE

Carbamide products

UREA

4042R	Cream 100 mg per g (10%), 100 g	‡1	2	..	12.19	5.60	Aquacare H.P.	AG
					12.45	5.60	Urederm	HA
					12.77	5.60	Calmurid	OL

Other emollients and protectives

CARMELLOSE SODIUM with PECTIN and GELATIN

4518T	Paste 167 mg-167 mg-167 mg per g (16.7%-16.7%- 16.7%), 5 g	‡1	..	..	11.85	5.60	Orabase	SI

SKIN EMOLLIENT

4107E	Lotion 500 mL	‡1	2	..	17.35	5.60	Alpha Keri Lotion	MT
4122Y	Bath oil 500 mL	‡1	2	..	17.35	5.60	Alpha Keri Bath Oil	MT
					19.76	5.60	QV Bath Oil	EO
					19.85	5.60	Hamilton Skin Therapy Oil	HA

Protectives against UV-radiation

Protectives against UV-radiation for topical use

SUNSCREENS

4543D	Solid stick 4.5 g	‡1	2	..	12.30	5.60	Hamilton Solastick 30+	HA
4544E	Cream 100 g	‡1	2	..	16.07	5.60	Hamilton Sunscreen Family Sunscreen	HA

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
4546G	Lotion (non-alcoholic) 125 mL	1	2	..	15.98	5.60	Cream SPF 15	
					16.07	5.60	Aquasun Lotion SPF 18	PF
					16.99	5.60	Hamilton Sunscreen Family Sunscreen Milk SPF 15 SunSense Ultra SPF 30+	HA EO

Antipruritics, incl. antihistamines, anesthetics, etc.

Antipruritics, incl. antihistamines, anesthetics, etc.

Anesthetics for topical use

LIGNOCAINE HYDROCHLORIDE with CARBOXYMETHYLCELLULOSE								
4308R	Mucilage 20 mg-25 mg per mL (2%-2.5%), 200 mL	1	..	..	79.35	5.60	Xylocaine Viscous	AP

Other antipruritics

PINE TAR with TRIETHANOLAMINE LAURYL SULFATE

Note

For patients who have failed to respond to simple moisturising agents.

4408B	Solution 23 mg-60 mg per mL (2.3%-6%), 500 mL	1	2	..	20.73	5.60	Hamilton Pine Tar Solution	HA
					22.92	5.60	Pinetarsol	EO

Antipsoriatics

Antipsoriatics for topical use

Tars

ALLANTOIN with SULFUR, PHENOL, COAL TAR SOLUTION and MENTHOL								
4505D	Gel 25 mg-5 mg-5 mg-0.05 mL-7.5 mg per g (2.5%-0.5%-0.5%-5%-0.75%), 30 g	1	2	..	16.02	5.60	Egopsoryl-TA	EO

Antibiotics and chemotherapeutics for dermatological use

Antibiotics for topical use

Other antibiotics for topical use

MUPIROCIN

Restricted benefit

For the topical treatment of secondarily infected traumatic skin lesions.

4348W	Cream 20 mg (as calcium) per g (2%), 15 g	1	..	..	16.28	5.60	Bactroban	GK
4350Y	Ointment 20 mg per g (2%), 15 g	1	..	..	16.28	5.60	Bactroban	GK

Chemotherapeutics for topical use

Antivirals

PODOPHYLLOTOXIN

Authority required

For the treatment of ano-genital warts.

4390C	Cream 1.5 mg per g (0.15%), 5 g	1	..	..	52.66	5.60	Wartec Cream	GK
4566H	Paint 5 mg per mL (0.5%), 3.5 mL (with 30 swabs)	1	..	..	39.75	5.60	Condyline Paint	HA

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Corticosteroids, dermatological preparations								
Corticosteroids, plain								
Corticosteroids, potent (group III)								
BETAMETHASONE VALERATE								
4131K	Cream 1 mg (base) per g (0.1%), 30 g	#1	2	..	22.43	5.60	Betnovate	SI
4132L	Ointment 1 mg (base) per g (0.1%), 30 g	#1	2	..	22.43	5.60	Betnovate	SI
MOMETASONE FUROATE								
<u>Note</u>								
Application to large areas of skin for longer than four weeks is not recommended.								
4342M	Cream 1 mg per g (0.1%), 45 g	#1	..	..	30.78	5.60	Elocon	SH
4343N	Ointment 1 mg per g (0.1%), 45 g	#1	..	..	30.78	5.60	Elocon	SH
Corticosteroids, combinations with antibiotics								
Corticosteroids, moderately potent, combinations with antibiotics								
TRIAMCINOLONE ACETONIDE with NEOMYCIN SULFATE, GRAMICIDIN and NYSTATIN								
<u>Caution</u>								
For the short-term treatment of localised infective eczema only.								
4482X	Ointment 1 mg-2.5 mg (base)-250 micrograms- 100,000 units per g (0.1%-0.25% (base)- 0.025%- 100,000 units in 1 g), 15 g	#1	..	..	19.09	5.60	Kenacomb	SI

Antiseptics and disinfectants								
Antiseptics and disinfectants								
Iodine products								
POVIDONE-IODINE								
4411E	Solution 100 mg per mL (10%), 100 mL	#1	..	..	22.11	5.60	Betadine Antiseptic Liquid	SW

Other dermatological preparations								
Other dermatological preparations								
Antihidrotics								
DIPHEMANIL METHYLSULFATE								
4191N	Dusting powder 20 mg per g (2%), 50 g	#1	1	..	17.74	5.60	Prantal	SH
Medicated shampoos								
PINE TAR with CADE OIL, COAL TAR SOLUTION, ARACHIS OIL EXTRACT OF CRUDE COAL TAR and OLEYL ALCOHOL								
4405W	Scalp cleanser 3 mg-3 mg-1 mg-3 mg-10 mg per mL (0.3%-0.3%-0.1%-0.3%-1%), 300 mL	#1	2	..	21.03	5.60	Polytar	GK
SALICYLIC ACID with COAL TAR SOLUTION								
4560B	Scalp cleanser 20 mg-50 mg per mL (2%-5%), 200 mL	#1	2	..	20.38	5.60	Ionil-T	GA
SALICYLIC ACID with COAL TAR SOLUTION and PINE TAR								
4447C	Scalp cleanser 20 mg-10 mg-10 mg per mL (2%- 1%-1%), 250 mL	#1	2	..	18.84	5.60	Sebitar	EO
SELENIUM SULFIDE								
4452H	Shampoo 25 mg per mL (2.5%), 125 mL	#1	..	..	14.14	5.60	Selsun	DO

Dermatologicals

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Wart and anti-corn preparations								
SALICYLIC ACID								
4389B	Gel 270 mg per g (27%), 15 g	‡1	..	..	19.54	5.60	Duofilm Gel	GK
SALICYLIC ACID with LACTIC ACID								
4386W	Liquid 167 mg-167 mg per g (16.7%-16.7%), 15 mL	‡1	..	..	18.15	5.60	Duofilm Solution	GK
Other dermatologicals								
ALLANTOIN with GLYCEROL and ICHTHAMMOL								
Note								
For patients who have failed to respond to simple moisturising agents.								
4280G	Ointment 5 mg-10 mg-10 mg per g (0.5%-1%- 1%), 50 g	‡1	2	..	18.10	5.60	Egoderm Ointment	EO
4281H	Cream 5 mg-10 mg-10 mg per g (0.5%-1%-1%), 50 g	‡1	2	..	18.10	5.60	Egoderm Cream	EO
CATIONIC CONDITIONER with PANTHENOL								
Note								
To be used in conjunction with the scalp cleanser salicylic acid with coal tar solution and pine tar (code 4447C).								
4510J	Cream 200 g	‡1	2	..	14.25	5.60	SebiRinse	EO
DICLOFENAC SODIUM								
Authority required								
For the management of actinic keratoses in patients where other standard treatments are inappropriate, and topical drug therapy is required as field treatment for clinically visible and subclinical lesions.								
Note								
Maximum quantity of four tubes (original + 3 repeats) in 12 months.								
4046Y	Gel 30 mg per g (3%), 25 g	‡1	3	..	58.19	5.60	Solaraze 3% Gel	CS
IMIQUIMOD								
Authority required								
Primary treatment of histopathologically confirmed superficial basal cell carcinoma where other standard treatments are inappropriate and topical drug therapy is required.								
4559Y	Cream 50 mg per g (5%), 250 mg single use sachets, 12	1	1	..	159.95	5.60	Aldara	IA
IMIQUIMOD								
Authority required								
Treatment of solar keratosis on the face and scalp in patients where other standard treatments are inappropriate and topical drug therapy is required as field treatment for clinically visible and subclinical lesions.								
4134N	Cream 50 mg per g (5%), 250 mg single use sachets, 12	1	1	..	159.95	5.60	Aldara	IA
SKIN CLEANSER								
4549K	Lotion 500 mL	‡1	2	..	20.74	5.60	Hamilton Skin Therapy Wash	HA
ZINC OXIDE with STARCH and CHLORPHENESIN								
4497Q	Dusting powder 100 g	‡1	1	..	12.26	5.60	Z.S.C.	SI

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Genito urinary system and sex hormones

Gynecological antiinfectives and antiseptics

Antiinfectives and antiseptics, excl. comb. with corticosteroids

Antibiotics

NYSTATIN

4012E	Cream pessaries 100,000 units, 15	£1	1	..	13.79	5.60	Nilstat	SI
4013F	Vaginal cream 100,000 units per dose, 15 doses, 75 g	£1	1	..	13.79	5.60	Nilstat	SI

Imidazole derivatives

CLOTRIMAZOLE

4016J	Vaginal cream 50 mg per 5 g (1%), 35 g	£1	..	..	15.08	5.60	APO-Clotrimazole 6 Day Cream	TX
4017K	Vaginal cream 100 mg per 5 g (2%), 20 g	£1	..	..	15.08	5.60	APO-Clotrimazole 3 Day Cream	TX

Other gynecologicals

Other gynecologicals

RICINOLEIC ACID with ACETIC ACID and HYDROXYQUINOLINE SULFATE

4434J	Vaginal jelly 7 mg-9.4 mg-250 micrograms per g (0.7%-0.94%-0.025%), 100 g	£1	..	..	32.90	5.60	Aci-Jel	CU

Sex hormones and modulators of the genital system

Estrogens

Natural and semisynthetic estrogens, plain

OESTRADIOL

Restricted benefit

Post-menopausal symptoms in women who have failed to respond using oral or topical oestrogens.

4365R	Implant 50 mg	1	..	..	83.34	5.60	Schering-Plough Pty Limited	SH
4366T	Implant 100 mg	1	..	..	121.58	5.60	Schering-Plough Pty Limited	SH

Urologicals

Other urologicals, incl. antispasmodics

Drugs used in erectile dysfunction

ALPROSTADIL

Authority required

Specific accepted war-caused or service-related disabilities for males with vasculogenic, psychogenic or neurogenic erectile dysfunction.

Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.

4579B	Intracavernosal injection 10 micrograms with diluent in single use syringe	6	3	..	*82.62	5.60	Caverject Impulse	PH
4580C	Intracavernosal injection 20 micrograms with diluent in single use syringe	6	3	..	*103.62	5.60	Caverject Impulse	PH

SILDENAFIL CITRATE

Authority required

Specific accepted war-caused or service-related disabilities for males with vasculogenic, psychogenic or neurogenic erectile dysfunction.

Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.

Genito urinary system and sex hormones

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer	
					\$	\$		
4584G	Tablet 25 mg (base)	4	5	..	60.68	5.60	Viagra	PF
4585H	Tablet 50 mg (base)	4	5	..	75.49	5.60	Viagra	PF
4586J	Tablet 100 mg (base)	4	5	..	81.12	5.60	Viagra	PF

TADALAFIL

Authority required

Specific accepted war-caused or service-related disabilities for males with vasculogenic, psychogenic or neurogenic erectile dysfunction.

Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.

4596X	Tablet 10 mg	4	5	..	79.69	5.60	Cialis	LY
4597Y	Tablet 20 mg	4	5	..	83.26	5.60	Cialis	LY

Other urologicals

SODIUM CITRO-TARTRATE

Restricted benefit

For relief of urinary symptoms when antibiotic or other therapy alone is inappropriate.

4049D	Sachets containing oral effervescent powder 4 g, 28	£1	4	..	13.55	5.60	Uracol	GM
							Ural Sachets	SI

Drugs used in benign prostatic hypertrophy

Alpha-adrenoreceptor antagonists

TAMSULOSIN HYDROCHLORIDE

Authority required

Treatment of benign prostatic hyperplasia where surgery is inappropriate, or where other drug treatment has failed or is contraindicated.

4070F	Tablet 400 micrograms (prolonged release)	30	5	..	63.36	5.60	Flomaxtra	CS

TERAZOSIN HYDROCHLORIDE

Authority required

Treatment of benign prostatic hyperplasia where surgery is inappropriate, or where other drug treatment has failed or is contraindicated.

4396J	Starter pack containing 7 tablets 1 mg and 7 tablets 2 mg	£1	..	..	20.05	5.60	Hytrin	AB
4397K	Tablet 2 mg	28	5	..	41.69	5.60	Hytrin	AB
4398L	Tablet 5 mg	28	5	..	58.19	5.60	Hytrin	AB
4399M	Tablet 10 mg	28	5	..	86.06	5.60	Hytrin	AB

Testosterone-5-alpha reductase inhibitors

FINASTERIDE

Authority required

Treatment of benign prostatic hyperplasia where surgery is inappropriate, or where other drug treatment has failed or is contraindicated.

4233T	Tablet 5 mg	30	5	..	102.12	5.60	^a Finasta	SZ
					111.69	5.60	^a Proscar	MK

Antiinfectives for systemic use

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

Antiinfectives for systemic use

Antibacterials for systemic use

Macrolides, lincosamides and streptogramins

Macrolides

AZITHROMYCIN

Restricted benefit

Upper and lower respiratory tract infections.

4115N	Tablet 500 mg	3	..	..	31.51	5.60	Zithromax	PF

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Antineoplastic and immunomodulating agents

Antineoplastic agents

Antimetabolites

Pyrimidine analogues

FLUOROURACIL							
4222F	Cream 50 mg per g (5%), 20 g	‡1	..	..	55.31	5.60	Efudix VT

Immunosuppressants

Immunosuppressants

Tumor necrosis factor alpha (TNF-alpha) inhibitors

INFLIXIMAB

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to the Veterans' Affairs Pharmaceutical Advisory Centre (VAPAC) on 1800 552 580.

Written applications for authority to prescribe infliximab should be forwarded to:

Reply Paid 9998
Veterans' Affairs Pharmaceutical Advisory Centre (VAPAC)
Department of Veterans' Affairs
GPO Box 9998
BRISBANE QLD 4001.

Authority required

Initial treatment, in combination with methotrexate, of specific accepted war-caused or service-related disability of refractory rheumatoid arthritis. Initial treatment may be prescribed by rheumatologists or consultant physicians for the reduction of signs and symptoms and prevention of structural joint damage in adult patients with active rheumatoid arthritis who satisfy all of the following criteria:

- (1) (a) Proven raised erythrocyte sedimentation rate (ESR) and/or C-reactive protein (CRP); and
- (1) (b) Proven erosive rheumatoid arthritis without end-stage disease;
- (2) Failure of an adequate trial of methotrexate and 2 other disease modifying anti-rheumatic drugs (such as sulfasalazine, hydroxychloroquine, leflunomide or cyclosporin) — unless these drugs were contraindicated or intolerance had developed;
- (3) No history of active tuberculosis requiring treatment in the last 3 years;
- (4) No history of opportunistic infection in the last 2 months;
- (5) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Infliximab (Remicade) RPBS Authority Application - Supporting Information form (contact the VAPAC on 1800 552 580 for a copy of the form).

Authority required

Continuing treatment, in combination with methotrexate, of specific accepted war-caused or service-related disability of refractory rheumatoid arthritis. Continuing treatment may be prescribed by rheumatologists or consultant physicians, following initial therapy of 3 doses, in patients who satisfy the following criteria:

- (1) There is improvement in ESR and/or CRP; and
- (2) An ACR20 (American College of Rheumatology) response is achieved by 14 weeks after the commencement of therapy.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Infliximab (Remicade) RPBS Authority Application - Supporting Information form (contact the VAPAC on 1800 552 580 for a copy of the form).

Note

Any queries concerning the arrangements to prescribe infliximab may be directed to the Veterans' Affairs Pharmaceutical Advisory Centre (VAPAC) on 1800 552 580.

Written applications for authority to prescribe infliximab should be forwarded to:

Reply Paid 9998
Veterans' Affairs Pharmaceutical Advisory Centre (VAPAC)
Department of Veterans' Affairs
GPO Box 9998
BRISBANE QLD 4001.

Antineoplastic and immunomodulating agents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty	Recordable Value for Safety Net	
					\$	\$	
4284L	Powder for I.V. infusion 100 mg	1	2	..	846.98	5.60	Remicade SH

Musculo-skeletal system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

Musculo-skeletal system

Antiinflammatory and antirheumatic products

Antiinflammatory and antirheumatic products, non-steroids

Acetic acid derivatives and related substances

DICLOFENAC SODIUM with MISOPROSTOL

Authority required

Patients requiring an NSAID in whom a risk of upper gastrointestinal complications is high or with a history of peptic ulcer disease.

4190M	Tablet 50 mg-200 micrograms	60	2	..	37.78	5.60	Arthrotec 50	PH

Topical products for joint and muscular pain

Topical products for joint and muscular pain

Preparations with salicylic acid derivatives

METHYL SALICYLATE

4022Q	Compound cream APF, 100 g	£1	1	..	14.02	5.60	Gold Cross	BI
4023R	Ointment BP, 100 g	£1	1	..	12.17	5.60	Gold Cross	BI
4026X	Liniment APF, 100 mL	£1	1	..	9.94	5.60	Gold Cross	BI

Drugs for treatment of bone diseases

Drugs affecting bone structure and mineralization

Bisphosphonates

RISEDRONATE SODIUM

Authority required

For preservation of bone mineral density in patients on long-term glucocorticoid therapy where patients are undergoing continuous treatment with a dose equal to or greater than 7.5 mg of prednisone or equivalent per day. Prescribers need to demonstrate that the patient has been on continuous therapy for 3 months or more and demonstrate that the patient is osteopenic (bone mineral density t-score of less than -1.0).

4443W	Tablet 5 mg	28	5	..	53.34	5.60	Actonel	SW
4444X	Tablet 35 mg	4	5	..	53.34	5.60	Actonel Once-a-Week	SW

Bisphosphonates, combinations

RISEDRONATE SODIUM and CALCIUM CARBONATE

Authority required

For preservation of bone mineral density in patients on long-term glucocorticoid therapy where patients are undergoing continuous treatment with a dose equal to or greater than 7.5 mg of prednisone or equivalent per day. Prescribers need to demonstrate that the patient has been on continuous therapy for 3 months or more and demonstrate that the patient is osteopenic (bone mineral density t-score of less than -1.0).

4059P	Pack containing 4 tablets risedronate sodium 35 mg and 24 tablets calcium carbonate 1.25 g (equivalent to 500 mg calcium)	£1	5	..	53.34	5.60	Actonel Combi	SW

RISEDRONATE SODIUM and CALCIUM CARBONATE with COLECALCIFEROL

Authority required

For preservation of bone mineral density in patients on long-term glucocorticoid therapy where patients are undergoing continuous treatment with a dose equal to or greater than 7.5 mg of prednisone or equivalent per day. Prescribers need to demonstrate that the patient has been on continuous therapy for 3 months or more and demonstrate that the patient is osteopenic (bone mineral density T-score of less than -1.0).

4380M	Pack containing 4 tablets risedronate sodium 35 mg and 24 sachets containing granules of calcium carbonate 2.5 g (equivalent to 1 g calcium) with colecalciferol 22 micrograms	£1	5	..	53.34	5.60	Actonel Combi D	SW

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer

Nervous system

Analgesics

Opioids

Natural opium alkaloids

MORPHINE SULFATE

Caution

The risk of drug dependence is high.

Restricted benefit

Chronic severe disabling pain not responding to non-narcotic analgesics.

Note

Authorities for increased maximum quantities and/or repeats will be granted only for

(i) chronic severe disabling pain associated with proven malignant neoplasia; or

(ii) chronic severe disabling pain where treatment has been initiated by a specialist with appropriate expertise in pain management.

4349X	Tablet 200 mg (controlled release)	20	..	..	91.28	5.60	MS Contin	MF

Diphenylpropylamine derivatives

DEXTROPROPOXYPHENE NAPSYLATE

Caution

Chronic use of this preparation is likely to cause drug dependence.

4081T	Capsule 100 mg	50	..	..	*22.27	5.60	Doloxene	AS

Other analgesics and antipyretics

Salicylic acid and derivatives

CODEINE PHOSPHATE with ASPIRIN

4061R	Tablet soluble 8 mg-300 mg	50	2	..	13.57	5.60	Aspalgin	FM

Anilides

CODEINE PHOSPHATE with PARACETAMOL

4170L	Tablet 15 mg-500 mg	20	2	..	9.73	5.60	Prodeine 15	SW
4171M	Tablet 8 mg-500 mg	50	2	..	11.74	5.60	Panamax Co.	SW
					12.86	5.60	Codalgin	FM

Other analgesics and antipyretics

GABAPENTIN

Authority required

To be approved for the treatment of refractory neuropathic pain not controlled by other drugs.

4591P	Capsule 100 mg	100	5	..	23.81	5.60	^a Gabatine 100	SI
							^a Gantin	AW
							^a Nupentin 100	AF
					24.81	5.60	^a Neurontin	PF
4592Q	Capsule 300 mg	100	5	..	62.01	5.60	^a DBL Gabapentin	HH
							^a Douglas	GN
							Gabapentin	
							300mg	
							^a Gabahexal 300mg	SZ
							^a Gabatine 300	SI
							^a Gantin	AW
							^a GenRx Gabapentin	GX
							^a Nupentin 300	AF
					63.00	5.60	^a Neurontin	PF

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	
4593R	Capsule 400 mg	100	5	..	82.24	5.60	^a DBL Gabapentin HH
							^a Douglas GN
							Gabapentin 400mg
							^a Gabahexal 400mg SZ
							^a Gabatine 400 SI
							^a Gantin AW
							^a GenRx Gabapentin GX
					83.24	5.60	^a Nupentin 400 AF
							^a Neurontin PF
4594T	Tablet 600 mg	100	5	..	127.77	5.60	Neurontin PF
4595W	Tablet 800 mg	100	5	..	166.86	5.60	^a Gantin AW
					167.86	5.60	^a Neurontin PF

PREGABALIN

Authority required

For the treatment of refractory neuropathic pain not controlled by other drugs.

4320J	Capsule 25 mg	56	5	..	42.65	5.60	Lyrica	PF
4322L	Capsule 75 mg	56	5	..	84.96	5.60	Lyrica	PF
4323M	Capsule 150 mg	56	5	..	124.24	5.60	Lyrica	PF
4324N	Capsule 300 mg	56	5	..	183.14	5.60	Lyrica	PF

Psycholeptics

Anxiolytics

Benzodiazepine derivatives

BROMAZEPAM

Authority required

Patients with terminal disease;

Patients with refractory phobic or anxiety states.

Note

For short-term use and palliative care. This drug should not be used as the first line of treatment. Other PBS-listed benzodiazepines should have been adequately tried and found to be ineffective or inappropriate. Authorities for increased quantities and/or repeats may be granted to patients with terminal disease, and other patients who have been shown to be dependent on this item by an unsuccessful attempt at gradual withdrawal.

4150K	Tablet 3 mg	60	..	..	*27.38	5.60	Lexotan	RO
4151L	Tablet 6 mg	60	..	..	*33.40	5.60	Lexotan	RO

Azaspirodecanedione derivatives

BUSPIRONE HYDROCHLORIDE

Authority required

For the short-term treatment of anxiety.

4144D	Tablet 5 mg	50	..	..	37.99	5.60	Buspar	SI
4145E	Tablet 10 mg	50	..	..	54.84	5.60	Buspar	SI

Hypnotics and sedatives

Benzodiazepine derivatives

FLUNITRAZEPAM

Authority required

Patients with terminal disease;

Patients with refractory phobic or anxiety states.

Note

For short-term use and palliative care. This drug should not be used as the first line of treatment. Other PBS-listed benzodiazepines should have been adequately tried and found to be ineffective or inappropriate. Authorities for increased quantities and/or repeats may be granted to patients with terminal disease, and other patients who have been shown to be dependent on this item by an unsuccessful attempt at gradual withdrawal.

Nervous system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$		
4216X	Tablet 1 mg	30	..	..	13.27	5.60	Hypnodorm	AF

Benzodiazepine related drugs

ZOPICLONE

Restricted benefit

For the short-term treatment of insomnia.

4522B	Tablet 7.5 mg	30	..	..	21.76	5.60 ^a	Imrest	AF
					24.92	5.60 ^a	Imovane	SW

Other nervous system drugs

Drugs used in addictive disorders

Drugs used in nicotine dependence

NICOTINE

Authority required

Patients who have indicated that they are ready to cease smoking and who have entered a support and counselling program.

Note

Studies have shown that successful therapy with this drug is enhanced by patient participation in a support and counselling program.

4571N	Transdermal patches releasing approximately 7 mg per 24 hours, 7	2	..	..	*51.38	5.60	QuitX	AF
4572P	Transdermal patches releasing approximately 14 mg per 24 hours, 7	2	..	..	*54.56	5.60	QuitX	AF
					*68.74	5.60	Nicabate CQ 14	GC
4573Q	Transdermal patches releasing approximately 21 mg per 24 hours, 7	2	2	..	*57.68	5.60	QuitX	AF
					*68.74	5.60	Nicabate CQ 21	GC
4576W	Transdermal patches releasing approximately 5 mg per 16 hours, 7	2	..	..	*50.82	5.60	Nicorette Patch	JT
4577X	Transdermal patches releasing approximately 10 mg per 16 hours, 7	2	..	..	*54.78	5.60	Nicorette Patch	JT
4578Y	Transdermal patches releasing approximately 15 mg per 16 hours, 7	2	2	..	*59.96	5.60	Nicorette Patch	JT

Antiparasitic products, insecticides and repellents

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

Antiparasitic products, insecticides and repellents

Anthelmintics

Antinematodal agents
Benzimidazole derivatives

MEBENDAZOLE							
4325P	Tablet 100 mg	6	..	..	14.92	5.60	Vermox BI

Respiratory system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	

Respiratory system

Nasal preparations

Decongestants and other nasal preparations for topical use

Sympathomimetics, plain

OXYMETAZOLINE HYDROCHLORIDE								
4378K	Nasal spray 500 micrograms per mL (0.05%), 15 mL	‡1	..	..	17.15	5.60	Drixine	SH
4379L	Nasal spray 500 micrograms per mL (0.05%), 18 mL	‡1	..	..	16.76	5.60	Logicin Rapid Relief	SI

Antiallergic agents, excl. corticosteroids

LEVOCABASTINE HYDROCHLORIDE								
4311X	Nasal spray 500 micrograms per mL (0.05%), 10 mL (100 doses)	‡1	2	..	18.24	5.60	Livostin	JT
SODIUM CROMOGLYCATE								
4468E	Nasal spray metered dose pump 20 mg per mL (2%), 26 mL	‡1	5	..	22.91	5.60	Rynacrom	SW

*Corticosteroids***BUDESONIDE**Restricted benefit

Severe intractable rhinitis.

4092J	Aqueous nasal spray (pump pack) 64 micrograms per dose (120 doses)	‡1	..	..	31.73	5.60	Budamax Aqueous	PM

*Other nasal preparations***IPRATROPIUM BROMIDE**Restricted benefit

Severe intractable rhinorrhoea, associated with perennial rhinitis, unresponsive to insufflated nasal steroids.

4089F	Aqueous nasal spray (pump pack) 21 micrograms (anhydrous) per dose (180 doses)	‡1	5	..	23.59	5.60	Atrovent Nasal Aqueous	BY
4090G	Aqueous nasal spray (pump pack) 42 micrograms (anhydrous) per dose (180 doses)	‡1	5	..	30.47	5.60	Atrovent Nasal Forte	BY

Nasal decongestants for systemic use

Sympathomimetics

PSEUDOEPHEDRINE HYDROCHLORIDE								
4029C	Tablet 60 mg	12	..	..	11.02	5.60	Logicin Sinus	SI

Cough and cold preparations

Expectorants, excl. combinations with cough suppressants

Expectorants

SENEGA and AMMONIA								
4074K	Mixture 200 mL	‡1	4	..	9.18	5.60	Gold Cross	BI

Cough suppressants, excl. combinations with expectorants

Opium alkaloids and derivatives

PHOLCODINE								
4071G	Linctus 1 mg per mL (0.1%), 100 mL	‡1	2	..	9.02	5.60	Gold Cross	BI

Respiratory system

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer	
					Max. Qty \$	Safety Net \$		
					14.61	5.60	Duro-Tuss	IA

Antihistamines for systemic use

Antihistamines for systemic use***Phenothiazine derivatives*****PROMETHAZINE HYDROCHLORIDE****Caution**

Significant side effects may occur.

4072H	Tablet 10 mg	50	2	..	14.67	5.60	Phenergan	SW
4073J	Tablet 25 mg	50	2	..	16.76	5.60	Phenergan	SW

Piperazine derivatives**CETIRIZINE HYDROCHLORIDE**

4175R	Tablet 10 mg	30	..	..	29.65	5.60	^a Alzene	AF
					32.87	5.60	Zilarex	SZ
					39.45	5.60	^a Zyrtec	JT

Other antihistamines for systemic use**FEXOFENADINE HYDROCHLORIDE**

4237B	Tablet 60 mg	60	..	..	*41.13	5.60	^a Fexal	SZ
					*54.99	5.60	^a Telfast	SW
4238C	Tablet 120 mg	30	..	..	29.42	5.60	^a Xergic	AF
					34.71	5.60	^a Fexal	SZ
					47.13	5.60	^a Telfast 120	SW

LORATADINE

4313B	Tablet 10 mg	30	..	..	32.99	5.60	^a Allereze	AF
					43.65	5.60	^a Lorano	SZ
					45.92	5.60	^a Claratyne	SH

Sensory organs

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	

Sensory organs

Ophthalmologicals

Decongestants and antiallergics

Sympathomimetics used as decongestants

ANTAZOLINE with NAPHAZOLINE								
4031E	Eye drops 5 mg (sulfate)-250 micrograms (nitrate) per mL (0.5%-0.025%), 10 mL	‡1	1	..	13.86	5.60	Antistine-Privine	NV
4032F	Eye drops 5 mg (phosphate)-500 micrograms (hydrochloride) per mL (0.5%-0.05%), 15 mL	‡1	1	..	14.80	5.60	Albalon-A	AG
NAPHAZOLINE HYDROCHLORIDE								
4035J	Eye drops 1 mg per mL (0.1%), 15 mL	‡1	1	..	15.09	5.60	Albalon Liquifilm	AG

Other antiallergics

LEVOCABASTINE HYDROCHLORIDE								
4310W	Eye drops 500 micrograms per mL (0.05%), 4 mL (120 doses)	‡1	1	..	18.24	5.60	Livostin	JT

Otologicals

Other otologicals

Indifferent preparations

CARBAMIDE PEROXIDE								
4176T	Ear drops 65 mg per mL (6.5%), 12 mL	‡1	..	..	16.08	5.60	Ear Clear for Ear Wax Removal	KY
DICHLOROBENZENE with CHLORBUTOL and TURPENTINE OIL								
4180B	Ear drops 20 mg-50 mg-0.1 mL per mL (2%-5%-10%), 10 mL	‡1	..	..	14.08	5.60	Cerumol	AC
DOCUSATE SODIUM								
4199B	Ear drops 5 mg per mL (0.5%), 10 mL	‡1	..	..	14.47	5.60	Waxsol	NE

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price	Maximum	Brand Name and Manufacturer
					for Max. Qty \$	Recordable Value for Safety Net \$	

Various

All other therapeutic products

All other therapeutic products
Drugs for treatment of hyperkalemia and hyperphosphatemia

SODIUM POLYSTYRENE SULFONATE							
4470G	Oral powder 454 g	‡1	2	..	71.12	5.60	Resonium-A SW

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

REPATRIATION PHARMACEUTICAL BENEFITS SCHEME (RPBS) WOUND ASSESSMENT AND DRESSING IDENTIFICATION

It is essential to define the aetiology of the wound before selecting a dressing. Recommendations are based on wound type, colour of wound base, depth of wound, and amount of exudate.

This wound chart adheres to the MOIST WOUND concept of healing and wound dressings are described below as ABSORBING or MOISTURE DONATING.

Most wound healing products are designed to remain in situ for several days, with the exception of those for infected wounds which should be changed daily. The quantities and repeats listed in the Repatriation Schedule are considered to be adequate to manage the treatment of a wound for two weeks to one month, when an assessment of the wound's healing process should be undertaken.

DRESSINGS

PINK EPITHELIALISING WOUND

Aim: To protect and promote epithelialisation. Epithelialising wounds normally are superficial and only produce a light exudate.

(A) Covering	• Film;	• Gauze—Paraffin;
	• Film Island	• Non-adherent
(B) Absorbing	• Foam (Light Exudate);	• Hydrocolloid (Superficial Wound—Light Exudate)
	• Hydroactive (Superficial Wound—Light Exudate)	

RED GRANULATING WOUND

Aims: (1) to protect the granulating tissue; (2) to encourage epithelialisation; (3) to absorb excess exudate.

LIGHT EXUDATE:	Superficial	Cavity
(A) Absorbing	• Foam (Light Exudate);	• Hydrocolloid (Cavity Wound)
	• Hydroactive (Superficial Wound—Light Exudate);	
	• Hydrocolloid (Superficial Wound—Light Exudate)	
(B) Moisture donating	• Hydrogel—Amorphous;	• Hydrogel—Amorphous
	• Hydrogel—Sheet	
HIGH EXUDATE:	Superficial	Cavity
(A) Absorbing	• Alginate (Superficial Wound);	• Alginate (Cavity Wound);
	• Foam—Heavy Exudate;	• Foam—Moderate Exudate (see “cavity conforming” product);
	• Hydroactive (Superficial Wound—Moderate Exudate);	• Hydroactive (Cavity Wound);
	• Hydrocolloid (Superficial Wound—Moderate/High Exudate)	• Hydrocolloid (Cavity Wound)
(B) Moisture donating	NOT APPROPRIATE	

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer

YELLOW SLOUGHY WOUND

Aims: (1) to remove slough; (2) to encourage granulation; (3) to absorb excess exudate.

LIGHT EXUDATE:	Superficial	Cavity
(A) Absorbing	• Cadexomer Iodine; • Foam—Light Exudate; • Foam with Charcoal; • Hydroactive (Superficial Wound—Moderate Exudate); • Hydrocolloid (Superficial Wound—Moderate Exudate)	• Cadexomer Iodine; • Hydrocolloid (Cavity Wound)
(B) Moisture Donating	• Hydrogel—Amorphous; • Hydrogel—Sheet	• Hydrogel—Amorphous

HIGH EXUDATE:	Superficial	Cavity
(A) Absorbing	• Alginate (Superficial Wound); • Cadexomer Iodine; • Foam—Heavy Exudate; • Hydroactive (Superficial Wound—Moderate/High Exudate); • Hydrocolloid (Superficial Wound—Moderate/High Exudate)	• Alginate (Cavity Wound); • Cadexomer Iodine; • Hydrocolloid (Cavity Wound)

(B) Moisture donating NOT APPROPRIATE

BLACK NECROTIC WOUND

Aim: To remove eschar by — (1) sharp debridement, e.g., scissor/scalpel and/or (2) rehydration and autolytic debridement. (These wounds usually produce a LIGHT EXUDATE.)

DRY / LIGHT EXUDATE:	Superficial	Cavity
(A) Absorbing	• Hydroactive (Superficial Wound—Light Exudate); • Hydrocolloid (Superficial Wound—Light/Moderate Exudate)	• Hydrocolloid (Cavity Wound)
(B) Moisture donating	• Hydrogel—Amorphous; • Hydrogel—Sheet	• Hydrogel—Amorphous; • Hydrogel—Sheet

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable Value for	Brand Name and Manufacturer
					Max. Qty \$	Safety Net \$	

INFECTED WOUNDS

Aims: (1) to clear the infection with systemic antibiotics; (2) to absorb excess exudate; (3) to remove slough if present; (4) to decrease bacterial burden - by applying a Silver dressing or Cadexomer Iodine dressing.

MALODOROUS WOUNDS

Aims: (1) to clear infection if present; (2) to remove slough if present; (3) to clear colonising odour-producing bacteria in slough — by applying metronidazole gel, a Silver dressing or a Cadexomer Iodine dressing; (4) to absorb excess exudate.

Products: Activated Charcoal; Alginate with Charcoal; Foam with Charcoal; Silver dressing; Cadexomer Iodine dressing.

MINOR SKIN TRAUMA

Aims: (1) to stop bleeding; (2) to prevent infection; (3) to minimise the surface defect; (4) to promote epithelialisation.

ORDERING HARTMANN PRODUCTS

Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.

ORDERING COLOPLAST PRODUCTS

Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

ORDERING MOLNLYCKE HEALTHCARE PRODUCTS

Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.

All other non-therapeutic products

All other non-therapeutic products

Other non-therapeutic auxiliary products

BANDAGE—ABSORBENT WOOL								
4653X	Bandage 10 cm x 3 m	6	..	..	20.32	5.60	Surepress 650948	CC
BANDAGE—CALICO								
4717G	Bandage, triangular, large	‡1	..	..	13.49	5.60	Handy 5608	BV
BANDAGE—COMPRESSION								
Note								
Treatment of varices and oedema associated with venous disease and lymphoedema; contraindicated in arterial disease.								
4654Y	Bandage, short stretch, 8 cm x 5 m	5	..	..	*77.77	5.60	Comprilan 1027	BV
4656C	Bandage, high stretch, 7.5 cm x 3.5 m	5	..	..	*68.37	5.60	Setopress 3504	SS
4657D	Bandage, high stretch, 10 cm x 3.5 m	5	..	..	*78.57	5.60	Setopress 3505	SS
4736G	Bandage, high stretch, 7.5 cm x 3 m	5	..	..	*94.32	5.60	Tensopress 66004347	BV
4748X	Bandage, high stretch, 10 cm x 3 m	5	..	..	*72.92	5.60	Surepress 650947	CC
					*126.62	5.60	Tensopress 66004348	BV

BANDAGE—COMPRESSION

Note

Treatment of varices and oedema associated with venous disease and lymphoedema; contraindicated in arterial disease.

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4598B	Bandage, four layer	5	..	..	*145.67	5.60	Profore Lite 66050415	SN
4658E	Bandage, four layer	5	..	..	*214.97	5.60	Profore 66050016	SN
BANDAGE—COMPRESSION								
Note Treatment of varices and oedema associated with venous disease and lymphoedema; contraindicated in arterial disease.								
Restricted benefit Initial treatment of venous ulcers.								
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4938X	Bandage, two layer, 18 cm-22 cm (red)	1	..	..	49.03	5.60	ProGuide 66000780	SN
4939Y	Bandage, two layer, 22 cm-28 cm (yellow)	1	..	..	49.03	5.60	ProGuide 66000781	SN
4940B	Bandage, two layer, 28 cm-32 cm (green)	1	..	..	49.03	5.60	ProGuide 66000782	SN
BANDAGE—COMPRESSION								
Note Treatment of varices and oedema associated with venous disease and lymphoedema; contraindicated in arterial disease.								
Restricted benefit Continuation of treatment of venous ulcers where patient's ability to tolerate dressing has been demonstrated.								
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4941C	Bandage, two layer, 18 cm-22 cm (red)	4	..	..	*174.10	5.60	ProGuide 66000780	SN
4942D	Bandage, two layer, 22 cm-28 cm (yellow)	4	..	..	*174.10	5.60	ProGuide 66000781	SN
4943E	Bandage, two layer, 28 cm-32 cm (green)	4	..	..	*174.10	5.60	ProGuide 66000782	SN
BANDAGE—RETENTION—COHESIVE—HEAVY								
4659F	Bandage 7.5 cm x 3 m	2	..	..	*20.06	5.60	Coplus 3629	BV
4660G	Bandage 10 cm x 2 m	2	..	..	*19.36	5.60	Coban 1584	MM
4811F	Bandage 5 cm x 1.3 m	2	..	..	*14.02	5.60	Peg 7420	BK
4812G	Bandage 7.5 cm x 1.3 m	2	..	..	*17.30	5.60	Peg 7422	BK
4813H	Bandage 10 cm x 1.3 m	2	..	..	*21.04	5.60	Peg 7423	BK
4814J	Bandage 15 cm x 1.3 m	2	..	..	*28.18	5.60	Peg 7425	BK
BANDAGE—RETENTION—COHESIVE—LIGHT								
4662J	Bandage 10 cm x 4 m	2	..	..	*17.14	5.60	Handygauze Cohesive 8635	BV
4718H	Bandages 2.5 cm x 4 m, 2	1	..	..	12.52	5.60	Handygauze Cohesive 8631	BV
4719J	Bandage 6 cm x 4 m	2	..	..	*14.70	5.60	Handygauze Cohesive 8633	BV

Various

					Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$		
Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium			Brand Name and Manufacturer	
BANDAGE—RETENTION—COTTON CREPE								
4727T	Bandage 5 cm x 2.3 m	2	..	..	*17.44	5.60	Telfa 8252F	KE
					*18.28	5.60	Tensocrepe 36300501	BV
4728W	Bandage 7.5 cm x 2.3 m	2	..	..	*22.22	5.60	Telfa 8253F	KE
					*22.40	5.60	Tensocrepe 36307501	BV
4729X	Bandage 10 cm x 2.3 m	2	..	..	*25.38	5.60	Telfa 8254F	KE
					*27.82	5.60	Tensocrepe 36301001	BV
BANDAGE—TUBULAR								
4663K	Bandage, straight, size C	‡1	..	..	15.38	5.60	Elastoplast 2225	BE
4664L	Bandage, straight, size D	‡1	..	..	15.38	5.60	Elastoplast 2226	BE
4665M	Bandage, straight, size E	‡1	..	..	15.38	5.60	Elastoplast 2227	BE
4855M	Bandage 6.25 cm x 1 m	‡1	..	..	18.17	5.60	Tubigrip B 1520	SS
4856N	Bandage 6.75 cm x 1 m	‡1	..	..	18.17	5.60	Tubigrip C 1545	SS
4857P	Bandage 7.5 cm x 1 m	‡1	..	..	18.17	5.60	Tubigrip D 1546	SS
4858Q	Bandage 8.75 cm x 1 m	‡1	..	..	18.17	5.60	Tubigrip E 1547	SS
4859R	Bandage 10 cm x 1 m	‡1	..	..	18.17	5.60	Tubigrip F 1548	SS
BANDAGE—TUBULAR (FINGER)								
4726R	Refill	‡1	..	..	13.73	5.60	Tubegauz 0501658	SS
4798M	Complete pack including applicator	‡1	..	..	17.77	5.60	Tubegauz 0501633	SS
BANDAGE—TUBULAR (LIGHTWEIGHT)								
4671W	Bandage, small limb size (red), 10 m	‡1	..	..	28.36	5.60	Tubifast 2434	SS
4672X	Bandage, medium limb size (green), 10 m	‡1	..	..	32.02	5.60	Tubifast 2436	SS
4673Y	Bandage, large limb size (blue), 10 m	‡1	..	..	35.58	5.60	Tubifast 2438	SS
BANDAGE—TUBULAR (LONG STOCKING)								
4674B	Bandage, small size	2	..	..	*40.18	5.60	Tubigrip 1482	SS
4675C	Bandage, XX/large size	2	..	..	*40.18	5.60	Tubigrip 1486	SS
4797L	Bandage, medium size	2	..	..	*40.18	5.60	Tubigrip 1483	SS
4799N	Bandage, large size	2	..	..	*40.18	5.60	Tubigrip 1484	SS
BANDAGE—TUBULAR (SHORT STOCKING)								
4661H	Bandage, small B/C size	2	..	..	*30.44	5.60	Tubigrip 1479	SS
4815K	Bandage, medium C/D size	2	..	..	*30.44	5.60	Tubigrip 1480	SS
4816L	Bandage, large D/E size	2	..	..	*30.44	5.60	Tubigrip 1481	SS
BANDAGE—ZINC PASTE								
Note								
Used as an adjunct in the management of leg ulceration and associated eczema and skin conditions.								
4668Q	Bandage 7.5 cm x 6 m	2	..	..	*29.20	5.60	Zincaband 3604	SS
4669R	Bandage 7.5 cm x 6 m	2	3	..	*29.66	5.60	Steripaste 3610	XP
4670T	Bandage 10 cm x 9.1 m	2	3	..	*28.78	5.60	Flexidress 650941	CC

BANDAGE—ZINC PASTE

Note

Used as an adjunct in the management of leg ulceration and associated eczema and skin conditions.

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4750B	Bandage 7.5 cm x 6 m	2	3	..	*70.64	5.60	Viscopaste 4948	SN
4760M	Bandages 80 cm (stockings), 4	£1	3	..	81.42	5.60	ZipZoc 66051550	SN
COTTON WOOL ROLL								
4701K	Roll 100 g	£1	2	..	10.51	5.60	JJ 02013	JJ
DRESSING—ACTIVATED CHARCOAL (MALODOROUS WOUND)								
4681J	Dressing 10.5 cm x 10.5 cm	10	..	..	*100.92	5.60	Actisorb Plus MAC031	JJ
4742N	Dressings 10 cm x 10 cm, 10	£1	..	..	78.98	5.60	CarboFLEX 403202	CC
4743P	Dressings 15 cm x 20 cm, 5	£1	..	..	89.87	5.60	CarboFLEX 403204	CC
DRESSING—ALGINATE (CAVITY WOUND)								
Note This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.								
4832H	Rope 2 g	10	..	..	*109.12	5.60	Sorbsan 1411	UM
					*115.26	5.60	Kaltostat 168117	CC
DRESSING—ALGINATE (CAVITY WOUND)								
Note This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.								
Note Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.								
4682K	Ropes 2 g (40 cm), 6	2	..	..	*137.72	5.60	Comfeel SeaSorb Filler 3740	CT
DRESSING—ALGINATE (SUPERFICIAL WOUND)								
Note This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.								
Note Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.								
4684M	Dressing 5 cm x 5 cm	10	1	..	*46.92	5.60	Comfeel SeaSorb Dressing 3705	CT
4831G	Dressing 10 cm x 10 cm	10	1	..	*84.42	5.60	Sorbsan 1410	UM
					*90.12	5.60	Comfeel SeaSorb Dressing 3710	CT
DRESSING—ALGINATE (SUPERFICIAL WOUND)								
Note This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.								
4683L	Dressings 7.5 cm x 12 cm, 10	£1	1	..	91.08	5.60	Kaltostat 168212	CC

DRESSING—ALGINATE (SUPERFICIAL WOUND)

Note

This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4691X	Dressings 15 cm x 20 cm, 10	‡1	1	..	225.99	5.60	Algisite M 66000521	SN
4699H	Dressings 5 cm x 5 cm, 10	‡1	1	..	49.36	5.60	Algisite M 66000519	SN
					49.40	5.60	Kaltostat 168210	CC
4700J	Dressings 10 cm x 10 cm, 10	‡1	1	..	94.05	5.60	Algisite M 66000520	SN
DRESSING—FILM								
4686P	Dressings 6 cm x 7 cm, 8	‡1	..	..	15.64	5.60	Nexcare Tegaderm Transparent H1624	MM
4687Q	Dressings 10 cm x 12 cm, 4	‡1	..	..	19.65	5.60	Nexcare Tegaderm Transparent H1626	MM
4688R	Dressing 15 cm x 20 cm	6	..	..	*30.66	5.60	Tegaderm Transparent 1628	MM
DRESSING—FILM								
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4893M	Dressings 10 cm x 12 cm, 10	‡1	..	..	31.26	5.60	Op-Site Flexigrid 4629	SN
DRESSING—FILM ISLAND								
4689T	Dressing 5 cm x 7 cm	10	..	..	*16.22	5.60	Tegaderm Transparent Island 3582	MM
4690W	Dressing 9 cm x 10 cm	10	..	..	*27.72	5.60	Tegaderm Transparent Island 3586	MM
DRESSING—FILM ISLAND								
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4898T	Dressings 5 cm x 7.2 cm, 5	2	..	..	*28.92	5.60	Cutifilm Plus 76309	SN
4899W	Dressings 8 cm x 10 cm, 5	2	..	..	*45.50	5.60	Cutifilm Plus 76308	SN
DRESSING—FOAM—HEAVY EXUDATE								
Note This dressing should remain in place until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.								
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4795J	Dressings 10 cm x 10 cm, 10	‡1	1	..	75.06	5.60	Lyof foam Extra 603088	XP
					120.97	5.60	Allevyn 66007637	SN

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
DRESSING—FOAM—HEAVY EXUDATE								
Note This dressing should remain in place until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.								
4880W	Dressings 20 cm x 15 cm, 10	1	1	..	189.11	5.60	Lyof foam Extra 603090	XP
DRESSING—FOAM—MODERATE EXUDATE								
Note This dressing should remain in place until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.								
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4590N	Dressings 12.5 cm x 12.5 cm, 10	1	..	..	117.66	5.60	Allevyn Adhesive 66000044	SN
4694C	Dressing, cavity, conforming, 20 g	1	1	..	84.82	5.60	Cavicare 4563	SN
DRESSING—FOAM—MODERATE EXUDATE								
Note This dressing should remain in place until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.								
4878R	Dressings 20 cm x 15 cm, 10	1	1	..	101.86	5.60	Lyof foam Flat 603095	XP
4890J	Dressings 7.5 cm x 7.5 cm, 10	1	1	..	42.84	5.60	Lyof foam Flat 603092	XP
4891K	Dressings 10 cm x 10 cm, 10	1	1	..	49.48	5.60	Lyof foam Flat 603093	XP
DRESSING—FOAM with CHARCOAL (MALODOROUS WOUND)								
Note This dressing should remain in place on wounds with odour until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.								
4892L	Dressings 10 cm x 10 cm, 10	2	..	..	*174.06	5.60	Lyof foam C 603025	SS
DRESSING—FOAM with SILICONE—HEAVY EXUDATE								
Note Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.								
4642H	Dressings 7.5 cm x 7.5 cm, 5	1	..	..	30.77	5.60	Mepilex Border 295200	MH
4643J	Dressings 10 cm x 10 cm, 5	1	..	..	42.68	5.60	Mepilex Border 295300	MH
DRESSING—FOAM with SILICONE—LIGHT EXUDATE								
Note Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.								
4644K	Dressings 6 cm x 8.5 cm, 5	1	..	..	28.06	5.60	Mepilex Lite 284000	MH
4645L	Dressings 10 cm x 10 cm, 5	1	..	..	38.21	5.60	Mepilex Lite 284100	MH

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
DRESSING—FOAM with SILICONE—MODERATE EXUDATE								
Note Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.								
4626L	Dressings 10 cm x 10 cm, 5	‡1	..	..	42.68	5.60	Mepilex 294100	MH
DRESSING—GAUZE (ABSORBENT PAD)								
4707R	Pads 5 cm x 5 cm, 100	‡1	..	..	13.88	5.60	Handy 5672	BV
4708T	Pads 10 cm x 10 cm, 100	‡1	..	..	27.40	5.60	Handy 5674	BV
DRESSING—GAUZE—EYE PAD								
4768Y	Pads, 12	‡1	..	..	12.83	5.60	Curity 4112	KE
DRESSING—GAUZE—PARAFFIN								
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4759L	Dressings 10 cm x 10 cm, 10	‡1	..	..	19.29	5.60	Jelonet 7404	SN
DRESSING—GAUZE—PARAFFIN with CHLORHEXIDINE ACETATE								
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4845B	Dressings 10 cm x 10 cm, 10	‡1	2	..	26.05	5.60	Bactigras 7457	SN
DRESSING—HYDROACTIVE (CAVITY WOUND)								
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4918W	Dressings 5 cm x 6 cm, 10	‡1	1	..	84.60	5.60	Allevyn Plus Cavity 66047571	SN
4919X	Dressings 10 cm x 10 cm, 5	2	1	..	*178.36	5.60	Allevyn Plus Cavity 66047573	SN
DRESSING—HYDROACTIVE (DEBRIDEMENT)								
Note Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.								
4948K	Dressings 5.5 cm, 8	‡1	..	..	68.66	5.60	TenderWet Active Cavity	HR
4949L	Dressings 4 cm, 8	‡1	..	..	67.90	5.60	TenderWet 24 Active	HR
4950M	Dressings 7.5 cm x 7.5 cm, 8	‡1	..	..	92.19	5.60	TenderWet 24 Active	HR
DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—HIGH EXUDATE)								
4692Y	Dressings (foam alternative) 10 cm x 10 cm, 10	‡1	..	..	54.90	5.60	CombiDERM 651031	CC
4693B	Dressings (foam alternative) 15 cm x 18 cm, 5	‡1	..	..	71.72	5.60	CombiDERM 651027	CC
4695D	Dressings, island, 11 cm x 11 cm, 10	‡1	..	..	111.24	5.60	Tielle MTL101E	JJ
4696E	Dressings, island, 18 cm x 18 cm, 5	‡1	..	..	135.84	5.60	Tielle MT2442	JJ

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer	
					Max. Qty \$	Value for Safety Net \$		

DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—HIGH EXUDATE)

Note

Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

4927H	Non-adhesive waterproof semi-permeable absorbent foam pads 10 cm x 10 cm, 10	#1	1	..	87.93	5.60	Biatain Non-adhesive 3410	CT
4928J	Non-adhesive waterproof semi-permeable absorbent foam pads 15 cm x 15 cm, 5	#1	2	..	86.45	5.60	Biatain Non-adhesive 3413	CT
4929K	Adhesive waterproof semi-permeable absorbent foam pads 12 cm x 12 cm, 10	#1	1	..	96.95	5.60	Biatain Adhesive 3420	CT
4930L	Adhesive waterproof semi-permeable absorbent foam pads 18 cm x 18 cm, 5	#1	2	..	93.82	5.60	Biatain Adhesive 3423	CT

DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—LIGHT EXUDATE)

Note

Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.

4905E	Dressings 5 cm x 6 cm, 10	#1	1	..	56.69	5.60	Allevyn Thin 66047576	SN
4906F	Dressings 10 cm x 10 cm, 5	2	1	..	*103.50	5.60	Allevyn Thin 66047578	SN

DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—MODERATE EXUDATE)

Note

Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.

4885D	Dressings 5 cm x 6 cm, 10	#1	1	..	46.13	5.60	Cutinova Hydro 66047441	SN
4886E	Dressings 10 cm x 10 cm, 5	2	1	..	*75.98	5.60	Cutinova Hydro 66047443	SN

DRESSING—HYDROCOLLOID (CAVITY WOUND)

Note

This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

4896Q	Paste 30 g	10	..	..	*145.12	5.60	DuoDERM Paste H7930	CC

DRESSING—HYDROCOLLOID (CAVITY WOUND)

Note

This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

Note

Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

4895P	Paste 50 g	2	3	..	*43.22	5.60	Comfeel Paste 4701	CT

DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—LIGHT EXUDATE)

Note

This dressing should be applied to a thickness of 3 mm to 5 mm. It should be covered with a hydrocolloid dressing and may be left in place for up to 7 days.

Note

Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
4888G	Dressings 5 cm x 7 cm, 10	‡1	1	..	41.72	5.60	Comfeel Plus Transparent 3530	CT
4889H	Dressings 9 cm x 14 cm, 10	‡1	1	..	84.52	5.60	Comfeel Plus Transparent 3536	CT
4924E	Dressings 10 cm x 10 cm, 10	‡1	1	..	69.78	5.60	Comfeel Plus Transparent 3533	CT

DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—LIGHT EXUDATE)

Note

This dressing should be applied to a thickness of 3 mm to 5 mm. It should be covered with a hydrocolloid dressing and may be left in place for up to 7 days.

4907G	Dressings 10 cm x 10 cm, 10	‡1	1	..	71.72	5.60	DuoDERM Extra Thin H7955	CC

DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—LIGHT EXUDATE)

Note

This dressing should be applied to a thickness of 3 mm to 5 mm. It should be covered with a hydrocolloid dressing and may be left in place for up to 7 days.

Note

Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.

4947J	Dressings 10 cm x 10 cm, 10	‡1	1	..	48.15	5.60	Hydrocoll Thin 900942/1	HR

DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—MODERATE EXUDATE)

Note

This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

4897R	Dressings 10 cm x 10 cm, 5	2	1	..	*81.40	5.60	DuoDERM CGF H7660	CC
4920Y	Dressings 20 cm x 20 cm, 5	2	1	..	*222.32	5.60	DuoDERM CGF H7662	CC

DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—MODERATE EXUDATE)

Note

This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

Note

Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.

4921B	Dressings 10 cm x 10 cm, 10	‡1	1	..	76.78	5.60	Repicare Ultra 66000434	SN

DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—MODERATE EXUDATE)

Note

This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

Note

Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.

4945G	Dressings 10 cm x 10 cm, 10	‡1	1	..	48.15	5.60	Hydrocoll 900938/1	HR
4946H	Dressings 15 cm x 15 cm, 10	‡1	1	..	89.91	5.60	Hydrocoll 900939/1	HR

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty	Maximum Recordable Value for Safety Net	Brand Name and Manufacturer
					\$	\$	

DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—MODERATE EXUDATE)

Note

This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

Note

Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

4678F	Butterfly shape 7 cm	5	..	..	*55.17	5.60	Comfeel Plus Pressure Relieving 3350	CT
4679G	Round 10 cm	5	..	..	*59.62	5.60	Comfeel Plus Pressure Relieving 3353	CT
4923D	Dressings with alginate 10 cm x 10 cm, 10	1	1	..	81.99	5.60	Comfeel Plus Ulcer Dressing 3110	CT

DRESSING—HYDROFIBRE (ALTERNATE TO ALGINATES)

4649Q	Dressings 10 cm x 10 cm, 10	1	1	..	100.98	5.60	Aquacel 177902	CC
4698G	Ropes 2 g (30 cm), 5	1	1	..	83.71	5.60	Aquacel 177904	CC
4922C	Dressings 15 cm x 15 cm, 5	2	1	..	*208.70	5.60	Aquacel 177903	CC

DRESSING—HYDROGEL—AMORPHOUS

Note

This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.

Note

Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

4912M	Tubes 15 g, 10	1	1	..	64.48	5.60	DuoDERM Gel H7990	CC
					72.09	5.60	Comfeel Purilon Gel 3900	CT

DRESSING—HYDROGEL—AMORPHOUS

Note

This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.

4913N	Tubes 30 g, 3	3	1	..	*97.11	5.60	DuoDERM Gel H7987	CC
4914P	Tube 50 g	3	3	..	*33.12	5.60	Solugel 10336	JJ

DRESSING—HYDROGEL—AMORPHOUS

Note

This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.

Note

Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.

4599C	Tube 50 g	3	3	..	*30.51	5.60	SoloSite Gel 36361338	SN
4894N	Tube 25 g	4	3	..	*59.54	5.60	Intrasite Gel 7313	SN

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
DRESSING—HYDROGEL—SHEET								
Note This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.								
4911L	Dressings 9.5 cm x 10.2 cm, 5	2	..	..	*83.20	5.60	Nu-Gel 2497	JJ
DRESSING—HYDROGEL—SHEET								
Note This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.								
Note Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.								
4806Y	Dressings 10 cm x 10 cm, 5	2	..	..	*53.28	5.60	Aquaclear 900796	HR
DRESSING—NON-ADHERENT								
Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4819P	Dressings 5 cm x 5 cm, 5	2	..	..	*15.80	5.60	Cutilin Non-Stick Wound Pad 76301	SN
4860T	Dressings 5 cm x 5 cm, 5	2	..	..	*16.48	5.60	Melolin 36361357	SN
4861W	Dressings 10 cm x 10 cm, 10	1	..	..	32.40	5.60	Melolin 66974933	SN
4862X	Dressings 10 cm x 10 cm, 5	2	..	..	*25.40	5.60	Cutilin Non-Stick Wound Pad 76300	SN
DRESSING—NON-ADHERENT								
4755G	Dressings 5 cm x 7.5 cm, 10	1	..	..	11.02	5.60	Telfa 1970C	KE
4758K	Dressings 7.5 cm x 10 cm, 6	1	..	..	11.23	5.60	Telfa 2140C	KE
4844Y	Dressings, self-adhesive, 7.5 cm x 10 cm, 6	1	2	..	12.02	5.60	Telfa 7650C	KE
DRESSING—NON-ADHERENT								
Note Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.								
4944F	Dressings 7.5 cm x 10 cm, 10	1	..	..	15.24	5.60	Atrauman 499513	HR
DRESSING—NON-ADHERENT								
Note Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.								
4243H	Dressings, non-woven, with silicone 5 cm x 7.5 cm, 10	1	..	..	63.62	5.60	Mepitel 290510	MH
4244J	Dressings, non-woven, with silicone 7.5 cm x 10 cm, 10	1	..	..	107.62	5.60	Mepitel 290710	MH
DRESSING—TULLE NON-GAUZE—PARAFFIN								
4909J	Dressing 7.6 cm x 7.6 cm	10	1	..	*15.72	5.60	Adaptic 2012	JJ

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for Max. Qty \$	Maximum Recordable Value for Safety Net \$	Brand Name and Manufacturer	
DRESSING with CADEXOMER IODINE								
<u>Note</u> Suitable for yellow sloughy infected and malodorous wounds.								
<u>Note</u> Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.								
4931M	Sachets 3 g, 7	¥1	2	..	64.48	5.60	Iodosorb Powder 66051070	SN
4932N	Tubes 10 g, 4	¥1	2	..	103.73	5.60	Iodosorb Ointment 66051240	SN
4933P	Tubes 20 g, 2	¥1	2	..	102.76	5.60	Iodosorb Ointment 66051230	SN
4935R	Sachets 5 g (6 cm x 4 cm), 5	¥1	2	..	98.05	5.60	Iodosorb 66051330	SN
4936T	Sachets 10 g (8 cm x 6 cm), 3	¥1	2	..	141.63	5.60	Iodosorb 66051340	SN
4937W	Sachets 17 g (10 cm x 8 cm), 2	¥1	..	..	149.26	5.60	Iodosorb 66051360	SN
DRESSING with SILVER								
<u>Authority required</u> For wounds where there is evidence of critical colonisation and for well-assessed chronic wounds that have not responded to conventional dressings.								
<u>Note</u> Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.								
4646M	Hydroactive dressings non-adhesive 10 cm x 10 cm, 5	¥1	..	..	175.89	5.60	Biatain Ag 9622	CT
4647N	Hydroactive dressings adhesive 12.5 cm x 12.5 cm, 5	¥1	..	..	191.29	5.60	Biatain Ag 9632	CT
DRESSING with SILVER								
<u>Authority required</u> For wounds where there is evidence of critical colonisation and for well-assessed chronic wounds that have not responded to conventional dressings.								
<u>Note</u> Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.								
4648P	Tulle dressings 10 cm x 10 cm, 3	¥1	..	..	43.78	5.60	Atrauman Ag 499572	HR
GAUZE and COTTON TISSUE (COMBINE ROLL)								
4761N	Wrapped pack 10 cm x 10 m	¥1	..	..	17.21	5.60	JJ 12010	JJ
4767X	Wrapped pack 9 cm x 10 m	¥1	..	..	15.39	5.60	BSN 2902165	BV
GLOVES PLASTIC (DISPOSABLE)								
4772E	Gloves, small, 100	¥1	..	..	12.20	5.60	Handy 4207	BV
4773F	Gloves, medium, 100	¥1	..	..	12.20	5.60	Handy 4208	BV
4774G	Gloves, large, 100	¥1	..	..	12.20	5.60	Handy 4209	BV
TAPES—NON-WOVEN RETENTION (POLYACRYLATE)								
4915Q	Roll 2.5 cm x 9.1 m	¥1	..	..	12.87	5.60	Medipore 2961	MM

TAPES—NON-WOVEN RETENTION (POLYACRYLATE)

Note

Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788

Various

Code	Name, Restriction, Manner of Administration and Form	Max. Qty	No. of Rpts	Premium	Dispensed Price for	Maximum Recordable	Brand Name and Manufacturer
					Max. Qty \$	Value for Safety Net \$	
855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.							
4917T	Roll 2.5 cm x 10 m	‡1	..	..	11.04	5.60	Mefix 310250 MH

TAPES—PLASTER ADHESIVE (WITH SILICONE)

Note

Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.

4239D	Roll 2 cm x 3 m	‡1	..	..	21.37	5.60	Mepitac 298300 MH
4240E	Roll 4 cm x 1.5 m	‡1	..	..	21.37	5.60	Mepitac 298400 MH

TAPES—PLASTER ADHESIVE ELASTIC

4780N	Roll 2.5 cm x 2.5 m	‡1	..	..	12.76	5.60	Leukoplast 1071 BV
4781P	Roll 5 cm x 2.5 m	‡1	..	..	18.56	5.60	Leukoplast 1072 BV
4782Q	Roll 7.5 cm x 2.5 m	‡1	..	..	22.13	5.60	Leukoplast 1073 BV

TAPES—PLASTER ADHESIVE HYPOALLERGENIC

4783R	Roll 1.25 cm x 5 m	‡1	..	..	10.34	5.60	Leukopor 2471 BV
4785W	Roll 1.25 cm x 5 m	‡1	..	..	10.62	5.60	Leukosilk 1021 BV
4787Y	Roll 2.5 cm x 5 m	‡1	..	..	13.24	5.60	Leukosilk 1022 BV
4788B	Stretch roll 5 cm x 5 m	‡1	..	..	17.21	5.60	Leukoflex 1124 BV
4789C	Roll 5 cm x 5 m	‡1	..	..	17.05	5.60	Leukosilk 1024 BV
4790D	Roll 5 cm x 5 m	‡1	..	..	16.21	5.60	Leukopor 2474 BV
4794H	Roll 2.5 cm x 5 m	‡1	..	..	12.73	5.60	Leukopor 2472 BV
4848E	Roll (dispenser) 1.9 cm x 5.4 m	‡1	..	..	11.05	5.60	Nexcare Durable Cloth First Aid Tape 799 MM
4849F	Roll (dispenser) 1.9 cm x 7.3 m	‡1	..	..	11.05	5.60	Nexcare Gentle Paper First Aid Tape 789 MM

Section 2

Standard Packs and Prices

NOTE—

Standard packs and prices (including mark-up, but without dispensing fee and dangerous drug fee) are for items against the price of which an asterisk () is shown in Section 1 of the Schedule.*

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$	Manufacturer
4579B	ALPROSTADIL	10 mcg	2@ 25.40	PH
4580C		20 mcg	2@ 32.40	PH
4118R	ALUMINIUM HYDROXIDE with MAGNESIUM HYDROXIDE and SIMETHICONE	400 mg-400 mg-30 mg per 5 mL, 500 mL	1@ 8.11	JT
4453J		400 mg-400 mg-40 mg	100@ 19.85	JT
4598B	BANDAGE—COMPRESSION	Four layer	1@ 27.85	SN
4654Y		8 cm x 5 m	1@ 14.27	BV
4656C		7.5 cm x 3.5 m	1@ 12.39	SS
4657D		10 cm x 3.5 m	1@ 14.43	SS
4658E		Four layer	1@ 41.71	SN
4736G		7.5 cm x 3 m	1@ 17.58	BV
4748X		10 cm x 3 m	1@ 24.04	BV
4941C		Two layer, 18 cm-22 cm	1@ 41.92	SN
4942D		Two layer, 22 cm-28 cm	1@ 41.92	SN
4943E		Two layer, 28 cm-32 cm	1@ 41.92	SN
4659F	BANDAGE—RETENTION—COHESIVE—HEAVY	7.5 cm x 3 m	1@ 6.82	BV
4660G		10 cm x 2 m	1@ 6.47	MM
4811F		5 cm x 1.3 m	1@ 3.80	BK
4812G		7.5 cm x 1.3 m	1@ 5.44	BK
4813H		10 cm x 1.3 m	1@ 7.31	BK
4814J		15 cm x 1.3 m	1@ 10.88	BK
4662J	BANDAGE—RETENTION—COHESIVE—LIGHT	10 cm x 4 m	1@ 5.36	BV
4719J		6 cm x 4 m	1@ 4.14	BV
4727T	BANDAGE—RETENTION—COTTON CREPE	5 cm x 2.3 m	1@ 5.93	BV
4728W		7.5 cm x 2.3 m	1@ 7.99	BV
4729X		10 cm x 2.3 m	1@ 10.70	BV
4674B	BANDAGE—TUBULAR (LONG STOCKING)	Small	1@ 16.88	SS
4675C		XX/large	1@ 16.88	SS
4797L		Medium	1@ 16.88	SS
4799N		Large	1@ 16.88	SS
4661H	BANDAGE—TUBULAR (SHORT STOCKING)	Small B/C	1@ 12.01	SS
4815K		Medium C/D	1@ 12.01	SS
4816L		Large D/E	1@ 12.01	SS
4668Q	BANDAGE—ZINC PASTE	7.5 cm x 6 m	1@ 11.39	SS
4669R		7.5 cm x 6 m	1@ 11.62	XP
4670T		10 cm x 9.1 m	1@ 11.18	CC
4750B		7.5 cm x 6 m	1@ 32.11	SN
4150K	BROMAZEPAM	3 mg	30@ 10.48	RO
4151L		6 mg	30@ 13.49	RO
4094L	CALCIUM	500 mg	60@ 6.01	IA
4142B		600 mg	120@ 7.89	PP
4333C		500 mg	60@ 6.01	IA
4055K	CALCIUM CARBONATE with GLYCINE	420 mg-180 mg	100@ 8.38	MM
4081T	DEXTROPROPOXYPHENE NAPSYLATE	100 mg	10@ 3.17	AS
4681J	DRESSING—ACTIVATED CHARCOAL (MALODOROUS WOUND)	10.5 cm x 10.5 cm	1@ 9.45	JJ
4682K	DRESSING—ALGINATE (CAVITY WOUND)	2 g (40 cm), 6	1@ 65.65	CT
4832H		2 g	5@ 54.42	CC
4684M	DRESSING—ALGINATE (SUPERFICIAL WOUND)	5 cm x 5 cm	1@ 4.05	CT
4831G		10 cm x 10 cm	1@ 7.80	UM
4688R	DRESSING—FILM	15 cm x 20 cm	1@ 4.04	MM
4689T	DRESSING—FILM ISLAND	5 cm x 7 cm	1@ .98	MM
4690W		9 cm x 10 cm	1@ 2.13	MM
4898T		5 cm x 7.2 cm, 5	1@ 11.25	SN
4899W		8 cm x 10 cm, 5	1@ 19.54	SN
4892L	DRESSING—FOAM with CHARCOAL (MALODOROUS WOUND)	10 cm x 10 cm, 10	1@ 83.82	SS
4919X	DRESSING—HYDROACTIVE (CAVITY WOUND)	10 cm x 10 cm, 5	1@ 85.97	SN
4906F	DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—LIGHT EXUDATE)	10 cm x 10 cm, 5	1@ 48.54	SN
4886E	DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—MODERATE EXUDATE)	10 cm x 10 cm, 5	1@ 34.78	SN
4895P	DRESSING—HYDROCOLLOID (CAVITY WOUND)	50 g	1@ 18.40	CT
4896Q		30 g	1@ 13.87	CC
4678F	DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—MODERATE EXUDATE)	7 cm	1@ 9.75	CT
4679G		10 cm	1@ 10.64	CT
4897R		10 cm x 10 cm, 5	1@ 37.49	CC
4920Y		20 cm x 20 cm, 5	1@ 107.95	CC
4922C	DRESSING—HYDROFIBRE (ALTERNATE TO ALGINATES)	15 cm x 15 cm, 5	1@ 101.14	CC
4599C	DRESSING—HYDROGEL—AMORPHOUS	50 g	1@ 8.03	SN
4894N		25 g	1@ 13.28	SN

(APPLY WASTAGE FACTOR IN CALCULATING BROKEN QUANTITY PRICES)

Code	Name	Form/Strength	Pack and Price \$		Manufacturer
4913N		30 g, 3	1@	30.23	CC
4914P		50 g	1@	8.90	JJ
4806Y	DRESSING—HYDROGEL—SHEET	10 cm x 10 cm, 5	1@	23.43	HR
4911L		9.5 cm x 10.2 cm, 5	1@	38.39	JJ
4819P	DRESSING—NON-ADHERENT	5 cm x 5 cm, 5	1@	4.69	SN
4860T		5 cm x 5 cm, 5	1@	5.03	SN
4862X		10 cm x 10 cm, 5	1@	9.49	SN
4909J	DRESSING—TULLE NON-GAUZE—PARAFFIN	7.6 cm x 7.6 cm	1@	.93	JJ
4237B	FEXOFENADINE HYDROCHLORIDE	60 mg	20@	16.19	SW
4246L	GLYCEROL	2.8 g, 12	1@	4.44	PP
6099B	MORPHINE SULFATE		20@	82.15	MF
4571N	NICOTINE	Approx. 7 mg per 24 hours, 7	1@	22.48	AF
4572P		Approx. 14 mg per 24 hours, 7	1@	31.16	GC
4573Q		Approx. 21 mg per 24 hours, 7	1@	31.16	GC
4576W		Approx. 5 mg per 16 hours, 7	1@	22.20	JT
4577X		Approx. 10 mg per 16 hours, 7	1@	24.18	JT
4578Y		Approx. 15 mg per 16 hours, 7	1@	26.77	JT

GENERIC/PROPRIETARY INDEX

3

3TC (VI)

.Section 100 565, 701

A

ABACAVIR

.Section 100 563, 700

ABACAVIR SULFATE with LAMIVUDINE and ZIDOVUDINE

.Section 100 567, 703

ABACAVIR with LAMIVUDINE

.Section 100 567, 704

ABATACEPT

.Section 100 576, 713

Abbocillin-V (SI)

.Antifungives for systemic use 179

.Dental 494

Abbocillin-VK Filmtab (SI)

.Antifungives for systemic use 178, 179

.Dental 493

ABCIXIMAB

.Blood and blood forming organs 98

Abilify (BQ)

.Nervous system 404

Abraxane (TS)

.Antineoplastic and immunomodulating agents 205

ACAMPROSATE CALCIUM

.Nervous system 428

ACARBOSE

.Alimentary tract and metabolism 92

Accomin Adult Tonic (WT)

.Repatriation Schedule 839

Accu-Chek Active (RD)

.Various 455, 456

Accu-Chek Advantage/Sensor Comfort (RD)

.Various 455, 456

Accu-Chek Go (RD)

.Various 455, 456

Accu-Chek Integra (RD)

.Various 455, 456

Accu-Chek Mobile (RD)

.Various 457

Accu-Chek Performa (RD)

.Various 455, 456

Accupril (PF)

.Cardiovascular system 125, 126

Accuretic 10/12.5mg (PF)

.Cardiovascular system 129

Accuretic 20/12.5mg (PF)

.Cardiovascular system 129

ACETAZOLAMIDE

.Sensory organs 442

Aci-Jel (CU)

.Repatriation Schedule 847

ACICLOVIR

.Antifungives for systemic use 194, 195

.Sensory organs 440

.Optometrical 516

Aciclovir 200 (CR)

.Antifungives for systemic use 194

Aciclovir 800 (CR)

.Antifungives for systemic use 195

Acihexal (SZ)

.Antifungives for systemic use 194, 195

Acimax Tablets (AL)

.Alimentary tract and metabolism 72, 73

ACITRETIN

.Dermatologicals 151

Aclasta (NV)

.Musculo-skeletal system 368

Aclin (AF)

.Musculo-skeletal system 359

.Palliative Care 477

.Dental 501

Aclin 200 (AF)

.Musculo-skeletal system 359

.Palliative Care 477

.Dental 502

Aclor 125 (SI)

.Antifungives for systemic use 183

.Dental 497

Aclor 250 (SI)

.Antifungives for systemic use 183

.Dental 497

Acquin 10 (SI)

.Cardiovascular system 125

Acquin 20 (SI)

.Cardiovascular system 126

Acquin 5 (SI)

.Cardiovascular system 125

Actemra (RO)

.Section 100 639, 776

Actilax (AF)

.Alimentary tract and metabolism 80

.Palliative Care 472

Actiq (OA)

.Palliative Care 481

Actisorb Plus MAC031 (JJ)

.Repatriation Schedule 866

Actonel (SW)

.Musculo-skeletal system 367

.Repatriation Schedule 852

Actonel Combi (SW)

.Musculo-skeletal system 371

.Repatriation Schedule 852

Actonel Combi D (SW)

.Musculo-skeletal system 371

.Repatriation Schedule 852

Actonel Once-a-Month (SW)

.Musculo-skeletal system 367

Actonel Once-a-Week (SW)

.Musculo-skeletal system 367

.Repatriation Schedule 852

Actos (LY)

.Alimentary tract and metabolism 92, 93

Actrapid (NO)

.Alimentary tract and metabolism 87

Actrapid Penfill 3 mL (NO)

.Alimentary tract and metabolism 87

Acycto-V 200 (AF)

.Antifungives for systemic use	194	.Respiratory system	433
<i>Acyclo-V 800 (AF)</i>		<i>Airomir Autohaler (IA)</i>	
.Antifungives for systemic use	195	.Respiratory system	433
<i>Adalat 10 (BN)</i>		<i>Akamin 100 (AF)</i>	
.Cardiovascular system	119	.Antifungives for systemic use	177
<i>Adalat 20 (BN)</i>		<i>Akamin 50 (AF)</i>	
.Cardiovascular system	120	.Antifungives for systemic use	177
<i>Adalat Oros 20mg (BN)</i>		<i>Akineton (LM)</i>	
.Cardiovascular system	120	.Nervous system	395
<i>Adalat Oros 30 (BN)</i>		<i>Albalon Liquifilm (AG)</i>	
.Cardiovascular system	120	.Repatriation Schedule	859
<i>Adalat Oros 60 (BN)</i>		<i>Albalon-A (AG)</i>	
.Cardiovascular system	120	.Repatriation Schedule	859
ADALIMUMAB		ALBENDAZOLE	
.Antineoplastic and immunomodulating agents 238, 242, 245, 248, 251, 254, 257, 263, 269, 274, 279, 282		.Antiparasitic products, insecticides and repellents ...	431, 432
.Section 100	582, 719	<i>Albex Bee Venom (HL)</i>	
<i>Adaptic 2012 (JJ)</i>		.Various	452
.Repatriation Schedule	873	<i>Albex Paper Wasp Venom (HL)</i>	
<i>add-ins (SB)</i>		.Various	452
.Various	463	<i>Albex Yellow Jacket Venom (HL)</i>	
<i>Addos XR 30 (SI)</i>		.Various	452
.Cardiovascular system	120	<i>Aldactone (PF)</i>	
<i>Addos XR 60 (SI)</i>		.Cardiovascular system	113
.Cardiovascular system	120	<i>Aldara (IA)</i>	
<i>Adefin 10 (AF)</i>		.Dermatologicals	155
.Cardiovascular system	119	.Repatriation Schedule	846
<i>Adefin 20 (AF)</i>		<i>Aldomet (AS)</i>	
.Cardiovascular system	119	.Cardiovascular system	110
<i>Adefin XL 30 (AF)</i>		<i>Alendro Once Weekly (SI)</i>	
.Cardiovascular system	120	.Musculo-skeletal system	365
<i>Adefin XL 60 (AF)</i>		<i>Alendrobell 70mg (BF)</i>	
.Cardiovascular system	120	.Musculo-skeletal system	365
ADEFOVIR DIPIVOXIL		<i>Alendronate Sandoz (SZ)</i>	
.Section 100	563, 700	.Musculo-skeletal system	365
ADRENALINE		ALENDRONATE SODIUM	
.Cardiovascular system	108, 109	.Musculo-skeletal system	365
.Respiratory system	437	ALENDRONATE SODIUM with COLECALCIFEROL	
.Dental	489, 511	.Musculo-skeletal system	368, 369
<i>Adriamycin (PF)</i>		ALENDRONATE SODIUM with COLECALCIFEROL and CALCIUM CARBONATE	
.Antineoplastic and immunomodulating agents	206	.Musculo-skeletal system	369
<i>Adriamycin Solution (PF)</i>		<i>Alendronate-GA (GM)</i>	
.Antineoplastic and immunomodulating agents	206	.Musculo-skeletal system	365
<i>Adronat (AF)</i>		<i>Alepam 15 (AF)</i>	
.Musculo-skeletal system	365	.Nervous system	409
<i>ADT Booster (CS)</i>		.Palliative Care	484
.Emergency Drug Supplies	64	.Dental	510
.Antifungives for systemic use	198	<i>Alepam 30 (AF)</i>	
<i>Advantage II (RD)</i>		.Nervous system	409
.Various	455, 456	.Palliative Care	484
<i>Advantan (CS)</i>		.Dental	510
.Dermatologicals	153	<i>Alfaré (NT)</i>	
<i>Aeron 250 (SI)</i>		.Various	460
.Respiratory system	436	<i>Algisite M 66000519 (SN)</i>	
<i>Aeron 500 (SI)</i>		.Repatriation Schedule	867
.Respiratory system	436	<i>Algisite M 66000520 (SN)</i>	
<i>Aggrastat (AS)</i>		.Repatriation Schedule	867
.Blood and blood forming organs	101	<i>Algisite M 66000521 (SN)</i>	
<i>Airomir (IA)</i>		.Repatriation Schedule	867
.Emergency Drug Supplies	65		

<i>Alimta (LY)</i>		
.Antineoplastic and immunomodulating agents	201	
<i>Alkeran (AS)</i>		
.Antineoplastic and immunomodulating agents	199	
<i>ALLANTOIN with GLYCEROL and ICHTHAMMOL</i>		
.Repatriation Schedule	846	
<i>ALLANTOIN with SULFUR, PHENOL, COAL TAR SOLUTION and MENTHOL</i>		
.Repatriation Schedule	844	
<i>Allegron (AS)</i>		
.Nervous system	412	
<i>Allereze (AF)</i>		
.Repatriation Schedule	858	
<i>Allevyn 66007637 (SN)</i>		
.Repatriation Schedule	867	
<i>Allevyn Adhesive 66000044 (SN)</i>		
.Repatriation Schedule	868	
<i>Allevyn Plus Cavity 66047571 (SN)</i>		
.Repatriation Schedule	869	
<i>Allevyn Plus Cavity 66047573 (SN)</i>		
.Repatriation Schedule	869	
<i>Allevyn Thin 66047576 (SN)</i>		
.Repatriation Schedule	870	
<i>Allevyn Thin 66047578 (SN)</i>		
.Repatriation Schedule	870	
<i>ALLOPURINOL</i>		
.Musculo-skeletal system	364	
<i>Allopurinol Sandoz (SZ)</i>		
.Musculo-skeletal system	364	
<i>Allosig (FM)</i>		
.Musculo-skeletal system	364	
<i>Alodorm (AF)</i>		
.Nervous system	388, 409, 410	
.Palliative Care	484, 485	
.Dental	510	
<i>Aloxi (TS)</i>		
.Alimentary tract and metabolism	78	
<i>Alpha Keri Bath Oil (MT)</i>		
.Repatriation Schedule	843	
<i>Alpha Keri Lotion (MT)</i>		
.Repatriation Schedule	843	
<i>Alphagan (AG)</i>		
.Sensory organs	442	
.Optometrical	516	
<i>Alphagan P 1.5 (AG)</i>		
.Sensory organs	441	
.Optometrical	517	
<i>Alphamox 125 (AF)</i>		
.Antiinfectives for systemic use	177	
.Dental	493	
<i>Alphamox 250 (AF)</i>		
.Antiinfectives for systemic use	177	
.Dental	492, 493	
<i>Alphamox 500 (AF)</i>		
.Antiinfectives for systemic use	178	
.Dental	492	
<i>Alphapress 25 (AF)</i>		
.Cardiovascular system	111	
<i>Alphapress 50 (AF)</i>		
.Cardiovascular system	111	
<i>Alphapril (AF)</i>		
.Cardiovascular system	122	
<i>Alprax 0.25 (SI)</i>		
.Nervous system	408	
<i>Alprax 0.5 (SI)</i>		
.Nervous system	408	
<i>Alprax 1 (SI)</i>		
.Nervous system	408	
<i>Alprax 2 (SI)</i>		
.Nervous system	408	
<i>ALPRAZOLAM</i>		
.Nervous system	407	
<i>Alprazolam Sandoz (SZ)</i>		
.Nervous system	408	
<i>Alprazolam-GA (GM)</i>		
.Nervous system	408	
<i>Alprim (AF)</i>		
.Antiinfectives for systemic use	184	
<i>ALPROSTADIL</i>		
.Repatriation Schedule	847	
<i>ALUMINIUM HYDROXIDE with MAGNESIUM HYDROXIDE</i>		
.Alimentary tract and metabolism	68	
<i>ALUMINIUM HYDROXIDE with MAGNESIUM HYDROXIDE and SIMETHICONE</i>		
.Repatriation Schedule	837	
<i>ALUMINIUM HYDROXIDE with MAGNESIUM TRISILICATE and MAGNESIUM HYDROXIDE</i>		
.Alimentary tract and metabolism	68	
<i>Alvesco 160 (NQ)</i>		
.Respiratory system	436	
<i>Alvesco 80 (NQ)</i>		
.Respiratory system	436	
<i>Alzene (AF)</i>		
.Repatriation Schedule	858	
<i>AMANTADINE HYDROCHLORIDE</i>		
.Nervous system	397	
<i>Amaryl (SW)</i>		
.Alimentary tract and metabolism	90	
<i>AMBRISENTAN</i>		
.Section 100	526, 658	
<i>AMILORIDE HYDROCHLORIDE</i>		
.Cardiovascular system	113	
<i>AMINO ACID FORMULA with FAT, CARBOHYDRATE, VITAMINS, MINERALS and TRACE ELEMENTS without PHENYLALANINE and TYROSINE, and supplemented with DOCOSAHEXANOIC ACID</i>		
.Various	462	
<i>AMINO ACID FORMULA with FAT, CARBOHYDRATE, VITAMINS, MINERALS, and TRACE ELEMENTS, without METHIONINE and supplemented with DOCOSAHEXANOIC ACID</i>		
.Various	462	
<i>AMINO ACID FORMULA with VITAMINS and MINERALS without LYSINE and low in TRYPTOPHAN</i>		
.Various	462, 463	
<i>AMINO ACID FORMULA with VITAMINS and MINERALS without METHIONINE</i>		
.Various	463	

<i>AMINO ACID FORMULA with VITAMINS and MINERALS without METHIONINE, THREONINE and VALINE and low in ISOLEUCINE</i>	
.Various.....	463
<i>AMINO ACID FORMULA with VITAMINS and MINERALS without PHENYLALANINE</i>	
.Various.....	463
<i>AMINO ACID FORMULA with VITAMINS and MINERALS without PHENYLALANINE and TYROSINE</i>	
.Various.....	464
<i>AMINO ACID FORMULA with VITAMINS and MINERALS without VALINE, LEUCINE and ISOLEUCINE</i>	
.Various.....	464
<i>AMINO ACID FORMULA with VITAMINS and MINERALS without VALINE, LEUCINE and ISOLEUCINE with FAT, CARBOHYDRATE and TRACE ELEMENTS and supplemented with DOCOSAHEXANOIC ACID</i>	
.Various.....	464
<i>AMINO ACID FORMULA with VITAMINS, MINERALS and LONG CHAIN POLYUNSATURATED FATTY ACIDS without PHENYLALANINE</i>	
.Various.....	462
<i>AMINO ACID FORMULA without PHENYLALANINE</i>	
.Various.....	462
<i>AMINO ACID SYNTHETIC FORMULA supplemented with LONG CHAIN POLYUNSATURATED FATTY ACIDS</i>	
.Various.....	458, 459
<i>AMINO ACIDS—SYNTHETIC, FORMULA</i>	
.Various.....	457, 458
<i>AMIODARONE HYDROCHLORIDE</i>	
.Cardiovascular system.....	108
<i>Amipride 400 (SI)</i>	
.Nervous system.....	403
<i>Amira 150 (AF)</i>	
.Nervous system.....	416
<i>Amira 300 (AF)</i>	
.Nervous system.....	416
<i>AMISULPRIDE</i>	
.Nervous system.....	403
<i>Amisulpride 100 Winthrop (WA)</i>	
.Nervous system.....	403
<i>Amisulpride 200 Winthrop (WA)</i>	
.Nervous system.....	403
<i>Amisulpride 400 Winthrop (WA)</i>	
.Nervous system.....	403
<i>Amisulpride Sandoz (SZ)</i>	
.Nervous system.....	403
<i>AMITRIPTYLINE HYDROCHLORIDE</i>	
.Nervous system.....	411
<i>Amlo 10 (ZP)</i>	
.Cardiovascular system.....	118
<i>Amlo 5 (ZP)</i>	
.Cardiovascular system.....	118
<i>AMLODIPINE</i>	
.Cardiovascular system.....	118
<i>AMLODIPINE BESYLATE with ATORVASTATIN CALCIUM</i>	
.Cardiovascular system.....	148
<i>Amlodipine generichealth (GQ)</i>	
.Cardiovascular system.....	118, 119
<i>Amlodipine Sandoz (SZ)</i>	
.Cardiovascular system.....	118, 119
<i>AMLODIPINE with VALSARTAN</i>	
.Cardiovascular system.....	132
<i>AMLODIPINE with VALSARTAN and HYDROCHLOROTHIAZIDE</i>	
.Cardiovascular system.....	133
<i>Amlodipine-DRLA (RZ)</i>	
.Cardiovascular system.....	118
<i>Amlodipine-GA (GM)</i>	
.Cardiovascular system.....	118, 119
<i>AMOROLFINE HYDROCHLORIDE</i>	
.Repatriation Schedule.....	842
<i>Amoxil (GK)</i>	
.Antiinfectives for systemic use.....	66, 67, 177, 178
.Dental.....	492, 493
<i>Amoxil Forte (GK)</i>	
.Antiinfectives for systemic use.....	178
.Dental.....	493
<i>AMOXYCILLIN</i>	
.Antiinfectives for systemic use.....	66, 177, 178
.Dental.....	492
<i>Amoxycillin Ranbaxy (RA)</i>	
.Antiinfectives for systemic use.....	177, 178
.Dental.....	492
<i>Amoxycillin Sandoz (BG)</i>	
.Antiinfectives for systemic use.....	178
<i>Amoxycillin Sandoz (SZ)</i>	
.Antiinfectives for systemic use.....	177, 178
.Dental.....	492, 493
<i>AMOXYCILLIN with CLAVULANIC ACID</i>	
.Antiinfectives for systemic use.....	180
.Dental.....	494
<i>Amoxycillin-GA (GM)</i>	
.Antiinfectives for systemic use.....	177, 178
.Dental.....	492
<i>Amoxycillin/ Clavulanic Acid 500/125 generichealth (GQ)</i>	
.Antiinfectives for systemic use.....	180
.Dental.....	495
<i>Amoxycillin/ Clavulanic Acid 875/125 generichealth (GQ)</i>	
.Antiinfectives for systemic use.....	180
.Dental.....	494
<i>AMPHOTERICIN</i>	
.Alimentary tract and metabolism.....	68
.Dental.....	486
<i>AMPICILLIN</i>	
.Antiinfectives for systemic use.....	178
.Dental.....	493
<i>AMYLOPECTIN, MODIFIED LONG CHAIN</i>	
.Various.....	461
<i>Anafranil 25 (NV)</i>	
.Nervous system.....	409, 411
<i>Anamorph (FM)</i>	
.Nervous system.....	378
.Dental.....	505
<i>Anandron (SW)</i>	
.Antineoplastic and immunomodulating agents.....	233
<i>Anapen (LM)</i>	
.Cardiovascular system.....	109
.Respiratory system.....	437
<i>Anapen Junior (LM)</i>	

.Cardiovascular system	109
.Respiratory system	437
<i>Anaprox 550 (RO)</i>	
.Musculo-skeletal system	361
.Palliative Care	478, 479
.Dental	503
<i>ANASTROZOLE</i>	
.Antineoplastic and immunomodulating agents	233
<i>Andriol Testocaps (SH)</i>	
.Genito urinary system and sex hormones	159
<i>Androcur (SC)</i>	
.Genito urinary system and sex hormones	164
.Antineoplastic and immunomodulating agents	232
<i>Androcur-100 (SC)</i>	
.Genito urinary system and sex hormones	164
.Antineoplastic and immunomodulating agents	232
<i>Androderm (HH)</i>	
.Genito urinary system and sex hormones	159
<i>Anginine Stabilised (SI)</i>	
.Cardiovascular system	109
.Dental	489
<i>Angiomax (CS)</i>	
.Blood and blood forming organs	102
<i>Anpec 40 (AF)</i>	
.Cardiovascular system	120
<i>Anpec 80 (AF)</i>	
.Cardiovascular system	120
<i>ANTAZOLINE with NAPHAZOLINE</i>	
.Repatriation Schedule	859
<i>Antenex 2 (AF)</i>	
.Nervous system	408
.Palliative Care	483, 484
.Dental	509
<i>Antenex 5 (AF)</i>	
.Nervous system	408
.Palliative Care	483, 484
.Dental	509
<i>Anthel 125 (AF)</i>	
.Antiparasitic products, insecticides and repellents	432
<i>Anthel 250 (AF)</i>	
.Antiparasitic products, insecticides and repellents	432
<i>Antistine-Privine (NV)</i>	
.Repatriation Schedule	859
<i>Antroquoril (EX)</i>	
.Dermatologicals	152, 153
<i>Anusol (JT)</i>	
.Repatriation Schedule	841
<i>Anzatax (HH)</i>	
.Antineoplastic and immunomodulating agents .	205, 206
<i>Anzemet (SW)</i>	
.Alimentary tract and metabolism	76
<i>Apidra (AV)</i>	
.Alimentary tract and metabolism	87
<i>Apidra (SW)</i>	
.Alimentary tract and metabolism	87
<i>Apidra SoloStar (SW)</i>	
.Alimentary tract and metabolism	87
<i>APO- Paracetamol/Codeine 500/30 (TX)</i>	
.Nervous system	375
.Dental	504
<i>APO-Alendronate (TX)</i>	
.Musculo-skeletal system	365
<i>APO-Amlodipine (TX)</i>	
.Cardiovascular system	118, 119
<i>APO-Amoxycillin (TX)</i>	
.Antiinfectives for systemic use	177, 178
.Dental	492
<i>APO-Amoxycillin/ Clavulanic Acid 500/125 (TX)</i>	
.Antiinfectives for systemic use	180
.Dental	495
<i>APO-Atenolol (TX)</i>	
.Cardiovascular system	115
<i>APO-Bicalutamide (TX)</i>	
.Antineoplastic and immunomodulating agents	232
<i>APO-Carvedilol (TX)</i>	
.Cardiovascular system	117
<i>APO-Cephalexin (TX)</i>	
.Antiinfectives for systemic use	181
.Dental	496
<i>APO-Citalopram (TX)</i>	
.Nervous system	412
<i>APO-Clarithromycin (TX)</i>	
.Antiinfectives for systemic use	185
<i>APO-Clopidogrel (TX)</i>	
.Blood and blood forming organs	99
<i>APO-Clotrimazole 3 Day Cream (TX)</i>	
.Repatriation Schedule	847
<i>APO-Clotrimazole 6 Day Cream (TX)</i>	
.Repatriation Schedule	847
<i>APO-Diclofenac (TX)</i>	
.Musculo-skeletal system	358
.Palliative Care	475, 476
.Dental	501
<i>APO-Escitalopram (TX)</i>	
.Nervous system	412, 413
<i>APO-Famciclovir (TX)</i>	
.Antiinfectives for systemic use	195, 196
<i>APO-Fluconazole (TX)</i>	
.Antiinfectives for systemic use	191
<i>APO-Fluvoxamine (TX)</i>	
.Nervous system	414
<i>APO-Fosinopril HCTZ 10/12.5 (TX)</i>	
.Cardiovascular system	128
<i>APO-Fosinopril HCTZ 20/12.5 (TX)</i>	
.Cardiovascular system	129
<i>APO-Gabapentin (TX)</i>	
.Nervous system	391
<i>APO-Gliclazide MR (TX)</i>	
.Alimentary tract and metabolism	89
<i>APO-Glimepiride (TX)</i>	
.Alimentary tract and metabolism	90
<i>APO-go (HH)</i>	
.Section 100	650, 787
<i>APO-Ipratropium (TX)</i>	
.Respiratory system	436
<i>APO-Lamotrigine (TX)</i>	
.Nervous system	391, 392
<i>APO-Lansoprazole (TX)</i>	
.Alimentary tract and metabolism	71, 72
<i>APO-Lercanidipine (TX)</i>	

.Cardiovascular system	119	<i>Apomine PFS (HH)</i>	
<i>APO-Levetiracetam (TX)</i>		.Section 100	650, 787
.Nervous system	393	<i>APOMORPHINE HYDROCHLORIDE</i>	
<i>APO-Lisinopril (TX)</i>		.Section 100	650, 787
.Cardiovascular system	123, 124	<i>APRACLONIDINE HYDROCHLORIDE</i>	
<i>APO-Memantine (TX)</i>		.Sensory organs.....	441
.Nervous system	426	<i>APREPITANT</i>	
<i>APO-Mirtazapine (TX)</i>		.Alimentary tract and metabolism.....	78
.Nervous system	418	<i>Aptivus (BY)</i>	
<i>APO-Nifedipine XR (TX)</i>		.Section 100	563, 699, 700
.Cardiovascular system	120	<i>Aquacare H.P. (AG)</i>	
<i>APO-Omeprazole (TX)</i>		.Repatriation Schedule	843
.Alimentary tract and metabolism.....	72, 73	<i>Aquacel 177902 (CC)</i>	
<i>APO-Ondansetron (TX)</i>		.Repatriation Schedule	872
.Alimentary tract and metabolism.....	77	<i>Aquacel 177903 (CC)</i>	
<i>APO-Oxazepam (TX)</i>		.Repatriation Schedule	872
.Nervous system	409	<i>Aquacel 177904 (CC)</i>	
.Palliative Care	484	.Repatriation Schedule	872
.Dental	510	<i>Aquaclear 900796 (HR)</i>	
<i>APO-Pantoprazole (TX)</i>		.Repatriation Schedule	873
.Alimentary tract and metabolism.....	73, 74	<i>Aquae (HA)</i>	
<i>APO-Paracetamol (TX)</i>		.Palliative Care	469
.Nervous system	384	.Repatriation Schedule	837
.Dental	508, 509	<i>Aquae Gel (HA)</i>	
<i>APO-Perindopril (TX)</i>		.Palliative Care	469, 470
.Cardiovascular system	124, 125	<i>Aquasun Lotion SPF 18 (PF)</i>	
<i>APO-Pravastatin (TX)</i>		.Repatriation Schedule	844
.Cardiovascular system	138, 139, 140	<i>Arabloc (AV)</i>	
<i>APO-Prochlorperazine (TX)</i>		.Antineoplastic and immunomodulating agents	237
.Alimentary tract and metabolism.....	79	.Musculo-skeletal system	363
.Dental	486	<i>Aranesp (AN)</i>	
<i>APO-Quinapril (TX)</i>		.Section 100	524, 656
.Cardiovascular system	125, 126	<i>Aranesp SureClick (AN)</i>	
<i>APO-Ramipril (TX)</i>		.Section 100	524, 656
.Cardiovascular system	126, 127	<i>Aratac 100 (AF)</i>	
<i>APO-Risperidone (TX)</i>		.Cardiovascular system	108
.Nervous system	405, 406, 407	<i>Aratac 200 (AF)</i>	
<i>APO-Roxithromycin (TX)</i>		.Cardiovascular system	108
.Antiinfectives for systemic use.....	186	<i>Arava (SW)</i>	
.Dental	498	.Antineoplastic and immunomodulating agents	237
<i>APO-Simvastatin (TX)</i>		.Musculo-skeletal system	363
.Cardiovascular system.....	140, 141, 142, 143, 144	<i>Aredia 15 mg (NV)</i>	
<i>APO-Sumatriptan (TX)</i>		.Musculo-skeletal system	366
.Nervous system	386	.Section 100	648, 785
<i>APO-Temazepam (TX)</i>		<i>Aredia 30 mg (NV)</i>	
.Nervous system	410	.Musculo-skeletal system	366
.Palliative Care	485	.Section 100	648, 785
.Dental	510	<i>Aredia 90 mg (NV)</i>	
<i>APO-Topiramate (TX)</i>		.Section 100	649, 786
.Nervous system	394, 395	<i>Arginine Amino Acid Supplement (VF)</i>	
<i>APO-Tramadol (TX)</i>		.Various.....	465
.Nervous system	382	<i>ARGININE with CARBOHYDRATE</i>	
.Dental	507	.Various.....	464
<i>APO-Tramadol SR (TX)</i>		<i>Aricept (PF)</i>	
.Nervous system	383	.Nervous system	422
.Dental	507, 508	<i>Arimidex (AP)</i>	
<i>APO-Trandolapril (TX)</i>		.Antineoplastic and immunomodulating agents	233
.Cardiovascular system	128	<i>ARIPIPRAZOLE</i>	
<i>Apomine (HH)</i>		.Nervous system	403
.Section 100	650, 787	<i>Aristocort 0.02% (SI)</i>	

.Dermatologicals.....	152	<i>Atacand Plus 32/12.5 (AP)</i>	
<i>Arixtra (GK)</i>		.Cardiovascular system	131
.Blood and blood forming organs	103	<i>Atacand Plus 32/25 (AP)</i>	
<i>Aromasin (PF)</i>		.Cardiovascular system	131
.Antineoplastic and immunomodulating agents	233	<i>ATAZANAVIR</i>	
<i>Aropax (GK)</i>		.Section 100	562, 698
.Nervous system	415	<i>ATENOLOL</i>	
<i>ARSENIC TRIOXIDE</i>		.Cardiovascular system	115
.Antineoplastic and immunomodulating agents	226	<i>Atenolol Sandoz (SZ)</i>	
<i>Artane (SI)</i>		.Cardiovascular system	115
.Nervous system	395	<i>Atenolol-GA (GN)</i>	
<i>ARTEMETHER with LUMEFANTRINE</i>		.Cardiovascular system	115
.Antiparasitic products, insecticides and repellents	431	<i>ATOMOXETINE HYDROCHLORIDE</i>	
<i>Arthrexin (AF)</i>		.Nervous system	419
.Musculo-skeletal system	358	<i>ATORVASTATIN</i>	
.Palliative Care	476	.Cardiovascular system	137
.Dental	501	<i>ATOVAQUONE</i>	
<i>Arthrotec 50 (PH)</i>		.Antiparasitic products, insecticides and repellents	430
.Repatriation Schedule	852	<i>ATOVAQUONE with PROGUANIL HYDROCHLORIDE</i>	
<i>Asasantin SR (BY)</i>		.Antiparasitic products, insecticides and repellents	430
.Blood and blood forming organs	100	<i>Atrauman 499513 (HR)</i>	
<i>Ascent Pharma Pty Ltd (GM)</i>		.Repatriation Schedule	873
.Cardiovascular system	121	<i>Atrauman Ag 499572 (HR)</i>	
<i>Ascent Pharmaceuticals Limited (GN)</i>		.Repatriation Schedule	874
.Alimentary tract and metabolism	88	<i>Atripia (GI)</i>	
.Antifectives for systemic use	187, 188	.Section 100	568, 704
<i>Asmol 2.5 uni-dose (AF)</i>		<i>ATROPINE SULFATE</i>	
.Emergency Drug Supplies	65	.Sensory organs.....	443
.Respiratory system	433	<i>Atropt (SI)</i>	
<i>Asmol 5 uni-dose (AF)</i>		.Sensory organs.....	443
.Emergency Drug Supplies	65	<i>Atrovent (BY)</i>	
.Respiratory system	434	.Respiratory system	436
<i>Asmol CFC-free (AL)</i>		<i>Atrovent Adult (BY)</i>	
.Emergency Drug Supplies	65	.Respiratory system	436
.Respiratory system	433	<i>Atrovent Nasal Aqueous (BY)</i>	
<i>Aspalgin (FM)</i>		.Repatriation Schedule	857
.Repatriation Schedule	853	<i>Atrovent Nasal Forte (BY)</i>	
<i>Aspen Ampicyn (AS)</i>		.Repatriation Schedule	857
.Antifectives for systemic use	178	<i>Augmentin (GK)</i>	
.Dental	493	.Antifectives for systemic use	180
<i>Aspen Pharmacare Australia Pty Limited (LN)</i>		.Dental	495
.Antifectives for systemic use	180	<i>Augmentin Duo (GK)</i>	
.Dental	494	.Antifectives for systemic use	180
<i>ASPIRIN</i>		.Dental	495
.Blood and blood forming organs	98	<i>Augmentin Duo 400 (GK)</i>	
.Nervous system	384	.Antifectives for systemic use	181
.Dental	508	.Dental	495
.Repatriation Schedule	840	<i>Augmentin Duo forte (GK)</i>	
<i>AstraZeneca Pty Ltd (AP)</i>		.Antifectives for systemic use	180
.Emergency Drug Supplies	64	.Dental	495
.Cardiovascular system	108	<i>AURANOFIN</i>	
.Respiratory system	437	.Musculo-skeletal system	362
.Dental	489, 511	<i>Aurorix (RO)</i>	
<i>Astrix (YN)</i>		.Nervous system	416
.Blood and blood forming organs	98	<i>Aurorix 300 mg (RO)</i>	
.Repatriation Schedule	840	.Nervous system	417
<i>Atacand (AP)</i>		<i>Auscap (SI)</i>	
.Cardiovascular system	130	.Nervous system	414
<i>Atacand Plus 16/12.5 (AP)</i>		<i>Ausfam 20 (SI)</i>	
.Cardiovascular system	131	.Alimentary tract and metabolism	68

<i>Ausfam 40 (SI)</i>	
.Alimentary tract and metabolism.....	69
<i>Ausgem (SI)</i>	
.Cardiovascular system.....	145
<i>Auspril (SI)</i>	
.Cardiovascular system.....	122
<i>Ausran (SI)</i>	
.Alimentary tract and metabolism.....	69
<i>Austrapen (LN)</i>	
.Antifungives for systemic use	178
.Dental	493
<i>Avandamet (GK)</i>	
.Alimentary tract and metabolism.....	91
<i>Avandia (GK)</i>	
.Alimentary tract and metabolism.....	93
<i>Avanza (SH)</i>	
.Nervous system	418
<i>Avanza SolTab (SH)</i>	
.Nervous system	418
<i>Avapro (BQ)</i>	
.Cardiovascular system.....	131
<i>Avapro HCT 150/12.5 (BQ)</i>	
.Cardiovascular system.....	132
<i>Avapro HCT 300/12.5 (BQ)</i>	
.Cardiovascular system.....	132
<i>Avapro HCT 300/25 (BQ)</i>	
.Cardiovascular system.....	132
<i>Avastin (RO)</i>	
.Antineoplastic and immunomodulating agents	209
<i>Avonex (BD)</i>	
.Antineoplastic and immunomodulating agents	235
<i>Axit 30 (AF)</i>	
.Nervous system	418
<i>Axit 15 (AF)</i>	
.Nervous system	418
<i>Aylide 1 (AF)</i>	
.Alimentary tract and metabolism.....	90
<i>Aylide 2 (AF)</i>	
.Alimentary tract and metabolism.....	90
<i>Aylide 3 (AF)</i>	
.Alimentary tract and metabolism.....	90
<i>Aylide 4 (AF)</i>	
.Alimentary tract and metabolism.....	90
<i>Azamun (GM)</i>	
.Antineoplastic and immunomodulating agents	356
<i>Azapin (SI)</i>	
.Antineoplastic and immunomodulating agents	357
<i>Azarga (AQ)</i>	
.Sensory organs	442
.Optometrical	517
AZATHIOPRINE	
.Antineoplastic and immunomodulating agents	356
<i>Azathioprine Sandoz (SZ)</i>	
.Antineoplastic and immunomodulating agents	357
AZITHROMYCIN	
.Antifungives for systemic use	185
.Sensory organs	440
.Section 100	561, 697
.Repatriation Schedule	849
<i>Azithromycin Sandoz (SZ)</i>	
.Antifungives for systemic use	185
.Sensory organs.....	440
<i>Azol 100 (AF)</i>	
.Genito urinary system and sex hormones	164
<i>Azol 200 (AF)</i>	
.Genito urinary system and sex hormones	164
<i>Azopt (AQ)</i>	
.Sensory organs.....	442
.Optometrical.....	517
B	
<i>B. Braun Australia Pty Ltd (BR)</i>	
.Blood and blood forming organs	105, 106
.Dental	488
BACLOFEN	
.Musculo-skeletal system	363
.Section 100	648, 785
<i>Bactigras 7457 (SN)</i>	
.Repatriation Schedule	869
<i>Bactrim (RO)</i>	
.Antifungives for systemic use	185
.Dental	498
<i>Bactrim DS (RO)</i>	
.Antifungives for systemic use	184
.Dental	498
<i>Bactroban (GK)</i>	
.Respiratory system	433
.Repatriation Schedule	844
BALSALAZIDE SODIUM	
.Alimentary tract and metabolism.....	83
BANDAGE—ABSORBENT WOOL	
.Repatriation Schedule	863
BANDAGE—CALICO	
.Repatriation Schedule	863
BANDAGE—COMPRESSION	
.Repatriation Schedule	863, 864
BANDAGE—RETENTION—COHESIVE—HEAVY	
.Repatriation Schedule	864
BANDAGE—RETENTION—COHESIVE—LIGHT	
.Repatriation Schedule	864
BANDAGE—RETENTION—COTTON CREPE	
.Repatriation Schedule	865
BANDAGE—TUBULAR	
.Repatriation Schedule	865
BANDAGE—TUBULAR (FINGER)	
.Repatriation Schedule	865
BANDAGE—TUBULAR (LIGHTWEIGHT)	
.Repatriation Schedule	865
BANDAGE—TUBULAR (LONG STOCKING)	
.Repatriation Schedule	865
BANDAGE—TUBULAR (SHORT STOCKING)	
.Repatriation Schedule	865
BANDAGE—ZINC PASTE	
.Repatriation Schedule	865
<i>Baraclude (BQ)</i>	
.Section 100	564, 701
<i>Barbloc 15 (AF)</i>	
.Cardiovascular system	114
<i>Barbloc 5 (AF)</i>	
.Cardiovascular system	114

Baxter Healthcare Pty Ltd (BX)

.Blood and blood forming organs	105, 106
.Antifungives for systemic use	189, 191
.Dental	488, 499
.Repatriation Schedule	840

BCG IMMUNOTHERAPEUTIC (Bacillus Calmette-Guérin/Connaught strain)

.Antineoplastic and immunomodulating agents	236

BCG-TICE (Bacillus Calmette-Guérin/ Tice strain)

.Antineoplastic and immunomodulating agents	236

BECLOMETHASONE DIPROPIONATE

.Respiratory system	435

BenPen (CS)

.Emergency Drug Supplies	64
.Antifungives for systemic use	178
.Dental	493

BENZATHINE BENZYL PENICILLIN

.Antifungives for systemic use	178
.Dental	493

BENZHEXOL HYDROCHLORIDE

.Nervous system	395

Benztrop (PL)

.Nervous system	395

BENZTROPINE MESYLATE

.Nervous system	395
.Dental	509

BENZYDAMINE HYDROCHLORIDE

.Alimentary tract and metabolism	68
.Palliative Care	469
.Dental	486

BENZYL PENICILLIN

.Antifungives for systemic use	178
.Dental	493

Bergoline 1 (SI)

.Nervous system	397

Bergoline 2 (SI)

.Nervous system	397

Betachek (NA)

.Various	455, 456

Betachek G5 (NA)

.Various	455, 456

Betadine Antiseptic Liquid (SW)

.Repatriation Schedule	845

Betaferon (SC)

.Antineoplastic and immunomodulating agents	236

Betaloc (AP)

.Cardiovascular system	116

BETAMETHASONE ACETATE with BETAMETHASONE**SODIUM PHOSPHATE**

.Systemic hormonal preparations, excl. sex hormones and insulins	168
.Dental	491

BETAMETHASONE DIPROPIONATE

.Dermatologicals	152

BETAMETHASONE VALERATE

.Dermatologicals	152, 153
.Repatriation Schedule	845

Betamin (SW)

.Alimentary tract and metabolism	95
.Repatriation Schedule	839

BETAXOLOL HYDROCHLORIDE

.Sensory organs	442
.Optometrical	517

BETHANECHOL CHLORIDE

.Nervous system	427

Betnovate (SI)

.Repatriation Schedule	845

Betnovate 1/2 (SI)

.Dermatologicals	153

Betnovate 1/5 (SI)

.Dermatologicals	153

Betoptic (AQ)

.Sensory organs	442
.Optometrical	517

Betoptic S (AQ)

.Sensory organs	442
.Optometrical	517

BetoQuin (IQ)

.Sensory organs	442
.Optometrical	517

BEVACIZUMAB

.Antineoplastic and immunomodulating agents	209

Bgramin (GM)

.Antifungives for systemic use	177
.Dental	493

Biatain Adhesive 3420 (CT)

.Repatriation Schedule	870

Biatain Adhesive 3423 (CT)

.Repatriation Schedule	870

Biatain Ag 9622 (CT)

.Repatriation Schedule	874

Biatain Ag 9632 (CT)

.Repatriation Schedule	874

Biatain Non-adhesive 3410 (CT)

.Repatriation Schedule	870

Biatain Non-adhesive 3413 (CT)

.Repatriation Schedule	870

Biaxig (AV)

.Antifungives for systemic use	186
.Dental	498

BICALUTAMIDE

.Antineoplastic and immunomodulating agents	232

Bicalutamide Ranbaxy (RA)

.Antineoplastic and immunomodulating agents	232

Bicalutamide-GA (GM)

.Antineoplastic and immunomodulating agents	232

Bicillin L-A (AS)

.Antifungives for systemic use	178
.Dental	493

Bicor (AL)

.Cardiovascular system	115

Biltricide (BN)

.Antiparasitic products, insecticides and repellents	431

BIMATOPROST

.Sensory organs	443
.Optometrical	518

BIMATOPROST with TIMOLOL MALEATE

.Sensory organs	443
.Optometrical	518

Biodone Forte (MW)

.Section 100	798	.Respiratory system	438
<i>Bion Tears (AQ)</i>		<i>Bricanyl Turbuhaler (AP)</i>	
.Sensory organs	448	.Respiratory system	434
.Optometrical	520	BRIMONIDINE TARTRATE	
<i>Bionime Rightest (QB)</i>		.Sensory organs.....	441
.Various.....	456	.Optometrical.....	516
BIPERIDEN HYDROCHLORIDE		BRIMONIDINE TARTRATE with TIMOLOL MALEATE	
.Nervous system	395	.Sensory organs.....	442
BISACODYL		.Optometrical.....	517
.Alimentary tract and metabolism.....	80, 81	BRINZOLAMIDE	
.Palliative Care	471, 473	.Sensory organs.....	442
<i>Bisalax (AS)</i>		.Optometrical.....	517
.Alimentary tract and metabolism.....	80, 81	BRINZOLAMIDE with TIMOLOL MALEATE	
.Palliative Care	471, 473	.Sensory organs.....	442
BISOPROLOL FUMARATE		.Optometrical.....	517
.Cardiovascular system.....	115	<i>BrinzoQuin (IQ)</i>	
<i>Bisoprolol Sandoz (SZ)</i>		.Sensory organs.....	442
.Cardiovascular system.....	115	.Optometrical.....	517
<i>Bispro 10 (AF)</i>		BROMAZEPAM	
.Cardiovascular system.....	115	.Repatriation Schedule	854
<i>Bispro 2.5 (AF)</i>		BROMOCRIPTINE MESYLATE	
.Cardiovascular system.....	115	.Genito urinary system and sex hormones	156
<i>Bispro 5 (AF)</i>		.Nervous system	397
.Cardiovascular system.....	115	<i>Brufen (AB)</i>	
BIVALIRUDIN TRIFLUOROACETATE		.Musculo-skeletal system	360
.Blood and blood forming organs	102	.Palliative Care	477
<i>Blenamax (SI)</i>		.Dental	502
.Antineoplastic and immunomodulating agents	66	BSN 2902165 (BV)	
<i>Blenoxane (BQ)</i>		.Repatriation Schedule	874
.Antineoplastic and immunomodulating agents	66	<i>Budamax Aqueous (PM)</i>	
BLEOMYCIN SULFATE		.Repatriation Schedule	857
.Antineoplastic and immunomodulating agents	66	BUDESONIDE	
<i>Bleph 10 (AG)</i>		.Respiratory system	436
.Sensory organs	440	.Repatriation Schedule	857
.Optometrical	516	BUDESONIDE with EFORMOTEROL FUMARATE DIHYDRATE	
<i>Blink Intensive Tears (AO)</i>		.Respiratory system	434
.Sensory organs	449	BUPRENORPHINE	
.Optometrical	520	.Nervous system	381
<i>Bondronat (HH)</i>		.Section 100	798
.Musculo-skeletal system	367	BUPRENORPHINE with NALOXONE	
.Section 100	649, 786	.Section 100	798
<i>Bonefos (SC)</i>		BUPROPION HYDROCHLORIDE	
.Musculo-skeletal system	368	.Nervous system	427
<i>Bonefos 800 mg (SC)</i>		<i>Buscopan (BY)</i>	
.Musculo-skeletal system	368	.Emergency Drug Supplies	64
BORTEZOMIB		.Palliative Care	470
.Antineoplastic and immunomodulating agents .	226, 228	.Repatriation Schedule	837
BOSENTAN MONOHYDRATE		<i>Buspar (SI)</i>	
.Section 100	531, 663	.Repatriation Schedule	854
<i>Botox (AG)</i>		BUSPIRONE HYDROCHLORIDE	
.Section 100	793	.Repatriation Schedule	854
BOTULINUM TOXIN TYPE A PURIFIED NEUROTOXIN COMPLEX		BUSULFAN	
.Section 100	793	.Antineoplastic and immunomodulating agents	199
<i>Brevinor (PF)</i>		<i>Butamol 2.5 (SI)</i>	
.Genito urinary system and sex hormones.....	157	.Emergency Drug Supplies	65
<i>Brevinor-1 (PF)</i>		.Respiratory system	433
.Genito urinary system and sex hormones.....	157, 158	<i>Butamol 5 (SI)</i>	
<i>Bricanyl (AP)</i>		.Emergency Drug Supplies	65
.Emergency Drug Supplies	65	.Respiratory system	434
		<i>Byetta 10 microgram (LY)</i>	

.Alimentary tract and metabolism.....	94
<i>Byetta 5 microgram (LY)</i>	
.Alimentary tract and metabolism.....	94
C	
<i>C-Flox 250 (AL)</i>	
.Antifungives for systemic use.....	187
<i>C-Flox 500 (AL)</i>	
.Antifungives for systemic use.....	188
<i>C-Flox 750 (AL)</i>	
.Antifungives for systemic use.....	188
<i>Cabaser (PF)</i>	
.Nervous system.....	397
CABERGOLINE	
.Genito urinary system and sex hormones.....	156
.Nervous system.....	397
<i>Caduet 10/10 (PF)</i>	
.Cardiovascular system.....	148
<i>Caduet 10/20 (PF)</i>	
.Cardiovascular system.....	148
<i>Caduet 10/40 (PF)</i>	
.Cardiovascular system.....	148
<i>Caduet 10/80 (PF)</i>	
.Cardiovascular system.....	148
<i>Caduet 5/10 (PF)</i>	
.Cardiovascular system.....	148
<i>Caduet 5/20 (PF)</i>	
.Cardiovascular system.....	148
<i>Caduet 5/40 (PF)</i>	
.Cardiovascular system.....	148
<i>Caduet 5/80 (PF)</i>	
.Cardiovascular system.....	148
<i>Caelyx (SH)</i>	
.Antineoplastic and immunomodulating agents.....	206
.Section 100.....	569, 706
<i>CAL-600 (PP)</i>	
.Repatriation Schedule.....	839
<i>Cal-Sup (IA)</i>	
.Alimentary tract and metabolism.....	95
.Repatriation Schedule.....	839
<i>Calci-Tab 600 (AE)</i>	
.Alimentary tract and metabolism.....	95
CALCIPOTRIOL	
.Dermatologicals.....	151
CALCIPOTRIOL with BETAMETHASONE DIPROPIONATE	
.Dermatologicals.....	151
CALCITRIOL	
.Alimentary tract and metabolism.....	95
.Musculo-skeletal system.....	371
<i>Calcitriol Sandoz (SZ)</i>	
.Alimentary tract and metabolism.....	95
.Musculo-skeletal system.....	371
<i>Calcitriol-DP (GN)</i>	
.Alimentary tract and metabolism.....	95
.Musculo-skeletal system.....	371
<i>Calcitriol-GA (GM)</i>	
.Alimentary tract and metabolism.....	95
.Musculo-skeletal system.....	371
CALCIUM	
.Alimentary tract and metabolism.....	95

.Repatriation Schedule.....	839
CALCIUM CARBONATE with GLYCINE	
.Repatriation Schedule.....	837
CALCIUM FOLINATE	
.Various.....	453
<i>Calcium Folate Ebewe (IT)</i>	
.Various.....	453
<i>Calmurid (OL)</i>	
.Repatriation Schedule.....	843
<i>Calutex (SI)</i>	
.Antineoplastic and immunomodulating agents.....	232
<i>Campral (AF)</i>	
.Nervous system.....	428
<i>Camptosar (PF)</i>	
.Antineoplastic and immunomodulating agents.....	229
CANDESARTAN CILEXETIL	
.Cardiovascular system.....	130
CANDESARTAN CILEXETIL with HYDROCHLOROTHIAZIDE	
.Cardiovascular system.....	131
CAPECITABINE	
.Antineoplastic and immunomodulating agents.....	202
<i>Capoten (SI)</i>	
.Cardiovascular system.....	121, 122
<i>Caprilon (SB)</i>	
.Various.....	461
CAPTOPRIL	
.Cardiovascular system.....	121, 122
<i>Captopril Sandoz (SZ)</i>	
.Cardiovascular system.....	121
<i>Carafate (AS)</i>	
.Alimentary tract and metabolism.....	75
CARBAMAZEPINE	
.Nervous system.....	388
.Dental.....	509
<i>Carbamazepine Sandoz (SZ)</i>	
.Nervous system.....	388
.Dental.....	509
CARBAMIDE PEROXIDE	
.Repatriation Schedule.....	859
CARBIMAZOLE	
.Systemic hormonal preparations, excl. sex hormones and insulins.....	171
<i>CarboFLEX 403202 (CC)</i>	
.Repatriation Schedule.....	866
<i>CarboFLEX 403204 (CC)</i>	
.Repatriation Schedule.....	866
<i>Carbohydrate Free Mixture (SB)</i>	
.Various.....	466
CARBOHYDRATE, FAT, VITAMINS, MINERALS and TRACE ELEMENTS	
.Various.....	465
CARBOMER	
.Sensory organs.....	445, 446
.Optometrical.....	518, 519
CARBOMER 974	
.Sensory organs.....	446
.Optometrical.....	519
CARBOPLATIN	
.Antineoplastic and immunomodulating agents.....	207
<i>Carboplatin Ebewe (IT)</i>	

.Antineoplastic and immunomodulating agents	207
<i>Cardinorm (SZ)</i>	
.Cardiovascular system	108
<i>Cardiprin 100 (RC)</i>	
.Repatriation Schedule	840
<i>Cardizem (SW)</i>	
.Cardiovascular system	121
<i>Cardizem CD (SW)</i>	
.Cardiovascular system	121
<i>Cardol (AF)</i>	
.Cardiovascular system	108, 114
<i>CareSens (LB)</i>	
.Various	455, 456
<i>CareSens N (LB)</i>	
.Various	455, 456
CARMELLOSE SODIUM	
.Sensory organs	446
.Palliative Care	469
.Optometrical	519
.Repatriation Schedule	837
CARMELLOSE SODIUM with GLYCERIN	
.Sensory organs	447
.Optometrical	519
CARMELLOSE SODIUM with PECTIN and GELATIN	
.Repatriation Schedule	843
CARMUSTINE	
.Antineoplastic and immunomodulating agents	199
<i>Cartia (GC)</i>	
.Repatriation Schedule	840
CARVEDILOL	
.Cardiovascular system	116
<i>Carvedilol generichealth (GQ)</i>	
.Cardiovascular system	117
<i>Carvedilol Sandoz (SZ)</i>	
.Cardiovascular system	117
<i>Catapres (BY)</i>	
.Cardiovascular system	110
<i>Catapres 100 (BY)</i>	
.Cardiovascular system	110
CATIONIC CONDITIONER with PANTHENOL	
.Repatriation Schedule	846
<i>Caverject Impulse (PH)</i>	
.Repatriation Schedule	847
<i>Cavicare 4563 (SN)</i>	
.Repatriation Schedule	868
<i>Ceclor (AS)</i>	
.Antiinfectives for systemic use	183
.Dental	497
<i>Ceclor CD (AS)</i>	
.Antiinfectives for systemic use	183
.Dental	497
CEFACLOR	
.Antiinfectives for systemic use	182
.Dental	496
<i>Cefaclor Sandoz (SZ)</i>	
.Antiinfectives for systemic use	183
.Dental	497
<i>Cefaclor-GA (GN)</i>	
.Antiinfectives for systemic use	182
.Dental	497
<i>Cefalexin Sandoz (SZ)</i>	
.Antiinfectives for systemic use	181, 182
.Dental	495, 496
CEFALOTIN	
.Antiinfectives for systemic use	181
.Dental	495
<i>Cefalotin Sandoz (SZ)</i>	
.Antiinfectives for systemic use	181
.Dental	495
<i>Cefazolin Sandoz (SZ)</i>	
.Antiinfectives for systemic use	182
CEFEPIME	
.Antiinfectives for systemic use	184
CEFOTAXIME	
.Antiinfectives for systemic use	183
.Dental	497
<i>Cefotaxime Sandoz (SZ)</i>	
.Antiinfectives for systemic use	183
.Dental	497
CEFTRIAXONE	
.Antiinfectives for systemic use	183, 184
<i>Ceftriaxone ICP (PP)</i>	
.Antiinfectives for systemic use	183, 184
<i>Ceftriaxone Sandoz (SZ)</i>	
.Antiinfectives for systemic use	184
CEFUROXIME AXETIL	
.Antiinfectives for systemic use	183
.Dental	497
<i>Celapram (AF)</i>	
.Nervous system	412
<i>Celebrex (PF)</i>	
.Musculo-skeletal system	362
CELECOXIB	
.Musculo-skeletal system	361
<i>Celestone Chronodose (SH)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	169
.Dental	491
<i>Celestone-M (SH)</i>	
.Dermatologicals	153
<i>Celica (RA)</i>	
.Nervous system	412
<i>CellCept (RO)</i>	
.Antineoplastic and immunomodulating agents	237
.Section 100	581, 719
<i>Cellufresh (AG)</i>	
.Sensory organs	447
.Optometrical	519
<i>Celluvisc (AG)</i>	
.Sensory organs	446
.Optometrical	519
<i>Celsentri (PF)</i>	
.Section 100	568, 705
<i>Cephalexin (BF)</i>	
.Antiinfectives for systemic use	181, 182
.Dental	495, 496
CEPHALEXIN	
.Antiinfectives for systemic use	181
.Dental	495
<i>Cephalexin generichealth (GQ)</i>	

.Antifectives for systemic use	181, 182
.Dental	496
<i>Cephadrust 250 (MI)</i>	
.Antifectives for systemic use	181
.Dental	496
<i>Cephadrust 500 (MI)</i>	
.Antifectives for systemic use	182
.Dental	496
CEPHAZOLIN	
.Antifectives for systemic use	182
<i>Cephazolin Alphapharm (AF)</i>	
.Antifectives for systemic use	182
<i>Certican (NV)</i>	
.Antineoplastic and immunomodulating agents	236
.Section 100	581, 718
CERTOLIZUMAB PEGOL	
.Antineoplastic and immunomodulating agents	285
<i>Cerumol (AC)</i>	
.Repatriation Schedule	859
CETIRIZINE HYDROCHLORIDE	
.Repatriation Schedule	858
CETRORELIX	
.Section 100	797
<i>Cetrotide (SG)</i>	
.Section 100	797
CETUXIMAB	
.Antineoplastic and immunomodulating agents	209
<i>Champix (PF)</i>	
.Nervous system	428
<i>Chem mart Aciclovir (CH)</i>	
.Antifectives for systemic use	194
<i>Chem mart Alendronate 70mg (CH)</i>	
.Musculo-skeletal system	365
<i>Chem mart Allopurinol (CH)</i>	
.Musculo-skeletal system	364
<i>Chem mart Alprazolam (CH)</i>	
.Nervous system	408
<i>Chem mart Amiodarone (CH)</i>	
.Cardiovascular system	108
<i>Chem mart Amlodipine (CH)</i>	
.Cardiovascular system	118, 119
<i>Chem mart Amoxycillin (CH)</i>	
.Antifectives for systemic use	177, 178
.Dental	492, 493
<i>Chem mart Amoxycillin and Clavulanic Acid (CH)</i>	
.Antifectives for systemic use	180
.Dental	494
<i>Chem mart Atenolol (CH)</i>	
.Cardiovascular system	115
<i>Chem mart Baclofen (CH)</i>	
.Musculo-skeletal system	363
<i>Chem mart Carvedilol 12.5 mg (CH)</i>	
.Cardiovascular system	117
<i>Chem mart Carvedilol 25 mg (CH)</i>	
.Cardiovascular system	117
<i>Chem mart Carvedilol 3.125 mg (CH)</i>	
.Cardiovascular system	117
<i>Chem mart Carvedilol 6.25 mg (CH)</i>	
.Cardiovascular system	117
<i>Chem mart Cefaclor (CH)</i>	

.Antifectives for systemic use	183
.Dental	497
<i>Chem mart Cefaclor CD (CH)</i>	
.Antifectives for systemic use	182
.Dental	497
<i>Chem mart Cephalexin (CH)</i>	
.Antifectives for systemic use	181, 182
.Dental	496
<i>Chem mart Citalopram (CH)</i>	
.Nervous system	412
<i>Chem mart Clarithromycin (CH)</i>	
.Antifectives for systemic use	185
<i>Chem mart Clomipramine (CH)</i>	
.Nervous system	409, 411
<i>Chem mart Clopidogrel (CH)</i>	
.Blood and blood forming organs	99
<i>Chem mart Diclofenac (CH)</i>	
.Musculo-skeletal system	358
.Palliative Care	475, 476
.Dental	501
<i>Chem mart Diltiazem (CH)</i>	
.Cardiovascular system	121
<i>Chem mart Diltiazem CD (CH)</i>	
.Cardiovascular system	121
<i>Chem mart Doxycycline (CH)</i>	
.Antifectives for systemic use	175, 176
.Dental	492
<i>Chem mart Enalapril (CH)</i>	
.Cardiovascular system	122
<i>Chem mart Escitalopram (CH)</i>	
.Nervous system	413
<i>Chem mart Famotidine (CH)</i>	
.Alimentary tract and metabolism	69
<i>Chem mart Fluoxetine (CH)</i>	
.Nervous system	414
<i>Chem mart Frusemide (CH)</i>	
.Cardiovascular system	112
<i>Chem mart Gemfibrozil (CH)</i>	
.Cardiovascular system	145
<i>Chem mart Gliclazide (CH)</i>	
.Alimentary tract and metabolism	89
<i>Chem mart Gliclazide MR (CH)</i>	
.Alimentary tract and metabolism	90
<i>Chem mart Indapamide (CH)</i>	
.Cardiovascular system	112
<i>Chem mart Isosorbide Mononitrate (CH)</i>	
.Cardiovascular system	110
<i>Chem mart Lercanidipine (CH)</i>	
.Cardiovascular system	119
<i>Chem mart Levetiracetam (CH)</i>	
.Nervous system	393
<i>Chem mart Lisinopril (CH)</i>	
.Cardiovascular system	123, 124
<i>Chem mart Meloxicam 15 mg (CH)</i>	
.Musculo-skeletal system	359
<i>Chem mart Meloxicam 7.5 mg (CH)</i>	
.Musculo-skeletal system	359
<i>Chem mart Metformin (CH)</i>	
.Alimentary tract and metabolism	88
<i>Chem mart Metoprolol (CH)</i>	

.Cardiovascular system.....	116
<i>Chem mart Mirtazapine (CH)</i>	
.Nervous system	418
<i>Chem mart Moclobemide (CH)</i>	
.Nervous system	416
<i>Chem mart Norfloxacin (CH)</i>	
.Antiinfectives for systemic use	188
<i>Chem mart Omeprazole (CH)</i>	
.Alimentary tract and metabolism.....	72, 73
<i>Chem mart Pantoprazole (CH)</i>	
.Alimentary tract and metabolism.....	73, 74
<i>Chem mart Paracetamol (CH)</i>	
.Nervous system	384
.Dental	508, 509
<i>Chem mart Paroxetine (CH)</i>	
.Nervous system	415
<i>Chem mart Perindopril (CH)</i>	
.Cardiovascular system.....	124, 125
<i>Chem mart Perindopril/ Indapamide 4/1.25 (CH)</i>	
.Cardiovascular system.....	129
<i>Chem mart Piroxicam (CH)</i>	
.Musculo-skeletal system	360
.Dental	502
<i>Chem mart Pravastatin (CH)</i>	
.Cardiovascular system.....	138, 139, 140
<i>Chem mart Prazosin (CH)</i>	
.Cardiovascular system.....	111
<i>Chem mart Ramipril (CH)</i>	
.Cardiovascular system.....	126, 127
<i>Chem mart Ranitidine (CH)</i>	
.Alimentary tract and metabolism.....	69
<i>Chem mart Roxithromycin (CH)</i>	
.Antiinfectives for systemic use	186
.Dental	498
<i>Chem mart Sertraline (CH)</i>	
.Nervous system	415
<i>Chem mart Simvastatin (CH)</i>	
.Cardiovascular system.....	140, 141, 142, 143, 144
<i>Chem mart Sotalol (CH)</i>	
.Cardiovascular system.....	108, 114
<i>Chem mart Sumatriptan (CH)</i>	
.Nervous system	386
<i>Chem mart Tramadol (CH)</i>	
.Nervous system	382
.Dental	507
<i>Chem mart Tramadol SR (CH)</i>	
.Nervous system	383
.Dental	507, 508
CHLORAMBUCIL	
.Antineoplastic and immunomodulating agents	199
CHLORAMPHENICOL	
.Sensory organs	440, 450
.Dental	512
.Optometrical	516
CHLORHEXIDINE GLUCONATE	
.Repatriation Schedule	837
<i>Chloromycetin (PF)</i>	
.Sensory organs	440, 450
.Dental	512
.Optometrical	516
CHLORPROMAZINE HYDROCHLORIDE	
.Nervous system	400
<i>Chlorsig (SI)</i>	
.Sensory organs.....	440
.Dental	512
.Optometrical.....	516
CHLORTHALIDONE	
.Cardiovascular system	112
<i>Chlorvescent (AS)</i>	
.Alimentary tract and metabolism.....	96
CHOLESTYRAMINE	
.Cardiovascular system	145, 146
<i>Cholstat 10 (AF)</i>	
.Cardiovascular system	138, 139
<i>Cholstat 20 (AF)</i>	
.Cardiovascular system	138, 139
<i>Cholstat 40 (AF)</i>	
.Cardiovascular system	138, 139
CHORIOGONADOTROPIN ALFA	
.Section 100	796
<i>Cialis (LY)</i>	
.Repatriation Schedule	848
<i>Ciazil (GM)</i>	
.Nervous system	412
CICLESONIDE	
.Respiratory system	436
CICLOPIROX OLAMINE	
.Repatriation Schedule	842
<i>Cicloral (SZ)</i>	
.Antineoplastic and immunomodulating agents	356
.Section 100	640, 776, 777
CIDOFOVIR	
.Section 100	561, 697
<i>Cifran (RA)</i>	
.Antiinfectives for systemic use	187, 188
<i>Cilamox (SI)</i>	
.Antiinfectives for systemic use	177, 178
.Dental	492, 493
<i>Cilex (GM)</i>	
.Antiinfectives for systemic use	181, 182
.Dental	496
<i>Cilicaine (SI)</i>	
.Emergency Drug Supplies	64
.Antiinfectives for systemic use	179
.Dental	494
<i>Cilicaine V (FM)</i>	
.Antiinfectives for systemic use	179
.Dental	493
<i>Cilicaine VK (FM)</i>	
.Antiinfectives for systemic use	178, 179
.Dental	493
<i>Cilopen VK (GM)</i>	
.Antiinfectives for systemic use	178, 179
.Dental	493
<i>CiloQuin (IQ)</i>	
.Sensory organs.....	441
<i>Ciloxan (AQ)</i>	
.Sensory organs.....	441, 451
CIMETIDINE	
.Alimentary tract and metabolism.....	68

<i>Cimzia (UC)</i>		
.Antineoplastic and immunomodulating agents	290	
<i>CINACALCET HYDROCHLORIDE</i>		
.Systemic hormonal preparations, excl. sex hormones and insulins	173	
.Section 100	559, 695	
<i>Cipramil (LU)</i>		
.Nervous system	412	
<i>CIPROFLOXACIN</i>		
.Antimicrobials for systemic use	187	
.Sensory organs	441, 450	
<i>Ciprofloxacin 500 (CR)</i>		
.Antimicrobials for systemic use	188	
<i>Ciprofloxacin 750 (CR)</i>		
.Antimicrobials for systemic use	188	
<i>Ciprofloxacin Sandoz (SZ)</i>		
.Antimicrobials for systemic use	187, 188	
<i>Ciprofloxacin-BW (BF)</i>		
.Antimicrobials for systemic use	188	
<i>Ciprofloxacin-DRLA (RZ)</i>		
.Antimicrobials for systemic use	187, 188	
<i>Ciprofloxacin-GA (GM)</i>		
.Antimicrobials for systemic use	188	
<i>Ciprol 250 (SI)</i>		
.Antimicrobials for systemic use	187	
<i>Ciprol 500 (SI)</i>		
.Antimicrobials for systemic use	188	
<i>Ciprol 750 (SI)</i>		
.Antimicrobials for systemic use	188	
<i>Ciproxin 250 (BN)</i>		
.Antimicrobials for systemic use	187	
<i>Ciproxin 500 (BN)</i>		
.Antimicrobials for systemic use	188	
<i>Ciproxin 750 (BN)</i>		
.Antimicrobials for systemic use	188	
<i>CISPLATIN</i>		
.Antineoplastic and immunomodulating agents	207	
<i>Cisplatin Ebewe (IT)</i>		
.Antineoplastic and immunomodulating agents	207	
<i>Citalobell (BF)</i>		
.Nervous system	412	
<i>Citalopram 20 (CR)</i>		
.Nervous system	412	
<i>Citalopram generichealth (GQ)</i>		
.Nervous system	412	
<i>CITALOPRAM HYDROBROMIDE</i>		
.Nervous system	412	
<i>Citalopram Sandoz (SZ)</i>		
.Nervous system	412	
<i>CLADRIBINE</i>		
.Antineoplastic and immunomodulating agents	201	
<i>Clamoxyl (AL)</i>		
.Antimicrobials for systemic use	180	
.Dental	495	
<i>Clamoxyl Duo (AL)</i>		
.Antimicrobials for systemic use	180	
.Dental	495	
<i>Clamoxyl Duo 400 (AL)</i>		
.Antimicrobials for systemic use	180	
.Dental	495	
<i>Clamoxyl Duo forte (AL)</i>		
.Antimicrobials for systemic use	180	
.Dental	495	
<i>Clarac (GM)</i>		
.Antimicrobials for systemic use	185	
<i>Claratyne (SH)</i>		
.Repatriation Schedule	858	
<i>Clarithexal (SZ)</i>		
.Antimicrobials for systemic use	185	
<i>Clarithro 250 (SI)</i>		
.Antimicrobials for systemic use	185	
<i>CLARITHROMYCIN</i>		
.Antimicrobials for systemic use	185	
.Section 100	561, 697	
<i>Clavycillin 875/125 (CR)</i>		
.Antimicrobials for systemic use	180	
.Dental	495	
<i>Cleocin (KR)</i>		
.Antimicrobials for systemic use	186	
.Dental	499	
<i>Clexane (SW)</i>		
.Blood and blood forming organs	97, 98	
<i>Climara 100 (SC)</i>		
.Genito urinary system and sex hormones	160	
<i>Climara 25 (SC)</i>		
.Genito urinary system and sex hormones	160	
<i>Climara 50 (SC)</i>		
.Genito urinary system and sex hormones	159	
<i>Climara 75 (SC)</i>		
.Genito urinary system and sex hormones	160	
<i>CLINDAMYCIN</i>		
.Antimicrobials for systemic use	186	
.Dental	499	
<i>Clinistix (BN)</i>		
.Various	455	
<i>Clinitest (BN)</i>		
.Various	454	
<i>Clobemix (GM)</i>		
.Nervous system	416	
<i>Clofen 10 (AF)</i>		
.Musculo-skeletal system	363	
<i>Clofen 25 (AF)</i>		
.Musculo-skeletal system	363	
<i>Clomid (SW)</i>		
.Genito urinary system and sex hormones	163	
<i>CLOMIPHENE CITRATE</i>		
.Genito urinary system and sex hormones	163	
<i>CLOMIPRAMINE HYDROCHLORIDE</i>		
.Nervous system	409, 411	
<i>Clonac 25 (SI)</i>		
.Musculo-skeletal system	358	
.Palliative Care	475	
.Dental	501	
<i>Clonac 50 (SI)</i>		
.Musculo-skeletal system	358	
.Palliative Care	475, 476	
.Dental	501	
<i>CLONAZEPAM</i>		
.Nervous system	388	
.Palliative Care	483	

<i>Clonea (AF)</i>		
.Dermatologicals.....	149	
.Repatriation Schedule	842	
CLONIDINE		
.Cardiovascular system.....	110	
CLOPIDOGREL		
.Blood and blood forming organs.....	98, 99	
.Repatriation Schedule	840	
<i>Clpidogrel Actavis (GQ)</i>		
.Blood and blood forming organs.....	99	
<i>Clpidogrel Sandoz (SZ)</i>		
.Blood and blood forming organs.....	99	
<i>Clpidogrel Winthrop (WA)</i>		
.Blood and blood forming organs.....	99	
CLOPIDOGREL with ASPIRIN		
.Blood and blood forming organs.....	99	
<i>Clpidogrel-GA (GM)</i>		
.Blood and blood forming organs.....	99	
<i>Clopine 100 (HH)</i>		
.Section 100.....	650, 787	
<i>Clopine 200 (HH)</i>		
.Section 100.....	650, 787	
<i>Clopine 25 (HH)</i>		
.Section 100.....	650, 787	
<i>Clopine 50 (HH)</i>		
.Section 100.....	650, 787	
<i>Clopine Suspension (HH)</i>		
.Section 100.....	650, 787	
<i>Clopixol Depot (LU)</i>		
.Nervous system.....	401	
CLOSTRIDIUM BOTULINUM TYPE A TOXIN— HAEMAGGLUTININ COMPLEX		
.Section 100.....	793	
CLOTRIMAZOLE		
.Dermatologicals.....	149	
.Repatriation Schedule	842, 847	
<i>Clovix 75 (SI)</i>		
.Blood and blood forming organs.....	99	
CLOZAPINE		
.Section 100.....	650, 787	
<i>Clozaril 100 (NV)</i>		
.Section 100.....	650, 787	
<i>Clozaril 25 (NV)</i>		
.Section 100.....	650, 787	
<i>Co-Diovan 160/12.5 (NV)</i>		
.Cardiovascular system.....	132	
<i>Co-Diovan 160/25 (NV)</i>		
.Cardiovascular system.....	132	
<i>Co-Diovan 320/12.5 (NV)</i>		
.Cardiovascular system.....	132	
<i>Co-Diovan 320/25 (NV)</i>		
.Cardiovascular system.....	132	
<i>Co-Diovan 80/12.5 (NV)</i>		
.Cardiovascular system.....	132	
COAL TAR - PREPARED		
.Dermatologicals.....	150	
<i>Coban 1584 (MM)</i>		
.Repatriation Schedule	864	
<i>Codalgin (FM)</i>		
.Repatriation Schedule	853	
<i>Codalgin Forte (FM)</i>		
.Nervous system.....	375	
.Dental	504	
<i>Codapane Forte (AL)</i>		
.Nervous system.....	375	
.Dental	504	
CODEINE PHOSPHATE		
.Nervous system.....	375	
.Respiratory system.....	439	
.Dental	504	
CODEINE PHOSPHATE with ASPIRIN		
.Repatriation Schedule	853	
CODEINE PHOSPHATE with PARACETAMOL		
.Nervous system.....	375	
.Dental	504	
.Repatriation Schedule	853	
<i>Cogentin (FK)</i>		
.Emergency Drug Supplies.....	64	
.Nervous system.....	395	
.Dental	509	
<i>Colazide (PK)</i>		
.Alimentary tract and metabolism.....	83	
COLCHICINE		
.Musculo-skeletal system.....	364	
<i>Colese (AF)</i>		
.Repatriation Schedule	837	
<i>Colestid (PF)</i>		
.Cardiovascular system.....	146	
COLESTIPOL HYDROCHLORIDE		
.Cardiovascular system.....	146	
<i>Colgout (AS)</i>		
.Musculo-skeletal system.....	364	
<i>Colifoam (AS)</i>		
.Alimentary tract and metabolism.....	83	
<i>Colofac (SM)</i>		
.Repatriation Schedule	837	
<i>Coloxyl 50 (FM)</i>		
.Repatriation Schedule	837	
<i>Coloxyl with Senna (FM)</i>		
.Repatriation Schedule	838	
<i>CombiDERM 651027 (CC)</i>		
.Repatriation Schedule	869	
<i>CombiDERM 651031 (CC)</i>		
.Repatriation Schedule	869	
<i>Combigan (AG)</i>		
.Sensory organs.....	442	
.Optometical.....	517	
<i>Combivir (VI)</i>		
.Section 100.....	567, 704	
<i>Comfarol Forte (SZ)</i>		
.Nervous system.....	375	
.Dental	504	
<i>Comfeel Paste 4701 (CT)</i>		
.Repatriation Schedule	870	
<i>Comfeel Plus Pressure Relieving 3350 (CT)</i>		
.Repatriation Schedule	872	
<i>Comfeel Plus Pressure Relieving 3353 (CT)</i>		
.Repatriation Schedule	872	
<i>Comfeel Plus Transparent 3530 (CT)</i>		
.Repatriation Schedule	871	

<i>Comfeel Plus Transparent 3533 (CT)</i>		
.Repatriation Schedule	871	
<i>Comfeel Plus Transparent 3536 (CT)</i>		
.Repatriation Schedule	871	
<i>Comfeel Plus Ulcer Dressing 3110 (CT)</i>		
.Repatriation Schedule	872	
<i>Comfeel Purilon Gel 3900 (CT)</i>		
.Repatriation Schedule	872	
<i>Comfeel SeaSorb Dressing 3705 (CT)</i>		
.Repatriation Schedule	866	
<i>Comfeel SeaSorb Dressing 3710 (CT)</i>		
.Repatriation Schedule	866	
<i>Comfeel SeaSorb Filler 3740 (CT)</i>		
.Repatriation Schedule	866	
<i>Comprilan 1027 (BV)</i>		
.Repatriation Schedule	863	
<i>Comtan (NV)</i>		
.Nervous system	399	
<i>Concerta (JC)</i>		
.Nervous system	420	
<i>Concorz (SZ)</i>		
.Nervous system	415	
<i>Condylone Paint (HA)</i>		
.Repatriation Schedule	844	
<i>Copaxone (SW)</i>		
.Antineoplastic and immunomodulating agents	236	
<i>CoPlavix (SW)</i>		
.Blood and blood forming organs	99	
<i>Coplus 3629 (BV)</i>		
.Repatriation Schedule	864	
COPPER SULFATE		
.Various.....	454	
<i>Coras (AF)</i>		
.Cardiovascular system	121	
<i>Corbeton 20 (AF)</i>		
.Cardiovascular system	114	
<i>Corbeton 40 (AF)</i>		
.Cardiovascular system	114	
<i>Cordarone X 100 (SW)</i>		
.Cardiovascular system	108	
<i>Cordarone X 200 (SW)</i>		
.Cardiovascular system	108	
<i>Cordilox 180 SR (KN)</i>		
.Cardiovascular system	120	
<i>Cordilox SR (KN)</i>		
.Cardiovascular system	120	
<i>Cortate (AS)</i>		
.Systemic hormonal preparations, excl. sex hormones and insulins	169	
<i>Cortef (VT)</i>		
.Dermatologicals.....	152	
.Dental	490	
<i>Cortic-DS 1% (FM)</i>		
.Dermatologicals.....	152	
.Dental	490	
CORTISONE ACETATE		
.Systemic hormonal preparations, excl. sex hormones and insulins	169	
<i>Cortival 1/2 (FM)</i>		
.Dermatologicals.....	153	
<i>Cortival 1/5 (FM)</i>		
.Dermatologicals.....	152	
<i>Cosamide (AF)</i>		
.Antineoplastic and immunomodulating agents	232	
<i>Cosopt (MK)</i>		
.Sensory organs.....	442	
.Optometrical.....	517	
<i>Cosudex (AP)</i>		
.Antineoplastic and immunomodulating agents	232	
COTTON WOOL ROLL		
.Repatriation Schedule	866	
<i>Coumadin (SI)</i>		
.Blood and blood forming organs	97	
<i>Coveram (SE)</i>		
.Cardiovascular system	130	
<i>Coversyl 10mg (SE)</i>		
.Cardiovascular system	125	
<i>Coversyl 2.5mg (SE)</i>		
.Cardiovascular system	124	
<i>Coversyl 5mg (SE)</i>		
.Cardiovascular system	125	
<i>Coversyl Plus 5mg/1.25mg (SE)</i>		
.Cardiovascular system	129	
<i>Coversyl Plus LD 2.5mg/0.625mg (SE)</i>		
.Cardiovascular system	129	
<i>Creon 10,000 (SM)</i>		
.Alimentary tract and metabolism	86	
<i>Creon 25,000 (SM)</i>		
.Alimentary tract and metabolism	86	
<i>Creon 40,000 (SM)</i>		
.Alimentary tract and metabolism	86	
<i>Creon 5000 (SM)</i>		
.Alimentary tract and metabolism	86	
<i>Creon Micro (SM)</i>		
.Alimentary tract and metabolism	86	
<i>Crestor (AP)</i>		
.Cardiovascular system	140	
<i>Crinone 8% (SG)</i>		
.Section 100	796	
<i>Crixivan 400 mg (MK)</i>		
.Section 100	562, 699	
<i>Cromolux (AE)</i>		
.Sensory organs.....	444	
.Optometrical.....	518	
<i>Crysanal (MD)</i>		
.Musculo-skeletal system	361	
.Palliative Care	478, 479	
.Dental	503	
<i>Curam (SZ)</i>		
.Anti-infectives for systemic use	180	
.Dental	495	
<i>Curam Duo (SZ)</i>		
.Anti-infectives for systemic use	180	
.Dental	495	
<i>Curam Duo 500/125 (SZ)</i>		
.Anti-infectives for systemic use	180	
.Dental	495	
<i>Curam Duo Forte 875/125 (SZ)</i>		
.Anti-infectives for systemic use	180	
.Dental	495	

<i>Curity 4112 (KE)</i>		
.Repatriation Schedule	869	
<i>Cutifilm Plus 76308 (SN)</i>		
.Repatriation Schedule	867	
<i>Cutifilm Plus 76309 (SN)</i>		
.Repatriation Schedule	867	
<i>Cutilin Non-Stick Wound Pad 76300 (SN)</i>		
.Repatriation Schedule	873	
<i>Cutilin Non-Stick Wound Pad 76301 (SN)</i>		
.Repatriation Schedule	873	
<i>Cutinova Hydro 66047441 (SN)</i>		
.Repatriation Schedule	870	
<i>Cutinova Hydro 66047443 (SN)</i>		
.Repatriation Schedule	870	
<i>Cycloblastin (PF)</i>		
.Antineoplastic and immunomodulating agents	199	
CYCLOPHOSPHAMIDE		
.Antineoplastic and immunomodulating agents	199	
CYCLOSPORIN		
.Antineoplastic and immunomodulating agents	355	
.Section 100	640, 776, 777	
<i>Cyklokapron (PF)</i>		
.Blood and blood forming organs	104	
<i>Cymbalta (LY)</i>		
.Nervous system	417	
<i>Cymevene (RO)</i>		
.Section 100	561, 697	
CYPROHEPTADINE HYDROCHLORIDE		
.Nervous system	386	
<i>Cyprohexal (SZ)</i>		
.Genito urinary system and sex hormones	163, 164	
.Antineoplastic and immunomodulating agents	232	
<i>Cyprone (AF)</i>		
.Genito urinary system and sex hormones	163, 164	
.Antineoplastic and immunomodulating agents	232	
<i>Cyprostat (SY)</i>		
.Genito urinary system and sex hormones	164	
.Antineoplastic and immunomodulating agents	232	
<i>Cyprostat-100 (SY)</i>		
.Genito urinary system and sex hormones	164	
.Antineoplastic and immunomodulating agents	232	
CYPROTERONE ACETATE		
.Genito urinary system and sex hormones	163, 164	
.Antineoplastic and immunomodulating agents	232	
<i>Cystine Amino Acid Supplement (VF)</i>		
.Various	465	
CYSTINE with CARBOHYDRATE		
.Various	465	
CYTARABINE		
.Antineoplastic and immunomodulating agents	202	
<i>Cytotec (PF)</i>		
.Alimentary tract and metabolism	70	
D		
<i>D-Penamine (AL)</i>		
.Musculo-skeletal system	362	
DABIGATRAN ETEXILATE		
.Blood and blood forming organs	102	
<i>Daivobet (CS)</i>		
.Dermatologicals	151	
<i>Daivonex (CS)</i>		
.Dermatologicals	151	
<i>Daktarin (JT)</i>		
.Dermatologicals	149	
.Repatriation Schedule	842	
<i>Dalacin C (PF)</i>		
.Antifectives for systemic use	186	
.Dental	499	
DALTEPARIN SODIUM (Low Molecular Weight Heparin Sodium—porcine mucous)		
.Blood and blood forming organs	97	
DANAZOL		
.Genito urinary system and sex hormones	164	
<i>Dantrium (PF)</i>		
.Musculo-skeletal system	364	
DANTROLENE SODIUM		
.Musculo-skeletal system	364	
<i>Daonil (SW)</i>		
.Alimentary tract and metabolism	89	
<i>Dapa-Tabs (AF)</i>		
.Cardiovascular system	112	
DAPSONE		
.Dermatologicals	155	
.Antifectives for systemic use	193	
<i>Daraprim (GK)</i>		
.Antiparasitic products, insecticides and repellents	430	
DARBEPOETIN ALFA		
.Section 100	524, 656	
DARUNAVIR		
.Section 100	562, 698	
DASATINIB		
.Antineoplastic and immunomodulating agents	210, 211, 212	
<i>DBL Aspirin 100 mg (YT)</i>		
.Blood and blood forming organs	98	
<i>DBL Ceftriaxone (HH)</i>		
.Antifectives for systemic use	184	
<i>DBL Fluconazole (HH)</i>		
.Antifectives for systemic use	191	
<i>DBL Gabapentin (HH)</i>		
.Nervous system	390, 391	
.Repatriation Schedule	853, 854	
<i>DBL Gemcitabine for Injection (HH)</i>		
.Antineoplastic and immunomodulating agents	202, 203	
<i>DBL Metronidazole Intravenous Infusion (HH)</i>		
.Antifectives for systemic use	189	
.Dental	499	
<i>DBL Oxaliplatin Concentrate (HH)</i>		
.Antineoplastic and immunomodulating agents	208	
<i>Deca-Durabolin (SH)</i>		
.Alimentary tract and metabolism	96	
DEFERASIROX		
.Section 100	654, 791	
DEFERIPRONE		
.Section 100	654, 791	
DEGARELIX		
.Antineoplastic and immunomodulating agents	234	
DENOSUMAB		
.Musculo-skeletal system	372	
<i>Depo-Medrol (PF)</i>		

.Systemic hormonal preparations, excl. sex hormones and insulins	170
.Dental	491
<i>Depo-Nisalone (KR)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	170
.Dental	491
<i>Depo-Provera (PF)</i>	
.Genito urinary system and sex hormones	158
<i>Depo-Ralovera (KR)</i>	
.Genito urinary system and sex hormones	158
<i>Deptran 10 (AF)</i>	
.Nervous system	411
<i>Deptran 25 (AF)</i>	
.Nervous system	411
<i>Deptran 50 (AF)</i>	
.Nervous system	411
<i>Deralin 10 (AF)</i>	
.Cardiovascular system	114
<i>Deralin 160 (AF)</i>	
.Cardiovascular system	114
<i>Deralin 40 (AF)</i>	
.Cardiovascular system	114
<i>Deseril (LM)</i>	
.Nervous system	385
<i>Desferal 2 g (NV)</i>	
.Section 100	654, 791
<i>Desferal 500 mg (NV)</i>	
.Section 100	654, 791
<i>DESFERRIOXAMINE MESYLATE</i>	
.Section 100	654, 791
<i>DESMOPRESSIN ACETATE</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	167, 168
<i>DESVENLAFAXINE SUCCINATE</i>	
.Nervous system	417
<i>DEXAMETHASONE</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	169
.Sensory organs	441
<i>DEXAMETHASONE SODIUM PHOSPHATE</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	169
<i>DEXAMETHASONE with FRAMYCETIN SULFATE and GRAMICIDIN</i>	
.Sensory organs	451
<i>DEXAMPHETAMINE SULFATE</i>	
.Nervous system	419
<i>Dexmethsone (AS)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	169
<i>DEXTROPROPOXYPHENE NAPSYLATE</i>	
.Repatriation Schedule	853
<i>Diabex (AL)</i>	
.Alimentary tract and metabolism	89
<i>Diabex 1000 (AL)</i>	
.Alimentary tract and metabolism	89
<i>Diabex 850 (AL)</i>	
.Alimentary tract and metabolism	88
<i>Diabex XR (AL)</i>	
.Alimentary tract and metabolism	89
<i>Diabex XR 1000 (AL)</i>	
.Alimentary tract and metabolism	89
<i>Diaformin (AF)</i>	
.Alimentary tract and metabolism	89
<i>Diaformin 1000 (AF)</i>	
.Alimentary tract and metabolism	89
<i>Diaformin 850 (AF)</i>	
.Alimentary tract and metabolism	88
<i>Diaformin XR (AF)</i>	
.Alimentary tract and metabolism	89
<i>Dialamine (SB)</i>	
.Various	465
<i>Diamicron 60mg MR (SE)</i>	
.Alimentary tract and metabolism	90
<i>Diamox (SI)</i>	
.Sensory organs	442
<i>Diapride 1 (SI)</i>	
.Alimentary tract and metabolism	90
<i>Diapride 2 (SI)</i>	
.Alimentary tract and metabolism	90
<i>Diapride 3 (SI)</i>	
.Alimentary tract and metabolism	90
<i>Diapride 4 (SI)</i>	
.Alimentary tract and metabolism	90
<i>Diastix (BN)</i>	
.Various	455
<i>DIAZEPAM</i>	
.Nervous system	408
.Palliative Care	483, 484
.Dental	509
<i>Diazepam-GA (GM)</i>	
.Nervous system	408
.Palliative Care	483, 484
.Dental	509
<i>Dibenyline (GH)</i>	
.Cardiovascular system	114
.Genito urinary system and sex hormones	165
<i>Dibenzyliline (GH)</i>	
.Cardiovascular system	114
.Genito urinary system and sex hormones	165
<i>Dicarz (AF)</i>	
.Cardiovascular system	117
<i>DICHLOROBENZENE with CHLORBUTOL and TURPENTINE OIL</i>	
.Repatriation Schedule	859
<i>Diclocil (BQ)</i>	
.Antiinfectives for systemic use	179
.Dental	494
<i>Diclofenac Sandoz (SZ)</i>	
.Musculo-skeletal system	358
.Palliative Care	475, 476
.Dental	501
<i>DICLOFENAC SODIUM</i>	
.Musculo-skeletal system	358
.Palliative Care	475, 476
.Dental	501
.Repatriation Schedule	846
<i>DICLOFENAC SODIUM with MISOPROSTOL</i>	
.Repatriation Schedule	852

.Antineoplastic and immunomodulating agents . 204, 205	
DOCUSATE SODIUM	
.Repatriation Schedule	837, 859
DOCUSATE SODIUM with SENNA	
.Repatriation Schedule	838
Dolapril (CO)	
.Nervous system	375
.Dental	504
Dolapril 0.5 (SI)	
.Cardiovascular system	128
Dolapril 1 (SI)	
.Cardiovascular system	128
Dolapril 2 (SI)	
.Cardiovascular system	128
Dolapril 4 (SI)	
.Cardiovascular system	128
DOLASETRON MESYLATE	
.Alimentary tract and metabolism	76
Doloxene (AS)	
.Repatriation Schedule	853
DOMPERIDONE	
.Alimentary tract and metabolism	76
DONEPEZIL HYDROCHLORIDE	
.Nervous system	421
DORNASE ALFA	
.Section 100	651, 788
Doryx (YN)	
.Antiinfectives for systemic use	175, 176
.Dental	492
DORZOLAMIDE HYDROCHLORIDE	
.Sensory organs	442
.Optometrical	517
DORZOLAMIDE HYDROCHLORIDE with TIMOLOL MALEATE	
.Sensory organs	442
.Optometrical	517
Dostinex (PF)	
.Genito urinary system and sex hormones	156, 157
Dothep 25 (AF)	
.Nervous system	411
Dothep 75 (AF)	
.Nervous system	411
DOTHIEPIN HYDROCHLORIDE	
.Nervous system	411
Douglas Cefaclor-CD (GM)	
.Antiinfectives for systemic use	182
.Dental	497
Douglas Gabapentin 300mg (GN)	
.Repatriation Schedule	853
Douglas Gabapentin 400mg (GN)	
.Nervous system	390
.Repatriation Schedule	854
DOXEPIN HYDROCHLORIDE	
.Nervous system	411
Doxorubicin Ebewe (IT)	
.Antineoplastic and immunomodulating agents	206
DOXORUBICIN HYDROCHLORIDE	
.Antineoplastic and immunomodulating agents	206
DOXORUBICIN HYDROCHLORIDE, PEGYLATED LIPOSOMAL	
.Antineoplastic and immunomodulating agents	206
.Section 100	569, 706
Doxsig (SI)	
.Antiinfectives for systemic use	175, 176
.Dental	492
Doxy-100 (GM)	
.Antiinfectives for systemic use	175, 176
.Dental	492
Doxy-50 (GM)	
.Antiinfectives for systemic use	175
DOXYCYCLINE	
.Antiinfectives for systemic use	175, 176
.Dental	492
Doxyhexal (SZ)	
.Antiinfectives for systemic use	175, 176
.Dental	492
Doxylin 100 (AF)	
.Antiinfectives for systemic use	175, 176
.Dental	492
Doxylin 50 (AF)	
.Antiinfectives for systemic use	175
DRESSING with CADEXOMER IODINE	
.Repatriation Schedule	874
DRESSING with SILVER	
.Repatriation Schedule	874
DRESSING—ACTIVATED CHARCOAL (MALODOROUS WOUND)	
.Repatriation Schedule	866
DRESSING—ALGINATE (CAVITY WOUND)	
.Repatriation Schedule	866
DRESSING—ALGINATE (SUPERFICIAL WOUND)	
.Repatriation Schedule	866
DRESSING—FILM	
.Repatriation Schedule	867
DRESSING—FILM ISLAND	
.Repatriation Schedule	867
DRESSING—FOAM with CHARCOAL (MALODOROUS WOUND)	
.Repatriation Schedule	868
DRESSING—FOAM with SILICONE—HEAVY EXUDATE	
.Repatriation Schedule	868
DRESSING—FOAM with SILICONE—LIGHT EXUDATE	
.Repatriation Schedule	868
DRESSING—FOAM with SILICONE—MODERATE EXUDATE	
.Repatriation Schedule	869
DRESSING—FOAM—HEAVY EXUDATE	
.Repatriation Schedule	867, 868
DRESSING—FOAM—MODERATE EXUDATE	
.Repatriation Schedule	868
DRESSING—GAUZE (ABSORBENT PAD)	
.Repatriation Schedule	869
DRESSING—GAUZE—EYE PAD	
.Repatriation Schedule	869
DRESSING—GAUZE—PARAFFIN	
.Repatriation Schedule	869
DRESSING—GAUZE—PARAFFIN with CHLORHEXIDINE ACETATE	
.Repatriation Schedule	869
DRESSING—HYDROACTIVE (CAVITY WOUND)	
.Repatriation Schedule	869
DRESSING—HYDROACTIVE (DEBRIDEMENT)	
.Repatriation Schedule	869

<i>DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—HIGH EXUDATE)</i>	
.Repatriation Schedule	869, 870
<i>DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—LIGHT EXUDATE)</i>	
.Repatriation Schedule	870
<i>DRESSING—HYDROACTIVE (SUPERFICIAL WOUND—MODERATE EXUDATE)</i>	
.Repatriation Schedule	870
<i>DRESSING—HYDROCOLLOID (CAVITY WOUND)</i>	
.Repatriation Schedule	870
<i>DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—LIGHT EXUDATE)</i>	
.Repatriation Schedule	870, 871
<i>DRESSING—HYDROCOLLOID (SUPERFICIAL WOUND—MODERATE EXUDATE)</i>	
.Repatriation Schedule	871, 872
<i>DRESSING—HYDROFIBRE (ALTERNATE TO ALGINATES)</i>	
.Repatriation Schedule	872
<i>DRESSING—HYDROGEL—AMORPHOUS</i>	
.Repatriation Schedule	872
<i>DRESSING—HYDROGEL—SHEET</i>	
.Repatriation Schedule	873
<i>DRESSING—NON-ADHERENT</i>	
.Repatriation Schedule	873
<i>DRESSING—TULLE NON-GAUZE—PARAFFIN</i>	
.Repatriation Schedule	873
<i>Drixine (SH)</i>	
.Repatriation Schedule	857
<i>Dronalen Plus (GM)</i>	
.Musculo-skeletal system	369
<i>Dronalen Plus D-Cal (FR)</i>	
.Musculo-skeletal system	370
<i>DROTRECOGIN ALFA (ACTIVATED)</i>	
.Blood and blood forming organs	101
<i>Dulcolax (BY)</i>	
.Alimentary tract and metabolism	80
.Palliative Care	471
<i>DULOXETINE HYDROCHLORIDE</i>	
.Nervous system	417
<i>Duocal (SB)</i>	
.Various	466
<i>DuoCover (BQ)</i>	
.Blood and blood forming organs	99
<i>DuoDERM CGF H7660 (CC)</i>	
.Repatriation Schedule	871
<i>DuoDERM CGF H7662 (CC)</i>	
.Repatriation Schedule	871
<i>DuoDERM Extra Thin H7955 (CC)</i>	
.Repatriation Schedule	871
<i>DuoDERM Gel H7987 (CC)</i>	
.Repatriation Schedule	872
<i>DuoDERM Gel H7990 (CC)</i>	
.Repatriation Schedule	872
<i>DuoDERM Paste H7930 (CC)</i>	
.Repatriation Schedule	870
<i>Duofilm Gel (GK)</i>	
.Repatriation Schedule	846
<i>Duofilm Solution (GK)</i>	
.Repatriation Schedule	846
<i>Duotrav (AQ)</i>	
.Sensory organs	443
.Optometrical	518
<i>Duphalac (SM)</i>	
.Alimentary tract and metabolism	80
.Palliative Care	472
<i>Duphaston (SM)</i>	
.Genito urinary system and sex hormones	161
<i>Duratears (AQ)</i>	
.Sensory organs	448, 449
.Optometrical	520
<i>Duride (AF)</i>	
.Cardiovascular system	110
<i>Duro-K (NM)</i>	
.Alimentary tract and metabolism	95
<i>Duro-Tuss (IA)</i>	
.Repatriation Schedule	858
<i>Durogesic 100 (JC)</i>	
.Nervous system	381
<i>Durogesic 12 (JC)</i>	
.Nervous system	381
<i>Durogesic 25 (JC)</i>	
.Nervous system	381
<i>Durogesic 50 (JC)</i>	
.Nervous system	381
<i>Durogesic 75 (JC)</i>	
.Nervous system	381
<i>Durotram XR (IA)</i>	
.Nervous system	383
.Dental	507
<i>DYDROGESTERONE</i>	
.Genito urinary system and sex hormones	161
<i>Dysport (IS)</i>	
.Section 100	793
E	
<i>E-Mycin (AF)</i>	
.Anti-infectives for systemic use	186
.Dental	498
<i>E-Mycin 200 (AF)</i>	
.Anti-infectives for systemic use	186
.Dental	498
<i>E-Mycin 400 (AF)</i>	
.Anti-infectives for systemic use	186
.Dental	498
<i>E.E.S. 200 (LM)</i>	
.Anti-infectives for systemic use	186
.Dental	498
<i>E.E.S. 400 Filmtab (LM)</i>	
.Anti-infectives for systemic use	186
.Dental	498
<i>E.E.S. Granules (LM)</i>	
.Anti-infectives for systemic use	186
.Dental	498
<i>EAA Supplement (VF)</i>	
.Various	465
<i>Ear Clear for Ear Wax Removal (KY)</i>	
.Repatriation Schedule	859
<i>Easiphen (SB)</i>	
.Various	464

<i>Ebixa (LU)</i>		
.Nervous system	426	
<i>Edecrin (FK)</i>		
.Cardiovascular system	113	
<i>Edronax (PF)</i>		
.Nervous system	418	
<i>EFAVIRENZ</i>		
.Section 100	566, 703	
<i>Efexor-XR (WX)</i>		
.Nervous system	418	
<i>Effient (LY)</i>		
.Blood and blood forming organs	100	
<i>EFORMOTEROL FUMARATE DIHYDRATE</i>		
.Respiratory system	433	
<i>Efudix (VT)</i>		
.Repatiation Schedule	850	
<i>Egoderm Cream (EO)</i>		
.Repatiation Schedule	846	
<i>Egoderm Ointment (EO)</i>		
.Repatiation Schedule	846	
<i>Egopsoryl-TA (EO)</i>		
.Repatiation Schedule	844	
<i>Elastoplast 2225 (BE)</i>		
.Repatiation Schedule	865	
<i>Elastoplast 2226 (BE)</i>		
.Repatiation Schedule	865	
<i>Elastoplast 2227 (BE)</i>		
.Repatiation Schedule	865	
<i>Eldepryl (AS)</i>		
.Nervous system	399	
<i>EleCare (AB)</i>		
.Various.....	457, 458	
<i>EleCare LCP (AB)</i>		
.Various.....	459	
<i>ELECTROLYTE REPLACEMENT (ORAL)</i>		
.Alimentary tract and metabolism.....	82	
<i>ELECTROLYTE REPLACEMENT SOLUTION</i>		
.Blood and blood forming organs	105	
<i>ELETRIPTAN</i>		
.Nervous system	385	
<i>Eleuphrat (EX)</i>		
.Dermatologicals.....	152	
<i>Eleva 100 (AF)</i>		
.Nervous system	415, 416	
<i>Eleva 50 (AF)</i>		
.Nervous system	415, 416	
<i>Elidel (NV)</i>		
.Dermatologicals.....	154	
<i>Eligard 1 month (HH)</i>		
.Antineoplastic and immunomodulating agents	231	
<i>Eligard 3 month (HH)</i>		
.Antineoplastic and immunomodulating agents	231	
<i>Eligard 4 month (HH)</i>		
.Antineoplastic and immunomodulating agents	231	
<i>Eligard 6 month (HH)</i>		
.Antineoplastic and immunomodulating agents	231	
<i>Elocon (SH)</i>		
.Dermatologicals.....	153, 154	
.Repatiation Schedule	845	
<i>Eloxatin (SW)</i>		
.Antineoplastic and immunomodulating agents	208	
<i>Emend (MK)</i>		
.Alimentary tract and metabolism.....	79	
<i>EMTRICITABINE</i>		
.Section 100	564, 700	
<i>Emtriva (GI)</i>		
.Section 100	564, 700	
<i>Enahexal (HX)</i>		
.Cardiovascular system	122	
<i>Enalabell (BF)</i>		
.Cardiovascular system	122	
<i>ENALAPRIL</i>		
.Cardiovascular system	122	
<i>Enalapril generichealth (GQ)</i>		
.Cardiovascular system	122	
<i>ENALAPRIL MALEATE with HYDROCHLOROTHIAZIDE</i>		
.Cardiovascular system	128	
<i>Enalapril Sandoz (SZ)</i>		
.Cardiovascular system	122, 123	
<i>Enalapril Winthrop (WA)</i>		
.Cardiovascular system	122, 123	
<i>Enalapril-DP 10mg (GN)</i>		
.Cardiovascular system	122	
<i>Enalapril-DP 20mg (GN)</i>		
.Cardiovascular system	122	
<i>Enalapril-DP 5mg (GN)</i>		
.Cardiovascular system	122	
<i>Enalapril-GA (GM)</i>		
.Cardiovascular system	122	
<i>Enalapril/HCT Sandoz (SZ)</i>		
.Cardiovascular system	128	
<i>Enbrel (WX)</i>		
.Antineoplastic and immunomodulating agents 295, 298,		
301, 304, 307, 310, 315, 318, 322, 325		
.Section 100	590, 593, 728, 730	
<i>Endep 10 (AF)</i>		
.Nervous system	411	
<i>Endep 25 (AF)</i>		
.Nervous system	411	
<i>Endep 50 (AF)</i>		
.Nervous system	411	
<i>Endone (SI)</i>		
.Nervous system	380	
.Dental	506	
<i>Endoxan (BX)</i>		
.Antineoplastic and immunomodulating agents	199	
<i>Energivit (SB)</i>		
.Various.....	465	
<i>ENFUVIRTIDE</i>		
.Section 100	568, 705	
<i>Enidin (PE)</i>		
.Sensory organs.....	442	
.Optometrical.....	516	
<i>ENOXAPARIN SODIUM</i>		
.Blood and blood forming organs	97	
<i>ENTACAPONE</i>		
.Nervous system	399	
<i>ENTECAVIR MONOHYDRATE</i>		
.Section 100	564, 700, 701	
<i>Epilim (SW)</i>		

.Nervous system	389	.Cardiovascular system	131
<i>Epilim EC (SW)</i>		<i>EPTIFIBATIDE ACETATE</i>	
.Nervous system	389	.Blood and blood forming organs	100
<i>Epilim Liquid (SW)</i>		<i>Erbix (SG)</i>	
.Nervous system	389	.Antineoplastic and immunomodulating agents	209
<i>Epilim Syrup (SW)</i>		<i>ERLOTINIB</i>	
.Nervous system	389	.Antineoplastic and immunomodulating agents	213
<i>EpiPen (AL)</i>		<i>Eryc (YN)</i>	
.Cardiovascular system	109	.Antiinfectives for systemic use	185
.Respiratory system	437	.Dental	498
<i>EpiPen Jr. (AL)</i>		<i>Erythrocin-I.V. (LM)</i>	
.Cardiovascular system	109	.Antiinfectives for systemic use	186
.Respiratory system	437	.Dental	498
<i>Epiramax 100 (SI)</i>		<i>ERYTHROMYCIN</i>	
.Nervous system	395	.Antiinfectives for systemic use	185
<i>Epiramax 200 (SI)</i>		.Dental	498
.Nervous system	395	<i>ERYTHROMYCIN ETHYL SUCCINATE</i>	
<i>Epiramax 25 (SI)</i>		.Antiinfectives for systemic use	186
.Nervous system	394	.Dental	498
<i>Epiramax 50 (SI)</i>		<i>ERYTHROMYCIN LACTOBIONATE</i>	
.Nervous system	394	.Antiinfectives for systemic use	186
<i>Epirubicin Ebewe (IT)</i>		.Dental	498
.Antineoplastic and immunomodulating agents	206	<i>ESCITALOPRAM OXALATE</i>	
<i>EPIRUBICIN HYDROCHLORIDE</i>		.Nervous system	412, 413
.Antineoplastic and immunomodulating agents	206	<i>Esipram (GM)</i>	
<i>EPLERENONE</i>		.Nervous system	413
.Cardiovascular system	113	<i>Esitalo (SZ)</i>	
<i>EPOETIN ALFA</i>		.Nervous system	413
.Section 100	524, 656	<i>Eskazole (GK)</i>	
<i>EPOETIN BETA</i>		.Antiparasitic products, insecticides and repellents	432
.Section 100	524, 656	<i>ESOMEPRAZOLE MAGNESIUM TRIHYDRATE</i>	
<i>EPOETIN LAMBDA</i>		.Alimentary tract and metabolism	70, 71
.Section 100	525, 657	<i>ESOMEPRAZOLE MAGNESIUM TRIHYDRATE and</i> <i>CLARITHROMYCIN and AMOXYCILLIN</i>	
<i>EPOPROSTENOL SODIUM</i>		.Alimentary tract and metabolism	75
.Section 100	538, 670, 675	<i>Essential Amino Acid Mix (SB)</i>	
<i>Eprex 1000 (JC)</i>		.Various	465
.Section 100	524, 656	<i>ESSENTIAL AMINO ACIDS FORMULA</i>	
<i>Eprex 10000 (JC)</i>		.Various	465
.Section 100	524, 656	<i>ESSENTIAL AMINO ACIDS FORMULA with MINERALS and</i> <i>VITAMIN C</i>	
<i>Eprex 20,000 (JC)</i>		.Various	465
.Section 100	524, 656	<i>ESSENTIAL AMINO ACIDS FORMULA with VITAMINS and</i> <i>MINERALS</i>	
<i>Eprex 2000 (JC)</i>		.Various	465
.Section 100	524, 656	<i>Estalis continuous 50/140 (NV)</i>	
<i>Eprex 30,000 (JC)</i>		.Genito urinary system and sex hormones	161
.Section 100	524, 656	<i>Estalis continuous 50/250 (NV)</i>	
<i>Eprex 3000 (JC)</i>		.Genito urinary system and sex hormones	161
.Section 100	524, 656	<i>Estalis sequi 50/140 (NV)</i>	
<i>Eprex 40,000 (JC)</i>		.Genito urinary system and sex hormones	162
.Section 100	524, 656	<i>Estalis sequi 50/250 (NV)</i>	
<i>Eprex 4000 (JC)</i>		.Genito urinary system and sex hormones	162
.Section 100	524, 656	<i>Estraderm 100 (NV)</i>	
<i>Eprex 5000 (JC)</i>		.Genito urinary system and sex hormones	159
.Section 100	524, 656	<i>Estraderm 25 (NV)</i>	
<i>Eprex 6000 (JC)</i>		.Genito urinary system and sex hormones	159
.Section 100	524, 656	<i>Estraderm MX 100 (NV)</i>	
<i>Eprex 8000 (JC)</i>		.Genito urinary system and sex hormones	160
.Section 100	524, 656	<i>Estraderm MX 25 (NV)</i>	
<i>EPROSARTAN MESYLATE</i>			
.Cardiovascular system	130		
<i>EPROSARTAN MESYLATE with HYDROCHLOROTHIAZIDE</i>			

.Genito urinary system and sex hormones.....	160
<i>Estraderm MX 50 (NV)</i>	
.Genito urinary system and sex hormones.....	160
<i>Estradot 100 (NV)</i>	
.Genito urinary system and sex hormones.....	160
<i>Estradot 25 (NV)</i>	
.Genito urinary system and sex hormones.....	160
<i>Estradot 37.5 (NV)</i>	
.Genito urinary system and sex hormones.....	160
<i>Estradot 50 (NV)</i>	
.Genito urinary system and sex hormones.....	160
<i>Estradot 75 (NV)</i>	
.Genito urinary system and sex hormones.....	160
ETANERCEPT	
.Antineoplastic and immunomodulating agents	290, 295, 298, 301, 304, 307, 310, 315, 319, 322
.Section 100	586, 590, 724, 728
ETHACRYNIC ACID	
.Cardiovascular system.....	112
ETHOSUXIMIDE	
.Nervous system	387
ETONOGESTREL	
.Genito urinary system and sex hormones.....	158
<i>Etopophos (BQ)</i>	
.Antineoplastic and immunomodulating agents	204
ETOPOSIDE	
.Antineoplastic and immunomodulating agents	204
<i>Etoposide Ebewe (IT)</i>	
.Antineoplastic and immunomodulating agents	204
ETRAVIRINE	
.Section 100	566, 703
<i>Eucerin (BE)</i>	
.Repatriation Schedule	843
<i>Eulexin (SH)</i>	
.Antineoplastic and immunomodulating agents	233
<i>Eutroxsig (FM)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	171
EVEROLIMUS	
.Antineoplastic and immunomodulating agents	236
.Section 100	580, 718
<i>Evista (LY)</i>	
.Genito urinary system and sex hormones.....	165
.Musculo-skeletal system	372
<i>Exelon (NV)</i>	
.Nervous system	425
<i>Exelon Patch 10 (NV)</i>	
.Nervous system	424
<i>Exelon Patch 5 (NV)</i>	
.Nervous system	424
EXEMESTANE	
.Antineoplastic and immunomodulating agents	233
EXENATIDE	
.Alimentary tract and metabolism.....	94
<i>Exforge 10/160 (NV)</i>	
.Cardiovascular system.....	133
<i>Exforge 10/320 (NV)</i>	
.Cardiovascular system.....	132
<i>Exforge 5/160 (NV)</i>	
.Cardiovascular system.....	133
<i>Exforge 5/320 (NV)</i>	
.Cardiovascular system	132
<i>Exforge 5/80 (NV)</i>	
.Cardiovascular system	132
<i>Exforge HCT 10/160/12.5 (NV)</i>	
.Cardiovascular system	133
<i>Exforge HCT 10/160/25 (NV)</i>	
.Cardiovascular system	133
<i>Exforge HCT 10/320/25 (NV)</i>	
.Cardiovascular system	133
<i>Exforge HCT 5/160/12.5 (NV)</i>	
.Cardiovascular system	133
<i>Exforge HCT 5/160/25 (NV)</i>	
.Cardiovascular system	133
<i>Exjade (NV)</i>	
.Section 100	654, 791
<i>Exorex (GM)</i>	
.Dermatologicals	150
<i>Extine 20 (SI)</i>	
.Nervous system	415
EZETIMIBE	
.Cardiovascular system	146
EZETIMIBE with SIMVASTATIN	
.Cardiovascular system	147
<i>Ezetrol (MK)</i>	
.Cardiovascular system	147
<i>Ezovir (AF)</i>	
.Antiinfectives for systemic use	195, 196, 197
F	
FAMCICLOVIR	
.Antiinfectives for systemic use	195, 196
<i>Famciclovir Sandoz (SZ)</i>	
.Antiinfectives for systemic use	195, 196
<i>Famohexal (HX)</i>	
.Alimentary tract and metabolism.....	69
FAMOTIDINE	
.Alimentary tract and metabolism.....	68
<i>Famotidine Sandoz (SZ)</i>	
.Alimentary tract and metabolism.....	69
<i>Famvir (NV)</i>	
.Antiinfectives for systemic use	195, 196, 197
<i>Fareston (SH)</i>	
.Antineoplastic and immunomodulating agents	232
<i>Farine (WQ)</i>	
.Antineoplastic and immunomodulating agents	201
<i>Fasigyn (PF)</i>	
.Antiinfectives for systemic use	190
.Antiparasitic products, insecticides and repellents....	430
<i>Faverin 100 (SI)</i>	
.Nervous system	414
<i>Faverin 50 (SI)</i>	
.Nervous system	414
<i>Favic 125 (SI)</i>	
.Antiinfectives for systemic use	195
<i>Favic 250 (SI)</i>	
.Antiinfectives for systemic use	195, 196
<i>Favic 500 (SI)</i>	
.Antiinfectives for systemic use	196, 197
<i>Fawns and McAllan Proprietary Limited (FM)</i>	

.Antifungals for systemic use	193
.Nervous system	375, 387
.Respiratory system	439
.Dental	504
<i>Febidol (GM)</i>	
.Nervous system	384
.Dental	508, 509
<i>Feldene (PF)</i>	
.Musculo-skeletal system	360
.Dental	502
<i>Feldene-D (PF)</i>	
.Musculo-skeletal system	360
.Dental	502
<i>Felodil XR 10 (SI)</i>	
.Cardiovascular system	119
<i>Felodil XR 5 (SI)</i>	
.Cardiovascular system	119
<i>FELODIPINE</i>	
.Cardiovascular system	119
<i>Felodur ER 10 mg (AL)</i>	
.Cardiovascular system	119
<i>Felodur ER 2.5 mg (AL)</i>	
.Cardiovascular system	119
<i>Felodur ER 5 mg (AL)</i>	
.Cardiovascular system	119
<i>Femara 2.5 mg (NV)</i>	
.Antineoplastic and immunomodulating agents	234
<i>Femoston 2/10 (SM)</i>	
.Genito urinary system and sex hormones	162
<i>Fenac (AF)</i>	
.Musculo-skeletal system	358
.Palliative Care	475, 476
.Dental	501
<i>Fenac 25 (AF)</i>	
.Musculo-skeletal system	358
.Palliative Care	475, 476
.Dental	501
<i>FENOFIBRATE</i>	
.Cardiovascular system	144
<i>FENTANYL</i>	
.Nervous system	380
.Palliative Care	481
<i>Ferriprox (OA)</i>	
.Section 100	654, 791
<i>Ferro-f-tab (AE)</i>	
.Blood and blood forming organs	104
<i>Ferro-Liquid (AE)</i>	
.Blood and blood forming organs	104
<i>Ferrosig (SI)</i>	
.Blood and blood forming organs	104
<i>FERROUS FUMARATE with FOLIC ACID</i>	
.Blood and blood forming organs	104
<i>FERROUS SULFATE</i>	
.Blood and blood forming organs	104
<i>Ferrum H (AS)</i>	
.Blood and blood forming organs	104
<i>Fexal (SZ)</i>	
.Repatriation Schedule	858
<i>FEXOFENADINE HYDROCHLORIDE</i>	
.Repatriation Schedule	858
<i>Fibsol 10 (SI)</i>	
.Cardiovascular system	123
<i>Fibsol 20 (SI)</i>	
.Cardiovascular system	124
<i>Fibsol 5 (SI)</i>	
.Cardiovascular system	123
<i>FILGRASTIM</i>	
.Section 100	569, 706
<i>Filpril (AL)</i>	
.Cardiovascular system	125, 126
<i>Finasta (SZ)</i>	
.Repatriation Schedule	848
<i>FINASTERIDE</i>	
.Repatriation Schedule	848
<i>Firmagon 120mg (FP)</i>	
.Antineoplastic and immunomodulating agents	234
<i>Firmagon 80mg (FP)</i>	
.Antineoplastic and immunomodulating agents	234
<i>Flagyl (SW)</i>	
.Antifungals for systemic use	189
.Antiparasitic products, insecticides and repellents	430
.Dental	499
<i>Flagyl S (SW)</i>	
.Antifungals for systemic use	190
.Antiparasitic products, insecticides and repellents	430
.Dental	499
<i>Flamazine (SN)</i>	
.Dermatologicals	152
<i>Flarex (AQ)</i>	
.Sensory organs	441
.Optometrical	516
<i>FLECAINIDE ACETATE</i>	
.Cardiovascular system	107
<i>Flecatab (AF)</i>	
.Cardiovascular system	107
<i>Flexidress 650941 (CC)</i>	
.Repatriation Schedule	865
<i>Flixotide (GK)</i>	
.Respiratory system	436
<i>Flixotide Accuhaler (GK)</i>	
.Respiratory system	436
<i>Flixotide Junior (GK)</i>	
.Respiratory system	436
<i>Flixotide Junior Accuhaler (GK)</i>	
.Respiratory system	436
<i>Folan (GK)</i>	
.Section 100	543, 674, 679
<i>Flomaxtra (CS)</i>	
.Repatriation Schedule	848
<i>Flopen (AS)</i>	
.Antifungals for systemic use	179, 180
.Dental	494
<i>Florinef (SI)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	168
<i>Fluanxol Concentrated Depot (LU)</i>	
.Nervous system	401
<i>Fluanxol Depot (LU)</i>	
.Nervous system	401
<i>Flubiclox (TS)</i>	

.Antifungives for systemic use	179	.Nervous system	401
.Dental	494	FLUPHENAZINE DECANOATE	
Flucil (AS)		.Nervous system	400
.Antifungives for systemic use	179	FLURBIPROFEN SODIUM	
.Dental	494	.Sensory organs.....	441
FLUCLOXACILLIN		.Optometrical.....	516
.Antifungives for systemic use	179	FLUTAMIDE	
.Dental	494	.Antineoplastic and immunomodulating agents	232
Flucon (AQ)		Flutamin (AF)	
.Sensory organs	441	.Antineoplastic and immunomodulating agents	233
.Optometrical	516	FLUTICASONE PROPIONATE	
FLUCONAZOLE		.Respiratory system	436
.Antifungives for systemic use	191	FLUTICASONE PROPIONATE with SALMETEROL XINAFOATE	
Fluconazole Hexal (HX)		.Respiratory system	435
.Antifungives for systemic use	191	FLUVASTATIN	
Fluconazole Hexal (SZ)		.Cardiovascular system	137
.Antifungives for systemic use	191	FLUVOXAMINE MALEATE	
Fluconazole Sandoz (SZ)		.Nervous system	414
.Antifungives for systemic use	191	Fluzole 200 (SI)	
Fluconazole Winthrop (WA)		.Antifungives for systemic use	191
.Antifungives for systemic use	191	Fluzole 50 (SI)	
Fluconazole-Clarix (AE)		.Antifungives for systemic use	191
.Antifungives for systemic use	191	FML Liquifilm (AG)	
Fludara (GZ)		.Sensory organs.....	441
.Antineoplastic and immunomodulating agents	201	.Optometrical.....	516
Fludarabine Actavis (GQ)		FOLIC ACID	
.Antineoplastic and immunomodulating agents	201	.Blood and blood forming organs	104, 105
Fludarabine Ebewe (IT)		FOLLITROPIN ALFA	
.Antineoplastic and immunomodulating agents	201	.Genito urinary system and sex hormones	162
FLUDARABINE PHOSPHATE		.Section 100	796
.Antineoplastic and immunomodulating agents	201	FOLLITROPIN BETA	
FLUDROCORTISONE ACETATE		.Genito urinary system and sex hormones	162
.Systemic hormonal preparations, excl. sex hormones and insulins	168	.Section 100	796
FLUNITRAZEPAM		FONDAPARINUX SODIUM	
.Repatriation Schedule	854	.Blood and blood forming organs	103
Fluohexal (SZ)		Foradile (NV)	
.Nervous system	414	.Respiratory system	433
FLUOROMETHOLONE		Formet 1000 (SI)	
.Sensory organs	441	.Alimentary tract and metabolism	89
.Optometrical	516	Formet 500 (SI)	
FLUOROMETHOLONE ACETATE		.Alimentary tract and metabolism	89
.Sensory organs	441	Formet 850 (SI)	
.Optometrical	516	.Alimentary tract and metabolism	88
FLUOROURACIL		Forteo (LY)	
.Antineoplastic and immunomodulating agents	202	.Systemic hormonal preparations, excl. sex hormones and insulins.....	173
.Repatriation Schedule	850	.Musculo-skeletal system	373
Fluorouracil Ebewe (IT)		Fosamax 40 mg (MK)	
.Antineoplastic and immunomodulating agents	202	.Musculo-skeletal system	365
Fluoxebell (BF)		Fosamax Once Weekly (MK)	
.Nervous system	414	.Musculo-skeletal system	365
Fluoxetine 20 (CR)		Fosamax Plus (MK)	
.Nervous system	414	.Musculo-skeletal system	369
Fluoxetine generichealth (GQ)		Fosamax Plus 70 mg/140 mcg (MK)	
.Nervous system	414	.Musculo-skeletal system	369
FLUOXETINE HYDROCHLORIDE		Fosamax Plus D-Cal (MK)	
.Nervous system	414	.Musculo-skeletal system	370
Fluoxetine-GA (GM)		FOSAMPRENAVIR	
.Nervous system	414	.Section 100	562, 698
FLUPENTHIXOL DECANOATE		FOSCARNET SODIUM	

.Section 100	562, 698
<i>Foscavir (AP)</i>	
.Section 100	562, 698
<i>Fosetic 20/12.5 (ZP)</i>	
.Cardiovascular system	129
<i>Fosinopril Sandoz (SZ)</i>	
.Cardiovascular system	123
FOSINOPRIL SODIUM	
.Cardiovascular system	123
FOSINOPRIL SODIUM with HYDROCHLOROTHIAZIDE	
.Cardiovascular system	128
<i>Fosinopril/HCT Sandoz 10mg/12.5mg (SZ)</i>	
.Cardiovascular system	128
<i>Fosinopril/HCT Sandoz 20mg/12.5mg (SZ)</i>	
.Cardiovascular system	129
<i>Fosinopril/HCTZ-GA 10/12.5 (GM)</i>	
.Cardiovascular system	128
<i>Fosinopril/HCTZ-GA 20/12.5 (GM)</i>	
.Cardiovascular system	129
<i>Fosipril 10 (SI)</i>	
.Cardiovascular system	123
<i>Fosipril 20 (SI)</i>	
.Cardiovascular system	123
<i>Fosrenol (ZI)</i>	
.Various.....	452
.Section 100	654, 791
FOTEMUSTINE	
.Antineoplastic and immunomodulating agents	199
<i>Fragmin (PF)</i>	
.Blood and blood forming organs	97
<i>Frakas (SI)</i>	
.Antimicrobials for systemic use	175
FRAMYCETIN SULFATE	
.Sensory organs	451
.Optometrical	521
<i>FreeStyle Lite (MS)</i>	
.Various.....	456
<i>Freestyle Papillon (MS)</i>	
.Various.....	456
<i>Fresenius Kabi Australia Pty Limited (PK)</i>	
.Blood and blood forming organs	105, 106
.Dental	488
FRUSEMIDE	
.Cardiovascular system	112
<i>Frusemide Sandoz (SZ)</i>	
.Emergency Drug Supplies	64
.Cardiovascular system	112
<i>Frusemide-Clarix (AE)</i>	
.Emergency Drug Supplies	64
.Cardiovascular system	112
<i>Frusid (GM)</i>	
.Cardiovascular system	112
<i>Fucidin (CS)</i>	
.Antimicrobials for systemic use	189
<i>Fungilin (SI)</i>	
.Alimentary tract and metabolism	68
.Dental	486
FUSIDIC ACID	
.Antimicrobials for systemic use	189
<i>Fuzeon (RO)</i>	

.Section 100	568, 705
<i>Fybogel (RC)</i>	
.Repatriation Schedule	838
G	
<i>GA gel (VF)</i>	
.Various.....	462
<i>GA Tramadol 50mg (GM)</i>	
.Nervous system	382
.Dental	507
<i>GA Tramadol SR 100mg (GM)</i>	
.Nervous system	383
.Dental	507
<i>GA Tramadol SR 150mg (GM)</i>	
.Nervous system	383
.Dental	507
<i>GA Tramadol SR 200mg (GM)</i>	
.Nervous system	383
.Dental	508
<i>GA-Amclav 500/125 (GM)</i>	
.Antimicrobials for systemic use	180
.Dental	495
<i>GA-Amclav Forte 875/125 (GM)</i>	
.Antimicrobials for systemic use	180
.Dental	495
<i>GA1 Anamix infant (SB)</i>	
.Various.....	462
<i>Gabahexal 300mg (SZ)</i>	
.Repatriation Schedule	853
<i>Gabahexal 400mg (SZ)</i>	
.Repatriation Schedule	854
GABAPENTIN	
.Nervous system	390
.Repatriation Schedule	853
<i>Gabapentin 300 (CR)</i>	
.Nervous system	390
<i>Gabapentin 400 (CR)</i>	
.Nervous system	390
<i>Gabapentin Sandoz (SZ)</i>	
.Nervous system	390
<i>Gabapentin-GA (GM)</i>	
.Nervous system	390
<i>Gabaran (RA)</i>	
.Nervous system	390, 391
<i>Gabazine 100 (SI)</i>	
.Nervous system	391
.Repatriation Schedule	853
<i>Gabazine 300 (SI)</i>	
.Nervous system	390
.Repatriation Schedule	853
<i>Gabazine 400 (SI)</i>	
.Nervous system	390
.Repatriation Schedule	854
<i>Gabazine 600 (SI)</i>	
.Nervous system	391
<i>Gabazine 800 (SI)</i>	
.Nervous system	391
<i>Gabitril (OA)</i>	
.Nervous system	390
GALANTAMINE HYDROBROMIDE	

.Nervous system	422	<i>Genotropin GoQuick (PF)</i>	
<i>Galantyl (AF)</i>		.Section 100	795
.Nervous system	423	<i>Genotropin MiniQuick (PF)</i>	
<i>Galvus (NV)</i>		.Section 100	794, 795
.Alimentary tract and metabolism.....	94	<i>Genox 10 (AF)</i>	
GANCICLOVIR		.Antineoplastic and immunomodulating agents	231
.Section 100	561, 653, 697, 790	<i>Genox 20 (AF)</i>	
<i>Ganfort 0.3/5 (AG)</i>		.Antineoplastic and immunomodulating agents	231
.Sensory organs	443	<i>GenRx Aciclovir (GX)</i>	
.Optometrical	518	.Antiinfectives for systemic use	194, 195
GANIRELIX		<i>GenRx Allopurinol (GX)</i>	
.Section 100	797	.Musculo-skeletal system	364
<i>Gantin (AW)</i>		<i>GenRx Alprazolam (GX)</i>	
.Nervous system	390, 391	.Nervous system	408
.Repatriation Schedule	853, 854	<i>GenRx Amiodarone (GX)</i>	
<i>Gastro-Stop Loperamide (AS)</i>		.Cardiovascular system	108
.Alimentary tract and metabolism.....	82	<i>GenRx Amoxycillin (GX)</i>	
<i>Gastrogel (FM)</i>		.Antiinfectives for systemic use	177, 178
.Alimentary tract and metabolism.....	68	.Dental	492, 493
GAUZE and COTTON TISSUE (COMBINE ROLL)		<i>GenRx Amoxycillin and Clavulanic Acid (GX)</i>	
.Repatriation Schedule	874	.Antiinfectives for systemic use	180
<i>Gaviscon P (RC)</i>		.Dental	495
.Alimentary tract and metabolism.....	75	<i>GenRx Azathioprine (GX)</i>	
GEFITINIB		.Antineoplastic and immunomodulating agents	357
.Antineoplastic and immunomodulating agents	213	<i>GenRx Baclofen (GX)</i>	
GELATIN - SUCCINYLATED		.Musculo-skeletal system	363
.Blood and blood forming organs	105	<i>GenRx Calcitriol (GX)</i>	
<i>Gelofusine (BR)</i>		.Alimentary tract and metabolism	95
.Blood and blood forming organs	105	.Musculo-skeletal system	371
<i>GelTears (BU)</i>		<i>GenRx Captopril (GX)</i>	
.Sensory organs	446	.Cardiovascular system	121, 122
.Optometrical	518	<i>GenRx Carvedilol (GX)</i>	
GEMCITABINE		.Cardiovascular system	117, 118
.Antineoplastic and immunomodulating agents .	202, 203	<i>GenRx Cefaclor (GX)</i>	
<i>Gemcitabine Actavis (GQ)</i>		.Antiinfectives for systemic use	183
.Antineoplastic and immunomodulating agents .	202, 203	.Dental	497
<i>Gemcitabine Ebewe (IT)</i>		<i>GenRx Cefaclor CD (GX)</i>	
.Antineoplastic and immunomodulating agents .	202, 203	.Antiinfectives for systemic use	182
<i>Gemcitabine Kabi (PK)</i>		.Dental	497
.Antineoplastic and immunomodulating agents .	202, 203	<i>GenRx Cephalexin (GX)</i>	
<i>Gemcitabine Sun (ZF)</i>		.Antiinfectives for systemic use	181, 182
.Antineoplastic and immunomodulating agents .	202, 203	.Dental	496
<i>Gemcite (ZP)</i>		<i>GenRx Cimetidine (GX)</i>	
.Antineoplastic and immunomodulating agents .	202, 203	.Alimentary tract and metabolism	68
GEMFIBROZIL		<i>GenRx Ciprofloxacin (GX)</i>	
.Cardiovascular system.....	145	.Antiinfectives for systemic use	187, 188
<i>Gemhexal (SZ)</i>		<i>GenRx Citalopram (GX)</i>	
.Cardiovascular system.....	145	.Nervous system	412
<i>Gemzar (LY)</i>		<i>GenRx Clarithromycin (GX)</i>	
.Antineoplastic and immunomodulating agents .	202, 203	.Antiinfectives for systemic use	185
<i>Generic Health Pty Ltd (GQ)</i>		<i>GenRx Clomipramine (GX)</i>	
.Nervous system	384	.Nervous system	409, 411
.Dental	508, 509	<i>GenRx Cyproterone Acetate (GX)</i>	
<i>Genlac (SI)</i>		.Genito urinary system and sex hormones	164
.Alimentary tract and metabolism.....	80	.Antineoplastic and immunomodulating agents	232
.Palliative Care	472	<i>GenRx Diltiazem (GX)</i>	
<i>Genoptic (AG)</i>		.Cardiovascular system	121
.Sensory organs	440	<i>GenRx Diltiazem CD (GX)</i>	
<i>Genotropin (PF)</i>		.Cardiovascular system	121
.Section 100	794, 795	<i>GenRx Doxycycline (GX)</i>	

.Antifungals for systemic use	175, 176	.Cardiovascular system	111
.Dental	492	<i>GenRx Ramipril (GX)</i>	
<i>GenRx Enalapril (GX)</i>		.Cardiovascular system	127
.Cardiovascular system	122, 123	<i>GenRx Ranitidine (GX)</i>	
<i>GenRx Famotidine (GX)</i>		.Alimentary tract and metabolism	69
.Alimentary tract and metabolism	69	<i>GenRx Salbutamol (GX)</i>	
<i>GenRx Fluoxetine (GX)</i>		.Emergency Drug Supplies	65
.Nervous system	414	.Respiratory system	433, 434
<i>GenRx Fosinopril (GX)</i>		<i>GenRx Sertraline (GX)</i>	
.Cardiovascular system	123	.Nervous system	415
<i>GenRx Frusemide (GX)</i>		<i>GenRx Simvastatin (GX)</i>	
.Cardiovascular system	112	.Cardiovascular system	140, 141, 142, 143, 144
<i>GenRx Gabapentin (GX)</i>		<i>GenRx Sotalol (GX)</i>	
.Nervous system	390, 391	.Cardiovascular system	108, 114, 115
.Repatriation Schedule	853, 854	<i>GenRx Tamoxifen (GX)</i>	
<i>GenRx Gemfibrozil (GX)</i>		.Antineoplastic and immunomodulating agents	231
.Cardiovascular system	145	<i>GenRx Terbinafine (GX)</i>	
<i>GenRx Gliclazide (GX)</i>		.Dermatologicals	150
.Alimentary tract and metabolism	89	<i>GenRx Tramadol (GX)</i>	
<i>GenRx Indapamide (GX)</i>		.Nervous system	382
.Cardiovascular system	112	.Dental	507
<i>GenRx Isosorbide Mononitrate (GX)</i>		GENTAMICIN SULFATE	
.Cardiovascular system	110	.Antifungals for systemic use	187
<i>GenRx Isotretinoin (GX)</i>		.Sensory organs	440
.Dermatologicals	154	<i>Genteal (NV)</i>	
<i>GenRx Lactulose (GX)</i>		.Sensory organs	447
.Alimentary tract and metabolism	80	.Optometrical	519
.Palliative Care	472	<i>Genteal gel (NV)</i>	
<i>GenRx Lamotrigine (GX)</i>		.Sensory organs	448
.Nervous system	391, 392	.Optometrical	520
<i>GenRx Lisinopril (GX)</i>		GESTRINONE	
.Cardiovascular system	123, 124	.Genito urinary system and sex hormones	164
<i>GenRx Meloxicam (GX)</i>		GLATIRAMER ACETATE	
.Musculo-skeletal system	359	.Antineoplastic and immunomodulating agents	236
<i>GenRx Metformin (GX)</i>		<i>Gladel (OA)</i>	
.Alimentary tract and metabolism	88, 89	.Antineoplastic and immunomodulating agents	199
<i>GenRx Metoprolol (GX)</i>		GLIBENCLAMIDE	
.Cardiovascular system	116	.Alimentary tract and metabolism	89
<i>GenRx Mirtazapine (GX)</i>		GLICLAZIDE	
.Nervous system	418	.Alimentary tract and metabolism	89
<i>GenRx Moclobemide (GX)</i>		<i>Glimel (AF)</i>	
.Nervous system	416	.Alimentary tract and metabolism	89
<i>GenRx Nifedipine (GX)</i>		GLIMEPIRIDE	
.Cardiovascular system	120	.Alimentary tract and metabolism	90
<i>GenRx Norfloxacin (GX)</i>		<i>Glimepiride Sandoz (SZ)</i>	
.Antifungals for systemic use	188	.Alimentary tract and metabolism	90
<i>GenRx Omeprazole (GX)</i>		GLIPIZIDE	
.Alimentary tract and metabolism	72, 73	.Alimentary tract and metabolism	90
<i>GenRx Paroxetine (GX)</i>		<i>Glivec (NV)</i>	
.Nervous system	415	.Antineoplastic and immunomodulating agents	214, 216, 217, 218, 219, 220, 221, 222
<i>GenRx Perindopril (GX)</i>		GLOVES PLASTIC (DISPOSABLE)	
.Cardiovascular system	124, 125	.Repatriation Schedule	874
<i>GenRx Perindopril/ Indapamide 4/1.25 (GX)</i>		<i>GlucaGen Hypokit (NO)</i>	
.Cardiovascular system	129	.Emergency Drug Supplies	64
<i>GenRx Piroxicam (GX)</i>		.Systemic hormonal preparations, excl. sex hormones and insulins	172
.Musculo-skeletal system	360	.Dental	491
.Dental	502	GLUCAGON HYDROCHLORIDE	
<i>GenRx Pravastatin (GX)</i>			
.Cardiovascular system	138, 139		
<i>GenRx Prazosin (GX)</i>			

.Systemic hormonal preparations, excl. sex hormones and insulins	172	.Cardiovascular system	128
.Dental	491	GOSERELIN ACETATE	
<i>Glucobay 100 (BN)</i>		.Antineoplastic and immunomodulating agents	230
.Alimentary tract and metabolism.....	92	GOSERELIN ACETATE and BICALUTAMIDE	
<i>Glucobay 50 (BN)</i>		.Antineoplastic and immunomodulating agents	230
.Alimentary tract and metabolism.....	92	GRANISETRON HYDROCHLORIDE	
<i>Glucocard 01 Sensor (OZ)</i>		.Alimentary tract and metabolism.....	76
.Various.....	456	<i>Granisetron Kabi (PK)</i>	
<i>Glucoflex-R (NA)</i>		.Alimentary tract and metabolism.....	76
.Various.....	455, 456	<i>Granocyte 13 (HH)</i>	
<i>Glucohexal (HX)</i>		.Section 100	570, 708
.Alimentary tract and metabolism.....	88, 89	<i>Granocyte 34 (HH)</i>	
<i>GlucoOz (OZ)</i>		.Section 100	570, 708
.Various.....	455, 456	GRISEOFULVIN	
<i>Glucophage (MQ)</i>		.Dermatologicals	150
.Alimentary tract and metabolism.....	88, 89	<i>Grisovin (SI)</i>	
GLUCOSE		.Dermatologicals	150
.Blood and blood forming organs	105	<i>Grisovin 500 (SI)</i>	
.Dental	488	.Dermatologicals	150
<i>Glucose 5% Freeflex (PK)</i>		H	
.Blood and blood forming organs	105	<i>Haemacel (AE)</i>	
GLUCOSE and KETONE INDICATOR—URINE		.Blood and blood forming organs	105
.Various.....	454	<i>Haldol decanoate (JC)</i>	
GLUCOSE INDICATOR—BLOOD		.Nervous system	401
.Various.....	455, 456, 457	HALOPERIDOL	
GLUCOSE INDICATOR—URINE		.Nervous system	400
.Various.....	455	HALOPERIDOL DECANOATE	
<i>Glucostix (BN)</i>		.Nervous system	401
.Various.....	455, 456	<i>Hamilton Pine Tar Solution (HA)</i>	
<i>Glucovance 250mg/1.25mg (AL)</i>		.Repatriation Schedule	844
.Alimentary tract and metabolism.....	91	<i>Hamilton Skin Therapy Oil (HA)</i>	
<i>Glucovance 500mg/2.5mg (AL)</i>		.Repatriation Schedule	843
.Alimentary tract and metabolism.....	90	<i>Hamilton Skin Therapy Wash (HA)</i>	
<i>Glucovance 500mg/5mg (AL)</i>		.Repatriation Schedule	846
.Alimentary tract and metabolism.....	90	<i>Hamilton Solastick 30+ (HA)</i>	
<i>Glyade (AF)</i>		.Repatriation Schedule	843
.Alimentary tract and metabolism.....	89	<i>Hamilton Sunscreen Family Sunscreen Cream SPF 15 (HA)</i>	
<i>Glyade MR (AF)</i>		.Repatriation Schedule	843
.Alimentary tract and metabolism.....	90	<i>Hamilton Sunscreen Family Sunscreen Milk SPF 15 (HA)</i>	
GLYCEROL		.Repatriation Schedule	844
.Alimentary tract and metabolism.....	81	<i>Handy 4207 (BV)</i>	
.Palliative Care	474	.Repatriation Schedule	874
.Repatriation Schedule	838	<i>Handy 4208 (BV)</i>	
GLYCERYL TRINITRATE		.Repatriation Schedule	874
.Cardiovascular system.....	109	<i>Handy 4209 (BV)</i>	
.Dental	489	.Repatriation Schedule	874
<i>Glycosade (VF)</i>		<i>Handy 5608 (BV)</i>	
.Various.....	461	.Repatriation Schedule	863
<i>GN-Carvedilol (GM)</i>		<i>Handy 5672 (BV)</i>	
.Cardiovascular system.....	117, 118	.Repatriation Schedule	869
<i>Gold Cross (BI)</i>		<i>Handy 5674 (BV)</i>	
.Repatriation Schedule	852, 857	.Repatriation Schedule	869
GOLIMUMAB		<i>Handygauze Cohesive 8631 (BV)</i>	
.Antineoplastic and immunomodulating agents 325, 329, 333, 336, 339, 342		.Repatriation Schedule	864
<i>Gonal-f Pen (SG)</i>		<i>Handygauze Cohesive 8633 (BV)</i>	
.Genito urinary system and sex hormones.....	162	.Repatriation Schedule	864
.Section 100	796	<i>Handygauze Cohesive 8635 (BV)</i>	
<i>Gopten (AB)</i>		.Repatriation Schedule	864
		<i>HCU Anamix infant (SB)</i>	

.Various.....	463	.Antineoplastic and immunomodulating agents	242, 245, 248, 251, 254, 257, 263, 269, 274, 279, 282, 285
<i>HCU Anamix junior LQ (SB)</i>		.Section 100	586, 724
.Various.....	462	<i>Humulin 30/70 (LY)</i>	
<i>HCU Cooler (VF)</i>		.Alimentary tract and metabolism.....	88
.Various.....	463	<i>Humulin NPH (LY)</i>	
<i>HCU express (VF)</i>		.Alimentary tract and metabolism.....	87
.Various.....	463	<i>Humulin R (LY)</i>	
<i>HCU gel (VF)</i>		.Alimentary tract and metabolism.....	87
.Various.....	463	<i>Hycamtin (GK)</i>	
HEPARIN SODIUM		.Antineoplastic and immunomodulating agents	230
.Blood and blood forming organs	98	<i>Hycor (SI)</i>	
<i>Hepsera (GI)</i>		.Sensory organs.....	441
.Section 100	564, 700	.Optometrical.....	516
<i>Herceptin (RO)</i>		<i>Hydopa (AF)</i>	
.Section 100	799, 800	.Cardiovascular system	110
HEXAMINE HIPPURATE		<i>HYDRALAZINE HYDROCHLORIDE</i>	
.Antifectives for systemic use	190	.Cardiovascular system	111
HIGH FAT FORMULA with VITAMINS, MINERALS and TRACE ELEMENTS and low in PROTEIN and CARBOHYDRATE		<i>Hydrea (BQ)</i>	
.Various.....	465	.Antineoplastic and immunomodulating agents	229
<i>Hiprex (IA)</i>		<i>Hydrene 25/50 (AF)</i>	
.Antifectives for systemic use	190	.Cardiovascular system	113
<i>Holoxan (BX)</i>		HYDROCHLOROTHIAZIDE	
.Antineoplastic and immunomodulating agents	199	.Cardiovascular system	111
HOMATROPINE HYDROBROMIDE		HYDROCHLOROTHIAZIDE with AMILORIDE HYDROCHLORIDE	
.Sensory organs	443	.Cardiovascular system	113
<i>Hospira Pty Limited (HH)</i>		HYDROCHLOROTHIAZIDE with TRIAMTERENE	
.Emergency Drug Supplies	64, 65	.Cardiovascular system	113
.Antineoplastic and immunomodulating agents ..	66, 200, 202, 203, 204, 206, 207, 208, 229, 357	<i>Hydrocoll 900938/1 (HR)</i>	
.Blood and blood forming organs	98	.Repatriation Schedule	871
.Systemic hormonal preparations, excl. sex hormones and insulins	169	<i>Hydrocoll 900939/1 (HR)</i>	
.Antifectives for systemic use ...	179, 181, 182, 183, 187, 188, 189	.Repatriation Schedule	871
.Nervous system	377, 379, 408	<i>Hydrocoll Thin 900942/1 (HR)</i>	
.Respiratory system	439	.Repatriation Schedule	871
.Dental	486, 494, 495, 497, 499, 505, 510	HYDROCORTISONE	
.Section 100	558, 559, 654, 695, 791	.Systemic hormonal preparations, excl. sex hormones and insulins.....	169
HPMC PAA (NM)		HYDROCORTISONE ACETATE	
.Sensory organs	448	.Alimentary tract and metabolism.....	82
.Optometrical	520	.Dermatologicals	152
<i>Humalog (LY)</i>		.Sensory organs.....	441
.Alimentary tract and metabolism.....	87	.Dental	490
<i>Humalog KwikPen (KP)</i>		.Optometrical.....	516
.Alimentary tract and metabolism.....	87	HYDROCORTISONE SODIUM SUCCINATE	
<i>Humalog Mix25 (LY)</i>		.Systemic hormonal preparations, excl. sex hormones and insulins.....	169
.Alimentary tract and metabolism.....	87	.Dental	491
<i>Humalog Mix25 KwikPen (KP)</i>		HYDROCORTISONE with CINCHOCAINE HYDROCHLORIDE	
.Alimentary tract and metabolism.....	88	.Repatriation Schedule	841
<i>Humalog Mix50 (LY)</i>		HYDROMORPHONE HYDROCHLORIDE	
.Alimentary tract and metabolism.....	88	.Nervous system	375, 376
<i>Humalog Mix50 KwikPen (KP)</i>		.Dental	504
.Alimentary tract and metabolism.....	88	<i>Hydroxo-B12 (AS)</i>	
HUMAN CHORIONIC GONADOTROPHIN		.Blood and blood forming organs	104
.Genito urinary system and sex hormones	163	HYDROXOCOBALAMIN	
.Section 100	797	.Blood and blood forming organs	104
<i>Humatrope (LY)</i>		HYDROXYCHLOROQUINE SULFATE	
.Section 100	794, 795	.Musculo-skeletal system	362
<i>Humira (AB)</i>		HYDROXYETHYL STARCH 130/0.4	
		.Blood and blood forming organs	105

HYDROXYUREA	
.Antineoplastic and immunomodulating agents	229
Hyforil (RA)	
.Cardiovascular system	129
Hygroton 25 (LM)	
.Cardiovascular system	112
HYOSCINE BUTYLBROMIDE	
.Palliative Care	470
.Repatriation Schedule	837
Hypnodorm (AF)	
.Repatriation Schedule	855
HYPROMELLOSE	
.Sensory organs	447
.Palliative Care	469, 470
.Optometrical	519
HYPROMELLOSE with CARBOMER 980	
.Sensory organs	448
.Optometrical	519
HYPROMELLOSE with DEXTRAN	
.Sensory organs	448
.Optometrical	520
Hypurin Isophane (AS)	
.Alimentary tract and metabolism	87
Hypurin Neutral (AS)	
.Alimentary tract and metabolism	87
Hysone 20 (AF)	
.Systemic hormonal preparations, excl. sex hormones and insulins	169
Hysone 4 (AF)	
.Systemic hormonal preparations, excl. sex hormones and insulins	169
Hytrin (AB)	
.Repatriation Schedule	848
I	
Iallex (LN)	
.Antiinfectives for systemic use	181, 182
.Dental	496
IBANDRONIC ACID	
.Musculo-skeletal system	367
.Section 100	649, 786
Ibilex 125 (AF)	
.Antiinfectives for systemic use	181
.Dental	496
Ibilex 250 (AF)	
.Antiinfectives for systemic use	181, 182
.Dental	496
Ibilex 500 (AF)	
.Antiinfectives for systemic use	182
.Dental	496
Ibimicyn (TS)	
.Antiinfectives for systemic use	178
.Dental	493
IBUPROFEN	
.Musculo-skeletal system	360
.Palliative Care	477
.Dental	502
IDARUBICIN HYDROCHLORIDE	
.Antineoplastic and immunomodulating agents	207
IFOSFAMIDE	
.Antineoplastic and immunomodulating agents	199
Ikorel (SW)	
.Cardiovascular system	110
ILOPROST TROMETAMOL	
.Section 100	543, 679
IMATINIB	
.Antineoplastic and immunomodulating agents	214, 216, 217, 218, 219, 220, 221
Imdur 120 mg (AP)	
.Cardiovascular system	110
Imdur Durule (AP)	
.Cardiovascular system	110
Imigran (GK)	
.Nervous system	386
Imigran FDT (GK)	
.Nervous system	386
IMIPRAMINE HYDROCHLORIDE	
.Nervous system	412
IMIQUIMOD	
.Dermatologicals	155
.Repatriation Schedule	846
ImmuCyst (SW)	
.Antineoplastic and immunomodulating agents	236
Imodium (JT)	
.Alimentary tract and metabolism	82
Imovane (SW)	
.Repatriation Schedule	855
Implanon (SH)	
.Genito urinary system and sex hormones	158
Improvil 28 Day (KR)	
.Genito urinary system and sex hormones	158
Imrest (AF)	
.Repatriation Schedule	855
Imtrate 60 mg (GM)	
.Cardiovascular system	110
Imukin (BY)	
.Section 100	572, 710
Imuran (AS)	
.Antineoplastic and immunomodulating agents	357
In a Wink Moisturising (NM)	
.Sensory organs	447
.Optometrical	519
INDAPAMIDE HEMIHYDRATE	
.Cardiovascular system	112
Indapamide Sandoz (SZ)	
.Cardiovascular system	112
Indapamide-GA (GM)	
.Cardiovascular system	112
Inderal (AP)	
.Cardiovascular system	114
INDINAVIR	
.Section 100	562, 699
Indocid (AS)	
.Musculo-skeletal system	358
.Palliative Care	476, 477
.Dental	501
INDOMETHACIN	
.Musculo-skeletal system	358
.Palliative Care	476, 477
.Dental	501

<i>Indopril 2 (SI)</i>		
.Cardiovascular system	124	
<i>Indopril 4 (SI)</i>		
.Cardiovascular system	125	
<i>Indopril 8 (SI)</i>		
.Cardiovascular system	125	
<i>INFLIXIMAB</i>		
.Section 100 .593, 597, 602, 607, 620, 624, 627, 730, 734, 739, 745, 757, 761, 764		
.Repatriation Schedule	850	
<i>INSECT ALLERGEN EXTRACT—HONEY BEE VENOM</i>		
.Various.....	452	
<i>INSECT ALLERGEN EXTRACT—PAPER WASP VENOM</i>		
.Various.....	452	
<i>INSECT ALLERGEN EXTRACT—YELLOW JACKET VENOM</i>		
.Various.....	452	
<i>Insig (SI)</i>		
.Cardiovascular system	112	
<i>Inspra (PF)</i>		
.Cardiovascular system	113	
<i>INSULIN ASPART</i>		
.Alimentary tract and metabolism.....	87	
<i>INSULIN ASPART—INSULIN ASPART PROTAMINE SUSPENSION</i>		
.Alimentary tract and metabolism.....	87	
<i>INSULIN DETEMIR</i>		
.Alimentary tract and metabolism.....	88	
<i>INSULIN GLARGINE</i>		
.Alimentary tract and metabolism.....	88	
<i>INSULIN GLULISINE</i>		
.Alimentary tract and metabolism.....	87	
<i>INSULIN ISOPHANE (N.P.H.)</i>		
.Alimentary tract and metabolism.....	87	
<i>INSULIN LISPRO</i>		
.Alimentary tract and metabolism.....	87	
<i>INSULIN LISPRO—INSULIN LISPRO PROTAMINE SUSPENSION</i>		
.Alimentary tract and metabolism.....	87	
<i>INSULIN NEUTRAL</i>		
.Alimentary tract and metabolism.....	87	
<i>INSULIN NEUTRAL—INSULIN ISOPHANE (N.P.H.), (MIXED) (Biphasic Isophane)</i>		
.Alimentary tract and metabolism.....	88	
<i>Intal CFC-Free (SW)</i>		
.Respiratory system	437	
<i>Intal Forte CFC-Free (SW)</i>		
.Respiratory system	437	
<i>Intal Spincaps (GM)</i>		
.Respiratory system	437	
<i>Integrilin (SH)</i>		
.Blood and blood forming organs	100	
<i>Intence (JC)</i>		
.Section 100	566, 703	
<i>INTERFERON ALFA-2a</i>		
.Antineoplastic and immunomodulating agents	234	
.Section 100	571, 709	
<i>INTERFERON ALFA-2b</i>		
.Antineoplastic and immunomodulating agents	235	
.Section 100	571, 709	
<i>INTERFERON BETA-1a</i>		
.Antineoplastic and immunomodulating agents	235	
<i>INTERFERON BETA-1b</i>		
.Antineoplastic and immunomodulating agents	235	
<i>INTERFERON GAMMA-1b</i>		
.Section 100	572, 710	
<i>Intrasite Gel 7313 (SN)</i>		
.Repatriation Schedule	872	
<i>Intron A (SH)</i>		
.Section 100	572, 710	
<i>Intron A Redipen (SH)</i>		
.Antineoplastic and immunomodulating agents	235	
.Section 100	572, 709, 710	
<i>Invega (JC)</i>		
.Nervous system	404	
<i>Invirase (RO)</i>		
.Section 100	563, 699	
<i>Inza 250 (AF)</i>		
.Musculo-skeletal system	361	
.Palliative Care	477, 478	
.Dental	502	
<i>Inza 500 (AF)</i>		
.Musculo-skeletal system	361	
.Palliative Care	477, 478	
.Dental	502	
<i>Iodosorb 66051330 (SN)</i>		
.Repatriation Schedule	874	
<i>Iodosorb 66051340 (SN)</i>		
.Repatriation Schedule	874	
<i>Iodosorb 66051360 (SN)</i>		
.Repatriation Schedule	874	
<i>Iodosorb Ointment 66051230 (SN)</i>		
.Repatriation Schedule	874	
<i>Iodosorb Ointment 66051240 (SN)</i>		
.Repatriation Schedule	874	
<i>Iodosorb Powder 66051070 (SN)</i>		
.Repatriation Schedule	874	
<i>Ionil-T (GA)</i>		
.Repatriation Schedule	845	
<i>Iopidine 0.5% (AQ)</i>		
.Sensory organs.....	441	
<i>Ipratrin (AF)</i>		
.Respiratory system	436	
<i>Ipratrin Adult (AF)</i>		
.Respiratory system	436	
<i>IPRATROPIUM BROMIDE</i>		
.Respiratory system	436	
.Repatriation Schedule	857	
<i>Ipravent (PF)</i>		
.Respiratory system	436	
<i>IRBESARTAN</i>		
.Cardiovascular system	131	
<i>IRBESARTAN with HYDROCHLOROTHIAZIDE</i>		
.Cardiovascular system	131	
<i>Ircal (PE)</i>		
.Sensory organs.....	448, 449	
.Optometrical.....	520	
<i>Iressa (AP)</i>		
.Antineoplastic and immunomodulating agents	214	
<i>Irinotecan Actavis (GQ)</i>		
.Antineoplastic and immunomodulating agents	229	
<i>Irinotecan Alphapharm (AF)</i>		

.Antineoplastic and immunomodulating agents	229
<i>Irinotecan Ebewe (IT)</i>	
.Antineoplastic and immunomodulating agents	229
<i>IRINOTECAN HYDROCHLORIDE TRIHYDRATE</i>	
.Antineoplastic and immunomodulating agents	229
<i>Irinotecan Sandoz (SZ)</i>	
.Antineoplastic and immunomodulating agents	229
<i>IRON POLYMALTOSE COMPLEX</i>	
.Blood and blood forming organs	104
<i>IRON SUCROSE</i>	
.Blood and blood forming organs	104
<i>Iscover (BQ)</i>	
.Blood and blood forming organs	99
.Repatriation Schedule	840
<i>Isentress (MK)</i>	
.Section 100	568, 705
<i>Isoleucine 1000 Amino Acid Supplement (VF)</i>	
.Various.....	465
<i>Isoleucine Amino Acid Supplement (VF)</i>	
.Various.....	465
<i>ISOLEUCINE with CARBOHYDRATE</i>	
.Various.....	465
<i>Isomonit (SZ)</i>	
.Cardiovascular system	110
<i>ISONIAZID</i>	
.Antifungives for systemic use	193
<i>Isoptin (AB)</i>	
.Emergency Drug Supplies	65
.Cardiovascular system	120
<i>Isoptin 180 SR (AB)</i>	
.Cardiovascular system	120
<i>Isoptin SR (AB)</i>	
.Cardiovascular system	120
<i>Isopto Carpine (AQ)</i>	
.Sensory organs	442
.Optometrical	517
<i>Isopto Homatropine (AQ)</i>	
.Sensory organs	443
<i>Isordil Sublingual (SI)</i>	
.Cardiovascular system	109
<i>ISOSORBIDE DINITRATE</i>	
.Cardiovascular system	109
<i>ISOSORBIDE MONONITRATE</i>	
.Cardiovascular system	110
<i>ISOTRETINOIN</i>	
.Dermatologicals.....	154
<i>ISPAGHULA HUSK</i>	
.Repatriation Schedule	838
<i>ITRACONAZOLE</i>	
.Antifungives for systemic use	192
<i>IVERMECTIN</i>	
.Antiparasitic products, insecticides and repellents....	432
J	
<i>Janumet (MK)</i>	
.Alimentary tract and metabolism.....	91
<i>Januvia (MK)</i>	
.Alimentary tract and metabolism.....	93
<i>Jelonet 7404 (SN)</i>	
.Repatriation Schedule	869

<i>Jezil (GN)</i>	
.Cardiovascular system	145
<i>JJ 02013 (JJ)</i>	
.Repatriation Schedule	866
<i>JJ 12010 (JJ)</i>	
.Repatriation Schedule	874
<i>Jurnista (JC)</i>	
.Nervous system	376
.Dental	505
K	
<i>Kaletra (AB)</i>	
.Section 100	567, 704
<i>Kalixocin (AF)</i>	
.Antifungives for systemic use	185
<i>Kalma 0.25 (AF)</i>	
.Nervous system	408
<i>Kalma 0.5 (AF)</i>	
.Nervous system	408
<i>Kalma 1 (AF)</i>	
.Nervous system	408
<i>Kalma 2 (AF)</i>	
.Nervous system	408
<i>Kaltostat 168117 (CC)</i>	
.Repatriation Schedule	866
<i>Kaltostat 168210 (CC)</i>	
.Repatriation Schedule	867
<i>Kaltostat 168212 (CC)</i>	
.Repatriation Schedule	866
<i>Kaluril (AF)</i>	
.Cardiovascular system	113
<i>Kapanol (GK)</i>	
.Nervous system	378
.Dental	506
<i>Karicare De-Lact (NU)</i>	
.Various.....	461
<i>Karlor CD (LN)</i>	
.Antifungives for systemic use	182
.Dental	497
<i>Karvea (SW)</i>	
.Cardiovascular system	131
<i>Karvezide 150/12.5 (SW)</i>	
.Cardiovascular system	132
<i>Karvezide 300/12.5 (SW)</i>	
.Cardiovascular system	132
<i>Karvezide 300/25 (SW)</i>	
.Cardiovascular system	132
<i>Keflex (AS)</i>	
.Antifungives for systemic use	181, 182
.Dental	496
<i>Keflin Neutral (AS)</i>	
.Antifungives for systemic use	181
.Dental	495
<i>Keflor (AF)</i>	
.Antifungives for systemic use	183
.Dental	497
<i>Keflor CD (AF)</i>	
.Antifungives for systemic use	182
.Dental	497
<i>Kefzol (AS)</i>	

.Antifungives for systemic use	182
<i>Kenacomb (SI)</i>	
.Repatriation Schedule	845
<i>Kenacomb Otic (SI)</i>	
.Sensory organs	451
<i>Kenacort-A10 (SI)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	170
.Dental	491
<i>Kepecet (GM)</i>	
.Nervous system	393
<i>Keppra (UC)</i>	
.Nervous system	393, 394
<i>Keto-Diabur- Test 5000 (RD)</i>	
.Various.....	454, 455
<i>Keto-Diastix (BN)</i>	
.Various.....	454, 455
<i>KetoCal (SB)</i>	
.Various.....	465
KETOCONAZOLE	
.Dermatologicals.....	149
.Antifungives for systemic use	190
.Repatriation Schedule	842
KETOPROFEN	
.Musculo-skeletal system	360
.Dental	502
<i>Kevtam (AF)</i>	
.Nervous system	393
<i>Kindergen (SB)</i>	
.Various.....	467
<i>Kinson (AF)</i>	
.Nervous system	396
<i>Kivexa (VI)</i>	
.Section 100.....	567, 704
<i>Klacid (AB)</i>	
.Antifungives for systemic use	185
.Section 100.....	561, 697
<i>Kosteo (SI)</i>	
.Alimentary tract and metabolism.....	95
.Musculo-skeletal system	372
<i>Kredex (MD)</i>	
.Cardiovascular system.....	117, 118
<i>Krypton 10 (AF)</i>	
.Genito urinary system and sex hormones.....	156
.Nervous system	397
<i>Krypton 2.5 (AF)</i>	
.Genito urinary system and sex hormones.....	156
.Nervous system	397
<i>Krypton 5 (AF)</i>	
.Genito urinary system and sex hormones.....	156
.Nervous system	397
<i>Kytril (HH)</i>	
.Alimentary tract and metabolism.....	76
L	
LABETALOL HYDROCHLORIDE	
.Cardiovascular system.....	118
<i>Lac-Dol (GM)</i>	
.Alimentary tract and metabolism.....	80
.Palliative Care	472

LACOSAMIDE	
.Nervous system	391
<i>Lacri-Lube (AG)</i>	
.Sensory organs.....	448, 449
.Optometrical.....	520
<i>Lactocur (SZ)</i>	
.Alimentary tract and metabolism.....	80
.Palliative Care	472
LACTULOSE	
.Alimentary tract and metabolism.....	80
.Palliative Care	472
<i>Lamictal (GK)</i>	
.Nervous system	392
<i>Lamidus (RA)</i>	
.Nervous system	391, 392
<i>Lamisil (NC)</i>	
.Dermatologicals.....	150
.Repatriation Schedule	842
<i>Lamisil (NV)</i>	
.Dermatologicals.....	150
.Repatriation Schedule	843
<i>Lamisil DermGel (NC)</i>	
.Repatriation Schedule	842
LAMIVUDINE	
.Section 100.....	565, 701
LAMIVUDINE with ZIDOVUDINE	
.Section 100.....	567, 704
<i>Lamogine (AF)</i>	
.Nervous system	391, 392
LAMOTRIGINE	
.Nervous system	391
<i>Lamotrigine generichealth (GQ)</i>	
.Nervous system	392
<i>Lamotrigine Sandoz (SZ)</i>	
.Nervous system	392
<i>Lamotrigine-DP (GM)</i>	
.Nervous system	392
<i>Lamotrigine-GA (GN)</i>	
.Nervous system	392
<i>Lamotrust 100 (MI)</i>	
.Nervous system	392
<i>Lamotrust 200 (MI)</i>	
.Nervous system	392
<i>Lamotrust 25 (MI)</i>	
.Nervous system	392
<i>Lamotrust 50 (MI)</i>	
.Nervous system	392
<i>Lanoxin (SI)</i>	
.Cardiovascular system	107
<i>Lanoxin-PG (SI)</i>	
.Cardiovascular system	107
LANREOTIDE ACETATE	
.Section 100.....	558, 694
LANSOPRAZOLE	
.Alimentary tract and metabolism.....	71
LANTHANUM	
.Various.....	452
.Section 100.....	654, 791
<i>Lantus (SW)</i>	
.Alimentary tract and metabolism.....	88

<i>Lantus SoloStar (AV)</i>		
.Alimentary tract and metabolism.....	88	
<i>Lanvis (AS)</i>		
.Antineoplastic and immunomodulating agents	202	
<i>Lanzopran (RA)</i>		
.Alimentary tract and metabolism.....	71, 72	
<i>LAPATINIB</i>		
.Antineoplastic and immunomodulating agents	222	
<i>Largactil (SW)</i>		
.Emergency Drug Supplies	64	
.Nervous system	400	
<i>Lasix (SW)</i>		
.Emergency Drug Supplies	64	
.Cardiovascular system	112	
<i>Lasix-M (SW)</i>		
.Cardiovascular system	112	
<i>LATANOPROST</i>		
.Sensory organs	443	
.Optomterical	518	
<i>LATANOPROST with TIMOLOL MALEATE</i>		
.Sensory organs	443	
.Optomterical	518	
<i>Lax-Tab (AE)</i>		
.Alimentary tract and metabolism.....	80	
.Palliative Care	471	
<i>LEFLUNOMIDE</i>		
.Antineoplastic and immunomodulating agents .	236, 237	
.Musculo-skeletal system	363	
<i>LENALIDOMIDE</i>		
.Section 100	640, 777	
<i>Lengout (LN)</i>		
.Musculo-skeletal system	364	
<i>LENOGRASTIM</i>		
.Section 100	570, 707	
<i>Lercan (SI)</i>		
.Cardiovascular system	119	
<i>LERCANIDIPINE HYDROCHLORIDE</i>		
.Cardiovascular system	119	
<i>LERCANIDIPINE HYDROCHLORIDE with ENALAPRIL MALEATE</i>		
.Cardiovascular system	129	
<i>Lercanidipine Sandoz (SZ)</i>		
.Cardiovascular system	119	
<i>Lescol (NV)</i>		
.Cardiovascular system	137	
<i>Lescol XL (NV)</i>		
.Cardiovascular system	137	
<i>LETROZOLE</i>		
.Antineoplastic and immunomodulating agents	233	
<i>Leucovorin Calcium (Hospira Pty Limited) (HH)</i>		
.Various.....	453	
<i>Leucovorin Calcium (Pfizer Australia Pty Ltd) (PF)</i>		
.Various.....	453	
<i>Leukeran (AS)</i>		
.Antineoplastic and immunomodulating agents	199	
<i>Leukoflex 1124 (BV)</i>		
.Repatriation Schedule	875	
<i>Leukoplast 1071 (BV)</i>		
.Repatriation Schedule	875	
<i>Leukoplast 1072 (BV)</i>		
.Repatriation Schedule	875	
<i>Leukoplast 1073 (BV)</i>		
.Repatriation Schedule	875	
<i>Leukopor 2471 (BV)</i>		
.Repatriation Schedule	875	
<i>Leukopor 2472 (BV)</i>		
.Repatriation Schedule	875	
<i>Leukopor 2474 (BV)</i>		
.Repatriation Schedule	875	
<i>Leukosilk 1021 (BV)</i>		
.Repatriation Schedule	875	
<i>Leukosilk 1022 (BV)</i>		
.Repatriation Schedule	875	
<i>Leukosilk 1024 (BV)</i>		
.Repatriation Schedule	875	
<i>LEUPRORELIN ACETATE</i>		
.Antineoplastic and immunomodulating agents	231	
<i>Leustatin (JC)</i>		
.Antineoplastic and immunomodulating agents	201	
<i>Levecetam 1000 (RZ)</i>		
.Nervous system	393	
<i>Levecetam 250 (RZ)</i>		
.Nervous system	393	
<i>Levecetam 500 (RZ)</i>		
.Nervous system	393	
<i>Levemir FlexPen (NF)</i>		
.Alimentary tract and metabolism	88	
<i>Levemir Penfill (NO)</i>		
.Alimentary tract and metabolism	88	
<i>LEVETIRACETAM</i>		
.Nervous system	392, 393	
<i>Levetiracetam generichealth (GQ)</i>		
.Nervous system	393	
<i>Levetiracetam SZ (SZ)</i>		
.Nervous system	393	
<i>Levitam 1000 (SI)</i>		
.Nervous system	393	
<i>Levitam 250 (SI)</i>		
.Nervous system	393	
<i>Levitam 500 (SI)</i>		
.Nervous system	393	
<i>Levlen ED (SY)</i>		
.Genito urinary system and sex hormones	157	
<i>Levo/Carbidopa Sandoz (SZ)</i>		
.Nervous system	396	
<i>LEVOCABASTINE HYDROCHLORIDE</i>		
.Repatriation Schedule	857, 859	
<i>LEVODOPA with BENSERAZIDE</i>		
.Nervous system	396	
<i>LEVODOPA with CARBIDOPA</i>		
.Nervous system	396	
<i>LEVODOPA with CARBIDOPA and ENTACAPONE</i>		
.Nervous system	396	
<i>LEVONORGESTREL</i>		
.Genito urinary system and sex hormones	156, 158	
<i>LEVONORGESTREL with ETHINYLOESTRADIOL</i>		
.Genito urinary system and sex hormones	157, 158	
<i>Lexam 10 (SI)</i>		
.Nervous system	413	
<i>Lexam 20 (SI)</i>		
.Nervous system	413	

<i>Lexapro (LU)</i>		
.Nervous system	413, 414	
<i>Lexotan (RO)</i>		
.Repatriation Schedule	854	
<i>Lifeline Attest (OI)</i>		
.Variou.....	456	
LIGNOCAINE HYDROCHLORIDE		
.Cardiovascular system	107	
.Dental	489	
LIGNOCAINE HYDROCHLORIDE with CARBOXYMETHYLCELLULOSE		
.Repatriation Schedule	844	
<i>Lincocin (PF)</i>		
.Anti-infectives for systemic use	186	
.Dental	499	
LINCOMYCIN		
.Anti-infectives for systemic use	186	
.Dental	499	
<i>Link Medical Products Pty Ltd (LM)</i>		
.Dermatologicals.....	155	
.Anti-infectives for systemic use	193	
<i>Lioresal 10 (NV)</i>		
.Musculo-skeletal system	363	
<i>Lioresal 25 (NV)</i>		
.Musculo-skeletal system	363	
<i>Lioresal Intrathecal (NV)</i>		
.Section 100.....	648, 785	
LIOTHYRONINE SODIUM		
.Systemic hormonal preparations, excl. sex hormones and insulins	171	
<i>Lipazil 600 mg (GM)</i>		
.Cardiovascular system	145	
<i>Lipex 10 (FR)</i>		
.Cardiovascular system	141, 143	
<i>Lipex 20 (FR)</i>		
.Cardiovascular system	141, 143	
<i>Lipex 40 (FR)</i>		
.Cardiovascular system	142, 144	
<i>Lipex 80 (FR)</i>		
.Cardiovascular system.....	142, 144	
<i>Lipidil (SM)</i>		
.Cardiovascular system.....	144	
<i>Lipigem (AF)</i>		
.Cardiovascular system.....	145	
<i>Lipitor (PF)</i>		
.Cardiovascular system.....	137	
<i>Lipostat 10 (SI)</i>		
.Cardiovascular system.....	138, 139	
<i>Lipostat 20 (SI)</i>		
.Cardiovascular system.....	138, 139	
<i>Lipostat 40 (SI)</i>		
.Cardiovascular system.....	138, 140	
<i>Lipostat 80 (SI)</i>		
.Cardiovascular system.....	139, 140	
<i>Liprace (GM)</i>		
.Cardiovascular system.....	123, 124	
<i>Liquifilm Forte (AG)</i>		
.Sensory organs	450	
.Optomelrical	521	
<i>Liquifilm Tears (AG)</i>		
.Sensory organs.....	450	
.Optomelrical.....	521	
<i>Liquigen (SB)</i>		
.Variou.....	457	
LISINOPRIL		
.Cardiovascular system	123	
<i>Lisinopril 10 (CR)</i>		
.Cardiovascular system	123	
<i>Lisinopril 20 (CR)</i>		
.Cardiovascular system	124	
<i>Lisinopril 5 (CR)</i>		
.Cardiovascular system	123	
<i>Lisinopril generichealth (GQ)</i>		
.Cardiovascular system	123, 124	
<i>Lisinopril Hexal (HX)</i>		
.Cardiovascular system	123	
<i>Lisinopril Ranbaxy (RA)</i>		
.Cardiovascular system	123, 124	
<i>Lisinopril Sandoz (SZ)</i>		
.Cardiovascular system	123, 124	
<i>Lisinopril Winthrop (WA)</i>		
.Cardiovascular system	123, 124	
<i>Lisinopril-DRLA (RZ)</i>		
.Cardiovascular system	123, 124	
<i>Lisinopril-GA (GN)</i>		
.Cardiovascular system	123, 124	
<i>Lisodur (AF)</i>		
.Cardiovascular system	123, 124	
<i>Litak (OA)</i>		
.Antineoplastic and immunomodulating agents	201	
<i>Lithicarb (AS)</i>		
.Nervous system	403, 417	
LITHIUM CARBONATE		
.Nervous system	403, 417	
<i>Livostin (JT)</i>		
.Repatriation Schedule	857, 859	
<i>Locasol (SB)</i>		
.Variou.....	462	
<i>Loceryl (GA)</i>		
.Repatriation Schedule	842	
<i>Locilan 28 Day (KR)</i>		
.Genito urinary system and sex hormones	158	
<i>Lodam 50 (ZP)</i>		
.Nervous system	382	
.Dental	507	
<i>Lodam SR 100 (ZP)</i>		
.Nervous system	383	
.Dental	507	
<i>Lodam SR 150 (ZP)</i>		
.Nervous system	383	
.Dental	508	
<i>Lodam SR 200 (ZP)</i>		
.Nervous system	383	
.Dental	508	
<i>Lofenoxal (HC)</i>		
.Alimentary tract and metabolism	82	
<i>Logicin Rapid Relief (SI)</i>		
.Repatriation Schedule	857	
<i>Logicin Sinus (SI)</i>		
.Repatriation Schedule	857	

<i>Logynon ED (SY)</i>	
.Genito urinary system and sex hormones.....	158
<i>Lomotil (BI)</i>	
.Alimentary tract and metabolism.....	82
<i>Loniten (PF)</i>	
.Cardiovascular system.....	111
<i>LOPERAMIDE HYDROCHLORIDE</i>	
.Alimentary tract and metabolism.....	82
<i>Lophlex (SB)</i>	
.Various.....	464
<i>Lopid (PF)</i>	
.Cardiovascular system.....	145
<i>LOPINA VIR with RITONAVIR</i>	
.Section 100.....	567, 704
<i>Lopresor 100 (NV)</i>	
.Cardiovascular system.....	116
<i>Lopresor 50 (NV)</i>	
.Cardiovascular system.....	116
<i>Lorano (SZ)</i>	
.Repatriation Schedule.....	858
<i>LORATADINE</i>	
.Repatriation Schedule.....	858
<i>Losec Tablets (AP)</i>	
.Alimentary tract and metabolism.....	72, 73
<i>Lovan (AL)</i>	
.Nervous system.....	414
<i>Lovan 20 Tab (AL)</i>	
.Nervous system.....	414
<i>Lovir (GM)</i>	
.Anti-infectives for systemic use.....	194, 195
<i>LoxaLate (AF)</i>	
.Nervous system.....	413
<i>LPV (AS)</i>	
.Anti-infectives for systemic use.....	178, 179
.Dental.....	493
<i>Lucentis (NV)</i>	
.Sensory organs.....	444
<i>Lucrin Depot 3 Month PDS (AB)</i>	
.Antineoplastic and immunomodulating agents.....	231
<i>Lucrin Depot 4 Month PDS (AB)</i>	
.Antineoplastic and immunomodulating agents.....	231
<i>Lucrin Depot 7.5mg PDS (AB)</i>	
.Antineoplastic and immunomodulating agents.....	231
<i>Lumigan (AG)</i>	
.Sensory organs.....	443
.Optometical.....	518
<i>Lumin 10 (AF)</i>	
.Nervous system.....	417
<i>Lumin 20 (AF)</i>	
.Nervous system.....	417
<i>Luvox (SM)</i>	
.Nervous system.....	414
<i>Lycinate (FM)</i>	
.Cardiovascular system.....	109
.Dental.....	489
<i>Lyclear (JT)</i>	
.Antiparasitic products, insecticides and repellents....	432
<i>Lyof foam C 603025 (SS)</i>	
.Repatriation Schedule.....	868
<i>Lyof foam Extra 603088 (XP)</i>	
.Repatriation Schedule.....	867
<i>Lyof foam Extra 603090 (XP)</i>	
.Repatriation Schedule.....	868
<i>Lyof foam Flat 603092 (XP)</i>	
.Repatriation Schedule.....	868
<i>Lyof foam Flat 603093 (XP)</i>	
.Repatriation Schedule.....	868
<i>Lyof foam Flat 603095 (XP)</i>	
.Repatriation Schedule.....	868
<i>Lyrica (PF)</i>	
.Repatriation Schedule.....	854
M	
<i>Mabthera (RO)</i>	
.Antineoplastic and immunomodulating agents .	209, 210
.Section 100.....	647, 783
<i>Macro dantin (PF)</i>	
.Anti-infectives for systemic use.....	190
<i>MACROGOL 3350</i>	
.Alimentary tract and metabolism.....	80
.Palliative Care.....	472
<i>Madopar (RO)</i>	
.Nervous system.....	396
<i>Madopar 125 (RO)</i>	
.Nervous system.....	396
<i>Madopar 62.5 (RO)</i>	
.Nervous system.....	396
<i>Madopar HBS (RO)</i>	
.Nervous system.....	396
<i>Madopar Rapid 125 (RO)</i>	
.Nervous system.....	396
<i>Madopar Rapid 62.5 (RO)</i>	
.Nervous system.....	396
<i>Mag-Sup (PP)</i>	
.Repatriation Schedule.....	839
<i>Magicul 200 (AF)</i>	
.Alimentary tract and metabolism.....	68
<i>Magicul 400 (AF)</i>	
.Alimentary tract and metabolism.....	68
<i>Magicul 800 (AF)</i>	
.Alimentary tract and metabolism.....	68
<i>Magmin (BB)</i>	
.Repatriation Schedule.....	839
<i>MAGNESIUM ASPARTATE</i>	
.Repatriation Schedule.....	839
<i>Malarone (GK)</i>	
.Antiparasitic products, insecticides and repellents....	431
<i>MARAVIROC</i>	
.Section 100.....	568, 705
<i>Marevan (FM)</i>	
.Blood and blood forming organs.....	97
<i>Max Pharma Pty Ltd (XF)</i>	
.Anti-infectives for systemic use.....	184
<i>Maxalt (MK)</i>	
.Nervous system.....	385
<i>Maxamox (SZ)</i>	
.Anti-infectives for systemic use.....	178
.Dental.....	493
<i>Maxidex (AQ)</i>	
.Sensory organs.....	441

<i>Maxipime (BQ)</i>		
.Antifungives for systemic use	184	
<i>Maxolon (VT)</i>		
.Emergency Drug Supplies	64	
.Alimentary tract and metabolism.....	76	
.Dental	486	
<i>Mayne Pharma Doxycycline (YT)</i>		
.Antifungives for systemic use	175, 176	
.Dental	492	
<i>Mayne Pharma Erythromycin (YT)</i>		
.Antifungives for systemic use	185	
.Dental	498	
<i>MCT Oil (SB)</i>		
.Various.....	457	
<i>MCT Pro-Cal (VF)</i>		
.Various.....	466	
<i>MEBENDAZOLE</i>		
.Repatriation Schedule	856	
<i>MEBEVERINE HYDROCHLORIDE</i>		
.Repatriation Schedule	837	
<i>MediHealth ClearLax (ON)</i>		
.Alimentary tract and metabolism.....	81	
.Palliative Care	472	
<i>Medipore 2961 (MM)</i>		
.Repatriation Schedule	874	
<i>Medroxyhexal (SZ)</i>		
.Genito urinary system and sex hormones.....	161	
<i>MEDROXYPROGESTERONE ACETATE</i>		
.Genito urinary system and sex hormones.....	158, 161	
.Antineoplastic and immunomodulating agents	230	
<i>MEFENAMIC ACID</i>		
.Musculo-skeletal system	361	
<i>Mefix 310250 (MH)</i>		
.Repatriation Schedule	875	
<i>Megace (SI)</i>		
.Antineoplastic and immunomodulating agents	230	
<i>Megafol 0.5 (AF)</i>		
.Blood and blood forming organs	104	
<i>Megafol 5 (AF)</i>		
.Blood and blood forming organs	105	
<i>MEGESTROL ACETATE</i>		
.Antineoplastic and immunomodulating agents	230	
<i>Melizide (AF)</i>		
.Alimentary tract and metabolism.....	90	
<i>Mellihexal (SZ)</i>		
.Alimentary tract and metabolism.....	89	
<i>Melolin 36361357 (SN)</i>		
.Repatriation Schedule	873	
<i>Melolin 66974933 (SN)</i>		
.Repatriation Schedule	873	
<i>Meloxibell (BF)</i>		
.Musculo-skeletal system	359	
<i>MELOXICAM</i>		
.Musculo-skeletal system	359	
<i>Meloxicam Ranbaxy (RA)</i>		
.Musculo-skeletal system	359	
<i>Meloxicam Sandoz (SZ)</i>		
.Musculo-skeletal system	359	
<i>Meloxicam Winthrop (WA)</i>		
.Musculo-skeletal system	359	
<i>Meloxicam-GA (GM)</i>		
.Musculo-skeletal system	359	
<i>MELPHALAN</i>		
.Antineoplastic and immunomodulating agents	199	
<i>MEMANTINE HYDROCHLORIDE</i>		
.Nervous system	426	
<i>Memnax (SI)</i>		
.Nervous system	426	
<i>Mepilex 294100 (MH)</i>		
.Repatriation Schedule	869	
<i>Mepilex Border 295200 (MH)</i>		
.Repatriation Schedule	868	
<i>Mepilex Border 295300 (MH)</i>		
.Repatriation Schedule	868	
<i>Mepilex Lite 284000 (MH)</i>		
.Repatriation Schedule	868	
<i>Mepilex Lite 284100 (MH)</i>		
.Repatriation Schedule	868	
<i>Mepitac 298300 (MH)</i>		
.Repatriation Schedule	875	
<i>Mepitac 298400 (MH)</i>		
.Repatriation Schedule	875	
<i>Mepitel 290510 (MH)</i>		
.Repatriation Schedule	873	
<i>Mepitel 290710 (MH)</i>		
.Repatriation Schedule	873	
<i>Meprazol (SZ)</i>		
.Alimentary tract and metabolism	72, 73	
<i>MERCAPTOPURINE</i>		
.Antineoplastic and immunomodulating agents	202	
<i>MESALAZINE</i>		
.Alimentary tract and metabolism	83, 84	
<i>Mesasal (GK)</i>		
.Alimentary tract and metabolism	83	
<i>MESNA</i>		
.Various.....	453	
<i>Mestinon (VT)</i>		
.Nervous system	427	
<i>Mestinon Timespan (VT)</i>		
.Nervous system	427	
<i>Metalyse (BY)</i>		
.Blood and blood forming organs	102	
<i>Metamucil Regular (PY)</i>		
.Repatriation Schedule	838	
<i>Metamucil Smooth Texture Orange (PY)</i>		
.Repatriation Schedule	838	
<i>Metex XR (SI)</i>		
.Alimentary tract and metabolism	89	
<i>Metformin 500 (CR)</i>		
.Alimentary tract and metabolism	89	
<i>Metformin 850 (CR)</i>		
.Alimentary tract and metabolism	88	
<i>Metformin generichealth (GQ)</i>		
.Alimentary tract and metabolism	88, 89	
<i>Metformin generichealth 1000 (GQ)</i>		
.Alimentary tract and metabolism	89	
<i>METFORMIN HYDROCHLORIDE</i>		
.Alimentary tract and metabolism	88	
<i>METFORMIN HYDROCHLORIDE with GLIBENCLAMIDE</i>		
.Alimentary tract and metabolism	90	

<i>Metformin Ranbaxy (RA)</i>	
.Alimentary tract and metabolism.....	88, 89
<i>Metformin Ranbaxy 1000 (RA)</i>	
.Alimentary tract and metabolism.....	89
<i>Metformin Sandoz (SZ)</i>	
.Alimentary tract and metabolism.....	88, 89
<i>Metformin-GA (GM)</i>	
.Alimentary tract and metabolism.....	88, 89
<i>METHADONE HYDROCHLORIDE</i>	
.Nervous system	381
.Palliative Care	482
.Section 100	798
<i>Methoblastin (PF)</i>	
.Antineoplastic and immunomodulating agents .	200, 357
<i>Methopt (SI)</i>	
.Sensory organs	447
.Optometrical	519
<i>METHOTREXATE</i>	
.Antineoplastic and immunomodulating agents .	200, 357
<i>Methotrexate Ebewe (IT)</i>	
.Antineoplastic and immunomodulating agents	200
<i>METHOXY POLYETHYLENE GLYCOL-EPOETIN BETA</i>	
.Section 100	525, 657
<i>METHYL SALICYLATE</i>	
.Repatriation Schedule	852
<i>METHYLDOPA</i>	
.Cardiovascular system	110
<i>METHYLNALTREXONE</i>	
.Palliative Care	473
<i>METHYLPHENIDATE HYDROCHLORIDE</i>	
.Nervous system	419, 420
<i>METHYLPREDNISOLONE ACEPONATE</i>	
.Dermatologicals	153
<i>METHYLPREDNISOLONE ACETATE</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	170
.Dental	491
<i>METHYLPREDNISOLONE SODIUM SUCCINATE</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	170
<i>METHYSERGIDE</i>	
.Nervous system	385
<i>METOCLOPRAMIDE HYDROCHLORIDE</i>	
.Alimentary tract and metabolism.....	76
.Dental	486
<i>Metohexal (SZ)</i>	
.Cardiovascular system	116
<i>METOPROLOL SUCCINATE</i>	
.Cardiovascular system	115
<i>METOPROLOL TARTRATE</i>	
.Cardiovascular system	116
<i>Metrogyl 200 (AF)</i>	
.Antiinfectives for systemic use	189
.Antiparasitic products, insecticides and repellents	430
.Dental	499
<i>Metrogyl 400 (AF)</i>	
.Antiinfectives for systemic use	189
.Antiparasitic products, insecticides and repellents	430
.Dental	499
<i>Metrol 100 (SI)</i>	
.Cardiovascular system	116
<i>Metrol 50 (SI)</i>	
.Cardiovascular system	116
<i>METRONIDAZOLE</i>	
.Antiinfectives for systemic use	189
.Antiparasitic products, insecticides and repellents	430
.Dental	499
<i>METRONIDAZOLE BENZOATE</i>	
.Antiinfectives for systemic use	190
.Antiparasitic products, insecticides and repellents	430
.Dental	499
<i>Metronidazole Sandoz (SZ)</i>	
.Antiinfectives for systemic use	189
.Dental	499
<i>Metronide 200 (AV)</i>	
.Antiinfectives for systemic use	189
.Antiparasitic products, insecticides and repellents	430
.Dental	499
<i>Metronide 400 (AV)</i>	
.Antiinfectives for systemic use	189
.Antiparasitic products, insecticides and repellents	430
.Dental	499
<i>Mezavant (ZI)</i>	
.Alimentary tract and metabolism	84
<i>Miacalcic 100 (NV)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins.....	173
<i>Miacalcic 50 (NV)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins.....	173
<i>MIANSERIN HYDROCHLORIDE</i>	
.Nervous system	417
<i>Micardis (BY)</i>	
.Cardiovascular system	131
<i>Micardis Plus 40/12.5 mg (BY)</i>	
.Cardiovascular system	132
<i>Micardis Plus 80/12.5 mg (BY)</i>	
.Cardiovascular system	132
<i>Micardis Plus 80/25 mg (BY)</i>	
.Cardiovascular system	132
<i>Micolette (AE)</i>	
.Alimentary tract and metabolism.....	81
.Palliative Care	473
<i>MICONAZOLE</i>	
.Dermatologicals	149
.Repatriation Schedule	842
<i>MICONAZOLE NITRATE</i>	
.Dermatologicals	149
.Repatriation Schedule	842
<i>Microgynon 30 ED (SC)</i>	
.Genito urinary system and sex hormones	157
<i>Microgynon 50 ED (SC)</i>	
.Genito urinary system and sex hormones	157
<i>Microlax (JT)</i>	
.Alimentary tract and metabolism	81
.Palliative Care	473
.Repatriation Schedule	838
<i>Microlut 28 (SC)</i>	
.Genito urinary system and sex hormones	158
<i>Micronor (JC)</i>	

.Genito urinary system and sex hormones.....	158	.Alimentary tract and metabolism.....	88
MILK POWDER—LACTOSE FREE FORMULA		<i>Mixtard 50/50 Penfill 3 mL (NO)</i>	
.Various.....	461	.Alimentary tract and metabolism.....	88
MILK POWDER—LACTOSE MODIFIED		<i>MMA/PA Anamix infant (SB)</i>	
.Various.....	461	.Various.....	463
MILK POWDER—SYNTHETIC		<i>MMA/PA express (VF)</i>	
.Various.....	462	.Various.....	463
MILK PROTEIN and FAT FORMULA with VITAMINS and MINERALS—CARBOHYDRATE FREE		<i>MMA/PA gel (VF)</i>	
.Various.....	466	.Various.....	463
<i>Minax 100 (AF)</i>		<i>Mobic (BY)</i>	
.Cardiovascular system.....	116	.Musculo-skeletal system.....	359
<i>Minax 50 (AF)</i>		<i>Mobilis 10 (AF)</i>	
.Cardiovascular system.....	116	.Musculo-skeletal system.....	360
<i>Minidiab (PF)</i>		.Dental.....	502
.Alimentary tract and metabolism.....	90	<i>Mobilis 20 (AF)</i>	
<i>Minipress (PF)</i>		.Musculo-skeletal system.....	360
.Cardiovascular system.....	111	.Dental.....	502
<i>Minirin (FP)</i>		<i>Mobilis D-10 (AF)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins.....	167	.Musculo-skeletal system.....	360
<i>Minirin Melt (FP)</i>		.Dental.....	502
.Systemic hormonal preparations, excl. sex hormones and insulins.....	168	<i>Mobilis D-20 (AF)</i>	
<i>Minirin Nasal Spray (FP)</i>		.Musculo-skeletal system.....	360
.Systemic hormonal preparations, excl. sex hormones and insulins.....	167, 168	.Dental.....	502
<i>Minitran 10 (IA)</i>		MOCLOBEMIDE	
.Cardiovascular system.....	109	.Nervous system.....	416
<i>Minitran 15 (IA)</i>		<i>Moclobemide Sandoz (SZ)</i>	
.Cardiovascular system.....	109	.Nervous system.....	416
<i>Minitran 5 (IA)</i>		MODAFINIL	
.Cardiovascular system.....	109	.Nervous system.....	420
MINOCYCLINE		<i>Modavigil (CS)</i>	
.Antifectives for systemic use.....	177	.Nervous system.....	421
<i>Minomycin-50 (SI)</i>		<i>Modecate (BQ)</i>	
.Antifectives for systemic use.....	177	.Nervous system.....	400
MINOXIDIL		<i>Moduretic (AS)</i>	
.Cardiovascular system.....	111	.Cardiovascular system.....	113
<i>Mircera (RO)</i>		<i>Mogadon (VT)</i>	
.Section 100.....	525, 657	.Nervous system.....	388, 410
<i>Mirena (SC)</i>		.Palliative Care.....	484, 485
.Genito urinary system and sex hormones.....	156	.Dental.....	510
MIRTAZAPINE		<i>Mohexal (HX)</i>	
.Nervous system.....	417	.Nervous system.....	416
<i>Mirtazapine Sandoz (SZ)</i>		MOMETASONE FUROATE	
.Nervous system.....	418	.Dermatologicals.....	153
<i>Mirtazapine-DP (GM)</i>		.Repatriation Schedule.....	845
.Nervous system.....	418	<i>Momex SR 10 (SI)</i>	
<i>Mirtazon (SI)</i>		.Nervous system.....	378
.Nervous system.....	418	.Dental.....	506
MISOPROSTOL		<i>Momex SR 100 (SI)</i>	
.Alimentary tract and metabolism.....	70	.Nervous system.....	378
<i>Mitozantrone Ebewe (IT)</i>		.Dental.....	506
.Antineoplastic and immunomodulating agents.....	207	<i>Momex SR 30 (SI)</i>	
MITOZANTRONE HYDROCHLORIDE		.Nervous system.....	378
.Antineoplastic and immunomodulating agents.....	207	.Dental.....	506
<i>Mixtard 30/70 InnoLet (NI)</i>		<i>Monace 10 (AF)</i>	
.Alimentary tract and metabolism.....	88	.Cardiovascular system.....	123
<i>Mixtard 30/70 Penfill 3 mL (NO)</i>		<i>Monace 20 (AF)</i>	
		.Cardiovascular system.....	123

<i>Monodur 120 mg (PM)</i>		.Nervous system	379
.Cardiovascular system	110		
<i>Monodur 60 mg (PM)</i>		.Nervous system	378
.Cardiovascular system	110	.Dental	506
<i>Monofeme 28 (WX)</i>		<i>MS Contin Suspension 30 mg (MF)</i>	
.Genito urinary system and sex hormones	157	.Nervous system	378
<i>Monogen (SB)</i>		.Dental	506
.Various	460	<i>MS Contin Suspension 60 mg (MF)</i>	
<i>Monoplus 10/12.5 (BQ)</i>		.Nervous system	378
.Cardiovascular system	129	.Dental	506
<i>Monoplus 20/12.5 (BQ)</i>		<i>MS Mono (MF)</i>	
.Cardiovascular system	129	.Nervous system	378, 379
<i>Monopril (BQ)</i>		.Dental	505
.Cardiovascular system	123	<i>MSUD AID III (SB)</i>	
MONTELUKAST SODIUM		.Various	464
.Respiratory system	438	<i>MSUD Anamix infant (SB)</i>	
MORPHINE HYDROCHLORIDE		.Various	464
.Nervous system	376	<i>MSUD Anamix Junior (SB)</i>	
.Dental	505	.Various	464
MORPHINE SULFATE		<i>MSUD Anamix Junior LQ (SB)</i>	
.Nervous system	377, 378, 379	.Various	464
.Palliative Care	480	<i>MSUD Cooler (VF)</i>	
.Dental	505	.Various	464
.Repatriation Schedule	853	<i>MSUD Express (VF)</i>	
MORPHINE TARTRATE		.Various	464
.Nervous system	379	<i>MSUD Maxamaid (SB)</i>	
<i>Motilium (JC)</i>		.Various	464
.Alimentary tract and metabolism	76	<i>MSUD Maxamum (SB)</i>	
<i>Movalis 15 (SI)</i>		.Various	464
.Musculo-skeletal system	359, 360	<i>MSUD-gel (VF)</i>	
<i>Movalis 7.5 (SI)</i>		.Various	464
.Musculo-skeletal system	359	<i>Muphoran (SE)</i>	
<i>Movicol (NE)</i>		.Antineoplastic and immunomodulating agents	199
.Alimentary tract and metabolism	81	MUPIROCIN	
.Palliative Care	472	.Respiratory system	433
<i>Movox 100 (AF)</i>		.Repatriation Schedule	844
.Nervous system	414	<i>Murelax (FM)</i>	
<i>Movox 50 (AL)</i>		.Nervous system	409
.Nervous system	414	.Palliative Care	484
<i>Moxicam 15 (AF)</i>		.Dental	510
.Musculo-skeletal system	359	<i>MWD Pen Sensor Strips (WF)</i>	
<i>Moxicam 7.5 (AF)</i>		.Various	455, 456
.Musculo-skeletal system	359	<i>Mycobutin (PF)</i>	
<i>Moxiclav Duo 500/125 (SI)</i>		.Section 100	561, 697
.Antifungives for systemic use	180	MYCOPHENOLATE MOFETIL	
.Dental	495	.Antineoplastic and immunomodulating agents	237
<i>Moxiclav Duo Forte 875/125 (SI)</i>		.Section 100	581, 718
.Antifungives for systemic use	180	MYCOPHENOLATE SODIUM	
.Dental	495	.Antineoplastic and immunomodulating agents	237
MOXONIDINE		.Section 100	581, 719
.Cardiovascular system	111	<i>Mycostatin (FM)</i>	
<i>MS Contin (MF)</i>		.Alimentary tract and metabolism	68
.Nervous system	378, 379	.Dermatologicals	149
.Palliative Care	480	.Dental	486
.Dental	505, 506	.Repatriation Schedule	842
.Repatriation Schedule	853	<i>Myfortic (NV)</i>	
<i>MS Contin Suspension 100 mg (MF)</i>		.Antineoplastic and immunomodulating agents	237
.Nervous system	378	.Section 100	581, 719
.Dental	506	<i>MyGlucoHealth (EH)</i>	
<i>MS Contin Suspension 200 mg (MF)</i>		.Various	456

<i>Mylanta Double Strength (JT)</i>	
.Repatriation Schedule	837
<i>Mylanta P (JT)</i>	
.Alimentary tract and metabolism.....	68
<i>Myleran (AS)</i>	
.Antineoplastic and immunomodulating agents	199
<i>Myocrisin (SW)</i>	
.Musculo-skeletal system	362
<i>Mysoline (LM)</i>	
.Nervous system	387

N

<i>NAB PACLITAXEL</i>	
.Antineoplastic and immunomodulating agents	205
<i>NAFARELIN ACETATE</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	168
<i>NALOXONE HYDROCHLORIDE</i>	
.Various.....	452
.Dental	513
<i>Naloxone Min-I-Jet (CS)</i>	
.Emergency Drug Supplies	65
.Various.....	452
.Dental	513
<i>Naltrexone generichealth (GQ)</i>	
.Nervous system	428
<i>NALTREXONE HYDROCHLORIDE</i>	
.Nervous system	428
<i>Naltrexone QP (XF)</i>	
.Nervous system	428
<i>NANDROLONE DECANOATE</i>	
.Alimentary tract and metabolism.....	96
<i>NAPHAZOLINE HYDROCHLORIDE</i>	
.Repatriation Schedule	859
<i>Naprosyn (RO)</i>	
.Musculo-skeletal system	361
.Palliative Care	477, 478
.Dental	502, 503
<i>Naprosyn SR1000 (RO)</i>	
.Musculo-skeletal system	361
.Palliative Care	478
.Dental	503
<i>Naprosyn SR750 (RO)</i>	
.Musculo-skeletal system	361
.Palliative Care	478
.Dental	503
<i>NAPROXEN</i>	
.Musculo-skeletal system	360, 361
.Palliative Care	477, 478
.Dental	502
<i>NAPROXEN SODIUM</i>	
.Musculo-skeletal system	361
.Palliative Care	478, 479
.Dental	503
<i>Naramig (GK)</i>	
.Nervous system	66, 67
<i>NARATRIPTAN</i>	
.Nervous system	66
<i>Nardil (LM)</i>	
.Nervous system	416

<i>NATALIZUMAB</i>	
.Section 100	581, 719
<i>Natrilix (SE)</i>	
.Cardiovascular system	112
<i>Natrilix SR (SE)</i>	
.Cardiovascular system	112
<i>Navelbine (FB)</i>	
.Antineoplastic and immunomodulating agents .	203, 204
<i>Navoban (NV)</i>	
.Alimentary tract and metabolism.....	78
<i>Nebilet (CS)</i>	
.Cardiovascular system	116
<i>NEBIVOLOL</i>	
.Cardiovascular system	116
<i>NEDOCROMIL SODIUM</i>	
.Respiratory system	437
<i>Nemdyn (HA)</i>	
.Sensory organs.....	451
<i>Neo-B12 (HH)</i>	
.Blood and blood forming organs	104
<i>Neo-Mercazole (LM)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins.....	171
<i>Neocate (SB)</i>	
.Various.....	458
<i>Neocate Advance (SB)</i>	
.Various.....	458
<i>Neocate Advance Tropical Flavour (SB)</i>	
.Various.....	458
<i>Neocate LCP (SB)</i>	
.Various.....	458, 459
<i>NEOMYCIN SULFATE</i>	
.Alimentary tract and metabolism.....	82
<i>NEOMYCIN UNDECENOATE with BACITRACIN ZINC</i>	
.Sensory organs.....	451
<i>Neoral (NV)</i>	
.Antineoplastic and immunomodulating agents	356
.Section 100	640, 776
<i>Neoral 10 (NV)</i>	
.Antineoplastic and immunomodulating agents	356
.Section 100	640, 776
<i>Neoral 100 (NV)</i>	
.Antineoplastic and immunomodulating agents	356
.Section 100	640, 777
<i>Neoral 25 (NV)</i>	
.Antineoplastic and immunomodulating agents	356
.Section 100	640, 777
<i>Neoral 50 (NV)</i>	
.Antineoplastic and immunomodulating agents	356
.Section 100	640, 777
<i>NeoRecormon (RO)</i>	
.Section 100	524, 525, 656, 657
<i>Neosulf (AF)</i>	
.Alimentary tract and metabolism.....	82
<i>Neotigason (TA)</i>	
.Dermatologicals.....	151
<i>Neulactil (SW)</i>	
.Nervous system	400
<i>Neulasta (AN)</i>	
.Section 100	571, 709

<i>Neupogen (AN)</i>		
. Section 100	570, 707	
<i>Neurontin (PF)</i>		
. Nervous system	390, 391	
. Repatriation Schedule	853, 854	
<i>NEVIRAPINE</i>		
. Section 100	567, 703	
<i>Nexavar (BN)</i>		
. Antineoplastic and immunomodulating agents	225	
<i>Nexcare Durable Cloth First Aid Tape 799 (MM)</i>		
. Repatriation Schedule	875	
<i>Nexcare Gentle Paper First Aid Tape 789 (MM)</i>		
. Repatriation Schedule	875	
<i>Nexcare Tegaderm Transparent H1624 (MM)</i>		
. Repatriation Schedule	867	
<i>Nexcare Tegaderm Transparent H1626 (MM)</i>		
. Repatriation Schedule	867	
<i>Nexium (AP)</i>		
. Alimentary tract and metabolism	70, 71	
<i>Nexium Hp7 (AP)</i>		
. Alimentary tract and metabolism	75	
<i>Nicabate CQ 14 (GC)</i>		
. Repatriation Schedule	855	
<i>Nicabate CQ 21 (GC)</i>		
. Repatriation Schedule	855	
<i>NICORANDIL</i>		
. Cardiovascular system	110	
<i>Nicorette Patch (JT)</i>		
. Nervous system	427	
. Repatriation Schedule	855	
<i>NICOTINE</i>		
. Nervous system	427	
. Repatriation Schedule	855	
<i>Nidem (SI)</i>		
. Alimentary tract and metabolism	89	
<i>NIFEDIPINE</i>		
. Cardiovascular system	119, 120	
<i>Nifehexal (SZ)</i>		
. Cardiovascular system	120	
<i>NILOTINIB</i>		
. Antineoplastic and immunomodulating agents .	222, 224	
<i>Nilstat (SI)</i>		
. Alimentary tract and metabolism	68, 82	
. Dental	486, 487	
. Repatriation Schedule	847	
<i>NILUTAMIDE</i>		
. Antineoplastic and immunomodulating agents	233	
<i>NITRAZEPAM</i>		
. Nervous system	388, 409, 410	
. Palliative Care	484, 485	
. Dental	510	
<i>Nitro-Dur 10 (SH)</i>		
. Cardiovascular system	109	
<i>Nitro-Dur 15 (SH)</i>		
. Cardiovascular system	109	
<i>Nitro-Dur 5 (SH)</i>		
. Cardiovascular system	109	
<i>NITROFURANTOIN</i>		
. Antiinfectives for systemic use	190	
<i>Nitrolingual Pumpspray (SW)</i>		
. Emergency Drug Supplies	64	
. Cardiovascular system	109	
<i>Nizac (LN)</i>		
. Alimentary tract and metabolism	69	
<i>NIZATIDINE</i>		
. Alimentary tract and metabolism	69	
<i>Nizoral (JC)</i>		
. Antiinfectives for systemic use	190	
<i>Nizoral 1% (JT)</i>		
. Dermatologicals	149	
<i>Nizoral 2% (JT)</i>		
. Dermatologicals	149	
. Repatriation Schedule	842	
<i>Nizoral 2% Cream (JT)</i>		
. Dermatologicals	149	
<i>Nolvadex-D (AP)</i>		
. Antineoplastic and immunomodulating agents	231	
<i>Nordette 28 (WY)</i>		
. Genito urinary system and sex hormones	157	
<i>Nordip (AF)</i>		
. Cardiovascular system	118, 119	
<i>Norditropin NordiFlex (NO)</i>		
. Section 100	794	
<i>Norditropin SimpleXx (NO)</i>		
. Section 100	794	
<i>NORETHISTERONE</i>		
. Genito urinary system and sex hormones	158, 161	
<i>NORETHISTERONE with ETHINYLOESTRADIOL</i>		
. Genito urinary system and sex hormones	157, 158	
<i>NORETHISTERONE with MESTRANOL</i>		
. Genito urinary system and sex hormones	158	
<i>Norflohexal (SZ)</i>		
. Antiinfectives for systemic use	188	
<i>NORFLOXACIN</i>		
. Antiinfectives for systemic use	188	
<i>Noriday 28 Day (PF)</i>		
. Genito urinary system and sex hormones	158	
<i>Norimin 28 Day (KR)</i>		
. Genito urinary system and sex hormones	157	
<i>Norimin-1 28 Day (KR)</i>		
. Genito urinary system and sex hormones	158	
<i>Norinyl-1 (PF)</i>		
. Genito urinary system and sex hormones	158	
<i>Norinyl-1/28 (PF)</i>		
. Genito urinary system and sex hormones	158	
<i>Normacol Plus (NE)</i>		
. Alimentary tract and metabolism	80	
. Palliative Care	471	
. Repatriation Schedule	838	
<i>Normison (SI)</i>		
. Nervous system	410	
. Palliative Care	485	
. Dental	510	
<i>Noroxin (MK)</i>		
. Antiinfectives for systemic use	188	
<i>Norprolac (FP)</i>		
. Genito urinary system and sex hormones	157	
<i>Norspan (MF)</i>		
. Nervous system	382	
<i>NORTRIPTYLINE HYDROCHLORIDE</i>		

.Nervous system	412
<i>Norvapine (GN)</i>	
.Cardiovascular system	118, 119
<i>Norvasc (PF)</i>	
.Cardiovascular system	118, 119
<i>Norvir (AB)</i>	
.Section 100	563, 699
<i>Noten (AF)</i>	
.Cardiovascular system	115
<i>Novasone (EX)</i>	
.Dermatologicals	153, 154
<i>Novicrit (NV)</i>	
.Section 100	525, 657
<i>NovoMix 30 FlexPen (NF)</i>	
.Alimentary tract and metabolism	87
<i>NovoMix 30 Penfill 3 mL (NO)</i>	
.Alimentary tract and metabolism	87
<i>NovoRapid (NO)</i>	
.Alimentary tract and metabolism	87
<i>NovoRapid FlexPen (NF)</i>	
.Alimentary tract and metabolism	87
<i>NovoRapid Penfill 3 mL (NO)</i>	
.Alimentary tract and metabolism	87
<i>Noxafil (SH)</i>	
.Antifungives for systemic use	192
<i>Nu-Gel 2497 (JJ)</i>	
.Repatriation Schedule	873
<i>Nucloxx (SI)</i>	
.Repatriation Schedule	838
<i>Nuelin (IA)</i>	
.Respiratory system	438
<i>Nuelin-SR 200 (IA)</i>	
.Respiratory system	438
<i>Nuelin-SR 250 (IA)</i>	
.Respiratory system	438
<i>Nuelin-SR 300 (IA)</i>	
.Respiratory system	438
<i>Nufloxib (AF)</i>	
.Antifungives for systemic use	188
<i>Nupentin 100 (AF)</i>	
.Nervous system	391
.Repatriation Schedule	853
<i>Nupentin 300 (AF)</i>	
.Nervous system	390
.Repatriation Schedule	853
<i>Nupentin 400 (AF)</i>	
.Nervous system	390
.Repatriation Schedule	854
<i>NutropinAq (IS)</i>	
.Section 100	795
<i>Nyogel (NV)</i>	
.Sensory organs	443
.Optometrical	517
NYSTATIN	
.Alimentary tract and metabolism	68, 82
.Dermatologicals	149
.Dental	486, 487
.Repatriation Schedule	842, 847

<i>O.R.S. (AS)</i>	
.Alimentary tract and metabolism	82
OCTREOTIDE	
.Section 100	558, 559, 694, 695
<i>Octreotide MaxRx (XF)</i>	
.Section 100	559, 695
<i>Ocufen (AG)</i>	
.Sensory organs	441
.Optometrical	516
<i>Ocuflox (AG)</i>	
.Sensory organs	441
OESTRADIOL	
.Genito urinary system and sex hormones	159, 160
.Repatriation Schedule	847
OESTRADIOL and OESTRADIOL with DYDROGESTERONE	
.Genito urinary system and sex hormones	162
OESTRADIOL and OESTRADIOL with NORETHISTERONE ACETATE	
.Genito urinary system and sex hormones	162
OESTRADIOL VALERATE	
.Genito urinary system and sex hormones	160
OESTRADIOL with NORETHISTERONE ACETATE	
.Genito urinary system and sex hormones	161
OESTRIOL	
.Genito urinary system and sex hormones	160
OFLOXACIN	
.Sensory organs	441
OLANZAPINE	
.Nervous system	402
OLMESARTAN MEDOXOMIL	
.Cardiovascular system	131
OLMESARTAN MEDOXOMIL with HYDROCHLOROTHIAZIDE	
.Cardiovascular system	132
OLMESARTAN with AMLODIPINE	
.Cardiovascular system	133
<i>Olmetec (SH)</i>	
.Cardiovascular system	131
<i>Olmetec Plus (SH)</i>	
.Cardiovascular system	132
OLSALAZINE SODIUM	
.Alimentary tract and metabolism	85
<i>Omegapharm Irinotecan (OE)</i>	
.Antineoplastic and immunomodulating agents	229
<i>Omegapharm Pty Ltd (OE)</i>	
.Antifungives for systemic use	184
<i>Omepral (PM)</i>	
.Alimentary tract and metabolism	72, 73
OMEPRAZOLE	
.Alimentary tract and metabolism	72, 73
OMEPRAZOLE and CLARITHROMYCIN and AMOXICILLIN	
.Alimentary tract and metabolism	75
<i>Omeprazole generichealth (GQ)</i>	
.Alimentary tract and metabolism	72, 73
<i>Omeprazole Ranbaxy (RA)</i>	
.Alimentary tract and metabolism	72, 73
<i>Omeprazole Winthrop (WA)</i>	
.Alimentary tract and metabolism	72, 73
<i>Omeprazole-GA (GM)</i>	
.Alimentary tract and metabolism	72, 73
<i>Omepro-GA (GM)</i>	

.Alimentary tract and metabolism.....	72, 73
<i>Omnitest EZ (BR)</i>	
.Various.....	455, 456
<i>Omnitrope (SZ)</i>	
.Section 100.....	795
<i>On-Call Plus (PZ)</i>	
.Various.....	456
<i>OncoTICE (SH)</i>	
.Antineoplastic and immunomodulating agents	236
ONDANSETRON	
.Alimentary tract and metabolism.....	76, 77
<i>Ondansetron-Clarix (AE)</i>	
.Alimentary tract and metabolism.....	77
<i>Ondaz (SZ)</i>	
.Alimentary tract and metabolism.....	77
<i>Ondaz Zydis (SZ)</i>	
.Alimentary tract and metabolism.....	77, 78
<i>OneTouch Verio (JJ)</i>	
.Various.....	455, 456
<i>Onkotrone (BX)</i>	
.Antineoplastic and immunomodulating agents	207
<i>Onsetron (ZP)</i>	
.Alimentary tract and metabolism.....	77
<i>Onsetron 4 (ZP)</i>	
.Alimentary tract and metabolism.....	77
<i>Onsetron 8 (ZP)</i>	
.Alimentary tract and metabolism.....	77
<i>Op-Site Flexigrid 4629 (SN)</i>	
.Repatriation Schedule	867
<i>Opticrom (SW)</i>	
.Sensory organs	444
.Optometrical	518
<i>Optium glucose (MS)</i>	
.Various.....	455, 456
<i>Optium Omega (MS)</i>	
.Various.....	455, 456
<i>Optive (AG)</i>	
.Sensory organs	447
.Optometrical	519
<i>Orabase (SI)</i>	
.Repatriation Schedule	843
<i>Oratane (GM)</i>	
.Dermatologicals.....	154
<i>Ordine 10 (MF)</i>	
.Nervous system	377
.Dental	505
<i>Ordine 2 (MF)</i>	
.Nervous system	377
.Dental	505
<i>Ordine 5 (MF)</i>	
.Nervous system	377
.Dental	505
<i>Orencia (BQ)</i>	
.Section 100.....	580, 718
<i>Orgalutran (SH)</i>	
.Section 100.....	797
<i>Orion Laboratories Pty Ltd (ON)</i>	
.Section 100.....	796
ORLISTAT	
.Repatriation Schedule	838
<i>Oroxine (SI)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins.....	171
<i>Orphan Australia Pty Ltd (OA)</i>	
.Nervous system	429
<i>Orudis (SW)</i>	
.Musculo-skeletal system	360
.Dental	502
<i>Orudis SR 200 (SW)</i>	
.Musculo-skeletal system	360
.Dental	502
<i>Oruvail SR (AV)</i>	
.Musculo-skeletal system	360
.Dental	502
<i>OsmoLax (KY)</i>	
.Alimentary tract and metabolism.....	81
.Palliative Care	472
<i>Ospolot (PL)</i>	
.Nervous system	394
<i>Ossmax 70mg (RA)</i>	
.Musculo-skeletal system	365
<i>Otocomb Otic (FM)</i>	
.Sensory organs.....	451
<i>Otodex (AV)</i>	
.Sensory organs.....	451
<i>Ovestin (SH)</i>	
.Genito urinary system and sex hormones	160
<i>Ovestin Ovula (SH)</i>	
.Genito urinary system and sex hormones	160
<i>Ovidrel (SG)</i>	
.Section 100	796
<i>Oxalatin (ZP)</i>	
.Antineoplastic and immunomodulating agents	208
OxALIPLATIN	
.Antineoplastic and immunomodulating agents .	207, 208
<i>Oxaliplatin Actavis (GQ)</i>	
.Antineoplastic and immunomodulating agents	208
<i>Oxaliplatin Alphapharm (AF)</i>	
.Antineoplastic and immunomodulating agents	208
<i>Oxaliplatin Ebewe (IT)</i>	
.Antineoplastic and immunomodulating agents	208
<i>Oxaliplatin Kabi (PK)</i>	
.Antineoplastic and immunomodulating agents	208
<i>Oxaliplatin Link (PK)</i>	
.Antineoplastic and immunomodulating agents	208
OXAZEPAM	
.Nervous system	409
.Palliative Care	484
.Dental	510
OXCARBAZEPINE	
.Nervous system	389
<i>Oxis Turbuhaler (AP)</i>	
.Respiratory system	433
OXPRENOLOL HYDROCHLORIDE	
.Cardiovascular system	114
OXYBUTYNIN	
.Genito urinary system and sex hormones	165
OXYBUTYNIN HYDROCHLORIDE	
.Genito urinary system and sex hormones	165
<i>Oxybutynin Sandoz (SZ)</i>	

.Genito urinary system and sex hormones.....	165
<i>Oxybutynin Winthrop (WA)</i>	
.Genito urinary system and sex hormones.....	165
OXYCODONE	
.Nervous system	379
.Dental	506
OXYCODONE HYDROCHLORIDE	
.Nervous system	379, 380
.Dental	506
<i>OxyContin (MF)</i>	
.Nervous system	380
.Dental	507
OXYMETAZOLINE HYDROCHLORIDE	
.Repatriation Schedule	857
<i>OxyNorm (MF)</i>	
.Nervous system	380
.Dental	506
<i>OxyNorm Liquid 5mg/5mL (MF)</i>	
.Nervous system	380
.Dental	506
<i>Oxytrol (HH)</i>	
.Genito urinary system and sex hormones.....	165
<i>Ozapace (RA)</i>	
.Cardiovascular system.....	124, 125
<i>Ozcef (RA)</i>	
.Antifungives for systemic use	183
.Dental	497
<i>Oziclide MR (RA)</i>	
.Alimentary tract and metabolism.....	90
<i>Ozidal (RA)</i>	
.Nervous system	405, 406, 407
<i>Ozlodip (RA)</i>	
.Cardiovascular system.....	118, 119
<i>Ozmep (ZP)</i>	
.Alimentary tract and metabolism.....	72, 73
<i>Ozole (RA)</i>	
.Antifungives for systemic use	191
<i>Ozpan (RA)</i>	
.Alimentary tract and metabolism.....	73, 74
<i>Ozvir (RA)</i>	
.Antifungives for systemic use	195
P	
<i>PAA (NM)</i>	
.Sensory organs	446
.Optometrical	518
PACLITAXEL	
.Antineoplastic and immunomodulating agents	205
<i>Paclitaxel Actavis (GQ)</i>	
.Antineoplastic and immunomodulating agents .	205, 206
<i>Paclitaxel Ebewe (IT)</i>	
.Antineoplastic and immunomodulating agents .	205, 206
<i>Paclitaxel Kabi (PK)</i>	
.Antineoplastic and immunomodulating agents .	205, 206
<i>Paediatric Seravit (SB)</i>	
.Various.....	467
PALIPERIDONE	
.Nervous system	404
PALONOSETRON	
.Alimentary tract and metabolism.....	78

<i>Pamacid 20 (AF)</i>	
.Alimentary tract and metabolism.....	69
<i>Pamacid 40 (AF)</i>	
.Alimentary tract and metabolism.....	69
<i>Pamisol (HH)</i>	
.Musculo-skeletal system	366
.Section 100	648, 649, 785, 786
<i>Panadeine Forte (SW)</i>	
.Nervous system	375
.Dental	504
<i>Panadol (GC)</i>	
.Palliative Care	482
<i>Panadol Osteo (GC)</i>	
.Nervous system	385
.Palliative Care	482
<i>Panafcort (AS)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins.....	170
<i>Panafcortelone (AS)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins.....	170
<i>Panamax (SW)</i>	
.Nervous system	384
.Dental	508, 509
<i>Panamax 240 Elixir (SW)</i>	
.Nervous system	384
.Dental	508
<i>Panamax Co. (SW)</i>	
.Repatriation Schedule	853
PANCREATIC EXTRACT	
.Alimentary tract and metabolism.....	85, 86
PANCRELIPASE	
.Alimentary tract and metabolism.....	86
<i>Panto (NZ)</i>	
.Alimentary tract and metabolism.....	73, 74
<i>Pantofast 20 (RZ)</i>	
.Alimentary tract and metabolism.....	74
<i>Pantofast 40 (RZ)</i>	
.Alimentary tract and metabolism.....	73, 74
<i>Pantoloc (NH)</i>	
.Alimentary tract and metabolism.....	73, 74
<i>Pantoprazole generichealth (GQ)</i>	
.Alimentary tract and metabolism.....	74
<i>Pantoprazole Sandoz (SZ)</i>	
.Alimentary tract and metabolism.....	74
PANTOPRAZOLE SODIUM SESQUIHYDRATE	
.Alimentary tract and metabolism.....	73, 74
<i>Pantoprazole-GA (GM)</i>	
.Alimentary tract and metabolism.....	74
<i>Panzytrat 25000 (TM)</i>	
.Alimentary tract and metabolism.....	86
PARACETAMOL	
.Nervous system	384, 385
.Palliative Care	482
.Dental	508, 509
<i>Paracetamol Sandoz (SZ)</i>	
.Nervous system	384
.Dental	508, 509
PARAFFIN	
.Sensory organs.....	448

.Optometrical	520	.Musculo-skeletal system	362
<i>Paralgin (FM)</i>		<i>Pentasa (FP)</i>	
.Nervous system	384	.Alimentary tract and metabolism	84, 85
.Dental	508, 509	<i>Penthrox (NQ)</i>	
<i>Pariet (JC)</i>		.Emergency Drug Supplies	64
.Alimentary tract and metabolism	75	<i>Pepcidine (MK)</i>	
<i>Parlodel (NV)</i>		.Alimentary tract and metabolism	69
.Genito urinary system and sex hormones	156	<i>Pepcidine M (MK)</i>	
.Nervous system	397	.Alimentary tract and metabolism	69
<i>Parnate (GH)</i>		<i>Pepti-Junior Gold (NU)</i>	
.Nervous system	416	.Various	460
PAROXETINE		<i>Pepzan (GM)</i>	
.Nervous system	414	.Alimentary tract and metabolism	69
<i>Paroxetine 20 (CR)</i>		PERGOLIDE MESYLATE	
.Nervous system	415	.Nervous system	398
<i>Paroxetine generichealth (GQ)</i>		PERHEXILINE MALEATE	
.Nervous system	415	.Cardiovascular system	110
<i>Paroxetine Sandoz (SZ)</i>		<i>Periactin (AS)</i>	
.Nervous system	415	.Nervous system	386
<i>Paroxetine-DP (GM)</i>		PERICYAZINE	
.Nervous system	415	.Nervous system	400
<i>Paroxetine-GA (GN)</i>		<i>Perindo (AF)</i>	
.Nervous system	415	.Cardiovascular system	124, 125
<i>Paxam 0.5 (AF)</i>		<i>Perindo Combi 4/1.25 (AF)</i>	
.Nervous system	388	.Cardiovascular system	129
.Palliative Care	483	PERINDOPRIL	
<i>Paxam 2 (AF)</i>		.Cardiovascular system	124, 125
.Nervous system	388	<i>Perindopril 2 (CR)</i>	
.Palliative Care	483	.Cardiovascular system	124
<i>Paxtine (AF)</i>		<i>Perindopril 4 (CR)</i>	
.Nervous system	415	.Cardiovascular system	125
<i>Peg 7420 (BK)</i>		<i>Perindopril 8 (CR)</i>	
.Repatriation Schedule	864	.Cardiovascular system	125
<i>Peg 7422 (BK)</i>		PERINDOPRIL with AMLODIPINE	
.Repatriation Schedule	864	.Cardiovascular system	130
<i>Peg 7423 (BK)</i>		PERINDOPRIL with INDAPAMIDE HEMIHYDRATE	
.Repatriation Schedule	864	.Cardiovascular system	129
<i>Peg 7425 (BK)</i>		<i>Perindopril-DP (GN)</i>	
.Repatriation Schedule	864	.Cardiovascular system	124, 125
<i>PEG-Intron Redipen (SH)</i>		<i>Perindopril-GA (GM)</i>	
.Section 100	573, 711	.Cardiovascular system	124, 125
<i>Pegasys (RO)</i>		<i>Perivasc (AL)</i>	
.Section 100	572, 710	.Cardiovascular system	118, 119
<i>Pegasys RBV (RO)</i>		<i>Permax (AS)</i>	
.Section 100	574, 712	.Nervous system	398
<i>Pegatron (SH)</i>		PERMETHRIN	
.Section 100	575, 713	.Antiparasitic products, insecticides and repellents	432
PEGFILGRASTIM		<i>Persantin SR (BY)</i>	
.Section 100	570, 708	.Blood and blood forming organs	100
<i>PEGINTERFERON ALFA-2a</i>		<i>Petrus Bisacodyl Suppositories (PP)</i>	
.Section 100	572, 710	.Alimentary tract and metabolism	80
<i>PEGINTERFERON ALFA-2b</i>		.Palliative Care	471
.Section 100	573, 710	<i>Petrus Pharmaceuticals Pty Ltd (PP)</i>	
PEMETREXED DISODIUM		.Alimentary tract and metabolism	81
.Antineoplastic and immunomodulating agents	200	.Palliative Care	474
<i>Pemzo (SI)</i>		.Repatriation Schedule	838
.Alimentary tract and metabolism	72, 73	<i>Pexsig (SI)</i>	
<i>Pendine 800 (AF)</i>		.Cardiovascular system	110
.Nervous system	391	<i>Pfizer Australia Pty Ltd (PF)</i>	
PENICILLAMINE		.Emergency Drug Supplies	64, 65

.Alimentary tract and metabolism.....	77, 78
.Blood and blood forming organs.....	98
.Cardiovascular system.....	107
.Antiinfectives for systemic use.....	187
.Antineoplastic and immunomodulating agents	200, 202, 203, 207
.Respiratory system.....	434
.Various.....	467
.Dental.....	489, 513
<i>Pharmacor Amlodipine 10 (CR)</i>	
.Cardiovascular system.....	119
<i>Pharmacor Amlodipine 5 (CR)</i>	
.Cardiovascular system.....	118
<i>Pharmacor Gemfibrozil 600 (CR)</i>	
.Cardiovascular system.....	145
<i>Pharmacor Meloxicam 15 (CR)</i>	
.Musculo-skeletal system.....	359
<i>Pharmacor Meloxicam 7.5 (CR)</i>	
.Musculo-skeletal system.....	359
<i>Pharmacor Metformin 1000 (CR)</i>	
.Alimentary tract and metabolism.....	89
<i>Pharmacor Omeprazole 20 (CR)</i>	
.Alimentary tract and metabolism.....	72, 73
<i>Pharmacor Paroxo 20 (CR)</i>	
.Nervous system.....	415
<i>Pharmacor Quinapril 10 (CR)</i>	
.Cardiovascular system.....	125
<i>Pharmacor Quinapril 20 (CR)</i>	
.Cardiovascular system.....	126
<i>Pharmacor Quinapril 5 (CR)</i>	
.Cardiovascular system.....	125
<i>Pharmacor Ramipril 1.25 (CR)</i>	
.Cardiovascular system.....	126
<i>Pharmacor Ramipril 10 (CR)</i>	
.Cardiovascular system.....	127
<i>Pharmacor Ramipril 2.5 (CR)</i>	
.Cardiovascular system.....	126
<i>Pharmacor Ramipril 5 (CR)</i>	
.Cardiovascular system.....	127
<i>Pharmacor Salbutamol 2.5 (CR)</i>	
.Emergency Drug Supplies.....	65
.Respiratory system.....	434
<i>Pharmacor Salbutamol 5 (CR)</i>	
.Emergency Drug Supplies.....	65
.Respiratory system.....	434
<i>Pharmacor Simvastatin 10 (MI)</i>	
.Cardiovascular system.....	141, 142
<i>Pharmacor Simvastatin 20 (MI)</i>	
.Cardiovascular system.....	141, 143
<i>Pharmacor Simvastatin 40 (MI)</i>	
.Cardiovascular system.....	141, 143
<i>Pharmacor Simvastatin 80 (MI)</i>	
.Cardiovascular system.....	142, 144
<i>Pharmacor Sumatriptan 50 (CR)</i>	
.Nervous system.....	386
<i>Pharmacy Choice Paracetamol (YM)</i>	
.Nervous system.....	384
.Dental.....	508, 509
<i>Pharmorubicin Solution (PF)</i>	
.Antineoplastic and immunomodulating agents.....	206
<i>Phenasen (PL)</i>	
.Antineoplastic and immunomodulating agents.....	226
<i>PHENELZINE SULFATE</i>	
.Nervous system.....	416
<i>Phenergan (SW)</i>	
.Palliative Care.....	470, 471
.Repatriation Schedule.....	858
<i>Phenex-2 (AB)</i>	
.Various.....	463
<i>PHENOBARBITONE</i>	
.Nervous system.....	386
<i>PHENOBARBITONE SODIUM</i>	
.Nervous system.....	387
<i>PHENOXYBENZAMINE HYDROCHLORIDE</i>	
.Cardiovascular system.....	114
.Genito urinary system and sex hormones.....	165
<i>PHENOXYMETHYLPENICILLIN</i>	
.Antiinfectives for systemic use.....	178, 179
.Dental.....	493
<i>Phenylalanine Amino Acid Supplement (VF)</i>	
.Various.....	466
<i>PHENYLALANINE with CARBOHYDRATE</i>	
.Various.....	466
<i>PHENYTOIN</i>	
.Nervous system.....	387
<i>PHENYTOIN SODIUM</i>	
.Nervous system.....	387
<i>Phlexy-10 (SB)</i>	
.Various.....	462
<i>Phlexy-10 Drink Mix (SB)</i>	
.Various.....	462
<i>PHOLCODINE</i>	
.Repatriation Schedule.....	857
<i>Phosphate Sandoz (NV)</i>	
.Various.....	453
<i>Physeptone (SI)</i>	
.Nervous system.....	381
<i>Physiotens (SM)</i>	
.Cardiovascular system.....	111
<i>Piax (AF)</i>	
.Blood and blood forming organs.....	99
<i>PILOCARPINE HYDROCHLORIDE</i>	
.Sensory organs.....	442
.Optometrical.....	517
<i>PIMECROLIMUS</i>	
.Dermatologicals.....	154
<i>PINDOLOL</i>	
.Cardiovascular system.....	114
<i>PINE TAR with CADE OIL, COAL TAR SOLUTION, ARACHIS OIL EXTRACT OF CRUDE COAL TAR and OLEYL ALCOHOL</i>	
.Repatriation Schedule.....	845
<i>PINE TAR with TRIETHANOLAMINE LAURYL SULFATE</i>	
.Repatriation Schedule.....	844
<i>Pinetarsol (EO)</i>	
.Repatriation Schedule.....	844
<i>PIOGLITAZONE</i>	
.Alimentary tract and metabolism.....	92
<i>PIROXICAM</i>	
.Musculo-skeletal system.....	360
.Dental.....	502

<i>PIZOTIFEN MALATE</i>		
.Nervous system	386	
<i>PKU Anamix infant (SB)</i>		
.Variou.....	462	
<i>PKU Anamix Junior (SB)</i>		
.Variou.....	464	
<i>PKU Anamix Junior LQ (SB)</i>		
.Variou.....	464	
<i>PKU Cooler 10 (VF)</i>		
.Variou.....	463	
<i>PKU Cooler 15 (VF)</i>		
.Variou.....	464	
<i>PKU Cooler 20 (VF)</i>		
.Variou.....	463	
<i>PKU Lophlex LQ 10 (SB)</i>		
.Variou.....	464	
<i>PKU Lophlex LQ 20 (SB)</i>		
.Variou.....	464	
<i>PKU-Express (VF)</i>		
.Variou.....	463	
<i>PKU-gel (VF)</i>		
.Variou.....	463	
<i>Placil (AF)</i>		
.Nervous system	409, 411	
<i>Plaqacide (OB)</i>		
.Repatriation Schedule	837	
<i>Plaquenil (SW)</i>		
.Musculo-skeletal system	362	
<i>Plasma-Lyte 148 (BX)</i>		
.Blood and blood forming organs	105	
<i>Plavix (SW)</i>		
.Blood and blood forming organs	99	
.Repatriation Schedule	840	
<i>Plaxel (WQ)</i>		
.Antineoplastic and immunomodulating agents .	205, 206	
<i>Plendil ER (AP)</i>		
.Cardiovascular system	119	
<i>PNEUMOCOCCAL VACCINE, POLYVALENT</i>		
.Anti-infectives for systemic use	198	
<i>Pneumovax 23 (CS)</i>		
.Anti-infectives for systemic use	198	
<i>PODOPHYLLOTOXIN</i>		
.Repatriation Schedule	844	
<i>Poly Gel (AQ)</i>		
.Sensory organs	446	
.Optomotriral	519	
<i>Poly Visc (IQ)</i>		
.Sensory organs	448, 449	
.Optomotriral	520	
<i>POLY-L-LACTIC ACID</i>		
.Variou.....	454	
<i>Poly-Tears (IQ)</i>		
.Sensory organs	448	
.Optomotriral	520	
<i>POLYETHYLENE GLYCOL 400</i>		
.Sensory organs	449	
.Optomotriral	520	
<i>POLYETHYLENE GLYCOL 400 with PROPYLENE GLYCOL</i>		
.Sensory organs	449, 450	
.Optomotriral	520	
<i>POLYGELINE</i>		
.Blood and blood forming organs	105	
<i>Polytar (GK)</i>		
.Repatriation Schedule	845	
<i>POLYVINYL ALCOHOL</i>		
.Sensory organs.....	450	
.Optomotriral.....	520	
<i>Ponstan (PD)</i>		
.Musculo-skeletal system	361	
<i>POSACONAZOLE</i>		
.Anti-infectives for systemic use	192	
<i>POTASSIUM CHLORIDE</i>		
.Alimentary tract and metabolism	95	
<i>POTASSIUM CHLORIDE with POTASSIUM BICARBONATE</i>		
.Alimentary tract and metabolism	96	
<i>POVIDONE-IODINE</i>		
.Repatriation Schedule	845	
<i>Pradaxa (BY)</i>		
.Blood and blood forming organs	102	
<i>Pramin (AF)</i>		
.Alimentary tract and metabolism	76	
.Dental	486	
<i>PRAMIPEXOLE HYDROCHLORIDE</i>		
.Nervous system	398, 399	
<i>Prantal (SH)</i>		
.Repatriation Schedule	845	
<i>PRASUGREL</i>		
.Blood and blood forming organs	100	
<i>Pravachol (FM)</i>		
.Cardiovascular system	138, 139, 140	
<i>PRAVASTATIN</i>		
.Cardiovascular system	138, 139	
<i>Pravastatin 10 (CR)</i>		
.Cardiovascular system	138, 139	
<i>Pravastatin 20 (CR)</i>		
.Cardiovascular system	138, 139	
<i>Pravastatin 40 (CR)</i>		
.Cardiovascular system	138, 140	
<i>Pravastatin generichealth (GQ)</i>		
.Cardiovascular system	138, 139, 140	
<i>Pravastatin Sandoz (SZ)</i>		
.Cardiovascular system	138, 139, 140	
<i>Pravastatin Winthrop (WA)</i>		
.Cardiovascular system	138, 139, 140	
<i>Pravastatin-GA 10 (GM)</i>		
.Cardiovascular system	138, 139	
<i>Pravastatin-GA 20 (GM)</i>		
.Cardiovascular system	138, 139	
<i>Pravastatin-GA 40 (GM)</i>		
.Cardiovascular system	138, 140	
<i>Pravastatin-GA 80 (GM)</i>		
.Cardiovascular system	139, 140	
<i>PRAZIQUANTEL</i>		
.Antiparasitic products, insecticides and repellents	431	
<i>PRAZOSIN HYDROCHLORIDE</i>		
.Cardiovascular system	111	
<i>PredMix (LN)</i>		
.Systemic hormonal preparations, excl. sex hormones and insulins.....	170	
<i>Prednefrin Forte (AG)</i>		

.Sensory organs	441	.Musculo-skeletal system	364
PREDNISOLONE		PROBENECID	
.Systemic hormonal preparations, excl. sex hormones and insulins	170	.Musculo-skeletal system	364
PREDNISOLONE ACETATE with PHENYLEPHRINE HYDROCHLORIDE		Probitor (SZ)	
.Sensory organs	441	.Alimentary tract and metabolism	72, 73
PREDNISOLONE SODIUM PHOSPHATE		Probitor Hp7 (SZ)	
.Alimentary tract and metabolism	83	.Alimentary tract and metabolism	75
.Systemic hormonal preparations, excl. sex hormones and insulins	170	PROCAINE PENICILLIN	
PREDNISONE		.Antiinfectives for systemic use	179
.Systemic hormonal preparations, excl. sex hormones and insulins	170	.Dental	494
Predsol (SI)		ProCalm (SI)	
.Alimentary tract and metabolism	83	.Alimentary tract and metabolism	79
Predsolone (LN)		.Dental	486
.Systemic hormonal preparations, excl. sex hormones and insulins	170	PROCHLORPERAZINE	
Predstone (LN)		.Alimentary tract and metabolism	79
.Systemic hormonal preparations, excl. sex hormones and insulins	170	.Dental	486
PREGABALIN		Prochlorperazine-GA (GM)	
.Repatriation Schedule	854	.Alimentary tract and metabolism	79
Pregnyl (SH)		.Dental	486
.Genito urinary system and sex hormones	163	Proctosedyl (SW)	
.Section 100	797	.Repatriation Schedule	841
Presolol 100 (AF)		Procur (GM)	
.Cardiovascular system	118	.Genito urinary system and sex hormones	164
Presolol 200 (AF)		.Antineoplastic and immunomodulating agents	232
.Cardiovascular system	118	Procur 100 (GM)	
Prexaton (AF)		.Genito urinary system and sex hormones	164
.Nervous system	427	.Antineoplastic and immunomodulating agents	232
Prezista (JC)		Prodeine 15 (SW)	
.Section 100	562, 698	.Repatriation Schedule	853
Prilace 1.25 (SI)		Prodeine Forte (AV)	
.Cardiovascular system	126	.Nervous system	375
Prilace 10 (SI)		.Dental	504
.Cardiovascular system	127	Profloxin (HX)	
Prilace 2.5 (SI)		.Antiinfectives for systemic use	187
.Cardiovascular system	126	Profore 66050016 (SN)	
Prilace 5 (SI)		.Repatriation Schedule	864
.Cardiovascular system	127	Profore Lite 66050415 (SN)	
PRIMIDONE		.Repatriation Schedule	864
.Nervous system	387	PROGESTERONE	
Primolut N (SC)		.Section 100	796
.Genito urinary system and sex hormones	161	Progut 100 (AF)	
Primoteston Depot (SC)		.Musculo-skeletal system	364
.Genito urinary system and sex hormones	159	Progut 300 (AF)	
Prinivil 10 (MK)		.Musculo-skeletal system	364
.Cardiovascular system	124	Prograf (JC)	
Prinivil 20 (MK)		.Antineoplastic and immunomodulating agents	356
.Cardiovascular system	124	.Section 100	640, 777
Prinivil 5 (MK)		Prograf XL (JC)	
.Cardiovascular system	123	.Antineoplastic and immunomodulating agents	356
Pristiq (WX)		.Section 100	640, 777
.Nervous system	417	ProGuide 66000780 (SN)	
Pro-Banthine (SI)		.Repatriation Schedule	864
.Genito urinary system and sex hormones	165	ProGuide 66000781 (SN)	
Pro-Cid (PL)		.Repatriation Schedule	864
		ProGuide 66000782 (SN)	
		.Repatriation Schedule	864
		Progynova (SC)	
		.Genito urinary system and sex hormones	160
		Proladone (PL)	
		.Nervous system	379

.Dental	506	.Respiratory system	436
<i>Prolia (AN)</i>		<i>Pulmicort Turbuhaler (AP)</i>	
.Musculo-skeletal system	372	.Respiratory system	436
<i>PROMETHAZINE HYDROCHLORIDE</i>		<i>Pulmozyme (RO)</i>	
.Respiratory system	439	.Section 100	652, 789
.Palliative Care	470	<i>Puregon 300 IU/0.36 mL (SH)</i>	
.Dental	486	.Genito urinary system and sex hormones	163
.Repatriation Schedule	858	.Section 100	796
<i>PROPANTHELINE BROMIDE</i>		<i>Puregon 600 IU/0.72 mL (SH)</i>	
.Genito urinary system and sex hormones	165	.Genito urinary system and sex hormones	163
<i>PROPRANOLOL HYDROCHLORIDE</i>		.Section 100	796
.Cardiovascular system	114	<i>Puregon 900 IU/1.08 mL (SH)</i>	
<i>PROPYLTHIOURACIL</i>		.Genito urinary system and sex hormones	163
.Systemic hormonal preparations, excl. sex hormones and insulins	171	.Section 100	797
<i>Proscar (MK)</i>		<i>Purinethol (AS)</i>	
.Repatriation Schedule	848	.Antineoplastic and immunomodulating agents	202
<i>Protaphane (NO)</i>		<i>PVA Forte (PE)</i>	
.Alimentary tract and metabolism	87	.Sensory organs	450
<i>Protaphane InnoLet (NI)</i>		.Optometrical	521
.Alimentary tract and metabolism	87	<i>PVA Tears (PE)</i>	
<i>Protaphane NovoLet 3 mL (NL)</i>		.Sensory organs	450
.Alimentary tract and metabolism	87	.Optometrical	521
<i>Protaphane Penfill 3 mL (NO)</i>		<i>Pyralin EN (KR)</i>	
.Alimentary tract and metabolism	87	.Alimentary tract and metabolism	85
<i>PROTEIN HYDROLYSATE FORMULA with MEDIUM CHAIN TRIGLYCERIDES</i>		<i>PYRANTEL EMBONATE</i>	
.Various	459, 460	.Antiparasitic products, insecticides and repellents	432
<i>Prothiaden (AB)</i>		<i>PYRIDOSTIGMINE BROMIDE</i>	
.Nervous system	411	.Nervous system	427
<i>Protos 2 g (SE)</i>		<i>PYRIMETHAMINE</i>	
.Musculo-skeletal system	372	.Antiparasitic products, insecticides and repellents	430
<i>Provera (PF)</i>		Q	
.Genito urinary system and sex hormones	161	<i>Qpril 10 (AF)</i>	
.Antineoplastic and immunomodulating agents	230	.Cardiovascular system	125
<i>Proxen SR 1000 (MD)</i>		<i>Qpril 20 (AF)</i>	
.Musculo-skeletal system	361	.Cardiovascular system	126
.Palliative Care	478	<i>Qpril 5 (AF)</i>	
.Dental	503	.Cardiovascular system	125
<i>Proxen SR 750 (MD)</i>		<i>Questran Lite (SI)</i>	
.Musculo-skeletal system	360	.Cardiovascular system	145, 146
.Palliative Care	478	<i>QUETIAPINE</i>	
.Dental	503	.Nervous system	402
<i>Prozac 20 (LY)</i>		<i>Quilonum SR (GK)</i>	
.Nervous system	414	.Nervous system	403, 417
<i>Prozac Tab (LY)</i>		<i>QUINAGOLIDE HYDROCHLORIDE</i>	
.Nervous system	414	.Genito urinary system and sex hormones	157
<i>ProZero (VF)</i>		<i>QUINAPRIL</i>	
.Various	466	.Cardiovascular system	125
<i>PSEUDOEPHEDRINE HYDROCHLORIDE</i>		<i>Quinapril generichealth (GQ)</i>	
.Repatriation Schedule	857	.Cardiovascular system	125, 126
<i>PSYLLIUM HYDROPHILIC MUCILLOID</i>		<i>QUINAPRIL HYDROCHLORIDE with HYDROCHLOROTHIAZIDE</i>	
.Repatriation Schedule	838	.Cardiovascular system	129
<i>PSYLLIUM HYDROPHILIC MUCILLOID with HIGH AMYLOSE MAIZE STARCH</i>		<i>Quinapril Sandoz (SZ)</i>	
.Repatriation Schedule	838	.Cardiovascular system	125, 126
<i>PTU (PL)</i>		<i>Quinapril-DP (GN)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	171	.Cardiovascular system	125
<i>Pulmicort Respules (AP)</i>		<i>Quinapril-GA (GM)</i>	
		.Cardiovascular system	126
		<i>Quinate (AS)</i>	
		.Antiparasitic products, insecticides and repellents	431

QUININE SULFATE

.Antiparasitic products, insecticides and repellents.... 431

QuitX (AF)

.Repatriation Schedule 855

QV Bath Oil (EO)

.Repatriation Schedule 843

Qvar 100 (IA)

.Respiratory system 435

Qvar 100 Autohaler (IA)

.Respiratory system 435

Qvar 50 (IA)

.Respiratory system 435

Qvar 50 Autohaler (IA)

.Respiratory system 435

R**RABEPRAZOLE SODIUM**

.Alimentary tract and metabolism..... 75

Ralovera (KR)

.Genito urinary system and sex hormones..... 161

RALOXIFENE HYDROCHLORIDE

.Genito urinary system and sex hormones..... 165

.Musculo-skeletal system 372

RALTEGRAVIR

.Section 100 568, 705

RALTITREXED

.Antineoplastic and immunomodulating agents 201

Ramace 1.25 mg (AV)

.Cardiovascular system..... 126

Ramace 10 mg (AV)

.Cardiovascular system..... 127

Ramace 2.5 mg (AV)

.Cardiovascular system..... 126

Ramace 5 mg (AV)

.Cardiovascular system..... 127

RAMIPRIL

.Cardiovascular system..... 126, 127, 128

Ramipril generichealth (GQ)

.Cardiovascular system..... 126, 127

Ramipril Sandoz (SZ)

.Cardiovascular system..... 126, 127

Ramipril Winthrop (WA)

.Cardiovascular system..... 126, 127

RAMIPRIL with FELODIPINE

.Cardiovascular system..... 130

Ramipril-DP (GN)

.Cardiovascular system..... 126, 127

Ramipril-GA (GM)

.Cardiovascular system..... 126, 127

Rancef (RA)

.Antinfectives for systemic use 181, 182

.Dental 496

Rani 2 (AF)

.Alimentary tract and metabolism..... 69

RANIBIZUMAB

.Sensory organs 444

Ranihexal (HX)

.Alimentary tract and metabolism..... 70

RANITIDINE HYDROCHLORIDE

.Alimentary tract and metabolism..... 69, 70

Ranitidine Sandoz (SZ)

.Alimentary tract and metabolism..... 69, 70

Ranmoxy (RA)

.Antinfectives for systemic use 177, 178

.Dental 493

Ranoxyl (GM)

.Alimentary tract and metabolism..... 70

Ransim (RA)

.Cardiovascular system 141, 142, 143, 144

Ranzepam (RA)

.Nervous system 408

.Palliative Care 483, 484

.Dental 509

Rapamune (WX)

.Antineoplastic and immunomodulating agents 238

.Section 100 582, 719

Rapilysin 10 U (TA)

.Blood and blood forming organs 101

RBX Topiramate (RA)

.Nervous system 394, 395

RCF (AB)

.Various..... 466

Reandron 1000 (SC)

.Genito urinary system and sex hormones 159

Rebif 44 (SG)

.Antineoplastic and immunomodulating agents 235

REBOXETINE MESILATE

.Nervous system 418

Redipred (AS)

.Systemic hormonal preparations, excl. sex hormones
and insulins..... 170

Refresh Liquigel (AG)

.Sensory organs..... 446

.Optometrical..... 519

Refresh Tears Plus (AG)

.Sensory organs..... 446

.Optometrical..... 519

Relistor (WX)

.Palliative Care 473, 474

Relpax (PF)

.Nervous system 385

Remicade (SH)

.Section 100 .597, 602, 607, 620, 624, 627, 634, 734, 739,
744, 757, 761, 764, 771

.Repatriation Schedule 851

Reminyl (JC)

.Nervous system 423

Renagel (GZ)

.Various..... 453

.Section 100 655, 792

RenaStart (VF)

.Various..... 467

Renitec (MK)

.Cardiovascular system 122

Renitec 20 (MK)

.Cardiovascular system 122

Renitec M (MK)

.Cardiovascular system 123

Renitec Plus 20/6 (MK)

.Cardiovascular system 128

<i>ReoPro (LY)</i>		
.Blood and blood forming organs	98	
<i>Repalyte New Formulation (SW)</i>		
.Alimentary tract and metabolism.....	82	
<i>Replicare Ultra 66000434 (SN)</i>		
.Repatriation Schedule	871	
<i>Resdone 0.5 (CR)</i>		
.Nervous system	405	
<i>Resdone 1 (CR)</i>		
.Nervous system	405, 406	
<i>Resdone 2 (CR)</i>		
.Nervous system	406, 407	
<i>Resdone 3 (CR)</i>		
.Nervous system	406	
<i>Resdone 4 (CR)</i>		
.Nervous system	406	
<i>Resonium-A (SW)</i>		
.Repatriation Schedule	860	
<i>Resprim (AF)</i>		
.Anti-infectives for systemic use	184	
.Dental	497	
<i>Resprim Forte (AF)</i>		
.Anti-infectives for systemic use	184	
.Dental	498	
<i>restore O.R.S. (GM)</i>		
.Alimentary tract and metabolism.....	82	
<i>RETEPLASE (Recombinant plasminogen activator)</i>		
.Blood and blood forming organs	101	
<i>Retrovir (GK)</i>		
.Section 100	566, 703	
<i>Revatio (PF)</i>		
.Section 100	552, 689	
<i>ReVia (BQ)</i>		
.Nervous system	428	
<i>Revlimid (CI)</i>		
.Section 100	642, 779	
<i>Reyataz (BQ)</i>		
.Section 100	562, 698	
<i>Riamet (NV)</i>		
.Antiparasitic products, insecticides and repellents....	431	
<i>Riamet 20mg/120mg Dispersible (NV)</i>		
.Antiparasitic products, insecticides and repellents....	431	
<i>RIBAVIRIN and PEGINTERFERON ALFA-2a</i>		
.Section 100	573, 711	
<i>RIBAVIRIN and PEGINTERFERON ALFA-2b</i>		
.Section 100	574, 712	
<i>RICINOLEIC ACID with ACETIC ACID and HYDROXYQUINOLINE SULFATE</i>		
.Repatriation Schedule	847	
<i>Ridaura (GH)</i>		
.Musculo-skeletal system	362	
<i>RIFABUTIN</i>		
.Section 100	561, 697	
<i>Rifadin (SW)</i>		
.Anti-infectives for systemic use	194	
<i>RIFAMPICIN</i>		
.Anti-infectives for systemic use	193, 194	
<i>Rilutek (SW)</i>		
.Nervous system	429	
<i>RILUZOLE</i>		
.Nervous system	428	
<i>Rimycin 150 (AF)</i>		
.Anti-infectives for systemic use	194	
<i>Rimycin 300 (AF)</i>		
.Anti-infectives for systemic use	194	
<i>RISEDRONATE SODIUM</i>		
.Musculo-skeletal system	367	
.Repatriation Schedule	852	
<i>RISEDRONATE SODIUM and CALCIUM CARBONATE</i>		
.Musculo-skeletal system	370	
.Repatriation Schedule	852	
<i>RISEDRONATE SODIUM and CALCIUM CARBONATE with COLECALCIFEROL</i>		
.Musculo-skeletal system	371	
.Repatriation Schedule	852	
<i>Rispa (SI)</i>		
.Nervous system	405, 406, 407	
<i>Risperdal (JC)</i>		
.Nervous system	405, 406, 407	
<i>Risperdal Consta (JC)</i>		
.Nervous system	407	
<i>Risperdal Quicklet (JC)</i>		
.Nervous system	405, 406, 407	
<i>RISPERIDONE</i>		
.Nervous system	404, 405, 407	
<i>Risperidone generichealth (GQ)</i>		
.Nervous system	405, 406, 407	
<i>Risperidone-DRLA (RZ)</i>		
.Nervous system	405, 406, 407	
<i>Risperidone-GA (GM)</i>		
.Nervous system	405, 406, 407	
<i>Ritalin 10 (NV)</i>		
.Nervous system	420	
<i>Ritalin LA (NV)</i>		
.Nervous system	420	
<i>Rithmik 100 (SI)</i>		
.Cardiovascular system	108	
<i>Rithmik 200 (SI)</i>		
.Cardiovascular system	108	
<i>RITONAVIR</i>		
.Section 100	563, 699	
<i>RITUXIMAB</i>		
.Antineoplastic and immunomodulating agents	209	
.Section 100	642, 779	
<i>RIVAROXABAN</i>		
.Blood and blood forming organs	103	
<i>RIVASTIGMINE</i>		
.Nervous system	423	
<i>RIVASTIGMINE HYDROGEN TARTRATE</i>		
.Nervous system	424	
<i>Rivotril (RO)</i>		
.Emergency Drug Supplies	64	
.Nervous system	388	
.Palliative Care	483	
<i>Rixadone (AF)</i>		
.Nervous system	405, 406, 407	
<i>RIZATRIPTAN</i>		
.Nervous system	385	
<i>Roaccutane (RO)</i>		
.Dermatologicals	154	

<i>Sebivo (NV)</i>		
. Section 100	565, 702	
<i>Sebizole (GM)</i>		
. Repatriation Schedule	842	
<i>SELEGILINE HYDROCHLORIDE</i>		
. Nervous system	399	
<i>SELENIUM SULFIDE</i>		
. Repatriation Schedule	845	
<i>Selgene (AF)</i>		
. Nervous system	399	
<i>Selsun (DQ)</i>		
. Repatriation Schedule	845	
<i>SENEGA and AMMONIA</i>		
. Repatriation Schedule	857	
<i>SENNA STANDARDISED</i>		
. Repatriation Schedule	838	
<i>Senokot (RC)</i>		
. Repatriation Schedule	838	
<i>Sensipar (AN)</i>		
. Systemic hormonal preparations, excl. sex hormones and insulins	173, 174	
. Section 100	560, 696	
<i>SensoCard (PX)</i>		
. Various	455, 456	
<i>Septrin (SI)</i>		
. Anti-infectives for systemic use	185	
. Dental	498	
<i>Septrin Forte (SI)</i>		
. Anti-infectives for systemic use	185	
. Dental	498	
<i>Serenace (SI)</i>		
. Emergency Drug Supplies	64	
. Nervous system	401	
<i>Serepax (SI)</i>		
. Nervous system	409	
. Palliative Care	484	
. Dental	510	
<i>Seretide Accuhaler 100/50 (GK)</i>		
. Respiratory system	435	
<i>Seretide Accuhaler 250/50 (GK)</i>		
. Respiratory system	435	
<i>Seretide Accuhaler 500/50 (GK)</i>		
. Respiratory system	435	
<i>Seretide MDI 125/25 (GK)</i>		
. Respiratory system	435	
<i>Seretide MDI 250/25 (GK)</i>		
. Respiratory system	435	
<i>Seretide MDI 50/25 (GK)</i>		
. Respiratory system	435	
<i>Serevent Accuhaler (GK)</i>		
. Respiratory system	434	
<i>Serophene (SG)</i>		
. Genito urinary system and sex hormones	163	
<i>Seroquel (AP)</i>		
. Nervous system	402, 403	
<i>Seroquel XR (AP)</i>		
. Nervous system	402, 403	
<i>Sertra 100 (SI)</i>		
. Nervous system	415	
<i>Sertra 50 (SI)</i>		
. Nervous system	415	
<i>Sertraline 100 (CR)</i>		
. Nervous system	415	
<i>Sertraline 50 (CR)</i>		
. Nervous system	415	
<i>Sertraline generichealth (GQ)</i>		
. Nervous system	415	
<i>SERTRALINE HYDROCHLORIDE</i>		
. Nervous system	415, 416	
<i>Sertraline Winthrop (WA)</i>		
. Nervous system	415	
<i>Sertraline-GA (GM)</i>		
. Nervous system	415	
<i>Setopress 3504 (SS)</i>		
. Repatriation Schedule	863	
<i>Setopress 3505 (SS)</i>		
. Repatriation Schedule	863	
<i>Setrona (RA)</i>		
. Nervous system	415	
<i>SEVELAMER HYDROCHLORIDE</i>		
. Various	452	
. Section 100	654, 792	
<i>Sevikar 20/5 (SH)</i>		
. Cardiovascular system	133	
<i>Sevikar 40/10 (SH)</i>		
. Cardiovascular system	133	
<i>Sevikar 40/5 (SH)</i>		
. Cardiovascular system	133	
<i>Sevredol (MF)</i>		
. Nervous system	377	
. Palliative Care	480	
<i>Sical (AF)</i>		
. Alimentary tract and metabolism	95	
. Musculo-skeletal system	372	
<i>Sifrol (BY)</i>		
. Nervous system	398, 399	
<i>Sifrol ER (BY)</i>		
. Nervous system	399	
<i>Sigma Pharmaceuticals (Australia) Pty Ltd (SI)</i>		
. Antineoplastic and immunomodulating agents	199	
. Nervous system	387, 419	
. Palliative Care	482	
. Section 100	798	
<i>Sigmacort (SI)</i>		
. Dermatologicals	152	
. Dental	490	
<i>Sigmaxin (FM)</i>		
. Cardiovascular system	107	
<i>Sigmaxin-PG (FM)</i>		
. Cardiovascular system	107	
<i>SILDENAFIL CITRATE</i>		
. Section 100	548, 684	
. Repatriation Schedule	847	
<i>Silic 15 (EO)</i>		
. Repatriation Schedule	843	
<i>SILVER SULFADIAZINE</i>		
. Dermatologicals	151	
<i>Simplotan (GP)</i>		
. Anti-infectives for systemic use	190	
. Antiparasitic products, insecticides and repellents	430	

<i>Simponi (SH)</i>	
.Antineoplastic and immunomodulating agents	329, 332, 336, 338, 339, 342, 345
<i>Simvahexal (SZ)</i>	
.Cardiovascular system	141, 142, 143, 144
<i>Simvar 10 (SI)</i>	
.Cardiovascular system	141, 143
<i>Simvar 20 (SI)</i>	
.Cardiovascular system	141, 143
<i>Simvar 40 (SI)</i>	
.Cardiovascular system	141, 143
<i>Simvar 80 (SI)</i>	
.Cardiovascular system	142, 144
<i>SIMVASTATIN</i>	
.Cardiovascular system	140, 142
<i>Simvastatin generichealth (GQ)</i>	
.Cardiovascular system	141, 142, 143, 144
<i>Simvastatin Winthrop (WA)</i>	
.Cardiovascular system	141, 142, 143, 144
<i>Simvastatin-DP (GM)</i>	
.Cardiovascular system	141, 142, 143, 144
<i>Simvastatin-GA 10 (GN)</i>	
.Cardiovascular system	141, 143
<i>Simvastatin-GA 20 (GN)</i>	
.Cardiovascular system	141, 143
<i>Simvastatin-GA 40 (GN)</i>	
.Cardiovascular system	142, 143
<i>Simvastatin-GA 80 (GN)</i>	
.Cardiovascular system	142, 144
<i>Simvastatin-Spirit 10 (ZP)</i>	
.Cardiovascular system	141, 143
<i>Simvastatin-Spirit 20 (ZP)</i>	
.Cardiovascular system	141, 143
<i>Simvastatin-Spirit 40 (ZP)</i>	
.Cardiovascular system	142, 143
<i>Simvastatin-Spirit 80 (ZP)</i>	
.Cardiovascular system	142, 144
<i>Simvasyn (CR)</i>	
.Cardiovascular system	141, 142, 143, 144
<i>Sinemet (MK)</i>	
.Nervous system	396
<i>Sinemet 100/25 (MK)</i>	
.Nervous system	396
<i>Sinemet CR (MK)</i>	
.Nervous system	396
<i>Sinequan (PF)</i>	
.Nervous system	411
<i>Singulair (MK)</i>	
.Respiratory system	438
<i>SIROLIMUS</i>	
.Antineoplastic and immunomodulating agents	237
.Section 100	581, 719
<i>SITAGLIPTIN</i>	
.Alimentary tract and metabolism	93
<i>SITAGLIPTIN with METFORMIN</i>	
.Alimentary tract and metabolism	91
<i>SITAXENTAN SODIUM</i>	
.Section 100	552, 689
<i>Skelid (SW)</i>	
.Musculo-skeletal system	368
<i>SKIN CLEANSER</i>	
.Repatriation Schedule	846
<i>SKIN EMOLLIENT</i>	
.Repatriation Schedule	843
<i>Slow-K (NV)</i>	
.Alimentary tract and metabolism	96
<i>Sodibic (AS)</i>	
.Genito urinary system and sex hormones	166
<i>SODIUM ACID PHOSPHATE</i>	
.Various	453
<i>SODIUM ALGINATE with CALCIUM CARBONATE and SODIUM BICARBONATE</i>	
.Alimentary tract and metabolism	75
<i>SODIUM AUROTHIOMALATE</i>	
.Musculo-skeletal system	362
<i>SODIUM BICARBONATE</i>	
.Genito urinary system and sex hormones	166
<i>SODIUM CHLORIDE</i>	
.Blood and blood forming organs	105
.Various	467
.Dental	488, 513
.Repatriation Schedule	840
<i>Sodium Chloride 0.9% Freeflex (PK)</i>	
.Blood and blood forming organs	106
<i>SODIUM CHLORIDE COMPOUND</i>	
.Blood and blood forming organs	106
<i>SODIUM CHLORIDE with GLUCOSE</i>	
.Blood and blood forming organs	106
.Dental	488
<i>SODIUM CITRO-TARTRATE</i>	
.Repatriation Schedule	848
<i>SODIUM CLODRONATE TETRAHYDRATE</i>	
.Musculo-skeletal system	367
<i>SODIUM CROMOGLYCATE</i>	
.Respiratory system	437
.Sensory organs	444
.Optometrical	518
.Repatriation Schedule	857
<i>SODIUM LACTATE COMPOUND</i>	
.Blood and blood forming organs	106
<i>SODIUM POLYSTYRENE SULFONATE</i>	
.Repatriation Schedule	860
<i>SODIUM VALPROATE</i>	
.Nervous system	389
<i>Sodium Valproate Sandoz (SZ)</i>	
.Nervous system	389
<i>Soflax (GM)</i>	
.Repatriation Schedule	838
<i>Sofradex (SW)</i>	
.Sensory organs	451
<i>Soframycin (SW)</i>	
.Sensory organs	451
.Optometrical	521
<i>Solaraze 3% Gel (CS)</i>	
.Repatriation Schedule	846
<i>Solavert (SI)</i>	
.Cardiovascular system	108, 114, 115
<i>Solian 100 (SW)</i>	
.Nervous system	403
<i>Solian 200 (SW)</i>	

.Nervous system	403
<i>Solian 400 (SW)</i>	
.Nervous system	403
<i>Solian Solution (SW)</i>	
.Nervous system	403
<i>Solone (FM)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	170
<i>SoloSite Gel 36361338 (SN)</i>	
.Repatriation Schedule	872
<i>Solprin (RC)</i>	
.Blood and blood forming organs	98
.Nervous system	384
.Dental	508
<i>Solu-Cortef (PF)</i>	
.Emergency Drug Supplies	64
.Systemic hormonal preparations, excl. sex hormones and insulins	169
.Dental	491
<i>Solu-Medrol (PF)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	170
<i>Solugel 10336 (JJ)</i>	
.Repatriation Schedule	872
<i>Somac (NQ)</i>	
.Alimentary tract and metabolism	74, 75
<i>SOMATROPIN (Recombinant human growth hormone)</i>	
.Section 100	794
<i>Somatuline Autogel (IS)</i>	
.Section 100	558, 694
<i>Somatuline LA (IS)</i>	
.Section 100	558, 694
<i>Sone (FM)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	170
<i>SORAFENIB</i>	
.Antineoplastic and immunomodulating agents	224
<i>Sorbidin (AF)</i>	
.Cardiovascular system	109
<i>SORBITOL with SODIUM CITRATE and SODIUM LAURYL SULFOACETATE</i>	
.Alimentary tract and metabolism	81
.Palliative Care	473
.Repatriation Schedule	838
<i>Sorbsan 1410 (UM)</i>	
.Repatriation Schedule	866
<i>Sorbsan 1411 (UM)</i>	
.Repatriation Schedule	866
<i>Sotacor (FM)</i>	
.Cardiovascular system	108, 115
<i>SOTALOL HYDROCHLORIDE</i>	
.Cardiovascular system	108, 114
<i>Sotalol Sandoz (SZ)</i>	
.Cardiovascular system	108, 115
<i>SOY LECITHIN</i>	
.Sensory organs	450
.Optometrical	521
<i>SOY PROTEIN and FAT FORMULA with VITAMINS and MINERALS—CARBOHYDRATE FREE</i>	
.Various	466
<i>Sozol (SI)</i>	
.Alimentary tract and metabolism	74, 75
<i>Span-K (AS)</i>	
.Alimentary tract and metabolism	95
<i>Spiractin 100 (AF)</i>	
.Cardiovascular system	113
<i>Spiractin 25 (AF)</i>	
.Cardiovascular system	113
<i>Spiriva (BY)</i>	
.Respiratory system	437
<i>SPIRONOLACTONE</i>	
.Cardiovascular system	113
<i>Sporanox (JC)</i>	
.Antiinfectives for systemic use	192
<i>Sprycel (BQ)</i>	
.Antineoplastic and immunomodulating agents	211, 212, 213
<i>Stalevo 100/25/200mg (NV)</i>	
.Nervous system	397
<i>Stalevo 125/31.25/200mg (NV)</i>	
.Nervous system	397
<i>Stalevo 150/37.5/200mg (NV)</i>	
.Nervous system	397
<i>Stalevo 200/50/200mg (NV)</i>	
.Nervous system	397
<i>Stalevo 50/12.5/200mg (NV)</i>	
.Nervous system	396
<i>Stalevo 75/18.75/200mg (NV)</i>	
.Nervous system	397
<i>Staphylex 250 (AF)</i>	
.Antiinfectives for systemic use	180
.Dental	494
<i>Staphylex 500 (AF)</i>	
.Antiinfectives for systemic use	180
.Dental	494
<i>STAVUDINE</i>	
.Section 100	565, 701
<i>Stelara (JC)</i>	
.Antineoplastic and immunomodulating agents	350, 355
<i>Stelax 10 (SI)</i>	
.Musculo-skeletal system	363
<i>Stelax 25 (SI)</i>	
.Musculo-skeletal system	363
<i>Stelazine (GH)</i>	
.Nervous system	400
<i>Stemetil (SW)</i>	
.Emergency Drug Supplies	64
.Alimentary tract and metabolism	79
.Dental	486
<i>Stemzine (AV)</i>	
.Alimentary tract and metabolism	79
.Dental	486
<i>STERCULIA with FRANGULA BARK</i>	
.Alimentary tract and metabolism	80
.Palliative Care	471
.Repatriation Schedule	838
<i>Steripaste 3610 (XP)</i>	
.Repatriation Schedule	865
<i>Stieprox Liquid (GK)</i>	
.Repatriation Schedule	842

<i>Stocrin (MK)</i>		
.Section 100	566, 703	
<i>Strattera (LY)</i>		
.Nervous system	419	
<i>Stromectol (MK)</i>		
.Antiparasitic products, insecticides and repellents	432	
<i>STRONTIUM RANELATE</i>		
.Musculo-skeletal system	372	
<i>Suboxone (RC)</i>		
.Section 100	798	
<i>Subutex (RC)</i>		
.Section 100	798	
<i>SUCRALFATE</i>		
.Alimentary tract and metabolism	75	
<i>SULFACETAMIDE SODIUM</i>		
.Sensory organs	440	
.Optometrical	516	
<i>SULFASALAZINE</i>		
.Alimentary tract and metabolism	85	
<i>SULINDAC</i>		
.Musculo-skeletal system	358	
.Palliative Care	477	
.Dental	501	
<i>Sulprix (AF)</i>		
.Nervous system	403	
<i>SULTHIAME</i>		
.Nervous system	394	
<i>Sumagran 50 (SI)</i>		
.Nervous system	386	
<i>Sumatab (AF)</i>		
.Nervous system	386	
<i>SUMATRIPTAN</i>		
.Nervous system	386	
<i>SUNITINIB</i>		
.Antineoplastic and immunomodulating agents	225	
<i>SUNSCREENS</i>		
.Repatriation Schedule	843	
<i>SunSense Ultra SPF 30+ (EO)</i>		
.Repatriation Schedule	844	
<i>Surepress 650947 (CC)</i>		
.Repatriation Schedule	863	
<i>Surepress 650948 (CC)</i>		
.Repatriation Schedule	863	
<i>Surgam (SW)</i>		
.Musculo-skeletal system	361	
<i>Sustanon 250 (SH)</i>		
.Genito urinary system and sex hormones	159	
<i>Sutent (PF)</i>		
.Antineoplastic and immunomodulating agents .	225, 226	
<i>Symbicort Turbuhaler 100/6 (AP)</i>		
.Respiratory system	434	
<i>Symbicort Turbuhaler 200/6 (AP)</i>		
.Respiratory system	434	
<i>Symbicort Turbuhaler 400/12 (AP)</i>		
.Respiratory system	435	
<i>Symmetrel 100 (NV)</i>		
.Nervous system	397	
<i>Synacthen Depot 1 mg/1 mL (NV)</i>		
.Systemic hormonal preparations, excl. sex hormones and insulins	167	
<i>Synarel (PF)</i>		
.Systemic hormonal preparations, excl. sex hormones and insulins	168	
<i>Synphasic (PF)</i>		
.Genito urinary system and sex hormones	158	
<i>Systane (AQ)</i>		
.Sensory organs	449, 450	
.Optometrical	520	
T		
<i>Tacidine (AF)</i>		
.Alimentary tract and metabolism	69	
<i>TACROLIMUS</i>		
.Antineoplastic and immunomodulating agents	356	
.Section 100	640, 777	
<i>Tacrolimus Sandoz (SZ)</i>		
.Antineoplastic and immunomodulating agents	356	
.Section 100	640, 777	
<i>TADALAFIL</i>		
.Repatriation Schedule	848	
<i>Talam (SI)</i>		
.Nervous system	412	
<i>Tamate (AF)</i>		
.Nervous system	394, 395	
<i>Tambocor (IA)</i>		
.Cardiovascular system	107	
<i>Tamosin (SI)</i>		
.Antineoplastic and immunomodulating agents	231	
<i>Tamoxen 20 mg (GM)</i>		
.Antineoplastic and immunomodulating agents	231	
<i>TAMOXIFEN CITRATE</i>		
.Antineoplastic and immunomodulating agents	231	
<i>Tamoxifen Sandoz (SZ)</i>		
.Antineoplastic and immunomodulating agents	231	
<i>Tamsil (SI)</i>		
.Dermatologicals	150	
.Repatriation Schedule	843	
<i>TAMSULOSIN HYDROCHLORIDE</i>		
.Repatriation Schedule	848	
<i>TAPES—NON-WOVEN RETENTION (POLYACRYLATE)</i>		
.Repatriation Schedule	874	
<i>TAPES—PLASTER ADHESIVE (WITH SILICONE)</i>		
.Repatriation Schedule	875	
<i>TAPES—PLASTER ADHESIVE ELASTIC</i>		
.Repatriation Schedule	875	
<i>TAPES—PLASTER ADHESIVE HYPOALLERGENIC</i>		
.Repatriation Schedule	875	
<i>Tarceva (RO)</i>		
.Antineoplastic and immunomodulating agents	213	
<i>Tarka 2/180 (AB)</i>		
.Cardiovascular system	130	
<i>Tarka 4/240 (AB)</i>		
.Cardiovascular system	130	
<i>Tasigna (NV)</i>		
.Antineoplastic and immunomodulating agents .	223, 224	
<i>Taxol (BQ)</i>		
.Antineoplastic and immunomodulating agents .	205, 206	
<i>Taxotere (SW)</i>		
.Antineoplastic and immunomodulating agents .	204, 205	
<i>Tazac (AS)</i>		

.Alimentary tract and metabolism.....	69	TEMAZEPAM	
<i>Tears Naturale (AQ)</i>		.Nervous system	410
.Sensory organs	448	.Palliative Care	485
.Optometrical	520	.Dental	510
<i>tearsagain (RB)</i>		Temodal (SH)	
.Sensory organs	450	.Antineoplastic and immunomodulating agents	200
.Optometrical	521	TEMOZOLOMIDE	
<i>Tecan (WQ)</i>		.Antineoplastic and immunomodulating agents	200
.Antineoplastic and immunomodulating agents	229	Temtabs (FM)	
<i>Tegaderm Transparent 1628 (MM)</i>		.Nervous system	410
.Repatriation Schedule	867	.Palliative Care	485
<i>Tegaderm Transparent Island 3582 (MM)</i>		.Dental	510
.Repatriation Schedule	867	TenderWet 24 Active (HR)	
<i>Tegaderm Transparent Island 3586 (MM)</i>		.Repatriation Schedule	869
.Repatriation Schedule	867	TenderWet Active Cavity (HR)	
<i>Tegretol 100 (NV)</i>		.Repatriation Schedule	869
.Nervous system	389	TENECTEPLASE	
.Dental	509	.Blood and blood forming organs	101
<i>Tegretol 200 (NV)</i>		TENOFOVIR	
.Nervous system	388	.Section 100	565, 702
.Dental	509	TENOFOVIR with EMTRICITABINE	
<i>Tegretol CR 200 (NV)</i>		.Section 100	567, 704
.Nervous system	389	TENOFOVIR with EMTRICITABINE and EFAVIRENZ	
.Dental	509	.Section 100	568, 704
<i>Tegretol CR 400 (NV)</i>		Tenopt (SI)	
.Nervous system	389	.Sensory organs.....	443
.Dental	509	.Optometrical.....	517
<i>Tegretol Liquid (NV)</i>		Tenormin (AP)	
.Nervous system	389	.Cardiovascular system	115
.Dental	509	Tensig (SI)	
TELBIVUDINE		.Cardiovascular system	115
.Section 100	565, 702	Tensocrepe 36300501 (BV)	
<i>Telfa 1970C (KE)</i>		.Repatriation Schedule	865
.Repatriation Schedule	873	Tensocrepe 36301001 (BV)	
<i>Telfa 2140C (KE)</i>		.Repatriation Schedule	865
.Repatriation Schedule	873	Tensocrepe 36307501 (BV)	
<i>Telfa 7650C (KE)</i>		.Repatriation Schedule	865
.Repatriation Schedule	873	Tensopress 66004347 (BV)	
<i>Telfa 8252F (KE)</i>		.Repatriation Schedule	863
.Repatriation Schedule	865	Tensopress 66004348 (BV)	
<i>Telfa 8253F (KE)</i>		.Repatriation Schedule	863
.Repatriation Schedule	865	TERAZOSIN HYDROCHLORIDE	
<i>Telfa 8254F (KE)</i>		.Repatriation Schedule	848
.Repatriation Schedule	865	Terbihexal (SZ)	
<i>Telfast (SW)</i>		.Dermatologicals	150
.Repatriation Schedule	858	.Repatriation Schedule	843
<i>Telfast 120 (SW)</i>		TERBINAFINE	
.Repatriation Schedule	858	.Dermatologicals	149, 150
TELMISARTAN		.Repatriation Schedule	842, 843
.Cardiovascular system	131	Terbinafine 250 (CR)	
TELMISARTAN with HYDROCHLOROTHIAZIDE		.Dermatologicals	150
.Cardiovascular system	132	Terbinafine-DP (GN)	
<i>Telzir (GK)</i>		.Repatriation Schedule	843
.Section 100	562, 699	Terbinafine-DRLA (RZ)	
<i>Telzir (VI)</i>		.Dermatologicals	150
.Section 100	562, 699	Terbinafine-GA (GM)	
<i>Temaze (AF)</i>		.Dermatologicals	150
.Nervous system	410	Terbix 250 (MI)	
.Palliative Care	485	.Dermatologicals	150
.Dental	510	TERBUTALINE SULFATE	

.Respiratory system	434, 438	.Cardiovascular system	121
<i>Teril (AF)</i>		<i>Terry White Chemists Diltiazem CD (TW)</i>	
.Nervous system	388	.Cardiovascular system	121
.Dental	509	<i>Terry White Chemists Doxycycline (TW)</i>	
TERIPARATIDE		.Antinfectives for systemic use	175, 176
.Systemic hormonal preparations, excl. sex hormones		.Dental	492
and insulins	172	<i>Terry White Chemists Enalapril (TW)</i>	
.Musculo-skeletal system	372	.Cardiovascular system	122, 123
<i>Terry White Chemists Aciclovir (TW)</i>		<i>Terry White Chemists Escitalopram (TW)</i>	
.Antinfectives for systemic use	195	.Nervous system	413
<i>Terry White Chemists Alendronate 70mg (TW)</i>		<i>Terry White Chemists Famotidine (TW)</i>	
.Musculo-skeletal system	365	.Alimentary tract and metabolism	69
<i>Terry White Chemists Allopurinol (TW)</i>		<i>Terry White Chemists Fluoxetine (TW)</i>	
.Musculo-skeletal system	364	.Nervous system	414
<i>Terry White Chemists Alprazolam (TW)</i>		<i>Terry White Chemists Frusemide (TW)</i>	
.Nervous system	408	.Cardiovascular system	112
<i>Terry White Chemists Amiodarone (TW)</i>		<i>Terry White Chemists Gemfibrozil (TW)</i>	
.Cardiovascular system	108	.Cardiovascular system	145
<i>Terry White Chemists Amlodipine (TW)</i>		<i>Terry White Chemists Gliclazide (TW)</i>	
.Cardiovascular system	118, 119	.Alimentary tract and metabolism	89
<i>Terry White Chemists Amoxycillin (TW)</i>		<i>Terry White Chemists Gliclazide MR (TW)</i>	
.Antinfectives for systemic use	177, 178	.Alimentary tract and metabolism	90
.Dental	492, 493	<i>Terry White Chemists Indapamide (TW)</i>	
<i>Terry White Chemists Amoxycillin and Clavulanic Acid (TW)</i>		.Cardiovascular system	112
.Antinfectives for systemic use	180	<i>Terry White Chemists Isosorbide Mononitrate (TW)</i>	
.Dental	495	.Cardiovascular system	110
<i>Terry White Chemists Atenolol (TW)</i>		<i>Terry White Chemists Lercanidipine (TW)</i>	
.Cardiovascular system	115	.Cardiovascular system	119
<i>Terry White Chemists Baclofen (TW)</i>		<i>Terry White Chemists Levetiracetam (TW)</i>	
.Musculo-skeletal system	363	.Nervous system	393
<i>Terry White Chemists Carvedilol 12.5 mg (TW)</i>		<i>Terry White Chemists Lisinopril (TW)</i>	
.Cardiovascular system	117	.Cardiovascular system	123, 124
<i>Terry White Chemists Carvedilol 25 mg (TW)</i>		<i>Terry White Chemists Meloxicam 15 mg (TW)</i>	
.Cardiovascular system	118	.Musculo-skeletal system	359
<i>Terry White Chemists Carvedilol 3.125 mg (TW)</i>		<i>Terry White Chemists Meloxicam 7.5 mg (TW)</i>	
.Cardiovascular system	117	.Musculo-skeletal system	359
<i>Terry White Chemists Carvedilol 6.25 mg (TW)</i>		<i>Terry White Chemists Metformin (TW)</i>	
.Cardiovascular system	117	.Alimentary tract and metabolism	88, 89
<i>Terry White Chemists Cefaclor (TW)</i>		<i>Terry White Chemists Metoprolol (TW)</i>	
.Antinfectives for systemic use	183	.Cardiovascular system	116
.Dental	497	<i>Terry White Chemists Mirtazapine (TW)</i>	
<i>Terry White Chemists Cefaclor CD (TW)</i>		.Nervous system	418
.Antinfectives for systemic use	183	<i>Terry White Chemists Moclobemide (TW)</i>	
.Dental	497	.Nervous system	416, 417
<i>Terry White Chemists Cephalexin (TW)</i>		<i>Terry White Chemists Norfloxacin (TW)</i>	
.Antinfectives for systemic use	181, 182	.Antinfectives for systemic use	188
.Dental	496	<i>Terry White Chemists Omeprazole (TW)</i>	
<i>Terry White Chemists Citalopram (TW)</i>		.Alimentary tract and metabolism	72, 73
.Nervous system	412	<i>Terry White Chemists Pantoprazole (TW)</i>	
<i>Terry White Chemists Clarithromycin (TW)</i>		.Alimentary tract and metabolism	74, 75
.Antinfectives for systemic use	185	<i>Terry White Chemists Paracetamol (TW)</i>	
<i>Terry White Chemists Clomipramine (TW)</i>		.Nervous system	384
.Nervous system	409, 411	.Dental	508, 509
<i>Terry White Chemists Clopidogrel (TW)</i>		<i>Terry White Chemists Paroxetine (TW)</i>	
.Blood and blood forming organs	99	.Nervous system	415
<i>Terry White Chemists Diclofenac (TW)</i>		<i>Terry White Chemists Perindopril (TW)</i>	
.Musculo-skeletal system	358	.Cardiovascular system	124, 125
.Palliative Care	475, 476	<i>Terry White Chemists Perindopril/ Indapamide 4/1.25 (TW)</i>	
.Dental	501	.Cardiovascular system	129
<i>Terry White Chemists Diltiazem (TW)</i>		<i>Terry White Chemists Piroxicam (TW)</i>	

.Musculo-skeletal system	360
.Dental	502
<i>Terry White Chemists Pravastatin (TW)</i>	
.Cardiovascular system	138, 139, 140
<i>Terry White Chemists Prazosin (TW)</i>	
.Cardiovascular system	111
<i>Terry White Chemists Ramipril (TW)</i>	
.Cardiovascular system	126, 127
<i>Terry White Chemists Ranitidine (TW)</i>	
.Alimentary tract and metabolism	69, 70
<i>Terry White Chemists Roxithromycin (TW)</i>	
.Antifectives for systemic use	186
.Dental	498
<i>Terry White Chemists Sertraline (TW)</i>	
.Nervous system	415
<i>Terry White Chemists Simvastatin (TW)</i>	
.Cardiovascular system	141, 142, 143, 144
<i>Terry White Chemists Sotalol (TW)</i>	
.Cardiovascular system	108, 115
<i>Terry White Chemists Sumatriptan (TW)</i>	
.Nervous system	386
<i>Terry White Chemists Tramadol (TW)</i>	
.Nervous system	382
.Dental	507
<i>Terry White Chemists Tramadol SR (TW)</i>	
.Nervous system	383
.Dental	507, 508
<i>Tertroxin (SI)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	171
<i>Testogel (SC)</i>	
.Genito urinary system and sex hormones	159
TESTOSTERONE	
.Genito urinary system and sex hormones	158
TESTOSTERONE ENANTHATE	
.Genito urinary system and sex hormones	159
TESTOSTERONE ESTERS	
.Genito urinary system and sex hormones	159
TESTOSTERONE UNDECANOATE	
.Genito urinary system and sex hormones	159
TETRABENAZINE	
.Nervous system	429
TETRACOSACTRIN	
.Systemic hormonal preparations, excl. sex hormones and insulins	167
<i>Teveten (SM)</i>	
.Cardiovascular system	130
<i>Teveten Plus 600/12.5 (SM)</i>	
.Cardiovascular system	131
THALIDOMIDE	
.Section 100	647, 784
<i>Thalomid (CJ)</i>	
.Section 100	647, 784
<i>Thelin (PF)</i>	
.Section 100	557, 693
THEOPHYLLINE	
.Respiratory system	438
<i>TheraTears (CX)</i>	
.Sensory organs	447
.Optometrical	519

THIAMINE HYDROCHLORIDE	
.Alimentary tract and metabolism	95
.Repatriation Schedule	839
THIOGUANINE	
.Antineoplastic and immunomodulating agents	202
<i>Thioprine (AF)</i>	
.Antineoplastic and immunomodulating agents	357
THIOTEPA	
.Antineoplastic and immunomodulating agents	199
<i>Thyrogen (GZ)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	167
THYROTROPIN ALFA	
.Systemic hormonal preparations, excl. sex hormones and insulins	167
THYROXINE SODIUM	
.Systemic hormonal preparations, excl. sex hormones and insulins	171
TIAGABINE HYDROCHLORIDE	
.Nervous system	389
TIAPROFENIC ACID	
.Musculo-skeletal system	361
TICARCILLIN with CLAVULANIC ACID	
.Antifectives for systemic use	181
.Dental	495
TICLOPIDINE HYDROCHLORIDE	
.Blood and blood forming organs	100
<i>Tielle MT2442 (JJ)</i>	
.Repatriation Schedule	869
<i>Tielle MTL101E (JJ)</i>	
.Repatriation Schedule	869
<i>Tilade CFC-Free (SW)</i>	
.Respiratory system	437
<i>Tilodene (AF)</i>	
.Blood and blood forming organs	101
TILUDRONATE DISODIUM	
.Musculo-skeletal system	368
<i>Timentin (GK)</i>	
.Antifectives for systemic use	181
.Dental	495
TIMOLOL MALEATE	
.Sensory organs	443
.Optometrical	517
<i>Timoptol (FR)</i>	
.Sensory organs	443
.Optometrical	517
<i>Timoptol XE (MK)</i>	
.Sensory organs	443
.Optometrical	517
<i>Tinaderm (SH)</i>	
.Repatriation Schedule	842
<i>Tinexa (SI)</i>	
.Genito urinary system and sex hormones	157
TINIDAZOLE	
.Antifectives for systemic use	190
.Antiparasitic products, insecticides and repellents	430
TIOTROPIUM BROMIDE MONOHYDRATE	
.Respiratory system	437
TIPRANAVIR	
.Section 100	563, 699

<i>TIROFIBAN HYDROCHLORIDE</i>		
.Blood and blood forming organs	101	
<i>Titralac (MM)</i>		
.Repatriation Schedule	837	
<i>Tobra-Day (PL)</i>		
.Antifungives for systemic use	187	
<i>TOBRAMYCIN</i>		
.Sensory organs	440	
<i>TOBRAMYCIN SULFATE</i>		
.Antifungives for systemic use	187	
<i>Tobrex (AQ)</i>		
.Sensory organs	440	
<i>TOCILIZUMAB</i>		
.Section 100	634, 771	
<i>Tofranil 10 (LM)</i>		
.Nervous system	412	
<i>Tofranil 25 (LM)</i>		
.Nervous system	412	
<i>Toleraide 10 (PQ)</i>		
.Nervous system	412	
<i>Toleraide 25 (PQ)</i>		
.Nervous system	412	
<i>TOLNAFTATE</i>		
.Repatriation Schedule	842	
<i>Tolvon (SH)</i>		
.Nervous system	417	
<i>Tomudex (HH)</i>		
.Antineoplastic and immunomodulating agents	201	
<i>Topamax (JC)</i>		
.Nervous system	394, 395	
<i>Topamax Sprinkle (JC)</i>		
.Nervous system	395	
<i>TOPIRAMATE</i>		
.Nervous system	394, 395	
<i>Topiramate Sandoz (SZ)</i>		
.Nervous system	394, 395	
<i>Topiramate-GA (GM)</i>		
.Nervous system	394, 395	
<i>TOPOTECAN HYDROCHLORIDE</i>		
.Antineoplastic and immunomodulating agents	229	
<i>Toprol-XL 190 (AP)</i>		
.Cardiovascular system	116	
<i>Toprol-XL 23.75 (AP)</i>		
.Cardiovascular system	115	
<i>Toprol-XL 47.5 (AP)</i>		
.Cardiovascular system	115	
<i>Toprol-XL 95 (AP)</i>		
.Cardiovascular system	116	
<i>TOREMIFENE CITRATE</i>		
.Antineoplastic and immunomodulating agents	232	
<i>Tracleer (AT)</i>		
.Section 100	538, 670	
<i>TRAMADOL HYDROCHLORIDE</i>		
.Nervous system	382, 383, 384	
.Dental	507, 508	
<i>Tramadol Sandoz (SZ)</i>		
.Nervous system	382	
.Dental	507	
<i>Tramadol Sandoz SR (SZ)</i>		
.Nervous system	383	
.Dental	508	
<i>Tramahexal (SZ)</i>		
.Emergency Drug Supplies	65	
.Nervous system	384	
.Dental	508	
<i>Tramahexal SR (SZ)</i>		
.Nervous system	383	
.Dental	507, 508	
<i>Tramal (CS)</i>		
.Nervous system	382, 383	
.Dental	507	
<i>Tramal 100 (CS)</i>		
.Emergency Drug Supplies	65	
.Nervous system	384	
.Dental	508	
<i>Tramal SR 100 (CS)</i>		
.Nervous system	383	
.Dental	507	
<i>Tramal SR 150 (CS)</i>		
.Nervous system	383	
.Dental	508	
<i>Tramal SR 200 (CS)</i>		
.Nervous system	383	
.Dental	508	
<i>Tramal SR 50 (CS)</i>		
.Nervous system	383	
.Dental	507	
<i>Tramedo (AF)</i>		
.Nervous system	382	
.Dental	507	
<i>Tramedo SR 100 (AF)</i>		
.Nervous system	383	
.Dental	507	
<i>Tramedo SR 150 (AF)</i>		
.Nervous system	383	
.Dental	508	
<i>Tramedo SR 200 (AF)</i>		
.Nervous system	383	
.Dental	508	
<i>Tranalpha (AF)</i>		
.Cardiovascular system	128	
<i>Trandate (SI)</i>		
.Cardiovascular system	118	
<i>TRANDOLAPRIL</i>		
.Cardiovascular system	128	
<i>Trandolapril generichealth (GQ)</i>		
.Cardiovascular system	128	
<i>TRANDOLAPRIL with VERAPAMIL HYDROCHLORIDE</i>		
.Cardiovascular system	130	
<i>Trandolapril-DP (GN)</i>		
.Cardiovascular system	128	
<i>TRANEXAMIC ACID</i>		
.Blood and blood forming organs	104	
<i>Transiderm-Nitro 25 (NV)</i>		
.Cardiovascular system	109	
<i>Transiderm-Nitro 50 (NV)</i>		
.Cardiovascular system	109	
<i>TRANLYCPROMINE SULFATE</i>		
.Nervous system	416	
<i>TRASTUZUMAB</i>		

.Section 100	799, 800	.Cardiovascular system	126
<i>Travatan (AQ)</i>		<i>Tritace 5 mg (SW)</i>	
.Sensory organs	443	.Cardiovascular system	127
.Optometrical	518	<i>Tritace Titration Pack (SW)</i>	
<i>TRAVOPROST</i>		.Cardiovascular system	128
.Sensory organs	443	<i>Trizivir (VI)</i>	
.Optometrical	518	.Section 100	567, 703
<i>TRAVOPROST with TIMOLOL MALEATE</i>		<i>TROPISETRON HYDROCHLORIDE</i>	
.Sensory organs	443	.Alimentary tract and metabolism	78
.Optometrical	518	<i>TrueTrack (NX)</i>	
<i>TRIAMCINOLONE ACETONIDE</i>		.Various	456
.Dermatologicals	152	<i>Trusopt (MK)</i>	
.Systemic hormonal preparations, excl. sex hormones and insulins	170	.Sensory organs	442
.Dental	491	.Optometrical	517
<i>TRIAMCINOLONE ACETONIDE with NEOMYCIN SULFATE, GRAMICIDIN and NYSTATIN</i>		<i>Truvada (GI)</i>	
.Sensory organs	451	.Section 100	567, 704
.Repatriation Schedule	845	<i>Tryzan Caps 1.25 (AF)</i>	
<i>Triasyn 2.5/2.5 (SW)</i>		.Cardiovascular system	126
.Cardiovascular system	130	<i>Tryzan Caps 10 (AF)</i>	
<i>Triasyn 5.0/5.0 (SW)</i>		.Cardiovascular system	127
.Cardiovascular system	130	<i>Tryzan Caps 2.5 (AF)</i>	
<i>Tricortone (FM)</i>		.Cardiovascular system	127
.Dermatologicals	152	<i>Tryzan Caps 5 (AF)</i>	
<i>Trifeme 28 (WX)</i>		.Cardiovascular system	127
.Genito urinary system and sex hormones	158	<i>Tryzan Tabs 1.25 (AF)</i>	
<i>TRIFLUOPERAZINE HYDROCHLORIDE</i>		.Cardiovascular system	126
.Nervous system	400	<i>Tryzan Tabs 10 (AF)</i>	
<i>TRIGLYCERIDES, LONG CHAIN with GLUCOSE POLYMER</i>		.Cardiovascular system	127
.Various	466	<i>Tryzan Tabs 2.5 (AF)</i>	
<i>TRIGLYCERIDES, MEDIUM CHAIN</i>		.Cardiovascular system	126
.Various	457	<i>Tryzan Tabs 5 (AF)</i>	
<i>TRIGLYCERIDES, MEDIUM CHAIN and LONG CHAIN with GLUCOSE POLYMER</i>		.Cardiovascular system	127
.Various	466	<i>Tubegauz 0501633 (SS)</i>	
<i>TRIGLYCERIDES—MEDIUM CHAIN, FORMULA</i>		.Repatriation Schedule	865
.Various	460, 466	<i>Tubegauz 0501658 (SS)</i>	
<i>Trileptal (NV)</i>		.Repatriation Schedule	865
.Nervous system	389	<i>Tubifast 2434 (SS)</i>	
<i>TRIMETHOPRIM</i>		.Repatriation Schedule	865
.Antifungives for systemic use	184	<i>Tubifast 2436 (SS)</i>	
<i>TRIMETHOPRIM with SULFAMETHOXAZOLE</i>		.Repatriation Schedule	865
.Antifungives for systemic use	184	<i>Tubifast 2438 (SS)</i>	
.Dental	497	.Repatriation Schedule	865
<i>Triphasil 28 (WY)</i>		<i>Tubigrip 1479 (SS)</i>	
.Genito urinary system and sex hormones	158	.Repatriation Schedule	865
<i>Triprim (SI)</i>		<i>Tubigrip 1480 (SS)</i>	
.Antifungives for systemic use	184	.Repatriation Schedule	865
<i>TRIPTORELIN</i>		<i>Tubigrip 1481 (SS)</i>	
.Antineoplastic and immunomodulating agents	231	.Repatriation Schedule	865
<i>Triquilar ED (SC)</i>		<i>Tubigrip 1482 (SS)</i>	
.Genito urinary system and sex hormones	158	.Repatriation Schedule	865
<i>Tritace (SW)</i>		<i>Tubigrip 1483 (SS)</i>	
.Cardiovascular system	127	.Repatriation Schedule	865
<i>Tritace 1.25 mg (SW)</i>		<i>Tubigrip 1484 (SS)</i>	
.Cardiovascular system	126	.Repatriation Schedule	865
<i>Tritace 10 mg (SW)</i>		<i>Tubigrip 1486 (SS)</i>	
.Cardiovascular system	127	.Repatriation Schedule	865
<i>Tritace 2.5 mg (SW)</i>		<i>Tubigrip B 1520 (SS)</i>	
		.Repatriation Schedule	865
		<i>Tubigrip C 1545 (SS)</i>	
		.Repatriation Schedule	865

<i>Tubigrip D 1546 (SS)</i>	
.Repatriation Schedule	865
<i>Tubigrip E 1547 (SS)</i>	
.Repatriation Schedule	865
<i>Tubigrip F 1548 (SS)</i>	
.Repatriation Schedule	865
<i>Tykerb (GK)</i>	
.Antineoplastic and immunomodulating agents	222
<i>TYR Anamix infant (SB)</i>	
.Various.....	464
<i>TYR Anamix Junior (SB)</i>	
.Various.....	464
<i>TYR Anamix junior LQ (SB)</i>	
.Various.....	462
<i>TYR Cooler (VF)</i>	
.Various.....	464
<i>TYR Express (VF)</i>	
.Various.....	464
<i>TYR gel (VF)</i>	
.Various.....	464
<i>Tyrosine Amino Acid Supplement (VF)</i>	
.Various.....	467
<i>TYROSINE with CARBOHYDRATE</i>	
.Various.....	467
<i>Tysabri (BD)</i>	
.Section 100.....	581, 719

U

<i>Ulcaid (RA)</i>	
.Alimentary tract and metabolism.....	69, 70
<i>Ulcylte (AF)</i>	
.Alimentary tract and metabolism.....	75
<i>Uracol (GM)</i>	
.Repatriation Schedule	848
<i>Ural Sachets (SI)</i>	
.Repatriation Schedule	848
<i>UREA</i>	
.Repatriation Schedule	843
<i>Urederm (HA)</i>	
.Repatriation Schedule	843
<i>Uremide (AF)</i>	
.Cardiovascular system.....	112
<i>Urex (FM)</i>	
.Cardiovascular system.....	112
<i>Urex-Forte (FM)</i>	
.Cardiovascular system.....	112
<i>Urex-M (FM)</i>	
.Cardiovascular system.....	112
<i>Uro-Carb (HA)</i>	
.Nervous system	427
<i>Uromitexan (BX)</i>	
.Various.....	453
<i>URSODEOXYCHOLIC ACID</i>	
.Alimentary tract and metabolism.....	79
<i>Ursafalk (OA)</i>	
.Alimentary tract and metabolism.....	79
<i>USTEKINUMAB</i>	
.Antineoplastic and immunomodulating agents .	345, 350

V

<i>Vagifem (NO)</i>	
.Genito urinary system and sex hormones	160
<i>VALACICLOVIR HYDROCHLORIDE</i>	
.Antiinfectives for systemic use	197
.Section 100	561, 698
<i>Valcyte (RO)</i>	
.Section 100	562, 698
<i>VALGANCICLOVIR HYDROCHLORIDE</i>	
.Section 100	562, 698
<i>Valine 1000 Amino Acid Supplement (VF)</i>	
.Various.....	467
<i>Valine Amino Acid Supplement (VF)</i>	
.Various.....	467
<i>VALINE with CARBOHYDRATE</i>	
.Various.....	467
<i>Valium (RO)</i>	
.Nervous system	408
.Palliative Care	483, 484
.Dental	509
<i>Valpam 2 (SI)</i>	
.Nervous system	408
.Palliative Care	483, 484
.Dental	509
<i>Valpam 5 (SI)</i>	
.Nervous system	408
.Palliative Care	483, 484
.Dental	509
<i>Valprease 200 (SI)</i>	
.Nervous system	389
<i>Valprease 500 (SI)</i>	
.Nervous system	389
<i>Valpro 200 (AF)</i>	
.Nervous system	389
<i>Valpro 500 (AF)</i>	
.Nervous system	389
<i>Valproate Winthrop EC 200 (WA)</i>	
.Nervous system	389
<i>Valproate Winthrop EC 500 (WA)</i>	
.Nervous system	389
<i>VALSARTAN</i>	
.Cardiovascular system.....	131
<i>VALSARTAN with HYDROCHLOROTHIAZIDE</i>	
.Cardiovascular system	132
<i>Valtrex (GK)</i>	
.Antiinfectives for systemic use	197
.Section 100	561, 698
<i>Vancocin (AS)</i>	
.Alimentary tract and metabolism.....	82
<i>Vancocin CP (AS)</i>	
.Antiinfectives for systemic use	189
.Dental	499
<i>VANCOMYCIN</i>	
.Alimentary tract and metabolism.....	82
.Antiinfectives for systemic use	188, 189
.Dental	499
<i>Vancomycin Sandoz (SZ)</i>	
.Antiinfectives for systemic use	188, 189
.Dental	499
<i>VARENICLINE</i>	
.Nervous system	428

<i>Vasocardol (AV)</i>		.Nervous system	391
.Cardiovascular system	121		
<i>Vasocardol CD (AV)</i>		<i>VINBLASTINE SULFATE</i>	
.Cardiovascular system	121	.Antineoplastic and immunomodulating agents	203
<i>Vastin (NM)</i>		<i>VINCISTINE SULFATE</i>	
.Cardiovascular system	137	.Antineoplastic and immunomodulating agents	203
<i>Vastoran (RA)</i>		<i>Vinorelbine Ebewe (IT)</i>	
.Cardiovascular system	138, 139, 140	.Antineoplastic and immunomodulating agents	204
<i>Vedilol 12.5 (SI)</i>		<i>Vinorelbine Link (PK)</i>	
.Cardiovascular system	117	.Antineoplastic and immunomodulating agents	204
<i>Vedilol 25 (SI)</i>		<i>VINORELBINE TARTRATE</i>	
.Cardiovascular system	118	.Antineoplastic and immunomodulating agents .	203, 204
<i>Vedilol 3.125 (SI)</i>		<i>Viramune (BY)</i>	
.Cardiovascular system	117	.Section 100	567, 703
<i>Vedilol 6.25 (SI)</i>		<i>Viread (GI)</i>	
.Cardiovascular system	117	.Section 100	566, 702
<i>Velcade (JC)</i>		<i>Viscopaste 4948 (SN)</i>	
.Antineoplastic and immunomodulating agents .	228, 229	.Repatriation Schedule	866
<i>VENLAFAXINE HYDROCHLORIDE</i>		<i>Viscotears (NV)</i>	
.Nervous system	418	.Sensory organs.....	446
<i>Venofer (AS)</i>		.Optometrical.....	518
.Blood and blood forming organs	104	<i>Viscotears Gel PF (NV)</i>	
<i>Ventavis (SC)</i>		.Sensory organs.....	446
.Section 100	548, 684	.Optometrical.....	519
<i>Ventolin (GK)</i>		<i>Visken 15 (NV)</i>	
.Respiratory system	437	.Cardiovascular system	114
<i>Ventolin CFC-free (GK)</i>		<i>Visken 5 (NV)</i>	
.Emergency Drug Supplies	65	.Cardiovascular system	114
.Respiratory system	433	<i>Vistide (GI)</i>	
<i>Ventolin Nebules (GK)</i>		.Section 100	561, 697
.Emergency Drug Supplies	65	<i>Vistil (AE)</i>	
.Respiratory system	434	.Sensory organs.....	450
<i>Ventolin Rotacaps (GK)</i>		.Optometrical.....	521
.Respiratory system	433	<i>Vistil Forte (AE)</i>	
<i>Vepesid (BQ)</i>		.Sensory organs.....	450
.Antineoplastic and immunomodulating agents	204	.Optometrical.....	521
<i>Veracaps SR (SI)</i>		<i>Visudyne (NV)</i>	
.Cardiovascular system	120	.Sensory organs.....	445
<i>VERAPAMIL HYDROCHLORIDE</i>		<i>VITAMIN B GROUP COMPLEX</i>	
.Cardiovascular system	120	.Repatriation Schedule	839
<i>Vermox (BI)</i>		<i>VITAMINS, MINERALS and TRACE ELEMENTS with</i>	
.Repatriation Schedule	856	<i>CARBOHYDRATE</i>	
<i>VERTEPORFIN</i>		.Various.....	467
.Sensory organs	444	<i>Vitrasert (BU)</i>	
<i>Vfend (PF)</i>		.Section 100	653, 790
.Antifungives for systemic use	193	<i>Volibris (GK)</i>	
<i>Viagra (PF)</i>		.Section 100	531, 663
.Repatriation Schedule	848	<i>Voltaren 100 (NV)</i>	
<i>Vibra-Tabs (PF)</i>		.Musculo-skeletal system	358
.Antifungives for systemic use	175	.Palliative Care	475, 476
<i>Vibramycin (PF)</i>		.Dental	501
.Antifungives for systemic use	175, 176	<i>Voltaren 25 (NV)</i>	
.Dental	492	.Musculo-skeletal system	358
<i>Videx EC (BQ)</i>		.Palliative Care	475, 476
.Section 100	564, 700	.Dental	501
<i>VIGABATRIN</i>		<i>Voltaren 50 (NV)</i>	
.Nervous system	390	.Musculo-skeletal system	358
<i>VILDAGLIPTIN</i>		.Palliative Care	475, 476
.Alimentary tract and metabolism.....	93	.Dental	501
<i>Vimpat (UC)</i>		<i>Voluven 6% (PK)</i>	
		.Blood and blood forming organs	105

VORICONAZOLE

.Antifungives for systemic use 192, 193

Voxam (SZ)

.Nervous system 414

Vytorin (MK)

.Cardiovascular system 147, 148

W**WARFARIN SODIUM**

.Blood and blood forming organs 97

Wartec Cream (GK)

.Repatriation Schedule 844

WaveSense Jazz (HE)

.Various 456

Waxsol (NE)

.Repatriation Schedule 859

Wellvone (GK)

.Antiparasitic products, insecticides and repellents 430

WHEY PROTEIN FORMULA supplemented with AMINO ACIDS, LONG CHAIN POLYUNSATURATED FATTY ACIDS, VITAMINS and MINERALS, and low in PROTEIN, PHOSPHATE, POTASSIUM and LACTOSE

.Various 467

WHEY PROTEIN FORMULA supplemented with AMINO ACIDS, VITAMINS and MINERALS, and low in PROTEIN, PHOSPHATE, POTASSIUM and LACTOSE

.Various 467

Winthrop Oxaliplatin (WA)

.Antineoplastic and immunomodulating agents 208

WOOL ALCOHOLS

.Repatriation Schedule 843

X**Xalacom (PF)**

.Sensory organs 443

.Optometrical 518

Xalatan (PF)

.Sensory organs 443

.Optometrical 518

Xalox (WQ)

.Antineoplastic and immunomodulating agents 208

Xanax (PF)

.Nervous system 408

Xanax Tri-Score (PF)

.Nervous system 408

Xarelto (BN)

.Blood and blood forming organs 103

Xeloda (RO)

.Antineoplastic and immunomodulating agents 202

Xenical (RO)

.Repatriation Schedule 839

Xergic (AF)

.Repatriation Schedule 858

Xigris (LY)

.Blood and blood forming organs 101

XLYS, LOW TRY Maxamaid (SB)

.Various 463

XMET Maxamaid (SB)

.Various 463

XMET Maxamum (SB)

.Various 463

XMTVI Maxamaid (SB)

.Various 463

XMTVI Maxamum (SB)

.Various 463

XP Maxamaid (SB)

.Various 463

XP Maxamum (SB)

.Various 463, 464

XPhen, Tyr Maxamaid (SB)

.Various 464

XPhen, Tyr Maxamum (SB)

.Various 464

Xydep 100 (GN)

.Nervous system 415, 416

Xydep 50 (GN)

.Nervous system 415, 416

Xylocaine Viscous (AP)

.Repatriation Schedule 844

Xylocard 500 (AP)

.Cardiovascular system 107

Z**Z.S.C. (SI)**

.Repatriation Schedule 846

Zabel (AF)

.Dermatologicals 150

.Repatriation Schedule 843

Zactin (AF)

.Nervous system 414

Zan-Extra 10/10 (SM)

.Cardiovascular system 129

Zan-Extra 10/20 (SM)

.Cardiovascular system 129

Zanidip (SM)

.Cardiovascular system 119

Zantac (GK)

.Alimentary tract and metabolism 69, 70

Zantac Syrup (GK)

.Alimentary tract and metabolism 70

Zarontin (PF)

.Nervous system 387

Zavedos (PF)

.Antineoplastic and immunomodulating agents 207

Zavedos Solution (PF)

.Antineoplastic and immunomodulating agents 207

Zedace (AF)

.Cardiovascular system 121, 122

Zeffix (GK)

.Section 100 565, 701

Zeldox (PF)

.Nervous system 401

Zentel (GK)

.Antiparasitic products, insecticides and repellents 432

Zerit (BQ)

.Section 100 565, 702

Zestril (AP)

.Cardiovascular system 123, 124

Ziagen (VI)

.Section 100 563, 700

ZIDOVUDINE	
.Section 100	566, 702
Zilarex (SZ)	
.Repatriation Schedule	858
Zimstat (AF)	
.Cardiovascular system	141, 142, 143, 144
ZINC OXIDE	
.Repatriation Schedule	841
ZINC OXIDE with STARCH and CHLORPHENESIN	
.Repatriation Schedule	846
Zincaband 3604 (SS)	
.Repatriation Schedule	865
Zinnat (GK)	
.Antifectives for systemic use	183
.Dental	497
ZIPRASIDONE HYDROCHLORIDE	
.Nervous system	401
ZipZoc 66051550 (SN)	
.Repatriation Schedule	866
Zithromax (PF)	
.Antifectives for systemic use	185
.Sensory organs	440
.Section 100	561, 697
.Repatriation Schedule	849
Zocor (MK)	
.Cardiovascular system	141, 142, 143, 144
Zofran (GK)	
.Alimentary tract and metabolism	77, 78
Zofran syrup 50 mL (GK)	
.Alimentary tract and metabolism	77, 78
Zofran Zydys (GK)	
.Alimentary tract and metabolism	77, 78
ZolaCos CP 10.8/50(28) (AP)	
.Antineoplastic and immunomodulating agents	230
ZolaCos CP 10.8/50(84) (AP)	
.Antineoplastic and immunomodulating agents	231
ZolaCos CP 3.6/50 (AP)	
.Antineoplastic and immunomodulating agents	230
Zoladex 10.8 Implant (AP)	
.Antineoplastic and immunomodulating agents	230
Zoladex Implant (AP)	
.Antineoplastic and immunomodulating agents	230
ZOLEDRONIC ACID	
.Musculo-skeletal system	368
.Section 100	649, 786
ZOLMITRIPTAN	
.Nervous system	67
Zoloft (PF)	
.Nervous system	415, 416
Zomacton (FP)	
.Section 100	794
Zometa (NV)	
.Section 100	649, 786
Zomig (AP)	
.Nervous system	67
ZOPICLONE	
.Repatriation Schedule	855
Zopral (AF)	
.Alimentary tract and metabolism	71, 72
Zoton FasTabs (WX)	
.Alimentary tract and metabolism	71, 72
Zovirax (GK)	
.Sensory organs	440
.Optometrical	516
Zovirax 200 mg (GK)	
.Antifectives for systemic use	194, 195
Zovirax 800 mg (GK)	
.Antifectives for systemic use	195
ZUCLOPENTHIXOL DECANOATE	
.Nervous system	401
Zumenon (SM)	
.Genito urinary system and sex hormones	160
Zyban (GK)	
.Nervous system	427
Zydol (SI)	
.Nervous system	382, 383
.Dental	507
Zydol SR 100 (SI)	
.Nervous system	383
.Dental	507
Zydol SR 150 (SI)	
.Nervous system	383
.Dental	508
Zydol SR 200 (SI)	
.Nervous system	383
.Dental	508
Zyloprim (SI)	
.Musculo-skeletal system	364
Zyprexa (LY)	
.Nervous system	402
Zyprexa Relprevv (LY)	
.Nervous system	402
Zyprexa Zydys (LY)	
.Nervous system	402
Zyrtec (JT)	
.Repatriation Schedule	858

THERAPEUTIC GROUP PREMIUM POLICY

PHARMACEUTICAL BENEFIT ITEMS WHICH HAVE A THERAPEUTIC GROUP PREMIUM WITH EFFECT FROM 1 January 2011

The Schedule of Pharmaceutical Benefits shows differences in price in some therapeutic groups where alternative drugs may have a therapeutic group premium.

The Therapeutic Group Premium Policy applies within narrowly defined therapeutic sub-groups where the drugs concerned are of similar safety and health outcomes.

The Australian Government, through the PBS, subsidises up to the price of the lowest priced drug in the group. This means that consumers may have to pay for more expensive drugs (those with a therapeutic group premium). This extra amount does not count towards their PBS safety net threshold.

Therapeutic group premiums apply where a prescriber has prescribed a drug within a therapeutic group that attracts a therapeutic group premium and has not sought an exemption from Medicare Australia on clinical grounds.

The exemption provisions are:

- adverse effects occurring with all of the base-priced drugs; or
- drug interactions occurring with all of the base-priced drugs; or
- drug interactions expected to occur with all of the base-priced drugs; or
- transfer to a base-priced drug would cause patient confusion resulting in problems with compliance.

The premiums are not a Government charge but reflect the fact that the supplier(s) of the drug charge a price higher than the Government is willing to subsidise.

Under the Therapeutic Group Premium Policy drug substitution by pharmacists is not permitted.

For ease of prescribing and dispensing, and in the interests of your patients, the following list shows those PBS drugs that attract a therapeutic group premium.

Premium Priced Brand	Form and Strength	Max Qty	Therapeutic Group Premium \$

H₂-RECEPTOR ANTAGONISTS

<i>Zantac</i>	Effervescent tablet 150 mg (base)	60	3.26
<i>Zantac Syrup</i>	Syrup 150 mg (base) per 10 mL, 300 mL	2	2.20

The base-priced drugs in this therapeutic group are cimetidine, famotidine, nizatidine, and ranitidine hydrochloride (except ranitidine hydrochloride effervescent tablet 150 mg (base) and syrup 150 mg (base) per 10 mL, 300 mL).

DIHYDROPYRIDINE-DERIVATIVE CALCIUM CHANNEL BLOCKERS

<i>Adalat Oros 20mg</i>	Tablet 20 mg (controlled release)	30	2.26

The base-priced drugs in this therapeutic group are amlodipine, felodipine, lercanidipine hydrochloride and nifedipine (except nifedipine controlled release tablet 20 mg).

ANGIOTENSIN II ANTAGONISTS

<i>Teveten</i>	Tablet 400 mg (base)	56	2.00

The base-priced drugs in this therapeutic group are candesartan cilexetil, irbesartan, olmesartan medoxomil, telmisartan, valsartan and eprosartan mesylate (except eprosartan mesylate tablet 400 mg (base)).

BRAND PREMIUM POLICY

BRANDS OF PHARMACEUTICAL BENEFIT ITEMS WHICH HAVE A BRAND PREMIUM AND THAT MAY BE SUBSTITUTED WITH EFFECT FROM 1 January 2011

The Schedule of Pharmaceutical Benefits shows differences in price between some alternative brands of the same drug product.

Manufacturers can develop generic equivalents and apply to have them listed on the PBS. In doing this, manufacturers need to ensure that they comply with the relevant legislation applicable to patents. These brands are clinically equivalent and must undergo the same strict quality controls. Although these brands are designed to act on the body in exactly the same way, they are usually cheaper than the originator brands.

The Australian Government, through the PBS, subsidises up to the price of the lowest priced brand (except in those instances where the lowest priced brand has, as part of its price, a therapeutic group premium). This means that consumers may have to pay extra for more expensive brands (those with a brand premium). This extra amount does not count towards their PBS safety net threshold.

Brand substitution by pharmacists without reference to the prescriber is permitted for PBS prescriptions where:

- the patient agrees to the substitution;
- the brands are identified in the Schedule of Pharmaceutical Benefits as being interchangeable;
- the prescriber has not indicated on the prescription form that substitution is not to occur; and
- substitution is permitted under the relevant State or Territory legislation.

Prescription forms supplied by Medicare Australia contain a box to be ticked where brand substitution is not to take place.

Prescribers not using these prescription forms should endorse the prescription if brand substitution is not permitted. Where a stamp is used for this purpose, the prescriber will be required to initial the stamped statement.

For ease of prescribing and dispensing, and in the interests of your patients, the following list shows those PBS drugs that attract a brand premium and that can be substituted where permitted. They are listed alphabetically, by brand name, with the brand premium and benchmark brand(s) cited in the last column.

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Abbacillin-V</i>	Oral suspension 150 mg (as benzathine) per 5 mL, 100 mL	2	2.00	Cilicaine V
<i>Accupril</i>	Tablet 5 mg (as hydrochloride)	30	0.49	Acquin 5; APO-Quinapril; Filpril; Pharmacor Quinapril 5; Qpril 5; Quinapril-DP; Quinapril generichealth; Quinapril Sandoz
	Tablet 10 mg (as hydrochloride)	30	0.66	Acquin 10; APO-Quinapril; Filpril; Pharmacor Quinapril 10; Qpril 10; Quinapril-DP; Quinapril generichealth
	Tablet 20 mg (as hydrochloride)	30	1.00	Acquin 20; APO-Quinapril; Filpril; Pharmacor Quinapril 20; Qpril 20; Quinapril-GA; Quinapril generichealth; Quinapril Sandoz
<i>Adalat 10</i>	Tablet 10 mg	60	1.17	Adefin 10
<i>Adalat 20</i>	Tablet 20 mg	60	2.21	Adefin 20; GenRx Nifedipine; Nifehexal
<i>Adalat Oros 30</i>	Tablet 30 mg (controlled release)	30	2.54	Addos XR 30; Adefin XL 30; APO- Nifedipine XR
<i>Adalat Oros 60</i>	Tablet 60 mg (controlled release)	30	2.81	Addos XR 60; Adefin XL 60; APO- Nifedipine XR
<i>Aldactone</i>	Tablet 25 mg	100	1.84	Spiractin 25
	Tablet 100 mg	100	2.53	Spiractin 100
<i>Aldomet</i>	Tablet 250 mg	100	2.65	Hydopa
<i>Alphagan</i>	Eye drops 2 mg per mL (0.2%), 5 mL	1	1.66	Enidin
<i>Amaryl</i>	Tablet 1 mg	30	2.81	APO-Glimepiride; Aylide 1; Diapride 1; Dimirel; Glimepiride Sandoz
	Tablet 2 mg	30	2.81	APO-Glimepiride; Aylide 2; Diapride 2; Dimirel; Glimepiride Sandoz
	Tablet 3 mg	30	2.80	APO-Glimepiride; Aylide 3; Diapride 3; Dimirel; Glimepiride Sandoz
	Tablet 4 mg	30	2.80	APO-Glimepiride; Aylide 4; Diapride 4; Dimirel; Glimepiride Sandoz
<i>Amoxil</i>	Capsule 250 mg	20	0.79	Alphamox 250; Amoxycillin-GA; Amoxycillin Ranbaxy; Amoxycillin Sandoz; APO-Amoxycillin; Chem mart Amoxycillin; Cilamox; GenRx Amoxycillin; Terry White Chemists Amoxycillin
	Capsule 500 mg	20	0.80	Alphamox 500; Amoxycillin-GA; Amoxycillin Ranbaxy; Amoxycillin Sandoz; APO-Amoxycillin; Chem mart Amoxycillin; Cilamox; GenRx Amoxycillin; Terry White Chemists Amoxycillin
	Powder for syrup 125 mg per 5 mL, 100 mL	1	0.95	Alphamox 125; Amoxycillin Sandoz; Bgramin; Chem mart Amoxycillin; GenRx Amoxycillin; Ranmoxy; Terry White Chemists Amoxycillin
<i>Amoxil Forte</i>	Powder for syrup 250 mg per 5 mL, 100 mL	1	0.79	Alphamox 250; Amoxycillin Sandoz; Bgramin; Chem mart Amoxycillin; Cilamox; GenRx Amoxycillin; Ranmoxy; Terry White Chemists Amoxycillin
<i>Anafranil 25</i>	Tablet 25 mg	50	3.26	Chem mart Clomipramine; GenRx Clomipramine; Placil; Terry White Chemists Clomipramine
<i>Anaprox 550</i>	Tablet 550 mg	50	2.29	Crysanal
<i>Androcur</i>	Tablet 50 mg	20	3.13	Cyprohexal; Cyprone; Cyprostat; GenRx Cyproterone Acetate; Procur
	Tablet 50 mg	100	3.00	Cyprohexal; Cyprone; Cyprostat; GenRx Cyproterone Acetate; Procur
<i>Androcur-100</i>	Tablet 100 mg	50	1.65	Cyprohexal; Cyprostat-100; GenRx Cyproterone Acetate; Procur 100
<i>Anginine Stabilised</i>	Tablets 600 micrograms, 100	1	3.00	Lycinate
<i>Aristocort 0.02%</i>	Cream 200 micrograms per g (0.02%), 100 g	2	3.84	Tricortone
	Ointment 200 micrograms per g (0.02%), 100 g	2	3.84	Tricortone
<i>Aropax</i>	Tablet 20 mg (as hydrochloride)	30	0.86	Chem mart Paroxetine; Extine 20; GenRx Paroxetine; Paroxetine 20; Paroxetine-DP; Paroxetine-GA; Paroxetine Sandoz; Paxtine; Terry White Chemists Paroxetine
<i>Astrix</i>	Tablet 100 mg	112	1.32	DBL Aspirin 100 mg

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Atrovent</i>	Nebuliser solution single dose units 250 micrograms (anhydrous) in 1 mL, 30	2	0.72	Aeron 250; APO-Ipratropium; Ipratrin; Ipravent
<i>Atrovent Adult</i>	Nebuliser solution single dose units 500 micrograms (anhydrous) in 1 mL, 30	2	0.62	Aeron 500; APO-Ipratropium; Ipratrin Adult; Ipravent
<i>Augmentin</i>	Powder for syrup 125 mg-31.25 mg per 5 mL, 75 mL	1	1.51	Clamoxyl; Curam
<i>Augmentin Duo</i>	Tablet 500 mg-125 mg	10	1.55	Amoxycillin/ Clavulanic Acid 500/125 generichealth; APO-Amoxycillin/ Clavulanic Acid 500/125; Clamoxyl Duo; Curam Duo 500/125; GA-Amclav 500/125; Moxiclav Duo 500/125
<i>Augmentin Duo 400</i>	Powder for syrup 400 mg-57 mg per 5 mL, 60 mL	1	1.54	Clamoxyl Duo 400; Curam Duo
<i>Augmentin Duo forte</i>	Tablet 875 mg-125 mg	10	2.04	Amoxycillin/ Clavulanic Acid 875/125 generichealth; Chem mart Amoxycillin and Clavulanic Acid; Clamoxyl Duo forte; Clavycillin 875/125; Curam Duo Forte 875/125; GA-Amclav Forte 875/125; GenRx Amoxycillin and Clavulanic Acid; Moxiclav Duo Forte 875/125; Terry White Chemists Amoxycillin and Clavulanic Acid
<i>Aurorix</i>	Tablet 150 mg	60	0.72	Amira 150; Chem mart Moclobemide; Clobemix; GenRx Moclobemide; Moclobemide Sandoz; Moheal; Terry White Chemists Moclobemide
<i>Aurorix 300 mg</i>	Tablet 300 mg	60	1.45	Amira 300; Chem mart Moclobemide; Clobemix; GenRx Moclobemide; Moclobemide Sandoz; Terry White Chemists Moclobemide
<i>Avanza</i>	Tablet 30 mg	30	3.09	Axit 30; Chem mart Mirtazapine; GenRx Mirtazapine; Mirtazapine-DP; Mirtazapine Sandoz; Mirtazon; Terry White Chemists Mirtazapine
	Tablet 45 mg	30	2.91	APO-Mirtazapine; Chem mart Mirtazapine; Mirtazapine Sandoz; Mirtazon; Terry White Chemists Mirtazapine
<i>Azopt</i>	Eye drops 10 mg per mL (1%), 5 mL	1	1.18	BrinzoQuin
<i>Betaloc</i>	Tablet 50 mg	100	3.25	Chem mart Metoprolol; GenRx Metoprolol; Metohexal; Metrol 50; Minax 50; Terry White Chemists Metoprolol
	Tablet 100 mg	60	3.25	Chem mart Metoprolol; GenRx Metoprolol; Metohexal; Metrol 100; Minax 100; Terry White Chemists Metoprolol
<i>Betnovate 1/2</i>	Cream 500 micrograms (base) per g (0.05%), 15 g	1	3.00	Cortival 1/2
	Ointment 500 micrograms (base) per g (0.05%), 15 g	1	3.00	Cortival 1/2
<i>Betnovate 1/5</i>	Cream 200 micrograms (base) per g (0.02%), 100 g	2	7.00	Cortival 1/5
<i>Betoptic</i>	Eye drops, solution, 5 mg (base) per mL (0.5%), 5 mL	1	2.09	BetoQuin
<i>Blenoxane</i>	Powder for injection 15,000 i.u. (solvent required)	10	72.32	Blenamax; Hospira Pty Limited
	Pack containing 21 tablets 500 micrograms-35 micrograms and 7 inert tablets	4	7.88	Norimin 28 Day
	Pack containing 21 tablets 1 mg-35 micrograms and 7 inert tablets	4	7.88	Norimin-1 28 Day
<i>Capoten</i>	Tablet 25 mg	90	5.00	Ascent Pharma Pty Ltd; Captopril Sandoz; GenRx Captopril; Zedace
	Tablet 50 mg	90	5.00	Ascent Pharma Pty Ltd; Captopril Sandoz; GenRx Captopril; Zedace
<i>Carafate</i>	Tablet equivalent to 1 g anhydrous sucralfate	120	2.11	Ulcylte
<i>Ceclor</i>	Powder for oral suspension 125 mg per 5 mL, 100 mL	1	4.18	Aclor 125; Cefaclor Sandoz; Chem mart Cefaclor; GenRx Cefaclor; Keflor; Ozcef; Terry White Chemists Cefaclor
	Powder for oral suspension 250 mg per 5 mL, 75 mL	1	4.37	Aclor 250; Cefaclor Sandoz; Chem mart Cefaclor; GenRx Cefaclor; Keflor; Ozcef; Terry White Chemists Cefaclor
<i>Ceclor CD</i>	Tablet 375 mg (sustained release)	10	5.20	Cefaclor-GA; Chem mart Cefaclor CD; Douglas Cefaclor-CD; GenRx Cefaclor CD; Karlor CD; Keflor CD; Ozcef; Terry White Chemists Cefaclor CD
<i>Celestone-M</i>	Cream 200 micrograms (base) per g (0.02%), 100 g	2	2.50	Antroquoril

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Cipramil</i>	Ointment 200 micrograms (base) per g (0.02%), 100 g	2	2.50	Antroquoril
	Eye drops 3 mg per mL (0.3%), 5 mL	2	2.06	CiloQuin
	Tablet 20 mg (base)	28	4.45	AP0-Citalopram; Celapram; Celica; Chem mart Citalopram; Ciazil; Citalobell; Citalopram 20; Citalopram generichealth; Citalopram Sandoz; GenRx Citalopram; Talam; Terry White Chemists Citalopram
<i>Ciproxin 250</i>	Tablet 250 mg	14	1.45	C-Flox 250; Cifran; Ciprofloxacin-DRLA; Ciprofloxacin Sandoz; Ciprol 250; GenRx Ciprofloxacin; Profloxin
<i>Ciproxin 500</i>	Tablet 500 mg	14	1.26	Ascent Pharmaceuticals Limited; C-Flox 500; Cifran; Ciprofloxacin 500; Ciprofloxacin-BW; Ciprofloxacin-DRLA; Ciprofloxacin-GA; Ciprofloxacin Sandoz; Ciprol 500; GenRx Ciprofloxacin
<i>Ciproxin 750</i>	Tablet 750 mg	14	1.37	Ascent Pharmaceuticals Limited; C-Flox 750; Cifran; Ciprofloxacin 750; Ciprofloxacin-BW; Ciprofloxacin-DRLA; Ciprofloxacin-GA; Ciprofloxacin Sandoz; Ciprol 750; GenRx Ciprofloxacin
<i>Colgout</i>	Tablet 500 micrograms	30	0.87	Lengout
<i>Dalacin C</i>	Capsule 150 mg	24	1.40	Cleocin
<i>Daonil</i>	Tablet 5 mg	100	1.44	Glimel
<i>Depo-Medrol</i>	Injection 40 mg in 1 mL	5	0.73	Depo-Nisolone
<i>Depo-Provera</i>	Injection 150 mg in 1 mL	1	3.26	Depo-Ralovera
<i>Diabex</i>	Tablet 500 mg	100	1.79	Ascent Pharmaceuticals Limited; Chem mart Metformin; Diaformin; Formet 500; GenRx Metformin; Glucohexal; Glucophage; Metformin 500; Metformin-GA; Metformin generichealth; Metformin Ranbaxy; Metformin Sandoz; Terry White Chemists Metformin
<i>Diabex 1000</i>	Tablet 1 g	90	1.80	Diaformin 1000; Formet 1000; Glucohexal; Metformin-GA; Metformin generichealth 1000; Metformin Ranbaxy 1000; Metformin Sandoz; Pharmacor Metformin 1000
<i>Diabex 850</i>	Tablet 850 mg	60	1.79	Ascent Pharmaceuticals Limited; Chem mart Metformin; Diaformin 850; Formet 850; GenRx Metformin; Glucohexal; Glucophage; Metformin 850; Metformin-GA; Metformin generichealth; Metformin Ranbaxy; Metformin Sandoz; Terry White Chemists Metformin
<i>Diprosone</i>	Cream 500 micrograms (base) per g (0.05%), 15 g	1	2.50	Eleuphrat
<i>Doryx</i>	Ointment 500 micrograms (base) per g (0.05%), 15 g	1	2.50	Eleuphrat
	Capsule 100 mg (as hydrochloride)	7	1.15	Mayne Pharma Doxycycline
	Capsule 50 mg (as hydrochloride)	25	1.31	Mayne Pharma Doxycycline
<i>Dulcolax</i>	Capsule 100 mg (as hydrochloride)	28	4.60	Mayne Pharma Doxycycline
	Capsule 100 mg (as hydrochloride)	21	2.09	Mayne Pharma Doxycycline
	Suppositories 10 mg, 10	3	1.08	Petrus Bisacodyl Suppositories
<i>Duphalac</i>	Mixture 3.34 g per 5 mL, 500 mL	1	1.67	Actilax; Genlac; GenRx Lactulose; Lac-Dol; Lactocur
	Mixture 3.34 g per 5 mL, 500 mL	3	5.01	Actilax; Genlac; GenRx Lactulose; Lac-Dol; Lactocur
<i>Duratears</i>	Compound eye ointment 3.5 g	2	2.16	Poly Visc
<i>E.E.S. 200</i>	Powder for oral liquid 200 mg (base) per 5 mL, 100 mL	1	2.86	E-Mycin 200
<i>E.E.S. 400 Filmtab</i>	Tablet 400 mg (base)	25	2.80	E-Mycin
<i>E.E.S. Granules</i>	Powder for oral liquid 400 mg (base) per 5 mL, 100 mL	1	2.89	E-Mycin 400
<i>Elocon</i>	Cream 1 mg per g (0.1%), 15 g	1	2.50	Novasone
<i>Epilim EC</i>	Ointment 1 mg per g (0.1%), 15 g	1	2.50	Novasone
	Lotion 1 mg per g (0.1% w/w), 30 mL	1	2.50	Novasone
	Tablet 200 mg (enteric coated)	200	1.42	Sodium Valproate Sandoz; Valprease 200; Valpro 200; Valproate Winthrop EC 200
	Tablet 500 mg (enteric coated)	200	1.40	Sodium Valproate Sandoz; Valprease

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
				500; Valpro 500; Valproate Winthrop EC 500
<i>Eryc</i>	Capsule 250 mg	25	1.35	Mayne Pharma Erythromycin
<i>Fasigyn</i>	Tablet 500 mg	4	2.46	Simplotan
<i>Feldene</i>	Capsule 10 mg	50	2.66	Chem mart Piroxicam; GenRx Piroxicam; Mobilis 10; Terry White Chemists Piroxicam
	Capsule 20 mg	25	2.64	Chem mart Piroxicam; GenRx Piroxicam; Mobilis 20; Terry White Chemists Piroxicam
<i>Feldene-D</i>	Dispersible tablet 20 mg	25	2.64	Mobilis D-20
<i>Flagyl</i>	Tablet 400 mg	21	2.30	Metrogyl 400; Metronide 400
	Tablet 200 mg	21	2.30	Metrogyl 200; Metronide 200
<i>Fosamax Once Weekly</i>	Tablet equivalent to 70 mg alendronic acid	4	2.00	Adronat; Alendrobell 70mg; Alendronate-GA; Alendronate Sandoz; Alendro Once Weekly; APO-Alendronate; Chem mart Alendronate 70mg; Ossmax 70mg; Terry White Chemists Alendronate 70mg
<i>Genteal</i>	Eye drops 3 mg per mL (0.3%), 15 mL (contains sodium perborate as preservative)	1	1.79	In a Wink Moisturising
<i>Genteal gel</i>	Ocular lubricating gel 3 mg-2 mg per g (0.3%-0.2%), 10 g	1	1.79	HPMC PAA
<i>Glucophage</i>	Tablet 500 mg	100	1.10	Ascent Pharmaceuticals Limited; Chem mart Metformin; Diabex; Diaformin; Formet 500; GenRx Metformin; Glucohexal; Metformin 500; Metformin-GA; Metformin generichealth; Metformin Ranbaxy; Metformin Sandoz; Terry White Chemists Metformin
	Tablet 850 mg	60	1.10	Ascent Pharmaceuticals Limited; Chem mart Metformin; Diabex 850; Diaformin 850; Formet 850; GenRx Metformin; Glucohexal; Metformin 850; Metformin-GA; Metformin generichealth; Metformin Ranbaxy; Metformin Sandoz; Terry White Chemists Metformin
<i>Gopten</i>	Capsule 500 micrograms	28	1.87	APO-Trandolapril; Dolapril 0.5; Tranalpha; Trandolapril-DP; Trandolapril generichealth
	Capsule 1 mg	28	1.87	APO-Trandolapril; Dolapril 1; Tranalpha; Trandolapril-DP; Trandolapril generichealth
	Capsule 2 mg	28	1.87	APO-Trandolapril; Dolapril 2; Tranalpha; Trandolapril-DP; Trandolapril generichealth
	Capsule 4 mg	28	1.88	APO-Trandolapril; Dolapril 4; Tranalpha; Trandolapril-DP; Trandolapril generichealth
<i>Imdur 120 mg</i>	Tablet 120 mg (sustained release)	30	3.00	Monodur 120 mg
<i>Imdur Durule</i>	Tablet 60 mg (sustained release)	30	2.84	Chem mart Isosorbide Mononitrate; Duride; GenRx Isosorbide Mononitrate; Imtrate 60 mg; Isomonit; Monodur 60 mg; Terry White Chemists Isosorbide Mononitrate
<i>Imodium</i>	Capsule 2 mg	12	0.91	Gastro-Stop Loperamide
<i>Indocid</i>	Capsule 25 mg	100	2.16	Arthrexin
<i>Isoptin</i>	Tablet 40 mg	100	0.76	Anpec 40
	Tablet 80 mg	100	0.76	Anpec 80
<i>Isoptin 180 SR</i>	Tablet 180 mg (sustained release)	30	2.27	Cordilox 180 SR
<i>Isoptin SR</i>	Tablet 240 mg (sustained release)	30	2.26	Cordilox SR
<i>Keflex</i>	Capsule 250 mg	20	3.30	Cefalexin Sandoz; Cephabell; Cephalexin generichealth; Cephatrust 250; Chem mart Cephalexin; Cilex; GenRx Cephalexin; Ialex; Ibilex 250; Rancef; Terry White Chemists Cephalexin
	Capsule 500 mg	20	4.41	Cefalexin Sandoz; Cephabell; Cephalexin generichealth; Cephatrust

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
				500; Chem mart Cephalixin; Cilex; GenRx Cephalixin; Ialex; Ibilex 500; Rancef; Terry White Chemists Cephalixin
	Granules for syrup 125 mg per 5 mL, 100 mL	1	3.55	APO-Cephalixin; Cefalexin Sandoz; Chem mart Cephalixin; Cilex; GenRx Cephalixin; Ialex; Ibilex 125; Terry White Chemists Cephalixin
	Granules for syrup 250 mg per 5 mL, 100 mL	1	4.38	APO-Cephalixin; Cefalexin Sandoz; Chem mart Cephalixin; Cilex; GenRx Cephalixin; Ialex; Ibilex 250; Terry White Chemists Cephalixin
<i>Kenacomb Otic</i>	Ear drops 1 mg-2.5 mg (base)- 250 micrograms- 100,000 units per g (0.1%-0.25%-0.025%-100,000 units per g), 7.5 mL	1	2.00	Otocomb Otic
	Ear ointment 1 mg-2.5 mg (base)- 250 micrograms- 100,000 units per g (0.1%-0.25%-0.025%-100,000 units per g), 5 g	1	2.01	Otocomb Otic
<i>Klacid</i>	Tablet 250 mg	14	1.96	APO-Clarithromycin; Chem mart Clarithromycin; Clarac; Clarihexal; Clarithro 250; GenRx Clarithromycin; Kalixocin; Terry White Chemists Clarithromycin
<i>Lacri-Lube</i>	Pack containing 2 tubes compound eye ointment 3.5 g	1	2.16	Ircal
<i>Lamictal</i>	Tablet 5 mg	56	0.76	Lamogine; Seaze 5
	Tablet 25 mg	56	0.77	APO-Lamotrigine; GenRx Lamotrigine; Lamidus; Lamogine; Lamotrigine-DP; Lamotrigine-GA; Lamotrigine generichealth; Lamotrigine Sandoz; Lamotrust 25; Seaze 25
	Tablet 50 mg	56	0.66	APO-Lamotrigine; GenRx Lamotrigine; Lamidus; Lamogine; Lamotrigine-DP; Lamotrigine-GA; Lamotrigine generichealth; Lamotrigine Sandoz; Lamotrust 50; Seaze 50
	Tablet 100 mg	56	0.73	APO-Lamotrigine; GenRx Lamotrigine; Lamidus; Lamogine; Lamotrigine-DP; Lamotrigine-GA; Lamotrigine generichealth; Lamotrigine Sandoz; Lamotrust 100; Seaze 100
	Tablet 200 mg	56	0.72	APO-Lamotrigine; GenRx Lamotrigine; Lamidus; Lamogine; Lamotrigine-DP; Lamotrigine-GA; Lamotrigine generichealth; Lamotrigine Sandoz; Lamotrust 200; Seaze 200
<i>Lamisil</i>	Tablet 250 mg (as hydrochloride)	42	1.46	GenRx Terbinafine; Sebifin 250; Tamsil; Terbihexal; Terbinafine 250; Terbinafine-DRLA; Terbinafine-GA; Terbix 250; Zabel
<i>Lanoxin</i>	Tablet 250 micrograms	100	3.00	Sigmaxin
<i>Lanoxin-PG</i>	Tablet 62.5 micrograms	200	3.00	Sigmaxin-PG
<i>Lasix</i>	Tablet 40 mg	100	2.40	Chem mart Frusemide; Frusemide Sandoz; Frusid; GenRx Frusemide; Terry White Chemists Frusemide; Uremide
<i>Lasix-M</i>	Tablet 20 mg	100	1.90	Chem mart Frusemide; Frusid; GenRx Frusemide; Terry White Chemists Frusemide
<i>Lexapro</i>	Tablet 10 mg (base)	28	4.69	APO-Escitalopram; Chem mart Escitalopram; Esipram; Esitalo; Lexam 10; LoxaLate; Terry White Chemists Escitalopram
	Tablet 20 mg (base)	28	6.86	APO-Escitalopram; Chem mart Escitalopram; Esipram; Esitalo; Lexam 20; LoxaLate; Terry White Chemists Escitalopram
<i>Lioresal 10</i>	Tablet 10 mg	100	2.26	Chem mart Baclofen; Clofen 10; GenRx Baclofen; Stelax 10; Terry White Chemists Baclofen
<i>Lioresal 25</i>	Tablet 25 mg	100	2.18	Chem mart Baclofen; Clofen 25; GenRx Baclofen; Stelax 25; Terry

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Lipex 10</i>	Tablet 10 mg	30	3.51	White Chemists Baclofen APO-Simvastatin; Chem mart Simvastatin; GenRx Simvastatin; Pharmacor Simvastatin 10; Ransim; Simvahexal; Simvar 10; Simvastatin- DP; Simvastatin-GA 10; Simvastatin generichealth; Simvastatin-Spirit 10; Simvastatin Winthrop; Simvasyn; Terry White Chemists Simvastatin; Zimstat; Zocor
<i>Lipex 20</i>	Tablet 20 mg	30	3.50	APO-Simvastatin; Chem mart Simvastatin; GenRx Simvastatin; Pharmacor Simvastatin 20; Ransim; Simvahexal; Simvar 20; Simvastatin- DP; Simvastatin-GA 20; Simvastatin generichealth; Simvastatin-Spirit 20; Simvastatin Winthrop; Simvasyn; Terry White Chemists Simvastatin; Zimstat; Zocor
<i>Lipex 40</i>	Tablet 40 mg	30	3.50	APO-Simvastatin; Chem mart Simvastatin; GenRx Simvastatin; Pharmacor Simvastatin 40; Ransim; Simvahexal; Simvar 40; Simvastatin- DP; Simvastatin-GA 40; Simvastatin generichealth; Simvastatin-Spirit 40; Simvastatin Winthrop; Simvasyn; Terry White Chemists Simvastatin; Zimstat; Zocor
<i>Lipex 80</i>	Tablet 80 mg	30	3.50	APO-Simvastatin; Chem mart Simvastatin; GenRx Simvastatin; Pharmacor Simvastatin 80; Ransim; Simvahexal; Simvar 80; Simvastatin- DP; Simvastatin-GA 80; Simvastatin generichealth; Simvastatin-Spirit 80; Simvastatin Winthrop; Simvasyn; Terry White Chemists Simvastatin; Zimstat; Zocor
<i>Liquifilm Forte</i>	Eye drops 30 mg per mL (3%), 15 mL	1	5.71	PVA Forte
<i>Liquifilm Tears</i>	Eye drops 14 mg per mL (1.4%), 15 mL	1	1.64	PVA Tears
<i>Lomotil</i>	Tablet 2.5 mg-25 micrograms	20	1.76	Lofenoxal
<i>Lopid</i>	Tablet 600 mg	60	2.95	Ausgem; Chem mart Gemfibrozil; Gemhexal; GenRx Gemfibrozil; Jezil; Lipazil 600 mg; Lipigem; Pharmacor Gemfibrozil 600; Terry White Chemists Gemfibrozil
<i>Losec Tablets</i>	Tablet 20 mg (as magnesium)	30	3.75	Acimax Tablets; Omepral
<i>Luvox</i>	Tablet 50 mg	30	2.96	APO-Fluvoxamine; Faverin 50; Movox 50; Voxam
	Tablet 100 mg	30	2.95	APO-Fluvoxamine; Faverin 100; Movox 100; Voxam
<i>Maxamox</i>	Tablet 1 g	14	1.17	Amoxycillin Sandoz
<i>Microgynon 30 ED</i>	Pack containing 21 tablets 150 micrograms-30 micrograms and 7 inert tablets	4	13.87	Levlen ED
<i>Minidiab</i>	Tablet 5 mg	100	3.91	Melizide
<i>Minipress</i>	Tablet 1 mg (base)	100	2.95	Chem mart Prazosin; GenRx Prazosin; Terry White Chemists Prazosin
	Tablet 2 mg (base)	100	3.04	Chem mart Prazosin; GenRx Prazosin; Terry White Chemists Prazosin
	Tablet 5 mg (base)	100	3.29	Chem mart Prazosin; GenRx Prazosin; Terry White Chemists Prazosin
<i>Minomycin-50</i>	Tablet 50 mg	60	2.00	Akamin 50
<i>Mobic</i>	Tablet 7.5 mg	30	1.30	Chem mart Meloxicam 7.5 mg; GenRx Meloxicam; Meloxibell; Meloxicam- GA; Meloxicam Ranbaxy; Meloxicam Sandoz; Meloxicam Winthrop; Movalis 7.5; Moxicam 7.5; Pharmacor Meloxicam 7.5; Terry White Chemists Meloxicam 7.5 mg
	Tablet 15 mg	30	1.30	Chem mart Meloxicam 15 mg; GenRx Meloxicam; Meloxibell; Meloxicam- GA; Meloxicam Ranbaxy; Meloxicam

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
				Sandoz; Meloxicam Winthrop; Movalis 15; Moxicam 15; Pharmacor Meloxicam 15; Terry White Chemists Meloxicam 15 mg
<i>Mogadon</i>	Tablet 5 mg	50	3.06	Alodorm
	Tablet 5 mg	25	1.53	Alodorm
<i>Naprosyn</i>	Tablet 250 mg	100	2.34	Inza 250
	Tablet 500 mg	50	1.36	Inza 500
<i>Naprosyn SR1000</i>	Tablet 1 g (sustained release)	28	1.34	Proxen SR 1000
<i>Naprosyn SR750</i>	Tablet 750 mg (sustained release)	28	1.27	Proxen SR 750
<i>Natrilix</i>	Tablet 2.5 mg	90	2.56	Chem mart Indapamide; Dapa-Tabs; GenRx Indapamide; Indapamide-GA; Indapamide Sandoz; Insig; Terry White Chemists Indapamide
<i>Neoral 100</i>	Capsule 100 mg	60	2.16	Cicloral
<i>Neoral 25</i>	Capsule 25 mg	60	2.20	Cicloral
<i>Neoral 50</i>	Capsule 50 mg	60	2.38	Cicloral
<i>Neurontin</i>	Capsule 100 mg	100	1.00	APO-Gabapentin; DBL Gabapentin; Gabatine 100; Gantin; Nupentin 100
	Capsule 300 mg	100	0.99	DBL Gabapentin; Gabapentin 300; Gabapentin-GA; Gabapentin Sandoz; Gabatine 300; Gantin; GenRx Gabapentin; Nupentin 300
	Capsule 400 mg	100	1.00	DBL Gabapentin; Douglas Gabapentin 400mg; Gabapentin 400; Gabapentin Sandoz; Gabatine 400; Gantin; GenRx Gabapentin; Nupentin 400
	Tablet 600 mg	100	1.00	Gabaran; Gabatine 600; GenRx Gabapentin
	Tablet 800 mg	100	1.00	Gabaran; Gabatine 800; Gantin; GenRx Gabapentin; Pendine 800
<i>Nolvadex-D</i>	Tablet 20 mg (base)	60	3.82	Genox 20; GenRx Tamoxifen; Tamosin; Tamoxen 20 mg; Tamoxifen Sandoz
<i>Nordette 28</i>	Pack containing 21 tablets 150 micrograms-30 micrograms and 7 inert tablets	4	13.82	Monofeme 28
<i>Noriday 28 Day</i>	Tablets 350 micrograms, 28	4	4.00	Locilan 28 Day
<i>Normison</i>	Tablet 10 mg	25	1.50	APO-Temazepam; Temaze; Temtabs
	Tablet 10 mg	50	3.00	APO-Temazepam; Temaze; Temtabs
<i>Noroxin</i>	Tablet 400 mg	14	4.12	Ascent Pharmaceuticals Limited; Chem mart Norfloxacin; GenRx Norfloxacin; Norfloxacin; Nufloxib; Roxin; Terry White Chemists Norfloxacin
<i>Norvasc</i>	Tablet 5 mg (as besylate)	30	3.91	Amlodipine-DRLA; Amlodipine-GA; Amlodipine generichealth; Amlodipine Sandoz; APO-Amlodipine; Chem mart Amlodipine; Nordip; Norvapine; Ozlodip; Perivasc; Pharmacor Amlodipine 5; Terry White Chemists Amlodipine
	Tablet 10 mg (as besylate)	30	5.69	Amlodipine-DRLA; Amlodipine-GA; Amlodipine generichealth; Amlodipine Sandoz; APO-Amlodipine; Chem mart Amlodipine; Nordip; Norvapine; Ozlodip; Perivasc; Pharmacor Amlodipine 10; Terry White Chemists Amlodipine
<i>Oroxine</i>	Tablet equivalent to 50 micrograms anhydrous thyroxine sodium	200	2.25	Eutroxsig
	Tablet equivalent to 75 micrograms anhydrous thyroxine sodium	200	2.32	Eutroxsig
	Tablet equivalent to 100 micrograms anhydrous thyroxine sodium	200	2.25	Eutroxsig
	Tablet equivalent to 200 micrograms anhydrous thyroxine sodium	200	2.25	Eutroxsig
<i>Orudis SR 200</i>	Capsule 200 mg (sustained release)	28	2.21	Oruvail SR
<i>Panadeine Forte</i>	Tablet 30 mg-500 mg	20	2.80	APO- Paracetamol/Codeine 500/30; Codalgine Forte; Codapane Forte; Comfarol Forte; Dolaforte; Prodeine Forte

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
	Tablet 30 mg-500 mg	60	8.40	APO- Paracetamol/Codeine 500/30; Codalgin Forte; Codapane Forte; Comfarol Forte; Dolaforte; Prodeine Forte
<i>Panafcort</i>	Tablet 1 mg	100	0.63	Predsone
<i>Panafcortelone</i>	Tablet 1 mg	100	0.46	Predsolone
<i>Parlodel</i>	Tablet 2.5 mg (base)	30	2.84	Krypton 2.5
	Tablet 2.5 mg (base)	60	2.92	Krypton 2.5
	Capsule 5 mg (base)	60	2.91	Krypton 5
	Capsule 10 mg (base)	100	3.08	Krypton 10
<i>Pepcidine</i>	Tablet 40 mg	30	5.41	Ausfam 40; Chem mart Famotidine; Famotidine Sandoz; GenRx Famotidine; Pamacid 40; Pepzan; Terry White Chemists Famotidine
<i>Pepcidine M</i>	Tablet 20 mg	60	4.95	Ausfam 20; Chem mart Famotidine; Famohexal; Famotidine Sandoz; GenRx Famotidine; Pamacid 20; Pepzan; Terry White Chemists Famotidine
<i>Plendil ER</i>	Tablet 2.5 mg (extended release)	30	4.99	Felodur ER 2.5 mg
	Tablet 5 mg (extended release)	30	5.00	Felodil XR 5; Felodur ER 5 mg
	Tablet 10 mg (extended release)	30	5.00	Felodil XR 10; Felodur ER 10 mg
<i>Pravachol</i>	Tablet containing pravastatin sodium 10 mg	30	3.99	APO-Pravastatin; Chem mart Pravastatin; Cholstat 10; GenRx Pravastatin; Lipostat 10; Pravastatin 10; Pravastatin-GA 10; Pravastatin generichealth; Pravastatin Sandoz; Pravastatin Winthrop; Terry White Chemists Pravastatin
	Tablet containing pravastatin sodium 20 mg	30	4.00	APO-Pravastatin; Chem mart Pravastatin; Cholstat 20; GenRx Pravastatin; Lipostat 20; Pravastatin 20; Pravastatin-GA 20; Pravastatin generichealth; Pravastatin Sandoz; Pravastatin Winthrop; Terry White Chemists Pravastatin; Vastoran
	Tablet containing pravastatin sodium 40 mg	30	4.00	APO-Pravastatin; Chem mart Pravastatin; Cholstat 40; GenRx Pravastatin; Lipostat 40; Pravastatin 40; Pravastatin-GA 40; Pravastatin generichealth; Pravastatin Sandoz; Pravastatin Winthrop; Terry White Chemists Pravastatin; Vastoran
	Tablet containing pravastatin sodium 80 mg	30	4.00	APO-Pravastatin; Chem mart Pravastatin; Lipostat 80; Pravastatin- GA 80; Pravastatin generichealth; Pravastatin Sandoz; Terry White Chemists Pravastatin
<i>Prinivil 10</i>	Tablet 10 mg	30	3.96	APO-Lisinopril; Chem mart Lisinopril; Fibsol 10; GenRx Lisinopril; Liprace; Lisinopril 10; Lisinopril-DRLA; Lisinopril-GA; Lisinopril generichealth; Lisinopril Hexal; Lisinopril Ranbaxy; Lisinopril Winthrop; Lisodur; Terry White Chemists Lisinopril; Zestril
<i>Prinivil 20</i>	Tablet 20 mg	30	3.95	APO-Lisinopril; Chem mart Lisinopril; Fibsol 20; GenRx Lisinopril; Liprace; Lisinopril 20; Lisinopril-DRLA; Lisinopril-GA; Lisinopril generichealth; Lisinopril Ranbaxy; Lisinopril Sandoz; Lisinopril Winthrop; Lisodur; Terry White Chemists Lisinopril; Zestril
<i>Prinivil 5</i>	Tablet 5 mg	30	3.95	APO-Lisinopril; Chem mart Lisinopril; Fibsol 5; GenRx Lisinopril; Liprace; Lisinopril 5; Lisinopril-DRLA; Lisinopril- GA; Lisinopril generichealth; Lisinopril Ranbaxy; Lisinopril Sandoz; Lisinopril Winthrop; Lisodur; Terry White Chemists Lisinopril; Zestril
<i>Prothiaden</i>	Capsule 25 mg	50	1.58	Dothep 25
	Tablet 75 mg	30	0.81	Dothep 75

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Provera</i>	Tablet 10 mg	100	1.57	Ralovera
	Tablet 5 mg	56	1.68	Ralovera
	Tablet 10 mg	30	1.68	Medroxyhexal; Ralovera
<i>Prozac 20</i>	Capsule 20 mg (base)	28	4.13	Auscap; Chem mart Fluoxetine; Fluohexal; Fluoxebell; Fluoxetine 20; Fluoxetine-GA; Fluoxetine generichealth; GenRx Fluoxetine; Lovan; Terry White Chemists Fluoxetine; Zactin
<i>Prozac Tab</i>	Tablet 20 mg (base) (dispersible)	28	4.13	Lovan 20 Tab
<i>Redipred</i>	Oral solution equivalent to 5 mg prednisolone per mL, 30 mL	1	1.86	PredMix
<i>Renitec</i>	Tablet containing enalapril maleate 10 mg	30	4.90	Alphapril; Auspril; Chem mart Enalapril; Enahexal; Enalabell; Enalapril-DP 10mg; Enalapril-GA; Enalapril generichealth; Enalapril Sandoz; Enalapril Winthrop; GenRx Enalapril; Terry White Chemists Enalapril
<i>Renitec 20</i>	Tablet containing enalapril maleate 20 mg	30	4.90	Alphapril; Auspril; Chem mart Enalapril; Enahexal; Enalabell; Enalapril-DP 20mg; Enalapril-GA; Enalapril generichealth; Enalapril Sandoz; GenRx Enalapril; Terry White Chemists Enalapril
<i>Renitec M</i>	Tablet containing enalapril maleate 5 mg	30	4.90	Alphapril; Auspril; Chem mart Enalapril; Enahexal; Enalabell; Enalapril-DP 5mg; Enalapril-GA; Enalapril generichealth; Enalapril Sandoz; Enalapril Winthrop; GenRx Enalapril; Terry White Chemists Enalapril
<i>Rivotril</i>	Tablet 500 micrograms	100	1.79	Paxam 0.5
	Tablet 500 micrograms	200	3.58	Paxam 0.5
	Tablet 2 mg	100	2.04	Paxam 2
	Tablet 2 mg	200	4.08	Paxam 2
<i>Rulide</i>	Capsule 20 mg	60	2.29	GenRx Isotretinoin; Oratane
	Tablet 150 mg	10	2.60	APO-Roxithromycin; Biaxsig; Chem mart Roxithromycin; Roxar 150; Roxide; Roximycin; Roxithromycin-GA; Terry White Chemists Roxithromycin
	Tablet 300 mg	5	2.60	APO-Roxithromycin; Biaxsig; Chem mart Roxithromycin; Roxar 300; Roxide; Roximycin; Roxithromycin-GA; Terry White Chemists Roxithromycin
<i>Salazopyrin-EN</i>	Tablet 500 mg (enteric coated)	200	1.96	Pyralin EN
<i>Seprtin Forte</i>	Tablet 160 mg-800 mg	10	1.54	Bactrim DS; Resprim Forte
<i>Serepax</i>	Tablet 15 mg	25	2.75	Alepam 15
	Tablet 15 mg	50	5.50	Alepam 15
	Tablet 30 mg	25	2.75	Alepam 30; APO-Oxazepam; Murelax
	Tablet 30 mg	50	5.50	Alepam 30; APO-Oxazepam; Murelax
<i>Sigmacort</i>	Cream 10 mg per g (1%), 30 g	1	2.75	Cortic-DS 1%
	Cream 10 mg per g (1%), 50 g	1	2.76	Cortic-DS 1%
	Topical ointment 10 mg per g (1%), 30 g	1	2.75	Cortic-DS 1%
	Topical ointment 10 mg per g (1%), 50 g	1	2.76	Cortic-DS 1%
<i>Sinemet</i>	Tablet 250 mg-25 mg	100	2.98	Levo/Carbidopa Sandoz
<i>Sinemet 100/25</i>	Tablet 100 mg-25 mg	100	5.20	Kinson
<i>Slow-K</i>	Tablet 600 mg (sustained release)	200	2.72	Duro-K
<i>Sofradex</i>	Ear drops 500 micrograms-5 mg-50 micrograms per mL, 8 mL	1	1.91	Otodex
<i>Sotacor</i>	Tablet 80 mg	60	5.00	GenRx Sotalol; Solavert; Sotalol Sandoz
	Tablet 160 mg	60	5.00	Cardol; Chem mart Sotalol; GenRx Sotalol; Solavert; Sotalol Sandoz; Terry White Chemists Sotalol
<i>Stemetil</i>	Tablet containing prochlorperazine maleate 5 mg	25	3.45	APO-Prochlorperazine; ProCalm; Prochlorperazine-GA; Stemizine
<i>Synphasic</i>	Pack containing 12 tablets 500 micrograms-35 micrograms, 9 tablets 1 mg-35 micrograms and 7 inert tablets	4	7.88	Improvil 28 Day

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
<i>Tazac</i>	Capsule 150 mg	60	5.60	Nizac; Tacidine
	Capsule 300 mg	30	5.60	Nizac; Tacidine
<i>Tears Naturale</i>	Eye drops 3 mg-1 mg per mL (0.3%-0.1%), 15 mL	1	1.77	Poly-Tears
<i>Tegretol 100</i>	Tablet 100 mg	200	2.56	Carbamazepine Sandoz
<i>Tegretol 200</i>	Tablet 200 mg	200	2.74	Carbamazepine Sandoz; Teril
<i>Tenormin</i>	Tablet 50 mg	30	3.54	APO-Atenolol; Atenolol-GA; Atenolol Sandoz; Chem mart Atenolol; Noten; Tensig; Terry White Chemists
				Atenolol
<i>Timoptol</i>	Eye drops 2.5 mg (base) per mL (0.25%), 5 mL	1	3.20	Tenopt
	Eye drops 5 mg (base) per mL (0.5%), 5 mL	1	3.20	Tenopt
<i>Tofranil 10</i>	Tablet 10 mg	50	2.85	Tolerade 10
<i>Tofranil 25</i>	Tablet 25 mg	50	2.85	Tolerade 25
<i>Tolvon</i>	Tablet 10 mg	50	1.91	Lumin 10
	Tablet 20 mg	50	2.85	Lumin 20
<i>Tramal</i>	Capsule 50 mg	20	2.42	APO-Tramadol; Chem mart Tramadol; GA Tramadol 50mg; GenRx Tramadol; Lodam 50; Terry White Chemists
				Tramadol; Tramadol Sandoz; Tramedo; Zydol
<i>Tramal SR 100</i>	Tablet 100 mg (twice daily sustained release)	20	4.51	APO-Tramadol SR; Chem mart Tramadol SR; GA Tramadol SR 100mg; Lodam SR 100; Terry White Chemists
				Tramadol SR; Tramaheal SR; Tramedo SR 100; Zydol SR 100
	Tablet 100 mg (twice daily sustained release)	40	9.02	APO-Tramadol SR; Chem mart Tramadol SR; GA Tramadol SR 100mg; Lodam SR 100; Terry White Chemists
				Tramadol SR; Tramaheal SR; Tramedo SR 100; Zydol SR 100
<i>Tramal SR 150</i>	Tablet 150 mg (twice daily sustained release)	20	5.37	APO-Tramadol SR; Chem mart Tramadol SR; GA Tramadol SR 150mg; Lodam SR 150; Terry White Chemists
				Tramadol SR; Tramadol Sandoz SR; Tramedo SR 150; Zydol SR 150
	Tablet 150 mg (twice daily sustained release)	40	10.74	APO-Tramadol SR; Chem mart Tramadol SR; GA Tramadol SR 150mg; Lodam SR 150; Terry White Chemists
				Tramadol SR; Tramadol Sandoz SR; Tramedo SR 150; Zydol SR 150
<i>Tramal SR 200</i>	Tablet 200 mg (twice daily sustained release)	20	6.09	APO-Tramadol SR; Chem mart Tramadol SR; GA Tramadol SR 200mg; Lodam SR 200; Terry White Chemists
				Tramadol SR; Tramaheal SR; Tramedo SR 200; Zydol SR 200
	Tablet 200 mg (twice daily sustained release)	40	11.90	APO-Tramadol SR; Chem mart Tramadol SR; GA Tramadol SR 200mg; Lodam SR 200; Terry White Chemists
				Tramadol SR; Tramaheal SR; Tramedo SR 200; Zydol SR 200
<i>Trandate</i>	Tablet 100 mg	100	3.20	Presolol 100
	Tablet 200 mg	100	3.20	Presolol 200
<i>Triphasil 28</i>	Pack containing 6 tablets 50 micrograms-30 micrograms, 5 tablets 75 micrograms-40 micrograms, 10 tablets 125 micrograms-30 micrograms and 7 inert tablets	4	13.82	Trifeme 28
<i>Triprim</i>	Tablet 300 mg	7	2.00	Alprim
<i>Triquilar ED</i>	Pack containing 6 tablets 50 micrograms-30 micrograms, 5 tablets 75 micrograms-40 micrograms, 10 tablets 125 micrograms-30 micrograms and 7 inert tablets	4	13.87	Logynon ED
<i>Valium</i>	Tablet 2 mg	50	0.86	Antenex 2; Valpam 2
	Tablet 2 mg	100	1.72	Antenex 2; Valpam 2
	Tablet 5 mg	50	0.88	Antenex 5; Diazepam-GA; Ranzepam; Valpam 5
	Tablet 5 mg	100	1.76	Antenex 5; Diazepam-GA; Ranzepam; Valpam 5
<i>Vastin</i>	Capsule 20 mg (as sodium)	28	3.13	Lescol
	Capsule 40 mg (as sodium)	28	3.44	Lescol
<i>Ventolin CFC-Free</i>	Oral pressurised inhalation 100 micrograms (base) per dose (200 doses), CFC-free formulation	2	1.18	Airomir; Asmol CFC-free
<i>Ventolin Nebules</i>	Nebuliser solution single dose units 2.5 mg (base) in 2.5 mL, 30	2	1.46	Asmol 2.5 uni-dose; Butamol 2.5; GenRx Salbutamol; Pfizer Australia

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
	Nebuliser solution single dose units 5 mg (base) in 2.5 mL, 30	2	1.46	Pty Ltd; Pharmacor Salbutamol 2.5; Salbutamol-GA; Salbutamol Sandoz Asmol 5 uni-dose; Butamol 5; GenRx Salbutamol; Pharmacor Salbutamol 5; Salbutamol-GA; Salbutamol Sandoz
<i>Vibramycin</i>	Tablet 100 mg (as hydrochloride)	7	1.19	Doxsig; Doxy-100; Doxylin 100
	Tablet 100 mg (as hydrochloride)	28	4.76	Doxsig; Doxy-100; Doxylin 100
	Tablet 100 mg (as hydrochloride)	21	3.57	Doxsig; Doxy-100; Doxylin 100
<i>Vibra-Tabs</i>	Tablet 50 mg (as hydrochloride)	25	1.25	Doxy-50; Doxylin 50
<i>Viscotears</i>	Eye gel 2 mg per g (0.2%), 10 g	1	0.96	PAA
<i>Visken 15</i>	Tablet 15 mg	50	2.69	Barbloc 15
<i>Visken 5</i>	Tablet 5 mg	100	2.71	Barbloc 5
<i>Voltaren 25</i>	Tablet 25 mg (enteric coated)	100	1.96	APO-Diclofenac; Chem mart Diclofenac; Clonac 25; Diclofenac-GA; Diclofenac Sandoz; Fenac 25; Terry White Chemists Diclofenac
<i>Voltaren 50</i>	Tablet 50 mg (enteric coated)	50	1.96	APO-Diclofenac; Chem mart Diclofenac; Clonac 50; Diclofenac-GA; Diclofenac Sandoz; Fenac; Terry White Chemists Diclofenac
<i>Xanax</i>	Tablet 250 micrograms	50	1.04	Alprax 0.25; Alprazolam Sandoz; Kalma 0.25
	Tablet 500 micrograms	50	1.12	Alprax 0.5; Alprazolam Sandoz; Kalma 0.5
	Tablet 1 mg	50	1.32	Alprax 1; Alprazolam-GA; Alprazolam Sandoz; Chem mart Alprazolam; GenRx Alprazolam; Kalma 1; Terry White Chemists Alprazolam
<i>Xanax Tri-Score</i>	Tablet 2 mg	50	1.60	Alprax 2; Alprazolam-GA; Alprazolam Sandoz; Chem mart Alprazolam; GenRx Alprazolam; Kalma 2; Terry White Chemists Alprazolam
<i>Zanidip</i>	Tablet 10 mg	28	1.95	APO-Lercanidipine; Chem mart Lercanidipine; Lercan; Lercanidipine Sandoz; Terry White Chemists Lercanidipine
	Tablet 20 mg	28	3.44	APO-Lercanidipine; Chem mart Lercanidipine; Lercan; Lercanidipine Sandoz; Terry White Chemists Lercanidipine
<i>Zantac</i>	Tablet 150 mg (base)	60	1.71	Ausran; Chem mart Ranitidine; GenRx Ranitidine; Rani 2; Ranihexal; Ranitidine Sandoz; Ranoxyl; Terry White Chemists Ranitidine; Ulcaid
	Tablet 300 mg (base)	30	1.71	Ausran; Chem mart Ranitidine; GenRx Ranitidine; Rani 2; Ranitidine Sandoz; Terry White Chemists Ranitidine; Ulcaid
<i>Zestril</i>	Tablet 5 mg	30	2.08	APO-Lisinopril; Chem mart Lisinopril; Fibsol 5; GenRx Lisinopril; Liprace; Lisinopril 5; Lisinopril-DRLA; Lisinopril-GA; Lisinopril generichealth; Lisinopril Ranbaxy; Lisinopril Sandoz; Lisinopril Winthrop; Lisodur; Prinivil 5; Terry White Chemists Lisinopril
	Tablet 10 mg	30	2.08	APO-Lisinopril; Chem mart Lisinopril; Fibsol 10; GenRx Lisinopril; Liprace; Lisinopril 10; Lisinopril-DRLA; Lisinopril-GA; Lisinopril generichealth; Lisinopril Hexal; Lisinopril Ranbaxy; Lisinopril Winthrop; Lisodur; Prinivil 10; Terry White Chemists Lisinopril
	Tablet 20 mg	30	2.08	APO-Lisinopril; Chem mart Lisinopril; Fibsol 20; GenRx Lisinopril; Liprace; Lisinopril 20; Lisinopril-DRLA; Lisinopril-GA; Lisinopril generichealth; Lisinopril Ranbaxy; Lisinopril Sandoz; Lisinopril Winthrop; Lisodur; Prinivil 20; Terry White Chemists Lisinopril
<i>Zocor</i>	Tablet 5 mg	30	3.50	Simvahexal; Simvasyn; Zimstat
	Tablet 10 mg	30	3.51	APO-Simvastatin; Chem mart

Premium Priced Brand	Form and Strength	Max. Qty	Brand Premium \$	Benchmark Priced Brands
	Tablet 20 mg	30	3.50	Simvastatin; GenRx Simvastatin; Lipex 10; Pharmacor Simvastatin 10; Ransim; Simvahexal; Simvar 10; Simvastatin-DP; Simvastatin-GA 10; Simvastatin generichealth; Simvastatin-Spirit 10; Simvastatin Winthrop; Simvasyn; Terry White Chemists Simvastatin; Zimstat APO-Simvastatin; Chem mart Simvastatin; GenRx Simvastatin; Lipex 20; Pharmacor Simvastatin 20; Ransim; Simvahexal; Simvar 20; Simvastatin-DP; Simvastatin-GA 20; Simvastatin generichealth; Simvastatin-Spirit 20; Simvastatin Winthrop; Simvasyn; Terry White Chemists Simvastatin; Zimstat APO-Simvastatin; Chem mart Simvastatin; GenRx Simvastatin; Lipex 40; Pharmacor Simvastatin 40; Ransim; Simvahexal; Simvar 40; Simvastatin-DP; Simvastatin-GA 40; Simvastatin generichealth; Simvastatin-Spirit 40; Simvastatin Winthrop; Simvasyn; Terry White Chemists Simvastatin; Zimstat APO-Simvastatin; Chem mart Simvastatin; GenRx Simvastatin; Lipex 80; Pharmacor Simvastatin 80; Ransim; Simvahexal; Simvar 80; Simvastatin-DP; Simvastatin-GA 80; Simvastatin generichealth; Simvastatin-Spirit 80; Simvastatin Winthrop; Simvasyn; Terry White Chemists Simvastatin; Zimstat
	Tablet 40 mg	30	3.50	Chem mart Sertraline; Concorz; Eleva 50; GenRx Sertraline; Sertra 50; Sertraline 50; Sertraline-GA; Sertraline generichealth; Sertraline Winthrop; Setrona; Terry White Chemists Sertraline; Xydep 50
	Tablet 80 mg	30	3.50	Chem mart Sertraline; Concorz; Eleva 100; GenRx Sertraline; Sertra 100; Sertraline 100; Sertraline-GA; Sertraline generichealth; Setrona; Terry White Chemists Sertraline; Xydep 100
	Tablet 50 mg (base)	30	1.49	Acihexal; Acyclo-V 200; GenRx Aciclovir; Lovir
	Tablet 100 mg (base)	30	1.49	Aciclovir 200; Acihexal; Acyclo-V 200; Chem mart Aciclovir; GenRx Aciclovir; Lovir; Ozvir; Terry White Chemists Aciclovir
	Tablet 200 mg	50	4.34	Aciclovir 800; Acihexal; Acyclo-V 800; GenRx Aciclovir
Zovirax 200 mg	Tablet 200 mg	90	3.21	Prexaton
Zovirax 800 mg	Tablet 800 mg	35	1.56	Prexaton
Zyban	Tablet 150 mg (sustained release)	30	0.81	Allopurinol Sandoz; Allosig; Chem mart Allopurinol; GenRx Allopurinol; Progout 100; Terry White Chemists Allopurinol
Zyloprim	Tablet 150 mg (sustained release)	90	0.84	Allopurinol Sandoz; Allosig; Chem mart Allopurinol; GenRx Allopurinol; Progout 300; Terry White Chemists Allopurinol
	Tablet 100 mg	200	3.01	
	Tablet 300 mg	60	3.00	