



Australian Government

Department of Health



Schedule of Pharmaceutical Benefits

Summary of Changes

Effective 1 December 2016



Fees, Patient Contributions and Safety Net Thresholds

The following fees, patient contributions and safety net thresholds apply as at 1 December 2016 and are included, where applicable, in prices published in the Schedule —

Dispensing Fees:	Ready-prepared	\$7.02
	Dangerous drug fee	\$2.95
	Extemporaneously-prepared	\$9.06
	Allowable additional patient charge*	\$4.33
Additional Fees (for safety net prices):	Ready-prepared	\$1.19
	Extemporaneously-prepared	\$1.55
Patient Co-payments:	General	\$38.30
	Concessional	\$6.20
Safety Net Thresholds:	General	\$1475.70
	Concessional	\$372.00
Safety Net Card Issue Fee:		\$9.61

* The allowable additional patient charge is a discretionary charge to general patients if a pharmaceutical item has a dispensed price for maximum quantity less than the general patient co-payment. The pharmacist may charge general patients the allowable additional fee but the fee cannot take the cost of the prescription above the general patient co-payment for the medicine. This fee does not count towards the Safety Net threshold.

Summary of Changes

These changes to the Schedule of Pharmaceutical Benefits are effective from 1 December 2016. The Schedule is updated on the first day of each month and is available on the internet at www.pbs.gov.au.

General Pharmaceutical Benefits

Additions

Addition – Item

10944B	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
10945C	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL cartridges (<i>Humira</i>)
10955N	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL cartridges (<i>Humira</i>)
10960W	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
10961X	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL cartridges (<i>Humira</i>)
10972L	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL syringes (<i>Humira</i>)
10958R	EVOLOCUMAB , evolocumab 140 mg/mL injection, 1 mL injection device (<i>Repatha</i>)
2742H	FLUTAMIDE , flutamide 250 mg tablet, 30 (<i>Flutamide MYLAN</i>)
10952K	LENVATINIB , lenvatinib 4 mg capsule, 30 (<i>Lenvima</i>)
10965D	LENVATINIB , lenvatinib 10 mg capsule, 30 (<i>Lenvima</i>)
10962Y	LEUPRORELIN (&) INERT SUBSTANCE (&) BICALUTAMIDE , leuprorelin acetate 7.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe] (&) bicalutamide 50 mg tablet [28], 1 pack (<i>Bi ELIGARD CP</i>)
10963B	LEUPRORELIN (&) INERT SUBSTANCE (&) BICALUTAMIDE , leuprorelin acetate 22.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe] (&) bicalutamide 50 mg tablet [28], 1 pack (<i>Bi ELIGARD CP</i>)
10969H	LEUPRORELIN (&) INERT SUBSTANCE (&) BICALUTAMIDE , leuprorelin acetate 22.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe] (&) bicalutamide 50 mg tablet [84], 1 pack (<i>Bi ELIGARD CP</i>)
10947E	OCRIPLASMIN , ocriplasmin 500 microgram/0.2 mL injection, 0.2 mL vial (<i>Jetrea</i>)
10951J	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes (<i>Actemra Subcutaneous Injection</i>)
10954M	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes (<i>Actemra Subcutaneous Injection</i>)

Addition – Brand

2600W	<i>Progout 100, AF</i> – ALLOPURINOL , allopurinol 100 mg tablet, 200
2751T	<i>Amlodipine GH, GQ</i> – AMLODIPINE , amlodipine 5 mg tablet, 30
2752W	<i>Amlodipine GH, GQ</i> – AMLODIPINE , amlodipine 10 mg tablet, 30
8717T	<i>Aripiprazole GH, GQ</i> – ARIPIRAZOLE , aripiprazole 10 mg tablet, 30
8717T	<i>Aripiprazole Sandoz, SZ</i> – ARIPIRAZOLE , aripiprazole 10 mg tablet, 30
8718W	<i>Aripiprazole GH, GQ</i> – ARIPIRAZOLE , aripiprazole 15 mg tablet, 30

8718W	<i>Aripiprazole Sandoz, SZ</i> – ARIPIRAZOLE , aripiprazole 15 mg tablet, 30
8719X	<i>Aripiprazole GH, GQ</i> – ARIPIRAZOLE , aripiprazole 20 mg tablet, 30
8719X	<i>Aripiprazole Sandoz, SZ</i> – ARIPIRAZOLE , aripiprazole 20 mg tablet, 30
8720Y	<i>Aripiprazole GH, GQ</i> – ARIPIRAZOLE , aripiprazole 30 mg tablet, 30
8720Y	<i>Aripiprazole Sandoz, SZ</i> – ARIPIRAZOLE , aripiprazole 30 mg tablet, 30
8844L	<i>Bivalirudin APOTEX, TX</i> – BIVALIRUDIN , bivalirudin 250 mg injection, 1 vial
8295N	<i>CANDESAN, RF</i> – CANDESARTAN , candesartan cilexetil 4 mg tablet, 30
8296P	<i>CANDESAN, RF</i> – CANDESARTAN , candesartan cilexetil 8 mg tablet, 30
8297Q	<i>CANDESAN, RF</i> – CANDESARTAN , candesartan cilexetil 16 mg tablet, 30
8889W	<i>CANDESAN, RF</i> – CANDESARTAN , candesartan cilexetil 32 mg tablet, 30
8504N	<i>CANDESAN COMBI 16/12.5, RF</i> – CANDESARTAN + HYDROCHLOROTHIAZIDE , candesartan cilexetil 16 mg + hydrochlorothiazide 12.5 mg tablet, 30
9314F	<i>CANDESAN COMBI 32/12.5, RF</i> – CANDESARTAN + HYDROCHLOROTHIAZIDE , candesartan cilexetil 32 mg + hydrochlorothiazide 12.5 mg tablet, 30
9315G	<i>CANDESAN COMBI 32/25, RF</i> – CANDESARTAN + HYDROCHLOROTHIAZIDE , candesartan cilexetil 32 mg + hydrochlorothiazide 25 mg tablet, 30
10915L	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 100 mg capsule, 60
10918P	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 100 mg capsule, 60
10920R	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 100 mg capsule, 60
10924Y	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 100 mg capsule, 60
10940T	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 100 mg capsule, 60
10941W	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 100 mg capsule, 60
10942X	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 100 mg capsule, 60
10916M	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 400 mg capsule, 30
10917N	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 400 mg capsule, 30
10921T	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 400 mg capsule, 30
10925B	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 400 mg capsule, 30
10933K	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 400 mg capsule, 30
10935M	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 400 mg capsule, 30
10939R	<i>Imatinib-APOTEX, TX</i> – IMATINIB , imatinib 400 mg capsule, 30
8196J	<i>APO-Itraconazole, TX</i> – ITRACONAZOLE , itraconazole 100 mg capsule, 60
8196J	<i>ITRANOX, RW</i> – ITRACONAZOLE , itraconazole 100 mg capsule, 60
8196J	<i>Itracap, AF</i> – ITRACONAZOLE , itraconazole 100 mg capsule, 60
2848X	<i>LAMITAN, RF</i> – LAMOTRIGINE , lamotrigine 25 mg tablet, 56
2849Y	<i>LAMITAN, RF</i> – LAMOTRIGINE , lamotrigine 50 mg tablet, 56
2850B	<i>LAMITAN, RF</i> – LAMOTRIGINE , lamotrigine 100 mg tablet, 56
2851C	<i>LAMITAN, RF</i> – LAMOTRIGINE , lamotrigine 200 mg tablet, 56
9435N	<i>Blooms the Chemist Metformin XR 500, IB</i> – METFORMIN , metformin hydrochloride 500 mg modified release tablet, 120
9435N	<i>Metformin XR 500 APOTEX, GX</i> – METFORMIN , metformin hydrochloride 500 mg modified release tablet, 120
3439B	<i>Blooms the Chemist Metformin XR 1000, IB</i> – METFORMIN , metformin hydrochloride 1 g modified release tablet, 60
3439B	<i>Chem mart Metformin XR 1000, CH</i> – METFORMIN , metformin hydrochloride 1 g modified release tablet, 60
3439B	<i>Terry White Chemists Metformin XR 1000, TW</i> – METFORMIN , metformin hydrochloride 1 g modified release tablet, 60
2147B	<i>Olmesartan - MYL, AF</i> – OLMESARTAN MEDOXOMIL , olmesartan medoxomil 20 mg tablet, 30

2148C	<i>Olmesartan - MYL, AF</i> – OLMESARTAN MEDOXOMIL , olmesartan medoxomil 40 mg tablet, 30
5292M	<i>Olmesartan/Amlodipine - MYL 20/5, AF</i> – OLMESARTAN MEDOXOMIL + AMLODIPINE , olmesartan medoxomil 20 mg + amlodipine 5 mg tablet, 30
5293N	<i>Olmesartan/Amlodipine - MYL 40/5, AF</i> – OLMESARTAN MEDOXOMIL + AMLODIPINE , olmesartan medoxomil 40 mg + amlodipine 5 mg tablet, 30
5294P	<i>Olmesartan/Amlodipine - MYL 40/10, AF</i> – OLMESARTAN MEDOXOMIL + AMLODIPINE , olmesartan medoxomil 40 mg + amlodipine 10 mg tablet, 30
2161R	<i>Olmesartan HCT - MYL 20/12.5, AF</i> – OLMESARTAN MEDOXOMIL + HYDROCHLOROTHIAZIDE , olmesartan medoxomil 20 mg + hydrochlorothiazide 12.5 mg tablet, 30
2166B	<i>Olmesartan HCT - MYL 40/12.5, AF</i> – OLMESARTAN MEDOXOMIL + HYDROCHLOROTHIAZIDE , olmesartan medoxomil 40 mg + hydrochlorothiazide 12.5 mg tablet, 30
2170F	<i>Olmesartan HCT - MYL 40/25, AF</i> – OLMESARTAN MEDOXOMIL + HYDROCHLOROTHIAZIDE , olmesartan medoxomil 40 mg + hydrochlorothiazide 25 mg tablet, 30
5458G	<i>QUEPINE XR, RW</i> – QUETIAPINE , quetiapine 150 mg modified release tablet, 60
2477J	<i>Rivastigmine Patch 5, AF</i> – RIVASTIGMINE , rivastigmine 4.6 mg/24 hours patch, 30
9161E	<i>Rivastigmine Patch 5, AF</i> – RIVASTIGMINE , rivastigmine 4.6 mg/24 hours patch, 30
2551G	<i>Rivastigmine Patch 10, AF</i> – RIVASTIGMINE , rivastigmine 9.5 mg/24 hours patch, 30
9162F	<i>Rivastigmine Patch 10, AF</i> – RIVASTIGMINE , rivastigmine 9.5 mg/24 hours patch, 30
8378Y	<i>Temozolomide Amneal, ED</i> – TEMOZOLOMIDE , temozolomide 5 mg capsule, 5
8819E	<i>Temozolomide Amneal, ED</i> – TEMOZOLOMIDE , temozolomide 5 mg capsule, 5
8379B	<i>Temozolomide Amneal, ED</i> – TEMOZOLOMIDE , temozolomide 20 mg capsule, 5
8820F	<i>Temozolomide Amneal, ED</i> – TEMOZOLOMIDE , temozolomide 20 mg capsule, 5
8380C	<i>Temozolomide Amneal, ED</i> – TEMOZOLOMIDE , temozolomide 100 mg capsule, 5
8821G	<i>Temozolomide Amneal, ED</i> – TEMOZOLOMIDE , temozolomide 100 mg capsule, 5
9361Q	<i>Temozolomide Amneal, ED</i> – TEMOZOLOMIDE , temozolomide 140 mg capsule, 5
9362R	<i>Temozolomide Amneal, ED</i> – TEMOZOLOMIDE , temozolomide 140 mg capsule, 5
8381D	<i>Temozolomide Amneal, ED</i> – TEMOZOLOMIDE , temozolomide 250 mg capsule, 5
10173K	<i>Vzole, RW</i> – VORICONAZOLE , voriconazole 50 mg tablet, 56
9363T	<i>Vzole, RW</i> – VORICONAZOLE , voriconazole 50 mg tablet, 56
10198R	<i>Vzole, RW</i> – VORICONAZOLE , voriconazole 200 mg tablet, 56
9364W	<i>Vzole, RW</i> – VORICONAZOLE , voriconazole 200 mg tablet, 56

Addition – Equivalence Indicator

8717T	<i>Abilify, OS</i> – ARIPIPRAZOLE , aripiprazole 10 mg tablet, 30
8718W	<i>Abilify, OS</i> – ARIPIPRAZOLE , aripiprazole 15 mg tablet, 30
8719X	<i>Abilify, OS</i> – ARIPIPRAZOLE , aripiprazole 20 mg tablet, 30
8720Y	<i>Abilify, OS</i> – ARIPIPRAZOLE , aripiprazole 30 mg tablet, 30
8844L	<i>Angiomax, XM</i> – BIVALIRUDIN , bivalirudin 250 mg injection, 1 vial
8196J	<i>Sporanox, JC</i> – ITRACONAZOLE , itraconazole 100 mg capsule, 60
2147B	<i>Olmetec, MK</i> – OLMESARTAN MEDOXOMIL , olmesartan medoxomil 20 mg tablet, 30
2148C	<i>Olmetec, MK</i> – OLMESARTAN MEDOXOMIL , olmesartan medoxomil 40 mg tablet, 30
5292M	<i>Sevikar 20/5, MK</i> – OLMESARTAN MEDOXOMIL + AMLODIPINE , olmesartan medoxomil 20 mg + amlodipine 5 mg tablet, 30
5293N	<i>Sevikar 40/5, MK</i> – OLMESARTAN MEDOXOMIL + AMLODIPINE , olmesartan medoxomil 40 mg + amlodipine 5 mg tablet, 30
5294P	<i>Sevikar 40/10, MK</i> – OLMESARTAN MEDOXOMIL + AMLODIPINE , olmesartan medoxomil 40 mg + amlodipine 10 mg tablet, 30

2161R	<i>Olmetec Plus, MK</i> – OLMESARTAN MEDOXOMIL + HYDROCHLOROTHIAZIDE , olmesartan medoxomil 20 mg + hydrochlorothiazide 12.5 mg tablet, 30
2166B	<i>Olmetec Plus, MK</i> – OLMESARTAN MEDOXOMIL + HYDROCHLOROTHIAZIDE , olmesartan medoxomil 40 mg + hydrochlorothiazide 12.5 mg tablet, 30
2170F	<i>Olmetec Plus, MK</i> – OLMESARTAN MEDOXOMIL + HYDROCHLOROTHIAZIDE , olmesartan medoxomil 40 mg + hydrochlorothiazide 25 mg tablet, 30
5458G	<i>Seroquel XR, AP</i> – QUETIAPINE , quetiapine 150 mg modified release tablet, 60
2477J	<i>Exelon Patch 5, NV</i> – RIVASTIGMINE , rivastigmine 4.6 mg/24 hours patch, 30
9161E	<i>Exelon Patch 5, NV</i> – RIVASTIGMINE , rivastigmine 4.6 mg/24 hours patch, 30
2551G	<i>Exelon Patch 10, NV</i> – RIVASTIGMINE , rivastigmine 9.5 mg/24 hours patch, 30
9162F	<i>Exelon Patch 10, NV</i> – RIVASTIGMINE , rivastigmine 9.5 mg/24 hours patch, 30

Deletions

Deletion – Item

2014B	ALGINATE SODIUM + CALCIUM CARBONATE + BICARBONATE , alginate sodium 500 mg/10 mL + calcium carbonate 160 mg/10 mL + sodium bicarbonate 267 mg/10 mL oral liquid, 500 mL (<i>Gaviscon P</i>)
1397M	ERYTHROMYCIN LACTOBIONATE , erythromycin (as lactobionate) 1 g injection, 1 vial (<i>Erythrocin-I.V.</i>)
5088T	ERYTHROMYCIN LACTOBIONATE , erythromycin (as lactobionate) 1 g injection, 1 vial (<i>Erythrocin-I.V.</i>) (Dental)
10063P	HOMATROPINE , homatropine hydrobromide 2% eye drops, 15 mL (<i>Isopto Homatropine</i>) (Optometrical)
2541R	HOMATROPINE , homatropine hydrobromide 2% eye drops, 15 mL (<i>Isopto Homatropine</i>)
5492C	OLMESARTAN MEDOXOMIL , olmesartan medoxomil 20 mg tablet, 30 (<i>Olmetec</i>)
5493D	OLMESARTAN MEDOXOMIL , olmesartan medoxomil 40 mg tablet, 30 (<i>Olmetec</i>)

Deletion – Brand

1003T	<i>Lovir, GN</i> – ACICLOVIR , aciclovir 200 mg tablet, 25
2751T	<i>Amlodipine generichealth, GQ</i> – AMLODIPINE , amlodipine 5 mg tablet, 30
2752W	<i>Amlodipine generichealth, GQ</i> – AMLODIPINE , amlodipine 10 mg tablet, 30
1886G	<i>Bgramin, FM</i> – AMOXYCILLIN , amoxycillin 125 mg/5 mL powder for oral liquid, 100 mL
3302T	<i>Bgramin, FM</i> – AMOXYCILLIN , amoxycillin 125 mg/5 mL powder for oral liquid, 100 mL (Dental)
1887H	<i>Bgramin, FM</i> – AMOXYCILLIN , amoxycillin 250 mg/5 mL powder for oral liquid, 100 mL
3393N	<i>Bgramin, FM</i> – AMOXYCILLIN , amoxycillin 250 mg/5 mL powder for oral liquid, 100 mL (Dental)
8213G	<i>Atorvastatin Pfizer, FZ</i> – ATORVASTATIN , atorvastatin 10 mg tablet, 30
9230T	<i>Atorvastatin Pfizer, FZ</i> – ATORVASTATIN , atorvastatin 10 mg tablet, 30
8214H	<i>Atorvastatin Pfizer, FZ</i> – ATORVASTATIN , atorvastatin 20 mg tablet, 30
9231W	<i>Atorvastatin Pfizer, FZ</i> – ATORVASTATIN , atorvastatin 20 mg tablet, 30
8215J	<i>Atorvastatin Pfizer, FZ</i> – ATORVASTATIN , atorvastatin 40 mg tablet, 30
9232X	<i>Atorvastatin Pfizer, FZ</i> – ATORVASTATIN , atorvastatin 40 mg tablet, 30
8521L	<i>Atorvastatin Pfizer, FZ</i> – ATORVASTATIN , atorvastatin 80 mg tablet, 30
9233Y	<i>Atorvastatin Pfizer, FZ</i> – ATORVASTATIN , atorvastatin 80 mg tablet, 30
8362D	<i>Capecitabine GH, GQ</i> – CAPECITABINE , capecitabine 500 mg tablet, 120
8439E	<i>Kudeq, FZ</i> – CELECOXIB , celecoxib 100 mg capsule, 60
8440F	<i>Kudeq, FZ</i> – CELECOXIB , celecoxib 200 mg capsule, 30
8220P	<i>Citalopram generichealth, GQ</i> – CITALOPRAM , citalopram 20 mg tablet, 28
10103R	<i>Exemestane Pfizer, FZ</i> – EXEMESTANE , exemestane 25 mg tablet, 30
8506Q	<i>Exemestane Pfizer, FZ</i> – EXEMESTANE , exemestane 25 mg tablet, 30

1182F	<i>Monopril, BQ</i> – FOSINOPRIL , fosinopril sodium 10 mg tablet, 30
1183G	<i>Monopril, BQ</i> – FOSINOPRIL , fosinopril sodium 20 mg tablet, 30
8401E	<i>Monoplus 20/12.5, BQ</i> – FOSINOPRIL + HYDROCHLOROTHIAZIDE , fosinopril sodium 20 mg + hydrochlorothiazide 12.5 mg tablet, 30
5552F	<i>Latanoprost Pfizer, FZ</i> – LATANOPROST , latanoprost 0.005% eye drops, 2.5 mL (Optometrical)
8243W	<i>Latanoprost Pfizer, FZ</i> – LATANOPROST , latanoprost 0.005% eye drops, 2.5 mL
5553G	<i>Latanocom, FZ</i> – LATANOPROST + TIMOLOL , latanoprost 0.005% + timolol 0.5% eye drops, 2.5 mL (Optometrical)
8895E	<i>Latanocom, FZ</i> – LATANOPROST + TIMOLOL , latanoprost 0.005% + timolol 0.5% eye drops, 2.5 mL
1945J	<i>Ramipril RBX Tabs, RA</i> – RAMIPRIL , ramipril 2.5 mg tablet, 30
1946K	<i>Ramipril RBX Tabs, RA</i> – RAMIPRIL , ramipril 5 mg tablet, 30
1316G	<i>Ramipril RBX Tabs, RA</i> – RAMIPRIL , ramipril 10 mg tablet, 30
8787L	<i>Risperidone GH, GQ</i> – RISPERIDONE , risperidone 500 microgram tablet, 60
8869T	<i>Risperidone GH, GQ</i> – RISPERIDONE , risperidone 500 microgram tablet, 60
2091C	<i>Microlax, JT</i> – SORBITOL + CITRIC ACID + LAURYL SULFOACETATE SODIUM , sorbitol 3.125 g/5 mL + citrate sodium dihydrate 450 mg/5 mL + lauryl sulfoacetate sodium 45 mg/5 mL enema, 12 x 5 mL
8868R	<i>Altven, FZ</i> – VENLAFAXINE , venlafaxine 37.5 mg modified release capsule, 28
8301X	<i>Altven, FZ</i> – VENLAFAXINE , venlafaxine 75 mg modified release capsule, 28
8302Y	<i>Altven, FZ</i> – VENLAFAXINE , venlafaxine 150 mg modified release capsule, 28

Deletion – Equivalence Indicator

2091C	<i>Micolette, AE</i> – SORBITOL + CITRIC ACID + LAURYL SULFOACETATE SODIUM , sorbitol 3.125 g/5 mL + citrate sodium dihydrate 450 mg/5 mL + lauryl sulfoacetate sodium 45 mg/5 mL enema, 12 x 5 mL
-------	---

Alterations

Alteration – Restriction

The following items have additions, deletions or alterations to restrictions, notes and/or cautions.

2168D	AFLIBERCEPT , aflibercept 4 mg/0.1 mL injection, 0.1 mL vial (<i>Eylea</i>)
10138N	RANIBIZUMAB , ranibizumab 1.65 mg/0.165 mL injection, 1 x 0.165 mL syringe (<i>Lucentis</i>)
1382R	RANIBIZUMAB , ranibizumab 2.3 mg/0.23 mL injection, 0.23 mL vial (<i>Lucentis</i>)
10767Q	USTEKINUMAB , ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial (<i>Stelara</i>)

Alteration – Manufacturer Code

		<i>From</i>	<i>To</i>
1007B	<i>Lovir</i> – ACICLOVIR , aciclovir 200 mg tablet, 90	GN	EA
2876J	<i>Xylocard 500</i> – LIGNOCAINE , lignocaine hydrochloride anhydrous 500 mg/5 mL injection, 10 x 5 mL ampoules	AP	AS
9454N	<i>Oxytrol</i> – OXYBUTYNIN , oxybutynin 3.9 mg/24 hours patch, 8	AG	GN
9160D	<i>Lamisil</i> – TERBINAFINE , terbinafine hydrochloride 1% cream, 15 g	NC	GK
8460G	<i>Androderm</i> – TESTOSTERONE , testosterone 2.5 mg/24 hours patch, 60	AG	GN
8619P	<i>Androderm</i> – TESTOSTERONE , testosterone 5 mg/24 hours patch, 30	AG	GN

Advance Notices

1 January 2017

Deletion – Brand

1002R	<i>AciVision, DZ</i> – ACICLOVIR , aciclovir 3% eye ointment, 4.5 g
1007B	<i>Chem mart Aciclovir, CH</i> – ACICLOVIR , aciclovir 200 mg tablet, 90
1007B	<i>Terry White Chemists Aciclovir, TW</i> – ACICLOVIR , aciclovir 200 mg tablet, 90
5501M	<i>AciVision, DZ</i> – ACICLOVIR , aciclovir 3% eye ointment, 4.5 g (Optometrical)
3408J	<i>Anapen Junior, LM</i> – ADRENALINE , adrenaline 150 microgram/0.3 mL injection, 1 dose
3409K	<i>Anapen, LM</i> – ADRENALINE , adrenaline 300 microgram/0.3 mL injection, 1 dose
8295N	<i>Candesartan RBX, RA</i> – CANDESARTAN , candesartan cilexetil 4 mg tablet, 30
9155W	<i>Deotine 30, SZ</i> – DULOXETINE , duloxetine 30 mg enteric capsule, 28
9156X	<i>Deotine 60, SZ</i> – DULOXETINE , duloxetine 60 mg enteric capsule, 28
1524F	<i>Flubiclox, JU</i> – FLUCLOXACILLIN , flucloxacillin 500 mg injection, 5 vials
1525G	<i>Flubiclox, JU</i> – FLUCLOXACILLIN , flucloxacillin 1 g injection, 5 vials
1525G	<i>Hospira Pty Limited, HH</i> – FLUCLOXACILLIN , flucloxacillin 1 g injection, 5 vials
5094D	<i>Flubiclox, JU</i> – FLUCLOXACILLIN , flucloxacillin 500 mg injection, 5 vials (Dental)
5095E	<i>Flubiclox, JU</i> – FLUCLOXACILLIN , flucloxacillin 1 g injection, 5 vials (Dental)
5095E	<i>Hospira Pty Limited, HH</i> – FLUCLOXACILLIN , flucloxacillin 1 g injection, 5 vials (Dental)
8400D	<i>Monoplus 10/12.5, BQ</i> – FOSINOPRIL + HYDROCHLOROTHIAZIDE , fosinopril sodium 10 mg + hydrochlorothiazide 12.5 mg tablet, 30
2712R	<i>Camino Pro Restore, QH</i> – GLYCOMACROPEPTIDE AND ESSENTIAL AMINO ACIDS , glycomacropeptide and essential amino acids oral liquid, 12 x 500 mL bottles
1801T	<i>Metformin generichealth, GQ</i> – METFORMIN , metformin hydrochloride 850 mg tablet, 60
2277W	<i>Metronidazole-Clarix, AE</i> – METRONIDAZOLE , metronidazole 500 mg/100 mL (0.5%) injection, 5 x 100 mL bags
2298Y	<i>Metronidazole-Clarix, AE</i> – METRONIDAZOLE , metronidazole 500 mg/100 mL (0.5%) injection, 5 x 100 mL bags (Dental)
2357C	<i>Digestelact, SJ</i> – MILK POWDER LACTOSE MODIFIED PREDIGESTED , milk powder lactose modified predigested powder for oral liquid, 900 g
2358D	<i>Digestelact, SJ</i> – MILK POWDER LACTOSE MODIFIED PREDIGESTED , milk powder lactose modified predigested powder for oral liquid, 900 g
3381Y	<i>Olanzapine RBX ODT, RA</i> – OLANZAPINE , OLANZAPINE Tablet 5 mg (orally disintegrating), 28
3382B	<i>Olanzapine RBX ODT, RA</i> – OLANZAPINE , OLANZAPINE Tablet 10 mg (orally disintegrating), 28
3384D	<i>Chem mart Olanzapine ODT, CH</i> – OLANZAPINE , olanzapine 15 mg tablet, 28
3384D	<i>Terry White Chemists Olanzapine ODT, TW</i> – OLANZAPINE , olanzapine 15 mg tablet, 28
3385E	<i>Chem mart Olanzapine ODT, CH</i> – OLANZAPINE , olanzapine 20 mg tablet, 28
3385E	<i>Terry White Chemists Olanzapine ODT, TW</i> – OLANZAPINE , olanzapine 20 mg tablet, 28
8185T	<i>Olanzapine GH, GQ</i> – OLANZAPINE , olanzapine 5 mg tablet, 28
8186W	<i>Olanzapine GH, GQ</i> – OLANZAPINE , olanzapine 7.5 mg tablet, 28
8187X	<i>Olanzapine GH, GQ</i> – OLANZAPINE , olanzapine 10 mg tablet, 28
9197C	<i>Paroxetine generichealth, GQ</i> – PAROXETINE , paroxetine 20 mg tablet, 30
10622C	<i>Hymenoptera Yellow Jacket Venom, DE</i> – VESPUA SPP VENOM , vespuia spp venom 550 microgram injection [1 vial] (&) inert substance diluent [9 mL vial], 1 pack

1 February 2017

Deletion – Brand

8118G	<i>Kalma 2, AF</i> – ALPRAZOLAM , alprazolam 2 mg tablet, 50
8118G	<i>GenRx Alprazolam, GX</i> – ALPRAZOLAM , alprazolam 2 mg tablet, 50
8118G	<i>Alprax 2, QA</i> – ALPRAZOLAM , alprazolam 2 mg tablet, 50
8289G	<i>Avonex, BD</i> – INTERFERON BETA-1A , interferon beta-1a 6 million units (30 microgram) injection [4 vials] (&) inert substance diluent [4 x 1.1 mL syringes], 1 pack
1574W	<i>Nizoral 2%, JT</i> – KETOCONAZOLE , ketoconazole 2% shampoo, 60 mL
9024Y	<i>Nizoral 2% Cream, JT</i> – KETOCONAZOLE , ketoconazole 2% cream, 30 g
9025B	<i>Nizoral 1%, JT</i> – KETOCONAZOLE , ketoconazole 1% shampoo, 100 mL
9027D	<i>Daktarin, JT</i> – MICONAZOLE , miconazole nitrate 2% cream, 30 g
9028E	<i>Daktarin, JT</i> – MICONAZOLE , miconazole nitrate 2% cream, 70 g
9029F	<i>Daktarin, JT</i> – MICONAZOLE , miconazole nitrate 2% dusting powder, 30 g
9031H	<i>Daktarin Tincture, JT</i> – MICONAZOLE , miconazole 2% solution, 30 mL
10634Q	<i>Pegasys RBV, RO</i> – PEGINTERFERON ALFA-2A (&) RIBAVIRIN , peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [168 tablets], 1 pack
10636T	<i>Pegasys RBV, RO</i> – PEGINTERFERON ALFA-2A (&) RIBAVIRIN , peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [140 tablets], 1 pack

Palliative Care Additions

Addition – Item

10957Q	BUPRENORPHINE , buprenorphine 5 microgram/hour patch, 2 (<i>Norspan</i>)
10948F	BUPRENORPHINE , buprenorphine 10 microgram/hour patch, 2 (<i>Norspan</i>)
10953L	BUPRENORPHINE , buprenorphine 15 microgram/hour patch, 2 (<i>Norspan</i>)
10970J	BUPRENORPHINE , buprenorphine 20 microgram/hour patch, 2 (<i>Norspan</i>)
10964C	BUPRENORPHINE , buprenorphine 25 microgram/hour patch, 2 (<i>Norspan</i>)
10949G	BUPRENORPHINE , buprenorphine 30 microgram/hour patch, 2 (<i>Norspan</i>)
10959T	BUPRENORPHINE , buprenorphine 40 microgram/hour patch, 2 (<i>Norspan</i>)

Deletions

Deletion – Brand

5331N	<i>Microlax, JT</i> – SORBITOL + CITRIC ACID + LAURYL SULFOACETATE SODIUM , sorbitol 3.125 g/5 mL + citrate sodium dihydrate 450 mg/5 mL + lauryl sulfoacetate sodium 45 mg/5 mL enema, 12 x 5 mL
-------	--

Deletion – Equivalence Indicator

5331N	<i>Micolette, AE</i> – SORBITOL + CITRIC ACID + LAURYL SULFOACETATE SODIUM , sorbitol 3.125 g/5 mL + citrate sodium dihydrate 450 mg/5 mL + lauryl sulfoacetate sodium 45 mg/5 mL enema, 12 x 5 mL
-------	---

Highly Specialised Drugs Program (Private Hospital)

Additions

Addition – Item

- 10971K **APOMORPHINE**, apomorphine hydrochloride 50 mg/10 mL injection, 5 x 10 mL syringes (*Movapo PFS*)
- 10956P **OMALIZUMAB**, omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe (*Xolair*)
- 10968G **OMALIZUMAB**, omalizumab 150 mg/mL injection, 1 mL syringe (*Xolair*)

Alterations

Alteration – Restriction

The following items have additions, deletions or alterations to restrictions, notes and/or cautions.

- 10184B **INFLIXIMAB**, infliximab 100 mg injection, 1 vial (*Inflixtra, Remicade*)
- 10398G **VEDOLIZUMAB**, vedolizumab 300 mg injection, 1 vial (*Entyvio*)

Advance Notices

1 February 2017

Deletion – Brand

- 10662E *Pegasys RBV, RO* – **PEGINTERFERON ALFA-2A (&) RIBAVIRIN**, peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [168 tablets], 1 pack
- 10674T *Pegasys RBV, RO* – **PEGINTERFERON ALFA-2A (&) RIBAVIRIN**, peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [140 tablets], 1 pack
- 6392K *Pegasys RBV, RO* – **PEGINTERFERON ALFA-2A (&) RIBAVIRIN**, peginterferon alfa-2a 135 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [168 tablets], 1 pack
- 6394M *Pegasys RBV, RO* – **PEGINTERFERON ALFA-2A (&) RIBAVIRIN**, peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [112 tablets], 1 pack
- 6395N *Pegasys RBV, RO* – **PEGINTERFERON ALFA-2A (&) RIBAVIRIN**, peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [140 tablets], 1 pack
- 6396P *Pegasys RBV, RO* – **PEGINTERFERON ALFA-2A (&) RIBAVIRIN**, peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [168 tablets], 1 pack

Highly Specialised Drugs Program (Public Hospital)

Additions

Addition – Item

- 10950H **APOMORPHINE**, apomorphine hydrochloride 50 mg/10 mL injection, 5 x 10 mL syringes (*Movapo PFS*)
- 10967F **OMALIZUMAB**, omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe (*Xolair*)
- 10973M **OMALIZUMAB**, omalizumab 150 mg/mL injection, 1 mL syringe (*Xolair*)

Alterations

Alteration – Restriction

The following items have additions, deletions or alterations to restrictions, notes and/or cautions.

- 10196P **INFLIXIMAB**, infliximab 100 mg injection, 1 vial (*Inflixtra, Remicade*)
- 10384M **VEDOLIZUMAB**, vedolizumab 300 mg injection, 1 vial (*Entyvio*)

Advance Notices

1 February 2017

Deletion – Brand

10655T	<i>Pegasys RBV, RO</i> – PEGINTERFERON ALFA-2A (&) RIBAVIRIN , peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [168 tablets], 1 pack
10664G	<i>Pegasys RBV, RO</i> – PEGINTERFERON ALFA-2A (&) RIBAVIRIN , peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [140 tablets], 1 pack
9524G	<i>Pegasys RBV, RO</i> – PEGINTERFERON ALFA-2A (&) RIBAVIRIN , peginterferon alfa-2a 135 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [168 tablets], 1 pack
9525H	<i>Pegasys RBV, RO</i> – PEGINTERFERON ALFA-2A (&) RIBAVIRIN , peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [112 tablets], 1 pack
9526J	<i>Pegasys RBV, RO</i> – PEGINTERFERON ALFA-2A (&) RIBAVIRIN , peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [140 tablets], 1 pack
9527K	<i>Pegasys RBV, RO</i> – PEGINTERFERON ALFA-2A (&) RIBAVIRIN , peginterferon alfa-2a 180 microgram/0.5 mL injection [4 x 0.5 mL syringes] (&) ribavirin 200 mg tablet [168 tablets], 1 pack

Highly Specialised Drugs Program (Community Access)

Additions

Addition – Item

10946D	TENOFOVIR + EMTRICITABINE , tenofovir alafenamide 10mg + emtricitabine 200 mg tablet, 30 (<i>Descovy 10/200</i>)
10966E	TENOFOVIR + EMTRICITABINE , tenofovir alafenamide 25mg + emtricitabine 200 mg tablet, 30 (<i>Descovy 25/200</i>)

Addition – Brand

10279B	<i>ENTECAVIR RBX, RA</i> – ENTECAVIR , entecavir monohydrate 500 microgram tablet, 30
10279B	<i>ENTECLUDE, RW</i> – ENTECAVIR , entecavir monohydrate 500 microgram tablet, 30
10279B	<i>Entecavir APOTEX, TX</i> – ENTECAVIR , entecavir monohydrate 500 microgram tablet, 30
10279B	<i>Entecavir Amneal, EA</i> – ENTECAVIR , entecavir monohydrate 500 microgram tablet, 30
10279B	<i>Entecavir Mylan, AF</i> – ENTECAVIR , entecavir monohydrate 500 microgram tablet, 30
10279B	<i>Entecavir Sandoz, SZ</i> – ENTECAVIR , entecavir monohydrate 500 microgram tablet, 30
10353X	<i>ENTECAVIR RBX, RA</i> – ENTECAVIR , entecavir monohydrate 1 mg tablet, 30
10353X	<i>ENTECLUDE, RW</i> – ENTECAVIR , entecavir monohydrate 1 mg tablet, 30
10353X	<i>Entecavir APOTEX, TX</i> – ENTECAVIR , entecavir monohydrate 1 mg tablet, 30
10353X	<i>Entecavir Amneal, EA</i> – ENTECAVIR , entecavir monohydrate 1 mg tablet, 30
10353X	<i>Entecavir Mylan, AF</i> – ENTECAVIR , entecavir monohydrate 1 mg tablet, 30
10353X	<i>Entecavir Sandoz, SZ</i> – ENTECAVIR , entecavir monohydrate 1 mg tablet, 30

Addition – Equivalence Indicator

10279B	<i>Baraclude, BQ</i> – ENTECAVIR , entecavir monohydrate 500 microgram tablet, 30
10353X	<i>Baraclude, BQ</i> – ENTECAVIR , entecavir monohydrate 1 mg tablet, 30

Advance Notices

1 January 2017

Deletion – Brand

- 10350R *Videx EC, BQ* – **DIDANOSINE**, didanosine 125 mg enteric capsule, 30
10351T *Videx EC, BQ* – **DIDANOSINE**, didanosine 200 mg enteric capsule, 30

Growth Hormone Program

Advance Notices

1 January 2017

Deletion – Brand

- 10433D *Saizen 8 mg click.easy, SG* – **SOMATROPIN**, SOMATROPIN (Recombinant human growth hormone) Injection 8 mg (24 i.u.) vial with 1.37 mL diluent cartridge (with preservative) (for use with one.click auto-injector), 1
10471D *Saizen 8 mg click.easy, SG* – **SOMATROPIN**, SOMATROPIN (Recombinant human growth hormone) Injection 8 mg (24 i.u.) vial with 1.37 mL diluent cartridge (with preservative) (for use with one.click auto-injector), 1
6329D *Saizen 8 mg click.easy, SG* – **SOMATROPIN**, SOMATROPIN (Recombinant human growth hormone) Injection 8 mg (24 i.u.) vial with 1.37 mL diluent cartridge (with preservative) (for use with one.click auto-injector), 1

IVF Program

Advance Notices

1 January 2017

Deletion – Brand

- 9631X *Ovidrel, SG* – **CHORIOGONADOTROPIN ALFA**, choriogonadotropin alfa 250 microgram/0.5 mL injection, 0.5 mL syringe

Repatriation Pharmaceutical Benefits

Deletions

Deletion – Item

- 4755G **DRESSING NON ADHERENT**, dressing non adherent 5 cm x 7.5 cm dressing, 10 (*Telfa 1970C*)
4758K **DRESSING NON ADHERENT**, dressing non adherent 7.5 cm x 10 cm dressing, 6 (*Telfa 2140C*)
4844Y **DRESSING NON ADHERENT**, dressing non adherent 7.5 cm x 10 cm dressing, 6 (*Telfa 7650C*)
4008Y **KETOCONAZOLE**, ketoconazole 2% shampoo, 60 mL (*Nizoral 2%*)
4454K **MICONAZOLE**, miconazole nitrate 2% cream, 30 g (*Daktarin*)
4341L **MICONAZOLE**, miconazole 2% solution, 30 mL (*Daktarin Tincture*)

Alterations

Alteration – Manufacturer Code

		<i>From</i>	<i>To</i>
4115N	<i>Azithromycin-GA</i> – AZITHROMYCIN , azithromycin 500 mg tablet, 3	UA	EA
4049D	<i>Uracol</i> – BICARBONATE + CITRIC ACID + TARTARIC ACID , sodium bicarbonate 1.76 g + sodium citrate sodium 630 mg + citric acid 720 mg + tartaric acid 890 mg powder for oral liquid, 28 x 4 g sachets	GN	EA
4028B	<i>Soflax</i> – DOCUSATE + SENNOSIDE B , docusate sodium 50 mg + sennoside B 8 mg tablet, 100	GN	EA
4007X	<i>Sebizole</i> – KETOCONAZOLE , ketoconazole 2% shampoo, 100 mL	GN	EA
4308R	<i>Xylocaine Viscous</i> – LIGNOCAINE , lignocaine hydrochloride anhydrous 2% oral liquid, 200 mL	AP	QA
4584G	<i>Sildenafil Actavis</i> – SILDENAFIL , sildenafil 25 mg tablet, 4	UA	EA
4585H	<i>Sildenafil Actavis</i> – SILDENAFIL , sildenafil 50 mg tablet, 4	UA	EA
4586J	<i>Sildenafil Actavis</i> – SILDENAFIL , sildenafil 100 mg tablet, 4	UA	EA
4473K	<i>Lamisil</i> – TERBINAFINE , terbinafine hydrochloride 1% cream, 15 g	NC	GK

General Pharmaceutical Benefits

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of infliximab, vedolizumab and adalimumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either infliximab, vedolizumab or adalimumab at any one time. From 1 December 2016, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial each of PBS-subsidised infliximab, vedolizumab or adalimumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with infliximab, vedolizumab or adalimumab while they continue to show a response to therapy. A patient who received PBS-subsidised infliximab, vedolizumab or adalimumab treatment prior to 1 December 2016 is considered to be in their first cycle as of 1 December 2016. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised infliximab, vedolizumab or adalimumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised infliximab, vedolizumab or adalimumab treatment in the most recent cycle to the date of the first application for initial treatment with infliximab, vedolizumab or adalimumab under the new treatment cycle. A patient who has failed fewer than 3 trials of either infliximab, vedolizumab or adalimumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with infliximab, vedolizumab and adalimumab after therapy after 1 December 2016 .

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) an adult patient has received no prior PBS-subsidised treatment with infliximab, vedolizumab or adalimumab in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) an adult patient has received prior PBS-subsidised (initial or continuing) infliximab, vedolizumab or adalimumab therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) an adult patient wishes to re-commence treatment with infliximab, vedolizumab or adalimumab following a break in PBS-subsidised therapy with the same agent (Initial 2).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab , 14 weeks of therapy for infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab, and this assessment must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist. For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is provided to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with infliximab, vedolizumab or adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment. Assessments of response to a course of PBS-subsidised therapy must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate infliximab, vedolizumab or adalimumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with infliximab, vedolizumab or adalimumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction. To avoid confusion, an application for a patient who wishes to swap to the alternate infliximab, vedolizumab or adalimumab therapy should be accompanied by the approved authority prescription or remaining repeats for

the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Mayo clinic score or partial Mayo clinic score submitted with the first authority application for infliximab, vedolizumab or adalimumab. However, prescribers may provide new baseline measurements any time other than when an initial treatment authority application is provided within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised infliximab, vedolizumab or adalimumab therapy of at least 5 years, must requalify for initial treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the Mayo score is measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for moderate to severe ulcerative colitis prior to 1 December 2016 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a continuing patient.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg (for a child, 1 to 2 mg/kg up to 40 mg) prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score); OR
- Patient must have a Paediatric Ulcerative Colitis Activity Index (PUCAI) Score greater than or equal to 30 if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Applications for authorisation of initial treatment must be in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and

(iii) the signed patient acknowledgement or guardian acknowledgement.

For patients weighing 40 kg or greater, a maximum quantity and number of repeats to provide for an initial 16 weeks course of this drug consisting of a 160 mg dose at week 0, 80 mg dose at week 2 and 40 mg dose at weeks 4, 6, 8, 10, 12 and 14 will be authorised.

For patients weighing less than 40 kg, a maximum quantity and number of repeats to provide for an initial 16 weeks of this drug consisting of a 80 mg dose at week 0, 40 mg dose at week 2 and a 20 mg dose at weeks 4, 6, 8, 10, 12 and 14 will be authorised.

Two completed authority prescriptions must be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription must be written for 2 doses of 40 mg and 2 repeats.

For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic, partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) score must be no more than 1 month old at the time of application.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 within the first 12 weeks of receiving this drug for ulcerative colitis, or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or have failed to maintain a PUCAI score less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated.

The patient or guardian (required if patient is aged 6 to 17 years) must have signed a patient acknowledgement indicating that he or she understands and acknowledges that the PBS-subsidised treatment will cease if he or she does not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Note At the time of the authority application, medical practitioners should request sufficient quantity for up to 16 weeks of treatment under this restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Change or Re-commencement of treatment after a break of less than 5 years in therapy (Initial 2)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with infliximab, vedolizumab or adalimumab for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised treatment with adalimumab for this condition in the current treatment cycle.

Population criteria:

- Patient must be 6 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of this drug within the timelines specified in the relevant restriction. If the response assessment to the previous course of this drug is not submitted as detailed in the relevant restriction, the patient will be deemed to have failed therapy with this drug.

Applications for authorisation of initial treatment must be in writing and must include: (a) a completed authority prescription form; and (b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following: (i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition.

For patients weighing 40 kg or greater, a maximum quantity and number of repeats to provide for an initial 16 weeks course of this drug consisting of a 160 mg dose at week 0, 80 mg dose at week 2 and 40 mg dose at weeks 4, 6, 8, 10, 12 and 14 will be authorised.

For patients weighing less than 40 kg, a maximum quantity and number of repeats to provide for an initial 16 weeks of this drug consisting of a 80 mg dose at week 0, 40 mg dose at week 2 and a 20 mg dose at weeks 4, 6, 8, 10, 12 and 14 will be authorised.

Two completed authority prescriptions must be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription must be written for 2 doses of 40 mg and 2 repeats.

For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Note At the time of the authority application, medical practitioners should request sufficient quantity for up to 16 weeks of treatment under this restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No applications for increased repeats will be authorised.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply for Initial 1 and Initial 2

Clinical criteria:

- Patient must have received insufficient treatment with this drug under the Initial 1 (new patient or recommencement of treatment after more than 5 years break in therapy) restriction to complete 16 weeks of treatment; OR
- Patient must have received insufficient treatment with this drug under the Initial 2 (Change or Re-commencing of treatment after less than 5 years break in therapy) to complete 16 weeks of treatment.

Population criteria:

- Patient must be 6 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Authority approval for sufficient therapy to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting the Department of Human Services.

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL cartridges

10955N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1508.65	38.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

10944B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1508.65	38.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL cartridges

10945C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	4298.62	38.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL syringes

10972L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	4298.62	38.30	Humira [VE]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of infliximab, vedolizumab and adalimumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either infliximab, vedolizumab or adalimumab at any one time. From 1 December 2016, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial each of PBS-subsidised infliximab, vedolizumab or adalimumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with infliximab, vedolizumab or adalimumab while they continue to show a response to therapy. A patient who received PBS-subsidised infliximab, vedolizumab or adalimumab treatment prior to 1 December 2016 is considered to be in their first cycle as of 1 December 2016. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised infliximab, vedolizumab or adalimumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised infliximab, vedolizumab or adalimumab treatment in the most recent cycle to the date of the first application for initial treatment with infliximab, vedolizumab or adalimumab under the new treatment cycle. A patient who has failed fewer than 3 trials of either infliximab, vedolizumab or adalimumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with infliximab, vedolizumab and adalimumab after therapy after 1 December 2016 .

(a) Initial treatment. Applications for initial treatment should be made where:

(i) an adult patient has received no prior PBS-subsidised treatment with infliximab, vedolizumab or adalimumab in this treatment cycle and wishes to commence such therapy (Initial 1); or

- (ii) an adult patient has received prior PBS-subsidised (initial or continuing) infliximab, vedolizumab or adalimumab therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) an adult patient wishes to re-commence treatment with infliximab, vedolizumab or adalimumab following a break in PBS-subsidised therapy with the same agent (Initial 2).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab, and this assessment must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist. For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is provided to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with infliximab, vedolizumab or adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment. Assessments of response to a course of PBS-subsidised therapy must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate infliximab, vedolizumab or adalimumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with infliximab, vedolizumab or adalimumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction. To avoid confusion, an application for a patient who wishes to swap to the alternate infliximab, vedolizumab or adalimumab therapy should be accompanied by the approved authority prescription or remaining repeats for the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Mayo clinic score or partial Mayo clinic score submitted with the first authority application for infliximab, vedolizumab or adalimumab. However, prescribers may provide new baseline measurements any time other than when an initial treatment authority application is provided within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised infliximab, vedolizumab or adalimumab therapy of at least 5 years, must requalify for initial treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the Mayo score is measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for moderate to severe ulcerative colitis prior to 1 December 2016 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a continuing patient.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 while receiving treatment with this drug if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Patients who have failed to maintain a partial Mayo clinic score of less than or equal to 2, with no subscore greater than 1, or, patients who have failed to maintain a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

All applications for continuing treatment with this drug must include a measurement of response to the most recent course of PBS-subsidised therapy. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with this drug, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

Where a response assessment is not submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

At the time of the authority application, medical practitioners should request sufficient quantity for up to 24 weeks of treatment under this restriction.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No applications for increased repeats will be authorised.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial 3 (Grandfathered patient)

Clinical criteria:

- Patient must have a documented history of moderate to severe ulcerative colitis, **AND**
- Patient must have previously received non-PBS-subsidised treatment with this drug for this condition prior to 1 December 2016, **AND**
- Patient must be receiving treatment with this drug at the time of application, **AND**
- Patient must have a Mayo score greater than or equal to 6 prior to commencing treatment with this drug; OR
- Patient must have a partial Mayo score is greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo score) prior to commencing treatment with this drug; OR
- Patient must have a Paediatric Ulcerative Colitis Activity Index (PUCAI) Score greater than or equal to 30 prior to commencing treatment with this drug, if aged 6 to 17 years; OR
- Patient must have a documented history of moderate to severe refractory ulcerative colitis prior to having commenced treatment with this drug where a Mayo clinic, partial Mayo clinic or PUCAI baseline assessment is not available, if aged 6 to 17 years, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 while receiving treatment with this drug if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Applications for authorisation of initial treatment must be in writing and must include: (a) a completed authority prescription form; and (b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following: (i) the completed current and baseline Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheets including the dates of assessment of the patient's condition; and (ii) the date of commencement of this drug; and (iii) the signed patient or guardian acknowledgement

The current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) assessment must be no more than 1 month old at the time of application. The baseline assessment must be from immediately prior to commencing treatment with this drug. Where a baseline assessment is not available the prescriber must contact the Department of Human Services to discuss.

At the time of the authority application, medical practitioners should request sufficient quantity for up to 24 weeks of treatment under this restriction.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

Note The patient or guardian (required if patient is aged 6 to 17 years) must have signed a patient acknowledgement indicating that he or she understands and acknowledges that the PBS-subsidised treatment will cease if he or she does not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply for Continuing treatment and Initial 3 (Grandfathered patients)

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment; OR
- Patient must have received insufficient treatment with this drug to complete 24 weeks of treatment under the Initial 3 (Grandfathered patients).

Population criteria:

- Patient must be 6 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services.

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL cartridges

10961X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1508.65	38.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

10960W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1508.65	38.30	Humira [VE]

■ AFLIBERCEPT

Note Special Pricing Arrangements apply.

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be due to age-related macular degeneration (AMD), **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated by an ophthalmologist.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing or by telephone.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

A telephone application must be made following submission by facsimile of a copy of a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form and a copy of the optical coherence tomography or fluorescein angiogram report. The original documentation must be submitted to the Chief Executive Medicare by post after the application has been authorised.

Note The first authority application may be faxed to the Department of Human Services on 1300 093 177 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). The Department will then contact the prescriber by telephone.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be due to age-related macular degeneration (AMD), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously been granted an authority prescription for the same eye.

Treatment criteria:

- Must be treated by an ophthalmologist.

Note Authority approvals will be administered by the Complex Drugs Unit of the Department of Human Services.

Note Authority applications for continuing treatment in the same eye may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required

Branch retinal vein occlusion with macular oedema

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have visual impairment due to macular oedema secondary to branched retinal vein occlusion (BRVO), **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 73 and 20 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/40 to 20/400), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated by an ophthalmologist.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing or by telephone.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

A telephone application must be made following submission by facsimile of a copy of a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form and a copy of the optical coherence tomography or fluorescein angiogram report. The original documentation must be submitted to the Chief Executive Medicare by post after the application has been authorised.

Note The first authority application may be faxed to the Department of Human Services on 1300 093 177 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). The Department will then contact the prescriber by telephone.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Branch retinal vein occlusion with macular oedema

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated by an ophthalmologist.

Note Authority applications for continuing treatment in the same eye may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Central retinal vein occlusion with macular oedema

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have visual impairment due to macular oedema secondary to central retinal vein occlusion (CRVO), **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 73 and 24 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/40 to 20/320), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated by an ophthalmologist.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing or by telephone.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

A telephone application must be made following submission by facsimile of a copy of a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form and a copy of the optical coherence tomography or fluorescein angiogram report. The original documentation must be submitted to the Chief Executive Medicare by post after the application has been authorised.

Note The first authority application may be faxed to the Department of Human Services on 1300 093 177 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). The Department will then contact the prescriber by telephone.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Central retinal vein occlusion with macular oedema

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated by an ophthalmologist.

Note Authority applications for continuing treatment in the same eye may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

aflibercept 4 mg/0.1 mL injection, 0.1 mL vial

2168D

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	2	..	1148.91	38.30	Eylea [BN]

▪ **EVOLUCUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Familial homozygous hypercholesterolaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- The condition must have been confirmed by genetic testing; OR
- The condition must have been confirmed by a Dutch Lipid Clinic Network Score of at least 7, **AND**
- Patient must have an LDL cholesterol level in excess of 3.3 millimoles per litre after at least 3 months of treatment at a maximum tolerated dose of an HMG CoA reductase inhibitor (statin), in conjunction with dietary therapy and exercise; OR

- Patient must have an LDL cholesterol level in excess of 3.3 millimoles per litre after having developed a clinically important product-related adverse event during treatment with an HMG CoA reductase inhibitor (statin) necessitating a withdrawal of statin treatment; OR
- Patient must have an LDL cholesterol level in excess of 3.3 millimoles per litre and must be one in whom treatment with an HMG CoA reductase inhibitor (statin) is contraindicated.

Treatment criteria:

- Must be treated by a consultant physician or in consultation with a consultant physician.

A clinically important product-related adverse event is defined as follows:

- (i) Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or
- (ii) Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- (iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

The date of the consultation with a consultant physician must be no more than 6 months prior to the application for a PBS authority. The full name of the consultant physician consulted and the date of consultation are to be provided at the time of application.

The qualifying LDL cholesterol level must be provided at the time of application and must be no more than 2 months old. With the exception of patients contraindicated to a statin, the agent, dose and duration of statin treatment must be provided at the time of application.

The authority application must be made in writing and must include:

- a) A completed authority prescription form; and
- b) A completed Familial homozygous hypercholesterolaemia Initial PBS Authority Application - Supporting Information Form; and
- c) The date of consultation and the full name of the consultant physician; and
- d) A copy of the qualifying Dutch Lipid Clinic Network Score or a copy of the result of genetic testing; and
- e) The result of LDL cholesterol level and one of the following where appropriate: statin treatment details including agent, dose and treatment duration; or details of adverse event or contraindication to treatment with a statin as defined in the TGA-approved Product Information.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Familial homozygous hypercholesterolaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in conjunction with dietary therapy and exercise.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Familial homozygous hypercholesterolaemia

Treatment Phase: Grandfathering treatment

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment with this drug for this condition prior to 1 December 2016, **AND**
- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- The condition must have been confirmed by genetic testing; OR
- The condition must have been confirmed by a Dutch Lipid Clinic Network Score of at least 7, **AND**
- Patient must have had an LDL cholesterol level in excess of 3.3 millimoles per litre after at least 3 months of treatment at a maximum tolerated dose of an HMG CoA reductase inhibitor (statin), in conjunction with dietary therapy and exercise, prior to initiation of treatment with this drug; OR
- Patient must have had an LDL cholesterol level in excess of 3.3 millimoles per litre after having developed a clinically important product-related adverse event during treatment with an HMG CoA reductase inhibitor (statin) necessitating a withdrawal of statin treatment, prior to initiation of treatment with this drug; OR
- Patient must have had an LDL cholesterol level in excess of 3.3 millimoles per litre prior to initiation of treatment with this drug, and must be contraindicated to treatment with an HMG CoA reductase inhibitor (statin).

Treatment criteria:

- Must be treated by a consultant physician or in consultation with a consultant physician.

The date of the consultation with a consultant physician must be no more than 6 months prior to the application for a PBS authority. The full name of the consultant physician consulted and the date of consultation are to be provided at the time of application.

The qualifying LDL cholesterol level prior to initiation of treatment with this drug must be provided at the time of application. With the exception of patients contraindicated to a statin, the agent, dose and duration of statin treatment must be provided at the time of application.

A clinically important product-related adverse event is defined as follows:

- (i) Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or
- (ii) Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- (iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

The authority application must be made in writing and must include:

- a) A completed authority prescription form; and
- b) A completed Familial homozygous hypercholesterolaemia Initial PBS 'Grandfather' Authority Application - Supporting Information Form; and
- c) The date of consultation and the full name of the consultant physician; and
- d) A copy of the qualifying Dutch Lipid Clinic Network Score or a copy of the result of genetic testing; and
- e) The result of LDL cholesterol level and one of the following where appropriate: statin treatment details including agent, dose and treatment duration; or details of adverse event or contraindication to treatment with a statin as defined in the TGA-approved Product Information.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

evolocumab 140 mg/mL injection, 1 mL injection device

10958R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*942.63	38.30	Repatha [AN]

FLUTAMIDE

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6611

Metastatic (stage D) carcinoma of the prostate

Clinical criteria:

- The treatment must be in combination with GnRH (LH-RH) analogue therapy.

flutamide 250 mg tablet, 30

2742H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*153.78	38.30	Flutamide MYLAN [AF]

LENVATINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

6604

Locally advanced or metastatic differentiated thyroid cancer

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be refractory to radioactive iodine, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have symptomatic progressive disease prior to treatment; OR
- Patient must have progressive disease at critical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures, **AND**
- Patient must have thyroid stimulating hormone adequately repressed, **AND**
- Patient must be one in whom surgery is inappropriate, **AND**
- Patient must not be a candidate for radiotherapy with curative intent, **AND**

- Patient must have a WHO performance status of 2 or less.
- Radioactive iodine refractory is defined as:
- a lesion without iodine uptake on a radioactive iodine (RAI) scan; or
 - having received a cumulative RAI dose of greater than or equal to 600 mCi; or
 - progression within 12 months of a single RAI treatment; or
 - progression after two RAI treatments administered within 12 months of each other.

Authority required (STREAMLINED)

6578

Locally advanced or metastatic differentiated thyroid cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be refractory to radioactive iodine, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST).

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Authority required (STREAMLINED)

6577

Locally advanced or metastatic differentiated thyroid cancer

Treatment Phase: Grandfathering treatment

Clinical criteria:

- The condition must be refractory to radioactive iodine, **AND**
- Patient must have previously received non-PBS-subsidised treatment with this drug for this condition prior to 1 December 2016, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have thyroid stimulating hormone adequately repressed.

lenvatinib 10 mg capsule, 30

10965D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*6457.88	38.30	Lenvima [EI]

lenvatinib 4 mg capsule, 30

10952K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3302.88	38.30	Lenvima [EI]

▪ LEUPRORELIN (&) INERT SUBSTANCE (&) BICALUTAMIDE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Carcinoma of the prostate

Clinical criteria:

- The condition must be metastatic (stage D), **AND**
- Patient must require a combination of an antiandrogen and a GnRH (LH-RH) agonist.

leuprorelin acetate 22.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe] (&) bicalutamide 50 mg tablet [28], 1 pack

10963B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	1122.85	38.30	Bi ELIGARD CP [TL]

leuprorelin acetate 22.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe] (&) bicalutamide 50 mg tablet [84], 1 pack

10969H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	1268.41	38.30	Bi ELIGARD CP [TL]

leuprorelin acetate 7.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe] (&) bicalutamide 50 mg tablet [28], 1 pack

10962Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	5	..	471.68	38.30	Bi ELIGARD CP [TL]

■ OCRIPLASMIN

Note Where authority approval for treatment for both eyes simultaneously is being sought, a maximum quantity of 2 vials may be requested.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Vitreomacular traction syndrome

Clinical criteria:

- Patient must have visual impairment due to vitreomacular traction (VMT) without a full thickness macular hole (FTMH); OR
- Patient must have visual impairment due to vitreomacular traction (VMT) with a full thickness macular hole (FTMH) of a diameter of less than or equal to 400 micrometres, **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score of approximate Snellen equivalent 20/25 or worse in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography, **AND**
- The condition must have a vitreomacular adhesion diameter less than or equal to 1500 micrometres, **AND**
- Patient must not have an epiretinal membrane attached to the vitreomacular traction, **AND**
- The condition must be previously untreated in the eye proposed for treatment, **AND**
- Patient must not have received prior vitrectomy in the eye proposed for treatment, **AND**
- Patient must be symptomatic.

Treatment criteria:

- Must be treated by an ophthalmologist.

The prescriber must state which eye(s) is being treated at the time of application.

ocriplasmin 500 microgram/0.2 mL injection, 0.2 mL vial

10947E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	4147.88	38.30	Jetrea [NV]

■ RANIBIZUMAB

Note Special Pricing Arrangements apply.

Note Pharmaceutical benefits that have the form ranibizumab 0.165 mL injection syringe and pharmaceutical benefits that have the form ranibizumab 0.23 mL injection vial are equivalent for the purposes of substitution.

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be due to age-related macular degeneration (AMD), **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated by an ophthalmologist.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing or by telephone.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

A telephone application must be made following submission by facsimile of a copy of a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form and a copy of the optical coherence tomography or fluorescein angiogram report. The original documentation must be submitted to the Chief Executive Medicare by post after the application has been authorised.

Note The first authority application may be faxed to the Department of Human Services on 1300 093 177 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). The Department will then contact the prescriber by telephone.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be due to age-related macular degeneration (AMD), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously been granted an authority prescription for the same eye.

Treatment criteria:

- Must be treated by an ophthalmologist.

Note Authority applications for continuing treatment in the same eye may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Branch retinal vein occlusion with macular oedema

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have visual impairment due to macular oedema secondary to branched retinal vein occlusion (BRVO), **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 73 and 20 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/40 to 20/400), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated by an ophthalmologist.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing or by telephone.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

A telephone application must be made following submission by facsimile of a copy of a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form and a copy of the optical coherence tomography or fluorescein angiogram report. The original documentation must be submitted to the Chief Executive Medicare by post after the application has been authorised.

Note The first authority application may be faxed to the Department of Human Services on 1300 093 177 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). The Department will then contact the prescriber by telephone.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Branch retinal vein occlusion with macular oedema

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated by an ophthalmologist.

Note Authority applications for continuing treatment in the same eye may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Central retinal vein occlusion with macular oedema

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have visual impairment due to macular oedema secondary to central retinal vein occlusion (CRVO), **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 73 and 24 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/40 to 20/320), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated by an ophthalmologist.
- Authority approval for initial treatment of each eye must be sought.
The first authority application for each eye must be made in writing or by telephone.
A written application must include:

- a) a completed authority prescription form;
- b) a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

A telephone application must be made following submission by facsimile of a copy of a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form and a copy of the optical coherence tomography or fluorescein angiogram report. The original documentation must be submitted to the Chief Executive Medicare by post after the application has been authorised.

Note The first authority application may be faxed to the Department of Human Services on 1300 093 177 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). The Department will then contact the prescriber by telephone.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Central retinal vein occlusion with macular oedema

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated by an ophthalmologist.

Note Authority applications for continuing treatment in the same eye may be made by telephone on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

ranibizumab 1.65 mg/0.165 mL injection, 1 x 0.165 mL syringe

10138N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1148.91	38.30	^a Lucentis [NV]

ranibizumab 2.3 mg/0.23 mL injection, 0.23 mL vial

1382R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1148.91	38.30	^a Lucentis [NV]

■ TOCILIZUMAB

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe active rheumatoid arthritis, **AND**
- Patient must have demonstrated an adequate response to treatment with tocilizumab, **AND**
- Patient must have received tocilizumab as their most recent course of PBS-subsidised biological disease modifying anti-rheumatic drug (bDMARD) treatment, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

For the purposes of this restriction bDMARD means abatacept, adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, rituximab, tocilizumab or tofacitinib.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

All applications for continuing treatment with tocilizumab must include a measurement of response to the prior course of therapy. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with tocilizumab, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with an initial treatment course.

Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab.

If a patient fails to demonstrate a response to treatment with tocilizumab under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying antirheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alpha antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus-associated kinase (JAK) inhibitor (tofacitinib).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,

- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and

- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 24 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 24 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or

(ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 24 months (Initial 1); or

(iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or

(iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab,

etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to the Department of Human Services within 4 weeks.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Abatacept patients:

Patients are eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. For these patients two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the pre-filled syringes, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 24 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

Abatacept patients:

Patients swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

the Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major

joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment – balance of supply

Clinical criteria:

- Patient must have received insufficient tocilizumab therapy under the Continuing Treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

10954M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	952.70	38.30	Actemra Subcutaneous Injection [RO]

▪ **TOCILIZUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient or patient recommencing treatment after break of more than 24 months)

Clinical criteria:

- Patient must have severe active rheumatoid arthritis, **AND**
- Patient must have received no PBS-subsidised treatment with a biological disease modifying anti-rheumatic drug (bDMARD) for this condition in the previous 24 months, **AND**
- Patient must not have failed previous PBS-subsidised treatment with tocilizumab for this condition, and have not already failed, or ceased to respond to, PBS-subsidised bDMARD treatment for this condition 5 times, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

For the purposes of this restriction bDMARD means abatacept, adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, rituximab or tocilizumab.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The authority application must be made in writing and must include: (1) completed authority prescription form(s); and (2) a completed Rheumatoid Arthritis initial PBS Authority Application - Supporting Information Form; and (3) a signed patient acknowledgement. If a patient fails to demonstrate a response to treatment with tocilizumab under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application: an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active joints from the following list of major joints: (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth). The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied. Where the baseline joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP is provided with the initial application, the same marker will be used to determine response.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying antirheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alpha antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus-associated kinase (JAK) inhibitor (tofacitinib).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 24 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 24 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or

-
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 24 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to the Department of Human Services within 4 weeks.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Abatacept patients:

Patients are eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. For these patients two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the pre-filled syringes, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient has had a break in therapy of more than 24 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

Abatacept patients:

Patients swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the

baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after break of less than 24 months)

Clinical criteria:

- Patient must have a documented history of severe active rheumatoid arthritis, **AND**
- Patient must have received prior PBS-subsidised biological disease modifying anti-rheumatic drug (bDMARD) treatment for this condition and are eligible to receive further bDMARD therapy, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

For the purposes of this restriction bDMARD means abatacept, adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, rituximab, tocilizumab or tofacitinib.

The authority application must be made in writing and must include: (a) completed authority prescription form(s); and (b) a completed Rheumatoid Arthritis continuing PBS Authority Application - Supporting Information Form. Applications for a patient who has received PBS-subsidised treatment with tocilizumab and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised tocilizumab treatment, within the timeframes specified below. Where the most recent course of PBS-subsidised tocilizumab treatment was approved under either of the initial 1 or 2 treatment restrictions, the patient must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must be submitted no later than 4 weeks from the date that course was ceased. Where the most recent course of PBS-subsidised tocilizumab treatment was approved under the continuing treatment criteria, the patient must have been assessed for response, and the assessment must be submitted no later than 4 weeks from the date that course was ceased. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab. If a patient fails to demonstrate a response to a treatment with tocilizumab under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. If a patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying antirheumatic drugs (bDMARDs) for adults with severe active rheumatoid arthritis. Where the term bDMARD appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alpha antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus-associated kinase (JAK) inhibitor (tofacitinib).

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological disease modifying anti-rheumatic drugs

at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised bDMARD therapy may swap to an alternate bDMARD without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised bDMARD while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised bDMARD more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised bDMARDs for the treatment of rheumatoid arthritis.

For patients who have failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has a break in therapy of less than 24 months may commence a further course of treatment with a bDMARD without having to requalify under the Initial 1 treatment restriction. A patient who has failed fewer than 5 bDMARDs and who has had a break in therapy of longer than 24 months must requalify for treatment under the Initial 1 treatment restriction.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised bDMARD treatment is stopped to the date of the new application for treatment with a bDMARD.

(1) How to prescribe PBS-subsidised bDMARD therapy after 1 August 2010.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised bDMARD treatment and wishes to commence such therapy, excluding rituximab (Initial 1); or
- (ii) a patient wishes to re-commence treatment with a bDMARD following a break in PBS-subsidised therapy of more than 24 months (Initial 1); or
- (iii) a patient has received prior PBS-subsidised (initial or continuing) bDMARD therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iv) a patient wishes to re-commence treatment with a specific bDMARD following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2).

Initial applications for new or re-commencing patients (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

A patient must be assessed for response to any course of initial PBS-subsidised treatment (excluding rituximab) following a minimum of 12 weeks of therapy and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Rituximab patients must be assessed following a minimum of 12 weeks after the first infusion, and this assessment must be submitted to the Department of Human Services within 4 weeks.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

For second and subsequent courses of PBS-subsidised bDMARD (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Abatacept patients:

Patients are eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. For these patients two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the pre-filled syringes, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services 24 weeks after the first infusion. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific bDMARD (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing bDMARD treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted bDMARD supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMARD.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMARD is approved, a patient may swap to an alternate bDMARD without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-bDMARD therapy requirements, except if the patient

has had a break in therapy of more than 24 months. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each bDMARD trialled.

Patients who are not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent.

A patient may trial an alternate bDMARD at any time, regardless of whether they are receiving therapy (initial or continuing) with a bDMARD at the time of the application. However, they cannot swap to a particular bDMARD if they have failed to respond to prior treatment with that drug.

Abatacept patients:

Patients swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological agent during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate bDMARD. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate bDMARD may do so without having to have any treatment-free period.

To avoid confusion, an application for a patient who wishes to swap to an alternate bDMARD should be accompanied by the approved authority prescription or remaining repeats for the bDMARD the patient is ceasing.

(3) Baseline measurements to determine response.

the Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a bDMARD. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Except as specified under the Initial 1 treatment restriction, a baseline joint count and ESR and/or CRP should be performed whilst the patient is still on treatment or within 1 month of ceasing prior treatment. Applications under the Initial 1 treatment restriction for new or re-commencing patients must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient or patient recommencing treatment after a break of more than 24 months) or Initial 2 (change or recommencement of treatment after break of less than 24 months) – balance of supply

Clinical criteria:

- Patient must have received insufficient tocilizumab therapy under the Initial 1 (new patient or patient recommencing treatment after break of more than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient tocilizumab therapy under the Initial 2 (change or recommencement of treatment after break of less than 24 months) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Note Authority approval for sufficient therapy to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

10951J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	952.70	38.30	Actemra Subcutaneous Injection [RO]

▪ USTEKINUMAB

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (initial PBS-subsidised supply for continuing treatment in a patient commenced on non-PBS-subsidised therapy)

Clinical criteria:

- Patient must have a documented history of severe active psoriatic arthritis, **AND**
- Patient must have been receiving treatment with this drug for this condition prior to 1 May 2016, **AND**
- Patient must be receiving treatment with this drug for this condition at the time of application, **AND**

- Patient must have demonstrated a response to treatment as specified in the criteria for continuing PBS-subsidised treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be an adult.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

For the purposes of this restriction 'biological agent' means adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab or ustekinumab.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form; and
- (3) a signed patient acknowledgement.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

Note The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to the Department of Human Services no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Note Patients who fail to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this Treatment Cycle. Patients may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe active psoriatic arthritis, **AND**
- Patient must have received this drug as their most recent course of PBS-subsidised treatment with a biological agent for this condition in the current Treatment Cycle, **AND**
- Patient must demonstrate, at the time of application, an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be an adult.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

For the purposes of this restriction 'biological agent' means adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab or ustekinumab.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

All applications for continuing treatment with this drug must include a measurement of response to the most recent course of PBS-subsidised therapy. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with this drug, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

Where a response assessment is not submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Note Patients who fail to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this Treatment Cycle. Patients may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time.

Where the term 'biological agents' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab only.

Patients receiving PBS-subsidised treatment for psoriatic arthritis are able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle. There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 1 or Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and secukinumab, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients - ustekinumab and secukinumab only.

For patients who commenced treatment with ustekinumab for psoriatic arthritis prior to 1 May 2016 and for patients who commenced treatment with secukinumab for psoriatic arthritis prior to 1 October 2016, applications for initial PBS-subsidised treatment as continuing therapy may be made under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Within a Treatment Cycle patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the

PBS; and

(iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment application is submitted within a treatment Cycle and these revised baseline measurements will be used to assess response.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial 3 (initial PBS-subsidised supply for continuing treatment in a patient commenced on non-PBS-subsidised therapy) or Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 3 (initial PBS-subsidised supply for continuing treatment in a patient commenced on non-PBS-subsidised therapy) restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

No increase in the maximum number of repeats may be authorised.

Special Pricing Arrangements apply.

ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial

10767Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	4380.19	38.30	Stelara [JC]

Palliative Care

■ BUPRENORPHINE

Caution The risk of drug dependence is high.

Note Telephone approvals are limited to 1 month's therapy.

Authority required

Chronic severe disabling pain

Clinical criteria:

- Patient must be receiving palliative care, **AND**
- The condition must be unresponsive to non-opioid analgesics.

buprenorphine 10 microgram/hour patch, 2

10948F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*65.79	38.30	Norspan [MF]

buprenorphine 15 microgram/hour patch, 2

10953L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*79.97	38.30	Norspan [MF]

buprenorphine 20 microgram/hour patch, 2

10970J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*94.17	38.30	Norspan [MF]

buprenorphine 25 microgram/hour patch, 2

10964C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*107.61	38.30	Norspan [MF]

buprenorphine 30 microgram/hour patch, 2

10949G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*121.05	38.30	Norspan [MF]

buprenorphine 40 microgram/hour patch, 2

10959T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*147.93	38.30	Norspan [MF]

buprenorphine 5 microgram/hour patch, 2

10957Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*42.55	38.30	Norspan [MF]

Highly Specialised Drugs Program (Private Hospital)

■ APOMORPHINE

Authority required

Parkinson disease

Clinical criteria:

- Patient must have experienced severely disabling motor fluctuations which have not responded to other therapy.

apomorphine hydrochloride 50 mg/10 mL injection, 5 x 10 mL syringes

10971K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	36	5	..	*8215.02	Movapo PFS [TD]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of infliximab, vedolizumab and adalimumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either infliximab, vedolizumab or adalimumab at any one time. From 1 December 2016, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial each of PBS-subsidised infliximab, vedolizumab or adalimumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with infliximab, vedolizumab or adalimumab while they continue to show a response to therapy. A patient who received PBS-subsidised infliximab, vedolizumab or adalimumab treatment prior to 1 December 2016 is considered to be in their first cycle as of 1 December 2016. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised infliximab, vedolizumab or adalimumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised infliximab, vedolizumab or adalimumab treatment in the most recent cycle to the date of the first application for initial treatment with infliximab, vedolizumab or adalimumab under the new treatment cycle. A patient who has failed fewer than 3 trials of either infliximab, vedolizumab or adalimumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with infliximab, vedolizumab and adalimumab after therapy after 1 December 2016 .

(a) Initial treatment. Applications for initial treatment should be made where:

- an adult patient has received no prior PBS-subsidised treatment with infliximab, vedolizumab or adalimumab in this treatment cycle and wishes to commence such therapy (Initial 1); or
- an adult patient has received prior PBS-subsidised (initial or continuing) infliximab, vedolizumab or adalimumab therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- an adult patient wishes to re-commence treatment with infliximab, vedolizumab or adalimumab following a break in PBS-subsidised therapy with the same agent (Initial 2).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab , 14 weeks of therapy for infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab, and this assessment must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist. For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is provided to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with infliximab, vedolizumab or adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment. Assessments of response to a course of PBS-subsidised therapy must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate infliximab, vedolizumab or adalimumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with infliximab, vedolizumab or adalimumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction. To avoid confusion, an application for a patient who wishes to swap to the alternate infliximab, vedolizumab or adalimumab therapy should be accompanied by the approved authority prescription or remaining repeats for the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Mayo clinic score or partial Mayo clinic score submitted with the first authority application for infliximab, vedolizumab or adalimumab. However, prescribers may provide new baseline measurements any time other than when an initial treatment authority application is provided within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised infliximab, vedolizumab or adalimumab therapy of at least 5 years, must requalify for initial treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the Mayo score is measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for moderate to severe ulcerative colitis prior to 1 December 2016 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a continuing patient.

Note Special Pricing Arrangements apply.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg (for a child, 1 to 2 mg/kg up to 40 mg) prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score); OR
- Patient must have a Paediatric Ulcerative Colitis Activity Index (PUCAI) Score greater than or equal to 30 if aged 6 to 17 years; OR
- Patient must have previously received induction therapy with this drug for an acute severe episode of ulcerative colitis in the last 4 months and demonstrated an adequate response to induction therapy by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a PUCAI score less than 10 (if aged 6 to 17 years).

Population criteria:

- Patient must be 6 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Applications for authorisation of initial treatment must be in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the signed patient acknowledgement or guardian acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, or to be administered at 8-weekly intervals for patients who have received prior treatment for an acute severe episode, will be authorised.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic, partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) score must be no more than 1 month old at the time of application.

Where treatment for an acute severe episode has occurred, an adequate response to induction therapy needs to be demonstrated by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 (if aged 6 to 17 years), within the first 12 weeks of receiving this drug for acute severe ulcerative colitis.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 within the first 12 weeks of receiving this drug for ulcerative colitis, or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or have failed to maintain a PUCAI score less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose for patients administered doses at weeks 0, 2 and 6 (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

The patient or guardian (required if patient is aged 6 to 17 years) must have signed a patient acknowledgement indicating that he or she understands and acknowledges that the PBS-subsidised treatment will cease if he or she does not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Note At the time of the authority application, medical practitioners should request sufficient quantity for up to 16 weeks of treatment under this restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 while receiving treatment with this drug, if aged 6 to 17 years.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Patients who have failed to maintain a partial Mayo clinic score of less than or equal to 2, with no subscore greater than 1, or, patients who have failed to maintain a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg.

Up to a maximum of 2 repeats will be authorised.

Note No applications for increased repeats will be authorised.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Change or Re-commencement of treatment after a break in therapy (Initial 2)

Clinical criteria:

- Patient must have previously been issued with an authority prescription for adalimumab, infliximab or vedolizumab for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with infliximab for this condition more than once in the current treatment cycle.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of this drug within the timelines specified in the relevant restriction. If the response assessment to the previous course of this drug is not submitted as detailed in the relevant restriction, the patient will be deemed to have failed therapy with this drug.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg.

Up to a maximum of 2 repeats will be authorised.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (Change or Recommencement of treatment after a break in therapy) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 and Initial 2 restrictions) or 2 repeats (Continuing restriction).

Population criteria:

- Patient must be 6 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services.

infliximab 100 mg injection, 1 vial

10184B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	604.86	^a Inflectra [HH]	^a Remicade [JC]

■ OMALIZUMAB

Note Special Pricing Arrangements apply.

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a diagnosis of asthma confirmed and documented by a paediatric respiratory physician, clinical immunologist, or allergist; or paediatrician or general physician experienced in the management of patients with severe asthma in consultation with a respiratory physician, defined by the following standard clinical features: forced expiratory volume (FEV1) reversibility or airway hyperresponsiveness or peak expiratory flow (PEF) variability, **AND**
- Patient must have a duration of asthma of at least 1 year, **AND**
- Patient must have past or current evidence of atopy, documented by skin prick testing or an in vitro measure of specific IgE, **AND**
- Patient must have total serum human immunoglobulin E greater than or equal to 30 IU/mL, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction, **AND**
- Patient must be under the care of the same physician for at least 6 months.

Population criteria:

- Patient must be aged 6 to less than 12 years.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

Optimised asthma therapy includes:

- (i) Adherence to optimal inhaled therapy, including high dose inhaled corticosteroid (ICS) and long-acting beta-2 agonist (LABA) therapy for at least six months. If LABA therapy is contraindicated, not tolerated or not effective, montelukast, cromoglycate or nedocromil may be used as an alternative; **AND**
- (ii) treatment with at least 2 courses of oral or IV corticosteroids (daily or alternate day maintenance treatment courses, or 3-5 day exacerbation treatment courses), in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications (including those specified in the relevant TGA-approved Product Information) and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The initial IgE assessment must be no more than 12 months old at the time of application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

- (a) An Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month (for children aged 6 to 10 years it is recommended that the Interviewer Administered version - the ACQ-IA be used), **AND**
- (b) while receiving optimised asthma therapy in the previous 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician. The Asthma Control Questionnaire (5 item version) or ACQ-IA assessment of the patient's response to this initial course of treatment, the assessment of oral corticosteroid dose, and the assessment of exacerbation rate must be made at around 22 to 26 weeks after the first dose so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with omalizumab.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate maximum quantity and number of repeats to provide for an initial course of omalizumab of up to 28 weeks, consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information) to be administered every 2 or 4 weeks.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Paediatric Severe Allergic Asthma Initial PBS Authority Application - Supporting Information form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (dosage, date of commencement and duration of therapy); and
 - (ii) details of severe exacerbation/s experienced in the past 12 months while receiving optimised asthma therapy (date and treatment); and
 - (iii) acknowledgement signed by a parent or authorised guardian; and

(c) a copy of the IgE pathology report; and

(d) a completed Asthma Control Questionnaire (ACQ-5) or the Asthma Control Questionnaire interviewer administered version (ACQ-IA) calculation sheet including the date of assessment of the patient's symptoms and is endorsed with the prescriber's signature.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note It is recommended that an application for continuing treatment is submitted at the time of the 22 to 26 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised omalizumab treatment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note TREATMENT OF PAEDIATRIC PATIENTS WITH UNCONTROLLED SEVERE ALLERGIC ASTHMA

Patients are eligible to commence an 'omalizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to omalizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised omalizumab therapy before they are eligible to commence the next cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised omalizumab treatment is stopped to the date of the first application for initial treatment with omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised omalizumab therapy.

(a) Initial treatment:

Applications for initial treatment should be made where a patient has received no prior PBS-subsidised omalizumab treatment in this treatment cycle and wishes to commence such therapy.

All applications for initial treatment for non-grandfathered patients will be limited to provide for a maximum of 28 weeks of therapy for omalizumab.

(b) Grandfather patients:

Paediatric patients who commenced treatment with omalizumab for uncontrolled severe allergic asthma prior to 1 December 2016 and who continue to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment with omalizumab will be authorised under this restriction. Approval will be based on the criteria included in the grandfather restriction. Following completion of the Initial PBS-subsidised course, further applications for treatment with omalizumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle (initial treatment course with or without continuing treatment course/s). If a 'Grandfathered' patient recommences on second and subsequent cycles after a treatment break, the 'Grandfathered' patient must re-qualify for Initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 6 month break in PBS-subsidised therapy' below for further details.

(c) Continuing treatment:

Following the completion of the initial treatment course with omalizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with omalizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing omalizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or ACQ-IA, systemic corticosteroid dose and time-adjusted exacerbation rate, submitted with the Initial authority application for omalizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and The Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised omalizumab therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (systemic corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score or ACQ-IA, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

(4) Monitoring of patients:

Anaphylaxis and anaphylactoid reactions have been reported following first or subsequent administration of omalizumab (see Product Information). Patients should be monitored post-injection, and medications for the treatment of anaphylactic reactions should be available for immediate use following administration of omalizumab. Patients should be informed that such reactions are possible and prompt medical attention should be sought if allergic reactions occur.

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma
Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe allergic asthma, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

An adequate response to omalizumab treatment is defined as:

- (a) a reduction in the Asthma Control Questionnaire (ACQ-5) or ACQ-IA score of at least 0.5 from baseline, OR
- (b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ-5 or ACQ-IA score from baseline, OR

- (c) a reduction in the time-adjusted exacerbation rates compared to the 12 months prior to baseline.

All applications for continuing treatment with omalizumab must include a measurement of response to the prior course of therapy. The Asthma Control Questionnaire (5 item version) or Asthma Control Questionnaire interviewer administered version (ACQ-IA) assessment of the patient's response to the prior course of treatment, the assessment of systemic corticosteroid dose, and the assessment of time-adjusted exacerbation rate must be made at around 18 to 22 weeks after the first dose of PBS-subsidised omalizumab so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

The first assessment should, where possible, be completed by the same physician who initiated treatment with omalizumab. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with omalizumab.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats to provide for a continuing course of omalizumab consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information), sufficient for 24 weeks of therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Paediatric Severe Allergic Asthma Continuing PBS Authority Application - Supporting Information form which includes details of maintenance oral corticosteroid dose; and
- (c) a completed Asthma Control Questionnaire (ACQ-5) or the Asthma Control Questionnaire interviewer administered version (ACQ-IA) calculation sheet including the date of assessment of the patient's symptoms and is endorsed with the signature of the prescriber.

Note If the same physician cannot assess the patient please call the Department of Human Services on 1800 700 270.

Note For copies of the ACQ please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note It is recommended that an application for continuing treatment is submitted at the time of the 18 to 22 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised omalizumab treatment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note TREATMENT OF PAEDIATRIC PATIENTS WITH UNCONTROLLED SEVERE ALLERGIC ASTHMA

Patients are eligible to commence an 'omalizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to omalizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised omalizumab therapy before they are eligible to commence the next cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised omalizumab treatment is stopped to the date of the first application for initial treatment with omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

- (1) How to prescribe PBS-subsidised omalizumab therapy.

- (a) Initial treatment:

Applications for initial treatment should be made where a patient has received no prior PBS-subsidised omalizumab treatment in this treatment cycle and wishes to commence such therapy.

All applications for initial treatment for non-grandfathered patients will be limited to provide for a maximum of 28 weeks of therapy for omalizumab.

- (b) Grandfather patients:

Paediatric patients who commenced treatment with omalizumab for uncontrolled severe allergic asthma prior to 1 December 2016 and who continue to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24

weeks of treatment with omalizumab will be authorised under this restriction. Approval will be based on the criteria included in the grandfather restriction. Following completion of the Initial PBS-subsidised course, further applications for treatment with omalizumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle (initial treatment course with or without continuing treatment course/s). If a 'Grandfathered' patient recommences on second and subsequent cycles after a treatment break, the 'Grandfathered' patient must re-qualify for Initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 6 month break in PBS-subsidised therapy' below for further details.

(c) Continuing treatment:

Following the completion of the initial treatment course with omalizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with omalizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing omalizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or ACQ-IA, systemic corticosteroid dose and time-adjusted exacerbation rate, submitted with the Initial authority application for omalizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and The Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised omalizumab therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (systemic corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score or ACQ-IA, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

(4) Monitoring of patients:

Anaphylaxis and anaphylactoid reactions have been reported following first or subsequent administration of omalizumab (see Product Information). Patients should be monitored post-injection, and medications for the treatment of anaphylactic reactions should be available for immediate use following administration of omalizumab. Patients should be informed that such reactions are possible and prompt medical attention should be sought if allergic reactions occur.

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial and continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction or Grandfather treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the Initial restriction or up to 24 weeks treatment available under the Continuing or Grandfather restrictions.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

Note Authority approval for sufficient therapy to complete a maximum of 28 weeks of treatment under the initial restriction or 24 weeks of treatment under the continuing or grandfather restrictions may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial treatment - Grandfather patients

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment with this drug for this condition prior to 1 December 2016, **AND**
- Patient must be receiving treatment with this drug for this condition at the time of application, **AND**
- Patient must have had, prior to commencement of omalizumab, a diagnosis of asthma confirmed and documented by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician, defined by the following standard clinical features: forced expiratory volume (FEV1) reversibility or airway hyperresponsiveness or peak expiratory flow (PEF) variability, **AND**
- Patient must have had a duration of asthma of at least 1 year prior to commencement of omalizumab, **AND**
- Patient must have past or current evidence of atopy, documented by skin prick testing or an in vitro measure of specific IgE, **AND**

- Patient must have failed to achieve adequate control with optimised asthma therapy prior to omalizumab therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- Patient must have demonstrated an adequate response to treatment, **AND**
- Patient must be under the care of the same physician for at least 6 months.

Population criteria:

- Patient must be aged 6 to less than 12 years.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

Optimised asthma therapy includes:

- (i) Adherence to optimal inhaled therapy, including high dose inhaled corticosteroid (ICS) and long-acting beta-2 agonist (LABA) therapy for at least six months. If LABA therapy is contraindicated, not tolerated or not effective, montelukast, cromoglycate or nedocromil may be used as an alternative; **AND**
- (ii) treatment with at least 2 courses of oral or IV corticosteroids (daily or alternate day maintenance treatment courses, or 3-5 day exacerbation treatment courses), in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications (including those specified in the relevant TGA-approved Product Information) and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

A review of the patient's records should be conducted to extract pre- and post-omalizumab data on symptoms, quality of life, medication doses, exacerbations and hospitalisations. Examples of parameters to establish response include:

- (i) a reduction in Asthma Control Questionnaire (ACQ-5) or Asthma Control Questionnaire Interviewer Administered (ACQ-IA) score of at least 0.5;
- (ii) maintenance oral corticosteroid dose reduced by at least 25% from baseline; and/or
- (iii) a reduction in the number of hospitalisations or severe exacerbations requiring use of systemic corticosteroids, compared to the 12 months prior to commencement of omalizumab.

The assessment of the patient's response to the initial PBS subsidised course of treatment must be made at around 18 to 22 weeks after the first dose so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed. The same parameters used to establish response to non-PBS-subsidised therapy with omalizumab should be used for the assessment.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with omalizumab.

Patients will be eligible to receive continuing courses of omalizumab treatment of up to 24 weeks providing they continue to demonstrate an adequate response to treatment.

Patients may qualify for PBS-subsidised treatment under this restriction once only.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats to provide for an initial course of omalizumab of up to 24 weeks, consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information) to be administered every 2 or 4 weeks.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Paediatric Grandfather Severe Allergic Asthma PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (dosage, date of commencement and duration of therapy); and
 - (ii) details of pre- and post-omalizumab data on symptoms, quality of life, medication doses, exacerbations and hospitalisations; and
 - (iii) acknowledgement signed by a parent or authorised guardian.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note It is recommended that an application for continuing treatment is submitted at the time of the 22 to 26 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised omalizumab treatment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note TREATMENT OF PAEDIATRIC PATIENTS WITH UNCONTROLLED SEVERE ALLERGIC ASTHMA

Patients are eligible to commence an 'omalizumab treatment cycle' (initial treatment course with or without continuing

treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction. Once a patient has either failed to achieve or maintain a response to omalizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised omalizumab therapy before they are eligible to commence the next cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised omalizumab treatment is stopped to the date of the first application for initial treatment with omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised omalizumab therapy.

(a) Initial treatment:

Applications for initial treatment should be made where a patient has received no prior PBS-subsidised omalizumab treatment in this treatment cycle and wishes to commence such therapy.

All applications for initial treatment for non-grandfathered patients will be limited to provide for a maximum of 28 weeks of therapy for omalizumab.

(b) Grandfather patients:

Paediatric patients who commenced treatment with omalizumab for uncontrolled severe allergic asthma prior to 1 December 2016 and who continue to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment with omalizumab will be authorised under this restriction. Approval will be based on the criteria included in the grandfather restriction. Following completion of the Initial PBS-subsidised course, further applications for treatment with omalizumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle (initial treatment course with or without continuing treatment course/s). If a 'Grandfathered' patient recommences on second and subsequent cycles after a treatment break, the 'Grandfathered' patient must re-qualify for Initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 6 month break in PBS-subsidised therapy' below for further details.

(c) Continuing treatment:

Following the completion of the initial treatment course with omalizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with omalizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing omalizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or ACQ-IA, systemic corticosteroid dose and time-adjusted exacerbation rate, submitted with the Initial authority application for omalizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and The Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised omalizumab therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (systemic corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score or ACQ-IA, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

(4) Monitoring of patients:

Anaphylaxis and anaphylactoid reactions have been reported following first or subsequent administration of omalizumab (see Product Information). Patients should be monitored post-injection, and medications for the treatment of anaphylactic reactions should be available for immediate use following administration of omalizumab. Patients should be informed that such reactions are possible and prompt medical attention should be sought if allergic reactions occur.

omalizumab 150 mg/mL injection, 1 mL syringe

10968G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	433.42	Xolair [NV]

omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe

10956P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	220.22	Xolair [NV]

▪ VEDOLIZUMAB

Note Special Pricing Arrangements apply.

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of infliximab, vedolizumab and adalimumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either infliximab, vedolizumab or adalimumab at any one time. From 1 December 2016, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial each of PBS-subsidised infliximab, vedolizumab or adalimumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with infliximab, vedolizumab or adalimumab while they continue to show a response to therapy. A patient who received PBS-subsidised infliximab, vedolizumab or adalimumab treatment prior to 1 December 2016 is considered to be in their first cycle as of 1 December 2016. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised infliximab, vedolizumab or adalimumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised infliximab, vedolizumab or adalimumab treatment in the most recent cycle to the date of the first application for initial treatment with infliximab, vedolizumab or adalimumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of either infliximab, vedolizumab or adalimumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with infliximab, vedolizumab and adalimumab after therapy after 1 December 2016 .

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) an adult patient has received no prior PBS-subsidised treatment with infliximab, vedolizumab or adalimumab in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) an adult patient has received prior PBS-subsidised (initial or continuing) infliximab, vedolizumab or adalimumab therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) an adult patient wishes to re-commence treatment with infliximab, vedolizumab or adalimumab following a break in PBS-subsidised therapy with the same agent (Initial 2).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab , 14 weeks of therapy for infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab, and this assessment must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist. For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is provided to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with infliximab, vedolizumab or adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment. Assessments of response to a course of PBS-subsidised therapy must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate infliximab, vedolizumab or adalimumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with infliximab, vedolizumab or adalimumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction. To avoid confusion, an application for a patient who wishes to swap to the alternate infliximab, vedolizumab or adalimumab therapy should be accompanied by the approved authority prescription or remaining repeats for the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Mayo clinic score or partial Mayo clinic score submitted with the first authority application for infliximab, vedolizumab or adalimumab. However, prescribers may provide new baseline measurements any time other than when an initial treatment authority application is provided within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised infliximab, vedolizumab or adalimumab therapy of at least 5 years, must requalify for initial treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the Mayo score is measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for moderate to severe ulcerative colitis prior to 1 December 2016 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a continuing patient.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment (new patient – Initial 1)

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score), **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Applications for authorisation of initial treatment must be in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic calculation sheet including the date of assessment of the patient's condition; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and

(iii) the signed patient acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic or partial Mayo clinic score must be no more than 1 month old at the time of application.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose for patients administered doses at weeks 0, 2 and 6 (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

Patients must have signed a patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Note Details of accepted toxicities including severity can be found on the Department of Human Services website at www.humanservices.gov.au.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR

- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]. Patients who have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug. Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response. At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide for a single infusion of 300 mg per dose. Up to a maximum of 2 repeats will be authorised.

Note No applications for increased repeats will be authorised.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Change or Re-commencement of treatment after a break in therapy (Initial 2)

Clinical criteria:

- Patient must have previously been issued with an authority prescription for adalimumab, infliximab or vedolizumab for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with vedolizumab for this condition more than once in the current treatment cycle, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of this drug within the timelines specified in the relevant restriction. If the response assessment to the previous course of this drug is not submitted as detailed in the relevant restriction, the patient will be deemed to have failed therapy with this drug.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide for a single infusion of 300 mg per dose.

Up to a maximum of 2 repeats will be authorised.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses of treatment may be requested by telephone by contacting the Department of Human Services.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (Change or Recommencement of treatment after a break in therapy) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 and Initial 2 restrictions) or 2 repeats (Continuing restriction), **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services.

vedolizumab 300 mg injection, 1 vial

10398G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3152.21	Entyvio [TK]

Highly Specialised Drugs Program (Public Hospital)

■ APOMORPHINE

Authority required (STREAMLINED)

4833

Parkinson disease

Clinical criteria:

- Patient must have experienced severely disabling motor fluctuations which have not responded to other therapy.

apomorphine hydrochloride 50 mg/10 mL injection, 5 x 10 mL syringes

10950H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	36	5	..	*8168.04	Movapo PFS [TD]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of infliximab, vedolizumab and adalimumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either infliximab, vedolizumab or adalimumab at any one time. From 1 December 2016, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial each of PBS-subsidised infliximab, vedolizumab or adalimumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with infliximab, vedolizumab or adalimumab while they continue to show a response to therapy. A patient who received PBS-subsidised infliximab, vedolizumab or adalimumab treatment prior to 1 December 2016 is considered to be in their first cycle as of 1 December 2016. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised infliximab, vedolizumab or adalimumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised infliximab, vedolizumab or adalimumab treatment in the most recent cycle to the date of the first application for initial treatment with infliximab, vedolizumab or adalimumab under the new treatment cycle. A patient who has failed fewer than 3 trials of either infliximab, vedolizumab or adalimumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with infliximab, vedolizumab and adalimumab after therapy after 1 December 2016 .

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) an adult patient has received no prior PBS-subsidised treatment with infliximab, vedolizumab or adalimumab in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) an adult patient has received prior PBS-subsidised (initial or continuing) infliximab, vedolizumab or adalimumab therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) an adult patient wishes to re-commence treatment with infliximab, vedolizumab or adalimumab following a break in PBS-subsidised therapy with the same agent (Initial 2).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab , 14 weeks of therapy for infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab, and this assessment must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist. For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is provided to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with infliximab, vedolizumab or adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment. Assessments of response to a course of PBS-subsidised therapy must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within

these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate infliximab, vedolizumab or adalimumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with infliximab, vedolizumab or adalimumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction. To avoid confusion, an application for a patient who wishes to swap to the alternate infliximab, vedolizumab or adalimumab therapy should be accompanied by the approved authority prescription or remaining repeats for the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Mayo clinic score or partial Mayo clinic score submitted with the first authority application for infliximab, vedolizumab or adalimumab. However, prescribers may provide new baseline measurements any time other than when an initial treatment authority application is provided within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised infliximab, vedolizumab or adalimumab therapy of at least 5 years, must requalify for initial treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the Mayo score is measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for moderate to severe ulcerative colitis prior to 1 December 2016 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a continuing patient.

Note Special Pricing Arrangements apply.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg (for a child, 1 to 2 mg/kg up to 40 mg) prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score); OR
- Patient must have a Paediatric Ulcerative Colitis Activity Index (PUCAI) Score greater than or equal to 30 if aged 6 to 17 years; OR
- Patient must have previously received induction therapy with this drug for an acute severe episode of ulcerative colitis in the last 4 months and demonstrated an adequate response to induction therapy by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a PUCAI score less than 10 (if aged 6 to 17 years).

Population criteria:

- Patient must be 6 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR

- Must be treated by a specialist paediatric gastroenterologist.

Applications for authorisation of initial treatment must be in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the signed patient acknowledgement or guardian acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, or to be administered at 8-weekly intervals for patients who have received prior treatment for an acute severe episode, will be authorised.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic, partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) score must be no more than 1 month old at the time of application.

Where treatment for an acute severe episode has occurred, an adequate response to induction therapy needs to be demonstrated by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 (if aged 6 to 17 years), within the first 12 weeks of receiving this drug for acute severe ulcerative colitis.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 within the first 12 weeks of receiving this drug for ulcerative colitis, or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or have failed to maintain a PUCAI score less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose for patients administered doses at weeks 0, 2 and 6 (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

The patient or guardian (required if patient is aged 6 to 17 years) must have signed a patient acknowledgement indicating that he or she understands and acknowledges that the PBS-subsidised treatment will cease if he or she does not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Note At the time of the authority application, medical practitioners should request sufficient quantity for up to 16 weeks of treatment under this restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 while receiving treatment with this drug, if aged 6 to 17 years.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Patients who have failed to maintain a partial Mayo clinic score of less than or equal to 2, with no subscore greater than 1, or, patients who have failed to maintain a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg.

Up to a maximum of 2 repeats will be authorised.

Note No applications for increased repeats will be authorised.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Change or Re-commencement of treatment after a break in therapy (Initial 2)

Clinical criteria:

- Patient must have previously been issued with an authority prescription for adalimumab, infliximab or vedolizumab for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with infliximab for this condition more than once in the current treatment cycle.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of this drug within the timelines specified in the relevant restriction. If the response assessment to the previous course of this drug is not submitted as detailed in the relevant restriction, the patient will be deemed to have failed therapy with this drug.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg.

Up to a maximum of 2 repeats will be authorised.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (Change or Recommencement of treatment after a break in therapy) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 and Initial 2 restrictions) or 2 repeats (Continuing restriction).

Population criteria:

- Patient must be 6 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services.

infliximab 100 mg injection, 1 vial

10196P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	574.85	^a Inflectra [HH]	^a Remicade [JC]

■ OMALIZUMAB

Note Special Pricing Arrangements apply.

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a diagnosis of asthma confirmed and documented by a paediatric respiratory physician, clinical immunologist, or allergist; or paediatrician or general physician experienced in the management of patients with severe asthma in consultation with a respiratory physician, defined by the following standard clinical features: forced expiratory volume (FEV1) reversibility or airway hyperresponsiveness or peak expiratory flow (PEF) variability, **AND**
- Patient must have a duration of asthma of at least 1 year, **AND**
- Patient must have past or current evidence of atopy, documented by skin prick testing or an in vitro measure of specific IgE, **AND**
- Patient must have total serum human immunoglobulin E greater than or equal to 30 IU/mL, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction, **AND**
- Patient must be under the care of the same physician for at least 6 months.

Population criteria:

- Patient must be aged 6 to less than 12 years.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

Optimised asthma therapy includes:

- (i) Adherence to optimal inhaled therapy, including high dose inhaled corticosteroid (ICS) and long-acting beta-2 agonist (LABA) therapy for at least six months. If LABA therapy is contraindicated, not tolerated or not effective, montelukast, cromoglycate or nedocromil may be used as an alternative; **AND**
- (ii) treatment with at least 2 courses of oral or IV corticosteroids (daily or alternate day maintenance treatment courses, or 3-5 day exacerbation treatment courses), in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications (including those specified in the relevant TGA-approved Product Information) and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The initial IgE assessment must be no more than 12 months old at the time of application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

- (a) An Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month (for children aged 6 to 10 years it is recommended that the Interviewer Administered version - the ACQ-IA be used), **AND**
- (b) while receiving optimised asthma therapy in the previous 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician. The Asthma Control Questionnaire (5 item version) or ACQ-IA assessment of the patient's response to this initial course of treatment, the assessment of oral corticosteroid dose, and the assessment of exacerbation rate must be made at around 22 to 26 weeks after the first dose so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with omalizumab.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate maximum quantity and number of repeats to provide for an initial course of omalizumab of up to 28 weeks, consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information) to be administered every 2 or 4 weeks.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Paediatric Severe Allergic Asthma Initial PBS Authority Application - Supporting Information form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (dosage, date of commencement and duration of therapy); and
 - (ii) details of severe exacerbation/s experienced in the past 12 months while receiving optimised asthma therapy (date and treatment); and
 - (iii) acknowledgement signed by a parent or authorised guardian; and

(c) a copy of the IgE pathology report; and

(d) a completed Asthma Control Questionnaire (ACQ-5) or the Asthma Control Questionnaire interviewer administered version (ACQ-IA) calculation sheet including the date of assessment of the patient's symptoms and is endorsed with the prescriber's signature.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note It is recommended that an application for continuing treatment is submitted at the time of the 22 to 26 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised omalizumab treatment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note TREATMENT OF PAEDIATRIC PATIENTS WITH UNCONTROLLED SEVERE ALLERGIC ASTHMA

Patients are eligible to commence an 'omalizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to omalizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised omalizumab therapy before they are eligible to commence the next cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised omalizumab treatment is stopped to the date of the first application for initial treatment with omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised omalizumab therapy.

(a) Initial treatment:

Applications for initial treatment should be made where a patient has received no prior PBS-subsidised omalizumab treatment in this treatment cycle and wishes to commence such therapy.

All applications for initial treatment for non-grandfathered patients will be limited to provide for a maximum of 28 weeks of therapy for omalizumab.

(b) Grandfather patients:

Paediatric patients who commenced treatment with omalizumab for uncontrolled severe allergic asthma prior to 1 December 2016 and who continue to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment with omalizumab will be authorised under this restriction. Approval will be based on the criteria included in the grandfather restriction. Following completion of the Initial PBS-subsidised course, further applications for treatment with omalizumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle (initial treatment course with or without continuing treatment course/s). If a 'Grandfathered' patient recommences on second and subsequent cycles after a treatment break, the 'Grandfathered' patient must re-qualify for Initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 6 month break in PBS-subsidised therapy' below for further details.

(c) Continuing treatment:

Following the completion of the initial treatment course with omalizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with omalizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing omalizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or ACQ-IA, systemic corticosteroid dose and time-adjusted exacerbation rate, submitted with the Initial authority application for omalizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and The Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised omalizumab therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (systemic corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score or ACQ-IA, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

(4) Monitoring of patients:

Anaphylaxis and anaphylactoid reactions have been reported following first or subsequent administration of omalizumab (see Product Information). Patients should be monitored post-injection, and medications for the treatment of anaphylactic reactions should be available for immediate use following administration of omalizumab. Patients should be informed that such reactions are possible and prompt medical attention should be sought if allergic reactions occur.

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma
Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe allergic asthma, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

An adequate response to omalizumab treatment is defined as:

- (a) a reduction in the Asthma Control Questionnaire (ACQ-5) or ACQ-IA score of at least 0.5 from baseline, OR
- (b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ-5 or ACQ-IA score from baseline, OR

- (c) a reduction in the time-adjusted exacerbation rates compared to the 12 months prior to baseline.

All applications for continuing treatment with omalizumab must include a measurement of response to the prior course of therapy. The Asthma Control Questionnaire (5 item version) or Asthma Control Questionnaire interviewer administered version (ACQ-IA) assessment of the patient's response to the prior course of treatment, the assessment of systemic corticosteroid dose, and the assessment of time-adjusted exacerbation rate must be made at around 18 to 22 weeks after the first dose of PBS-subsidised omalizumab so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

The first assessment should, where possible, be completed by the same physician who initiated treatment with omalizumab. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with omalizumab.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats to provide for a continuing course of omalizumab consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information), sufficient for 24 weeks of therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Paediatric Severe Allergic Asthma Continuing PBS Authority Application - Supporting Information form which includes details of maintenance oral corticosteroid dose; and
- (c) a completed Asthma Control Questionnaire (ACQ-5) or the Asthma Control Questionnaire interviewer administered version (ACQ-IA) calculation sheet including the date of assessment of the patient's symptoms and is endorsed with the signature of the prescriber.

Note If the same physician cannot assess the patient please call the Department of Human Services on 1800 700 270.

Note For copies of the ACQ please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note It is recommended that an application for continuing treatment is submitted at the time of the 18 to 22 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised omalizumab treatment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note TREATMENT OF PAEDIATRIC PATIENTS WITH UNCONTROLLED SEVERE ALLERGIC ASTHMA

Patients are eligible to commence an 'omalizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to omalizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised omalizumab therapy before they are eligible to commence the next cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised omalizumab treatment is stopped to the date of the first application for initial treatment with omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

- (1) How to prescribe PBS-subsidised omalizumab therapy.

- (a) Initial treatment:

Applications for initial treatment should be made where a patient has received no prior PBS-subsidised omalizumab treatment in this treatment cycle and wishes to commence such therapy.

All applications for initial treatment for non-grandfathered patients will be limited to provide for a maximum of 28 weeks of therapy for omalizumab.

- (b) Grandfather patients:

Paediatric patients who commenced treatment with omalizumab for uncontrolled severe allergic asthma prior to 1 December 2016 and who continue to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24

weeks of treatment with omalizumab will be authorised under this restriction. Approval will be based on the criteria included in the grandfather restriction. Following completion of the Initial PBS-subsidised course, further applications for treatment with omalizumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle (initial treatment course with or without continuing treatment course/s). If a 'Grandfathered' patient recommences on second and subsequent cycles after a treatment break, the 'Grandfathered' patient must re-qualify for Initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 6 month break in PBS-subsidised therapy' below for further details.

(c) Continuing treatment:

Following the completion of the initial treatment course with omalizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with omalizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing omalizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or ACQ-IA, systemic corticosteroid dose and time-adjusted exacerbation rate, submitted with the Initial authority application for omalizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and The Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised omalizumab therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (systemic corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score or ACQ-IA, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

(4) Monitoring of patients:

Anaphylaxis and anaphylactoid reactions have been reported following first or subsequent administration of omalizumab (see Product Information). Patients should be monitored post-injection, and medications for the treatment of anaphylactic reactions should be available for immediate use following administration of omalizumab. Patients should be informed that such reactions are possible and prompt medical attention should be sought if allergic reactions occur.

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial and continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction or Grandfather treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the Initial restriction or up to 24 weeks treatment available under the Continuing or Grandfather restrictions.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

Note Authority approval for sufficient therapy to complete a maximum of 28 weeks of treatment under the initial restriction or 24 weeks of treatment under the continuing or grandfather restrictions may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial treatment - Grandfather patients

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment with this drug for this condition prior to 1 December 2016, **AND**
- Patient must be receiving treatment with this drug for this condition at the time of application, **AND**
- Patient must have had, prior to commencement of omalizumab, a diagnosis of asthma confirmed and documented by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician, defined by the following standard clinical features: forced expiratory volume (FEV1) reversibility or airway hyperresponsiveness or peak expiratory flow (PEF) variability, **AND**
- Patient must have had a duration of asthma of at least 1 year prior to commencement of omalizumab, **AND**
- Patient must have past or current evidence of atopy, documented by skin prick testing or an in vitro measure of specific IgE, **AND**

- Patient must have failed to achieve adequate control with optimised asthma therapy prior to omalizumab therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- Patient must have demonstrated an adequate response to treatment, **AND**
- Patient must be under the care of the same physician for at least 6 months.

Population criteria:

- Patient must be aged 6 to less than 12 years.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

Optimised asthma therapy includes:

- (i) Adherence to optimal inhaled therapy, including high dose inhaled corticosteroid (ICS) and long-acting beta-2 agonist (LABA) therapy for at least six months. If LABA therapy is contraindicated, not tolerated or not effective, montelukast, cromoglycate or nedocromil may be used as an alternative; **AND**
- (ii) treatment with at least 2 courses of oral or IV corticosteroids (daily or alternate day maintenance treatment courses, or 3-5 day exacerbation treatment courses), in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications (including those specified in the relevant TGA-approved Product Information) and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

A review of the patient's records should be conducted to extract pre- and post-omalizumab data on symptoms, quality of life, medication doses, exacerbations and hospitalisations. Examples of parameters to establish response include:

- (i) a reduction in Asthma Control Questionnaire (ACQ-5) or Asthma Control Questionnaire Interviewer Administered (ACQ-IA) score of at least 0.5;
- (ii) maintenance oral corticosteroid dose reduced by at least 25% from baseline; and/or
- (iii) a reduction in the number of hospitalisations or severe exacerbations requiring use of systemic corticosteroids, compared to the 12 months prior to commencement of omalizumab.

The assessment of the patient's response to the initial PBS subsidised course of treatment must be made at around 18 to 22 weeks after the first dose so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed. The same parameters used to establish response to non-PBS-subsidised therapy with omalizumab should be used for the assessment.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with omalizumab.

Patients will be eligible to receive continuing courses of omalizumab treatment of up to 24 weeks providing they continue to demonstrate an adequate response to treatment.

Patients may qualify for PBS-subsidised treatment under this restriction once only.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats to provide for an initial course of omalizumab of up to 24 weeks, consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information) to be administered every 2 or 4 weeks.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Paediatric Grandfather Severe Allergic Asthma PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (dosage, date of commencement and duration of therapy); and
 - (ii) details of pre- and post-omalizumab data on symptoms, quality of life, medication doses, exacerbations and hospitalisations; and
 - (iii) acknowledgement signed by a parent or authorised guardian.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note It is recommended that an application for continuing treatment is submitted at the time of the 22 to 26 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised omalizumab treatment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note TREATMENT OF PAEDIATRIC PATIENTS WITH UNCONTROLLED SEVERE ALLERGIC ASTHMA

Patients are eligible to commence an 'omalizumab treatment cycle' (initial treatment course with or without continuing

treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction. Once a patient has either failed to achieve or maintain a response to omalizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised omalizumab therapy before they are eligible to commence the next cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised omalizumab treatment is stopped to the date of the first application for initial treatment with omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised omalizumab therapy.

(a) Initial treatment:

Applications for initial treatment should be made where a patient has received no prior PBS-subsidised omalizumab treatment in this treatment cycle and wishes to commence such therapy.

All applications for initial treatment for non-grandfathered patients will be limited to provide for a maximum of 28 weeks of therapy for omalizumab.

(b) Grandfather patients:

Paediatric patients who commenced treatment with omalizumab for uncontrolled severe allergic asthma prior to 1 December 2016 and who continue to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment with omalizumab will be authorised under this restriction. Approval will be based on the criteria included in the grandfather restriction. Following completion of the Initial PBS-subsidised course, further applications for treatment with omalizumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle (initial treatment course with or without continuing treatment course/s). If a 'Grandfathered' patient recommences on second and subsequent cycles after a treatment break, the 'Grandfathered' patient must re-qualify for Initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 6 month break in PBS-subsidised therapy' below for further details.

(c) Continuing treatment:

Following the completion of the initial treatment course with omalizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with omalizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing omalizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or ACQ-IA, systemic corticosteroid dose and time-adjusted exacerbation rate, submitted with the Initial authority application for omalizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and The Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised omalizumab therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (systemic corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score or ACQ-IA, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

(4) Monitoring of patients:

Anaphylaxis and anaphylactoid reactions have been reported following first or subsequent administration of omalizumab (see Product Information). Patients should be monitored post-injection, and medications for the treatment of anaphylactic reactions should be available for immediate use following administration of omalizumab. Patients should be informed that such reactions are possible and prompt medical attention should be sought if allergic reactions occur.

omalizumab 150 mg/mL injection, 1 mL syringe

10973M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	410.00	Xolair [NV]

omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe

10967F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	205.00	Xolair [NV]

▪ VEDOLIZUMAB

Note Special Pricing Arrangements apply.

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of infliximab, vedolizumab and adalimumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either infliximab, vedolizumab or adalimumab at any one time. From 1 December 2016, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial each of PBS-subsidised infliximab, vedolizumab or adalimumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with infliximab, vedolizumab or adalimumab while they continue to show a response to therapy. A patient who received PBS-subsidised infliximab, vedolizumab or adalimumab treatment prior to 1 December 2016 is considered to be in their first cycle as of 1 December 2016. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised infliximab, vedolizumab or adalimumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised infliximab, vedolizumab or adalimumab treatment in the most recent cycle to the date of the first application for initial treatment with infliximab, vedolizumab or adalimumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of either infliximab, vedolizumab or adalimumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with infliximab, vedolizumab and adalimumab after therapy after 1 December 2016 .

(a) Initial treatment. Applications for initial treatment should be made where:

(i) an adult patient has received no prior PBS-subsidised treatment with infliximab, vedolizumab or adalimumab in this treatment cycle and wishes to commence such therapy (Initial 1); or

(ii) an adult patient has received prior PBS-subsidised (initial or continuing) infliximab, vedolizumab or adalimumab therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or

(iii) an adult patient wishes to re-commence treatment with infliximab, vedolizumab or adalimumab following a break in PBS-subsidised therapy with the same agent (Initial 2).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab , 14 weeks of therapy for infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab, and this assessment must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist. For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is provided to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with infliximab, vedolizumab or adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment. Assessments of response to a course of PBS-subsidised therapy must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate infliximab, vedolizumab or adalimumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with infliximab, vedolizumab or adalimumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction. To avoid confusion, an application for a patient who wishes to swap to the alternate infliximab, vedolizumab or adalimumab therapy should be accompanied by the approved authority prescription or remaining repeats for the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Mayo clinic score or partial Mayo clinic score submitted with the first authority application for infliximab, vedolizumab or adalimumab. However, prescribers may provide new baseline measurements any time other than when an initial treatment authority application is provided within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised infliximab, vedolizumab or adalimumab therapy of at least 5 years, must requalify for initial treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the Mayo score is measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for moderate to severe ulcerative colitis prior to 1 December 2016 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a continuing patient.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment (new patient – Initial 1)

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score), **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Applications for authorisation of initial treatment must be in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic calculation sheet including the date of assessment of the patient's condition; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and

(iii) the signed patient acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic or partial Mayo clinic score must be no more than 1 month old at the time of application.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose for patients administered doses at weeks 0, 2 and 6 (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

Patients must have signed a patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Note Details of accepted toxicities including severity can be found on the Department of Human Services website at www.humanservices.gov.au.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR

- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]. Patients who have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug. Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response. At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide for a single infusion of 300 mg per dose. Up to a maximum of 2 repeats will be authorised.

Note No applications for increased repeats will be authorised.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Change or Re-commencement of treatment after a break in therapy (Initial 2)

Clinical criteria:

- Patient must have previously been issued with an authority prescription for adalimumab, infliximab or vedolizumab for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with vedolizumab for this condition more than once in the current treatment cycle, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of this drug within the timelines specified in the relevant restriction. If the response assessment to the previous course of this drug is not submitted as detailed in the relevant restriction, the patient will be deemed to have failed therapy with this drug.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide for a single infusion of 300 mg per dose.

Up to a maximum of 2 repeats will be authorised.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses of treatment may be requested by telephone by contacting the Department of Human Services.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (Change or Recommencement of treatment after a break in therapy) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 and Initial 2 restrictions) or 2 repeats (Continuing restriction), **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services.

vedolizumab 300 mg injection, 1 vial

10384M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3105.19	Entyvio [TK]

Highly Specialised Drugs Program (Community Access)

▪ TENOFOVIR + EMTRICITABINE

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

tenofovir alafenamide 10mg + emtricitabine 200 mg tablet, 30

10946D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1500.72	38.30	Descovy 10/200 [GI]

tenofovir alafenamide 25mg + emtricitabine 200 mg tablet, 30

10966E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1500.72	38.30	Descovy 25/200 [GI]