



Australian Government

Department of Health



Schedule of Pharmaceutical Benefits

Summary of Changes

Effective 1 September 2017



Fees, Patient Contributions and Safety Net Thresholds

The following fees, patient contributions and safety net thresholds apply as at 1 September 2017 and are included, where applicable, in prices published in the Schedule —

Dispensing Fees:	Ready-prepared	\$7.15
	Dangerous drug fee	\$3.01
	Extemporaneously-prepared	\$9.19
	Allowable additional patient charge*	\$4.38
Additional Fees (for safety net prices):	Ready-prepared	\$1.21
	Extemporaneously-prepared	\$1.57
Patient Co-payments:	General	\$38.80
	Concessional	\$6.30
Safety Net Thresholds:	General	\$1494.90
	Concessional	\$378.00
Safety Net Card Issue Fee:		\$9.73

* The allowable additional patient charge is a discretionary charge to general patients if a pharmaceutical item has a dispensed price for maximum quantity less than the general patient co-payment. The pharmacist may charge general patients the allowable additional fee but the fee cannot take the cost of the prescription above the general patient co-payment for the medicine. This fee does not count towards the Safety Net threshold.

Summary of Changes

These changes to the Schedule of Pharmaceutical Benefits are effective from 1 September 2017. The Schedule is updated on the first day of each month and is available on the internet at www.pbs.gov.au.

Prescriber Bag

Deletions

Deletion – Brand

3497C *Ventolin Nebules, GK* – **SALBUTAMOL**, salbutamol 5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules

Advance Notices

1 October 2017

Deletion – Brand

3496B *Ventolin Nebules, GK* – **SALBUTAMOL**, salbutamol 2.5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules

General Pharmaceutical Benefits

Additions

Addition – Item

- 11161K **AMINO ACID FORMULA WITH FAT, CARBOHYDRATE, VITAMINS, MINERALS, TRACE ELEMENTS AND MEDIUM CHAIN TRIGLYCERIDES**, amino acid formula with fat, carbohydrate, vitamins, minerals, trace elements and medium chain triglycerides powder for oral liquid, 400 g (*Neocate Junior*)
- 11183N **AMINO ACID FORMULA WITH FAT, CARBOHYDRATE, VITAMINS, MINERALS, TRACE ELEMENTS AND MEDIUM CHAIN TRIGLYCERIDES**, amino acid formula with fat, carbohydrate, vitamins, minerals, trace elements and medium chain triglycerides powder for oral liquid, 400 g (*Neocate Junior*)
- 11169W **CEFTRIAXONE**, ceftriaxone 2 g injection, 5 vials (*Ceftriaxone Alphapharm*)
- 11170X **IDELALISIB**, idelalisib 100 mg tablet, 60 (*Zydelig*)
- 11171Y **IDELALISIB**, idelalisib 100 mg tablet, 60 (*Zydelig*)
- 11162L **IDELALISIB**, idelalisib 150 mg tablet, 60 (*Zydelig*)
- 11165P **IDELALISIB**, idelalisib 150 mg tablet, 60 (*Zydelig*)
- 11180K **URSODEOXYCHOLIC ACID**, ursodeoxycholic acid 500 mg tablet, 100 (*Ursosalk*)
- 11178H **USTEKINUMAB**, ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial (*Stelara*)

Addition – Brand

- 10791Y *Momasone, QA* – **MOMETASONE**, mometasone furoate 0.1% ointment, 15 g
- 10793C *Momasone, QA* – **MOMETASONE**, mometasone furoate 0.1% ointment, 15 g
- 10812C *Momasone, QA* – **MOMETASONE**, mometasone furoate 0.1% ointment, 15 g
- 10814E *Momasone, QA* – **MOMETASONE**, mometasone furoate 0.1% ointment, 15 g
- 10828X *Momasone, QA* – **MOMETASONE**, mometasone furoate 0.1% ointment, 15 g
- 1915T *Momasone, QA* – **MOMETASONE**, mometasone furoate 0.1% ointment, 15 g

Addition – Equivalence Indicator

- 1581F *Pregnyl, MK* – **HUMAN CHORIONIC GONADOTROPHIN**, human chorionic gonadotrophin 1500 units injection [3 ampoules] (&) inert substance diluent [3 x 1 mL ampoules], 1 pack

11148R *Pregnyl, MK* – **HUMAN CHORIONIC GONADOTROPHIN**, human chorionic gonadotrophin 1500 units injection [3 vials] (&) inert substance diluent [3 x 1 mL vials], 1 pack

Deletions

Deletion – Item

8422G **HYDROMORPHONE**, hydromorphone hydrochloride 50 mg/5 mL injection, 5 x 5 mL ampoules (*Dilaudid-HP*)

Deletion – Brand

9012H *Alendronate D3 70 mg/70 microgram, EA* – **ALENDRONATE + COLECALCIFEROL**, alendronate 70 mg + colecalciferol 70 microgram tablet, 4

9183H *Alendronate D3 70 mg/140 microgram, EA* – **ALENDRONATE + COLECALCIFEROL**, alendronate 70 mg + colecalciferol 140 microgram tablet, 4

9351E *Alendronate Plus D3 Calcium Actavis, EA* – **ALENDRONATE + COLECALCIFEROL (&) CALCIUM CARBONATE**, alendronate 70 mg + colecalciferol 140 microgram tablet [4] (&) calcium (as carbonate) 500 mg tablet [48], 1 pack

9351E *Alendronate Plus D3 and Calcium Sandoz, SZ* – **ALENDRONATE + COLECALCIFEROL (&) CALCIUM CARBONATE**, alendronate 70 mg + colecalciferol 140 microgram tablet [4] (&) calcium (as carbonate) 500 mg tablet [48], 1 pack

1886G *Chem mart Amoxycillin, CH* – **AMOXYCILLIN**, amoxycillin 125 mg/5 mL powder for oral liquid, 100 mL

1886G *Terry White Chemists Amoxycillin, TW* – **AMOXYCILLIN**, amoxycillin 125 mg/5 mL powder for oral liquid, 100 mL

3302T *Chem mart Amoxycillin, CH* – **AMOXYCILLIN**, amoxycillin 125 mg/5 mL powder for oral liquid, 100 mL

3302T *Terry White Chemists Amoxycillin, TW* – **AMOXYCILLIN**, amoxycillin 125 mg/5 mL powder for oral liquid, 100 mL

1887H *Chem mart Amoxycillin, CH* – **AMOXYCILLIN**, amoxycillin 250 mg/5 mL powder for oral liquid, 100 mL

1887H *Terry White Chemists Amoxycillin, TW* – **AMOXYCILLIN**, amoxycillin 250 mg/5 mL powder for oral liquid, 100 mL

3393N *Chem mart Amoxycillin, CH* – **AMOXYCILLIN**, amoxycillin 250 mg/5 mL powder for oral liquid, 100 mL

3393N *Terry White Chemists Amoxycillin, TW* – **AMOXYCILLIN**, amoxycillin 250 mg/5 mL powder for oral liquid, 100 mL

8179L *Anastrozole RBX, RA* – **ANASTROZOLE**, anastrozole 1 mg tablet, 30

1785Y *Ceftriaxone Alphapharm, AF* – **CEFTRIAZONE**, ceftriaxone 2 g injection, 1 vial

2655R *Chem mart Cephalexin, CH* – **CEPHALEXIN**, cephalexin 250 mg capsule, 20

2655R *Terry White Chemists Cephalexin, TW* – **CEPHALEXIN**, cephalexin 250 mg capsule, 20

3058Y *Chem mart Cephalexin, CH* – **CEPHALEXIN**, cephalexin 250 mg capsule, 20

3058Y *Terry White Chemists Cephalexin, TW* – **CEPHALEXIN**, cephalexin 250 mg capsule, 20

3317N *Chem mart Cephalexin, CH* – **CEPHALEXIN**, cephalexin 250 mg capsule, 20

3317N *Terry White Chemists Cephalexin, TW* – **CEPHALEXIN**, cephalexin 250 mg capsule, 20

10778G *Chem mart Cephalexin, CH* – **CEPHALEXIN**, cephalexin 500 mg capsule, 20

10778G *Terry White Chemists Cephalexin, TW* – **CEPHALEXIN**, cephalexin 500 mg capsule, 20

3119E *Chem mart Cephalexin, CH* – **CEPHALEXIN**, cephalexin 500 mg capsule, 20

3119E *Terry White Chemists Cephalexin, TW* – **CEPHALEXIN**, cephalexin 500 mg capsule, 20

3318P *Chem mart Cephalexin, CH* – **CEPHALEXIN**, cephalexin 500 mg capsule, 20

3318P *Terry White Chemists Cephalexin, TW* – **CEPHALEXIN**, cephalexin 500 mg capsule, 20

8559L *Gabaran, RA* – **GABAPENTIN**, gabapentin 600 mg tablet, 100

8389M *Gabaran, RA* – **GABAPENTIN**, gabapentin 800 mg tablet, 100

8654L *Levitaccord, RA* – **LEVETIRACETAM**, levetiracetam 250 mg tablet, 60

8655M *Levitaccord, RA* – **LEVETIRACETAM**, levetiracetam 500 mg tablet, 60

1596B *Ondansetron-Clarix, AE* – **ONDANSETRON**, ondansetron 4 mg/2 mL injection, 2 mL ampoule

8226Y *Ondansetron-Clarix, AE* – **ONDANSETRON**, ondansetron 4 mg/2 mL injection, 2 mL ampoule

1597C *Ondansetron-Clarix, AE* – **ONDANSETRON**, ondansetron 8 mg/4 mL injection, 4 mL ampoule

8227B *Ondansetron-Clarix, AE* – **ONDANSETRON**, ondansetron 8 mg/4 mL injection, 4 mL ampoule

8399C *Chem mart Pantoprazole, CH* – **PANTOPRAZOLE**, pantoprazole 20 mg enteric tablet, 30

8399C	<i>Terry White Chemists Pantoprazole, TW</i> – PANTOPRAZOLE , pantoprazole 20 mg enteric tablet, 30
8007K	<i>Chem mart Pantoprazole, CH</i> – PANTOPRAZOLE , pantoprazole 40 mg enteric tablet, 30
8007K	<i>Terry White Chemists Pantoprazole, TW</i> – PANTOPRAZOLE , pantoprazole 40 mg enteric tablet, 30
8008L	<i>Chem mart Pantoprazole, CH</i> – PANTOPRAZOLE , pantoprazole 40 mg enteric tablet, 30
8008L	<i>Terry White Chemists Pantoprazole, TW</i> – PANTOPRAZOLE , pantoprazole 40 mg enteric tablet, 30
2001H	<i>Ventolin Nebules, GK</i> – SALBUTAMOL , salbutamol 5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules
8622T	<i>Chem mart Telmisartan HCTZ 40/12.5, CH</i> – TELMISARTAN + HYDROCHLOROTHIAZIDE , telmisartan 40 mg + hydrochlorothiazide 12.5 mg tablet, 28
8622T	<i>Terry White Chemists Telmisartan HCTZ 40/12.5, TW</i> – TELMISARTAN + HYDROCHLOROTHIAZIDE , telmisartan 40 mg + hydrochlorothiazide 12.5 mg tablet, 28
8623W	<i>Chem mart Telmisartan HCTZ 80/12.5, CH</i> – TELMISARTAN + HYDROCHLOROTHIAZIDE , telmisartan 80 mg + hydrochlorothiazide 12.5 mg tablet, 28
8623W	<i>Terry White Chemists Telmisartan HCTZ 80/12.5, TW</i> – TELMISARTAN + HYDROCHLOROTHIAZIDE , telmisartan 80 mg + hydrochlorothiazide 12.5 mg tablet, 28
9381R	<i>Chem mart Telmisartan HCTZ 80/25, CH</i> – TELMISARTAN + HYDROCHLOROTHIAZIDE , telmisartan 80 mg + hydrochlorothiazide 25 mg tablet, 28
9381R	<i>Terry White Chemists Telmisartan HCTZ 80/25, TW</i> – TELMISARTAN + HYDROCHLOROTHIAZIDE , telmisartan 80 mg + hydrochlorothiazide 25 mg tablet, 28

Alterations

Alteration – Restriction

The following items have additions, deletions or alterations to restrictions, notes and/or cautions.

1785Y	CEFTRIAXONE , ceftriaxone 2 g injection, 1 vial (<i>Ceftriaxone-AFT</i>)
10909E	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
11076Y	VEMURAFENIB , vemurafenib 240 mg tablet, 56 (<i>Zelboraf</i>)
11081F	VEMURAFENIB , vemurafenib 240 mg tablet, 56 (<i>Zelboraf</i>)

Advance Notices

1 October 2017

Deletion – Brand

1085D	<i>Cefotaxime Sandoz, SZ</i> – CEFOTAXIME , cefotaxime 1 g injection, 1 vial
5048Q	<i>Cefotaxime Sandoz, SZ</i> – CEFOTAXIME , cefotaxime 1 g injection, 1 vial
2593L	<i>Ferrum H, AS</i> – IRON POLYMALTOSE , iron (as polymaltose) 100 mg/2 mL injection, 5 x 2 mL ampoules
2805P	<i>Ferrum H, AS</i> – IRON POLYMALTOSE , iron (as polymaltose) 100 mg/2 mL injection, 5 x 2 mL ampoules
3010K	<i>Norfloxacin Sandoz, SZ</i> – NORFLOXACIN , norfloxacin 400 mg tablet, 14
8664B	<i>Riluzole Winthrop, WA</i> – RILUZOLE , riluzole 50 mg tablet, 56
2000G	<i>Ventolin Nebules, GK</i> – SALBUTAMOL , salbutamol 2.5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules
10062N	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 180 mg capsule, 5
2438H	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 180 mg capsule, 5
8378Y	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 5 mg capsule, 5
8379B	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 20 mg capsule, 5
8380C	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 100 mg capsule, 5
8381D	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 250 mg capsule, 5
8819E	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 5 mg capsule, 5
8820F	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 20 mg capsule, 5
8821G	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 100 mg capsule, 5
9361Q	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 140 mg capsule, 5
9362R	<i>Astromide, FR</i> – TEMOZOLOMIDE , temozolomide 140 mg capsule, 5

1 November 2017

Deletion – Brand

1210Q *Ciprofloxacin-BW, GQ* – **CIPROFLOXACIN**, ciprofloxacin 750 mg tablet, 14

1 December 2017

Deletion – Brand

8883M *Avanza, MK* – **MIRTAZAPINE**, mirtazapine 45 mg tablet, 30

Highly Specialised Drugs Program (Private Hospital)

Additions

Addition – Item

- 11163M **OMALIZUMAB**, omalizumab 150 mg/mL injection, 1 mL syringe (*Xolair*)
- 11175E **OMALIZUMAB**, omalizumab 150 mg/mL injection, 1 mL syringe (*Xolair*)
- 11167R **PEGVISOMANT**, pegvisomant 10 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack (*Somavert*)
- 11172B **PEGVISOMANT**, pegvisomant 15 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack (*Somavert*)
- 11166Q **PEGVISOMANT**, pegvisomant 20 mg injection [1 vial] (&) inert substance diluent [1 x 1 mL vial], 1 pack (*Somavert*)
- 11174D **PEGVISOMANT**, pegvisomant 20 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack (*Somavert*)
- 11164N **USTEKINUMAB**, ustekinumab 130 mg/26 mL injection, 26 mL vial (*Stelara*)

Addition – Brand

6357N *Valganciclovir Juno, JU* – **VALGANCICLOVIR**, valganciclovir 450 mg tablet, 60

Alterations

Alteration – Restriction

The following items have additions, deletions or alterations to restrictions, notes and/or cautions.

- 10184B **INFLIXIMAB**, infliximab 100 mg injection, 1 vial (*Inflectra, Remicade, Renflexis*)
- 6496X **INFLIXIMAB**, infliximab 100 mg injection, 1 vial (*Inflectra, Remicade, Renflexis*)
- 6332G **LANREOTIDE**, lanreotide 30 mg modified release injection [1 vial] (&) inert substance diluent [2 mL ampoule], 1 pack (*Somatuline LA*)
- 6423C **LANREOTIDE**, lanreotide 60 mg/0.5 mL injection, 0.5 mL syringe (*Somatuline Autogel*)
- 6424D **LANREOTIDE**, lanreotide 90 mg/0.5 mL injection, 0.5 mL syringe (*Somatuline Autogel*)
- 6425E **LANREOTIDE**, lanreotide 120 mg/0.5 mL injection, 0.5 mL syringe (*Somatuline Autogel*)
- 10549F **OCTREOTIDE**, octreotide 20 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (*Sandostatin LAR*)
- 10558Q **OCTREOTIDE**, octreotide 30 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (*Sandostatin LAR*)
- 10566D **OCTREOTIDE**, octreotide 10 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (*Sandostatin LAR*)
- 6227R **OCTREOTIDE**, octreotide 50 microgram/mL injection, 5 x 1 mL ampoules (*Hospira Pty Limited, Octreotide (SUN), Octreotide MaxRx, Sandostatin 0.05*)
- 6228T **OCTREOTIDE**, octreotide 100 microgram/mL injection, 5 x 1 mL ampoules (*Hospira Pty Limited, Octreotide (SUN), Octreotide MaxRx, Sandostatin 0.1*)
- 6229W **OCTREOTIDE**, octreotide 500 microgram/mL injection, 5 x 1 mL ampoules (*Hospira Pty Limited, Octreotide (SUN), Octreotide MaxRx, Sandostatin 0.5*)
- 10880P **PASIREOTIDE**, pasireotide 20 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (*Signifor LAR*)
- 10884W **PASIREOTIDE**, pasireotide 40 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (*Signifor LAR*)
- 10887B **PASIREOTIDE**, pasireotide 60 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (*Signifor LAR*)

Highly Specialised Drugs Program (Public Hospital)

Additions

Addition – Item

11168T	OMALIZUMAB , omalizumab 150 mg/mL injection, 1 mL syringe (<i>Xolair</i>)
11176F	OMALIZUMAB , omalizumab 150 mg/mL injection, 1 mL syringe (<i>Xolair</i>)
11179J	PEGVISOMANT , pegvisomant 10 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack (<i>Somavert</i>)
11173C	PEGVISOMANT , pegvisomant 15 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack (<i>Somavert</i>)
11177G	PEGVISOMANT , pegvisomant 20 mg injection [1 vial] (&) inert substance diluent [1 x 1 mL vial], 1 pack (<i>Somavert</i>)
11181L	PEGVISOMANT , pegvisomant 20 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack (<i>Somavert</i>)
11182M	USTEKINUMAB , ustekinumab 130 mg/26 mL injection, 26 mL vial (<i>Stelara</i>)

Addition – Brand

9569P	<i>Valganciclovir Juno, JU</i> – VALGANCICLOVIR , valganciclovir 450 mg tablet, 60
-------	---

Alterations

Alteration – Restriction

The following items have additions, deletions or alterations to restrictions, notes and/or cautions.

10196P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Remicade, Renflexis</i>)
5756Y	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Remicade, Renflexis</i>)
5776B	LANREOTIDE , lanreotide 30 mg modified release injection [1 vial] (&) inert substance diluent [2 mL ampoule], 1 pack (<i>Somatuline LA</i>)
5777C	LANREOTIDE , lanreotide 60 mg/0.5 mL injection, 0.5 mL syringe (<i>Somatuline Autogel</i>)
5778D	LANREOTIDE , lanreotide 90 mg/0.5 mL injection, 0.5 mL syringe (<i>Somatuline Autogel</i>)
5779E	LANREOTIDE , lanreotide 120 mg/0.5 mL injection, 0.5 mL syringe (<i>Somatuline Autogel</i>)
10533J	OCTREOTIDE , octreotide 20 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (<i>Sandostatin LAR</i>)
10543X	OCTREOTIDE , octreotide 10 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (<i>Sandostatin LAR</i>)
10550G	OCTREOTIDE , octreotide 30 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (<i>Sandostatin LAR</i>)
9508K	OCTREOTIDE , octreotide 50 microgram/mL injection, 5 x 1 mL ampoules (<i>Hospira Pty Limited, Octreotide (SUN), Octreotide MaxRx, Sandostatin 0.05</i>)
9509L	OCTREOTIDE , octreotide 100 microgram/mL injection, 5 x 1 mL ampoules (<i>Hospira Pty Limited, Octreotide (SUN), Octreotide MaxRx, Sandostatin 0.1</i>)
9510M	OCTREOTIDE , octreotide 500 microgram/mL injection, 5 x 1 mL ampoules (<i>Hospira Pty Limited, Octreotide (SUN), Octreotide MaxRx, Sandostatin 0.5</i>)
10882R	PASIREOTIDE , pasireotide 60 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (<i>Signifor LAR</i>)
10883T	PASIREOTIDE , pasireotide 40 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (<i>Signifor LAR</i>)
10886Y	PASIREOTIDE , pasireotide 20 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack (<i>Signifor LAR</i>)

Highly Specialised Drugs Program (Community Access)

Additions

Addition – Brand

10303G	<i>Nevirapine XR APOTEX, TX</i> – NEVIRAPINE , nevirapine 400 mg modified release tablet, 30
10306K	<i>Valganciclovir Juno, JU</i> – VALGANCICLOVIR , valganciclovir 450 mg tablet, 60

Addition – Equivalence Indicator

10303G Viramune XR, BY – **NEVIRAPINE**, nevirapine 400 mg modified release tablet, 30

Deletions

Deletion – Brand

10348P Lamivudine RBX, RA – **LAMIVUDINE**, lamivudine 150 mg tablet, 60

10311Q Lamivudine RBX, RA – **LAMIVUDINE**, lamivudine 300 mg tablet, 30

10304H Nevirapine RBX, RA – **NEVIRAPINE**, nevirapine 200 mg tablet, 60

IVF Program

Additions

Addition – Equivalence Indicator

6178E Pregnyl, MK – **HUMAN CHORIONIC GONADOTROPHIN**, human chorionic gonadotrophin 1500 units injection [3 ampoules] (&) inert substance diluent [3 x 1 mL ampoules], 1 pack

11154C Pregnyl, MK – **HUMAN CHORIONIC GONADOTROPHIN**, human chorionic gonadotrophin 1500 units injection [3 vials] (&) inert substance diluent [3 x 1 mL vials], 1 pack

11156E Pregnyl, MK – **HUMAN CHORIONIC GONADOTROPHIN**, human chorionic gonadotrophin 5000 units injection [1 vial] (&) inert substance diluent [1 mL vial], 1 pack

6181H Pregnyl, MK – **HUMAN CHORIONIC GONADOTROPHIN**, human chorionic gonadotrophin 5000 units injection [1 ampoule] (&) inert substance diluent [1 mL ampoule], 1 pack

Repatriation Pharmaceutical Benefits

Deletions

Deletion – Item

4453J **ALUMINIUM HYDROXIDE WITH MAGNESIUM HYDROXIDE AND SIMETHICONE**, ALUMINIUM HYDROXIDE with MAGNESIUM HYDROXIDE and SIMETHICONE Tablet 400 mg-400 mg-40 mg, 100 (*Mylanta Double Strength*)

4106D **CICLOPIROX**, ciclopirox olamine 1.5% shampoo, 60 mL (*Stieprox Liquid*)

Alterations

Alteration – Manufacturer Code

		From	To
4463X	Lamisil DermGel – TERBINAFINE , terbinafine 1% gel, 15 g	NC	GK

Advance Notices

1 October 2017

Deletion – Brand

4170L Prodeinextra, SW – **PARACETAMOL + CODEINE**, paracetamol 500 mg + codeine phosphate 15 mg tablet, 20

General Pharmaceutical Benefits

■ AMINO ACID FORMULA WITH FAT, CARBOHYDRATE, VITAMINS, MINERALS, TRACE ELEMENTS AND MEDIUM CHAIN TRIGLYCERIDES

Authority required

Cows' milk protein enteropathy

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides).

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Initial treatment for up to 3 months

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must require an amino acid based formula as a component of a dietary elimination program.

Population criteria:

- Patient must be 18 years of age or less.

Treatment with oral steroids should not be commenced during the period of initial treatment.

Eosinophilic oesophagitis is demonstrated by the following criteria:

- (i) Chronic symptoms of reflux that persisted despite a 2-month trial of a proton pump inhibitor or chronic dysphagia; and
- (ii) A lack of demonstrable anatomic abnormality with the exception of stricture, which can be attributable to eosinophilic oesophagitis; and
- (iii) Eosinophilic infiltration of the oesophagus, demonstrated by oesophageal biopsy specimens obtained by endoscopy and where the most densely involved oesophageal biopsy had 20 or more eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies.

The date of birth of the patient must be included in the authority application.

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

amino acid formula with fat, carbohydrate, vitamins, minerals, trace elements and medium chain triglycerides powder for oral liquid, 400 g

11161K

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8	5	..	*353.31	38.80	Neocate Junior [SB]

■ AMINO ACID FORMULA WITH FAT, CARBOHYDRATE, VITAMINS, MINERALS, TRACE ELEMENTS AND MEDIUM CHAIN TRIGLYCERIDES

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have been assessed at least once or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have had failure to thrive prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist at intervals not greater than 12 months.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides) prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Cows' milk anaphylaxis

Treatment criteria:

- Must be treated by a specialist allergist or clinical immunologist, or in consultation with a specialist allergist or clinical immunologist.

Population criteria:

- Patient must be up to the age of 24 months.

Anaphylaxis is defined as a severe and/or potentially life threatening allergic reaction.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe intestinal malabsorption including short bowel syndrome

Clinical criteria:

- Patient must have failed to respond to protein hydrolysate formulae; OR
- Patient must have been receiving parenteral nutrition.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must have responded to an initial course of PBS-subsidised treatment.

Population criteria:

- Patient must be 18 years of age or less.

Response to initial treatment is demonstrated by oesophageal biopsy specimens obtained by endoscopy, where the most densely involved oesophageal biopsy had 5 or less eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies. The response criteria will not be deemed to have been met if oral steroids were commenced during initial treatment.

amino acid formula with fat, carbohydrate, vitamins, minerals, trace elements and medium chain triglycerides powder for oral liquid, 400 g

11183N

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8	5	..	*353.31	38.80	Neocate Junior [SB]

▪ **CEFTRIAXONE**

Note Pharmaceutical benefits that have the form ceftriaxone 2 g injection, 1 vial and pharmaceutical benefits that have the form ceftriaxone 2 g injection, 5 vials are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

Restricted benefit

Septicaemia, suspected

Restricted benefit

Septicaemia, proven

ceftriaxone 2 g injection, 5 vials

11169W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	22.22	23.43	^a Ceftriaxone Alphapharm [AF]

ceftriaxone 2 g injection, 1 vial

1785Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	..	..	*22.25	23.46	^a Ceftriaxone-AFT [AE]	

■ CERTOLIZUMAB PEGOL

Note The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to the Department of Human Services no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Note Patients who fail to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this Treatment Cycle. Patients may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment – Initial 1 (new patient or patient recommencing treatment after a break of 5 years or more)

Clinical criteria:

- Patient must have severe active psoriatic arthritis, **AND**
- Patient must have received no prior PBS-subsidised treatment with a biological agent for this condition; OR
- Patient must have received no PBS-subsidised treatment with a biological agent for at least 5 years if they have previously received PBS-subsidised treatment with a biological agent for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be an adult.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

For the purposes of this restriction 'biological agent' means adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab or ustekinumab.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and
either

- (a) an active joint count of at least 20 active (swollen and tender) joints; or
 - (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).
- If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form; and
- (3) a signed patient acknowledgement.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment – Initial 2 (change or recommencement of treatment)

Clinical criteria:

- Patient must have a documented history of severe active psoriatic arthritis, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological agents within this Treatment Cycle, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with this drug during the current Treatment Cycle, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be an adult.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

For the purposes of this restriction 'biological agent' means adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab or ustekinumab.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Applications for a patient who has previously received PBS-subsidised treatment with this drug within this Treatment Cycle and who wishes to recommence therapy with this drug within this same Cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug.

Where the most recent course of PBS-subsidised treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised biological therapy or, under this restriction, for patients who have received previous PBS-subsidised biological therapy), the patient must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must have been submitted no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised treatment with this drug was approved under the continuing treatment criteria, the patient must have been assessed for response, and the assessment submitted no later than 4 weeks from the date that course was ceased.

Where a response assessment was not submitted within these timeframes, the patient will be deemed to have failed to respond to treatment.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time.

Where the term 'biological agents' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab only.

Patients receiving PBS-subsidised treatment for psoriatic arthritis are able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle.

There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and

(ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and

(iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 1 or Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and secukinumab, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients - ustekinumab and secukinumab only.

For patients who commenced treatment with ustekinumab for psoriatic arthritis prior to 1 May 2016 and for patients who commenced treatment with secukinumab for psoriatic arthritis prior to 1 October 2016, applications for initial PBS-subsidised treatment as continuing therapy may be made under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Within a Treatment Cycle patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

(i) they have not received PBS-subsidised treatment with that particular biological agent previously; or

(ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; and

(iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment application is submitted within a treatment Cycle and these revised baseline measurements will be used to assess response.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

10909E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*3549.97	38.80	Cimzia [UC]

■ IDELALISIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with rituximab, **AND**
- The condition must have relapsed or be refractory to at least one prior therapy, **AND**
- The condition must be CD20 positive, **AND**
- Patient must have a total cumulative illness rating scale (CIRS) score of greater than 6 (excluding CLL-induced illness or organ damage), **AND**
- Patient must be inappropriate for chemo-immunotherapy.

A patient can be considered inappropriate for chemo-immunotherapy when one or more of the following are experienced:

1. Severe neutropenia defined as absolute neutrophil count of less than or equal to $1.0 \times 10^9/L$; or
2. Severe thrombocytopenia defined as platelet count of less than or equal to $50 \times 10^9/L$; or
3. Evidence of one or more 17p chromosomal deletions demonstrated by fluorescence in situ hybridisation (FISH).

Full blood count results must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

- a) A completed authority prescription form;
- b) A completed CLL/SLL PBS Authority Application - Supporting information form; and
- c) Pathology report indicating that the patient can be considered inappropriate for chemo-immunotherapy due to one or more of the following:
 - 1) Recent severe neutropenia; or
 - 2) Recent severe thrombocytopenia; or
 - 3) Presence of 17p chromosomal deletion using fluorescence in situ hybridisation (FISH).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with rituximab, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

idelalisib 150 mg tablet, 60

11162L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5367.27	38.80	Zydelig [GI]

idelalisib 100 mg tablet, 60

11170X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5367.27	38.80	Zydelig [GI]

■ IDELALISIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Refractory follicular B-cell non-Hodgkin's lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be refractory to a prior therapy with rituximab, **AND**
- The condition must be refractory to a prior therapy with an alkylating agent, **AND**
- The treatment must be the sole PBS-subsidised treatment for this condition.

The condition is considered refractory to a prior therapy when the patient experiences less than a partial response or progression of disease within 6 months after completion of the prior therapy.

The condition is considered refractory to both rituximab and an alkylating agent if the agents were administered together or in successive treatment regimens.

The authority application must be made in writing and must include: a) A completed authority prescription form; and b) A completed Refractory follicular B-cell non-Hodgkin's lymphoma PBS Authority Application - Supporting information form which must include date of completion of prior therapies with rituximab and an alkylating agent.

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe this drug should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Refractory follicular B-cell non-Hodgkin's lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised treatment for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

idelalisib 150 mg tablet, 60

11165P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5367.27	38.80	Zydelig [GI]

idelalisib 100 mg tablet, 60

11171Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5367.27	38.80	Zydelig [GI]

■ URSODEOXYCHOLIC ACID

Note Not for use in the treatment of sclerosing cholangitis or cholelithiasis.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5757

Primary biliary cirrhosis

ursodeoxycholic acid 500 mg tablet, 100

11180K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	288.57	38.80	Ursofalk [OA]

■ USTEKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying drugs (bDMDs) for adult patients with severe Crohn disease. Where the term bDMDs appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab). Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological disease modifying drugs at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to be in their first cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised bDMD therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of bDMD therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of bDMD therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial 1 - new patients); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Initial 2 - Change or Re-commencement).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMD therapy.

For second and subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

Assessments of response to a course of PBS-subsidised therapy must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMD therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate bDMD therapy at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular bDMD therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to adalimumab, infliximab, vedolizumab or ustekinumab (where eligible in terms of disease severity) should be accompanied by the approved authority prescription or remaining repeats for the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised bDMD therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment (new patient - initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; OR
- Patient must have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to steroids, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more months or have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to this drug; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months or have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to this drug; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more months or have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to this drug, **AND**
- Patient must have severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220 if affected by extensive small intestine disease; OR
- Patient must have severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 if not affected by extensive small intestine disease, short gut syndrome or is an ostomy patient, **AND**

- Patient must have evidence of intestinal inflammation and have diagnostic imaging or surgical evidence of short gut syndrome if affected by the syndrome or has an ileostomy or colostomy; OR
- Patient must have radiological evidence of intestinal inflammation if the patient has extensive small intestinal disease affecting more than 50 cm of the small intestine; OR
- Patient must (a) have evidence of intestinal inflammation, including: (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or (ii) faeces: higher than normal lactoferrin or calprotectin level; or (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery; or (b) be assessed clinically as being in a high faecal output state; or (c) be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and

(iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and

(iv) the date of the most recent clinical assessment; and

(v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application.

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe Crohn disease

Treatment Phase: Change or Re-commencement of treatment (initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological disease modifying drug for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition in the current treatment cycle.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

-
- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following:
- (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or
 - (ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and
 - (iii) the date of clinical assessment; and
 - (iv) the details of prior biological disease modifying drug treatment including the details of date and duration of treatment.

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for 2 vials of 45 mg and no repeats.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological disease modifying drug (bDMD) therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised bDMD treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of bDMD treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of bDMD.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent first dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe Crohn disease

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment with this drug, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or
 - (ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and
 - (iii) the date of clinical assessment.

All assessments, pathology tests, and diagnostic imaging studies must be made within 1 month of the date of application.

If the application is the first application for continuing treatment with this drug, an assessment of the patient's response to the initial course of treatment must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to the Department of Human Services no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to the Department of Human Services within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats; up to 1 repeat will be authorised for patients whose dosing frequency is every 12 weeks. Up to a maximum of 2 repeats will be authorised for patients whose dosing frequency is every 8 weeks.

If fewer than the maximum stated repeats in the relevant treatment phase are requested at the time of the application, authority approvals for sufficient repeats to complete the balance of the stated repeats in the relevant treatment phase may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend the relevant treatment phase.

Note No increase in the maximum quantity or number of units may be authorised.

Note Increase in the maximum number of repeats of up to 2 may be authorised in patients whose dosing frequency is every 8 weeks.

Authority required

Severe Crohn disease

Treatment Phase: Initial PBS-subsidised treatment (Grandfather)

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have previously received non-PBS-subsidised therapy with this drug for this condition prior to 1 September 2017, **AND**
- Patient must be receiving treatment with ustekinumab at the time of application, **AND**
- Patient must have had a Crohn Disease Activity Index (CDAI) Score of greater than or equal to 300 prior to commencing treatment with this drug; OR
- Patient must have a documented history of intestinal inflammation and have diagnostic imaging or surgical evidence of short gut syndrome if affected by the syndrome or has an ileostomy or colostomy; OR
- Patient must have a documented history and radiological evidence of intestinal inflammation if the patient has extensive small intestinal disease affecting more than 50 cm of the small intestine, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Applications for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Crohn Disease Grandfathered PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and

(iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and

(iv) the date of the most recent clinical assessment; and

(v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to the Department of Human Services no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to the Department of Human Services within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats; up to 1 repeat will be authorised for patients whose dosing frequency is every 12 weeks. Up to a maximum of 2 repeats will be authorised for patients whose dosing frequency is every 8 weeks

If fewer than the maximum stated repeats in the relevant treatment phase are requested at the time of the application, authority approvals for sufficient repeats to complete the balance of the stated repeats in the relevant treatment phase may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend the relevant treatment phase.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

Note No applications for increased maximum quantities will be authorised.

Note Increase in the maximum number of repeats of up to two may be authorised in patients whose dosing frequency is every 8 weeks.

Authority required

Severe Crohn disease

Treatment Phase: Balance of supply for Initial treatment, Continuing treatment or Grandfathered treatment

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete 16 weeks of treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment; OR
- Patient must have received insufficient therapy with this drug under the Grandfathered treatment restriction to complete 24 weeks of treatment.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services

Note No increase in the maximum quantity or number of units may be authorised.

ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial

11178H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*8614.15	38.80	Stelara [JC]

▪ VEMURAFENIB

Note A patient who has had progressive disease when treated with another BRAF inhibitor is not eligible to receive PBS-subsidised treatment with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

6044

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be positive for a BRAF V600 mutation, **AND**
- The condition must not have been treated previously with PBS subsidised therapy; OR
- Patient must have developed intolerance to another BRAF inhibitor of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must have a WHO performance status of 2 or less.

vemurafenib 240 mg tablet, 56

11076Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	3	..	*8187.59	38.80	Zelboraf [RO]

▪ VEMURAFENIB

Note A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Note A patient who has had progressive disease when treated with another BRAF inhibitor is not eligible to receive PBS-subsidised treatment with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

6013

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must have stable or responding disease.

vemurafenib 240 mg tablet, 56

11081F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	..	*8187.59	38.80	Zelboraf [RO]

Highly Specialised Drugs Program (Private Hospital)

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of infliximab, vedolizumab and adalimumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either infliximab, vedolizumab or adalimumab at any one time. From 1 December 2016, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial each of PBS-subsidised infliximab, vedolizumab or adalimumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with infliximab, vedolizumab or adalimumab while they continue to show a response to therapy. A patient who received PBS-subsidised infliximab, vedolizumab or adalimumab treatment prior to 1 December 2016 is considered to be in their first cycle as of 1 December 2016. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised infliximab, vedolizumab or adalimumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised infliximab, vedolizumab or adalimumab treatment in the most recent cycle to the date of the first application for initial treatment with infliximab, vedolizumab or adalimumab under the new treatment cycle. A patient who has failed fewer than 3 trials of either infliximab, vedolizumab or adalimumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with infliximab, vedolizumab and adalimumab after therapy after 1 December 2016 .

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) an adult patient has received no prior PBS-subsidised treatment with infliximab, vedolizumab or adalimumab in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) an adult patient has received prior PBS-subsidised (initial or continuing) infliximab, vedolizumab or adalimumab therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) an adult patient wishes to re-commence treatment with infliximab, vedolizumab or adalimumab following a break in PBS-subsidised therapy with the same agent (Initial 2).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab , 14 weeks of therapy for infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab, and this assessment must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist. For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is provided to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with infliximab, vedolizumab or adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment. Assessments of response to a course of PBS-subsidised therapy must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate infliximab, vedolizumab or adalimumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with infliximab, vedolizumab or adalimumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction. To avoid confusion, an application for a patient who wishes to swap to the alternate infliximab, vedolizumab or adalimumab therapy should be accompanied by the approved authority prescription or remaining repeats for

the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Mayo clinic score or partial Mayo clinic score submitted with the first authority application for infliximab, vedolizumab or adalimumab. However, prescribers may provide new baseline measurements any time other than when an initial treatment authority application is provided within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised infliximab, vedolizumab or adalimumab therapy of at least 5 years, must requalify for initial treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the Mayo score is measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for moderate to severe ulcerative colitis prior to 1 December 2016 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a continuing patient.

Note TREATMENT OF PAEDIATRIC PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with infliximab or adalimumab for moderate to severe ulcerative colitis; and infliximab for acute severe ulcerative colitis.

Where the term 'tumour necrosis factor (TNF) alpha antagonist' appears in the following NOTES and restrictions, it refers to infliximab and adalimumab only. A patient is eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alpha antagonists at any one time. Infliximab and adalimumab are PBS-subsidised for moderate to severe disease while only infliximab is PBS-subsidised for acute severe disease. From 1 June 2017, under the PBS, all will be able to commence a treatment cycle where they may trial each PBS-subsidised TNF-alpha antagonist without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a TNF-alpha antagonist while they continue to show a response to therapy. A patient who received PBS-subsidised TNF-alpha antagonist treatment prior to 1 June 2017 is considered to be in their first cycle as of 1 June 2017. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alpha antagonist more than twice. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alpha antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alpha antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alpha antagonist under the new treatment cycle. A patient who has failed fewer than 3 trials of TNF-alpha antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle. A patient who has failed fewer than 3 trials of TNF-alpha antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle. There is no limit to the number of treatment cycles a patient may undertake in their lifetime. (1) How to prescribe PBS-subsidised TNF-alpha antagonist therapy after 1 June 2017. (a) Initial treatment. Applications for initial treatment should be made where: (i) a patient has received no prior PBS-subsidised TNF-alpha antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or (ii) a patient has received prior PBS-subsidised (initial or continuing) treatment with a TNF-alpha antagonist and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping treatment' below]; or (iii) a patient wishes to re-commence treatment with a specific TNF-alpha antagonist following a break in PBS-subsidised therapy with that agent (Initial 2). Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of treatment for adalimumab and 14 weeks of treatment for infliximab. From 1 June 2017, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alpha antagonist. For second and subsequent courses of PBS-subsidised TNF-alpha antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course. Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats. (b) Continuing treatment. Following the completion of an initial treatment course with a specific TNF-alpha antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alpha antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alpha antagonist supply. Assessments of response to a course of PBS-subsidised

treatment must be submitted the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist.(2) Swapping treatment. Once initial treatment with the first PBS-subsidised TNF-alfa antagonist is approved, a patient may swap if eligible to the alternate TNF-alfa antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Ulcerative Colitis Activity Index (PUCAI) Score, confirmation of ulcerative colitis disease), or the prior conventional therapies of corticosteroids or immunosuppressives. A patient may trial an alternate agent at any time, regardless of whether they are receiving treatment (initial or continuing) with infliximab or adalimumab at the time of the application. However, a patient cannot swap to a particular TNF-alfa antagonist if they have failed to respond to prior treatment with that drug two times within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these swapping arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction. To avoid confusion, an application for a patient who wishes to swap to the alternate TNF-alfa antagonist (where eligible in terms of disease severity) should be accompanied by the approved authority prescription or remaining repeats for the TNF-alfa antagonist the patient is ceasing.(3) Baseline measurements to determine response. The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PUCAI submitted with the first authority application for a TNF-alfa antagonist. However, prescribers may provide new baseline measurements any time other than when an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy. A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF-alfa antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the PUCAI score is measured.(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab. A patient who commenced treatment with adalimumab for moderate to severe ulcerative colitis prior to 1 June 2017 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction. 'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a continuing patient.

Note Special Pricing Arrangements apply.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment (new patient or Re commencement of treatment after more than 5 years break in therapy - Initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg (for a child, 1 to 2 mg/kg up to 40 mg) prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score); OR
- Patient must have a Paediatric Ulcerative Colitis Activity Index (PUCAI) Score greater than or equal to 30 if aged 6 to 17 years; OR
- Patient must have previously received induction therapy with this drug for an acute severe episode of ulcerative colitis in the last 4 months and demonstrated an adequate response to induction therapy by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a PUCAI score less than 10 (if aged 6 to 17 years).

Population criteria:

- Patient must be 6 years of age or older.

Applications for authorisation of initial treatment must be in writing and must include:

- (a) a completed authority prescription form; and

- (b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:
- (i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the signed patient acknowledgement or guardian acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, or to be administered at 8-weekly intervals for patients who have received prior treatment for an acute severe episode, will be authorised.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic, partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) score must be no more than 1 month old at the time of application.

Where treatment for an acute severe episode has occurred, an adequate response to induction therapy needs to be demonstrated by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 (if aged 6 to 17 years), within the first 12 weeks of receiving this drug for acute severe ulcerative colitis.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 within the first 12 weeks of receiving this drug for ulcerative colitis, or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or have failed to maintain a PUCAI score less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose for patients administered doses at weeks 0, 2 and 6 (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

The patient or guardian (required if patient is aged 6 to 17 years) must have signed a patient acknowledgement indicating that he or she understands and acknowledges that the PBS-subsidised treatment will cease if he or she does not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 while receiving treatment with this drug, if aged 6 to 17 years.

Patients who have failed to maintain a partial Mayo clinic score of less than or equal to 2, with no subscore greater than 1, or, patients who have failed to maintain a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg.

The authority application must be made in writing

Up to a maximum of 2 repeats will be authorised.

Note No applications for increased repeats will be authorised.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to

Friday).
Written application for authority approval should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Change or Re-commencement of treatment after a break in therapy (Initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with adalimumab, infliximab or vedolizumab for this condition in this treatment cycle; OR
- Patient must have previously received PBS-subsidised treatment with adalimumab or infliximab for this condition in this treatment cycle if aged 6 to 17 years, **AND**
- Patient must not have failed PBS-subsidised treatment with infliximab for this condition in the current treatment cycle; OR
- Patient must not have failed PBS-subsidised treatment with infliximab for this condition in the current treatment cycle more than once if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of this drug within the timelines specified in the relevant restriction. If the response assessment to the previous course of this drug is not submitted as detailed in the relevant restriction, the patient will be deemed to have failed therapy with this drug.

Applications for authorisation of initial treatment must be in writing and must include: (a) a completed authority prescription form; and (b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following: (i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy];

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg.

Up to a maximum of 2 repeats will be authorised.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (Change or Recommencement of treatment after a break in therapy) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 and Initial 2 restrictions) or 2 repeats (Continuing restriction).

Population criteria:

- Patient must be 6 years of age or older.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services.

infliximab 100 mg injection, 1 vial

10184B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	604.99	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB**Authority required**

Severe psoriatic arthritis

Treatment Phase: Initial treatment – Initial 1 (new patient or patient recommencing treatment after a break of 5 years or more)

Clinical criteria:

- Patient must have severe active psoriatic arthritis, **AND**
- Patient must have received no prior PBS-subsidised treatment with a biological agent for this condition; OR
- Patient must have received no PBS-subsidised treatment with a biological agent for at least 5 years if they have previously received PBS-subsidised treatment with a biological agent for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be an adult.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

For the purposes of this restriction 'biological agent' means adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab or ustekinumab.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and
either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form; and
- (3) a signed patient acknowledgement.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats may be authorised.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to the Department of Human Services no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Note Patients who fail to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this Treatment Cycle. Patients may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment – Initial 2 (change or recommencement of treatment)

Clinical criteria:

- Patient must have a documented history of severe active psoriatic arthritis, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological agents within this Treatment Cycle, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with this drug during the current Treatment Cycle, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be an adult.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

For the purposes of this restriction 'biological agent' means adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab or ustekinumab.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats may be authorised.

Applications for a patient who has previously received PBS-subsidised treatment with this drug within this Treatment Cycle and who wishes to recommence therapy with this drug within this same Cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug.

Where the most recent course of PBS-subsidised treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised biological therapy or, under this restriction, for patients who have received previous PBS-subsidised biological therapy), the patient must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must have been submitted no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised treatment with this drug was approved under the continuing treatment criteria, the patient must have been assessed for response, and the assessment submitted no later than 4 weeks from the date that course was ceased.

Where a response assessment was not submitted within these timeframes, the patient will be deemed to have failed to respond to treatment.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Note The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to the Department of Human Services no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Note Patients who fail to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this Treatment Cycle. Patients may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time.

Where the term 'biological agents' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab only.

Patients receiving PBS-subsidised treatment for psoriatic arthritis are able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle. There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 1 or Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and secukinumab, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients - ustekinumab and secukinumab only.

For patients who commenced treatment with ustekinumab for psoriatic arthritis prior to 1 May 2016 and for patients who commenced treatment with secukinumab for psoriatic arthritis prior to 1 October 2016, applications for initial PBS-subsidised treatment as continuing therapy may be made under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Within a Treatment Cycle patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment application is submitted within a treatment Cycle and these revised baseline measurements will be used to assess response.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient or patient recommencing treatment after a break of 5 years or more) or Initial 2 (change or recommencement of treatment) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient or patient recommencing treatment after a break of 5 years or more) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (change or recommencement of treatment) restriction to complete 22 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 22 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval for sufficient therapy to complete a maximum of 22 weeks of treatment should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe active psoriatic arthritis, **AND**
- Patient must have received this drug as their most recent course of PBS-subsidised treatment with a biological agent for this condition in the current Treatment Cycle, **AND**
- Patient must demonstrate, at the time of application, an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be an adult.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

For the purposes of this restriction 'biological agent' means adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab or ustekinumab.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

All applications for continuing treatment with this drug must include a measurement of response to the most recent course of PBS-subsidised therapy. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with this drug, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

Where a response assessment is not submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats may be authorised.

Note Patients who fail to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this Treatment Cycle. Patients may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time.

Where the term 'biological agents' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab only.

Patients receiving PBS-subsidised treatment for psoriatic arthritis are able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle. There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and

(ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and

(iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 1 or Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and secukinumab, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients - ustekinumab and secukinumab only.

For patients who commenced treatment with ustekinumab for psoriatic arthritis prior to 1 May 2016 and for patients who

commenced treatment with secukinumab for psoriatic arthritis prior to 1 October 2016, applications for initial PBS-subsidised treatment as continuing therapy may be made under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Within a Treatment Cycle patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment application is submitted within a treatment Cycle and these revised baseline measurements will be used to assess response.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment should be forwarded to:

infliximab 100 mg injection, 1 vial

6496X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	604.99	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ LANREOTIDE

Authority required

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (6 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

lanreotide 30 mg modified release injection [1 vial] (&) inert substance diluent [2 mL ampoule], 1 pack

6332G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*1472.15	Somatuline LA [IS]

■ LANREOTIDE

Authority required

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Authority required

Functional carcinoid tumour

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 120 mg every 28 days.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

lanreotide 60 mg/0.5 mL injection, 0.5 mL syringe

6423C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2602.65	Somatuline Autogel [IS]

lanreotide 90 mg/0.5 mL injection, 0.5 mL syringe

6424D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3448.15	Somatuline Autogel [IS]

lanreotide 120 mg/0.5 mL injection, 0.5 mL syringe

6425E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*4303.15	Somatuline Autogel [IS]

■ OCTREOTIDE**Authority required**

Acromegaly

Clinical criteria:

- The condition must be controlled with octreotide immediate release injections, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission

Authority required

Functional carcinoid tumour

Clinical criteria:

- Patient must have achieved symptom control on octreotide immediate release injections, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with octreotide immediate release injections.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

Authority required

Vasoactive intestinal peptide secreting tumour (VIPoma)

Clinical criteria:

- Patient must have achieved symptom control on octreotide immediate release injections, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with octreotide immediate release injections.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

octreotide 30 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10558Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*4402.07	Sandostatin LAR [NV]

octreotide 10 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10566D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2660.87	Sandostatin LAR [NV]

octreotide 20 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10549F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3526.77	Sandostatin LAR [NV]

■ OCTREOTIDE**Authority required**

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks, **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment at a dose of 100 micrograms 3 time daily, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission

Authority required

Functional carcinoid tumour

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months' therapy.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

Authority required

Vasoactive intestinal peptide secreting tumour (VIPoma)

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months' therapy.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

octreotide 100 microgram/mL injection, 5 x 1 mL ampoules

6228T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*1283.53	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.1 [NV]

octreotide 500 microgram/mL injection, 5 x 1 mL ampoules

6229W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*6241.63	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.5 [NV]

octreotide 50 microgram/mL injection, 5 x 1 mL ampoules

6227R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*651.01	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.05 [NV]

▪ **OMALIZUMAB**

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

Clinical criteria:

- The condition must be based on both physical examination and patient history (to exclude any factors that may be triggering the urticaria), **AND**
- Patient must have experienced itch and hives that persist on a daily basis for at least 6 weeks despite treatment with H1 antihistamines, **AND**
- Patient must have failed to achieve an adequate response after a minimum of 2 weeks treatment with a standard therapy, **AND**
- Patient must not receive more than 12 weeks of treatment under this restriction.

A standard therapy is defined as a combination of therapies that includes H1 antihistamines at maximally tolerated doses in accordance with clinical guidelines, and one of the following:

- 1) a H2 receptor antagonist (150 mg twice per day); or
- 2) a leukotriene receptor antagonist (LTRA) (10 mg per day); or
- 3) doxepin (up to 25 mg three times a day)

If the requirement for treatment with H1 antihistamines and a H2 receptor antagonist, or a leukotriene receptor antagonist or doxepin cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the authority application.

A failure to achieve an adequate response to standard therapy is defined as a current Urticaria Activity Score 7 (UAS7) score of equal to or greater than 28 with an itch score of greater than 8, as assessed while still on standard therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Chronic Spontaneous Urticaria Omalizumab Initial PBS Authority Application - Supporting Information Form which must include:
 - (i) demonstration of failure to achieve an adequate response to standard therapy; and
 - (ii) drug names and doses of standard therapies that the patient has failed; and
 - (iii) a signed patient acknowledgment that cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

omalizumab 150 mg/mL injection, 1 mL syringe

11175E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	2	..	*859.95	Xolair [NV]

■ OMALIZUMAB

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

Clinical criteria:

- Patient must have demonstrated a response to the most recent PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks per authorised course of treatment under this restriction.

Note A proportion of patients respond to 150 mg 4-weekly so where a substantial improvement has been obtained with a 300 mg dose it is reasonable to back-titrate dose after initial treatment.

Note Cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Grandfathering treatment

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 September 2017, **AND**
- Patient must have documented history of itch and hives that persisted on a daily basis for at least 6 weeks despite treatment with H1 antihistamines prior to commencing non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must have documented history of failure to achieve an adequate response after a minimum of 2 weeks treatment with a standard therapy prior to commencing non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

A standard therapy is defined as a combination of therapies that includes H1 antihistamines at maximally tolerated doses in accordance with clinical guidelines, and one of the following:

- 1) a H2 receptor antagonist (150 mg twice per day); or
- 2) a leukotriene receptor antagonist (LTRA) (10 mg per day); or
- 3) doxepin (up to 25 mg three times a day)

If the requirement for treatment with H1 antihistamines and a H2 receptor antagonist, or a leukotriene receptor antagonist or doxepin cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or

intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the authority application.

A failure to achieve an adequate response to standard therapy is defined as a current Urticaria Activity Score 7 (UAS7) score of equal to or greater than 28 with an itch score of greater than 8, as assessed while still on standard therapy.

A patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Chronic Spontaneous Urticaria Omalizumab Initial Grandfather PBS Authority Application - Supporting Information Form which must include:
 - (i) demonstration of failure to achieve an adequate response to standard therapy; and
 - (ii) drug names and doses of standard therapies that the patient has failed; and
 - (iii) a signed patient acknowledgment that cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

omalizumab 150 mg/mL injection, 1 mL syringe

11163M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*859.95	Xolair [NV]

■ PASIREOTIDE

Caution Careful monitoring of patients is mandatory due to high risk of developing hyperglycaemia

Note Special Pricing Arrangements apply.

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have a mean growth hormone (GH) level greater than 2.5 micrograms per litre, **AND**
- Patient must have an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) level greater than 1.3 times the upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

Population criteria:

- Patient must be aged 18 years or older.

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

Failure to achieve biochemical control is defined as:

- 1) Growth hormone level is greater than 2.5 mcg/L; and
- 2) IGF-1 level is greater than 1.3 times the age- and sex-adjusted ULN

In a patient treated with radiotherapy, pasireotide should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pasireotide should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as:

- 1) Growth hormone (GH) levels of less than 2.5 mcg/L; and
- 2) normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1)

The authority application must be made in writing and must include:

- a) a completed authority prescription form; and
- b) a completed Acromegaly PBS Authority Application - Supporting Information Form; and
- c) a signed patient acknowledgment; and
- d) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy must be provided; and a copy of GH and IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided; and
- e) a recent copy of GH and IGF-1 levels must be provided.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Grandfathering treatment

Clinical criteria:

- Patient must have received non-PBS treatment with this drug for this condition prior to 1 September 2016.

Population criteria:

- Patient must be aged 18 years or older.

In a patient treated with radiotherapy, pasireotide should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pasireotide should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as:

- 1) Growth hormone (GH) levels of less than 2.5 mcg/L; and
- 2) normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1)

The authority application must be made in writing and must include:

- a) a completed authority prescription form; and
- b) a completed Acromegaly PBS Authority Application - Supporting Information Form; and
- c) a signed patient acknowledgment; and
- d) in a patient who has previously been treated with radiotherapy for this condition, the date of completion of radiotherapy must be provided; and a copy of GH and IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

Population criteria:

- Patient must be aged 18 years or older.

In a patient treated with radiotherapy, pasireotide should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pasireotide should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as:

- 1) Growth hormone (GH) levels of less than 2.5 mcg/L; and
- 2) normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1)

In a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy and the GH and IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided at the time of approval.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Written applications for authorisation under this criterion should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

pasireotide 20 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10880P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7847.15	Signifor LAR [NV]

pasireotide 40 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10884W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7847.15	Signifor LAR [NV]

pasireotide 60 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10887B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7847.15	Signifor LAR [NV]

■ PEGVISOMANT

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Note Special Pricing Arrangements apply.

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must be an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) concentration greater than 1.3 times upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue.

Somatostatin analogues include octreotide, lanreotide and pasireotide

Failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide is defined as:

- 1) Growth hormone level greater than 2.5 mcg/L; and
- 2) IGF-1 level is greater than 1.3 times the age- and sex-adjusted ULN

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1).

Two completed authority prescriptions should be submitted with the initial application for this drug. One prescription should be for the loading dose of 80 mg for a quantity of 4 vials of 20 mg with no repeats. The second prescription should be for subsequent doses, starting from 10 mg daily, and allowing dose adjustments in increments of 5 mg based on serum IGF-1 levels measured every 4 to 6 weeks in order to maintain the serum IGF-1 level within the age-adjusted normal range based on the dosage recommendations in the TGA-approved Product Information.

The authority application must be made in writing and must include:

- a) two completed authority prescription forms ; and
- b) a completed Acromegaly Pegvisomant initial PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy, the date and result of IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy; and
- d) a recent result of the IGF-1 level and the date of assessment ; and
- e) demonstration of failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide

pegvisomant 20 mg injection [1 vial] (&) inert substance diluent [1 x 1 mL vial], 1 pack

11166Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	..	..	*586.59	Somavert [PF]

■ PEGVISOMANT

Note Special Pricing Arrangements apply.

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must be an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) concentration greater than 1.3 times upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue.

Somatostatin analogues include octreotide, lanreotide and pasireotide

Failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide is defined as:

- 1) Growth hormone level greater than 2.5 mcg/L; and
- 2) IGF-1 level is greater than 1.3 times the age- and sex-adjusted ULN

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1).

Two completed authority prescriptions should be submitted with the initial application for this drug. One prescription should be for the loading dose of 80 mg for a quantity of 4 vials of 20 mg with no repeats. The second prescription should be for subsequent doses, starting from 10 mg daily, and allowing dose adjustments in increments of 5 mg based on serum IGF-1 levels measured every 4 to 6 weeks in order to maintain the serum IGF-1 level within the age-adjusted normal range based on the dosage recommendations in the TGA-approved Product Information.

The authority application must be made in writing and must include:

- a) two completed authority prescription forms ; and
- b) a completed Acromegaly Pegvisomant initial PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy, the date and result of IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy; and
- d) a recent result of the IGF-1 level and the date of assessment ; and
- e) demonstration of failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs Programs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition, **AND**
 - The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue, **AND**
 - The treatment must cease if IGF-1 is not lower after 3 months of pegvisomant treatment at the maximum tolerated dose.
- Somatostatin analogues include octreotide, lanreotide and pasireotide

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1).

In a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy must be provided; and a copy of IGF-1 level taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided at the time of application.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Acromegaly

Treatment Phase: Grandfathering

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 September 2017, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue, **AND**
- Patient must have had a documented age- and sex- adjusted insulin- like factor 1 (IGF-1) concentration greater than 1.3 times upper limit of normal (ULN) prior to commencing non- PBS- subsidised treatment with this drug.

Somatostatin analogues include octreotide, lanreotide and pasireotide

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1). Treatment must be ceased if IGF-1 level is not lower after 3 months of pegvisomant treatment at the maximum tolerated dose.

A patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

The authority application must be made in writing and must include:

- a) a completed authority prescription form; and
- b) a completed Acromegaly Pegvisomant Grandfather PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy, the date and result of IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy; and
- d) a recent result of the IGF-1 level and the date of assessment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

pegvisomant 20 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack

11174D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4225.72	Somavert [PF]

pegvisomant 15 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack

11172B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4225.72	Somavert [PF]

pegvisomant 10 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack

11167R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4225.72	Somavert [PF]

■ USTEKINUMAB

Note It is recommended that an application for continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug.

Note No increase in the maximum number of repeats may be authorised.

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying drugs (bDMDs) for adult patients with severe Crohn disease. Where the term bDMDs appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological disease modifying drugs at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to be in their first cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised bDMD therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of bDMD therapy in a treatment cycle and who has a break in therapy of less

than 5 years, may commence a further course of treatment within the same treatment cycle.
 A patient who has failed fewer than 3 trials of bDMD therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.
 There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial 1 - new patients); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Initial 2 - Change or Re-commencement).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMD therapy.

For second and subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

Assessments of response to a course of PBS-subsidised therapy must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMD therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate bDMD therapy at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular bDMD therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to adalimumab, infliximab, vedolizumab or ustekinumab (where eligible in terms of disease severity) should be accompanied by the approved authority prescription or remaining repeats for the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised bDMD therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum

of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.
(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment (new patient - initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; OR
- Patient must have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to steroids, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more months or have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to this drug; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months or have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to this drug; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more months or have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to this drug, **AND**
- Patient must have severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220 if affected by extensive small intestine disease; OR
- Patient must have severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 if not affected by extensive small intestine disease, short gut syndrome or is an ostomy patient, **AND**
- Patient must have evidence of intestinal inflammation and have diagnostic imaging or surgical evidence of short gut syndrome if affected by the syndrome or has an ileostomy or colostomy; OR
- Patient must have radiological evidence of intestinal inflammation if the patient has extensive small intestinal disease affecting more than 50 cm of the small intestine; OR
- Patient must (a) have evidence of intestinal inflammation, including: (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or (ii) faeces: higher than normal lactoferrin or calprotectin level; or (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery; or (b) be assessed clinically as being in a high faecal output state; or (c) be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iv) the date of the most recent clinical assessment; and

(v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application.

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

Authority required

Severe Crohn disease

Treatment Phase: Change or Re-commencement of treatment (initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological disease modifying drug for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition in the current treatment cycle.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following:

(i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or

(ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and

(iii) the date of clinical assessment; and

(iv) the details of prior biological disease modifying drug treatment including the details of date and duration of treatment.

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for 2 vials of 45 mg and no repeats.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological disease modifying drug (bDMD) therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised bDMD treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of bDMD treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of bDMD.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent first dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

ustekinumab 130 mg/26 mL injection, 26 mL vial

11164N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±4	..	..	*16976.39	Stelara [JC]

Highly Specialised Drugs Program (Public Hospital)

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of infliximab, vedolizumab and adalimumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either infliximab, vedolizumab or adalimumab at any one time. From 1 December 2016, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial each of PBS-subsidised infliximab, vedolizumab or adalimumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with infliximab, vedolizumab or adalimumab while they continue to show a response to therapy. A patient who received PBS-subsidised infliximab, vedolizumab or adalimumab treatment prior to 1 December 2016 is considered to be in their first cycle as of 1 December 2016. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised infliximab, vedolizumab or adalimumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised infliximab, vedolizumab or adalimumab treatment in the most recent cycle to the date of the first application for initial treatment with infliximab, vedolizumab or adalimumab under the new treatment cycle. A patient who has failed fewer than 3 trials of either infliximab, vedolizumab or adalimumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with infliximab, vedolizumab and adalimumab after therapy after 1 December 2016 .

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) an adult patient has received no prior PBS-subsidised treatment with infliximab, vedolizumab or adalimumab in this treatment cycle and wishes to commence such therapy (Initial 1); or
- (ii) an adult patient has received prior PBS-subsidised (initial or continuing) infliximab, vedolizumab or adalimumab therapy and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; or
- (iii) an adult patient wishes to re-commence treatment with infliximab, vedolizumab or adalimumab following a break in PBS-subsidised therapy with the same agent (Initial 2).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab , 14 weeks of therapy for infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab, and this assessment must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alfa antagonist. For second and subsequent courses of PBS-subsidised TNF-alfa antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is provided to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(b) Continuing treatment.

Following the completion of an initial treatment course with infliximab, vedolizumab or adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment. Assessments of response to a course of PBS-subsidised therapy must be provided to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not provided to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate infliximab, vedolizumab or adalimumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with infliximab, vedolizumab or adalimumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction. To avoid confusion, an application for a patient who wishes to swap to the alternate infliximab, vedolizumab or adalimumab therapy should be accompanied by the approved authority prescription or remaining repeats for

the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Mayo clinic score or partial Mayo clinic score submitted with the first authority application for infliximab, vedolizumab or adalimumab. However, prescribers may provide new baseline measurements any time other than when an initial treatment authority application is provided within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised infliximab, vedolizumab or adalimumab therapy of at least 5 years, must requalify for initial treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the Mayo score is measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab.

A patient who commenced treatment with adalimumab for moderate to severe ulcerative colitis prior to 1 December 2016 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a continuing patient.

Note TREATMENT OF PAEDIATRIC PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with infliximab or adalimumab for moderate to severe ulcerative colitis; and infliximab for acute severe ulcerative colitis.

Where the term 'tumour necrosis factor (TNF) alpha antagonist' appears in the following NOTES and restrictions, it refers to infliximab and adalimumab only. A patient is eligible for PBS-subsidised treatment with only 1 of the 2 TNF-alpha antagonists at any one time. Infliximab and adalimumab are PBS-subsidised for moderate to severe disease while only infliximab is PBS-subsidised for acute severe disease. From 1 June 2017, under the PBS, all will be able to commence a treatment cycle where they may trial each PBS-subsidised TNF-alpha antagonist without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a TNF-alpha antagonist while they continue to show a response to therapy. A patient who received PBS-subsidised TNF-alpha antagonist treatment prior to 1 June 2017 is considered to be in their first cycle as of 1 June 2017. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised TNF-alpha antagonist more than twice. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised TNF-alpha antagonist therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised TNF-alpha antagonist treatment in the most recent cycle to the date of the first application for initial treatment with a TNF-alpha antagonist under the new treatment cycle. A patient who has failed fewer than 3 trials of TNF-alpha antagonists in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle. A patient who has failed fewer than 3 trials of TNF-alpha antagonists in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle. There is no limit to the number of treatment cycles a patient may undertake in their lifetime. (1) How to prescribe PBS-subsidised TNF-alpha antagonist therapy after 1 June 2017. (a) Initial treatment. Applications for initial treatment should be made where: (i) a patient has received no prior PBS-subsidised TNF-alpha antagonist treatment in this treatment cycle and wishes to commence such therapy (Initial 1); or (ii) a patient has received prior PBS-subsidised (initial or continuing) treatment with a TNF-alpha antagonist and wishes to trial an alternate agent (Initial 2) [further details are under 'Swapping treatment' below]; or (iii) a patient wishes to re-commence treatment with a specific TNF-alpha antagonist following a break in PBS-subsidised therapy with that agent (Initial 2). Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of treatment for adalimumab and 14 weeks of treatment for infliximab. From 1 June 2017, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF-alpha antagonist. For second and subsequent courses of PBS-subsidised TNF-alpha antagonist treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course. Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats. (b) Continuing treatment. Following the completion of an initial treatment course with a specific TNF-alpha antagonist, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing TNF-alpha antagonist treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted TNF-alpha antagonist supply. Assessments of response to a course of PBS-subsidised

treatment must be submitted the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that TNF- α antagonist.(2) Swapping treatment. Once initial treatment with the first PBS-subsidised TNF- α antagonist is approved, a patient may swap if eligible to the alternate TNF- α antagonist within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Ulcerative Colitis Activity Index (PUCAI) Score, confirmation of ulcerative colitis disease), or the prior conventional therapies of corticosteroids or immunosuppressives. A patient may trial an alternate agent at any time, regardless of whether they are receiving treatment (initial or continuing) with infliximab or adalimumab at the time of the application. However, a patient cannot swap to a particular TNF- α antagonist if they have failed to respond to prior treatment with that drug two times within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these swapping arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction. To avoid confusion, an application for a patient who wishes to swap to the alternate TNF- α antagonist (where eligible in terms of disease severity) should be accompanied by the approved authority prescription or remaining repeats for the TNF- α antagonist the patient is ceasing.(3) Baseline measurements to determine response. The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PUCAI submitted with the first authority application for a TNF- α antagonist. However, prescribers may provide new baseline measurements any time other than when an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy. A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised TNF- α antagonist therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the PUCAI score is measured.(5) Patients 'grandfathered' onto PBS-subsidised treatment with adalimumab. A patient who commenced treatment with adalimumab for moderate to severe ulcerative colitis prior to 1 June 2017 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment with adalimumab will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further applications for treatment with adalimumab will be assessed under the continuing treatment restriction. 'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a continuing patient.

Note Special Pricing Arrangements apply.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment (new patient or Re commencement of treatment after more than 5 years break in therapy - Initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg (for a child, 1 to 2 mg/kg up to 40 mg) prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score); OR
- Patient must have a Paediatric Ulcerative Colitis Activity Index (PUCAI) Score greater than or equal to 30 if aged 6 to 17 years; OR
- Patient must have previously received induction therapy with this drug for an acute severe episode of ulcerative colitis in the last 4 months and demonstrated an adequate response to induction therapy by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a PUCAI score less than 10 (if aged 6 to 17 years).

Population criteria:

- Patient must be 6 years of age or older.

Applications for authorisation of initial treatment must be in writing and must include:

- (a) a completed authority prescription form; and

- (b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:
- (i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the signed patient acknowledgement or guardian acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, or to be administered at 8-weekly intervals for patients who have received prior treatment for an acute severe episode, will be authorised.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic, partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) score must be no more than 1 month old at the time of application.

Where treatment for an acute severe episode has occurred, an adequate response to induction therapy needs to be demonstrated by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 (if aged 6 to 17 years), within the first 12 weeks of receiving this drug for acute severe ulcerative colitis.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 within the first 12 weeks of receiving this drug for ulcerative colitis, or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or have failed to maintain a PUCAI score less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose for patients administered doses at weeks 0, 2 and 6 (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

The patient or guardian (required if patient is aged 6 to 17 years) must have signed a patient acknowledgement indicating that he or she understands and acknowledges that the PBS-subsidised treatment will cease if he or she does not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 while receiving treatment with this drug, if aged 6 to 17 years.

Patients who have failed to maintain a partial Mayo clinic score of less than or equal to 2, with no subscore greater than 1, or, patients who have failed to maintain a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg.

Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing

Note No applications for increased repeats will be authorised.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to

Friday).
Written application for authority approval should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Change or Re-commencement of treatment after a break in therapy (Initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with adalimumab, infliximab or vedolizumab for this condition in this treatment cycle; OR
- Patient must have previously received PBS-subsidised treatment with adalimumab or infliximab for this condition in this treatment cycle if aged 6 to 17 years, **AND**
- Patient must not have failed PBS-subsidised treatment with infliximab for this condition in the current treatment cycle; OR
- Patient must not have failed PBS-subsidised treatment with infliximab for this condition in the current treatment cycle more than once if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of this drug within the timelines specified in the relevant restriction. If the response assessment to the previous course of this drug is not submitted as detailed in the relevant restriction, the patient will be deemed to have failed therapy with this drug.

Applications for authorisation of initial treatment must be in writing and must include: (a) a completed authority prescription form; and (b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following: (i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy];

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg.

Up to a maximum of 2 repeats will be authorised.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (Change or Recommencement of treatment after a break in therapy) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 and Initial 2 restrictions) or 2 repeats (Continuing restriction).

Population criteria:

- Patient must be 6 years of age or older.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services.

infliximab 100 mg injection, 1 vial

10196P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	574.85	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB**Authority required**

Severe psoriatic arthritis

Treatment Phase: Initial treatment – Initial 1 (new patient or patient recommencing treatment after a break of 5 years or more)

Clinical criteria:

- Patient must have severe active psoriatic arthritis, **AND**
- Patient must have received no prior PBS-subsidised treatment with a biological agent for this condition; OR
- Patient must have received no PBS-subsidised treatment with a biological agent for at least 5 years if they have previously received PBS-subsidised treatment with a biological agent for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be an adult.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

For the purposes of this restriction 'biological agent' means adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab or ustekinumab.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form; and
- (3) a signed patient acknowledgement.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats may be authorised.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to the Department of Human Services no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Note Patients who fail to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this Treatment Cycle. Patients may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment – Initial 2 (change or recommencement of treatment)

Clinical criteria:

- Patient must have a documented history of severe active psoriatic arthritis, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological agent for this condition in this Treatment Cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological agents within this Treatment Cycle, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with this drug during the current Treatment Cycle, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be an adult.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

For the purposes of this restriction 'biological agent' means adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab or ustekinumab.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 3 repeats may be authorised.

Applications for a patient who has previously received PBS-subsidised treatment with this drug within this Treatment Cycle and who wishes to recommence therapy with this drug within this same Cycle, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug.

Where the most recent course of PBS-subsidised treatment was approved under either of the initial treatment restrictions (i.e. for patients with no prior PBS-subsidised biological therapy or, under this restriction, for patients who have received previous PBS-subsidised biological therapy), the patient must have been assessed for response following a minimum of 12 weeks of therapy. This assessment must have been submitted no later than 4 weeks from the date that course was ceased.

Where the most recent course of PBS-subsidised treatment with this drug was approved under the continuing treatment criteria, the patient must have been assessed for response, and the assessment submitted no later than 4 weeks from the date that course was ceased.

Where a response assessment was not submitted within these timeframes, the patient will be deemed to have failed to respond to treatment.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Note The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to the Department of Human Services no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Note Patients who fail to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this Treatment Cycle. Patients may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time.

Where the term 'biological agents' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab only.

Patients receiving PBS-subsidised treatment for psoriatic arthritis are able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle. There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and
- (ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and
- (iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 1 or Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and secukinumab, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients - ustekinumab and secukinumab only.

For patients who commenced treatment with ustekinumab for psoriatic arthritis prior to 1 May 2016 and for patients who commenced treatment with secukinumab for psoriatic arthritis prior to 1 October 2016, applications for initial PBS-subsidised treatment as continuing therapy may be made under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Within a Treatment Cycle patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment application is submitted within a treatment Cycle and these revised baseline measurements will be used to assess response.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient or patient recommencing treatment after a break of 5 years or more) or Initial 2 (change or recommencement of treatment) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient or patient recommencing treatment after a break of 5 years or more) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (change or recommencement of treatment) restriction to complete 22 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 22 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval for sufficient therapy to complete a maximum of 22 weeks of treatment should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe active psoriatic arthritis, **AND**
- Patient must have received this drug as their most recent course of PBS-subsidised treatment with a biological agent for this condition in the current Treatment Cycle, **AND**
- Patient must demonstrate, at the time of application, an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be an adult.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

For the purposes of this restriction 'biological agent' means adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab or ustekinumab.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications.

All applications for continuing treatment with this drug must include a measurement of response to the most recent course of PBS-subsidised therapy. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with this drug, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

Where a response assessment is not submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg. Up to a maximum of 2 repeats may be authorised.

Note Patients who fail to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this Treatment Cycle. Patients may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological agent was approved in this Cycle and the date of the first application under the new Cycle.

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological agents adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab for adult patients with severe active psoriatic arthritis.

Patients are eligible for PBS-subsidised treatment with only 1 of the above biological agents at any 1 time.

Where the term 'biological agents' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, secukinumab and ustekinumab only.

Patients receiving PBS-subsidised treatment for psoriatic arthritis are able to commence a 'Biological Treatment Cycle' (Cycle), where they may trial biological agents without having to experience a disease flare when swapping to the alternate agent. Under these interchangeability arrangements, within a single Cycle, patients may receive long-term treatment with a biological agent as long as they sustain a response to therapy.

Following demonstration of response to initial treatment, these biological agents are available under the PBS for continuing treatment as set out in the continuing treatment restriction for each agent.

Once patients have either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single Cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological therapy before they are eligible to commence another Cycle [further details are under '(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last approval for PBS-subsidised treatment was granted in the most recent Cycle to the date of the first application for initial treatment with a biological agent under the new Cycle.

Within the same Cycle, patients are not allowed to fail, or cease to respond to, the same PBS-subsidised biological agent more than once. Therefore once a patient fails to meet the response criteria for any biological agent, they must change to an alternate agent which they have not previously failed, if they wish to continue PBS-subsidised biological treatment.

Patients for whom a break in PBS-subsidised therapy of less than 5 years has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, may commence a further course of treatment within that Cycle.

Patients for whom a break in PBS-subsidised therapy of 5 years or more has occurred, and, who have failed therapy fewer than 3 times within a particular treatment Cycle, as defined in the relevant restriction, are eligible to commence a new Cycle. There is no limit to the number of Biological Treatment Cycles a patient may undertake in their lifetime.

How to prescribe biological agents for the treatment of severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) patients have received no prior PBS-subsidised biological treatment and wish to commence such therapy (Initial 1); and

(ii) patients have received prior PBS-subsidised biological therapy and wish to trial an alternate agent (Initial 2) [further details are under 'Swapping therapy' below]; and

(iii) patients wish to re-commence treatment with a specific biological agent following a break in PBS-subsidised therapy with that specific agent (Initial 1 or Initial 2).

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept and golimumab and secukinumab, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that patients be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological agent supply.

Patients must be assessed for response to any course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

Grandfather patients - ustekinumab and secukinumab only.

For patients who commenced treatment with ustekinumab for psoriatic arthritis prior to 1 May 2016 and for patients who

commenced treatment with secukinumab for psoriatic arthritis prior to 1 October 2016, applications for initial PBS-subsidised treatment as continuing therapy may be made under the relevant initial treatment restriction (Initial 3). These patients access the PBS interchangeability arrangements in the same way as new patients who have not been treated with any biological agent prior to PBS listing of that agent.

Applications for initial PBS-subsidised treatment for grandfather patients will provide for a maximum of 24 weeks of treatment for all agents. Approval will be based on the criteria included in the relevant restriction.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological agent, patients may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. Patients are eligible to receive continuing biological treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Patients must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted within these timeframes, patients will be deemed to have failed to respond to treatment with that biological agent.

(3) Swapping therapy.

Once an authority for initial treatment with the first PBS-subsidised biological agent is approved, patients may swap to an alternate biological agent without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements.

Patients may swap to an alternate biological agent at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological agent at the time of the application or not.

Within a Treatment Cycle patients may alternate between therapy with any biological agent of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological agent previously; or
- (ii) they have demonstrated an adequate response to that particular biological agent if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this Treatment Cycle.

To ensure patients receive the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, applications for patients who wish to swap to an alternate biological agent should be accompanied by the approved authority prescription or remaining repeats for the biological agent the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological agent. However, prescribers may provide new baseline measurements any time that an initial treatment application is submitted within a treatment Cycle and these revised baseline measurements will be used to assess response.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be provided to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

Patients who wish to trial a second or subsequent treatment Cycle following a break in PBS-subsidised biological therapy of at least 5 years, must requalify for initial treatment with respect to both the indices of disease severity. Patients must have received treatment with methotrexate and sulfasalazine or leflunomide, at an adequate dose, for a minimum of 3 months at the time the ESR or CRP levels and the active joint counts are measured.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment should be forwarded to:

infliximab 100 mg injection, 1 vial

5756Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	574.85	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ LANREOTIDE

Authority required (STREAMLINED)

7042

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (6 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

lanreotide 30 mg modified release injection [1 vial] (&) inert substance diluent [2 mL ampoule], 1 pack

5776B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*1425.00	Somatuline LA [IS]

■ LANREOTIDE

Authority required (STREAMLINED)

7025

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Authority required (STREAMLINED)

4575

Functional carcinoid tumour

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 120 mg every 28 days.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

lanreotide 60 mg/0.5 mL injection, 0.5 mL syringe

5777C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2555.50	Somatuline Autogel [IS]

lanreotide 90 mg/0.5 mL injection, 0.5 mL syringe

5778D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3401.00	Somatuline Autogel [IS]

lanreotide 120 mg/0.5 mL injection, 0.5 mL syringe

5779E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*4256.00	Somatuline Autogel [IS]

■ OCTREOTIDE**Authority required (STREAMLINED)****7029**

Acromegaly

Clinical criteria:

- The condition must be controlled with octreotide immediate release injections, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission

Authority required (STREAMLINED)**5901**

Functional carcinoid tumour

Clinical criteria:

- Patient must have achieved symptom control on octreotide immediate release injections, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with octreotide immediate release injections.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

Authority required (STREAMLINED)**5906**

Vasoactive intestinal peptide secreting tumour (VIPoma)

Clinical criteria:

- Patient must have achieved symptom control on octreotide immediate release injections, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with octreotide immediate release injections.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

octreotide 30 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10550G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*4354.92	Sandostatin LAR [NV]

octreotide 10 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10543X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2613.72	Sandostatin LAR [NV]

octreotide 20 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10533J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3479.62	Sandostatin LAR [NV]

■ OCTREOTIDE**Authority required (STREAMLINED)****7028**

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**

- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks, **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment at a dose of 100 micrograms 3 time daily, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission

Authority required (STREAMLINED)

6390

Functional carcinoid tumour

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months' therapy.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

Authority required (STREAMLINED)

6369

Vasoactive intestinal peptide secreting tumour (VIPoma)

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months' therapy.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

octreotide 100 microgram/mL injection, 5 x 1 mL ampoules

9509L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*1236.42	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.1 [NV]

octreotide 500 microgram/mL injection, 5 x 1 mL ampoules

9510M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*6194.52	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.5 [NV]

octreotide 50 microgram/mL injection, 5 x 1 mL ampoules

9508K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*619.02	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.05 [NV]

▪ **OMALIZUMAB**

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

Clinical criteria:

- The condition must be based on both physical examination and patient history (to exclude any factors that may be triggering the urticaria), **AND**

- Patient must have experienced itch and hives that persist on a daily basis for at least 6 weeks despite treatment with H1 antihistamines, **AND**
- Patient must have failed to achieve an adequate response after a minimum of 2 weeks treatment with a standard therapy, **AND**

- Patient must not receive more than 12 weeks of treatment under this restriction.

A standard therapy is defined as a combination of therapies that includes H1 antihistamines at maximally tolerated doses in accordance with clinical guidelines, and one of the following:

- 1) a H2 receptor antagonist (150 mg twice per day); or
- 2) a leukotriene receptor antagonist (LTRA) (10 mg per day); or
- 3) doxepin (up to 25 mg three times a day)

If the requirement for treatment with H1 antihistamines and a H2 receptor antagonist, or a leukotriene receptor antagonist or doxepin cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the authority application.

A failure to achieve an adequate response to standard therapy is defined as a current Urticaria Activity Score 7 (UAS7) score of equal to or greater than 28 with an itch score of greater than 8, as assessed while still on standard therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Chronic Spontaneous Urticaria Omalizumab Initial PBS Authority Application - Supporting Information Form which must include:

- (i) demonstration of failure to achieve an adequate response to standard therapy; and
- (ii) drug names and doses of standard therapies that the patient has failed; and
- (iii) a signed patient acknowledgment that cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

omalizumab 150 mg/mL injection, 1 mL syringe

11176F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	2	..	*820.00	Xolair [NV]

■ OMALIZUMAB

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

Clinical criteria:

- Patient must have demonstrated a response to the most recent PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks per authorised course of treatment under this restriction.

Note A proportion of patients respond to 150 mg 4-weekly so where a substantial improvement has been obtained with a 300 mg dose it is reasonable to back-titrate dose after initial treatment.

Note Cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Grandfathering treatment

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 September 2017, **AND**
- Patient must have documented history of itch and hives that persisted on a daily basis for at least 6 weeks despite treatment with H1 antihistamines prior to commencing non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must have documented history of failure to achieve an adequate response after a minimum of 2 weeks treatment with a standard therapy prior to commencing non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

A standard therapy is defined as a combination of therapies that includes H1 antihistamines at maximally tolerated doses in accordance with clinical guidelines, and one of the following:

- 1) a H2 receptor antagonist (150 mg twice per day); or
- 2) a leukotriene receptor antagonist (LTRA) (10 mg per day); or
- 3) doxepin (up to 25 mg three times a day)

If the requirement for treatment with H1 antihistamines and a H2 receptor antagonist, or a leukotriene receptor antagonist or doxepin cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the authority application.

A failure to achieve an adequate response to standard therapy is defined as a current Urticaria Activity Score 7 (UAS7) score of equal to or greater than 28 with an itch score of greater than 8, as assessed while still on standard therapy.

A patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Chronic Spontaneous Urticaria Omalizumab Initial Grandfather PBS Authority Application - Supporting Information Form which must include:
 - (i) demonstration of failure to achieve an adequate response to standard therapy; and
 - (ii) drug names and doses of standard therapies that the patient has failed; and
 - (iii) a signed patient acknowledgment that cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

omalizumab 150 mg/mL injection, 1 mL syringe

11168T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*820.00	Xolair [NV]

■ PASIREOTIDE

Caution Careful monitoring of patients is mandatory due to high risk of developing hyperglycaemia

Note Special Pricing Arrangements apply.

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have a mean growth hormone (GH) level greater than 2.5 micrograms per litre, **AND**
- Patient must have an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) level greater than 1.3 times the upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

Population criteria:

- Patient must be aged 18 years or older.

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

Failure to achieve biochemical control is defined as:

- 1) Growth hormone level is greater than 2.5 mcg/L; and
- 2) IGF-1 level is greater than 1.3 times the age- and sex-adjusted ULN

In a patient treated with radiotherapy, pasireotide should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pasireotide should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as:

- 1) Growth hormone (GH) levels of less than 2.5 mcg/L; and
- 2) normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1)

The authority application must be made in writing and must include:

- a) a completed authority prescription form; and

-
- b) a completed Acromegaly PBS Authority Application - Supporting Information Form; and
 - c) a signed patient acknowledgment; and
 - d) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy must be provided; and a copy of GH and IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided; and
 - e) a recent copy of GH and IGF-1 levels must be provided.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Grandfathering treatment

Clinical criteria:

- Patient must have received non-PBS treatment with this drug for this condition prior to 1 September 2016.

Population criteria:

- Patient must be aged 18 years or older.

In a patient treated with radiotherapy, pasireotide should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pasireotide should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as:

- 1) Growth hormone (GH) levels of less than 2.5 mcg/L; and
- 2) normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1)

The authority application must be made in writing and must include:

- a) a completed authority prescription form; and
- b) a completed Acromegaly PBS Authority Application - Supporting Information Form; and
- c) a signed patient acknowledgment; and
- d) in a patient who has previously been treated with radiotherapy for this condition, the date of completion of radiotherapy must be provided; and a copy of GH and IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

Population criteria:

- Patient must be aged 18 years or older.

In a patient treated with radiotherapy, pasireotide should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pasireotide should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as:

- 1) Growth hormone (GH) levels of less than 2.5 mcg/L; and
- 2) normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1)

In a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy and the GH and IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided at the time of approval.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

pasireotide 20 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10886Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7800.00	Signifor LAR [NV]

pasireotide 40 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10883T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7800.00	Signifor LAR [NV]

pasireotide 60 mg injection: modified release [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10882R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7800.00	Signifor LAR [NV]

■ PEGVISOMANT

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Note Special Pricing Arrangements apply.

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) concentration greater than 1.3 times upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue.

Somatostatin analogues include octreotide, lanreotide and pasireotide

Failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide is defined as:

- 1) Growth hormone level greater than 2.5 mcg/L; and
- 2) IGF-1 level is greater than 1.3 times the age- and sex-adjusted ULN

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1).

Two completed authority prescriptions should be submitted with the initial application for this drug. One prescription should be for the loading dose of 80 mg for a quantity of 4 vials of 20 mg with no repeats. The second prescription should be for subsequent doses, starting from 10 mg daily, and allowing dose adjustments in increments of 5 mg based on serum IGF-1 levels measured every 4 to 6 weeks in order to maintain the serum IGF-1 level within the age-adjusted normal range based on the dosage recommendations in the TGA-approved Product Information.

The authority application must be made in writing and must include:

- a) two completed authority prescription forms ; and
- b) a completed Acromegaly Pegvisomant initial PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy, the date and result of IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy; and
- d) a recent result of the IGF-1 level and the date of assessment ; and
- e) demonstration of failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide

pegvisomant 20 mg injection [1 vial] (&) inert substance diluent [1 x 1 mL vial], 1 pack

11177G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	..	..	*557.16	Somavert [PF]

■ PEGVISOMANT

Note Special Pricing Arrangements apply.

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) concentration greater than 1.3 times upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue.

Somatostatin analogues include octreotide, lanreotide and pasireotide

Failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide is defined as:

- 1) Growth hormone level greater than 2.5 mcg/L; and
- 2) IGF-1 level is greater than 1.3 times the age- and sex-adjusted ULN

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1).

Two completed authority prescriptions should be submitted with the initial application for this drug. One prescription should be for the loading dose of 80 mg for a quantity of 4 vials of 20 mg with no repeats. The second prescription should be for subsequent doses, starting from 10 mg daily, and allowing dose adjustments in increments of 5 mg based on serum IGF-1 levels measured every 4 to 6 weeks in order to maintain the serum IGF-1 level within the age-adjusted normal range based on the dosage recommendations in the TGA-approved Product Information.

The authority application must be made in writing and must include:

- a) two completed authority prescription forms ; and
- b) a completed Acromegaly Pegvisomant initial PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy, the date and result of IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy; and
- d) a recent result of the IGF-1 level and the date of assessment ; and
- e) demonstration of failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition, **AND**
 - The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue, **AND**
 - The treatment must cease if IGF-1 is not lower after 3 months of pegvisomant treatment at the maximum tolerated dose.
- Somatostatin analogues include octreotide, lanreotide and pasireotide

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1).

In a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy must be provided; and a copy of IGF-1 level taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided at the time of application.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Acromegaly

Treatment Phase: Grandfathering

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 September 2017, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue, **AND**
- Patient must have had a documented age- and sex- adjusted insulin- like factor 1 (IGF-1) concentration greater than 1.3 times upper limit of normal (ULN) prior to commencing non- PBS- subsidised treatment with this drug.

Somatostatin analogues include octreotide, lanreotide and pasireotide

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1). Treatment must be ceased if IGF-1 level is not lower after 3 months of pegvisomant treatment at the maximum tolerated dose.

A patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

The authority application must be made in writing and must include:

- a) a completed authority prescription form; and
- b) a completed Acromegaly Pegvisomant Grandfather PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy, the date and result of IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy; and
- d) a recent result of the IGF-1 level and the date of assessment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

pegvisomant 20 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack

11181L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4178.57	Somavert [PF]

pegvisomant 15 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack

11173C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4178.57	Somavert [PF]

pegvisomant 10 mg injection [30 vials] (&) inert substance diluent [30 x 1 mL vials], 1 pack

11179J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4178.57	Somavert [PF]

▪ USTEKINUMAB

Note It is recommended that an application for continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug.

Note No increase in the maximum number of repeats may be authorised.

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological disease modifying drugs (bDMDs) for adult patients with severe Crohn disease. Where the term bDMDs appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab). Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological disease modifying drugs at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to be in their first cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment

cycle and they must have, at a minimum, a 5-year break in PBS-subsidised bDMD therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of bDMD therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of bDMD therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial 1 - new patients); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Initial 2 - Change or Re-commencement).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that bDMD therapy.

For second and subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

Assessments of response to a course of PBS-subsidised therapy must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that drug.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised bDMD therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial the alternate bDMD therapy at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular bDMD therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to adalimumab, infliximab, vedolizumab or ustekinumab (where eligible in terms of disease severity) should be accompanied by the approved authority prescription or remaining repeats for the therapy the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be provided for all subsequent continuing treatment

applications.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised bDMD therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment (new patient - initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period; OR
- Patient must have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to steroids, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more months or have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to this drug; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months or have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to this drug; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more months or have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contraindication to this drug, **AND**
- Patient must have severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220 if affected by extensive small intestine disease; OR
- Patient must have severity of disease activity which results in a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 if not affected by extensive small intestine disease, short gut syndrome or is an ostomy patient, **AND**
- Patient must have evidence of intestinal inflammation and have diagnostic imaging or surgical evidence of short gut syndrome if affected by the syndrome or has an ileostomy or colostomy; OR
- Patient must have radiological evidence of intestinal inflammation if the patient has extensive small intestinal disease affecting more than 50 cm of the small intestine; OR
- Patient must (a) have evidence of intestinal inflammation, including: (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or (ii) faeces: higher than normal lactoferrin or calprotectin level; or (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery; or (b) be assessed clinically as being in a high faecal output state; or (c) be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:

-
- (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iv) the date of the most recent clinical assessment; and
 - (v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application.

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

Authority required

Severe Crohn disease

Treatment Phase: Change or Re-commencement of treatment (initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological disease modifying drug for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition in the current treatment cycle.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following:

(i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or

(ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and

(iii) the date of clinical assessment; and

(iv) the details of prior biological disease modifying drug treatment including the details of date and duration of treatment.

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for 2 vials of 45 mg and no repeats.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological disease modifying drug (bDMD) therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised bDMD treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of bDMD treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of bDMD.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent first dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

ustekinumab 130 mg/26 mL injection, 26 mL vial

11182M

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
±4	..	..	*16929.24	Stelara [JC]